



TITLE:

# The GIT stability of polarized varieties via discrepancy

AUTHOR(S):

Odaka, Yuji

---

CITATION:

Odaka, Yuji. The GIT stability of polarized varieties via discrepancy. *Annals of Mathematics* 2013, 177(2): 645-661

ISSUE DATE:

2013

URL:

<http://hdl.handle.net/2433/173546>

RIGHT:

© 2013 Department of Mathematics, Princeton University.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。 ; This is not the published version. Please cite only the published version.

# THE GIT STABILITY OF POLARIZED VARIETIES VIA DISCREPANCY

YUJI ODAKA

*Dedicated to Professor Shigefumi Mori on his Kanreki (60th birthday)*

ABSTRACT. We prove that various GIT semistabilities of polarized varieties imply semi-log-canonicity.

## 1. INTRODUCTION

For the study of the moduli of polarized varieties, Geometric Invariant Theory [Mum65] (GIT, for short) is an important basis, because it constructs the moduli spaces as quotient schemes of the Hilbert schemes. In that theory, we must put restrictions on the objects to classify, which we call *stability*, the GIT stability. It is a quite difficult and interesting problem to explicitly understand the stability notion.

Let us recall that the compact moduli scheme of curves  $\bar{M}_g$  is constructed in GIT by permitting ordinary double points (nodes) to curves ([DM69], [KM76], [Mum77], [Gie82]), which is sometimes called the Deligne-Mumford compactification. We note that semistable polarized curves have only nodal singularities.

In this paper, we give its higher dimensional generalization and show that the general effect of singularities on stability is determined by the *discrepancy*, an invariant of singularity which was developed along the minimal model program. This is our new point of view. Recall that the discrepancy is defined under the following conditions, which ensure that the canonical divisor  $K_X$  or the canonical sheaf  $\omega_X$  is in a tractable class (cf. e.g., [Ale96]).

**Definition 1.1.** An algebraic scheme  $X$  is said to *satisfy*  $(*)$  when the following conditions hold.

- (i)  $X$  is equidimensional and reduced.
- (ii)  $X$  satisfies Serre's  $S_2$  condition.
- (iii) Codimension 1 points of  $X$  are Gorenstein.

---

2010 *Mathematics Subject Classification.* Primary 14L24; Secondary 14J17, 32Q15.

*Key words and phrases.* GIT-stability, Donaldson-Futaki invariants, semi-log-canonicity.

- (iv)  $K_X$  is  $\mathbb{Q}$ -Cartier, in the sense that  $\mathcal{O}_X(nK_X) := (\omega_X^{\otimes n})^{\vee\vee}$  is an invertible sheaf for some  $n \in \mathbb{Z}_{>0}$ , where  $\mathcal{F}^\vee := \text{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{O}_X)$ .

Then, our Main result is the following.

**Theorem 1.2.** *Let  $X$  be a projective scheme satisfying  $(*)$  and  $L$  be an ample line bundle on  $X$ . Then, if  $(X, L)$  is  $K$ -semistable,  $X$  has only semi-log-canonical singularities.*

We will also explain that Theorem 1.2 above should be the best possible as we will see in the statements of Theorem 1.5 below.

As already mentioned, the definition of *semi-log-canonicity* is based on the *discrepancy*. The theory of discrepancy originally stemmed out of a necessity in the way of extending minimal models for surfaces by the Italian school to higher dimensions after Mori [Mor82], as they should be allowed to have some mild singularities. Indeed, it forms a core notion in the minimal model program (the MMP, for short).

Along the development of the MMP, the semi-log-canonicity was first introduced by Kollár and Shepherd-Barron [KSB88] for surfaces and extended by Alexeev [Ale96] to higher dimensions. Their original purpose was to construct the compactified moduli spaces for varieties of general type *not* by GIT theory, but by MMP techniques. For the case of curves, semi-log-canonical singularities are simply smooth points or nodes. Semi-log-canonical surface singularities are classified by Kollár-Shepherd-Barron [KSB88, Theorem (4.24)].

Now, let us explain the other side i.e., the stability notion. While the GIT stability was originally intended to construct moduli spaces as mentioned at the beginning, the  $K$ -(*semi*)*stability* is a version of GIT-stability notion which was firstly introduced by Tian [Tia97] to describe when a Fano manifold has a Kähler-Einstein metric. Subsequently, Donaldson [Don02] extended the notion to general polarized varieties with an expectation of correspondence with the existence of Kähler metrics whose scalar curvature are constant (cscK, for short). We follow Donaldson's formulation [Don02] in this paper. We note that it is defined algebro-geometrically, although the introduction is motivated by differential geometry.

Thus roughly speaking, our Main theorem 1.2 bridges in a fresh way, these two theories in algebraic geometry, i.e., birational geometry and GIT stability (in a broader sense). In addition, due to the conjectural correspondence with metrics side, one could hope that stability or moduli problems have further connections with differential geometry.

We should make some comments on Theorem 1.2.

Firstly, we remark on the Fano case. In this paper,  $X$  is said to be  $(*)$ -Fano scheme if  $X$  is a projective scheme satisfying  $(*)$  and  $-K_X$  is ample (we do *not* a priori assume normality of  $X$ ). In this case, we can prove the following stronger result by slightly different arguments.

**Theorem 1.3.** *If  $X$  is a  $(*)$ -Fano scheme as above and  $(X, \mathcal{O}_X(-mK_X))$  is  $K$ -semistable with  $m \in \mathbb{Z}_{>0}$ , then  $X$  is log terminal. (In particular,  $X$  should be normal).*

Secondly, let us comment on other stability notions. Recall that Mumford and Gieseker studied *asymptotic (Chow and Hilbert) stabilities*, which were the original stability notions for polarized varieties ([Gie82], [Mum77] etc). It is well known that these asymptotic (Chow or Hilbert) semistabilities imply  $K$ -semistability (cf. [RT07, section 3]). Furthermore, there are more stability notions introduced recently by Donaldson ([Don10]), called  $\bar{K}$ -stability and b-stability. It seems that these two notions are expected to be equivalent at least for smooth case, and we can see that  $\bar{K}$ -semistability is also stronger than  $K$ -semistability. Therefore, we have

**Corollary 1.4.** (i) *Let  $X$  be a projective scheme satisfying  $(*)$  and  $L$  be an ample line bundle on  $X$ . Then, if  $(X, L)$  is asymptotically (Chow or Hilbert) semistable,  $X$  has only semi-log-canonical singularities.*

(ii) *Let  $X$  be a projective scheme satisfying  $(*)$  and  $L$  be an ample line bundle on  $X$ . Then, if  $(X, L)$  is  $\bar{K}$ -semistable,  $X$  has only semi-log-canonical singularities.*

(iii) *If  $X$  is a  $(*)$ -Fano scheme and  $(X, \mathcal{O}_X(-mK_X))$  with  $m \in \mathbb{Z}_{>0}$  is asymptotically (Chow or Hilbert) semistable or  $\bar{K}$ -semistable, then  $X$  is log terminal. (In particular,  $X$  should be normal).*

Final but an important remark about Theorem 1.2 is that the following converse has already been proved for Calabi-Yau case ([Od09]) and canonically polarized case ([Od11]). In this sense, Theorem 1.2 is the best possible as mentioned earlier.

**Theorem 1.5.** (i) ([Od09]) *A semi-log-canonical polarized variety  $(X, L)$  with numerically trivial canonical divisor  $K_X$  is  $K$ -semistable.*

(ii) ([Od11]) *A semi-log-canonical (pluri)canonically polarized variety  $(X, \mathcal{O}_X(mK_X))$  with  $m \in \mathbb{Z}_{>0}$  is  $K$ -stable.*

We make a caution that, on the other hand, the singularities do not determine stabilities *in general*, as it is well known that there are smooth but not semistable polarized manifolds.

We also remark that it has been known for a few decades that the asymptotic (semi)stability version of Theorem 1.5 does *not* hold (cf.

[She83], [Od11]). The author supposes that this phenomenon ought to be a major reason why the relation between discrepancy and stability of polarized varieties has been unexpected so far.

The bare structure of the proof of Theorem 1.2 is that, assuming non-semi-log-canonicity of  $X$  (i.e.,  $X$  has “bad” singularities), we construct a “de-stabilizing” one-parameter subgroup by making use of a certain birational model of  $X$  and  $X \times \mathbb{A}^1$ . On the way of the proof, we define *S-coefficient*, which is an invariant of certain ideals of  $X \times \mathbb{A}^1$ . Very roughly speaking, to those ideals we associate the one parameter subgroups.

The birational model of  $X$  which we shall use is the (relative) *semi-log canonical model* whose existence has been conjectured in the theory of the log minimal model program (LMMP, for short), at least for normal case. The existence is recently verified in [OX11].

In our standpoint, Shah [Sha81] introduced our key invariant S-coefficient for isolated singularities by an argument based on Eisenbud-Mumford’s local stability theory [Mum77], and applied it to give certain list of semistable surface singularities, which gave us one of the major inspirations for Theorem 1.2.

Our paper is organized as follows.

In the next section, we will review the basic stability notions for polarized varieties and some preparatory materials related to the log minimal model program. In section 3, we will formulate an invariant of polarized varieties (with an ideal of certain type attached), which we call the *S-coefficient*, as a generalization of “ $a_I$ ” in [Sha81]. Actually, the S-coefficient can be regarded as the leading coefficient of some series of the Donaldson-Futaki invariants, which can be calculated by formula 3.2 proven in [Od09], [Wan08]. After that, we give technical details to the (birational geometric part of) proof of Theorem 1.2, 1.3 in the following sections.

**Conventions.** Throughout, we work over an algebraically closed field  $k$  with characteristic 0.

A polarization means an *ample* invertible sheaf and a polarized scheme means an algebraic scheme  $X$  equipped with an ample invertible sheaf  $L$ .  $(X, L)$  always denotes a polarized scheme, and except in subsection 2.1 and a tiny part of subsection 3.1, it is assumed to satisfy (\*) as in the statement of Theorem 1.2. (For example, an arbitrary reduced projective hypersurface, or more generally, a (global) reduced complete intersection satisfies the conditions. )

$NN(X)$ ,  $NLC(X)$ ,  $NSLC(X)$  and  $NKLT(X)$  denote non-normal locus, non-log-canonical locus, non-semi-log-canonical locus, and non-kawamata-log-terminal locus of  $X$ , respectively.  $X^\nu$  denotes the normalization of a given variety  $X$ .

$a(E; X)$  denotes the discrepancy of a divisor  $E$  over a normal variety  $X$  and  $a(E; X, D)$  denotes the discrepancy of  $E$  over a normal pair  $(X, D)$  (i.e., a pair of a normal variety  $X$  and its Weil divisor  $D$  with  $\mathbb{Q}$ -Cartier  $K_X + D$ ).

**Acknowledgements.** This paper is an expanded version of the author's master thesis, submitted in January of 2009, to RIMS, Kyoto university. It essentially proved Theorem 1.2 modulo the existence of the (relative) semi-log-canonical model.

The author can not express enough his sincere and deep gratitude to his advisor Professor Shigefumi Mori for his helpful comments, reading the draft and the revision and for his warm encouragements. The author also would like to thank Professors Shigeru Mukai, Noboru Nakayama, Masayuki Kawakita, Osamu Fujino, Julius Ross, Xiaowei Wang, Yuji Sano, Yongnam Lee, Shiro Goto, Hisanori Ohashi and Mr. Kento Fujita for helpful comments and encouragements for pursuing the program, writing and revising the paper. For the revision, the anonymous referee considerably helped the author to improve the exposition, which he also greatly appreciate.

My big special thanks go to Professors János Kollár and Chenyang Xu. It is a great pleasure of the author to mention that, thanks to them, the technical problem of constructing the relative semi-log-canonical model is settled in [OX11] on which our proof depends.

The author is partially supported by the Grant-in-Aid for Scientific Research (KAKENHI No. 21-3748) and the Grant-in-Aid for JSPS fellows.

## 2. PRELIMINARIES

In this section, we review the basics for  $K$ -stability, discrepancy, and log canonical model.

**2.1.  $K$ -stability.** The  $K$ -stability was introduced first, under differential geometric background, by Tian in [Tia97], and reformulated and extended later by Donaldson [Don02]. Donaldson's version of  $K$ -stability and  $K$ -polystability have been slightly amended recently by [LX11], while the semistability notion remained the same (see also [Od12], [Stp11]). Recall that it is the motivation for introducing the  $K$ -(semi, poly)stability to seek the GIT-counterpart of the existence of special Kähler metric. Indeed, according to Professors Gang Tian

and Toshiki Mabuchi, the “K” in K-stability stands for the K-energy (Mabuchi energy), a functional on the space of Kähler metrics whose critical points are canonical Kähler metrics and at last the “K” in the K-energy came from “K”ähler.

For the definition of the stability, we need the concept of “test configuration” following Donaldson [Don02]. Our notation (and even expression) mostly follows [RT07].

**Definition 2.1.** A *test configuration* (resp. *semi test configuration*) for a polarized complete scheme  $(X, L)$  is a quasi-projective scheme  $\mathcal{X}$  with an invertible sheaf  $\mathcal{M}$  on it with:

- (i) a  $\mathbb{G}_m$  action on  $(\mathcal{X}, \mathcal{M})$
- (ii) a proper flat morphism  $\alpha: \mathcal{X} \rightarrow \mathbb{A}^1$

such that  $\alpha$  is  $\mathbb{G}_m$ -equivariant for the usual action on  $\mathbb{A}^1$ :

$$\begin{array}{ccc} \mathbb{G}_m \times \mathbb{A}^1 & \longrightarrow & \mathbb{A}^1 \\ (t, x) & \longmapsto & tx, \end{array}$$

$\mathcal{M}$  is relatively ample (resp. relatively semi ample), and  $(\mathcal{X}, \mathcal{M})|_{\alpha^{-1}(\mathbb{A}^1 \setminus \{0\})}$  is  $\mathbb{G}_m$ -equivariantly isomorphic to  $(X, L^{\otimes r}) \times (\mathbb{A}^1 \setminus \{0\})$  for some positive integer  $r$ , called *exponent*, with the natural action of  $\mathbb{G}_m$  on the latter and the trivial action on the former.

**Proposition 2.2** ([RT07, Proposition 3.7]). *In the above situation, a one-parameter subgroup of  $GL(H^0(X, L^{\otimes r}))$  is equivalent to the data of a test configuration  $(\mathcal{X}, \mathcal{M})$  of  $(X, L)$  with the polarization  $\mathcal{M}$  very ample (over  $\mathbb{A}^1$ ) and of exponent  $r$  for  $r \gg 0$ .*

In fact, let  $\lambda: \mathbb{G}_m \rightarrow GL(H^0(X, L^{\otimes r}))$  be a one-parameter subgroup. Then, consider the natural action  $\lambda \times \rho$  of  $\mathbb{G}_m$  on  $(\mathbb{P}(H^0(X, L^{\otimes r})) \times \mathbb{A}^1, \mathcal{O}(1))$  as a polarized variety, where  $\rho$  is the multiplication action on  $\mathbb{A}^1$ . Then the closure of the orbit  $\mathcal{X} := \overline{((\lambda \times \rho)(\mathbb{G}_m))(X \times \{1\})}$  is a test configuration with the natural polarization  $\mathcal{O}(1)|_{\mathcal{X}}$  and the restriction of the natural action on  $(\mathbb{P}(H^0(X, L^{\otimes r})) \times \mathbb{A}^1, \mathcal{O}(1))$ . This is called the *DeConcini-Procesi family* of  $\lambda$  by Mabuchi. The fact that any (very ample) test configuration can be obtained in this way follows from the fact that an arbitrary  $\mathbb{G}_m$ -equivariant vector bundle over  $\mathbb{A}^1$  should be equivariantly trivial (cf. [Don05, Lemma 2]).

Therefore, the test configuration can be regarded as *geometrization* of one-parameter subgroup.

Now, let us define the Donaldson-Futaki invariants for test configurations whose positivity define K-stability. As a preparation, let us note that the *total weight* of an action of  $\mathbb{G}_m$  on some finite-dimensional vector space will mean the sum of all weights in this



paper. Here the *weights* mean the exponents of eigenvalues which should be powers of  $t$ . Take a test configuration  $(\mathcal{X}, \mathcal{M})$  and suppose that the exponent  $r$  is 1. Otherwise, we can similarly proceed by considering  $(X, L^{\otimes r})$  instead of  $(X, L)$ . We denote the total weight of the induced action on  $(\alpha_* \mathcal{M}^{\otimes U})|_0$  by  $w(U)$  and  $\dim X$  as  $n$ . It is a polynomial of  $U$  of degree  $n + 1$ . On the other hand, we write  $P(u) := \dim H^0(X, L^{\otimes u})$ . Let us take  $rP(r)$ -th power of the action of  $\mathbb{G}_m$  on  $\mathcal{M}|_0$  and multiply suitable power of  $t$  so that the action on the vector space  $(\alpha_* \mathcal{M}^{\otimes r})|_{\{0\}}$  would be in the special linear group  $\mathrm{SL}((\alpha_* \mathcal{M}^{\otimes r})|_{\{0\}})$ . Then, the corresponding normalized weight on  $(\alpha_* \mathcal{M}^{\otimes U})|_0$  is  $\tilde{w}_{r,U} := w(u)rP(r) - w(r)uP(u)$ , where  $u := Ur$ . It is a polynomial of form  $\sum_{i=0}^{n+1} e_i(r)u^i$  of degree  $n + 1$  in  $u$  for  $u \gg 0$ . Further, the coefficients  $e_i(r)$  are again polynomials of degree  $n + 1$  in  $r$  for  $r \gg 0$ :  $e_i(r) = \sum_{j=0}^{n+1} e_{i,j}r^j$  for  $r \gg 0$ . Since the weight is normalized,  $e_{n+1,n+1} = 0$ .  $e_{n+1,n}$  is called the *Donaldson-Futaki invariant* of the test configuration, which we will denote by  $\mathrm{DF}(\mathcal{X}, \mathcal{M})$ .

Note that  $(n + 1)!e_{n+1}(r)r^{n+1}$  has meaning as the Chow weight of  $X \subset \mathbb{P}(H^0(X, L^{\otimes r}))$  with respect to the SL-normalization of the one parameter subgroups associated to  $(\mathcal{X}, \mathcal{M}^{\otimes r})$  via Proposition 2.2 for  $r \gg 0$  (cf. [Mum77, Lemma 2.11]).

For an arbitrary *semi* test configuration  $(\mathcal{X}, \mathcal{M})$  we can define the (normalized) Chow weight or the Donaldson-Futaki invariant in completely similar way from the total weights of the induced  $\mathbb{G}_m$ -action on  $(\alpha_* \mathcal{M}^{\otimes U})|_{\{0\}}$  for  $U \gg 0$ . Also note that the homogeneity  $\mathrm{DF}(\mathcal{X}, \mathcal{M}^{\otimes c}) = c^{2n} \mathrm{DF}(\mathcal{X}, \mathcal{M})$  easily follows from the definition.

Now, we can recall the definition of K-stability as follows.

**Definition 2.3** (cf. [Stp11], [Od12]). A test configuration  $(\mathcal{X}, \mathcal{L})$  is said to be *almost trivial* if  $\mathcal{X}$  is  $\mathbb{G}_m$ -equivariantly isomorphic to the product test configuration away from a closed subset of codimension at least 2.

**Definition 2.4.** (i) A polarized complete scheme  $(X, L)$  is *K-stable* (resp. *K-semistable*) if for any test configurations of  $(X, L)$  which are not almost trivial, with exponent  $r$ , the leading coefficient  $e_{n+1,n}$  of  $e_{n+1}(r)$  (the Donaldson-Futaki invariant) is positive (resp. non-negative).

(ii) A polarized complete scheme  $(X, L)$  is *K-polystable* if it is K-semistable and the Donaldson-Futaki invariant of a test configuration  $(\mathcal{X}, \mathcal{M})$  is 0 if and only if  $\mathcal{X}$  is isomorphic to  $X \times \mathbb{A}^1$  away from a closed subset of codimension at least 2.



Although we only use  $K$ -semistability in this paper, we make remarks on other notions. We should note that the original “ $K$ -stability” of [Don02] is what is called “ $K$ -polystability” in [RT07]. We follow the convention of [RT07] at this point. We further note that it is possible to re-define asymptotic stability by the quantities introduced above, associated to test configurations, due to Proposition 2.2.

About other stability notions, we only note that stability notions are related as follows, without giving their definitions and proofs. For the details, we refer to [RT07] and [Don10].

**Claim 2.5.** (i) *Asymptotically Chow stable*  $\Rightarrow$  *Asymptotically Hilbert stable*  $\Rightarrow$  *Asymptotically Hilbert semistable*  $\Rightarrow$  *Asymptotically Chow semistable*  $\Rightarrow$   *$K$ -semistable*.  
(ii)  $\bar{K}$ -stable  $\Rightarrow$   $\bar{K}$ -semistable  $\Rightarrow$   $K$ -semistable.

Hence, among these notions,  $K$ -semistability is the weakest notion. It is the reason why Corollary 1.4 should follow from Theorem 1.2, 1.3.

**2.2. Singularities via discrepancy.** We will now explain the discrepancy and some classes of mild singularities. Consult [KM98, section 2.3] and [Koletc92, Chapter 12] for the details. Let us first treat normal case. Let  $(X, D)$  be a normal pair, i.e., a pair of a normal variety  $X$  and an effective  $\mathbb{Q}$ -divisor  $D$  such that  $K_X + D$  is  $\mathbb{Q}$ -Cartier. Let  $\pi : X' \rightarrow X$  be a log resolution of  $D$ , i.e.,  $\pi$  is a proper birational morphism such that  $X'$  is smooth and  $\pi^{-1} \text{Supp}(D) \cup E$  has a simple normal crossing divisor support, where  $E$  is the exceptional divisor of  $\pi$ . Then, we denote

$$K_{X'} - \pi^*(K_X + D) = \sum_i a(E_i; (X, D))E_i,$$

where  $a(E_i; (X, D)) \in \mathbb{Q}$  and  $E_i$  run over the set of divisors of  $X'$  supported on the exceptional locus or the support  $\text{Supp}(\pi_*^{-1}D)$  of  $\pi_*^{-1}D$ , the strict transform of  $D$ . We sometimes simply write  $a(E_i; (X, D))$  as  $a(E_i; X)$  if  $D = 0$ , and write  $a(E_i)$  if the pair in concern is obvious from the context.

The pair  $(X, D)$  is called *log canonical* (resp. *kawamata log terminal*) if and only if  $a(E_i; (X, D)) \geq -1$  (resp.  $a(E_i; (X, D)) > -1$ ) for any  $E_i$ . These notions are independent of the choice of the log resolution. We simply call  $X$  is *log canonical* (resp. *log terminal*) when  $(X, 0)$  is log canonical (resp. kawamata log terminal).

The semi-log-canonicity is an extension to non-normal case of the notion of log-canonicity. We introduce those notions without divisors,

i.e. in non-log setting, at this stage of argument. (We will need some log versions as well later in section 5, where we introduce those definitions.)

Let  $X$  be a projective variety, which is reduced, equidimensional,  $\mathbb{Q}$ -Gorenstein, Gorenstein in codimension 1 and satisfies Serre condition  $S_2$  (as we assumed). Let  $\nu: X^\nu \rightarrow X$  be the normalization morphism and attach a conductor divisor  $\text{cond}(\nu)$  on  $X^\nu$  which is defined by  $K_{X^\nu} = \nu^*K_X + \text{cond}(\nu)$ . From the assumption,  $(X^\nu, \text{cond}(\nu))$  is a log pair (i.e.  $K_{X^\nu} + \text{cond}(\nu)$  is  $\mathbb{Q}$ -Cartier). Then, the *semi-log-canonicity* of  $X$  are defined simply as the log canonicity of the normalized pair,  $(X^\nu, \text{cond}(\nu))$ .

For curve case, the semi-log-canonicity is equivalent to that the curve is nodal (or smooth). For surface case, that class of singularities is also classified by Kollár-Shepherd-Barron [KSB88]. For higher dimensional case, it is well known that a semi-log-canonical variety has only normal crossing singularity in codimension 1, so that repeatedly taking general hyperplane section leads to a nodal curve.

**2.3. Log canonical model.** To construct “de-stabilizing” test configurations for non-semi-log-canonical polarized varieties, we need a birational model called *(relative) log canonical model*. The definition is as follows.

**Definition 2.6.** Let  $(X, D)$  be a normal pair, i.e.  $X$  is a normal variety attached with a  $\mathbb{Q}$ -divisor such that  $K_X + D$  is  $\mathbb{Q}$ -Cartier. We call that a birational projective morphism  $\pi: B \rightarrow (X, D)$  gives a *(relative) log canonical model* of  $(X, D)$  (or of  $X$  if  $D = 0$ ) if with the divisor  $E_{\text{red}}$ , which denotes the sum of  $\pi$ -exceptional prime divisors with coefficients 1, the pair  $(B, E_{\text{red}})$  satisfies

- (1)  $(B, E_{\text{red}})$  is a log canonical pair,
- (2)  $K_B + E_{\text{red}}$  is ample over  $X$ .

The existence is established in [OX11]. We used the variable  $B$  as we shall use this regarding it as a certain blow up of  $X$ . Indeed, this model is a log canonical model of a log resolution with a boundary supported on the exceptional set in the sense of log minimal model program.

### 3. THE S-COEFFICIENTS

In this section, we introduce the concept of *S-coefficients* which control asymptotic behaviors for Donaldson-Futaki invariants of certain series of test configurations, and establish some basic properties.

### 3.1. Review of the formula for Donaldson-Futaki invariants.

In this subsection, let us recall the formula for Donaldson-Futaki invariants we shall use from [Od09]. Note that a slightly different version of the formula had been also proved independently by Xiaowei Wang in [Wan08].

Firstly we define a class of ideals, which we shall use for our study of stability. Let  $(X, L)$  be an  $n$ -dimensional polarized complete variety (which is *not* necessarily normal).

**Definition 3.1.** A coherent ideal sheaf  $\mathcal{J}$  of  $X \times \mathbb{A}^1$  is called a *flag ideal* if  $\mathcal{J} = I_0 + I_1 t + \dots + I_{N-1} t^{N-1} + (t^N)$ , where  $I_0 \subseteq I_1 \subseteq \dots \subseteq I_{N-1} \subseteq \mathcal{O}_X$  is the sequence of coherent ideal sheaves. (It is equivalent to that the corresponding subscheme is supported on the central fiber  $X \times \{0\}$  and is  $\mathbb{G}_m$ -invariant under the natural action of  $\mathbb{G}_m$  on  $X \times \mathbb{A}^1$ .)

Let us introduce some notation. We set  $\bar{\mathcal{L}} := p_1^* L$  on  $X \times \mathbb{P}^1$  and its restriction  $\mathcal{L} := p_1^* L|_{(X \times \mathbb{A}^1)}$ , where  $p_i$  is the  $i$ -th projection morphism from  $X \times \mathbb{A}^1$  or  $X \times \mathbb{P}^1$ . Let us write the blow up  $\bar{\mathcal{B}} := Bl_{\mathcal{J}}(X \times \mathbb{P}^1) \rightarrow X \times \mathbb{P}^1$  or its restriction to  $\mathcal{B} := Bl_{\mathcal{J}}(X \times \mathbb{A}^1) \rightarrow X \times \mathbb{A}^1$  by  $\Pi$ . Its natural exceptional divisor will be written as  $E$ , i.e.  $\mathcal{O}_{\mathcal{B}}(-E') = \Pi^{-1} \mathcal{J}$ . (We shall use the symbol (prime)  $'$  for denoting exceptional divisors to indicate it they are exceptional divisors of  $(n+1)$ -dimensional variety  $X \times \mathbb{A}^1$ , not of  $X$ .)

Let us assume  $r$  is sufficiently large so that  $(\Pi^* \mathcal{L}^{\otimes r})(-E')$  is (relatively) semi-ample (over  $\mathbb{A}^1$ ). Consider the Donaldson-Futaki invariant of the (semi) test configuration  $(\mathcal{B}, (\Pi^* \mathcal{L})^{\otimes r}(-E'))$ . Let us recall our formula for that.

**Theorem 3.2** ([Od09, Theorem 3.2]). *Let  $(X, L)$  and  $\mathcal{B}, \mathcal{J}$  be as above. And we assume that exponent  $r = 1$ . (It is just to make the formula easier. For general  $r$ , put  $L^{\otimes r}$  and  $\bar{\mathcal{L}}^{\otimes r}$  to the place of  $L$  and  $\bar{\mathcal{L}}$ .) Furthermore, we assume that  $X$  is equidimensional, reduced, satisfying  $S_2$  condition, whose codimension 1 points are Gorenstein and having  $\mathbb{Q}$ -Cartier canonical divisor  $K_X$  and  $\mathcal{B}$  is Gorenstein in codimension 1. Then the corresponding Donaldson-Futaki invariant  $DF((Bl_{\mathcal{J}}(X \times \mathbb{A}^1), \mathcal{L}(-E')))$  is*

$$\frac{1}{2(n!)(n+1)!} \left\{ -n(L^{n-1} \cdot K_X)((\Pi^* \bar{\mathcal{L}})(-E'))^{n+1} \right. \\ \left. + (n+1)(L^n)((\Pi^* \bar{\mathcal{L}})(-E'))^n \cdot \Pi^*(p_1^* K_X) \right. \\ \left. + (n+1)(L^n)((\Pi^* \bar{\mathcal{L}})(-E'))^n \cdot K_{\bar{\mathcal{B}}/X \times \mathbb{P}^1} \right\}.$$

*In the above, the intersection numbers  $(L^{n-1} \cdot K_X)$  and  $(L^n)$  are taken on  $X$ . On the other hand,  $K_{\bar{\mathcal{B}}/X \times \mathbb{P}^1} := K_{\bar{\mathcal{B}}} - \Pi^* K_{X \times \mathbb{P}^1}$  is*

an exceptional divisor on  $\bar{\mathcal{B}}$  and thus  $((\Pi^*\bar{\mathcal{L}})(-E'))^n \cdot \Pi^*(p_1^*K_X)$  and  $((\Pi^*\bar{\mathcal{L}})(-E'))^n \cdot K_{\bar{\mathcal{B}}/X \times \mathbb{P}^1}$  are intersection numbers taken on  $\bar{\mathcal{B}}$ .

We call the sum of first two terms the *canonical divisor part* since they involve intersection numbers with  $K_X$  or its pullback, and the last term will be called the *discrepancy term* since it reflects discrepancies over  $X$ . We remark that although not all semi test configurations are of the form  $(\mathcal{B}, (\Pi^*\mathcal{L})^{\otimes r}(-E'))$ , it is sufficient for K-(semi)stability to check the Donaldson-Futaki invariants of the special semi test configurations ([Od09]).

**3.2. S-coefficient as a leading coefficient of Donaldson-Futaki invariants.** We define the S-coefficient, the key invariant as follows.

**Definition 3.3.** Let us fix  $(X, L)$  in Theorem 3.2 above and fix a flag ideal  $\mathcal{J}$ . Suppose that  $\bar{\mathcal{B}}$  is Gorenstein in codimension 1 so that the canonical divisor class  $K_{\bar{\mathcal{B}}}$  is well defined. Then, the *S-coefficient* for that flag ideal  $\mathcal{J}$  is defined as an intersection number  $(\mathcal{L}^s \cdot (-E')^{n-s} \cdot K_{\bar{\mathcal{B}}/(X \times \mathbb{P}^1)})$  taken on  $\bar{\mathcal{B}}$  and we denote it by  $S_{(X,L)}(\mathcal{J})$ , where  $s$  denotes the dimension of  $\text{Supp}(\mathcal{O}_{X \times \mathbb{A}^1}/\mathcal{J})$ . We note that homogeneity  $S_{(X,L^{\lambda_1})}(\mathcal{J}^{\lambda_2}) = \lambda_1^s \lambda_2^{n-s} S_{(X,L)}(\mathcal{J})$  follows from the definition.

The main motivation for above definition is the following meaning of S-coefficient, as leading coefficient of Donaldson-Futaki invariants.

**Proposition 3.4.** *Let  $(X, L)$  and  $\mathcal{J}$  be as above. Then, the following holds.*

- (i) *The sequence of Donaldson-Futaki invariants  $\text{DF}(Bl_{\mathcal{J}}(X \times \mathbb{A}^1), \mathcal{L}^{\otimes r}(-E'))$  for  $r \gg 0$ , forms a polynomial.*
- (ii) *Its coefficient of  $r^d$  is 0 for  $d > n + s$  and equals to*

$$\frac{\binom{n}{s}(L^n)}{2(n!)^2} S_{(X,L)}(\mathcal{J})$$

for  $d = n + s$ .

Hence, if  $S_{(X,L)}(\mathcal{J}) < 0$  for some flag ideal  $\mathcal{J}$ , then  $(X, L)$  is not K-semistable.

To prove Proposition 3.4 above and analyze the positivity of the S-coefficients later, we shall use the following general properties of intersection numbers. As it follows from a standard arguments, we omit the proof. However, we give statements here for the readers' convenience as it shall be a key for our estimation.

**Lemma 3.5.** *Let  $\mathcal{X}$  be an arbitrary  $n + 1$ -dimensional equidimensional complete scheme, and  $\pi: \bar{\mathcal{B}} \rightarrow \mathcal{X}$  a surjective, generically finite morphism. Then*

(i)

$$(\pi^* D_1 \cdots \pi^* D_s \cdot E'_1 \cdots E'_{n+1-s}) = 0$$

for arbitrary Cartier divisors  $D_1, \dots, D_s$  on  $\mathcal{X}$ , and arbitrary Cartier divisors  $E'_1, \dots, E'_{n+1-s}$  with  $\dim(\pi(\cap \text{Supp}(E'_i))) < s$ .

(ii)

$$(\pi^* D_1 \cdots \pi^* D_s \cdot E'_1 \cdots E'_{n+1-s}) > 0$$

for arbitrary ample Cartier divisors  $D_1, \dots, D_s$  on  $\mathcal{X}$ , arbitrary ample Cartier divisors  $E'_1, \dots, E'_{n-s}$  on  $\bar{\mathcal{B}}$  and an arbitrary effective Weil divisor  $E'_{n+1-s}$  on  $\bar{\mathcal{B}}$  with  $\dim(\pi(E'_{n+1-s})) = s$ .

*Proof of Proposition 3.4.* Replacing  $L$  by  $L^{\otimes r}$  and  $\mathcal{L}$  by  $\bar{\mathcal{L}}^{\otimes r}$  for the formula 3.2, we have the formula of  $\text{DF}(Bl_{\mathcal{J}}(X \times \mathbb{A}^1), \bar{\mathcal{L}}^{\otimes r}(-E'))$ . From that, Proposition 3.4 (i) easily follows.

Further, Lemma 3.5 (i) applied to  $\pi = \Pi: Bl_{\mathcal{J}}(X \times \mathbb{P}^1) \rightarrow X \times \mathbb{P}^1$  by taking  $D_i := H \times \mathbb{P}^1$  where  $H \in |L^{\otimes m}|$  ( $m \in \mathbb{Z}_{>0}$ ),  $E'_i = E'$  for  $i \leq n - s$  and  $E'_{n+1-s} = K_{\bar{\mathcal{B}}/(X \times \mathbb{P}^1)}$ , implies Proposition 3.4 (ii) straightforward.  $\square$

**3.3. S-coefficients and discrepancy.** In this subsection, we shall show a criterion on positivity of S-coefficients, which gives a relation with discrepancy.

Let us assume, from now on, that  $X$  is an equidimensional reduced projective variety, satisfies  $S_2$  condition and whose codimension 1 points are Gorenstein. Thus we can define the Weil divisor class  $K_X$  which we assume to be  $\mathbb{Q}$ -Cartier. If all codimension 1 points of  $\mathcal{B}$  are Gorenstein, we set  $K_{\bar{\mathcal{B}}/(X \times \mathbb{P}^1)} := K_{\bar{\mathcal{B}}} - \Pi^*(K_X \times \mathbb{P}^1) = \sum a(E'_i)E'_i$ .

**Proposition 3.6.** *Let  $X$  be as above and  $L$  be an ample line bundle on  $X$ . Moreover, assume that there is a flag ideal  $\mathcal{J}$  whose blow up  $\mathcal{B}$  is Gorenstein in codimension 1 as noted above. Furthermore, assume that the discrepancies  $a(E'_i)$  satisfy the following.  $a(E'_i) \leq 0$  for all the exceptional prime divisors  $E'_i$  on  $\mathcal{B}$  which dominate  $s$  (maximal)-dimensional components of  $\text{Supp}(\mathcal{O}/\mathcal{J})$  and moreover there exists at least one such  $i$  with  $a(E'_i) < 0$ . Then, we have  $S_{(X,L)}(\mathcal{J}) < 0$ .*

Hence, by combining with Proposition 3.4, we have the following criterion for when a polarized variety can be not K-semistable.

**Corollary 3.7.** *Let  $X$  be as above and assume that there is a flag ideal  $\mathcal{J}$  whose blow up  $\mathcal{B}$  is Gorenstein in codimension 1 and the discrepancies  $a(E_i)$  satisfy the following.  $a(E'_i) \leq 0$  for all the exceptional prime divisors  $E'_i$  on  $\mathcal{B}$  which dominate  $s$  (maximal)-dimensional components of  $\text{Supp}(\mathcal{O}/\mathcal{J})$  and moreover there exists at least one such  $i$  with  $a(E'_i) < 0$ . Then  $(X, L)$  is not K-semistable for an arbitrary polarization  $L$ .*

*Proof of Proposition 3.6.* We have

$$(1) \quad \begin{aligned} S_{(X,L)}(\mathcal{J}) &:= (\mathcal{L}^s \cdot (-E')^{n-s} \cdot K_{\bar{\mathcal{B}}/(X \times \mathbb{P}^1)}) \\ &= (\mathcal{L}^s \cdot (\bar{\mathcal{L}}^{\otimes r} - E')^{n-s} \cdot K_{\bar{\mathcal{B}}/(X \times \mathbb{P}^1)}). \end{aligned}$$

Indeed, the equality (1) follows from Lemma 3.5 (i) applied to  $\pi = \Pi: Bl_{\mathcal{J}}(X \times \mathbb{P}^1) \rightarrow X \times \mathbb{P}^1$  by taking  $D_i := H \times \mathbb{P}^1$  for  $i \leq s+1$  where  $H \in |L^{\otimes m}|$  ( $m \in \mathbb{Z}_{>0}$ ) and for  $i > s+1$ ,  $E'_i = \Pi^*(H \times \mathbb{P}^1)$  or  $E'_i = E'$  or  $E'_i = K_{\bar{\mathcal{B}}/(X \times \mathbb{P}^1)}$ .

Moreover, the last term  $(\mathcal{L}^s \cdot (\bar{\mathcal{L}}^{\otimes r} - E')^{n-s} \cdot K_{\bar{\mathcal{B}}/(X \times \mathbb{P}^1)})$  is positive due to Lemma 3.5 (ii) applied to  $\Pi: Bl_{\mathcal{J}}(X \times \mathbb{P}^1) \rightarrow X \times \mathbb{P}^1$  again by taking, this time,  $D_i := H \times \mathbb{P}^1$  for  $i \leq s$ ,  $E'_i$  ( $i \leq n-s$ ) to be an ample compactification of an ample divisor which belongs to  $|(\Pi^*\mathcal{L})^{\otimes r}(-E')|$  on  $\mathcal{B}$  to  $\bar{\mathcal{B}}$  with  $r > 1$ , and  $E'_{n+1-s} := K_{\bar{\mathcal{B}}/(X \times \mathbb{P}^1)}$ .  $\square$

#### 4. NORMAL CASE

As an application of the theory of S-coefficients prepared in the previous section, we partially prove Theorem 1.2 for normal case in this section. More precisely, let  $X$  be a normal variety of pure dimension  $n$ , having  $\mathbb{Q}$ -Cartier canonical divisor in this section.

*Proof of Theorem 1.2 for normal  $X$ .* Thanks to Corollary 3.7, it is sufficient to construct a flag ideal  $\mathcal{J}$  satisfying the following property.

**Property 4.1.** *The blow up  $\mathcal{B}$  of  $\mathcal{J}$  is normal. Furthermore, if we let  $K_{\mathcal{B}/X \times \mathbb{A}^1} = \sum a(E'_i)E'_i$ , then we have  $a(E'_i) < 0$  for the discrepancy for an arbitrary  $\Pi$ -exceptional divisor  $E'_i$ .*

We will construct such  $\mathcal{J}$  in the following 2 steps. Without loss of generality, we can assume that  $X$  is irreducible.

**Step 1.** Firstly, we construct a coherent ideal sheaf  $I$  of  $X$ , satisfying the following property. We denote the blow up of  $X$  along  $I$  by  $\pi: B = Bl_I(X) \rightarrow X$ .

**Property 4.2.** *The blow up  $B$  is normal. Furthermore, if let  $s$  be  $\dim(\text{Supp}(\mathcal{O}_X/I))$ , then, we have  $a(E_i; X) < -1$  for the discrepancy for an arbitrary  $\pi$ -exceptional divisor  $E_i$ .*

We construct such  $I$ , using the (relative) log canonical model (cf. subsection 2.3) as follows. Suppose  $\pi: B \rightarrow X$  is the (relative) log canonical model of  $X$ , which exists due to [OX11, Theorem 1.1]. Then, we take the coherent ideal sheaf  $I := (\pi)_*\mathcal{O}_B(m(K_{B/X} + E_{\text{red}}))$  for sufficiently divisible  $m \in \mathbb{Z}_{>0}$  and the total exceptional divisor  $E_{\text{red}}$ , then  $Bl_I(X) \simeq B$ . Therefore, this  $I$  satisfies Property 4.2.

**Step 2.** Next step starts with taking  $I$  constructed in the previous step. Using this, we will construct the flag ideal  $\mathcal{J}$  satisfying Property 4.1 as follows. From the construction, we have  $\dim(\text{Supp}(\mathcal{O}_X/I)) \leq \dim(X) - 2$ . Let us take sufficiently divisible positive integers  $m, N$  and let us define  $\mathcal{J} := \overline{(I + (t^m))}^N$  where the overline denotes the integral closure of the coherent ideal. Since it is an invariant ideal with respect to the natural  $\mathbb{G}_m$  action on  $X \times \mathbb{A}^1$ ,  $\mathcal{J}$  is a flag ideal as well. We note that  $\mathcal{C} := Bl_{I+(t)}(X \times \mathbb{A}^1)$  is *the deformation to the normal cone* (cf. [Ful84], [RT07]) but simply taking it is *not* sufficient for our purpose in general. Geometrically speaking, to take  $I + (t^m)$  as above, instead of the simplest  $I + (t)$ , corresponds to take base change of  $\mathcal{C}$  by  $m$ -th roots of  $t$  (i.e.  $s \mapsto t := s^m$ ) and  $\mathcal{B} := Bl_{\mathcal{J}}(X \times \mathbb{A}^1)$  is the normalization of the base change (cf. [Vas05]).

Let us think of the more detailed geometric structure of the deformation to the normal cone  $\mathcal{C}$  and its modification  $\mathcal{B}$ . We know that its central fiber consists of two parts: the strict transform of  $X \times \{0\}$  canonically isomorphic to  $B = Bl_I(X)$  (we will identify them from now on), and the exceptional divisors  $F'_i$  which intersect as  $F'_i \cap B = E_i$  whose generic points  $\eta_i$  are regular. Indeed, étale locally we can write  $t = xy^{c_i}$  with étale local coordinates (i.e., regular parameters)  $x, y$  such that  $(x = 0)$  corresponds to  $B$  and  $(y = 0)$  corresponds to  $F'_i$ .

Based on the above facts, we obtain an étale local description of  $\mathcal{B} \rightarrow \mathcal{C}$  explicitly around the generic point  $\eta_i$  of  $F'_i \cap B$  as follows. We can take an étale local coordinate system  $(u, y, z_1, \dots, z_{n-2}, s)$  of  $\mathcal{B}$  around  $\eta_i$ , and that of  $\mathcal{C}$ :  $(x, y, z_1, \dots, z_{n-2}, t)$  around the fiber of  $\eta_i$  which are connected by the following equations. Here,  $t$  denotes the original coordinate of  $\mathcal{C}$  corresponding to the  $\mathbb{A}^1$  direction.

$$x = u^{c_i}, t = s^m.$$

We denote the preimage of  $F'_i$  by  $E'_i$ , which is irreducible. Then, from the above local description, it directly follows that:

$$(2) \quad a(E'_i; X \times \mathbb{A}^1) = b_i(a(E_i; X) + 1),$$

where each  $b_i := \frac{m}{c_i}$  is a positive integer as  $m$  is sufficiently divisible. Therefore,  $a(E'_i; X \times \mathbb{A}^1) < 0$  follows from Property 4.2 in the previous step of construction. This completed the proof of Theorem 1.2 for normal varieties' case.

□

## 5. NON-NORMAL CASE

To give a proof of the main theorem 1.2 in full generality, we introduce a non-normal generalization of the (relative) log-canonical



model, which we used in the previous section for normal case. A reduced equidimensional variety  $X$  is called *demi-normal* if  $X$  is  $S_2$ , whose codimension 1 points are regular or ordinary nodes.

**Definition 5.1.** Let  $X$  be a demi-normal projective variety. We call a birational projective morphism  $\pi: B \rightarrow X$  a *(relative) semi-log-canonical model* if  $\pi$  is isomorphic over open locus of  $X$  with complement's codimension greater than 1, and satisfies the following two conditions. Here,  $E_{\text{red}}$  denotes the sum of  $\pi$ -exceptional prime divisors with coefficients 1.

- (1)  $(B, E_{\text{red}})$  is a semi-log-canonical pair.
- (2)  $K_B + E_{\text{red}}$  is ample over  $X$ .

The existence of such models for any  $X$  is again proven in [OX11, Corollary 1.3]. Given this birational model, the proof of Theorem 1.2 below is similar to the case where  $X$  is normal.

*Proof of Theorem 1.2.* Take the (relative) semi-log-canonical model  $\pi: B \rightarrow X$  of  $X$ , which exists due to [OX11, Corollary 1.3]. Here, we note that all the generic points of  $\pi$ -exceptional divisors are regular, by the definition of the model. Then, if we apply the negativity lemma [KM98, Lemma (3.39)] to these normalizations, we have  $a_i < -1$  for any  $i$ , where  $K_{B/X} = \sum a_i E_i$ . Therefore, if we take  $I := \pi_*(\omega_{B/X}^{[l]}(lE))^{**}$  with sufficiently divisible positive integer  $l$ , where  $E_{\text{red}} := \sum E_i$  denotes the total exceptional divisor of  $\pi$ , it would be a coherent ideal sheaf by Serre's  $S_2$ -condition of  $X$ . Further, it satisfies  $Bl_I(X) \cong B$  by the relative ampleness of  $K_{B/X} + E_{\text{red}}$ .

Let us consider a flag ideal  $\mathcal{J}' = I + (t^m)$  on  $X \times \mathbb{A}^1$  for sufficiently divisible positive integer  $m$ , its blow up  $\mathcal{C} = Bl_{\mathcal{J}'}(X \times \mathbb{A}^1)$  and its normalization  $\mathcal{C}^\nu \rightarrow \mathcal{C}$ . We denote  $\Pi: \mathcal{C}^\nu \rightarrow X^\nu \times \mathbb{A}^1$  the associated morphism. We can prove  $K_{\mathcal{C}^\nu} - \Pi^*(K_{X^\nu \times \mathbb{A}^1} + \text{cond}(\nu) \times \mathbb{A}^1) = \sum a'_i H'_i$  with  $a'_i = b_i(a_i + 1) < 0$  where  $b_i$  are some positive integers for each exceptional divisor  $H'_i$  and  $\text{cond}(\nu)$  is the conductor divisor of the normalization. The proof is in a completely similar manner as in the previous section, by taking  $(X^\nu, \text{cond}(\nu))$  instead of  $X$  with the normality assumption.

We use the *partial normalization*  $\mathcal{B}$  of  $\mathcal{C}$  which was introduced in the proof of [Od09, Proposition 3.8]. The definition is  $\mathcal{B} := \text{Spec}_{\mathcal{O}_{\mathcal{C}}}(i_* \mathcal{O}_{X \times (\mathbb{A}^1 \setminus \{0\})} \cap \mathcal{O}_{\mathcal{C}^\nu})$ , where  $i: X \times (\mathbb{A}^1 \setminus \{0\}) \hookrightarrow X \times \mathbb{A}^1$  is the open immersion. Let  $f$  be the associated morphism from  $\mathcal{C}^\nu$  to  $\mathcal{B}$ . Completely similarly as we argued in the former half of Step 2 of the previous section, we can take a flag ideal  $\mathcal{J}$  whose blow up is  $\Pi: \mathcal{B} = Bl_{\mathcal{J}}(X \times \mathbb{A}^1) \rightarrow X \times \mathbb{A}^1$ . Let us recall the following lemma.

**Lemma 5.2** ([Od09, Lemma 3.9]). *The morphism  $f: \mathcal{C}^\nu \rightarrow \mathcal{B}$  is an isomorphism over an open neighborhood of the generic points of the central fiber.*

Thus, similarly as in the comparison of discrepancies (2), we have  $K_{\mathcal{B}/X \times \mathbb{A}^1} = \sum a'_i E'_i$  with  $a'_i = b_i(a_i + 1) < 0$  where  $b_i$  are the positive integers introduced above, and  $E'_i$  is the strict transform of  $H'_i$ .

Therefore, we complete the proof Theorem 1.2 thanks to Corollary 3.7. □

## 6. FANO CASE

In this section, we prove Theorem 1.3.

*proof of Theorem 1.3.* In this section we do not use the notion of S-coefficients but the proof is done by analyzing the formula for the Donaldson-Futaki invariants 3.2 more directly.

Let us take a flag ideal  $\mathcal{J} := \overline{(I + (t))}^N$ , where  $I \subset \mathcal{O}_X$  corresponds to the reduced subscheme supported on  $\text{NN}(X)$ , the non-normal locus of  $X$ , and  $N$  is a sufficiently divisible positive integer. Here, we do not take parameter  $m$  into account. We note that  $\text{NN}(X)$  is purely codimension 1 in  $X$  and their generic points are ordinary double points. In the case of curves (i.e.,  $\dim(X) = 1$ ), this means  $X$  should be nodal. Consider the (semi) test configuration of the blow up type  $(\mathcal{B}, (\Pi^* \mathcal{L})^{\otimes r}(-E'))$  for that flag ideal  $\mathcal{J}$  as we did. Then, the S-coefficients become 0 and the leading coefficients of  $\text{DF}(\mathcal{B}, (\Pi^* \mathcal{L})^{\otimes r}(-E'))$  with respect to the variable  $r$  has the same signature as  $((\Pi^* \mathcal{L})^{n-1} \cdot E'^2)$ . This can be shown to be negative by cutting  $X$  for  $s := \dim(\text{Supp}(\mathcal{O}_{X \times \mathbb{A}^1} / \mathcal{J})) = n - 1$  times by general hypersurface sections in  $|L^{\otimes m}|$  for  $m \gg 0$ , reducing to the  $n = 1$  case.

Thus, we can assume that  $X$  is normal. Let us assume that  $X$  is log canonical but not log terminal (i.e., strictly log canonical) and derives a contradiction. In the sense of log minimal model program, a log resolution with kawamata-log-terminal boundary  $(\tilde{X}, (1 - \epsilon)E_{\text{red}})$  with  $0 < \epsilon \ll 1$  should have a log canonical model  $B$  over  $X$ , by [BCHM09, Theorem 1.2]. Note that  $B$  should be log terminal and so the morphism  $B \rightarrow X$  is not isomorphism, which is again a blow up of certain coherent ideal sheaf  $I$ . We further remark that the model of subsection 2.3 corresponds to the  $\epsilon = 0$  case. Similarly as in section 4, we construct a flag ideal  $\mathcal{J} := \overline{(I + (t^m))}^N$  where  $m, N$  are sufficiently divisible positive integers, and its blow up  $\mathcal{B} := \text{Bl}_{\mathcal{J}}(X \times \mathbb{A}^1)$ . Then,  $K_{\mathcal{B}/X \times \mathbb{A}^1} = 0$  so that the discrepancy term vanishes.

On the other hand, as  $s < n$ , the canonical divisor part of the formula 3.2 is negative by [Od11, proof of Theorem 2.13]. Hence,  $(X, L)$  should be not K-semistable. This completes the proof of Theorem 1.3.  $\square$

## REFERENCES

- [Ale96] V. Alexeev, Log canonical singularities and complete moduli of stable pairs, arXiv:alge-geom/960813 (1996).
- [BCHM09] C. Birkar, P. Cascini, C. Hacon, J. Mckernan, Existence of minimal models for varieties of log general type, *J. A. M. S.* vol. **23** (2009), 405-468.
- [Don02] S. K. Donaldson, Scalar curvature and stability of toric varieties, *J. Differential Geom.* vol. **62** (2002), 453-472.
- [Don05] S. K. Donaldson, Lower bounds on the Calabi functional, *J. Differential Geom.* vol. **70** (2005).
- [Don10] S. K. Donaldson, Stability, birational transformations and the Kähler-Einstein problem, arXiv:1007.4220 (2010).
- [DM69] P. Deligne, D. Mumford, The irreducibility of the space of curves of a given genus, *Publ. Math. I. H. E. S.* vol. **36** (1969).
- [Ful84] W. Fulton, *Intersection Theory*, Springer Verlag, 1984.
- [Gie82] D. Gieseker, *Lectures on moduli of curves*, Tata Institute of Fundamental Research Lectures on Mathematics and Physics. Springer-Verlag, (1982).
- [Har77] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics vol. **52**, Springer-Verlag.
- [KM76] F. F. Knudsen, D. Mumford, The projectivity of the moduli spaces of stable curves. I, *Math. Scand.* vol. **39** (1976), 19-50.
- [KM98] J. Kollár, S. Mori, *Birational Geometry of Algebraic Varieties*, Cambridge Tracts in Mathematics, vol. **134**, Cambridge University Press (1998).
- [KSB88] J. Kollár, N. I. Shepherd-Barron, Threefolds and deformations of surface singularities, *Invent. Math.* vol. **91** (1988), 299-338.
- [Koletc92] J. Kollár editor, *Flips and abundance for algebraic threefolds*, *Astérisque* **211** (1992).
- [Kol] J. Kollár, *Book on Moduli of Surfaces*, to appear. Chapter 1, 2, 3 is available in webpage: (URL) <http://www.math.princeton.edu/~kollar/>.
- [LX11] C. Li, C. Xu, Special test configurations and K-stability of  $\mathbb{Q}$ -Fano varieties, arXiv:1111.5398 (2011)
- [Mor82] S. Mori, Threefolds whose canonical divisors are not numerically effective, *Ann. of Math.* vol. **116** (1982).
- [Mum65] D. Mumford, *Geometric Invariant Theory*, *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, (1965).
- [Mum77] D. Mumford, Stability of Projective Varieties, *Enseignement Math.* vol. **23** (1977).
- [Od09] Y. Odaka, A generalization of the Ross-Thomas slope theory, to appear in *Osaka J. Math.*
- [Od11] Y. Odaka, The Calabi conjecture and K-stability, *International Mathematical Research Notices* no. **13** (2011).
- [Od12] Y. Odaka, On parametrization, optimization and triviality of test configurations, arXiv:1201.0692. (2012)

- [OX11] Y. Odaka, C. Xu, Log-canonical models of singular pairs and its applications, to appear in *Math. Res. Lett.*
- [RT07] J. Ross, R. Thomas, A study of the Hilbert Mumford criterion for the stability of projective varieties, *J. Algebraic Geom.* vol. **16** (2007), 201-255.
- [Sha81] J. Shah, Stability of two-dimensional local rings. I, *Invent. Math.* vol. **64** (1981), 297-343.
- [She83] N. I. Shepherd-Barron, Degenerations with numerically effective canonical divisor, *The birational geometry of degenerations* (Cambridge, Mass. , 1981), *Progr. Math.* , vol. **29**, Birkhäuser Boston, Boston, MA, (1983) 33-84.
- [Stp11] J. Stoppa, A note on the definition of K-stability, arXiv:1111.5826 (2011)
- [Tia97] G. Tian, Kähler-Einstein metrics with positive scalar curvatures, *Invent. Math.* vol. **130** (1997), 1-37.
- [Vas05] W. Vasconcelos, *Integral Closure: Rees Algebras, Multiplicities, Algorithms*, Springer Monographs in Mathematics. Springer-Verlag, (2005).
- [Wan08] X. Wang, Heights and GIT weights, preprint written in 2008, available in his webpage. (URL) <http://www.math.cuhk.edu.hk/~xiaowei/>

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES (RIMS), KYOTO UNIVERSITY, OIWAKE-CHO, KITASHIRAKAWA, SAKYO-KU, KYOTO 606-8502, JAPAN

*Current address:* IMPERIAL COLLEGE LONDON, SOUTH KENSINGTON CAMPUS, LONDON SW7 2AZ, ENGLAND

*E-mail address:* [y.odaka@imperial.ac.uk](mailto:y.odaka@imperial.ac.uk)