

# THE GREATEST PRIME DIVISOR OF A PRODUCT OF CONSECUTIVE INTEGERS

SHANTA LAISHRAM AND T. N. SHOREY

## 1. INTRODUCTION

Let  $k \geq 2$  and  $n \geq 1$  be integers. We denote by

$$\Delta(n, k) = n(n+1) \cdots (n+k-1).$$

For an integer  $\nu > 1$ , we denote by  $\omega(\nu)$  and  $P(\nu)$  the number of distinct prime divisors of  $\nu$  and the greatest prime factor of  $\nu$ , respectively, and we put  $\omega(1) = 0$ ,  $P(1) = 1$ .

A well known theorem of Sylvester [7] states that

$$(1) \quad P(\Delta(n, k)) > k \text{ if } n > k.$$

We observe that  $P(\Delta(1, k)) \leq k$  and therefore, the assumption  $n > k$  in (1) cannot be removed. For  $n > k$ , Moser [5] sharpened (1) to  $P(\Delta(n, k)) > \frac{11}{10}k$  and Hanson [3] to  $P(\Delta(n, k)) > 1.5k$  unless  $(n, k) = (3, 2), (8, 2), (6, 5)$ . Further Faulkner [2] proved that  $P(\Delta(n, k)) > 2k$  if  $n$  is greater than or equal to the least prime exceeding  $2k$  and  $(n, k) \neq (8, 2), (8, 3)$ . In this paper, we sharpen the results of Hanson and Faulkner. We shall not use these results in the proofs of our improvements. We prove

**Theorem 1.** *We have*

(a)

$$(2) \quad P(\Delta(n, k)) > 2k \text{ for } n > \max(k + 13, \frac{279}{262}k).$$

(b)

$$(3) \quad P(\Delta(n, k)) > 1.97k \text{ for } n > k + 13.$$

We observe that 1.97 in (3) cannot be replaced by 2 since there are arbitrary long chains of consecutive composite positive integers. The same reason implies that Theorem 1 (a) is not valid under the assumption  $n > k + 13$ . Further the assumption  $n > \frac{279}{262}k$  in Theorem 1 (a) is necessary since  $P(\Delta(279, 262)) \leq 2 \times 262$ .

---

2000 Mathematics Subject Classification: Primary 11A41, 11N05, 11N13.  
Keywords: Arithmetic Progressions, Primes.

Now we give a lower bound for  $P(\Delta(n, k)) > 2k$  which is valid for  $n > k > 2$  except for an explicitly given finite set. For this, we need some notation. For a pair  $(n, k)$  and a positive integer  $h$ , we write  $[n, k, h]$  for the set of all pairs  $(n, k), \dots, (n + h - 1, k)$  and we set  $[n, k] = [n, k, 1] = \{(n, k)\}$ . Let

$$A_{10} = \{58\}; \quad A_8 = A_{10} \cup \{59\}; \quad A_6 = A_8 \cup \{60\};$$

$$A_4 = A_6 \cup \{12, 16, 46, 61, 72, 93, 103, 109, 151, 163\};$$

$$A_2 = A_4 \cup \{4, 7, 10, 13, 17, 19, 25, 28, 32, 38, 43, 47, 62, 73, 94, 104, 110, 124, 152, 164, 269\}$$

and  $A_{2i+1} = A_{2i}$  for  $1 \leq i \leq 5$ . Further let

$$A_1 = A_2 \cup \{3, 5, 6, 8, 9, 11, 14, 15, 18, 20, 23, 26, 29, 33, 35, 39, 41, 44, 48, 50, 53, \\ 56, 63, 68, 74, 78, 81, 86, 89, 95, 105, 111, 125, 146, 153, 165, 173, 270\}.$$

Finally we denote  $B$  as the union of the sets  $[8, 3]$ ,  $[5, 4, 3]$ ,  $[14, 13, 3]$  and  $\{(k + 1, k) | k = 3, 5, 8, 11, 14, 18, 63\}$ . Then

**Theorem 2.** *We have*

$$(4) \quad P(\Delta(n, k)) > 1.95k \text{ for } n > k > 2$$

*unless and only unless  $(n, k) \in [k + 1, k, h]$  for  $k \in A_h$  with  $1 \leq h \leq 11$  or  $(n, k) = (8, 3)$ .*

If  $k = 2$ , we observe (see Lemma 7) that  $P(\Delta(n, k)) > 2k$  unless  $n = 3, 8$  and that  $P(\Delta(3, 2)) = P(\Delta(8, 2)) = 3$ . Thus the estimate (4) is valid for  $k = 2$  whenever  $n \neq 3, 8$ . We observe that  $P(\Delta(k + 1, k)) \leq 2k$  and therefore, 1.95 in (4) cannot be replaced by 2. There are few exceptions if 1.95 is replaced by 1.8 in Theorem 2. We derive from Theorem 2 the following result.

**Corollary 1.** *We have*

$$(5) \quad P(\Delta(n, k)) > 1.8k \text{ for } n > k > 2$$

*unless and only unless  $(n, k) \in B$ .*

## 2. LEMMAS

We begin with a well known result due to Levi ben Gerson on a particular case of Catalan equation.

**Lemma 1.** *The solutions of*

$$2^a - 3^b = \pm 1 \text{ in integers } a > 0, b > 0$$

*are given by  $(a, b) = (1, 1), (2, 1), (3, 2)$ .*

Next we state a result of Saradha and Shorey [6] on a lower bound for  $\omega(\Delta(n, k))$ .

**Lemma 2.** For  $n > k > 2$ , we have

$$\omega(\Delta(n, k)) \geq \pi(k) + \left\lfloor \frac{1}{3}\pi(k) \right\rfloor + 2$$

unless and only unless  $(n, k)$  belongs to the union of sets

$$\left\{ \begin{array}{l} [4, 3], [6, 3, 3], [16, 3], [6, 4], [6, 5, 4], [12, 5], [14, 5, 3], [23, 5, 2], \\ [7, 6, 2], [15, 6], [8, 7, 3], [12, 7], [14, 7, 2], [24, 7], [9, 8], [14, 8], \\ [14, 13, 3], [18, 13], [20, 13, 2], [24, 13], [15, 14], [20, 14], [20, 17]. \end{array} \right.$$

We shall use Lemma 2 only when  $k = 3$  or  $5 \leq k \leq 8$ . Let  $p_i$  denote the  $i$ -th prime number. Then

**Lemma 3.** We have

(6)

$$p_{i+1} - p_i < \begin{cases} 35 & \text{for } p_i \leq 5591 \\ 15 & \text{for } p_i \leq 1123, p_i \neq 523, 887, 1069 \\ 21 & \text{for } p_i = 523, 887, 1069 \\ 9 & \text{for } p_i \leq 361, p_i \neq 113, 139, 181, 199, 211, 241, 283, 293, 317, 337. \end{cases}$$

**Lemma 4.** Let  $\mathfrak{N}$  be a positive real number and  $k_0$  a positive integer. Let  $I(\mathfrak{N}, k_0) = \{i | p_{i+1} - p_i \geq k_0, p_i \leq \mathfrak{N}\}$ . Then

$$P(n(n+1) \cdots (n+k-1)) > 2k$$

for  $2k \leq n < \mathfrak{N}$  and  $k \geq k_0$  except possibly when  $p_i < n < n+k-1 < p_{i+1}$  for  $i \in I(\mathfrak{N}, k_0)$ .

*Proof.* Let  $2k \leq n < \mathfrak{N}$  and  $k > k_0$ . We may suppose that none of  $n, n+1, \dots, n+k-1$  is a prime, otherwise the result follows. Let  $p_i < n < n+k-1 < p_{i+1}$ . Then  $i = \pi(n)$  and  $p_{\pi(n)} < n < \mathfrak{N}$ . For  $\pi(n) \notin I(\mathfrak{N}, k_0)$ , we have

$$k-1 = n+k-1-n < p_{\pi(n)+1} - p_{\pi(n)} < k_0$$

which implies  $k-1 < k_0-1$ , a contradiction. Hence the assertion.  $\square$

The following result is on the estimates for primes due to Dusart [1, p.14].

**Lemma 5.** For  $\nu > 1$ , we have

$$\begin{aligned} (i) \quad \pi(\nu) &\leq \frac{\nu}{\log \nu} \left( 1 + \frac{1.2762}{\log \nu} \right) \\ (ii) \quad \pi(\nu) &\geq \frac{\nu}{\log \nu - 1} \text{ for } \nu \geq 5393. \end{aligned}$$

**Lemma 6.** *Let  $X > 0$  and  $0 < \theta < e - 1$  be real numbers. For  $l \geq 0$ , let*

$$X_0 = \max \left( \frac{5393}{1 + \theta}, \exp\left(\frac{\log(1 + \theta) + 0.2762}{\theta}\right) \right),$$

$$X_{l+1} = \max \left( \frac{5393}{1 + \theta}, \exp\left(\frac{\log(1 + \theta) + 0.2762}{\theta + \frac{1.2762(1 - \log(1 + \theta))}{\log^2 X_l}}\right) \right).$$

Then we have

$$\pi((1 + \theta)X) - \pi(X) > 0$$

for  $X > X_l$ .

*Proof.* Let  $l \geq 0$  and  $X > X_l$ . Then  $(1 + \theta)X \geq 5393$ . By Lemma 5, we have

$$\begin{aligned} \delta := \pi((1 + \theta)X) - \pi(X) &\geq \frac{(1 + \theta)X}{\log(1 + \theta)X - 1} - \frac{X}{\log X} \left( 1 + \frac{1.2762}{\log X} \right) \\ &\geq \frac{X}{\log(1 + \theta)X - 1} \left\{ 1 + \theta - \frac{\log(1 + \theta)X - 1}{\log X} \left( 1 + \frac{1.2762}{\log X} \right) \right\} \\ &\geq \frac{X}{\log(1 + \theta)X - 1} \left\{ 1 + \theta - \left( 1 - \frac{1 - \log(1 + \theta)}{\log X} \right) \left( 1 + \frac{1.2762}{\log X} \right) \right\} \\ &\geq \frac{X}{\log(1 + \theta)X - 1} \{F(X) + G(X)\} \end{aligned}$$

where  $F(X) = \theta - \frac{\log(1 + \theta) + 0.2762}{\log X}$  and  $G(X) = \frac{1.2762(1 - \log(1 + \theta))}{\log^2 X}$ . We see that  $G(X) > 0$  and decreasing since  $0 < \theta < e - 1$ . Further we observe that  $\{X_i\}$  is a non-increasing sequence. We notice that  $\delta > 0$  if  $F(X) + G(X) > 0$ . But  $F(X) + G(X) > F(X) > 0$  for  $X > X_0$  by the definition of  $X_0$ . Thus  $\delta > 0$  for  $X > X_0$ . Let  $X \leq X_0$ . Then  $F(X) + G(X) \geq F(X) + G(X_0)$  and  $F(X) + G(X_0) > 0$  if  $X > X_1$  by the definition of  $X_1$ . Hence  $\delta > 0$  for  $X > X_1$ . Now we proceed inductively as above to see that  $\delta > 0$  for  $X > X_l$  with  $l \geq 2$ .  $\square$

**Lemma 7.** *Let  $n > k$  and  $k \leq 16$ . Then*

$$(7) \quad P(\Delta(n, k)) \leq 2k$$

*implies that  $(n, k) \in \{(8, 2), (8, 3)\}$  or  $(n, k) \in [k + 1, k]$  for  $k \in \{2, 3, 5, 6, 8, 9, 11, 14, 15\}$  or  $(n, k) \in [k + 1, k, 3]$  for  $k \in \{4, 7, 10, 13\}$  or  $(n, k) \in [k + 1, k, 5]$  for  $k \in \{12, 16\}$ .*

*Proof.* We apply Lemma 1 to derive that (7) is possible only if  $n = 3, 8$  when  $k = 2$  and  $n = 5, 6, 7$  when  $k = 4$ . For the latter assertion, we apply Lemma 1 after securing  $P((n+i)(n+j)) \leq 3$  with  $0 \leq i < j \leq 3$  by deleting the terms divisible by 5 and 7 in  $n, n+1, n+2$  and  $n+3$ . For  $k = 3$  and  $5 \leq k \leq 8$ , the assertion follows from Lemma 2.

Thus we may assume that  $k \geq 9$ . Assume that (7) holds. Then there are at most  $1 + \lfloor \frac{k-1}{p} \rfloor$  terms divisible by the prime  $p$ . After removing all the terms divisible by  $p \geq 7$ , we are left with at least 4 terms only divisible by 2, 3 and 5. Further out of these terms, for each prime 2, 3 and 5, we remove a term in which the prime divides to a maximal power. Then we are left with a term  $n+i$  such that  $n \leq n+i \leq 8 \times 9 \times 5 = 360$ . Let  $n \geq 2k$ . We now apply Lemma 4 with  $\mathfrak{N} = 361, k_0 = 9$  and (6) to get  $P(\Delta(n, k)) > 2k$  for  $k \geq 9$  except possibly when  $p_i < n < n+k-1 < p_{i+1}$ ,  $p_i = 113, 139, 181, 199, 211, 241, 283, 293, 317, 337$ . For these values of  $n$ , we check that  $P(\Delta(n, k)) > 2k$  is valid for  $9 \leq k \leq 16$ . Thus it suffices to consider  $k < n < 2k$ . We calculate  $P(\Delta(n, k))$  for  $(n, k)$  with  $9 \leq k \leq 16$  and  $k < n < 2k$ . We find that (7) holds only if  $(n, k)$  is given in the statement of the Lemma 7.  $\square$

### 3. PROOF OF THEOREM 1 (a)

Let  $n > \max(k + 13, \frac{279}{262}k)$ . In view of Lemma 7, we may take  $k \geq 17$  since  $n \leq k + 5$  for the exceptions  $(n, k)$  given in Lemma 7. It suffices to prove (2) for  $k$  such that  $2k - 1$  is prime. Let  $k_1 < k_2$  be such that  $2k_1 - 1$  and  $2k_2 - 1$  are consecutive primes. Suppose (2) holds at  $k_1$ . Then for  $k_1 < k < k_2$ , we have

$$P(n(n+1) \cdots (n+k-1)) \geq P(n \cdots (n+k_1-1)) > 2k_1$$

implying  $P(\Delta(n, k)) \geq 2k_2 - 1 > 2k$ . Therefore we may suppose that  $k \geq 19$  since  $2k - 1$  with  $k = 17, 18$  are composites. We assume from now onward in the proof of Theorem 1 (a) that  $2k - 1$  is prime. We put  $x = n + k - 1$ . Then  $\Delta(n, k) = x(x-1) \cdots (x-k+1)$ . Let  $f_1 < f_2 < \cdots < f_\mu$  be all the integers in  $[0, k)$  such that

$$(8) \quad P((x-f_1) \cdots (x-f_\mu)) \leq k.$$

We derive as in the proof of [4, Lemma 4] to get

$$(9) \quad k! > x^{\mu-\pi(k)} \left(1 - \frac{k}{x}\right)^\mu.$$

We may suppose  $\omega(\Delta(n, k)) \leq \pi(2k)$  otherwise (2) follows. Then

$$(10) \quad \mu \geq k - \pi(2k) + \pi(k)$$

which we use as in [4, Lemma 4] to derive from (9) that

$$(11) \quad x < k^{\frac{3}{2}} \text{ for } k \geq 87; \quad x < k^{\frac{7}{4}} \text{ for } k \geq 40; \quad x < k^2 \text{ for } k \geq 19.$$

If  $x \geq 7k$  and  $k > 57$ , then we derive as in [4, Lemma 7] from (10) that  $x \geq k^{\frac{3}{2}}$ . Thus we get from (11) that  $x < 7k$  for  $k \geq 87$ . Putting back  $n = x - k + 1$ , we may assume that  $n < 6k + 1$  for  $k \geq 87$ ,  $n < k^{\frac{7}{4}} - k + 1$  for  $40 \leq k < 87$  and  $n < k^2 - k + 1$  for  $19 \leq k < 40$ .

Let  $k < 87$ . Suppose  $n \geq 2k$ . Then  $2k \leq n < k^{\frac{7}{4}} - k + 1$  for  $40 \leq k < 87$  and  $2k \leq n < k^2 - k + 1$  for  $19 \leq k < 40$ . Thus Lemma 4 with  $\mathfrak{N} = 87^{\frac{7}{4}} - 87 + 1, k_0 = 35$  and (6) implies that  $P(\Delta(n, k)) > 2k$  for  $k \geq 35$ . We note here that  $2k \leq n < \mathfrak{N}$  for  $35 \leq k < 40$ . Let  $k < 35$ . Taking  $\mathfrak{N} = 34^2 - 34 + 1, k_0 = 21$  for  $21 \leq k \leq 34$  and  $\mathfrak{N} = 19^2 - 19 + 1, k_0 = 19$  for  $k = 19$ , we see from Lemma 4 and (6) that  $P(\Delta(n, k)) > 2k$  for  $k \geq 19$ . Here the case  $k = 20$  is excluded since  $2k - 1$  is composite. Therefore we may assume that  $n < 2k$ . Further we observe that  $\pi(n + k - 1) - \pi(2k) \geq \pi(2k + 13) - \pi(2k)$  since  $n > k + 13$ . Next we check that  $\pi(2k + 13) - \pi(2k) > 0$ . This implies that  $[2k, n + k - 1]$  contains a prime.

Thus we may assume that  $k \geq 87$ . Then we write  $n = \alpha k + 1$  with  $\frac{279}{262} - \frac{1}{k} < \alpha \leq 6$  if  $k \geq 201$  and  $1 + \frac{12}{k} < \alpha \leq 6$  if  $k < 201$ . Further we consider  $\pi(n + k - 1) - \pi(\max(n - 1, 2k))$  which is

$$\begin{aligned} &= \pi((\alpha + 1)k) - \pi(\alpha k) \quad \text{for } \alpha \geq 2 \\ &\geq \pi\left(\left[\frac{541}{262}k\right]\right) - \pi(2k) \quad \text{for } \alpha < 2 \text{ and } k \geq 201 \\ &\geq \pi(2k + 13) - \pi(2k) \quad \text{for } \alpha < 2 \text{ and } k < 201. \end{aligned}$$

We check by using exact values of  $\pi$  function that  $\pi(2k + 13) - \pi(2k) > 0$  for  $k < 201$  and  $\pi\left(\left[\frac{541}{262}k\right]\right) - \pi(2k) > 0$  for  $201 \leq k \leq 2616$ . Thus we may suppose that  $k > 2616$  if  $\alpha < 2$ . Also  $\left[\frac{541}{262}k\right] \geq \frac{540}{262}k$  for  $k > 2616$ . Now we apply Lemma 6 with  $X = \alpha k, \theta = \frac{1}{\alpha}, l = 0$  if  $\alpha \geq 2$  and  $X = 2k, \theta = \frac{4}{131}, l = 1$  if  $\alpha < 2$  to get  $\pi(n + k - 1) - \pi(\max(n - 1, 2k)) > 0$  for  $X > X_0 = \frac{5393}{1 + \frac{1}{\alpha}}$  if  $\alpha \geq 2$  and  $X > X_1 = \frac{5393}{1 + \frac{4}{131}}$  if  $\alpha < 2$ . Further when  $\alpha < 2$ , we observe that  $X = 2k > X_1$  since  $k > 2616$ . Thus the assertion follows for  $n < 2k$ . It remains to consider the case  $\alpha \geq 2$  and  $X \leq 5393(1 + \frac{1}{\alpha})^{-1}$ . Then  $2k \leq n < n + k - 1 = X(1 + \frac{1}{\alpha}) \leq 5393$ . Now we apply Lemma 4 with  $\mathfrak{N} = 5393, k_0 = 35$  and (6) to conclude that  $P(\Delta(n, k)) > 2k$ .  $\square$

## 4. PROOF OF THEOREM 1 (b)

In view of Lemma 7 and Theorem 1 (a), we may assume that  $k \geq 17$  and  $k < n \leq \frac{279}{262}k$ . Let  $X = \frac{279}{262}k$ ,  $\theta = \frac{245}{279}$ ,  $l = 0$ . Then for  $k < n \leq X$ , we see from Lemma 6 that

$$\pi(2k) - \pi(n-1) \geq \pi((1+\theta)X) - \pi(X) > 0$$

for  $X > X_0 = 5393(1+\theta)^{-1}$  which is satisfied for  $k > 2696$  since  $(1+\theta)X = 2k$ . Thus we may suppose that  $k \leq 2696$ . Now we check with exact values of  $\pi$  function that  $\pi(2k) - \pi(\frac{279}{262}k) > 0$ . Therefore  $P(\Delta(n, k)) \geq P(n(n+1) \cdots 2k) \geq p_{\pi(2k)}$ . Further we apply Lemma 6 with  $X = 1.97k$ ,  $\theta = \frac{3}{197}$  and  $l = 25$ . We calculate that  $X_l \leq 284000$ . We conclude by Lemma 6 that

$$\pi(2k) - \pi(1.97k) = \pi((1+\theta)X) - \pi(X) > 0$$

for  $k > 145000$ . Let  $k \leq 145000$ . Then we check that  $\pi(2k) - \pi(1.97k) > 0$  is valid for  $k \geq 680$  by using the exact values of  $\pi$  function. Thus

$$(12) \quad p_{\pi(2k)} > 1.97k \text{ for } k \geq 680.$$

Therefore we may suppose that  $k < 680$ . Now we observe that for  $n > k+13$ ,  $\pi(n+k-1) - \pi(1.97k) \geq \pi(2k+13) - \pi(1.97k) > 0$ , the latter inequality can be checked by using exact values of  $\pi$  function. Hence the assertion follows since  $n < 1.97k$ .  $\square$

## 5. PROOF OF THEOREM 2

By Theorem 1 (b), we may assume that  $n \leq k+13$ . Also we may suppose that  $k < 680$  by (12). For  $k \leq 16$ , we calculate  $P(\Delta(n, k))$  for all the pairs  $(n, k)$  given in the statement of Lemma 7. We find that either  $P(\Delta(n, k)) > 1.95k$  or  $(n, k)$  is an exception stated in Theorem 1 (a). Thus we may suppose that  $k \geq 17$ . Now we check that  $\pi(n+k-1) - \pi(1.95k) > 0$  except for  $(n, k) \in [k+1, k, h]$  for  $k \in A_h$  with  $1 \leq h \leq 11$  and the assertion follows.  $\square$

## 6. PROOF OF COROLLARY 1

We calculate  $P(\Delta(n, k))$  for all  $(n, k)$  with  $k \leq 270$  and  $k+1 \leq n \leq k+11$ . This contains the set of exceptions given in Theorem 2. We find that  $P(\Delta(n, k)) > 1.8k$  unless  $(n, k) \in B$ . Hence the assertion (5) follows from Theorem 2.  $\square$

## REFERENCES

- [1] Pierre Dusart, *Autour de la fonction qui compte le nombre de nombres premiers*, Ph.D thesis, Université de Limoges, 1998.
- [2] M. Faulkner, *On a theorem of Sylvester and Schur*, J. Lond. Math. Soc., **41** (1966), 107-110.
- [3] D. Hanson, *On a theorem of Sylvester and Schur*, Canad. Math. Bull., **16** (1973), 195-199.
- [4] S. Laishram and T. N. Shorey, *Number of prime divisors in a product of consecutive integers*, Acta Arith. **113** (2004), 327-341.
- [5] L. Moser, *Insolvability of  $\binom{2n}{n} = \binom{2a}{a}\binom{2b}{b}$* , Canad. Math. Bull.(2), **6** (1963), 167-169.
- [6] N. Saradha and T. N. Shorey, *Almost squares and factorisations in consecutive integers*, Compositio Math. **138** (2003), 113-124.
- [7] J. J. Sylvester, *On arithmetical series*, Messenger of Mathematics, **XXI** (1892), 1-19, 87-120, and Mathematical Papers, **4**(1912), 687-731.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH,  
HOMI BHABHA ROAD, MUMBAI 400005, INDIA

*E-mail address:* shanta@math.tifr.res.in

*E-mail address:* shorey@math.tifr.res.in