



Published in final edited form as:

Nat Cell Biol. ; 13(8): 877–883. doi:10.1038/ncb2303.

The Hippo pathway in organ size control, tissue regeneration and stem cell self-renewal

Bin Zhao,

Life Sciences Institute, Zhejiang University, Hangzhou, Zhejiang 310058, China

Department of Pharmacology and Moores Cancer Center, University of California at San Diego, La Jolla, California 92093-0815, USA

Karen Tumaneng, and

Department of Pharmacology and Moores Cancer Center, University of California at San Diego, La Jolla, California 92093-0815, USA

Kun-Liang Guan

Department of Pharmacology and Moores Cancer Center, University of California at San Diego, La Jolla, California 92093-0815, USA

Kun-Liang Guan: kuguan@ucsd.edu

Abstract

Precise control of organ size is crucial during animal development and regeneration. In *Drosophila* and mammals, studies over the past decade have uncovered a critical role for the Hippo tumour-suppressor pathway in the regulation of organ size. Dysregulation of this pathway leads to massive overgrowth of tissue. The Hippo signalling pathway is highly conserved and limits organ size by phosphorylating and inhibiting the transcription co-activators YAP and TAZ in mammals and Yki in *Drosophila*, key regulators of proliferation and apoptosis. The Hippo pathway also has a critical role in the self-renewal and expansion of stem cells and tissue-specific progenitor cells, and has important functions in tissue regeneration. Emerging evidence shows that the Hippo pathway is regulated by cell polarity, cell adhesion and cell junction proteins. In this review we summarize current understanding of the composition and regulation of the Hippo pathway, and discuss how cell polarity and cell adhesion proteins inform the role of this pathway in organ size control and regeneration.

Organ size regulation is a highly coordinated process involving complex mechanisms in response to physiological cues. On the organismal level, circulating factors such as hormones and insulin-like growth factors (IGF) play important roles in promoting organ size¹. In contrast, physiological perturbations, such as prolonged starvation, cause profound reduction of organ size¹. Additionally, an intrinsic mechanism limits organ size, which was first demonstrated in salamander limbs by classical transplantation experiments¹. The underlying mechanism of organ-autonomous size determination remained largely unknown

until the past decade. Extensive research led to the identification of the Hippo tumour-suppressor pathway as a key regulator of organ size in *Drosophila* and mammals². It is also known that mutations of genes that are involved in patterning, cell polarity and cell adhesion cause marked alternations of organ size³. Thus, the recent finding that the Hippo pathway is regulated by cell polarity and cell adhesion proteins is a promising basis for the potential crosstalk of the Hippo pathway and cell polarity proteins in the regulation of organ size⁴. Several studies have also demonstrated important roles for the Hippo pathway in stem cell/progenitor cell expansion and tissue regeneration^{5–13}. These findings will be discussed here.

The Hippo pathway in *Drosophila*

In *Drosophila*, the first core components of the Hippo pathway to be identified, using genetic mosaic screens, were the tumour-suppressor genes *warts* (*wts*)^{14,15}, *hippo* (*hpo*)^{16–20} and *salvador* (*sav*)^{21,22}. These genes belong to the hyperplastic group of *Drosophila* tumour-suppressors. Mutation of these genes results in robust tissue overgrowth without alteration of cell fate determination or cell polarity. Biochemical studies revealed that Hpo directly interacts with Sav to phosphorylate and activate the complex formed by Wts and another core Hippo pathway protein, Mats^{16,23} (Fig. 1a). The kinase activity of Hpo is antagonized by a PP2A phosphatase complex, dSTRIPAK²⁴. The Hippo pathway is known to limit organ size partly by transcriptional regulation of *cyclin E* and *diap1* (refs 16,17,20,21,23), suggesting the existence of a transcriptional regulator as a downstream effector of the pathway. By performing a yeast two-hybrid screen using Wts as bait, the transcription co-activator Yorkie (Yki) was identified as a potent effector of the Hippo pathway²⁵. Subsequent biochemical studies showed that Wts directly phosphorylates and inhibits Yki²⁶.

Research in the past years has uncovered many proteins that act upstream in the *Drosophila* Hippo pathway. Two apical cytoskeleton-binding proteins, Merlin (Mer) and Expanded (Ex)²⁷, and their interacting protein Kibra^{28–30}, were found to activate the Hippo pathway. The Fat protocadherin, a cell-surface molecule, was also identified as an upstream regulator of the Hippo pathway^{31–35}. Fat activity is regulated by binding to another protocadherin, Dachshous (Ds)³⁶, and is modulated by several proteins, such as the casein kinase Discs overgrown (Dco)^{37,38}, the Golgiresident kinase Four-jointed (Fj)^{39–41} and the Fat/Ds-interacting protein Lowfat (Lft)⁴². Fat/Hippo pathway activity may also be influenced by Decapentaplegic (Dpp) and Wingless (Wg) morphogen gradients^{40,41,43}, which affect the expression of Fj and Ds. It has been proposed that Fat activates the Hippo pathway by regulating the protein level and localization of the protein Ex^{31–33,35}. Another study suggests that Fat may control the abundance of Wts through Dachs^{34,44}. Recently, dJub, a LIM-domain-containing protein that physically interacts with Wts and Sav, was shown to negatively regulate Hippo signalling, although the detailed mechanism has not been delineated⁴⁵. A number of proteins that determine cell polarity were also found to regulate the Hippo pathway. These include the Scribble (Scrib)–Discs large (Dlg)–Lethal giant larvae (Lgl) complex, atypical protein kinase C (aPKC) and Crumbs (Crb)^{46–49}, indicating a role of cell polarity in the regulation of Hippo signalling.

The Hippo pathway in mammals

The core components and downstream effectors of the *Drosophila* Hippo pathway are highly conserved in mammals: Mst1/2 (homologues of Hpo), Sav1 (Sav homologue), Lats1/2 (Wts homologues), MOBKL1A and MOBKL1B (collectively referred to as Mob1; homologues of Mats), and YAP and its paralogue TAZ (also called WWTR1; homologues of Yki) (Fig. 1b). Expression of human YAP, Lats1, Mst2 and Mob1 can rescue the phenotypes of their corresponding *Drosophila* mutants *in vivo*^{16,23,25,50}. The core components Mst1/2 are pro-apoptotic kinases that are activated by caspase cleavage under apoptotic stress⁵¹. Sav1 interacts with Mst1/2 through the SARAH domains present in both Sav1 and Mst1/2 (ref. 52). Although Sav1 has been shown to activate Mst1/2, the underlying mechanism is unclear, but might involve regulation of Mst1 nuclear translocation⁵³. Mst1/2 is also activated by binding to Ras association domain family (RASSF) proteins⁵⁴, possibly owing to alteration of Mst1/2 subcellular localization⁵⁵. In *Drosophila*, however, dRASSF inhibits Hpo possibly through competition with Sav for Hpo binding⁵⁶ and through recruitment of the dSTRIPAK–PP2A complex²⁴. Activation of Mst1/2 leads to phosphorylation and activation of their direct substrates, Lats1/2 (ref. 57). Mob1, which forms a complex with Lats1/2, is also phosphorylated by Mst1/2, resulting in an enhanced Lats1/2–Mob1 interaction⁵⁸. Activated Lats1/2 in turn phosphorylate and inhibit YAP/TAZ transcription co-activators^{26,59–62}.

Functions of the Hippo pathway in organ size determination and tumour suppression have been confirmed in genetically engineered mouse models. For instance, liver-specific overexpression of YAP results in enlarged livers that return to their normal size after cessation of YAP expression^{12,26}. However, sustained YAP overexpression leads to tumour formation²⁶. Genomic amplification of YAP is also observed in human cancers and a mouse model of breast cancer^{63,64}. Furthermore, elevated YAP protein levels and nuclear localization have been observed in multiple human cancers^{59,63,65}, and the alterations of YAP may have prognostic value for certain human cancers⁶⁶. Overexpression of TAZ, the paralogue of YAP, has been noted in human breast cancer samples and non-small-cell lung-cancer cell lines^{67,68}. Ablation of the Hippo pathway components *Mer* and *Sav* and double knockout of *Mst1/2* in mice also result in liver enlargement and tumour formation^{69–74}. Remarkably, loss of one or both copies of *YAP* can suppress liver expansion and tumorigenesis induced by *Mer* deficiency⁶⁹. Aberrant Mst1/2 and Lats1/2 expression and *Lats2*, *Sav1* and *Mob1* mutation were also observed in human cancers or cancer cell lines². Together, these studies highlight a significant role of the Hippo pathway in organ size regulation and tumorigenesis.

Mechanisms of YAP/TAZ/Yki inhibition

Activation of the Hippo pathway leads to phosphorylation and inhibition of YAP, TAZ and Yki transcription co-activators. In mammals, YAP and TAZ are phosphorylated by Lats1/2 *in vitro* and *in vivo*^{59,60,75}. The mechanism of inhibition by Hippo signalling involves phosphorylation of Ser 127 in YAP or the corresponding sites in TAZ and Yki, which promotes 14-3-3 binding and subsequent cytoplasmic sequestration and inactivation^{26,59,60,62,76} (Fig. 2a). Indeed, mutation of Ser 127 and disruption of 14-3-3

binding activate YAP⁵⁹, confirming the inhibitory nature of this phosphorylation. In *Drosophila*, Yki phosphorylation on two other sites by Wts similarly results in Yki inhibition, although the mechanism is yet to be determined⁷⁷.

Phosphorylation of YAP can also induce its degradation. Lats1/2 phosphorylates YAP at Ser 381, which primes YAP for subsequent phosphorylation by another kinase, possibly casein kinase 1 (CK1 δ/ϵ), activating a phosphorylation-dependent degradation motif termed a phosphodegron. Subsequently, the E3 ubiquitin ligase SCF β -TRCP is recruited to YAP, leading to its polyubiquitylation and degradation⁷⁵ (Fig. 2c). Consistently, decreased YAP phosphorylation in sparsely cultured NIH-3T3 cells, as well as in Mst1/2-deficient mouse liver, correlates with increased YAP protein levels^{73,75}. This mechanism is conserved in TAZ but not in Yki⁷⁸, which lacks a residue equivalent to Ser 381.

YAP, TAZ and Yki can also be inhibited through protein–protein interactions that result in their cytoplasmic sequestration (Fig. 2b). Yki contains two WW domains that can interact with PPXY motifs present in Mop⁷⁹ and the Hippo pathway components Ex, Wts and Hpo^{80,81}. Recently, YAP/TAZ and angiomin (AMOT) family proteins were shown to interact^{82–85}, resulting in YAP/TAZ localization to tight junctions and inhibition through phosphorylation-dependent and -independent mechanisms⁸². In addition, YAP and TAZ interact with another tight junction protein ZO-2, which was reported to increase nuclear localization of YAP and tight-junction localization of TAZ^{86,87}. It will be important to investigate the relationship between phosphorylation and these physical interactions in YAP regulation, and whether disruption of these interactions alters organ growth.

Transcriptional regulation of Hippo pathway target genes by YAP, TAZ and Yki

The TEAD family transcription factors were found to be critical partners of YAP and TAZ in the regulation of gene expression (the *Drosophila* TEAD homologue Scalloped (Sd) is partner of Yki)^{88–92}. Knockdown of TEADs or disruption of the YAP–TEAD interaction abolishes YAP-dependent gene transcription and largely diminishes YAP-induced cell proliferation, oncogenic transformation and the epithelial-to-mesenchymal transition (EMT)⁸⁸. In *Drosophila*, Sd was shown to genetically interact with Yki and to be required for Yki-induced target gene expression *in vivo*^{88,89,91,92}. Intriguingly, a mutation of TEAD1 Tyr 406, which forms a hydrogen bond with YAP, results in loss of interaction with YAP and leads to the human genetic disease Sveinsson’s chorioretinal atrophy^{93–96}. Precise regulation of YAP–TEAD interaction is therefore important in maintaining normal physiology.

Several direct target genes of YAP–/TAZ–TEAD and Yki–Sd have been identified, including *CTGF* and *Cyr61* in mammalian cells^{88,97}, and *diap1* and *dMyc* in *Drosophila*^{89,91,98,99}. *CTGF* was shown to have an important role in YAP-induced proliferation and anchorage-independent growth⁸⁸. In *Drosophila*, *diap1* is essential for Yki-induced overgrowth, but is not sufficient to explain all Yki phenotypes. Recently, Yki–Sd was shown to induce transcription of *dMyc*, a potent promoter of ribosome biogenesis and cell growth^{98,99}. *dMyc* expression also mediates a cell phenomenon induced by imbalance of

Hippo pathway activity, referred to as cell competition — wherein the contact between fast- and slow-growing cells in genetic mosaics favours the positive selection and clonal expansion of fast-dividing cells at the expense of slow-dividing cells^{98,99}. YAP also induces *Myc* in transgenic mouse liver²⁶, although the mechanism remains to be investigated.

Despite a major role for TEADs in YAP/TAZ function, other transcription factors containing PPXY motifs are known to interact with the WW domains of YAP/TAZ. These include Smad1, RUNX, ErbB4 and p73 for YAP^{100–104}, and RUNX, PPAR γ , Pax3, TBX5 and TTF-1 for TAZ^{105–109}. The interaction of YAP with Smad1 is believed to be important for BMP-mediated maintenance of pluripotency of mouse embryonic stem cells¹⁰⁴. YAP and TAZ also bind Smad2/3 through the coiled-coil region, and this interaction is believed to dictate the subcellular localization of Smad2/3 (refs 85,110). YAP also interacts with p73, a p53 family pro-apoptotic transcription factor, to induce expression of genes such as *Bax*, *puma* and *PML*¹¹¹. However, there are contradictory reports on the role of the Hippo pathway in activating¹¹² or inhibiting⁶¹ this activity. Recently, YAP was also shown to interact with β -catenin and induce expression of canonical Wnt target genes such as *Sox2* and *Snai2* in mouse heart tissue¹¹³.

bantam microRNA is a target gene of the Hippo pathway and promotes cell survival and proliferation^{114,115}. Homothorax (Hth) and Teashirt (Tsh) are two transcription factors mediating *bantam* expression anterior to the morphogenetic furrow¹¹⁶. In addition, the expression of *bantam* is also directly induced by a transcriptional complex formed by Yki and Mad, an effector of the *Drosophila* Dpp signalling pathway¹¹⁷. The existence of a *bantam* counterpart and the functions of Hth and Tsh homologues in the Hippo pathway in mammals remain to be investigated.

YAP, TAZ and Yki also induce many other genes directly or indirectly. In *Drosophila*, Yki induces: *cycE* (ref. 21) and *E2F1* (ref. 92), which may be involved in cell-autonomous regulation of cell proliferation; the EGFR (epidermal growth factor receptor) ligands *Vein*, *Keren* and *Spitz*^{11,118} and the Jak–Stat pathway ligands *Unpaired1/2/3* (*Upd1/2/3*)^{8–11}, which might mediate non-cell-autonomous functions of the Hippo pathway; and the Hippo pathway genes *Ex*, *Kibra*, *Crb*, and *Fj*^{27,29,34,119}, which may constitute a signal feedback loop. In mammals, YAP and TAZ also induce the expression of *AREG*¹¹⁸ and *FGF1* (ref. 60), which may also mediate non-cell-autonomous functions of the Hippo pathway. However, the mechanisms underlying the induction of these genes, including the responsible transcription factors, are mostly unclear.

Regulation of the Hippo pathway by cell polarity and cell adhesion complexes

In *Drosophila*, mutations of several genes that are involved in cell polarity and cell junction lead to massive overgrowth. The Dlg–Lgl–Scrib protein complex localizes to the basal–lateral membrane of epithelial cells, where it is required for the lateral exclusion of apical proteins, including the Par3–Par6–aPKC complex and the Crb–Stardust (Sdt)–PATJ complex. Interestingly, *Lgl* mutations lead to nuclear translocation of Yki and upregulation of Hippo pathway target genes in *Drosophila* epithelium⁴⁷. Expression of dominant-

negative aPKC rescued the tissue overgrowth in *Lgl*-mutant tissues⁴⁷. In zebrafish, Scrib was shown to interact genetically with and suppress the activity of the YAP homologue during embryonic kidney development¹²⁰. The tumour-suppressor function of the Dlg–Lgl–Scrib complex is possibly conserved in mammals, as depletion of Scrib in mammary epithelium results in disruption of apoptosis inhibition by cell polarity, and induction of dysplasia *in vivo* that progresses to tumours after long latency¹²¹. It would be interesting to determine whether the mammalian Hippo pathway mediates the tumour-suppressor function of the Dlg–Lgl–Scrib complex.

Crb is another cell polarity protein that regulates the *Drosophila* Hippo pathway^{46,48,49}. The intracellular domain of Crb contains a juxtamembrane FERM-binding motif (FBM) and a carboxy-terminal PDZ-binding motif (PBM). The PBM is important for polarity formation¹²², whereas the FBM regulates Hippo-pathway-dependent proliferation and apoptosis by promoting apical localization of the upstream component Ex^{46,48,49}. Thus, Crb regulates cell polarity and tissue growth through distinct mechanisms. In addition, it seems that the functions of the Dlg–Lgl–Scrib complex in cell polarity and tissue growth are also separable^{47,123}. It is therefore important to determine whether, and how, the two functions of these proteins are coupled to regulate tissue homeostasis.

In mammalian cells, Hippo pathway activation is triggered in part by cell–cell contact. In tissue culture, high cell density induces YAP phosphorylation and cytoplasmic translocation⁵⁹. And in mouse blastocysts, YAP is nuclear in outer layer cells, and cytoplasmic in the inner blastocyst layer cells¹²⁴. Consistently, it has been observed that disruption of cell–cell junctions in epithelium results in the nuclear localization of YAP and TAZ⁸⁵. Collectively, these studies suggest that maintenance of cell–cell junctions is important for mammalian Hippo pathway function.

Recent studies shed some light into the mechanisms of YAP/TAZ regulation by cell–cell contact. First, a tight-junction protein complex, composed of the AMOT family proteins, PALS1, PATJ/MPDZ and Lin7, was found to interact with YAP and TAZ^{82–85}. This interaction inhibits YAP and TAZ by promoting their localization to tight junctions and their phosphorylation by the Hippo pathway. In addition, α -catenin was shown to interact with YAP^{125,126}, possibly through a 14-3-3 protein, in a phosphorylation-dependent manner¹²⁶. This interaction may prevent YAP dephosphorylation by PP2A and results in YAP inhibition¹²⁶. Thus, it is possible that the tight junction and adherens junction are critically important for relaying cell contact signals to the Hippo pathway. Such a hypothesis needs to be further investigated.

The Hippo pathway in tissue regeneration, and stem cell self-renewal and expansion

The Hippo pathway was initially thought to limit organ size by inhibiting proliferation and promoting apoptosis^{16–20}. However, emerging evidence suggests that the Hippo pathway may also regulate stem cell and progenitor cell self-renewal and expansion. For instance, YAP and TAZ regulate embryonic stem cell self-renewal in response to TGF β /BMP (transforming growth factor beta/bone morphogenetic protein) signalling^{104,110}. In addition,

YAP is inactivated during mouse embryonic stem cell differentiation and activated in induced pluripotent stem (iPS) cells⁵. YAP knockdown in mouse embryonic stem cells leads to loss of pluripotency, whereas ectopic expression of YAP prevents embryonic stem cell differentiation⁵.

Additionally, the Hippo pathway also regulates tissue-specific progenitor cells. YAP expression is generally restricted to the progenitor cells in normal mouse intestines, and transgenic expression of YAP in mouse intestines causes a marked expansion of the progenitor cell compartment¹². Activation of YAP-TEAD also results in the expansion of neural progenitor cells in a chicken neural tube model¹³. Similarly, YAP expression expands basal epidermal progenitors in mouse skin and inhibits their terminal differentiation¹²⁷. In contrast, conditional knockout of YAP or knock-in of a TEAD-binding-deficient YAP mutant in mouse skin leads to decreased proliferation of basal cells, thinner epidermis and failure of skin expansion¹²⁶. Consistently, adult liver stem cells known as oval cells accumulate in *Mst1/2*-, *Sav1*- and *Mer*-knockout mice liver⁷⁰⁻⁷³. It should be noted that these genetic manipulations are applied at the whole organ level and not specifically to the progenitor cell compartment. However, the contribution of progenitor cell expansion in YAP-induced organ overgrowth is likely to be tissue-dependent. For instance, overgrown hearts induced by *Sav1* knockout showed excessive proliferation in cardiomyocytes but normal proliferation level of cardiac progenitors¹¹³. In addition, in certain cancers, such as a subtype of medulloblastomas, YAP expression is highly elevated in the perivascular cancer stem cell compartment¹²⁸.

The Hippo pathway was recently shown to be involved in tissue regeneration. In the *Drosophila* midgut, Yki expression is largely restricted to intestinal stem cells (ISC)¹⁰. Under resting conditions, Yki is mostly localized to the cytoplasm and seems to be inactive¹⁰. In contrast, Yki displays increased nuclear localization and reporter activity, and has an important and cell-autonomous role in ISC proliferation in response to injury^{9,10}. Interestingly, the Hippo pathway also has a non-cell-autonomous function during regeneration^{8,9,11}. In response to damage, the Hippo pathway is inactivated in enterocytes, a differentiated cell type in the *Drosophila* midgut, resulting in Yki activation and subsequent expression of Upd1/2/3 (refs 8,9,11), as well as EGFR ligands¹¹. This results in increased ISC proliferation in a non-cell-autonomous manner. Yki activation in enterocytes and in wing discs (where Yki also plays a role in regeneration⁶) seems to involve JNK signalling^{8,129}.

In mammals, there is also evidence for a role of YAP in tissue regeneration. Intestinal damage markedly induces YAP expression, and loss of YAP severely impairs dextran sodium sulfate-induced intestinal regeneration⁷. In the mouse liver, *Yap* knockout causes a defect in bile duct development⁶⁹. Interestingly, most adult mouse biliary ductal epithelial cells express Sox9 and these cells make a significant contribution to liver regeneration after injury as shown by lineage tracing¹³⁰. It remains to be determined whether ablation of *YAP* also results in compromised liver regeneration, and more importantly, whether the Hippo pathway activity is regulated during regeneration in mammals.

Conclusions and perspectives

Extensive studies in the past decade have elucidated the importance of the Hippo pathway in organ size control and regeneration in both *Drosophila* and mammals. Several mechanisms have been proposed, and it is clear that cell adhesion and polarity complexes play a key role in Hippo pathway regulation. YAP and Yki may promote organ size and regeneration by inducing stem cell and progenitor cell proliferation through both cell-autonomous and non-cell-autonomous mechanisms (Fig. 3a). In addition, inactivation of the Hippo pathway may block cell-cycle exit, leading to hyperplasia and differentiation defects⁵³ (Fig. 3b). The Hippo pathway can also inhibit proliferation and promote apoptosis in non-stem cells/non-progenitor cell types (Fig. 3c). Lastly, an imbalance of Hippo pathway activity in neighbouring cells may induce cell competition through differential expression of *dMyc* in *Drosophila*^{98,99} (Fig. 3d). How these mechanisms fit into organ size regulation and regeneration *in vivo* is yet to be determined.

Despite these insights into the critical role of this pathway in stem cell expansion and tissue regeneration, many important questions await answers. These include the role and mechanism of cell polarity and cell adhesion proteins in sensing organ size to regulate the Hippo pathway and the position of the Hippo pathway in the known signalling networks regulating cell proliferation, apoptosis and stem cell function. In addition, the mechanism by which upstream regulators of the Hippo pathway are integrated to initiate or terminate signalling is not yet fully understood. Importantly, Hippo pathway dysregulation in cancer remains to be fully elucidated. The Hippo–YAP pathway holds great promise as a target in cancer therapy and regenerative medicine. Insights into the upstream regulators and downstream targets of this pathway, and their mechanism of regulation, are crucial in translating our basic knowledge of this pathway into therapeutic designs.

Acknowledgments

We apologize for those primary works that we could not cite due to space constraints. Research in the lab of K.L.G. is supported by grants from NIH and CIRM.

References

1. Stanger BZ. Organ size determination and the limits of regulation. *Cell Cycle*. 2008; 7:318–324. [PubMed: 18235243]
2. Pan D. The Hippo signaling pathway in development and cancer. *Dev. Cell*. 2010; 19:491–505. [PubMed: 20951342]
3. Leever SJ, McNeill H. Controlling the size of organs and organisms. *Curr. Opin. Cell Biol*. 2005; 17:604–609. [PubMed: 16226450]
4. Halder G, Johnson RL. Hippo signaling: growth control and beyond. *Development*. 2011; 138:9–22. [PubMed: 21138973]
5. Lian I, et al. The role of YAP transcription coactivator in regulating stem cell self-renewal and differentiation. *Genes Dev*. 2010; 24:1106–1118. [PubMed: 20516196]
6. Grusche FA, Degoutin JL, Richardson HE, Harvey KF. The Salvador/Warts/Hippo pathway controls regenerative tissue growth in *Drosophila melanogaster*. *Dev. Biol*. 2010; 350:255–266. [PubMed: 21111727]
7. Cai J, et al. The Hippo signaling pathway restricts the oncogenic potential of an intestinal regeneration program. *Genes Dev*. 2010; 24:2383–2388. [PubMed: 21041407]

8. Staley BK, Irvine KD. Warts and Yorkie mediate intestinal regeneration by influencing stem cell proliferation. *Curr. Biol.* 2010; 20:1580–1587. [PubMed: 20727758]
9. Shaw RL, et al. The Hippo pathway regulates intestinal stem cell proliferation during *Drosophila* adult midgut regeneration. *Development.* 2010; 137:4147–4158. [PubMed: 21068063]
10. Karpowicz P, Perez J, Perrimon N. The Hippo tumor suppressor pathway regulates intestinal stem cell regeneration. *Development.* 2010; 137:4135–4145. [PubMed: 21098564]
11. Ren F, et al. Hippo signaling regulates *Drosophila* intestine stem cell proliferation through multiple pathways. *Proc. Natl Acad. Sci. USA.* 2010; 107:21064–21069. [PubMed: 21078993]
12. Camargo FD, et al. YAP1 increases organ size and expands undifferentiated progenitor cells. *Curr. Biol.* 2007; 17:2054–2060. [PubMed: 17980593]
13. Cao X, Pfaff SL, Gage FH. YAP regulates neural progenitor cell number via the TEA domain transcription factor. *Genes Dev.* 2008; 22:3320–3334. [PubMed: 19015275]
14. Justice RW, Zilian O, Woods DF, Noll M, Bryant PJ. The *Drosophila* tumor suppressor gene warts encodes a homolog of human myotonic dystrophy kinase and is required for the control of cell shape and proliferation. *Genes Dev.* 1995; 9:534–546. [PubMed: 7698644]
15. Xu T, Wang W, Zhang S, Stewart RA, Yu W. Identifying tumor suppressors in genetic mosaics: the *Drosophila lats* gene encodes a putative protein kinase. *Development.* 1995; 121:1053–1063. [PubMed: 7743921]
16. Wu S, Huang J, Dong J, Pan D. Hippo encodes a Ste-20 family protein kinase that restricts cell proliferation and promotes apoptosis in conjunction with salvador and warts. *Cell.* 2003; 114:445–456. [PubMed: 12941273]
17. Udan RS, Kango-Singh M, Nolo R, Tao C, Halder G. Hippo promotes proliferation arrest and apoptosis in the Salvador/Warts pathway. *Nat. Cell Biol.* 2003; 5:914–920. [PubMed: 14502294]
18. Pantalacci S, Tapon N, Leopold P. The Salvador partner Hippo promotes apoptosis and cell-cycle exit in *Drosophila*. *Nat. Cell Biol.* 2003; 5:921–927. [PubMed: 14502295]
19. Harvey KF, Pflieger CM, Hariharan IK. The *Drosophila* Mst ortholog, Hippo, restricts growth and cell proliferation and promotes apoptosis. *Cell.* 2003; 114:457–467. [PubMed: 12941274]
20. Jia J, Zhang W, Wang B, Trinko R, Jiang J. The *Drosophila* Ste20 family kinase dMST functions as a tumor suppressor by restricting cell proliferation and promoting apoptosis. *Genes Dev.* 2003; 17:2514–2519. [PubMed: 14561774]
21. Tapon N, et al. *salvador* promotes both cell cycle exit and apoptosis in *Drosophila* and is mutated in human cancer cell lines. *Cell.* 2002; 110:467–478. [PubMed: 12202036]
22. Kango-Singh M, et al. Shar-pei mediates cell proliferation arrest during imaginal disc growth in *Drosophila*. *Development.* 2002; 129:5719–5730. [PubMed: 12421711]
23. Lai ZC, et al. Control of cell proliferation and apoptosis by mob as tumor suppressor, mats. *Cell.* 2005; 120:675–685. [PubMed: 15766530]
24. Ribeiro PS, et al. Combined functional genomic and proteomic approaches identify a PP2A complex as a negative regulator of Hippo signaling. *Mol. Cell.* 2010; 39:521–534. [PubMed: 20797625]
25. Huang J, Wu S, Barrera J, Matthews K, Pan D. The Hippo signaling pathway coordinately regulates cell proliferation and apoptosis by inactivating Yorkie, the *Drosophila* Homolog of YAP. *Cell.* 2005; 122:421–434. [PubMed: 16096061]
26. Dong J, et al. Elucidation of a universal size-control mechanism in *Drosophila* and mammals. *Cell.* 2007; 130:1120–1133. [PubMed: 17889654]
27. Hamaratoglu F, et al. The tumour-suppressor genes NF2/Merlin and Expanded act through Hippo signalling to regulate cell proliferation and apoptosis. *Nat. Cell Biol.* 2006; 8:27–36. [PubMed: 16341207]
28. Baumgartner R, Poernbacher I, Buser N, Hafen E, Stocker H. The WW domain protein Kibra acts upstream of Hippo in *Drosophila*. *Dev. Cell.* 2010; 18:309–316. [PubMed: 20159600]
29. Genevet A, Wehr MC, Brain R, Thompson BJ, Tapon N. Kibra is a regulator of the Salvador/Warts/Hippo signaling network. *Dev. Cell.* 2010; 18:300–308. [PubMed: 20159599]
30. Yu J, et al. Kibra functions as a tumor suppressor protein that regulates Hippo signaling in conjunction with Merlin and expanded. *Dev. Cell.* 2010; 18:288–299. [PubMed: 20159598]

31. Tyler DM, Baker NE. Expanded and Fat regulate growth and differentiation in the *Drosophila* eye through multiple signaling pathways. *Dev. Biol.* 2007; 305:187–201. [PubMed: 17359963]
32. Willecke M, et al. The Fat cadherin acts through the Hippo tumor-suppressor pathway to regulate tissue size. *Curr. Biol.* 2006; 16:2090–2100. [PubMed: 16996265]
33. Silva E, Tsatskis Y, Gardano L, Tapon N, McNeill H. The tumor-suppressor gene Fat controls tissue growth upstream of expanded in the Hippo signaling pathway. *Curr. Biol.* 2006; 16:2081–2089. [PubMed: 16996266]
34. Cho E, et al. Delineation of a Fat tumor suppressor pathway. *Nat. Genet.* 2006; 38:1142–1150. [PubMed: 16980976]
35. Bennett FC, Harvey KF. Fat cadherin modulates organ size in *Drosophila* via the Salvador/Warts/Hippo signaling pathway. *Curr. Biol.* 2006; 16:2101–2110. [PubMed: 17045801]
36. Matakatsu H, Blair SS. Separating the adhesive and signaling functions of the Fat and Dachshous protocadherins. *Development.* 2006; 133:2315–2324. [PubMed: 16687445]
37. Sopko R, et al. Phosphorylation of the tumor suppressor Fat is regulated by its ligand Dachshous and the kinase discs overgrown. *Curr. Biol.* 2009; 19:1112–1117. [PubMed: 19540118]
38. Feng Y, Irvine KD. Processing and phosphorylation of the Fat receptor. *Proc. Natl Acad. Sci. USA.* 2009; 106:11989–11994. [PubMed: 19574458]
39. Simon MA, Xu A, Ishikawa HO, Irvine KD. Modulation of Fat:Dachshous binding by the cadherin domain kinase Four-jointed. *Curr Biol.* 2010; 20:811–817. [PubMed: 20434335]
40. Rogulja D, Rauskolb C, Irvine KD. Morphogen control of wing growth through the Fat signaling pathway. *Dev. Cell.* 2008; 15:309–321. [PubMed: 18694569]
41. Willecke M, Hamaratoglu F, Sansores-Garcia L, Tao C, Halder G. Boundaries of Dachshous cadherin activity modulate the Hippo signaling pathway to induce cell proliferation. *Proc. Natl Acad. Sci. USA.* 2008; 105:14897–14902. [PubMed: 18809931]
42. Mao Y, Kucuk B, Irvine KD. *Drosophila* lowfat, a novel modulator of Fat signaling. *Development.* 2009; 136:3223–3233. [PubMed: 19710173]
43. Zecca M, Struhl G. A feed-forward circuit linking wingless, Fat–Dachshous signaling, and the Warts–Hippo pathway to *Drosophila* wing growth. *PLoS Biol.* 2010; 8:e1000386. [PubMed: 20532238]
44. Feng Y, Irvine KD. Fat and expanded act in parallel to regulate growth through warts. *Proc. Natl Acad. Sci. USA.* 2007; 104:20362–20367. [PubMed: 18077345]
45. Das Thakur M, et al. Ajuba LIM proteins are negative regulators of the Hippo signaling pathway. *Curr. Biol.* 2010; 20:657–662. [PubMed: 20303269]
46. Chen CL, et al. The apical-basal cell polarity determinant Crumbs regulates Hippo signaling in *Drosophila*. *Proc. Natl Acad. Sci. USA.* 2010; 107:15810–15815. [PubMed: 20798049]
47. Grzeschik NA, Parsons LM, Allott ML, Harvey KF, Richardson HE. Lgl, aPKC, and Crumbs regulate the Salvador/Warts/Hippo pathway through two distinct mechanisms. *Curr. Biol.* 2010; 20:573–581. [PubMed: 20362447]
48. Ling C, et al. The apical transmembrane protein Crumbs functions as a tumor suppressor that regulates Hippo signaling by binding to Expanded. *Proc. Natl Acad. Sci. USA.* 2010; 107:10532–10537. [PubMed: 20498073]
49. Robinson BS, Huang J, Hong Y, Moberg KH. Crumbs regulates Salvador/Warts/Hippo signaling in *Drosophila* via the FERM-domain protein expanded. *Curr. Biol.* 2010; 20:582–590. [PubMed: 20362445]
50. Tao W, et al. Human homologue of the *Drosophila melanogaster* lats tumour suppressor modulates CDC2 activity. *Nat. Genet.* 1999; 21:177–181. [PubMed: 9988268]
51. Graves JD, et al. Caspase-mediated activation and induction of apoptosis by the mammalian Ste20-like kinase Mst1. *EMBO J.* 1998; 17:2224–2234. [PubMed: 9545236]
52. Callus BA, Verhagen AM, Vaux DL. Association of mammalian sterile twenty kinases, Mst1 and Mst2, with hSalvador via C-terminal coiled-coil domains, leads to its stabilization and phosphorylation. *FEBS J.* 2006; 273:4264–4276. [PubMed: 16930133]
53. Lee JH, et al. A crucial role of WW45 in developing epithelial tissues in the mouse. *EMBO J.* 2008; 27:1231–1242. [PubMed: 18369314]

54. Oh HJ, et al. Role of the tumor suppressor RASSF1A in Mst1-mediated apoptosis. *Cancer Res.* 2006; 66:2562–2569. [PubMed: 16510573]
55. Khokhlatchev A, et al. Identification of a novel Ras-regulated proapoptotic pathway. *Curr. Biol.* 2002; 12:253–265. [PubMed: 11864565]
56. Polesello C, Huelsmann S, Brown NH, Tapon N. The *Drosophila* RASSF homolog antagonizes the Hippo pathway. *Curr. Biol.* 2006; 16:2459–2465. [PubMed: 17174922]
57. Chan EH, et al. The Ste20-like kinase Mst2 activates the human large tumor suppressor kinase Lats1. *Oncogene.* 2005; 24:2076–2086. [PubMed: 15688006]
58. Praskova M, Xia F, Avruch J. MOBKL1A/MOBKL1B phosphorylation by MST1 and MST2 inhibits cell proliferation. *Curr. Biol.* 2008; 18:311–321. [PubMed: 18328708]
59. Zhao B, et al. Inactivation of YAP oncoprotein by the Hippo pathway is involved in cell contact inhibition and tissue growth control. *Genes Dev.* 2007; 21:2747–2761. [PubMed: 17974916]
60. Hao Y, Chun A, Cheung K, Rashidi B, Yang X. Tumor suppressor LATS1 is a negative regulator of oncogene YAP. *J. Biol. Chem.* 2008; 283:5496–5509. [PubMed: 18158288]
61. Oka T, Mazack V, Sudol M. Mst2 and Lats kinases regulate apoptotic function of Yes kinase-associated protein (YAP). *J. Biol. Chem.* 2008; 283:27534–27546. [PubMed: 18640976]
62. Lei QY, et al. TAZ promotes cell proliferation and epithelial–mesenchymal transition and is inhibited by the Hippo pathway. *Mol. Cell Biol.* 2008; 28:2426–2436. [PubMed: 18227151]
63. Zender L, et al. Identification and validation of oncogenes in liver cancer using an integrative oncogenomic approach. *Cell.* 2006; 125:1253–1267. [PubMed: 16814713]
64. Overholtzer M, et al. Transforming properties of YAP, a candidate oncogene on the chromosome 11q22 amplicon. *Proc. Natl Acad. Sci. USA.* 2006; 103:12405–12410. [PubMed: 16894141]
65. Steinhardt AA, et al. Expression of Yes-associated protein in common solid tumors. *Hum. Pathol.* 2008; 39:1582–1589. [PubMed: 18703216]
66. Xu MZ, et al. Yes-associated protein is an independent prognostic marker in hepatocellular carcinoma. *Cancer.* 2009; 115:4576–4585. [PubMed: 19551889]
67. Chan SW, et al. A role for TAZ in migration, invasion, and tumorigenesis of breast cancer cells. *Cancer Res.* 2008; 68:2592–2598. [PubMed: 18413727]
68. Zhou Z, et al. TAZ is a novel oncogene in non-small cell lung cancer. *Oncogene.* 2011; 30:2181–2186. [PubMed: 21258416]
69. Zhang N, et al. The Merlin/NF2 tumor suppressor functions through the YAP oncoprotein to regulate tissue homeostasis in mammals. *Dev. Cell.* 2010; 19:27–38. [PubMed: 20643348]
70. Benhamouche S, et al. Nf2/Merlin controls progenitor homeostasis and tumorigenesis in the liver. *Genes Dev.* 2010; 24:1718–1730. [PubMed: 20675406]
71. Lee KP, et al. The Hippo-Salvador pathway restrains hepatic oval cell proliferation, liver size and liver tumorigenesis. *Proc. Natl Acad. Sci. USA.* 2010; 107:8248–8253. [PubMed: 20404163]
72. Lu L, et al. Hippo signaling is a potent *in vivo* growth and tumor suppressor pathway in the mammalian liver. *Proc. Natl Acad. Sci. USA.* 2010; 107:1437–1442. [PubMed: 20080689]
73. Song H, et al. Mammalian Mst1 and Mst2 kinases play essential roles in organ size control and tumor suppression. *Proc. Natl Acad. Sci. USA.* 2010; 107:1431–1436. [PubMed: 20080598]
74. Zhou D, et al. Mst1 and Mst2 maintain hepatocyte quiescence and suppress hepatocellular carcinoma development through inactivation of the Yap1 oncogene. *Cancer Cell.* 2009; 16:425–438. [PubMed: 19878874]
75. Zhao B, Li L, Tumaneng K, Wang CY, Guan KL. A coordinated phosphorylation by Lats and CK1 regulates YAP stability through SCF^β-TRCP. *Genes Dev.* 2010; 24:72–85. [PubMed: 20048001]
76. Oh H, Irvine KD. *In vivo* regulation of Yorkie phosphorylation and localization. *Development.* 2008; 135:1081–1088. [PubMed: 18256197]
77. Ren F, Zhang L, Jiang J. Hippo signaling regulates Yorkie nuclear localization and activity through 14-3-3 dependent and independent mechanisms. *Dev. Biol.* 2009; 337:303–312. [PubMed: 19900439]
78. Liu CY, et al. The Hippo tumor pathway promotes TAZ degradation by phosphorylating a phosphodegron and recruiting the SCF^β-TRCP E3 ligase. *J. Biol. Chem.* 2010; 285:37159–37169. [PubMed: 20858893]

79. Gilbert MM, Tipping M, Veraksa A, Moberg KH. A screen for conditional growth suppressor genes identifies the *Drosophila* homolog of HD-PTP as a regulator of the oncoprotein Yorkie. *Dev. Cell.* 2011; 20:700–712. [PubMed: 21571226]
80. Badouel C, et al. The FERM-domain protein Expanded regulates Hippo pathway activity via direct interactions with the transcriptional activator Yorkie. *Dev. Cell.* 2009; 16:411–420. [PubMed: 19289086]
81. Oh H, Reddy BV, Irvine KD. Phosphorylation-independent repression of Yorkie in Fat-Hippo signaling. *Dev. Biol.* 2009; 335:188–197. [PubMed: 19733165]
82. Zhao B, et al. Angiomotin is a novel Hippo pathway component that inhibits YAP oncoprotein. *Genes Dev.* 2001; 25:51–63. [PubMed: 21205866]
83. Wang W, Huang J, Chen J. Angiomotin-like proteins associate with and negatively regulate YAP1. *J. Biol. Chem.* 2010; 286:4364–4370. [PubMed: 21187284]
84. Chan SW, et al. Hippo pathway-independent restriction of TAZ and YAP by angiomotin. *J. Biol. Chem.* 2011; 286:7018–7026. [PubMed: 21224387]
85. Varelas X, et al. The crumbs complex couples cell density sensing to Hippo-dependent control of the TGF- β -SMAD pathway. *Dev. Cell.* 2010; 19:831–844. [PubMed: 21145499]
86. Oka T, et al. Functional complexes between YAP2 and ZO-2 are PDZ domain-dependent, and regulate YAP2 nuclear localization and signalling. *Biochem. J.* 2010; 432:461–472. [PubMed: 20868367]
87. Remue E, et al. TAZ interacts with zonula occludens-1 and -2 proteins in a PDZ-1 dependent manner. *FEBS Lett.* 2010; 584:4175–4180. [PubMed: 20850437]
88. Zhao B, et al. TEAD mediates YAP-dependent gene induction and growth control. *Genes Dev.* 2008; 22:1962–1971. [PubMed: 18579750]
89. Zhang L, et al. The TEAD/TEF family of transcription factor Scalloped mediates Hippo signaling in organ size control. *Dev. Cell.* 2008; 14:377–387. [PubMed: 18258485]
90. Zhang H, et al. TEAD transcription factors mediate the function of TAZ in cell growth and epithelial–mesenchymal transition. *J. Biol. Chem.* 2009; 284:13355–13362. [PubMed: 19324877]
91. Wu S, Liu Y, Zheng Y, Dong J, Pan D. The TEAD/TEF family protein Scalloped mediates transcriptional output of the Hippo growth-regulatory pathway. *Dev. Cell.* 2008; 14:388–398. [PubMed: 18258486]
92. Goulev Y, et al. SCALLOPED interacts with YORKIE, the nuclear effector of the Hippo tumor-suppressor pathway in *Drosophila*. *Curr. Biol.* 2008; 18:435–441. [PubMed: 18313299]
93. Li Z, et al. Structural insights into the YAP and TEAD complex. *Genes Dev.* 2010; 24:235–240. [PubMed: 20123905]
94. Kitagawa M. A Sveinsson’s chorioretinal atrophy-associated missense mutation in mouse Tead1 affects its interaction with the co-factors YAP and TAZ. *Biochem. Biophys. Res. Commun.* 2007; 361:1022–1026. [PubMed: 17689488]
95. Chen L, et al. Structural basis of YAP recognition by TEAD4 in the Hippo pathway. *Genes Dev.* 2010; 24:290–300. [PubMed: 20123908]
96. Fossdal R, et al. A novel TEAD1 mutation is the causative allele in Sveinsson’s chorioretinal atrophy (helicoid peripapillary chorioretinal degeneration). *Hum. Mol. Genet.* 2004; 13:975–981. [PubMed: 15016762]
97. Lai D, Ho KC, Hao Y, Yang X. Taxol resistance in breast cancer cells is mediated by the Hippo pathway component TAZ and its downstream transcriptional targets Cyr61 and CTGF. *Cancer Res.* 2011; 71:2728–2738. [PubMed: 21349946]
98. Neto-Silva RM, de Beco S, Johnston LA. Evidence for a growth-stabilizing regulatory feedback mechanism between Myc and Yorkie, the *Drosophila* homolog of Yap. *Dev. Cell.* 2010; 19:507–520. [PubMed: 20951343]
99. Ziosi M, et al. dMyc functions downstream of Yorkie to promote the supercompetitive behavior of Hippo pathway mutant cells. *PLoS Genet.* 2010; 6:e1001140. [PubMed: 20885789]
100. Yagi R, Chen LF, Shigesada K, Murakami Y, Ito Y. A WW domain-containing yes-associated protein (YAP) is a novel transcriptional co-activator. *EMBO J.* 1999; 18:2551–2562. [PubMed: 10228168]

101. Strano S, et al. Physical interaction with Yes-associated protein enhances p73 transcriptional activity. *J. Biol. Chem.* 2001; 276:15164–15173. [PubMed: 11278685]
102. Komuro A, Nagai M, Navin NE, Sudol M. WW domain-containing protein YAP associates with ErbB-4 and acts as a co-transcriptional activator for the carboxyl-terminal fragment of ErbB-4 that translocates to the nucleus. *J. Biol. Chem.* 2003; 278:33334–33341. [PubMed: 12807903]
103. Omerovic J, et al. Ligand-regulated association of ErbB-4 to the transcriptional co-activator YAP65 controls transcription at the nuclear level. *Exp. Cell Res.* 2004; 294:469–479. [PubMed: 15023535]
104. Alarcon C, et al. Nuclear CDKs drive Smad transcriptional activation and turnover in BMP and TGF- β pathways. *Cell.* 2009; 139:757–769. [PubMed: 19914168]
105. Hong JH, et al. TAZ, a transcriptional modulator of mesenchymal stem cell differentiation. *Science.* 2005; 309:1074–1078. [PubMed: 16099986]
106. Murakami M, Nakagawa M, Olson EN, Nakagawa O. A WW domain protein TAZ is a critical coactivator for TBX5, a transcription factor implicated in Holt-Oram syndrome. *Proc. Natl Acad. Sci. USA.* 2005; 102:18034–18039. [PubMed: 16332960]
107. Cui CB, Cooper LF, Yang X, Karsenty G, Aukhil I. Transcriptional coactivation of bone-specific transcription factor Cbfa1 by TAZ. *Mol. Cell Biol.* 2003; 23:1004–1013. [PubMed: 12529404]
108. Park KS, et al. TAZ interacts with TTF-1 and regulates expression of surfactant protein-C. *J. Biol. Chem.* 2004; 279:17384–17390. [PubMed: 14970209]
109. Murakami M, et al. Transcriptional activity of Pax3 is co-activated by TAZ. *Biochem. Biophys. Res. Commun.* 2006; 339:533–539. [PubMed: 16300735]
110. Varelas X, et al. TAZ controls Smad nucleocytoplasmic shuttling and regulates human embryonic stem-cell self-renewal. *Nat. Cell Biol.* 2008; 10:837–848. [PubMed: 18568018]
111. Lapi E, et al. PML, YAP, and p73 are components of a proapoptotic autoregulatory feedback loop. *Mol. Cell.* 2008; 32:803–814. [PubMed: 19111660]
112. Matallanas D, et al. RASSF1A elicits apoptosis through an MST2 pathway directing proapoptotic transcription by the p73 tumor suppressor protein. *Mol. Cell.* 2007; 27:962–975. [PubMed: 17889669]
113. Heallen T, et al. Hippo pathway inhibits Wnt signaling to restrain cardiomyocyte proliferation and heart size. *Science.* 2011; 332:458–461. [PubMed: 21512031]
114. Thompson BJ, Cohen SM. The Hippo pathway regulates the bantam microRNA to control cell proliferation and apoptosis in *Drosophila*. *Cell.* 2006; 126:767–774. [PubMed: 16923395]
115. Nolo R, Morrison CM, Tao C, Zhang X, Halder G. The bantam microRNA is a target of the Hippo tumor-suppressor pathway. *Curr. Biol.* 2006; 16:1895–1904. [PubMed: 16949821]
116. Peng HW, Slattery M, Mann RS. Transcription factor choice in the Hippo signaling pathway: homothorax and yorkie regulation of the microRNA bantam in the progenitor domain of the *Drosophila* eye imaginal disc. *Genes Dev.* 2009; 23:2307–2319. [PubMed: 19762509]
117. Oh H, Irvine KD. Cooperative regulation of growth by Yorkie and Mad through bantam. *Dev. Cell.* 2011; 20:109–122. [PubMed: 21238929]
118. Zhang J, et al. YAP-dependent induction of amphiregulin identifies a non-cell-autonomous component of the Hippo pathway. *Nat. Cell Biol.* 2009; 11:1444–1450. [PubMed: 19935651]
119. Genevet A, et al. The Hippo pathway regulates apical-domain size independently of its growth-control function. *J. Cell Sci.* 2009; 122:2360–2370. [PubMed: 19531586]
120. Skouloudaki K, et al. Scribble participates in Hippo signaling and is required for normal zebrafish pronephros development. *Proc. Natl Acad. Sci. USA.* 2009; 106:8579–8584. [PubMed: 19439659]
121. Zhan L, et al. Dereglulation of scribble promotes mammary tumorigenesis and reveals a role for cell polarity in carcinoma. *Cell.* 2008; 135:865–878. [PubMed: 19041750]
122. Hong Y, Stronach B, Perrimon N, Jan LY, Jan YN. *Drosophila* Stardust interacts with Crumbs to control polarity of epithelia but not neuroblasts. *Nature.* 2001; 414:634–638. [PubMed: 11740559]

123. Grzeschik NA, Amin N, Secombe J, Brumby AM, Richardson HE. Abnormalities in cell proliferation and apico-basal cell polarity are separable in *Drosophila lgl* mutant clones in the developing eye. *Dev. Biol.* 2007; 311:106–123. [PubMed: 17870065]
124. Nishioka N, et al. The Hippo signaling pathway components Lats and Yap pattern Tead4 activity to distinguish mouse trophectoderm from inner cell mass. *Dev. Cell.* 2009; 16:398–410. [PubMed: 19289085]
125. Silvis MR, et al. α -catenin is a tumor suppressor that controls cell accumulation by regulating the localization and activity of the transcriptional coactivator Yap1. *Sci. Signal.* 2011; 4:ra33. [PubMed: 21610251]
126. Schlegelmilch K, et al. Yap1 acts downstream of α -catenin to control epidermal proliferation. *Cell.* 2011; 144:782–795. [PubMed: 21376238]
127. Zhang H, Pasolli HA, Fuchs E. Yes-associated protein (YAP) transcriptional coactivator functions in balancing growth and differentiation in skin. *Proc. Natl Acad. Sci. USA.* 2011; 108:2270–2275. [PubMed: 21262812]
128. Fernandez LA, et al. YAP1 is amplified and up-regulated in hedgehog-associated medulloblastomas and mediates Sonic hedgehog-driven neural precursor proliferation. *Genes Dev.* 2009; 23:2729–2741. [PubMed: 19952108]
129. Sun G, Irvine KD. Regulation of Hippo signaling by Jun kinase signaling during compensatory cell proliferation and regeneration, and in neoplastic tumors. *Dev. Biol.* 2011; 350:139–151. [PubMed: 21145886]
130. Furuyama K, et al. Continuous cell supply from a Sox9-expressing progenitor zone in adult liver, exocrine pancreas and intestine. *Nat. Genet.* 2010; 43:34–41. [PubMed: 21113154]




Figure 1. The Hippo pathway in *Drosophila* and mammals. Corresponding proteins in *Drosophila* (a) and mammals (b) are indicated by matching colours. Arrowed or blunted ends indicate activation or inhibition, respectively. Dashed lines indicate unknown mechanisms.




Figure 2. Mechanisms of YAP/TAZ/Yki inhibition by the Hippo pathway. **(a)** Phosphorylation-dependent cytoplasmic retention. Phosphorylation of YAP on Ser 127 by Lats1/2 induces 14-3-3 binding and cytoplasmic retention of YAP. The mechanism is conserved in TAZ and Yki. **(b)** Phosphorylation-independent cytoplasmic retention. Through WW domain–PPXY motif interactions, Yki binds to Mop, Ex, Hpo and Wts, and YAP/TAZ binds to AMOT family proteins. These interactions physically sequester Yki and YAP/TAZ in the cytoplasm. **(c)** Phosphorylation-induced ubiquitylation and degradation. Phosphorylation of

YAP on Ser 381 by Lats1/2 primes further phosphorylation of YAP by CK1 δ/ϵ , which induces interaction with SCF β -TRCP and finally leads to YAP ubiquitylation and degradation. The mechanism is conserved in TAZ.




Figure 3. Mechanisms of the Hippo pathway in regulation of organ size and regeneration. Hexagons denote differentiated cells and circles denote stem/progenitor cells. Blue colour indicates wild-type and yellow colour indicates Hippo-pathway mutant cells. **(a)** Hippo pathway inactivation leads to stem/progenitor cell expansion in both cell-autonomous and non-autonomous manners. **(b)** Hippo pathway inactivation leads to cell cycle exit defects in some cellular contexts. **(c)** Hippo pathway mutations promote proliferation and decrease

apoptosis in non-stem/progenitor cells. **(d)** Imbalance of Hippo pathway activity in neighbouring cells induces cell competition.