

# THE IDEMPOTENT-SEPARATING CONGRUENCES ON A REGULAR 0-BISIMPLE SEMIGROUP

by W. D. MUNN

(Received 1st September 1966)

A congruence  $\rho$  on a semigroup is said to be idempotent-separating if each  $\rho$ -class contains at most one idempotent. For any idempotent  $e$  of a semigroup  $S$  the set  $eSe$  is a subsemigroup of  $S$  with identity  $e$  and group of units  $H_e$ , the maximal subgroup of  $S$  containing  $e$ . The purpose of the present note is to show that if  $S$  is a regular 0-bisimple semigroup and  $e$  is a non-zero idempotent of  $S$  then there is a one-to-one correspondence between the idempotent-separating congruences on  $S$  and the subgroups  $N$  of  $H_e$  with the property that  $aN \subseteq Na$  for all right units  $a$  of  $eSe$  and  $Nb \subseteq bN$  for all left units  $b$  of  $eSe$ . Some special cases of this result are discussed and, in the final section, an application is made to the principal factors of the full transformation semigroup  $\mathcal{T}_X$  on a set  $X$ .

1. The notation of (1) will be used throughout. In particular, if  $\rho$  is an equivalence on a set  $X$  then  $x\rho$  denotes the  $\rho$ -class containing the element  $x$  of  $X$ . As in (1), an exception is made in the case of Green's equivalences  $\mathcal{R}$ ,  $\mathcal{L}$  and  $\mathcal{H}$  on a semigroup  $S$ : the corresponding classes containing the element  $a$  of  $S$  are denoted by  $R_a$ ,  $L_a$  and  $H_a$ .

Let  $S$  contain an idempotent  $e$ . Then  $eSe$  is a subsemigroup of  $S$  and we write

$$S_e = eSe, P_e = R_e \cap S_e, Q_e = L_e \cap S_e.$$

Since  $e$  is a left identity for  $R_e$  and a right identity for  $L_e$  we see that

$$P_e = \{x \in R_e : xe = x\}, Q_e = \{x \in L_e : ex = x\}.$$

In the first two lemmas we establish some basic properties of these sets.

**Lemma 1.** *Let  $e$  and  $f$  be  $\mathcal{D}$ -equivalent idempotents of a semigroup  $S$ . Then  $S_e \cong S_f$ .*

**Proof.** Since  $(e, f) \in \mathcal{D}$  there exists an element  $a$  in  $R_e \cap L_f$ . By ((1), Theorem 2.18), there is a unique element  $a'$  in  $R_f \cap L_e$  such that  $aa' = e$  and  $a'a = f$ . Let  $x \in S_e$ . Then

$$a'xa = a'exea = fa'xaf \in S_f.$$

Similarly, if  $y \in S_f$  then  $aya' \in S_e$ . Now let  $\theta: S_e \rightarrow S_f$  and  $\phi: S_f \rightarrow S_e$  be defined by  $x\theta = a'xa$  ( $x \in S_e$ ) and  $y\phi = aya'$  ( $y \in S_f$ ). Then  $x\theta\phi = exe = x$  for all  $x \in S_e$ . Similarly,  $y\phi\theta = y$  for all  $y \in S_f$ . Hence  $\theta$  and  $\phi$  are mutually

E.M.S.—Q

inverse bijections. Finally,  $a'xya = (a'xa)(a'ya)$  for all  $x, y \in S_e$  and so  $\theta$  is an isomorphism.

In particular, if  $S$  is a regular 0-bisimple semigroup then  $S_e \cong S_f$  for any non-zero idempotents  $e, f$ . In this case it can also be shown that  $S_e$  is 0-bisimple.

Let  $T$  be a semigroup with an identity  $e$ . By a *right unit* of  $T$  we mean an element  $a$  of  $T$  such that  $ax = e$  for some  $x$  in  $T$ . The set of all right units of  $T$  is readily seen to be a right cancellative subsemigroup of  $T$  (called the *right unit subsemigroup of  $T$* ). Left units are defined in a similar way; the set of all such elements is a left cancellative subsemigroup of  $T$  (the *left unit subsemigroup of  $T$* ). The elements of  $T$  that are both right units and left units are called *units*.

**Lemma 2.** *Let  $e$  be an idempotent of a semigroup  $S$ . Then  $P_e$  is the right unit subsemigroup of  $S_e$  and  $Q_e$  is the left unit subsemigroup of  $S_e$ . The group of units of  $S_e$  is  $H_e$ .*

**Proof.** Let  $a \in P_e$ . Then there exists  $x \in S^1$  such that  $ax = e$ . Write  $y = exe$ . Since  $ae = a$  we have that  $ay = axe = e^2 = e$ . Hence  $a$  is a right unit of  $S_e$ . Conversely, let  $a$  be a right unit of  $S_e$ . Then there exists  $y \in S_e$  such that  $ay = e$ . Also  $ea = a$  and so  $a \in R_e \cap S_e = P_e$ . Thus  $P_e$  is the right unit subsemigroup of  $S_e$ . Similarly,  $Q_e$  is the left unit subsemigroup of  $S_e$ .

Finally,  $P_e \cap Q_e = R_e \cap L_e \cap S_e = H_e$ .

2. By a *left normal divisor of  $P_e$*  we shall mean a subgroup  $N$  of  $H_e$  such that  $aN \subseteq Na$  for all  $a \in P_e$ . Since  $H_e \subseteq P_e$  it is clear that, in particular, a left normal divisor of  $P_e$  is a normal subgroup of  $H_e$ . Similarly, by a *right normal divisor of  $Q_e$*  we mean a subgroup  $N$  of  $H_e$  such that  $Nb \subseteq bN$  for all  $b \in Q_e$ . This terminology is due to Rees (5). In the next lemma we establish a connection between such subgroups of  $H_e$  and the congruences on  $S$  contained in  $\mathcal{H}$ .

**Lemma 3.** *Let  $e$  be an idempotent of a semigroup  $S$ . Let  $\rho$  be any congruence on  $S$  contained in  $\mathcal{H}$  and let  $N = e\rho$ . Then  $N$  is a left normal divisor of  $P_e$  and a right normal divisor of  $Q_e$ .*

**Proof.** Since  $\rho \cap (H_e \times H_e)$  is a congruence on  $H_e$  and  $\rho \subseteq \mathcal{H}$  it follows that  $N$  is a normal subgroup of  $H_e$ . Let  $a \in P_e$  and let  $b \in a\rho$ . Then since  $\rho \subseteq \mathcal{H}$  there exists  $x' \in S^1$  such that  $x'a = b$ . Write  $x = x'e$ ; then  $xa = x'ea = x'a = b$  and  $xe = x$ . Also since  $a\rho \subseteq H_a \subseteq R_e$  there exists  $y \in S^1$  such that  $by = e$ . Now  $(xby, xay) \in \rho$  since  $\rho$  is a congruence on  $S$ ; that is,  $(x, e) \in \rho$ . Thus  $x \in N$ . Hence, since  $b = xa$ , we have that  $a\rho \subseteq Na$ .

Next let  $z \in N$ . Then  $(az, ae) \in \rho$ . But  $ae = a$ ; hence  $aN \subseteq a\rho$ . Combining these results we see that  $aN \subseteq Na$ .

In the same way we can show that  $Nb \subseteq bN$  for all  $b \in Q_e$ .

**Corollary.** *If  $\mathcal{H}$  is a congruence on  $S$  then  $H_e$  is a left normal divisor of  $P_e$  and a right normal divisor of  $Q_e$ .*

Does every subgroup of  $H_e$  that is both a left normal divisor of  $P_e$  and a right normal divisor of  $Q_e$  arise, as in Lemma 3, from a congruence contained in  $\mathcal{H}$ ? This is answered for regular 0-bisimple semigroups in Lemma 6. As

a first step we establish

**Lemma 4.** *Let  $e$  be an idempotent of a semigroup  $S$ .*

(i) *Let  $a, b \in R_e$ . Then*

$$(a, b) \in \mathcal{H} \Leftrightarrow xa = b \text{ for some } x \in H_e.$$

(ii) *Let  $N$  be a left normal divisor of  $P_e$ . Define a relation  $\rho_R$  on  $R_e$  by the rule that*

$$(a, b) \in \rho_R \Leftrightarrow xa = b \text{ for some } x \in N.$$

*Then  $\rho_R$  is an equivalence on  $R_e$  contained in  $\mathcal{H}$ . Further, if  $(a, b) \in \rho_R$  then  $(ca, cb) \in \rho_R$  for all  $c \in P_e$ .*

**Proof.** (i) Since  $(e, a) \in \mathcal{R}$  we have  $ea = a$  and so  $H_e a = H_a$  ((1), Lemma 2.2). This gives the required result.

(ii) That  $\rho_R$  is an equivalence on  $R_e$  follows from the fact that  $N$  is a group whose identity  $e$  is a left identity for  $R_e$ . From (i) we see that  $\rho_R \subseteq \mathcal{H}$ .

Let  $(a, b) \in \rho_R$  and let  $c \in P_e$ . Since  $a \in R_e$  there exists  $z \in S^1$  such that  $az = e$ . Hence  $caz = ce = c$  and so  $(ca, c) \in \mathcal{R}$ ; that is,  $ca \in R_e$ . Similarly,  $cb \in R_e$ . Now  $xa = b$  for some  $x \in N$  and  $cN \subseteq Nc$ , by hypothesis. Hence

$$cb = cxa = yca$$

for some  $y \in N$ , which shows that  $(ca, cb) \in \rho_R$ .

Dually, for any right normal divisor  $N$  of  $Q_e$  we define a relation  $\rho_L$  on  $L_e$  by the rule that

$$(a, b) \in \rho_L \Leftrightarrow ax = b \text{ for some } x \in N.$$

Then  $\rho_L$  is an equivalence on  $L_e$  contained in  $\mathcal{H}$  and if  $(a, b) \in \rho_L$  then  $(ac, bc) \in \rho_L$  for all  $c \in Q_e$ .

**3.** In this section we restrict our attention to 0-bisimple semigroups. By ((1), Theorem 2.11) such a semigroup is regular if and only if it contains a non-zero idempotent.

**Lemma 5.** *Let  $S$  be a 0-bisimple semigroup and let  $a$  be an arbitrary but fixed non-zero element of  $S$ . Let  $\rho, \tau$  be congruences on  $S$  contained in  $\mathcal{H}$ . Then*

(i)  $\rho = \mathcal{H}$  if and only if  $a\rho = H_a$ ;

(ii)  $\rho \subseteq \tau$  if and only if  $a\rho \subseteq a\tau$ .

**Proof.** (i) Let  $a\rho = H_a$ . To establish (i) we need only prove that  $\mathcal{H} \subseteq \rho$ . First,  $H_0 = 0\rho = 0$ . Now let  $(b, c) \in \mathcal{H}$ , where  $b \neq 0, c \neq 0$ . Since  $S$  is 0-bisimple there exist elements  $s, s', t, t'$  in  $S^1$  such that  $b = sat, a = s'bt'$  and the mappings

$$x \rightarrow sxt \ (x \in H_a), \quad y \rightarrow s'yt' \ (y \in H_b)$$

are mutually inverse bijections from  $H_a$  to  $H_b$  and from  $H_b$  to  $H_a$  respectively ((1), Theorem 2.3). Thus, since  $(b, c) \in \mathcal{H}$ , we see that  $(s'bt', s'ct') \in \mathcal{H}$ . Since  $a = s'bt'$  and  $a\rho = H_a$  it follows that  $(a, s'ct') \in \rho$ . But  $\rho$  is a congruence on  $S$  and so  $(sat, ss'ct't) \in \rho$ ; that is,  $(b, c) \in \rho$ . Thus  $\mathcal{H} \subseteq \rho$ .

(ii) It is clear that if  $\rho \subseteq \tau$  then  $a\rho \subseteq a\tau$ . Suppose, conversely, that  $a\rho \subseteq a\tau$ . Let  $(b, c) \in \rho$ , where  $b \neq 0, c \neq 0$ . To prove (ii) it suffices to show that  $(b, c) \in \tau$ . As above, since  $(a, b) \in \mathcal{D}$  there exist elements  $s, s', t, t'$  in  $S^1$  such that  $b = sat, a = s'bt'$  and the mappings  $x \rightarrow sxt (x \in H_a), y \rightarrow s'yt' (y \in H_b)$  are mutually inverse bijections from  $H_a$  to  $H_b$  and from  $H_b$  to  $H_a$  respectively. Since  $\rho$  is a congruence,  $(s'bt', s'ct') \in \rho$ ; that is,  $(a, s'ct') \in \rho$ . But  $a\rho \subseteq a\tau$  and so  $(a, s'ct') \in \tau$ . Hence  $(sat, ss'ct't) \in \tau$  since  $\tau$  is a congruence. But  $(b, c) \in \mathcal{H}$  since  $\rho \subseteq \mathcal{H}$ . Thus  $ss'ct't = c$ . It follows that  $(b, c) \in \tau$ , as required.

In particular, from (ii) above,  $\rho = \tau$  if and only if  $a\rho = a\tau$  for any non-zero element  $a$  of  $S$ ; that is, a congruence contained in  $\mathcal{H}$  on a 0-bisimple semigroup is uniquely determined by any one of its non-zero classes.

We now come to the key result.

**Lemma 6.** *Let  $S$  be a regular 0-bisimple semigroup and let  $e$  be a non-zero idempotent of  $S$ . Let  $N$  be a subgroup of  $H_e$  that is both a left normal divisor of  $P_e$  and a right normal divisor of  $Q_e$ . Then there exists a congruence  $\rho$  on  $S$  contained in  $\mathcal{H}$  and such that  $e\rho = N$ .*

**Proof.** We construct  $\rho$  by defining  $\rho \cap (H \times H)$  for each non-zero  $\mathcal{H}$ -class  $H$  in terms of the equivalence  $\rho_R$  on  $R_e$  described in Lemma 4 (ii). The argument depends on several applications of the dual of ((1), Lemma 2.2) which we shall refer to below as Green's lemma.

Let  $(a, b) \in \mathcal{H}$ , where  $a \neq 0, b \neq 0$ . Since  $S$  is 0-bisimple there exists an element  $s \in S^1$  such that  $sa \in R_e \cap L_a$ . Then  $sb \in R_e \cap L_a$  by Green's lemma. Now let  $(sa, sb) \in \rho_R$ . We prove first that  $(za, zb) \in \rho_R$  for any  $z \in S^1$  such that  $za \in R_e \cap L_a$ . By Lemma 4 (i), since  $(sa, za) \in \mathcal{H}$ , there exists  $x \in H_e$  such that  $xsa = za$ . Also since  $(a, b) \in \mathcal{H}$ , there exists  $y \in S^1$  such that  $ay = b$ . Now  $(xsa, xsb) \in \rho_R$  by Lemma 4 (ii). But

$$xsb = xsay = zay = zb.$$

Hence  $(za, zb) \in \rho_R$ .

Let  $H$  be a non-zero  $\mathcal{H}$ -class of  $S$ . We define a relation  $\rho_H$  on  $H$  by the rule that

$$(a, b) \in \rho_H \Leftrightarrow (sa, sb) \in \rho_R, \quad (a, b \in H) \tag{1}$$

where  $s$  is any element of  $S^1$  such that  $sa \in R_e \cap L_a$ . Clearly  $\rho_H$  is reflexive and symmetric. To see that it is transitive, let  $(a, b) \in \rho_H$  and  $(b, c) \in \rho_H$ . Then there exists  $s \in S^1$  such that  $sa \in R_e \cap L_a$  and  $(sa, sb) \in \rho_R$ . Since  $sb \in R_e \cap L_a$  it follows that  $(sb, sc) \in \rho_R$  and so  $(sa, sc) \in \rho_R$  since  $\rho_R$  is transitive. Thus  $\rho_H$  is an equivalence on  $H$ .

The definition in (1) lacks left-right symmetry. We shall now show that we would arrive at the same equivalence on  $H$  by using the congruence  $\rho_L$  on  $L_e$  defined in the dual form of Lemma 4 (ii).

Again, let  $(a, b) \in \mathcal{H}$ , where  $a \neq 0$  and  $b \neq 0$ , let  $s \in S^1$  be such that  $sa \in R_e \cap L_a$  and let  $(sa, sb) \in \rho_R$ . Since  $S$  is 0-bisimple there exists  $t \in S^1$  such that  $at \in R_a \cap L_e$ . Then  $bt \in R_a \cap L_e$ . It will be sufficient to show that  $(at, bt) \in \rho_L$ . By the definition of  $\rho_R$  there exists  $g \in N$  such that  $gsa = sb$ .

Hence  $gsat = sbt$ . Now, by Green's lemma, the mapping

$$x \rightarrow sx \quad (x \in R_a)$$

is an  $\mathcal{L}$ -class-preserving bijection from  $R_a$  to  $R_e$  and so, since  $at \in R_a \cap L_e$ , it follows that  $sat \in R_e \cap L_e = H_e$ . Hence, since  $N$  is normal in  $H_e$ , there exists  $h \in N$  such that  $g(sat) = (sat)h$ . Thus

$$sath = sbt. \tag{2}$$

Now since  $(a, sa) \in \mathcal{L}$  there exists  $s' \in S^1$  such that  $s'sa = a$ . But  $(a, b) \in \mathcal{H}$  and so  $s'sb = b$ , by Green's lemma. Premultiplying both sides of (2) by  $s'$  we find that  $ath = bt$ . Hence  $(at, bt) \in \rho_L$ .

Next we define  $\rho \subseteq S \times S$  to be  $\rho^* \cup \{(0, 0)\}$ , where  $\rho^*$  is the union of all the subsets  $\rho_H$  of  $S \times S$  as  $H$  runs through all the non-zero  $\mathcal{H}$ -classes of  $S$ . Since  $\rho_H$  is an equivalence on  $H$  for each  $H$ , it follows that  $\rho$  is an equivalence on  $S$ . Moreover,  $\rho \subseteq \mathcal{H}$ . We prove that  $\rho$  is a congruence on  $S$ .

Let  $(a, b) \in \rho$  and let  $c \in S$ . It will be shown that  $(ca, cb) \in \rho$ . First suppose that  $ca = 0$ . Since  $\rho \subseteq \mathcal{H}$  there exists  $x \in S^1$  such that  $ax = b$ . Then  $cb = cax = 0$ . Hence  $(ca, cb) \in \rho$ . We therefore assume that  $ca \neq 0$  and  $cb \neq 0$ . Since  $S$  is 0-bisimple and  $a \neq 0$  there exist  $s, s' \in S^1$  such that  $sa \in R_e \cap L_a$  and  $s'sa = a$ . Then, as before,  $s'sb = b$ . From the definition of  $\rho$  we have that  $(sa, sb) \in \rho_R$ . Now  $esa = sa$  since  $sa \in R_e$ ; therefore

$$ca = cs'sa = cs'esa.$$

But  $ca \neq 0$ . Hence  $cs'e \neq 0$  and so there exist elements  $u, u' \in S^1$  such that

$$ucs'e \in R_e \cap L_{cs'e}, \quad u'ucs'e = cs'e.$$

Further,  $ucs'e \in P_e$  since  $(ucs'e)e = ucs'e$ . Then, applying Lemma 4 (ii), we find that

$$(ucs'e \cdot sa, ucs'e \cdot sb) \in \rho_R.$$

But  $ucs'esa = uca$  and  $ucs'esb = ucb$ ; thus

$$(uca, ucb) \in \rho_R. \tag{3}$$

Now

$$u'uca = u'ucs'esa = cs'esa = ca \tag{4}$$

and, similarly,  $u'ucb = cb$ . From (4),  $(uca, ca) \in \mathcal{L}$  and so, by Green's lemma, the mapping

$$x \rightarrow u'x \quad (x \in H_{uca})$$

is a bijection from  $H_{uca}$  to  $H_{ca}$ . Since  $\rho_R \subseteq \mathcal{H}$  we deduce from (3) that

$$(ca, cb) \in \mathcal{H}.$$

But  $u(ca) \in R_e \cap L_{ca}$ . It then follows from (3) that  $(ca, cb) \in \rho$ .

In the same way, using the alternative definition of  $\rho$  in terms of the equivalence  $\rho_L$  on  $L_e$ , we can show that  $(ac, bc) \in \rho$ . Thus  $\rho$  is a congruence on  $S$ .

Finally, let  $y \in H_e$ . Then  $(e, y) \in \rho$  if and only if  $(se, sy) \in \rho_R$  for any  $s \in S^1$  such that  $se \in R_e \cap L_e$ . In particular, taking  $s = e$ , we see that  $(e, y) \in \rho$  if and only if  $(e, y) \in \rho_R$ . But  $(e, y) \in \rho_R$  if and only if  $xe = y$  for some  $x \in N$ . Hence  $e\rho = N$ . This completes the proof.

**Corollary.** *Let  $S$  be a regular 0-bisimple semigroup and let  $e$  be any non-zero idempotent of  $S$ . Let  $H_e$  be a left normal divisor of  $P_e$  and a right normal divisor of  $Q_e$ . Then  $\mathcal{H}$  is a congruence on  $S$ .*

**Proof.** Take  $N = H_e$  in Lemma 6. Then there exists a congruence  $\rho$  on  $S$  contained in  $\mathcal{H}$  and such that  $e\rho = H_e$ . Then  $\rho = \mathcal{H}$  by Lemma 5 (i).

4. Lallement ((2), Theorem 2.3) has shown that the idempotent-separating congruences on a regular semigroup can be characterised as the congruences contained in  $\mathcal{H}$ . From Lemmas 3, 5 and 6 and the corollaries to Lemmas 3 and 6 we then obtain the following theorem concerning the idempotent-separating congruences on a regular 0-bisimple semigroup.

**Theorem.** *Let  $S$  be a regular 0-bisimple semigroup and let  $e$  be a non-zero idempotent of  $S$ . Let  $\Lambda$  denote the set of all idempotent-separating congruences on  $S$  and let  $\Delta$  denote the set of all subgroups of  $H_e$  that are left normal divisors of  $P_e$  and right normal divisors of  $Q_e$ . Then*

- (i)  $e\rho \in \Delta$  for all  $\rho \in \Lambda$ ;
- (ii)  $\rho \subseteq \tau$  if and only if  $e\rho \subseteq e\tau$  ( $\rho, \tau \in \Lambda$ );
- (iii) to each  $N$  in  $\Delta$  there corresponds  $\rho$  in  $\Lambda$  such that  $e\rho = N$ .

Furthermore,  $\mathcal{H}$  is a congruence on  $S$  if and only if  $H_e \in \Delta$ .

From ((2), Corollary 3.3) we see that  $\Lambda$  is a complete modular lattice. The greatest element  $\mu$  of  $\Lambda$  is the greatest congruence contained in  $\mathcal{H}$  and is characterised thus ((4), Lemma 1):

$$(a, b) \in \mu \Leftrightarrow (sat, sbt) \in \mathcal{H} \text{ for all } s, t \in S^1.$$

Let  $\Delta$  be partially ordered by inclusion. Then the theorem shows that

$$\rho \rightarrow e\rho$$

is an order-preserving bijection from  $\Lambda$  to  $\Delta$  whose inverse is also order-preserving. Hence  $\Delta$  is a complete modular lattice and  $\Delta \cong \Lambda$ . A direct calculation establishes that  $\Delta$  is a sublattice of the lattice of all normal subgroups of  $H_e$ .

It should also be noted that the theorem provides a description of the idempotent-separating congruences on a regular bisimple semigroup  $T$ ; for  $\rho \rightarrow \rho \cup \{(0, 0)\}$  is a bijection from the set of all such congruences on  $T$  to the set of all idempotent-separating congruences on the regular 0-bisimple semigroup  $T^0$ .

5. We now discuss two important classes of regular 0-bisimple semigroups.

First let  $S$  be a completely 0-simple semigroup. By ((1), Theorem 2.51),  $S$  is both regular and 0-bisimple. Let  $e$  be a non-zero idempotent of  $S$ . Then  $e$  is primitive and so  $S_e = H_e^0$  ((1), Lemma 2.47). Thus  $P_e = Q_e = H_e$ . The set  $\Delta$  in the theorem therefore consists of all normal subgroups of  $H_e$  and so there is a natural one-to-one correspondence between the idempotent-separating congruences on  $S$  and the normal subgroups of  $H_e$ . In particular,  $\mathcal{H}$  is a

congruence on  $S$ . These well-known results also follow immediately from the structure theorem for completely 0-simple semigroups ((1), Theorem 3.5).

Next, let  $S$  be a 0-bisimple inverse semigroup and let  $e$  be any non-zero idempotent in  $S$ . Let  $x \in S_e$ . Then,  $x = exe$  and so

$$x^{-1} = e^{-1}x^{-1}e^{-1} = ex^{-1}e \in S_e.$$

This shows that  $S_e$  is an inverse subsemigroup of  $S$ . Hence, by Lemma 2,  $Q_e$  consists of the inverses of the elements of  $P_e$ . Let  $N$  be a left normal divisor of  $P_e$ ; that is,  $aN \subseteq Na$  for all  $a \in P_e$ . Then  $Na^{-1} \subseteq a^{-1}N$  for all  $a \in P_e$  and so  $Nb \subseteq bN$  for all  $b \in Q_e$ . The set  $\Delta$  can therefore be taken as the set of all left normal divisors of  $P_e$ . For a bisimple inverse semigroup the theorem has been given in this form by Reilly and Clifford ((6), Theorem 2.4).

We deduce, in particular, that the idempotent-separating congruences on a bisimple inverse semigroup  $S$  with an identity are in one-to-one correspondence with the left normal divisors of the right unit subsemigroup of  $S$ . This result is due to Warne (7).

6. To conclude, we give an application of the theorem to the principal factors of the full transformation semigroup  $\mathcal{T}_X$  on a set  $X$ . It is easy to see that  $\mathcal{T}_X$  is regular ((1), p. 33, Exercise 1). The further properties required for our discussion—and outlined below—are established in ((1), § 2.2). We remark that Mal'cev (3) has determined a set of generators for the lattice of congruences on  $\mathcal{T}_X$ .

For  $\alpha \in \mathcal{T}_X$  the equivalence  $\alpha \circ \alpha^{-1}$  on  $X$  will be denoted by  $\pi_\alpha$ ; the cardinal of a set  $A$  will be denoted by  $|A|$ . Then the relations  $\mathcal{R}$ ,  $\mathcal{L}$  and  $\mathcal{D}$  on  $\mathcal{T}_X$  are characterised as follows:

$$\begin{aligned} (\alpha, \beta) \in \mathcal{R} &\Leftrightarrow \pi_\alpha = \pi_\beta, \\ (\alpha, \beta) \in \mathcal{L} &\Leftrightarrow X\alpha = X\beta, \\ (\alpha, \beta) \in \mathcal{D} &\Leftrightarrow |X\alpha| = |X\beta|. \end{aligned}$$

It is also easily verified that if  $\alpha, \varepsilon \in \mathcal{T}_X$  and  $\varepsilon^2 = \varepsilon$  then

$$\alpha\varepsilon = \alpha \Leftrightarrow X\alpha \subseteq X\varepsilon. \tag{1}$$

Now let  $|X| > 1$ . The principal factors of  $\mathcal{T}_X$  other than the kernel are of the form  $U_c/V_c$  where  $c$  is any cardinal such that  $|X| \geq c > 1$  and  $U_c, V_c$  are the ideals of  $\mathcal{T}_X$  defined by

$$U_c = \{\alpha \in \mathcal{T}_X : |X\alpha| \leq c\}, \quad V_c = \{\alpha \in \mathcal{T}_X : |X\alpha| < c\}.$$

We write  $T_c = U_c/V_c$ . Let  $\alpha$  be any element of  $\mathcal{T}_X$  of rank  $c$ . Then it can readily be shown that, since  $\mathcal{T}_X$  is regular, the  $\mathcal{R}$ -class  $R_\alpha$  of  $\mathcal{T}_X$  is also an  $\mathcal{R}$ -class of  $T_c$ ; similarly, the  $\mathcal{L}$ -class  $L_\alpha$  of  $\mathcal{T}_X$  is an  $\mathcal{L}$ -class of  $T_c$ . Hence  $T_c$  is a regular 0-bisimple semigroup. Moreover, by (1), for any non-zero idempotent  $\varepsilon$  of  $T_c$  we have that

$$P_\varepsilon = \{\alpha \in T_c \setminus 0 : \pi_\alpha = \pi_\varepsilon \text{ and } X\alpha \subseteq X\varepsilon\}. \tag{2}$$

Consider first the case where  $c$  is finite. Let  $\varepsilon, \eta$  be non-zero idempotents of

$T_c$  such that  $\varepsilon\eta = \eta = \eta\varepsilon$ . Since  $\eta = \eta\varepsilon$  it follows from (1) that  $X\eta \subseteq X\varepsilon$ . Thus  $X\eta = X\varepsilon$  since  $|X\eta| = |X\varepsilon| = c$ . Hence  $\varepsilon = \varepsilon\eta$  and so  $\varepsilon = \eta$ . This shows that  $T_c$  is completely 0-simple. Since  $H_c$  is isomorphic to the symmetric group of degree  $c$ , we see that, for  $c \geq 5$ ,  $T_c$  has exactly three distinct idempotent-separating congruences (corresponding to the three distinct normal subgroups of  $H_c$ ). Note that one of these congruences is  $\mathcal{H}$ .

Next let  $X$  be infinite and let  $c$  be an infinite cardinal such that  $|X| \geq c$ . We shall show that the only idempotent-separating congruence on  $T_c$  is the identity congruence. Let  $\varepsilon$  be a non-zero idempotent of  $T_c$  and let  $\gamma$  be an element of  $H_c$  distinct from  $\varepsilon$ . Then there exists  $y \in X$  such that  $y\gamma \neq y\varepsilon$ . Now  $\gamma \in L_\varepsilon$  and so  $y\gamma \in X\varepsilon$ . Since  $X\varepsilon$  is infinite there exists an element  $\theta$  in  $\mathcal{T}_X$  that induces a one-to-one mapping of  $X\varepsilon$  into  $X\varepsilon$  and is such that  $y\varepsilon \in (X\varepsilon)\theta$  and  $y\gamma \notin (X\varepsilon)\theta$ . Write  $\alpha = \varepsilon\theta$ . Then  $\alpha \in T_c \setminus \{0\}$ ; also  $\pi_\alpha = \pi_\varepsilon$  and  $X\alpha \subseteq X\varepsilon$ . Hence, by (2),  $\alpha \in P_\varepsilon$ . But there exists  $x \in X$  such that  $y\varepsilon = x\alpha$ . Therefore, since  $\varepsilon\gamma = \gamma$ , we have that

$$x\alpha\gamma = y\varepsilon\gamma = y\gamma \notin X\alpha.$$

In particular, this shows that  $\gamma$  cannot belong to a left normal divisor of  $P_\varepsilon$ . Hence the only left normal divisor of  $P_\varepsilon$  is the subgroup  $\{\varepsilon\}$  of  $H_c$ . It then follows from the theorem that the only idempotent-separating congruence on  $T_c$  is the identity congruence.

#### REFERENCES

- (1) A. H. CLIFFORD and G. B. PRESTON, *The Algebraic Theory of Semigroups*, vol. 1, Math. Surveys of the American Math. Soc. 7 (Providence, R.I., 1961).
- (2) G. LALLEMENT, Congruences et équivalences de Green sur un demi-groupe régulier, *C.R. Acad. Sc. Paris*, **262** (1966), 613-616.
- (3) A. I. MAL'CEV, Symmetric groupoids, *Mat. Sbornik*, **31** (73) (1952), 136-151.
- (4) G. B. PRESTON, Congruences on completely 0-simple semigroups, *Proc. London Math. Soc.* (3), **11** (1961), 557-576.
- (5) D. REES, On the ideal structure of a semi-group satisfying a cancellation law, *Quarterly J. Math.* Oxford Ser. **19** (1948), 101-108.
- (6) N. R. REILLY and A. H. CLIFFORD, Bisimple inverse semigroups as semigroups of ordered triples (to appear).
- (7) R. J. WARNE, The idempotent-separating congruences of a bisimple inverse semigroup with identity (to appear).

UNIVERSITY OF GLASGOW  
and  
UNIVERSITY OF STIRLING