

The Immunopathology of Sepsis: Pathogen Recognition, Systemic Inflammation, the Compensatory Anti-Inflammatory Response, and Regulatory T Cells

D.H. Lewis, D.L. Chan, D. Pinheiro, E. Armitage-Chan, and O.A. Garden

Sepsis, the systemic inflammatory response to infection, represents the major cause of death in critically ill veterinary patients. Whereas important advances in our understanding of the pathophysiology of this syndrome have been made, much remains to be elucidated. There is general agreement on the key interaction between pathogen-associated molecular patterns and cells of the innate immune system, and the amplification of the host response generated by pro-inflammatory cytokines. More recently, the concept of immunoparalysis in sepsis has also been advanced, together with an increasing recognition of the interplay between regulatory T cells and the innate immune response. However, the heterogeneous nature of this syndrome and the difficulty of modeling it *in vitro* or *in vivo* has both frustrated the advancement of new therapies and emphasized the continuing importance of patient-based clinical research in this area of human and veterinary medicine.

Key words: Clinical pathology; Critical care; Hematology; Immunology; Lymphocyte function; Multiple organ failure; Shock.

Sepsis, defined as the systemic inflammatory response to infection (Box 1), remains the major cause of death in critically ill human patients.^{1–6} Recent human studies estimate the annual incidence of sepsis to be 240–300 cases per 100,000 population, with associated costs of nearly \$17 billion in the United States^{7,8}; the rate of occurrence is also increasing at around 9% each year.⁸ Although large-scale veterinary epidemiological studies are uncommon, a substantial proportion of the critically ill veterinary population is estimated to be septic.^{9,10} The case fatality rate associated with sepsis in a variety of veterinary species is reported to approach 50%, emphasizing the need for a greater understanding of the pathophysiology of sepsis to improve therapeutic practices.^{11–13}

The pathophysiology of sepsis remains incompletely understood. A multitude of cell types, inflammatory mediators, and coagulation factors are involved and recent research has focused on the contributions of the innate immune system and T cells in this complex syndrome.^{2,14–20} This review will present an account of the current understanding of the function of the immune system in sepsis, with emphasis on the interaction of pathogens with innate components of the immune system and the key role of the endothelium in triggering and propagating a pro-inflammatory state.

From the Department of Veterinary Clinical Sciences, The Royal Veterinary College, Hatfield Campus, Hertfordshire, UK (Lewis, Chan, Garden); Regulatory T Cell Laboratory, The Royal Veterinary College, Camden Campus, London, NW1 0TU, UK (Pinheiro, Garden); Davies Veterinary Specialists, Manor Farm Business Park, Hertfordshire, SG5 3HR, UK (Armitage-Chan); and DHL's current address: Langford Veterinary Services, Small Animal Hospital, Langford House, Langford, Bristol, BS40 5DU, UK.

Corresponding author: O.A. Garden, Department of Veterinary Clinical Sciences, The Royal Veterinary College, Hawkshead Lane, North Mymms, Hatfield, Hertfordshire, AL9 7TA, UK; e-mail: ogarden@rvc.ac.uk.

Submitted November 30, 2010; Revised November 28, 2011; Accepted February 7, 2012.

Copyright © 2012 by the American College of Veterinary Internal Medicine

10.1111/j.1939-1676.2012.00905.x

Abbreviations:

AGE	advanced glycation end product
AP-1	activator protein 1
aPC	activated protein C
BK	bradykinin
CARD	common caspase activation and recruitment domain
CLARP	caspase-like apoptosis regulatory protein
DAMP	danger-associated molecular pattern
gp	glycoprotein
HMBG	high mobility group box protein
HSP	heat shock protein
Ig	immunoglobulin
IKK	inhibitor of κ B kinase
IPAF	interleukin-1 converting enzyme (ICE) protease-activating factor
IPS-1	interferon- β promoter stimulator-1 (also known as CARD adaptor inducing interferon- β [Cardif])
IRAK	IL-1R associated kinase
IRF	interferon regulatory factor
KK	kallikrein
LDL	low density lipoprotein
LGP2	laboratory of genetics and physiology 2
MAPK	mitogen-activated protein kinase
MDA5	melanoma differentiation-associated gene 5
MODS	multiple organ dysfunction syndrome
MyD88	myeloid differentiation primary response gene 88
NAIP	neuronal inhibitor of apoptosis
NEMO	NF- κ B essential modulator
NF- κ B	nuclear factor kappa B
NLR	nucleotide-binding domain, leucine-rich repeat containing protein
NLRB	NLR family, baculovirus inhibitor of apoptosis protein repeat domain containing
NLRC	NLR family, CARD containing
NLRP	NLR family, pyrin domain containing
NOD	nucleotide-binding oligomerization domain
PAMP	pathogen-associated molecular pattern
PAR	protease-activated receptor
PMN	polymorphonuclear cell
PRR	pattern-recognition receptor
RAGE	receptor for advanced glycation end products
RICK	RIP-like interacting CLARP kinase
RIG-1	retinoic acid-inducible gene-1

Abbreviations:

AGE	advanced glycation end product
AP-1	activator protein 1
aPC	activated protein C
BK	bradykinin
CARD	common caspase activation and recruitment domain
CLARP	caspase-like apoptosis regulatory protein
DAMP	danger-associated molecular pattern
gp	glycoprotein
HMBG	high mobility group box protein
HSP	heat shock protein
Ig	immunoglobulin
IKK	inhibitor of κ B kinase

Box 1

Definitions of SIRS, sepsis, and MODS^{1,4,6}

SIRS (systemic inflammatory response syndrome): clinical manifestation of the systemic response to injury.

Sepsis: SIRS in association with bacterial, viral, protozoal or fungal infection.

Severe sepsis: sepsis together with evidence of organ dysfunction, hypoperfusion, or hypotension.

Septic shock: sepsis with hypotension despite adequate fluid resuscitation.

MODS (multiple organ dysfunction syndrome): altered organ function in an acutely ill patient such that homeostasis cannot be maintained.

In addition, the complex interplay between pro- and anti-inflammatory cytokines and the spectrum of the host defense response will be discussed. Finally, the importance of regulatory T cells (Tregs) in maintaining the balance of host inflammatory mechanisms will be described.

Immunopathology of Sepsis

Living organisms face a constant barrage of potentially pathogenic microorganisms. Survival depends upon physical barriers to resist entry of pathogens, as well as the presence of a constitutive, or innate, immune system that can rapidly induce a defensive inflammatory response. Such a system can be found in virtually all species, suggesting that it is evolutionarily ancient and highly successful.^{21,22} The innate immune system further interacts with the adaptive immune system, based on a system of T and B lymphocytes that respond to specific epitopes.

Surface Barrier Mechanisms and Antimicrobial Peptides

Whereas the keratinized epithelium of the dermis and the mucous lining of the body cavities discourage

pathogenic colonization, various other innate defenses are employed to minimize penetration of the body wall by microorganisms, including antimicrobial peptides (AMPs; otherwise known as host-defense peptides) on mucosal surfaces.^{23,24} A comprehensive review of AMPs in veterinary species has recently been published.²⁵ In summary, AMPs comprise 3 main groups: digestive enzymes and peptides that disrupt the microbial cell membrane, peptides that bind essential elements, and peptides that act as decoys for microbial attachment. The 2 major classes of bactericidal AMPs in the mammalian immune system are the defensins and cathelicidins. Compared to eukaryotic cells, bacterial cell walls lack cholesterol—which acts to stabilize cell membranes—and negatively charged phospholipids. Instead, bacterial cell walls are rich in anions (basic amino acids) and thus attract the cationic defensins, which carpet the microbial membrane and institute channel formation.²⁶ These channels then lead to transmembrane pore formation, membrane destabilization, and microbial cell death, although recent studies suggest that their function could go beyond that of lipid bilayer perturbation.²⁷ Whereas a large number of AMPs form part of the non-oxidative killing mechanism within phagolysosomes in cells such as neutrophils and macrophages, a growing number are thought to be actively secreted onto epithelial surfaces of the gastrointestinal, respiratory and urinary tracts^{28–31}—including the ovine gastrointestinal tract³² and the canine testis.³³ Recent work has revealed that important amounts of these AMPs are secreted not only by immune cells such as neutrophils and alveolar macrophages but also by atypical defense cells such as type II pneumocytes.³⁴ This theme is universal throughout the processes of the innate immune system and challenges the traditional view of a “standing army” of immune defense cells, replacing it with the concept of a body-wide, integrated community of cells contributing to pathogen vigilance.³⁵

Recognition of Pathogens: PAMPs, MAMPs and DAMPs

Two factors are vital to the rapid ability of the innate immune system to respond to pathogen incursion: the presence of receptors against pathogen markers and the ubiquitous nature of these receptors in the body.³⁵ Individual receptors are genetically encoded and display strong homology within and between species.²² These receptors are not only expressed on many effector cells of the immune system—including macrophages, neutrophils, dendritic cells, and lymphocytes—but are also found on epithelial cells, endothelial cells, and myocytes^{36,37}; expression has also been detected in the bovine endometrium.³⁸ The major targets of these pattern recognition receptors (PRRs) are known as pathogen-associated molecular patterns (PAMPs), although the presence of these molecules in nonpathogenic and commensal bacteria has led to the suggestion that the term “microbial-associated molecular

patterns” (MAMPs) is more accurate. These molecules share certain core characteristics:

- PAMPs are produced only by microbial pathogens, not by the host (eg, peptidoglycans are produced by bacteria but not by eukaryotic cells): this confers automatic self/nonself discriminatory ability.
- PAMPs are generally invariant molecules shared by entire classes of microorganisms (eg, lipoarabinomannan is found on the cell wall of all *Mycobacteria*); this allows the evolutionary retention of a relatively small number of PRRs recognizing vast numbers of potential pathogens.
- PAMPs are usually structures vital to the survival or pathogenicity of the microorganism (eg, lipopolysaccharide [LPS] in the outer membrane of Gram-negative bacteria): this allows targeting of highly conserved molecules and obviates the need for variability in host PRRs.

Although bacterial cell surface components such as LPS in the outer cell membrane of Gram-negative bacteria and lipoteichoic acid (LTA) in the cell membrane of Gram-positive bacteria represent classic examples of PAMPs, recognition of “altered self” secondary to host cell colonization by viral pathogens is also likely. The innate immune system has thus developed the ability to detect markers of endogenous cell damage called “alarmins” or “danger-associated molecular pat-

terns” (DAMPs)^{39–41} Table 1 shows key PAMPs and their corresponding receptors,^{42,43} whereas Table 2 shows some of the confirmed interactions of DAMPs.^{39,40,43–46} Nonrodent sepsis models, genetic approaches, and immunological studies have demonstrated the presence of a large number of PRRs in clinical veterinary species (Table 3).^{38,47–55}

Initiation of an inflammatory reaction to necrotic, rather than apoptotic, cell death would appear to be useful in host defense; however, the interaction of DAMPs and PAMPs with their receptors leads to increased case fatality rates in sepsis.^{15,19,56} Owing to PRR cross-reactivity for both PAMPs and DAMPs, multiple positive feedback systems become established, leading to rapid progression of a global inflammatory response with consequent clinical signs (Fig 1).^{16,19,57} Functional interactions between PRRs, including synergy and cross-tolerance, also occur.^{58,59} If an inflammatory state persists, the very defensive mechanisms of the innate immune system designed to protect the host can lead to further tissue damage, as well as diminished antimicrobial activity that allows opportunistic secondary infections.^{15,20} Whereas the existence of such feedback systems in veterinary species can at present only be inferred, experimental data suggest PRR reactivity to both pathogen- and host-derived ligands in cattle, pigs, horses, and dogs.^{60–66} Furthermore, the blunted PAMP-induced TNF, IL-6 and IL-10 response of whole blood in dogs with lymphoma is thought to underlie their higher risk of sepsis.⁶⁷

Table 1. Key pathogen-associated molecular pattern ligands of the pattern recognition receptors implicated in sepsis.

PRR	Location	Cell Type	PAMP Recognized
TLR 1/CD281	pm	pbmc, np, u	triacyl lipopeptide
TLR 2/CD282	pm, el	pbmc, dc, mc, nkc	peptidoglycan, lipoprotein, lipopolysaccharide, glycosylphosphatidylinositols, mannans
TLR 3/CD283	pm, el	dc, epi, fb, blc, nkc	ssRNA, dsRNA, dsDNA
TLR 4/CD284	pm	pbmc, mc, np, epi	lipopolysaccharide, glycosylphosphatidylinositols, viral envelope proteins, mannans
TLR 5	pm	pbmc, dc, epi, nkc	flagellin
TLR 6/CD286	pm	mc, blc	diacyl lipopeptide
TLR 7	el	pbmc, dc, blc	ssRNA
TLR 8/CD288	el	nkc	ssRNA
TLR 9/CD289	el	dc, blc, nkc, epi	ssDNA, dsDNA
TLR 10/CD290	pm	pbmc, dc, blc	Unknown
TLR 11	pm	epi, dc, pbmc	profilin
NOD1/NLRC1	cyt, pm	epi, dc, pbmc	peptidoglycan
NOD2/NLRC2/CARD15	cyt, pm	epi, dc, pbmc, Paneth cells	muramyl dipeptide
NLRC4/IPAF	cyt	unknown	flagellin
NLRP1	cyt	unknown	muramyl dipeptide, <i>Bacillus anthracis</i> lethal toxin
NLRP3	cyt	unknown	bacterial & viral RNA, lipopolysaccharide, lipoteichoic acid, muramyl dipeptide
NLRB1/NAIP5	cyt	unknown	flagellin

PRR, pattern-recognition receptor; PAMP, pathogen-associated molecular pattern; TLR, Toll-like receptor; NOD, nucleotide-binding oligomerization domain; NLR, nucleotide-binding domain, leucine-rich repeat containing protein; NLRC, NLR family, CARD containing; IPAF, interleukin-1 converting enzyme (ICE) protease-activating factor; NLRP, NLR family, pyrin domain containing; NLRB, NLR family, baculovirus inhibitor of apoptosis protein repeat domain containing; NAIP, neuronal inhibitor of apoptosis; pm, plasma membrane; el, endolysosomes; cyt, cytoplasm; np, neutrophils; u, ubiquitous; dc, dendritic cells; mc, mast cells; nkc, natural killer cells; epi, epithelial cells; fb, fibroblasts; blc, B lymphocytes; ss, single-stranded; ds, double-stranded.

Table 2. Key danger-associated molecular pattern ligands of the pattern recognition receptors.

PRR	Location	Cell Type	DAMP Recognized
TLR 2/CD282	pm, el	pbmc, dc, mc, nkc	HMGB 1, necrotic cells, HSP-60, HSP-70, gp-96, biglycan, defensins
TLR 3/CD283	pm, el	dc, epi, fb, blc, nkc	endogenous mRNA
TLR 4/CD284	pm	pbmc, mc, np, epi	HSP-22, HSP-70, HSP-90, fibronectin, fibrinogen, heparan fragments, hyaluronate fragments, β -defensin 2, oxidized LDL, surfactant protein A, neutrophil elastase, HMGB 1, biglycan
TLR 9/CD289	el	dc, blc, nkc, epi	chromatin-IgG complex
NLRP3	cyt	unknown	uric acid crystals
RAGE	pm	u?	AGEs, HMGB 1, amyloid peptide, S100s

New abbreviations (for previous abbreviations see Table 1): DAMP, danger-associated molecular pattern; RAGE, receptor for advanced glycation end products; HMGB, high mobility group box protein; HSP, heat shock protein; gp, glycoprotein; LDL, low density lipoprotein; Ig, immunoglobulin; AGEs, advanced glycation end products; S100s, S100 proteins (calgranulins).

Table 3. Pattern-recognition receptors of those families implicated in sepsis identified in veterinary species to date.

Species	TLRs	NLRs
Dog	1–7, 9	NOD1, NOD2, NLRC4 NLRP 1–3, 5, 6, 8–10, 12–14
Cat	2–5, 7–9	
Horse	2–4, 6, 9	NLRC4
Cow	1–10	NOD1, NOD2, NLRC4 NLRP 1, 3, 5, 6, 8–10, 12–14
Sheep	1–10	NOD2
Goat	1–10	
Pig	1–10	NOD1, NOD2

For abbreviations, see footnote to Table 1.

Table 4. The spectrum of type I and type II acute phase proteins.

Acute Phase Protein	Role	Type
Serum amyloid A	Leukocyte recruitment and activation	I – induced by IL-1 and TNF
C-reactive protein	Enhance microbial phagocytosis and complement binding	
C3, C4, C4BP, C1inh	Complement components	
Haptoglobin (rat)	Binds free hemoglobin	
α 1-acid glycoprotein	Transport functions	
Fibrinogen	Hemostasis	II – induced by IL-6 and IL-6-like cytokines
Haptoglobin (man)	Binds free hemoglobin	
(Apo)Ferritin	Binds free iron	
α 1-antitrypsin	Protease inhibitor	
α 2-macroglobulin	Protease inhibitor	

Pattern Recognition Receptors: an Overview

Evolutionary pressure has resulted in the encoding of a number of different host proteins within three distinct families—the Toll-like receptors (TLRs), the nucleotide-binding domain, leucine-rich repeat containing proteins (NLRs; previously designated as the nucleo-

tide-binding oligomerization domain [NOD]-like receptors) and the retinoic acid-inducible gene-1 (RIG-1)-like receptors (RLRs).^{68–70} Cooperation between the RLRs—intracellular viral nucleic acid sensors—and the endosomal TLRs appears likely,^{71–73} although little is currently known about the involvement of RLRs in the pathogenesis of sepsis.¹⁴ This family of PRRs is therefore not considered further in this review.

Whereas PRRs form a heterogeneous group of proteins, certain characteristics—such as leucine-rich repeat domains, scavenger receptor cysteine-rich domains, and C-type lectin domains—can be commonly recognized.^{68,69}

The Toll-Like Receptors

The Toll gene was first identified in *Drosophila melanogaster* as encoding a transmembrane glycoprotein with a role in determination of dorsoventral polarity in the embryo.⁷⁴ Additional investigation revealed a large extracellular component with leucine-rich repeats, with a cytoplasmic portion described as the TIR (Toll/interleukin-1 receptor [L-1R]) domain owing to its similarity to the intracellular region of the mammalian IL-1R.⁷⁵ Not only was the Toll protein discovered to play a role in the innate immune response to fungal infection in *Drosophila*, but a subsequent series of studies served to highlight the link between TLRs and the mammalian immune response.^{76–79} To date, 10 human and 12 murine functional TLRs have been identified by a combination of immunological techniques and examination of genetic databases.^{42,71}

The cytoplasmic TIR domain of the TLRs interacts with a variety of TIR-domain-containing adaptors (Fig 2).^{80–83} PAMP/DAMP ligation of the respective PRR activates a chain of kinases in the IL-1R associated kinase (IRAK) family, ultimately resulting in activation of the inhibitor of κ B kinase (IKK) enzyme complex and the mitogen-activated protein kinase (MAPK) pathway (Fig 2).^{84–86} Whereas the connection of TLR activation to the NF- κ B and MAPK signaling pathways is well recognized, a number of other pathways are also triggered by TLR stimulation, including Protein Kinase R/eukaryotic translation initiation




Fig 1. The pathophysiology of sepsis—an overview. A primary infectious insult—of bacterial, viral, protozoal or fungal origin—damages host tissues. The ligation of pattern-recognition receptors (TLRs and NLRs) by PAMPs and DAMPs promotes the release of both pro- and anti-inflammatory cytokines, as well as acute phase proteins, with a number of pathophysiological sequelae that ultimately lead to organ hypoperfusion, tissue hypoxia, the generation of ROS and RNS, mitochondrial dysfunction and cell death by necrosis, apoptosis, pyroptosis, and autophagy. Further DAMPs are generated, helping to perpetuate the inflammatory response, and the death of immune cells leaves the organism vulnerable to secondary, or opportunistic, infections. Abbreviations: PAMPs = pathogen-associated molecular patterns; DAMPs = danger-associated molecular patterns; TLRs = Toll-like receptors; NLRs = nucleotide-binding domain, leucine-rich repeat containing protein; RLR = retinoic acid-inducible gene-1 (RIG-1)-like receptor; NFκB = nuclear factor kappa B; PMN = polymorphonuclear cell; ROS = reactive oxygen species; RNS = reactive nitrogen species.

factor 2- α kinase 2 (PKR/eIF2 α), Notch, phosphoinositide 3-kinases (PI-3K), and small GTPases.^{87–92}

TLRs are expressed by a number of different cells, including dendritic cells, macrophages, B cells, natural killer (NK) cells, endothelial cells, epithelial cells, and fibroblasts (Tables 1 and 2); their site of expression varies, with TLRs 1, 2, 4, 5, 6, and 11 present on the external plasma membrane and TLRs 3, 7, 8, and 9 in endosomes.^{71,93}

TLR 2 appears to be of key importance, owing to its ability to recognize PAMPs as diverse as lipoteichoic acid (from Gram-positive bacteria), peptidoglycan (from Gram-positive and Gram-negative bacteria),

hemagglutinin (from measles virus), polysaccharides (from yeasts), lipoproteins (from *E. coli*, *Borrelia burgdorferi*, *Mycoplasma* spp. and *Mycobacterium tuberculosis*), as well as complete pathogens such as *Clostridium* spp., *Chlamydomphila* spp., and herpes simplex virus, and a variety of endogenous ligands.^{94–97}

Much of its wide-ranging influence stems from the formation of heterodimers with TLRs 1 and 6.^{98–100} TLR 4 also has a significant role in triggering the innate immune response as it recognizes molecules such as LPS (from Gram-negative bacteria), various viral protein envelopes, and a large number of endogenous molecules (Table 2).^{57,101–103} The vital recognition of LPS by TLR




Fig 2. Toll-like receptor (TLR) signaling pathways. The cytoplasmic Toll/interleukin-1 receptor (TIR) domain of the TLRs interacts with TIR-domain-containing adaptors, such as myeloid differentiation primary response gene 88 (MyD88), TIR-containing adaptor protein (TIRAP), TIR-domain-containing adaptor molecule 1 (TICAM1, also known as TRIF) and TIR-domain-containing adaptor molecule 2 (TICAM2, also known as TRAM). PAMP binding to the respective receptor results in the activation of either the MyD88 or TICAM1/TRIF signaling pathways. These pathways involve a series of kinases in the IL-1R associated kinase (IRAK) family, whose action eventually results in activation of the inhibitor of κ B kinase (IKK) enzyme complex and the mitogen-activated protein kinase (MAPK) pathway. The IKK complex phosphorylates the inhibitory I κ B α protein, thus freeing the nuclear transcription factor nuclear factor kappa B (NF- κ B); triggering of the MAPK signaling pathway results in the activation of activator protein 1 (AP-1). Both NF- κ B and AP-1 then initiate the transcription of cytokine genes. Abbreviations: TRAF = TNF receptor-associated factor; RIP = receptor interacting protein; TAK = TGF- β -activated kinase; NEMO = NF- κ B essential modulator; TBK = TRAF family member-associated NF- κ B activator (TANK)-binding kinase; IRF = interferon regulatory factor.

4 appears to be dependent on formation of a complex with other PRRs, myeloid differentiation protein-2 (MD2), membrane-bound CD14 (mCD14), and lipopolysaccharide-binding protein (LBP).^{71,104} Both TLR 2 and 4 have been extensively researched in human septic patients, in whom early upregulation of these TLRs can exacerbate the severity of illness and mortality.^{19,57}

TLR 5 is expressed by epithelial cells of the respiratory and intestinal tract and is a PRR for flagellin, an important component of motile bacteria such as *Salmonella* spp.^{105,106} TLR 5 is incriminated in sepsis,¹⁰⁷ although exactly how the PAMP and PRR interact *in vivo* remains unclear.¹⁰⁸ The intracellular TLRs 3, 7, 8, and 9 all share high sequence homology and recognize nucleotides.^{71,109} TLR 3 is the only TLR yet discovered that does not initiate the MyD88 signaling pathway, interacting solely with TIR-domain-containing adaptor molecule 1 (TICAM1, also known as TRIF) (Fig 2).^{109,110} TLR 11, to date only identified in rats and mice, appears to recognize *Toxoplasma gondii* and certain pathogens of the urinary tract, although a role in sepsis has not so far been described.^{111,112}

The Nucleotide-Binding Domain, Leucine-Rich Repeat Containing Proteins, and Inflammasomes

The realization that TLRs could not account for the full range of PAMP recognition motivated the discovery of additional intracellular PRRs, including the NLRs. Of the NLRs, the 2 cytosolic receptors NOD1 and NOD2 were the first to be discovered; subsequent examination of genomic databases has suggested that there are at least 23 NLRs in humans and 34 in mice.^{113–117} Common to all NLRs is their structure, comprising a leucine-rich repeat domain (thought to be the PAMP receptor region), a central NOD domain, and an N-terminal effector domain responsible for downstream signaling.^{118–120} The NLRs are found in the cytosolic compartment of eukaryotic cells, although some recent evidence suggests that they can also be associated with the plasma membrane.^{121,122} The mechanisms underlying NLR contact with their respective PAMPs have yet to be confirmed *in vivo*,^{123–125} but recognition of the PAMP triggers the activation of a series of kinases leading to the phosphorylation of




Fig 3. Nucleotide-binding domain, leucine-rich repeat containing protein (NLR) signaling pathways. Upon PAMP ligation, NLRs recruit the serine-threonine kinase RIP-like interacting caspase-like apoptosis regulatory protein (CLARP) kinase (RICK), also known as receptor-interacting serine/threonine-protein kinase 2 (RIPK2 or RIP2), which binds to NF- κ B essential modulator (NEMO), a subunit of IKK, resulting in the phosphorylation of I κ B and the release of NF- κ B; RICK also mediates the recruitment of transforming growth factor β -activated kinase 1 (TAK1) and together these molecules stimulate activation of the mitogen-activated protein kinase (MAPK) signaling pathway. In common with the TLR family, NLR activation leads to the transcription of inflammatory cytokine genes via the mobilization of NF- κ B and AP-1. Another important consequence of NLR ligation is activation of the inflammasome, a macromolecular complex comprising pro-caspase-1 together with various adapter proteins. Caspase-1 is important in the synthesis of active IL-1 β and IL-18, and induces a type of programmed cell death called pyroptosis.

I κ B, mobilization of NF- κ B, and activation of MAPK signaling (Fig 3).^{126–129} NOD1 (NLR family, caspase activation, and recruitment domain [CARD] containing 1; NLRC1) and NOD2 (NLRC2, also known as CARD15) are known to recognize muopeptides derived from peptidoglycan, a major structural component of both Gram-positive and Gram-negative bacterial cell walls (Table 1).^{115,118} Whereas the role of NOD2 is well established in the pathogenesis of Crohn's disease,¹³⁰ little is known about the role of the NOD proteins in sepsis. However, synthetic NOD1 agonists administered to mice stimulate chemokine production, neutrophil recruitment,¹³¹ and—in one model—shock and multiple organ dysfunction,¹³² raising the possibility of NOD1 involvement in the pathogenesis of sepsis.

A number of studies have been performed on the role of inflammatory cysteinyl aspartate-specific proteinases, or caspases, in murine and human sepsis. Caspase-1 has been the focus of particular attention and is activated by a macromolecular complex of around 700 kDa—one of a number of inflamma-

somes¹³³—comprising pro-caspase-1 together with various adapter proteins, including apoptosis-associated speck-like protein containing a carboxy-terminal CARD (ASC).^{133–135} Activation of the currently identified inflammasomes depends upon the interaction of NLR family members (NLRC4; NLR family, pyrin domain containing 1 [NLRP1] and NLRP3) with their PAMP or DAMP ligands.^{136–138} Caspase-1 is thought to be the executioner caspase in inflammation, important in the synthesis of active IL-1 β and IL-18 and thus pathogen defense. Hence, mice deficient in caspase-1 are more susceptible to bacterial infections and sepsis^{139–141}; in contrast, mice deficient in caspase-12, which normally suppresses caspase-1 activity, show enhanced bacterial clearance and resistance to sepsis.¹⁴² Parallel observations have been made in human patients possessing the full-length caspase-12 proenzyme—uniquely present in approximately 20% of people of African descent—which attenuates the innate immune response to endotoxin and is thought to be a risk factor for the development of severe sepsis.¹⁴³ Whereas caspase-1 is essential for host defense against

pathogens, its activity needs to be tightly controlled. Excessive caspase-1 activity and thus endotoxic shock was induced in mice given high doses of LPS, but mice deficient in caspase-1,¹⁴⁴ or ASC in another study,¹⁴⁵ were resistant to the lethal effects of LPS. Inflammatory activity has also been linked to the phenomenon of immunoparalysis. Thus, the expression of mRNA encoding the key inflammasome components ASC, NALP1, and caspase-1 was decreased in human monocytes during early septic shock, thought to reflect monocyte deactivation; furthermore, NALP1 mRNA abundance was linked to survival in patients with sepsis.¹⁴⁶ Very little is known about inflammasomes in the veterinary species, although inflammasome assembly mRNAs have been analyzed in the ovine jejunum¹⁴⁷ and induction of inflammasome genes in the spleen was documented in septic pigs infected with either *Haemophilus parasuis*¹⁴⁸ or *Streptococcus suis*.¹⁴⁹

Pattern Recognition Receptors: Veterinary Species

After mapping of the gene sequences for human and murine PRRs, exploration of the genomic sequence of other species has allowed the identification of homologs of the majority of TLRs in dogs,^{150–154} cats,^{49,151,155,156} cattle,^{38,50,51,62,157–168} sheep,^{48,51,169–173} goats,^{52,174,175} horses,^{47,168,176–181} and pigs.^{169,182–191} Other PRRs have also been identified in these species, although less is known about them than the TLRs (Table 3).

Toll-like receptors 1–10 have been identified in the bovine genome,^{50,161} with numerous studies documenting their expression in tissues as varied as the endometrium,^{38,163} cornea,¹⁵⁸ mammary gland,^{167,192} skin,⁵¹ and lung.¹⁶⁸ Whereas most of these studies have utilized PCR techniques to identify TLR expression,^{38,51,158,163,167} an increasing number are employing flow cytometry^{192,193} or immunohistochemistry.¹⁶⁸ TLR signaling in cattle is similar to that described in mice and humans (Fig 2)^{162,192,194} and the PKR/eIF2 α pathway appears to be important in bovine viral diarrhea virus (BVDV) and rotavirus infections.^{195,196} Comparatively little is known about the NLR family in cattle, although mRNA encoding NOD 1 and 2 has been identified in bovine mammary tissue.^{197,198} Because of the financial implications of Johne's disease and the currently unconfirmed link between *Mycobacterium avium* ssp. *paratuberculosis* and Crohn's disease, a number of studies have examined the potential role of polymorphisms of candidate genes—including TLRs—in susceptibility to paratuberculosis.¹⁹⁹ Recently, TLRs 1–10 have also been cloned and sequenced in sheep⁴⁸ and goats.¹⁷⁴ Little is currently known about the implication of TLRs in sepsis of ruminants, although the pathways involved in triggering SIRS in sheep appear to be similar to those reported in other species.^{169,200}

The impact of infectious agents upon commercial viability of pigs and the contribution of this species to human disease modelling has helped advance the characterization of porcine PRRs.¹⁹¹ Thus, TLRs 1–10 have been cloned and sequenced^{201–208}; surface and endosomal TLRs have been detected—both by PCR

and immunohistochemistry—in a variety of porcine tissues and enhanced expression demonstrated in response to a number of infectious agents and PAMPs.^{168,183,209–211} Members of the NLR family have also been identified^{212–216} and current research suggests that porcine PRR signaling pathways are similar to those of other mammalian species.^{169,217}

Various members of the TLR family have been identified in horses, including TLR 2, 3, 4, 5, and 9.^{47,53} Whereas many of these have been characterized by PCR,^{176–179} TLRs 4 and 9 have also been localized by immunohistochemistry and immunogold electron microscopy^{168,179,181}—and a recent study demonstrated TLR 5 expression by equine neutrophils by flow cytometry.²¹⁸ Stimulation by PAMPs has increased gene expression of TLRs 2, 3, and 4 *in vitro*,¹⁷⁷ and TLR 9 *in vivo*,¹⁸¹ whereas clinical studies have reported increased TLR 4 gene expression in both foals and adult horses with SIRS/sepsis, but no differences in expression between survivors and nonsurvivors.^{219,220} Complementary studies have demonstrated increased plasma endothelin-1 concentrations and decreased long-term survival in horses with severe versus mild-to-moderate endotoxemia.²²¹ As for cattle and pigs, initial investigations have indicated that the downstream signaling pathways instigated by PAMP stimulation of equine TLRs 2–4 are similar to those identified in other species.²²² To date, there have been no published reports on members of the NLR family in the horse.

Less is currently known about PRRs in small animals.^{53–55} Various PCR studies have confirmed the presence of members of the TLR family in both dogs, including TLRs 2, 4, 7, and 9,^{61,151–154,223} and cats, including TLRs 1–9.^{49,151,155,156} Comparative examination of the canine genome has also identified the presence of genes encoding members of the NLRP family, integral to inflammasome and thus caspase activation.²²⁴ A number of canine TLRs have also been detected by immunohistochemical staining and flow cytometry.^{168,209,225,226} The use of feline models for investigation of human diseases as diverse as type 2 diabetes mellitus and human immunodeficiency virus infection has yielded additional data on PRRs in cats, including the expression of functional TLRs by the endocrine pancreas¹⁵⁵ and modulation of TLR signaling by retroviral pathogens.^{156,227} Enhanced expression of canine TLRs has been observed in clinical cases of osteoarthritis⁶⁰ and cystic endometrial hyperplasia/pyometra^{61,226}; however, the majority of published research in dogs concerns the expression of PRRs in the intestinal tract, particularly in relation to inflammatory bowel disease (IBD).^{150,223,228} Ongoing research is attempting to identify whether or not certain single nucleotide polymorphisms (SNPs) of PRRs can be related to the propensity for particular canine breeds to develop IBD and other immune-mediated diseases.^{229–231} The analysis of SNPs in PRR-encoding genetic sequences is also an exciting field of research in large animals,^{164,213,232} laying the foundation for the breeding of livestock with enhanced disease resistance and

the design of vaccines better able to target dendritic cells.^{191,233} Finally, recent work has documented the expression of an ortholog of human “triggering receptor expressed on myeloid cells-1” (TREM-1) by canine neutrophils.²³⁴ Expression of TREM-1 was upregulated by microbial agonists of TLR1/2, TLR2/6, and TLR4/MD2.²³⁴ This receptor, which has shown promise as a biomarker of sepsis in humans, amplifies pro-inflammatory responses to microbial products.^{59,235}

Inflammatory Mediators: Cytokines and Chemokines

PRR ligation triggers signaling cascades that culminate in the activation of NF- κ B and AP-1 via MyD88 or TICAM1/TRIF (Fig 2).^{236–238} NF- κ B and AP-1 enter the nucleus and activate transcription sites for a variety of genes, including acute phase proteins, inducible nitric oxide synthase (iNOS), coagulation factors, and pro-inflammatory cytokines and chemokines, such as tumor necrosis factor (TNF)- α and ILs-1, 6, 8, and 12. The TICAM1/TRIF pathway results in the phosphorylation of interferon regulatory factors 3 and 7 (IRF3, IRF7), which likewise enter the nucleus and stimulate the transcription of genes encoding interferon (IFN) α , IFN β , and other type 1 IFN-inducible genes.^{236–238}

Serum concentrations of TNF- α correlate with death in certain types of human sepsis.^{239,240} Studies of naturally occurring sepsis in veterinary patients have yielded similar results, although this pattern appears not to be universal: increased serum concentrations of TNF- α correlate with mortality in canine parvovirus and neonatal septicemia in cattle and horses, but not in septic cats.^{11,241,242} TNF- α is predominantly produced by activated macrophages and T cells—but also by mast cells, B cells, NK cells, neutrophils, endothelial cells, myocytes, osteoblasts, and fibroblasts—as a 26 kDa precursor (pro-TNF) expressed on the plasma membrane; there it is cleaved by TNF-converting enzyme (TACE/ADAM17) to yield a 17 kDa soluble form, both the soluble and membrane-bound forms appearing to be active.²⁴³ TNF- α exerts its effects by interaction with one of 2 receptors, TNF receptors 1 and 2 (TNFR1, TNFR2).^{244,245} Activation of TNFR1 appears to mediate the proinflammatory and apoptotic pathways associated with inflammation, whereas TNFR2 plays a role in the promotion of tissue repair and angiogenesis.²⁴⁵ However, the complexity of signaling networks operating in sepsis is underlined by the observation that NF- κ B activity induces molecules that block apoptosis mediated by TNFR1, suggesting that integration of the various signals occurs *in vivo*.²⁴⁶ Some “cooperation” between the 2 receptor types, particularly at low TNF- α concentrations, is also likely and stimulation of both TNFRs leads to further NF- κ B and AP-1 release.^{244,247}

Many of the classical features of inflammation can be attributed to the actions of TNF- α upon the endothelium, with increased production of iNOS and cyclo-oxygenase 2 (COX-2) leading to vasodilatation and local slowing of blood flow²⁴⁸; TNF- α also stim-

ulates the expression of endothelial adhesion molecules such as E-selectin, intercellular adhesion molecule 1 (ICAM-1), and vascular cell adhesion molecule 1 (VCAM-1).^{249,250} These 3 molecules lead to the tethering of leukocytes to the endothelial wall and their transmigration into the interstitium, accompanied by fluid and plasma macromolecules.¹⁹ There is also evidence for the upregulation of TNF- α and other proinflammatory cytokines in a canine model of sepsis involving the intravenous infusion of low doses of LPS: increased serum concentrations of TNF- α , IL-1 β , and IL-6^{251–253} were accompanied in one of the studies by increased expression of pulmonary E-selectin and ICAM-1 and the influx of neutrophils—although whether the LPS induced the expression of the adhesion molecules directly or via induction of the proinflammatory cytokines, or both, remained unclear.²⁵¹ A similar phenomenon of up-regulation of proinflammatory cytokines, including TNF- α , has also been observed in cats²⁵⁴ and horses^{255–257} treated with intravenous LPS, as well as the transcription of neutrophil chemoattractants by equine endothelial cells stimulated by Th2 cytokines.²⁵⁸ Furthermore, concentrations of TNF- α , IL-6, and endotoxin were all higher in the blood and peritoneal fluid of horses with colic than in a healthy control group.²⁵⁹ The stimulation of feline whole blood with various PAMPs has also elicited the synthesis of pro-inflammatory cytokines, including TNF, IL-1 β , and CXCL-8.²⁶⁰ Plasma nitrite and nitrate, oxidation products of NO, have been examined in canine sepsis, in which they were present at higher concentrations than in dogs with SIRS alone.²⁶¹ An additional study showed that the inflammatory response to the intravenous administration of LPS, examined by measuring serum concentrations of TNF- α , IL-1 and IL-6, was mitigated in dogs fed a diet rich in fish oils, suggesting that diet may be an adjunct to conventional anti-inflammatory treatment.²⁶² In addition to the upregulation of iNOS, COX-2, and adhesion molecules, TNF- α also induces the expression of procoagulant proteins such as tissue factor (TF)—and down-regulates anti-coagulant factors such as thrombomodulin—leading to activation of the coagulation cascade.²⁶³ Despite the important role for TNF- α in endothelial activation, experimental evidence suggests that direct stimulation of TLRs expressed by endothelial and vascular smooth muscle cells may provide an alternative pathway for the vascular dysfunction seen in sepsis.^{264–266}

In addition to TNF- α , NF- κ B activation results in the transcription of a number of proinflammatory interleukins, such as IL-1, IL-6, CXCL-8 (IL-8), and IL-12 (Fig 1). IL-1 acts in a synergistic manner with TNF- α in the “hyperacute” period after innate immune stimulation in sepsis.^{267,268} Two proinflammatory forms of IL-1 (IL-1 α and IL-1 β) have been identified and induce the synthesis of adhesion molecules and cytokines by endothelial cells, encouraging leukocyte activation, endothelial tethering, and transmigration into the interstitium.^{269,270} IL-1 also upregulates

iNOS and COX2 production, acts as the major endogenous pyrogen in fever, and increases corticosteroid release via hypothalamic effects.^{269,271}

Another important proinflammatory cytokine in sepsis is IL-6: as well as stimulatory effects upon leukocyte activation and myeloid progenitor cell proliferation, IL-6 also triggers the acute phase response and is a powerful pyrogen.^{272,273} Like TNF- α and IL-1, plasma concentrations of IL-6 are increased in sepsis and may be predictive of progression to multiple organ dysfunction and death.^{17,274,275} Whereas enhanced gene expression of IL-6 correlates with death in septic foals, the opposite relationship appears to hold for serum IL-6 concentrations.^{219,276} However, increased serum concentrations of IL-6 and IL-1 β do correlate with death in reports of naturally occurring sepsis in dogs and cats.^{11,13}

Serum concentrations of anti-inflammatory cytokines, most notably IL-10, also increase in sepsis.^{277,278} This acts not only to inhibit the release of TNF- α , IL-1 β , and IL-6 from monocytes and macrophages, but also to induce the production of IL-1 receptor antagonist protein (IRAP-1) and soluble TNFR, thus reducing circulating concentrations of these cytokines.²⁷⁹ The critical role of IL-10 in mediating the balance between pro- and anti-inflammatory processes can be seen in experimental models: IL-10 knockout mice are profoundly susceptible to sepsis, whereas IL-10 administration prevents these consequences.²⁸⁰ In clinical situations, however, the pattern appears to be more complex, with increased serum IL-10 concentrations associated with mortality in septic foals and humans,^{281,282} but a lower prevalence of feline infectious peritonitis in cats infected with feline coronavirus.²⁸³

Two additional cytokines have recently been identified as being critical in sepsis. Macrophage migratory inhibitory factor (MIF), produced by the anterior pituitary gland, is present at increased concentrations in SIRS and sepsis^{284,285}; serum concentrations have not only correlated with mortality, but inhibition of MIF appears to be protective.²⁸⁶⁻²⁸⁸ Although the trigger for MIF release *in vivo* is unclear, it is thought to delay apoptosis of activated monocytes and macrophages, thus helping to perpetuate a proinflammatory state.²⁸⁸ High mobility group box protein 1 (HMGB-1) is an endogenous protein involved in nuclear DNA stabilization.²⁸⁹ However, after necrotic or apoptotic cell death, it is released into the circulation where it has direct pro-inflammatory actions.²⁹⁰ In addition, HMGB-1 appears to potentiate the effect of certain PAMPs and DAMPs upon TLR-2, TLR-4, and the receptor for advanced glycation end products (RAGE).²⁹¹ HMGB-1 was initially reported as a late inflammatory mediator in sepsis,²⁹² although ongoing research has highlighted a number of roles in tissue repair and angiogenesis.²⁹³ Circulating concentrations of HMGB-1 appear to correlate with mortality in canine SIRS patients,²⁹⁴ but were unable to predict hospital mortality in 1 study of septic human patients.²⁹⁵

The Acute Phase Response

In addition to the release of cytokines and chemokines from activated immune cells, triggering of PRRs also bring about the release of large quantities of acute phase proteins (APPs) from hepatocytes; these proteins have a variety of functions designed to re-establish homeostasis, assisting in pathogen elimination and subduing inflammation.^{296,297} The acute phase response is characterized by fever, neutrophilia, activation of the coagulation, and complement cascades (classical, alternative, and mannose-binding lectin pathways), serum iron and zinc binding, enhanced gluconeogenesis, increased muscle catabolism, and altered lipid metabolism.²⁹⁷⁻²⁹⁹

In general, 2 groups of APPs are recognized: type I, induced by IL-1 α , IL-1 β , and TNF- α , and type II, induced by IL-6.²⁹⁸ As a consequence of the up-regulation of APP production, concentrations of other plasma proteins such as albumin, protein C, protein S, and antithrombin (collectively known as negative APPs) decrease.^{299,300} Granulocyte colony-stimulating factor (G-CSF) released from monocytes is also an important component of the acute phase response, thought to mediate a protective role against bacterial infection by virtue of its impact on neutrophils.³⁰¹ A number of APPs have been characterized in veterinary species, including dogs, cats, cattle, sheep, horses, and chicken,^{297,302} and they have been used as biomarkers of inflammation in both research and clinical arenas in these species.^{303,304} Recent studies have identified adiponectin and insulin-like growth factor-1 as negative acute phase proteins in a canine model of endotoxemia.³⁰⁵

Much research has been conducted to determine whether or not serum levels of APPs are predictive of survival in sepsis, with procalcitonin and C-reactive protein (CRP) the focus of particular attention.³⁰⁶ Plasma procalcitonin concentrations appear to correlate with bacteremia and organ dysfunction in human clinical studies.^{306,307} Whereas it also appears to be a canine APP, it does not allow the discrimination of inflammatory or infectious from neoplastic disease when measured as a whole blood PCR assay³⁰⁸; however, extrathyroidal procalcitonin gene expression was documented in dogs with SIRS but not in healthy animals in a preliminary observational study.³⁰⁹ The utility of CRP measurements has been assessed in a number of studies examining infectious,^{310,311} inflammatory,³¹²⁻³¹⁹ neoplastic,^{320,321} and endocrine³²² diseases in the dog, including critically ill dogs³²³; in general, serum CRP concentrations provide a sensitive but nonspecific means of measuring inflammation, offering diagnostic and prognostic information in some disorders but not others. A recent study of SIRS and sepsis in dogs demonstrated a correlation between decreasing serum CRP concentration and recovery from disease, suggesting its use as a prognostic biomarker in this context.³²⁴

The Interaction of Inflammation and Coagulation in Sepsis

Despite not classically considered part of the innate immune response, the prevalence of coagulation disorders and disseminated intravascular coagulation (DIC) in sepsis underlines the intimate link between the inflammatory and coagulation pathways.^{325,326} On a local scale, activation of coagulation may act defensively to impede the dispersal of pathogens and inflammatory mediators from the site of insult.³²⁷ Clinical studies have supported experimental models demonstrating increased activation of coagulation, as well as downregulation of anticoagulant mechanisms and reduced fibrinolysis, in human SIRS and sepsis patients.^{266,328} Similar findings have also been reported in dogs³²⁹ and horses.³³⁰ Whereas DIC³³¹ and pulmonary thromboembolism³³² have both been recorded in association with sepsis in cats, experimental models of endotoxin infusion in this species have yielded variable results,^{333,334} a recent study failing to elicit any biologically significant alterations in coagulation parameters,³³⁴ underlining the multifactorial pathogenesis of coagulopathies in clinical patients.

A detailed description of the complexities of the interaction between inflammation and coagulation is beyond the scope of this article; however, several comprehensive reviews of hemostasis in SIRS and sepsis in veterinary species have recently been published and an overview of key interactions is presented in Fig 4^{335–337} In brief, key to the triggering of the coagulation pathway in sepsis is tissue factor (TF), which initiates coagulation via the contact activation (extrinsic) pathway. In health, lack of TF exposure within the vascular system and the presence of various circulating proteins—such as protein C, antithrombin, and tissue factor plasminogen inhibitor—modulate coagulation by the prevention of TF activation.^{266,335} The expression of TF by monocytes or macrophages and tissue parenchymal cells is activated by various inflammatory cytokines, CRP, and PAMPs such as LPS^{335,338}—a phenomenon that has also been documented in cats³³⁹ and horses (earlier studies in this species citing “procoagulant activity” rather than TF per se).^{340–342} Although findings differ between species, large numbers of TF-expressing microparticles have been identified in blood samples from septic human patients and may correlate with mortality.^{343,344} These microparticles are released from a variety of activated or apoptotic cells—such as platelets, monocytes, erythrocytes, and endothelial cells—and their interaction with endothelial cells and platelets drives the coagulation pathway.^{345–347} Although only an indirect measure, plasma von Willebrand factor concentrations were higher in septic dogs than those in healthy control animals, suggesting that endothelial cell activation also occurs in canine sepsis.³⁴⁸ Interestingly, platelets enhanced endotoxin-induced equine monocyte TF activity *in vitro*³⁴¹—although whether microparticles derived from platelets or other sources play a role in the pathogenesis of equine sepsis *in vivo* remains unknown. Additional

work in an equine model of endotoxemia has shown that large volume resuscitation has no impact on coagulation parameters beyond the changes attributed to endotoxemia, providing useful additional data to inform the treatment of sepsis in this species.³⁴⁹

Whereas much research has been directed toward the inhibition of coagulation in sepsis, only activated protein C has shown any benefit in human clinical trials.^{350,351} The antithrombotic and anti-inflammatory properties of recombinant human activated protein C led to recommendations for its use in severe sepsis in 2004³⁵² and 2008,³⁵³ but a recent meta-analysis found no evidence in support of its administration in the treatment of severe sepsis or septic shock.³⁵⁴ Indeed, there is still debate about its mechanism of action and, as yet, little experience of its use in veterinary medicine, despite encouraging pharmacological data in experimental models.^{266,355,356}

In addition to the role of TF in initiating coagulation in the presence of inflammation, activated platelets and endothelial cells—as well as bacterial surfaces—also trigger the contact phase system, leading to the formation of kallikrein and bradykinin.^{357,358} Bradykinin in turn enhances vasodilatation and increases vascular permeability, as well as reducing platelet function³⁵⁹; kallikrein accelerates fibrinolysis by conversion of plasminogen to plasmin and causes additional activation of Factor XII, leading to stimulation of the classical complement pathway.^{360,361}

A final connection between coagulation and inflammation in sepsis has become apparent with the recent exploration of the role of “a disintegrin-like and metalloproteinase with thrombospondin type-I motifs-13” (ADAMTS-13). ADAMTS-13 is produced by the stellate (Ito) cells of the liver and acts to cleave ultra-large von Willebrand’s Factor (vWF) multimers into smaller multimers.³⁶² These ultra-large vWF multimers are released from endothelial stores after inflammation and lead to platelet activation and aggregation; the ensuing microthrombi further compromise tissue blood flow, leading to additional propagation of the pro-inflammatory state.³⁶³ Although not yet identified in clinical veterinary species, decreased plasma ADAMTS-13 activity is associated with a poor prognosis in human sepsis patients^{364–366}; decreased activity is attributed to both a diminution of hepatic production and an increase in breakdown by plasma proteases.³⁶⁶

The Compensatory Anti-Inflammatory Response Syndrome and Cell Death

Ongoing investigation of the molecular mechanisms of the SIRS response, as well as the notable failure of therapeutic blockade of proinflammatory mediators, led to the realization that the mortality associated with sepsis could not be explained solely by an uncontrollable “cytokine storm”.^{367,368} This resulted in the concept of an opposing “compensatory anti-inflammatory response syndrome” (CARS), thought to be an adaptive response to the excessive proinflammatory process in SIRS and sepsis.^{367,369} Whereas an appropriate balance is struck




Fig 4. Points of interaction of the coagulation cascade and sepsis. In the absence of inflammation, various anticoagulant mechanisms exist to prevent activation of coagulation. Thrombomodulin and endothelial protein C receptor interact with thrombin to activate protein C (aPC), which in turn interacts with protein S to inactivate factor V. Endothelial cells also express tissue factor pathway inhibitor (TFPI) and antithrombin on their luminal surface and secrete tissue plasminogen activator (tPA). However, the majority of these anticoagulant factors are negative acute phase proteins and thus plasma concentrations falling during sepsis. Furthermore, proinflammatory cytokines released during sepsis activate endothelial cells, platelets and circulating white blood cells, which begin to express tissue factor (TF) and lose surface anticoagulant proteins. Microparticle formation is increased, together with expression of functional adhesion molecules on platelets, microparticles, monocytes and endothelial cells. Plasma antithrombin, aPC, protein S, and TFPI concentrations fall; ADAMTS-13 activity also drops, leading to greater persistence of ultralarge von Willebrand factor, which further enhances platelet and microparticle activation and adhesion. After triggering of the contact activation (TF) pathway, the interaction of microparticles with endothelial cells and platelets further drives the coagulation pathway. The presence of coagulation pathway components (TF-VIIa complex, Xa, Va, thrombin) and other circulating proteases activates endothelial protease-activated receptors (PARs), which propagate the inflammatory process by stimulating the further release of mediators (TNF α , IL-6 and CXCL-8), platelet activation and cell death. Triggering of the contact phase system activates factor XII (Hageman factor), leading to the formation of kallikrein (KK) and bradykinin (BK).

in the vast majority of instances of host-defense challenge, this is lost in sepsis—leading either to an uncontrolled proinflammatory reaction to infection, resulting

in organ dysfunction, an undesirable compromise of the immune system permitting opportunistic infection (the “second hit”), or a combination of both.³⁶⁷

Rather than a sequential or compensatory change, as was initially proposed, SIRS and CARS appear to occur simultaneously, and act to balance the host's need to maintain defense while minimizing self-induced tissue damage.³⁷⁰ Serum concentrations of both pro- and anti-inflammatory mediators increase early on in sepsis³⁷⁰; likewise, concentrations of a variety of both types of mediators (eg, IL-6, IRAP-1) are predictive of septic morbidity or mortality.^{370–372} Thus the changes in cytokine profile are both dynamic and heterogeneous, indicating that prescriptive immunomodulatory therapies are unlikely to meet with success.^{370,373} Furthermore, apoptosis of lymphocytes, hepatocytes, gastrointestinal epithelial cells and endothelial cells is increased, whereas that of neutrophils is decreased.^{374,375} Neutrophil function is also altered: both the migration of neutrophils to infected tissues^{376,377} and their antimicrobial function³⁷⁸ is diminished; moreover, peritoneal neutrophils are a potential source of IL-10, suppressing inflammatory monocytes in a model of polymicrobial sepsis.³⁷⁹ Although monocyte survival appears unaltered, sepsis results in the production of molecules such as IL-1 receptor associated kinase M (IRAK-M), MyD88 short variant (MyD88s), and A20-binding inhibitor of NF- κ B activation3 (ABIN-3), which reduce activation of the NF- κ B signaling pathway and therefore dampen the response to PAMP recognition.^{380,381} Monocyte function is impaired in sepsis, with decreased expression of the MHC class II molecule human leukocyte antigen-DR (HLA-DR)^{382,383} and decreased inflammasome expression.¹⁴⁶ Longitudinal observational studies in human patients indicate that a failure to regain >70% normal monocytic expression of HLA-DR is associated with an increased risk of secondary bacterial infection and decreased survival.³⁸⁴ In addition to these changes in antigen-presenting function of monocytes, dendritic cells also undergo increased apoptosis in sepsis, further impairing the host's ability to respond to pathogens.³⁸⁵

Although apoptosis, or type 1 programmed cell death (PCD), is responsible for the majority of immune cell death in sepsis and is implicated in immunoparalysis, alternative pathways also play a role.^{386,387} Autophagy is a cellular mechanism that primarily acts as a cytoplasmic “clean-up” process, as well as assisting in delivery of proteins to antigen presentation pathways; however, it may also mediate type II PCD and interact with apoptosis.^{388,389} Much current interest has been directed toward the role of autophagy in trauma and sepsis, although whether it acts in a cytoprotective role or as a mechanism of PCD, or both, remains unclear.^{386,390} A final mechanism of interest in the pathogenesis of sepsis is pyroptosis, a term used to describe the process of caspase-1-mediated PCD, which is distinct from death mediated by the apoptotic caspases 3, 6, and 8.³⁹¹ Although some question whether pyroptosis is truly a unique cell death mechanism, or simply a special case of apoptosis or necrosis (oncosis),³⁹² it is characterized by rapid plasma membrane rupture and release of proinflamma-

tory intracellular contents, some of which may act as DAMPs.³⁹³ One of the targets of caspase-1 during sepsis is the glycolytic pathway.³⁹⁴

A number of studies also suggest an important role for apoptosis in the pathogenesis of SIRS, sepsis, and infectious disease in veterinary species. Apoptotic cells were observed in the liver, kidney, thymus, stomach, and lymphocyte population of endotoxemic piglets³⁹⁵ and the primary and secondary lymphoid organs of pigs infected with classical swine fever virus.³⁹⁶ LPS and TNF- α both induced apoptosis of bovine glomerular endothelial cells, modeling a potential pathomechanism of acute renal failure in Gram-negative sepsis³⁹⁷; this phenomenon could be potentially inhibited by glucocorticoids *in vitro*.³⁹⁸ *Haemophilus somnus*, a Gram-negative pathogen of cattle that causes sepsis and vasculitis, induces caspases 3 and 8, and subsequent apoptosis, of endothelial cells *in vitro*³⁹⁹; furthermore, the bacterium stimulates platelets that are in turn able to induce endothelial cell apoptosis by a contact-dependent mechanism involving the activation of caspases 8 and 9 and the synthesis of reactive oxygen species.⁴⁰⁰ The activity of matrix metalloproteinases 2 and 9 and the expression of phosphorylated Paxillin showed positive correlation with cardiomyocyte apoptosis in an ovine model of endotoxemic shock,⁴⁰¹ whereas apoptosis of peripheral blood mononuclear cells and splenocytes of sheep infected with bluetongue virus was thought to contribute to immunosuppression in this disease.⁴⁰² Ileal epithelial apoptosis was documented in one feline model of endotoxemia,⁴⁰³ although a second demonstrated lymphocyte apoptosis in the spleen and Peyer's patches.³³⁴ Apoptosis of T cells also contributes to the immunosuppression characteristic of feline immunodeficiency virus infection.⁴⁰⁴ Finally, apoptosis of intestinal epithelial cells was observed in a canine model of sepsis induced by the intravenous infusion of *E. coli*.⁴⁰⁵

Regulatory T Cells and Sepsis

One of the key features of the adaptive immune system is its ability to generate antigen-specific receptors with an enormous diversity of specificities, some of which may recognize host-derived epitopes.^{406,407} The majority of potentially autoaggressive thymocytes are deleted in the thymic medulla in a process called negative selection.^{408,409} However, this process of central tolerance is imperfect and underlines the importance of a number of peripheral tolerance mechanisms, including clonal deletion, functional inactivation (anergy), and phenotypic skewing.^{410–412} Although a subject of debate for many years, a population of Tregs is now known to play a major role in peripheral tolerance, complementing the preceding (intrinsic) mechanisms.^{413–415} Little is currently known about tolerance mechanisms in veterinary species,⁴¹⁶ but Tregs have been identified in cats,^{417–423} dogs,^{424–433} pigs,^{434–442} cows,^{443–446} sheep,^{447,448} and horses.^{449–454}

Both naturally occurring and peripherally induced Tregs have been characterized—the former the product of a pathway of thymic differentiation called altered

negative selection and the latter the product of peripheral activation of conventional T cells in the context of an environment rich in transforming growth factor β (TGF- β) or IL-10.^{455–457} Naturally occurring Tregs have been identified by their constitutive expression of the IL-2 receptor α chain (CD25) and Forkhead box P3 (FOXP3), a transcription factor that plays a pivotal role in both their ontogeny and peripheral function. FOXP3 acts to stabilize the Treg transcriptome by repressing a number of pro-inflammatory and growth-promoting genes—for example, *IL-2* and *IFNG*—while activating others encoding key molecules involved in Treg function—for example, *CTLA4* and *CD25*.^{458,459}

Naturally occurring Tregs are known to interact with cells of both the innate and adaptive immune systems—including monocytes, macrophages, natural killer cells, neutrophils, mast cells, dendritic cells, and both T and B cells—generally mediating a suppressive function to prevent the development of autoaggressive responses and maintain the population of peripheral CD4⁺ T cells, thus contributing to immune system homeostasis.^{456,460,461} Tregs are also known to express a variety of TLRs, stimulation of which has augmented or abolished regulatory function in various studies.^{462–465} The molecular mechanisms of immune suppression mediated by naturally occurring Tregs have not been fully elucidated, but involve cell contact-dependent interactions, induction of cell death, and secretion of immunosuppressive cytokines, including IL-10 and TGF- β .^{416,456}

Various studies have documented increased proportions of Tregs in human sepsis during the phase of immunoparalysis,^{466–469} but the role of this change remains unclear because depletion of Tregs in murine models of sepsis has yielded variable conclusions between models, either improving, enhancing, or bearing no influence on mortality.^{470–472} Given the ability of Tregs to induce the alternative activation pathway of macrophages⁴⁷³ and to inhibit the LPS-induced survival of monocytes through a proapoptotic mechanism involving the Fas/FasL pathway,⁴⁷⁴ they may make a potentially significant contribution to immune system dysfunction in human septic patients,^{468,469,475,476} although this is still a controversial area.⁴⁷⁷

Additional research is required to elucidate the role of Tregs in sepsis, as well as to identify their mechanisms of action in this context. Moreover, to the authors' knowledge no veterinary studies have interrogated Treg number or function in septic patients to date. Pharmacological manipulation of Treg activity continues to be explored experimentally and may eventually translate to clinical cases; however, therapeutic interventions to alter the resistance of immune cells to Treg suppression may prove an equally valid alternative approach.^{478,479}

Conclusions

As is apparent from Figure 1, a broad concept of the immunopathological mechanisms underlying sepsis is now generally accepted. However, many of the molecular details are both complex and incompletely

elucidated. Ongoing research has also indicated the existence of multiple redundant pathways within the innate immune response, potentially explaining the failure of many highly selective therapeutic interventions.^{14,19}

A single “magic bullet” for the treatment of sepsis is highly unlikely to exist: an individually tailored set of therapies, based on point-of-care assessment of the immunopathological status of the patient, is the likely future—albeit a distant one.³⁷³ Various human studies have resulted in the publication of consensus statements regarding the treatment of sepsis,^{352,353} but current veterinary evidence to substantiate these interventions is thin on the ground. What has become evident is that outside the experimental laboratory, the heterogeneous nature of the septic patient population means that clinical trials must be carefully designed to obtain meaningful data.^{480–482} Ongoing basic research into the immunopathology of sepsis in clinical veterinary species is equally important, to elucidate the underlying molecular mechanisms and thus direct clinical studies to those aspects of disease likely to benefit the greatest number of patients.

Acknowledgments

The authors acknowledge funding from the Biotechnology and Biological Sciences Research Council, Novartis Animal Health, the European College of Veterinary Internal Medicine, and the American College of Veterinary Emergency and Critical Care for work on canine regulatory T cells and sepsis.

References

1. Bone RC, Balk RA, Cerra FB, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest* 1992;101:1644–1655.
2. Hotchkiss RS, Karl IE. The pathophysiology and treatment of sepsis. *N Engl J Med* 2003;348:138–150.
3. Levy MM, Fink MP, Marshall JC, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med* 2003;31:1250–1256.
4. Levy MM, Fink MP, Marshall JC, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Intensive Care Med* 2003;29:530–538.
5. Vincent JL, Sakr Y, Sprung CL, et al. Sepsis in European intensive care units: Results of the SOAP study. *Crit Care Med* 2006;34:344–353.
6. Bone RC. The sepsis syndrome. Definition and general approach to management. *Clin Chest Med* 1996;17:175–181.
7. Angus DC, Linde-Zwirble WT, Lidicker J, et al. Epidemiology of severe sepsis in the United States: Analysis of incidence, outcome, and associated costs of care. *Crit Care Med* 2001;29:1303–1310.
8. Martin GS, Mannino DM, Eaton S, et al. The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med* 2003;348:1546–1554.
9. Brady CA, Otto CM. Systemic inflammatory response syndrome, sepsis, and multiple organ dysfunction. *Vet Clin North Am Small Anim Pract* 2001;31:1147–1162, v-vi.

10. Hoffman AM, Staempfli HR, Willan A. Prognostic variables for survival of neonatal foals under intensive care. *J Vet Intern Med* 1992;6:89–95.
11. Declue AE, Delgado C, Chang CH, et al. Clinical and immunologic assessment of sepsis and the systemic inflammatory response syndrome in cats. *J Am Vet Med Assoc* 2011;238:890–897.
12. Hurcombe SD, Toribio RE, Slovis NM, et al. Calcium regulating hormones and serum calcium and magnesium concentrations in septic and critically ill foals and their association with survival. *J Vet Intern Med* 2009;23:335–343.
13. Rau S, Kohn B, Richter C, et al. Plasma interleukin-6 response is predictive for severity and mortality in canine systemic inflammatory response syndrome and sepsis. *Vet Clin Pathol* 2007;36:253–260.
14. Chong DL, Sriskandan S. Pro-inflammatory mechanisms in sepsis. *Contrib Microbiol* 2011;17:86–107.
15. Cinel I, Opal SM. Molecular biology of inflammation and sepsis: A primer. *Crit Care Med* 2009;37:291–304.
16. Cohen J. The immunopathogenesis of sepsis. *Nature* 2002;420:885–891.
17. Ivady B, Beres BJ, Szabo D. Recent advances in sepsis research: Novel biomarkers and therapeutic targets. *Curr Med Chem* 2011;18:3211–3225.
18. Remick DG. Pathophysiology of sepsis. *Am J Pathol* 2007;170:1435–1444.
19. Sriskandan S, Altmann DM. The immunology of sepsis. *J Pathol* 2008;214:211–223.
20. Stearns-Kurosawa DJ, Osuchowski MF, Valentine C, et al. The pathogenesis of sepsis. *Annu Rev Pathol* 2011;6:19–48.
21. Ganz T, Weiss J. Antimicrobial peptides of phagocytes and epithelia. *Semin Hematol* 1997;34:343–354.
22. Hoffmann JA, Kafatos FC, Janeway CA, et al. Phylogenetic perspectives in innate immunity. *Science* 1999;284:1313–1318.
23. Zasloff M. Antimicrobial peptides in health and disease. *N Engl J Med* 2002;347:1199–1200.
24. Brogden KA, Ackermann M, McCray PB Jr, et al. Antimicrobial peptides in animals and their role in host defences. *Int J Antimicrob Agents* 2003;22:465–478.
25. Linde A, Ross CR, Davis EG, et al. Innate immunity and host defense peptides in veterinary medicine. *J Vet Intern Med* 2008;22:247–265.
26. Shai Y. Mechanism of the binding, insertion and destabilization of phospholipid bilayer membranes by alpha-helical antimicrobial and cell non-selective membrane-lytic peptides. *Biochim Biophys Acta* 1999;1462:55–70.
27. Wilmes M, Cammue BP, Sahl HG, et al. Antibiotic activities of host defense peptides: More to it than lipid bilayer perturbation. *Nat Prod Rep* 2011;28:1350–1358.
28. Ali AS, Townes CL, Hall J, et al. Maintaining a sterile urinary tract: The role of antimicrobial peptides. *J Urol* 2009;182:21–28.
29. Diamond G, Beckloff N, Ryan LK. Host defense peptides in the oral cavity and the lung: Similarities and differences. *J Dent Res* 2008;87:915–927.
30. Ham M, Kaunitz JD. Gastroduodenal mucosal defense. *Curr Opin Gastroenterol* 2008;24:665–673.
31. Laube DM, Yim S, Ryan LK, et al. Antimicrobial peptides in the airway. *Curr Top Microbiol Immunol* 2006;306:153–182.
32. Huttner KM, Brezinski-Caliguri DJ, Mahoney MM, et al. Antimicrobial peptide expression is developmentally regulated in the ovine gastrointestinal tract. *J Nutr* 1998;128:297S–299S.
33. Sang Y, Ortega MT, Blecha F, et al. Molecular cloning and characterization of three beta-defensins from canine testes. *Infect Immun* 2005;73:2611–2620.
34. Salaun B, de Saint-Vis B, Pacheco N, et al. CD208/dendritic cell-lysosomal associated membrane protein is a marker of normal and transformed type II pneumocytes. *Am J Pathol* 2004;164:861–871.
35. Matzinger P. Friendly and dangerous signals: Is the tissue in control? *Nat Immunol* 2007;8:11–13.
36. Frantz S, Ertl G, Bauersachs J. Toll-like receptor signaling in the ischemic heart. *Front Biosci* 2008;13:5772–5779.
37. Rakoff-Nahoum S, Paglino J, Eslami-Varzaneh F, et al. Recognition of commensal microflora by toll-like receptors is required for intestinal homeostasis. *Cell* 2004;118:229–241.
38. Davies D, Meade KG, Herath S, et al. Toll-like receptor and antimicrobial peptide expression in the bovine endometrium. *Reprod Biol Endocrinol* 2008;6:53.
39. Bianchi ME. DAMPs, PAMPs and alarmins: All we need to know about danger. *J Leukoc Biol* 2007;81:1–5.
40. Gallucci S, Matzinger P. Danger signals: SOS to the immune system. *Curr Opin Immunol* 2001;13:114–119.
41. Matzinger P. Tolerance, danger, and the extended family. *Annu Rev Immunol* 1994;12:991–1045.
42. Akira S, Uematsu S, Takeuchi O. Pathogen recognition and innate immunity. *Cell* 2006;124:783–801.
43. Miyake K. Innate immune sensing of pathogens and danger signals by cell surface Toll-like receptors. *Semin Immunol* 2007;19:3–10.
44. Chen GY, Tang J, Zheng P, et al. CD24 and Siglec-10 selectively repress tissue damage-induced immune responses. *Science* 2009;323:1722–1725.
45. Chun KH, Seong SY. CD14 but not MD2 transmit signals from DAMP. *Int Immunopharmacol* 2010;10:98–106.
46. Mollen KP, Anand RJ, Tsung A, et al. Emerging paradigm: Toll-like receptor 4-sentinel for the detection of tissue damage. *Shock* 2006;26:430–437.
47. Astakhova NM, Perelygin AA, Zharkikh AA, et al. Characterization of equine and other vertebrate TLR3, TLR7, and TLR8 genes. *Immunogenetics* 2009;61:529–539.
48. Chang JS, Russell GC, Jann O, et al. Molecular cloning and characterization of Toll-like receptors 1-10 in sheep. *Vet Immunol Immunopathol* 2009;127:94–105.
49. Ignacio G, Nordone S, Howard KE, et al. Toll-like receptor expression in feline lymphoid tissues. *Vet Immunol Immunopathol* 2005;106:229–237.
50. McGuire K, Jones M, Werling D, et al. Radiation hybrid mapping of all 10 characterized bovine Toll-like receptors. *Anim Genet* 2006;37:47–50.
51. Menzies M, Ingham A. Identification and expression of Toll-like receptors 1-10 in selected bovine and ovine tissues. *Vet Immunol Immunopathol* 2006;109:23–30.
52. Tirumurugan KG, Dhanasekaran S, Raj GD, et al. Differential expression of toll-like receptor mRNA in selected tissues of goat (*Capra hircus*). *Vet Immunol Immunopathol* 2010;133:296–301.
53. Turin L, Riva F. Toll-like receptor family in domestic animal species. *Crit Rev Immunol* 2008;28:513–538.
54. Werling D, Coffey TJ. Pattern recognition receptors in companion and farm animals The key to unlocking the door to animal disease? *Vet J* 2007;174:240–251.
55. Werling D, Jann OC, Offord V, et al. Variation matters: TLR structure and species-specific pathogen recognition. *Trends Immunol* 2009;30:124–130.
56. Williams DL, Ha T, Li C, et al. Modulation of tissue Toll-like receptor 2 and 4 during the early phases of polymicrobial sepsis correlates with mortality. *Crit Care Med* 2003;31:1808–1818.
57. Tsujimoto H, Ono S, Efron PA, et al. Role of Toll-like receptors in the development of sepsis. *Shock* 2008;29:315–321.
58. Sato S, Nomura F, Kawai T, et al. Synergy and cross-tolerance between Toll-like receptor (TLR) 2- and TLR4-mediated signaling pathways. *J Immunol* 2000;165:7096–7101.

59. Ford JW, McVicar DW. TREM and TREM-like receptors in inflammation and disease. *Curr Opin Immunol* 2009;21:38–46.
60. Kuroki K, Stoker AM, Sims HJ, et al. Expression of Toll-like receptors 2 and 4 in stifle joint synovial tissues of dogs with or without osteoarthritis. *Am J Vet Res* 2010;71:750–754.
61. Silva E, Leitao S, Henriques S, et al. Gene transcription of TLR2, TLR4, LPS ligands and prostaglandin synthesis enzymes are up-regulated in canine uteri with cystic endometrial hyperplasia-pyometra complex. *J Reprod Immunol* 2010;84:66–74.
62. Taraktsoglou M, Szalabska U, Magee DA, et al. Transcriptional profiling of immune genes in bovine monocyte-derived macrophages exposed to bacterial antigens. *Vet Immunol Immunopathol* 2011;140:130–139.
63. Gundersen Y, Vaagenes P, Thrane I, et al. Early time course of altered leukocyte response to lipopolysaccharide and peptidoglycan in porcine gunshot injury. *Acta Anaesthesiol Scand* 2008;52:1231–1237.
64. Deitschel SJ, Kerl ME, Chang CH, et al. Age-associated changes to pathogen-associated molecular pattern-induced inflammatory mediator production in dogs. *J Vet Emerg Crit Care (San Antonio)* 2010;20:494–502.
65. Benbarek H, Deby-Dupont G, Caudron I, et al. Interactions between lipopolysaccharides and blood factors on the stimulation of equine polymorphonuclear neutrophils. *Vet Immunol Immunopathol* 1998;64:313–322.
66. Declue AE, Johnson PJ, Day JL, et al. Pathogen associated molecular pattern motifs from Gram-positive and Gram-negative bacteria induce different inflammatory mediator profiles in equine blood. *Vet J* 2011;00:00–00.
67. Fowler BL, Axiak SM, Declue AE. Blunted pathogen-associated molecular pattern motif induced TNF, IL-6 and IL-10 production from whole blood in dogs with lymphoma. *Vet Immunol Immunopathol* 2011;144:167–171.
68. Ishii KJ, Koyama S, Nakagawa A, et al. Host innate immune receptors and beyond: Making sense of microbial infections. *Cell Host Microbe* 2008;3:352–363.
69. Medzhitov R, Janeway C, Jr. Innate immune recognition: Mechanisms and pathways. *Immunol Rev* 2000;173:89–97.
70. Zak DE, Aderem A. Systems biology of innate immunity. *Immunol Rev* 2009;227:264–282.
71. Kawai T, Akira S. Toll-like receptors and their crosstalk with other innate receptors in infection and immunity. *Immunity* 2011;34:637–650.
72. Koyama S, Ishii KJ, Kumar H, et al. Differential role of TLR- and RLR-signaling in the immune responses to influenza A virus infection and vaccination. *J Immunol* 2007;179:4711–4720.
73. Takahashi K, Kawai T, Kumar H, et al. Roles of caspase-8 and caspase-10 in innate immune responses to double-stranded RNA. *J Immunol* 2006;176:4520–4524.
74. Hashimoto C, Hudson KL, Anderson KV. The Toll gene of *Drosophila*, required for dorsal-ventral embryonic polarity, appears to encode a transmembrane protein. *Cell* 1988;52:269–279.
75. Gay NJ, Keith FJ. *Drosophila* Toll and IL-1 receptor. *Nature* 1991;351:355–356.
76. Brightbill HD, Libraty DH, Krutzik SR, et al. Host defense mechanisms triggered by microbial lipoproteins through Toll-like receptors. *Science* 1999;285:732–736.
77. Lemaitre B, Nicolas E, Michaut L, et al. The dorsoventral regulatory gene cassette spatzle/Toll/cactus controls the potent antifungal response in *Drosophila* adults. *Cell* 1996;86:973–983.
78. Medzhitov R, Preston-Hurlburt P, Janeway CA Jr. A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. *Nature* 1997;388:394–397.
79. Takeuchi O, Hoshino K, Kawai T, et al. Differential roles of TLR2 and TLR4 in recognition of gram-negative and gram-positive bacterial cell wall components. *Immunity* 1999;11:443–451.
80. Horng T, Barton GM, Medzhitov R. TIRAP: An adapter molecule in the Toll signaling pathway. *Nat Immunol* 2001;2:835–841.
81. Jenkins KA, Mansell A. TIR-containing adaptors in Toll-like receptor signalling. *Cytokine* 2010;49:237–244.
82. Medzhitov R, Preston-Hurlburt P, Kopp E, et al. MyD88 is an adaptor protein in the hToll/IL-1 receptor family signaling pathways. *Mol Cell* 1998;2:253–258.
83. O'Neill LA, Bowie AG. The family of five: TIR-domain-containing adaptors in Toll-like receptor signalling. *Nat Rev Immunol* 2007;7:353–364.
84. West AP, Koblansky AA, Ghosh S. Recognition and signaling by Toll-like receptors. *Annu Rev Cell Dev Biol* 2006;22:409–437.
85. Kawagoe T, Sato S, Matsushita K, et al. Sequential control of Toll-like receptor-dependent responses by IRAK1 and IRAK2. *Nat Immunol* 2008;9:684–691.
86. Sato S, Sanjo H, Takeda K, et al. Essential function for the kinase TAK1 in innate and adaptive immune responses. *Nat Immunol* 2005;6:1087–1095.
87. Aksoy E, Vanden Berghe W, Detienne S, et al. Inhibition of phosphoinositide 3-kinase enhances TRIF-dependent NF-kappa B activation and IFN-beta synthesis downstream of Toll-like receptor 3 and 4. *Eur J Immunol* 2005;35:2200–2209.
88. Liljeroos M, Vuolteenaho R, Morath S, et al. Bruton's tyrosine kinase together with PI 3-kinase are part of Toll-like receptor 2 multiprotein complex and mediate LTA induced Toll-like receptor 2 responses in macrophages. *Cell Signal* 2007;19:625–633.
89. Ojaniemi M, Glumoff V, Harju K, et al. Phosphatidylinositol 3-kinase is involved in Toll-like receptor 4-mediated cytokine expression in mouse macrophages. *Eur J Immunol* 2003;33:597–605.
90. Cabanski M, Steinmuller M, Marsh LM, et al. PKR regulates TLR2/TLR4-dependent signaling in murine alveolar macrophages. *Am J Respir Cell Mol Biol* 2008;38:26–31.
91. Lentschat A, Karahashi H, Michelsen KS, et al. Mastoparan, a G protein agonist peptide, differentially modulates TLR4- and TLR2-mediated signaling in human endothelial cells and murine macrophages. *J Immunol* 2005;174:4252–4261.
92. Skokos D, Nussenzweig MC. CD8- DCs induce IL-12-independent Th1 differentiation through Delta 4 Notch-like ligand in response to bacterial LPS. *J Exp Med* 2007;204:1525–1531.
93. Sabroe I, Parker LC, Dower SK, et al. The role of TLR activation in inflammation. *J Pathol* 2008;214:126–135.
94. Aldape MJ, Bryant AE, Katahira EJ, et al. Innate immune recognition of, and response to, *Clostridium sordellii*. *Anaerobe* 2010;16:125–130.
95. Netea MG, Kullberg BJ, Galama JM, et al. Non-LPS components of *Chlamydia pneumoniae* stimulate cytokine production through Toll-like receptor 2-dependent pathways. *Eur J Immunol* 2002;32:1188–1195.
96. Schwandner R, Dziarski R, Wesche H, et al. Peptidoglycan- and lipoteichoic acid-induced cell activation is mediated by toll-like receptor 2. *J Biol Chem* 1999;274:17406–17409.
97. Zahringer U, Lindner B, Inamura S, et al. TLR2 - promiscuous or specific? A critical re-evaluation of a receptor expressing apparent broad specificity. *Immunobiology* 2008;213:205–224.
98. Lorne E, Dupont H, Abraham E. Toll-like receptors 2 and 4: Initiators of non-septic inflammation in critical care medicine? *Intensive Care Med* 2010;36:1826–1835.
99. Takeuchi O, Kawai T, Muhlrath PF, et al. Discrimination of bacterial lipoproteins by Toll-like receptor 6. *Int Immunol* 2001;13:933–940.
100. Takeuchi O, Sato S, Horiuchi T, et al. Cutting edge: Role of Toll-like receptor 1 in mediating immune response to microbial lipoproteins. *J Immunol* 2002;169:10–14.

101. Hoshino K, Takeuchi O, Kawai T, et al. Cutting edge: Toll-like receptor 4 (TLR4)-deficient mice are hyporesponsive to lipopolysaccharide: Evidence for TLR4 as the Lps gene product. *J Immunol* 1999;162:3749–3752.
102. Lu YC, Yeh WC, Ohashi PS. LPS/TLR4 signal transduction pathway. *Cytokine* 2008;42:145–151.
103. Takeda K, Akira S. Toll-like receptors in innate immunity. *Int Immunol* 2005;17:1–14.
104. Park BS, Song DH, Kim HM, et al. The structural basis of lipopolysaccharide recognition by the TLR4-MD-2 complex. *Nature* 2009;458:1191–1195.
105. Andersen-Nissen E, Smith KD, Bonneau R, et al. A conserved surface on Toll-like receptor 5 recognizes bacterial flagellin. *J Exp Med* 2007;204:393–403.
106. Uematsu S, Fujimoto K, Jang MH, et al. Regulation of humoral and cellular gut immunity by lamina propria dendritic cells expressing Toll-like receptor 5. *Nat Immunol* 2008;9:769–776.
107. Liaudet L, Szabo C, Evgenov OV, et al. Flagellin from Gram-negative bacteria is a potent mediator of acute pulmonary inflammation in sepsis. *Shock* 2003;19:131–137.
108. Kumar H, Kawai T, Akira S. Pathogen recognition in the innate immune response. *Biochem J* 2009;420:1–16.
109. Herskovits AA, Auerbuch V, Portnoy DA. Bacterial ligands generated in a phagosome are targets of the cytosolic innate immune system. *PLoS Pathog* 2007;3:e51.
110. Yamamoto M, Sato S, Hemmi H, et al. Role of adaptor TRIF in the MyD88-independent Toll-like receptor signaling pathway. *Science* 2003;301:640–643.
111. Yarovinsky F, Zhang D, Andersen JF, et al. TLR11 activation of dendritic cells by a protozoan profilin-like protein. *Science* 2005;308:1626–1629.
112. Zhang D, Zhang G, Hayden MS, et al. A Toll-like receptor that prevents infection by uropathogenic bacteria. *Science* 2004;303:1522–1526.
113. Carneiro LA, Magalhaes JG, Tattoli I, et al. Nod-like proteins in inflammation and disease. *J Pathol* 2008;214:136–148.
114. Chamaillard M, Hashimoto M, Horie Y, et al. An essential role for NOD1 in host recognition of bacterial peptidoglycan containing diaminopimelic acid. *Nat Immunol* 2003;4:702–707.
115. Girardin SE, Boneca IG, Viala J, et al. Nod2 is a general sensor of peptidoglycan through muramyl dipeptide (MDP) detection. *J Biol Chem* 2003;278:8869–8872.
116. Jones JD, Dangl JL. The plant immune system. *Nature* 2006;444:323–329.
117. Kanneganti TD, Lamkanfi M, Nunez G. Intracellular NOD-like receptors in host defense and disease. *Immunity* 2007;27:549–559.
118. Franchi L, Warner N, Viani K, et al. Function of Nod-like receptors in microbial recognition and host defense. *Immunol Rev* 2009;227:106–128.
119. Inohara N, Chamaillard M, McDonald C, et al. NOD-LRR proteins: Role in host-microbial interactions and inflammatory disease. *Annu Rev Biochem* 2005;74:355–383.
120. Moore CB, Bergstralh DT, Duncan JA, et al. NLRX1 is a regulator of mitochondrial antiviral immunity. *Nature* 2008;451:573–577.
121. Barnich N, Aguirre JE, Reinecker HC, et al. Membrane recruitment of NOD2 in intestinal epithelial cells is essential for nuclear factor-kappaB activation in muramyl dipeptide recognition. *J Cell Biol* 2005;170:21–26.
122. Kufer TA, Kremmer E, Adam AC, et al. The pattern-recognition molecule Nod1 is localized at the plasma membrane at sites of bacterial interaction. *Cell Microbiol* 2008;10:477–486.
123. Marina-Garcia N, Franchi L, Kim YG, et al. Pannexin-1-mediated intracellular delivery of muramyl dipeptide induces caspase-1 activation via cryopyrin/NLRP3 independently of Nod2. *J Immunol* 2008;180:4050–4057.
124. Ratner AJ, Aguilar JL, Shchepetov M, et al. Nod1 mediates cytoplasmic sensing of combinations of extracellular bacteria. *Cell Microbiol* 2007;9:1343–1351.
125. Viala J, Chaput C, Boneca IG, et al. Nod1 responds to peptidoglycan delivered by the *Helicobacter pylori* cag pathogenicity island. *Nat Immunol* 2004;5:1166–1174.
126. Hasegawa M, Fujimoto Y, Lucas PC, et al. A critical role of RICK/RIP2 polyubiquitination in Nod-induced NF-kappaB activation. *EMBO J* 2008;27:373–383.
127. Inohara N, Koseki T, del Peso L, et al. Nod1, an Apaf-1-like activator of caspase-9 and nuclear factor-kappaB. *J Biol Chem* 1999;274:14560–14567.
128. Kim JY, Omori E, Matsumoto K, et al. TAK1 is a central mediator of NOD2 signaling in epidermal cells. *J Biol Chem* 2008;283:137–144.
129. Ogura Y, Inohara N, Benito A, et al. Nod2, a Nod1/Apaf-1 family member that is restricted to monocytes and activates NF-kappaB. *J Biol Chem* 2001;276:4812–4818.
130. Strober W, Watanabe T. NOD2, an intracellular innate immune sensor involved in host defense and Crohn's disease. *Mucosal Immunol* 2011;4:484–495.
131. Masumoto J, Yang K, Varambally S, et al. Nod1 acts as an intracellular receptor to stimulate chemokine production and neutrophil recruitment *in vivo*. *J Exp Med* 2006;203:203–213.
132. Cartwright N, Murch O, McMaster SK, et al. Selective NOD1 agonists cause shock and organ injury/dysfunction *in vivo*. *Am J Respir Crit Care Med* 2007;175:595–603.
133. Lamkanfi M, Dixit VM. Inflammasomes: Guardians of cytosolic sanctity. *Immunol Rev* 2009;227:95–105.
134. Lamkanfi M, Dixit VM. The inflammasomes. *PLoS Pathog* 2009;5:e1000510.
135. Martinon F, Burns K, Tschopp J. The inflammasome: A molecular platform triggering activation of inflammatory caspases and processing of proIL-beta. *Mol Cell* 2002;10:417–426.
136. Franchi L, Amer A, Body-Malapel M, et al. Cytosolic flagellin requires Ipaf for activation of caspase-1 and interleukin 1beta in salmonella-infected macrophages. *Nat Immunol* 2006;7:576–582.
137. Martinon F, Petrilli V, Mayor A, et al. Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature* 2006;440:237–241.
138. Sutterwala FS, Mijares LA, Li L, et al. Immune recognition of *Pseudomonas aeruginosa* mediated by the IPAF/NLRC4 inflammasome. *J Exp Med* 2007;204:3235–3245.
139. Tsuji NM, Tsutsui H, Seki E, et al. Roles of caspase-1 in *Listeria* infection in mice. *Int Immunol* 2004;16:335–343.
140. Mariathasan S, Weiss DS, Dixit VM, et al. Innate immunity against *Francisella tularensis* is dependent on the ASC/caspase-1 axis. *J Exp Med* 2005;202:1043–1049.
141. Sarkar A, Hall MW, Exline M, et al. Caspase-1 regulates *Escherichia coli* sepsis and splenic B cell apoptosis independently of interleukin-1beta and interleukin-18. *Am J Respir Crit Care Med* 2006;174:1003–1010.
142. Saleh M, Mathison JC, Wolinski MK, et al. Enhanced bacterial clearance and sepsis resistance in caspase-12-deficient mice. *Nature* 2006;440:1064–1068.
143. Saleh M, Vaillancourt JP, Graham RK, et al. Differential modulation of endotoxin responsiveness by human caspase-12 polymorphisms. *Nature* 2004;429:75–79.
144. Li P, Allen H, Banerjee S, et al. Mice deficient in IL-1 beta-converting enzyme are defective in production of mature IL-1 beta and resistant to endotoxic shock. *Cell* 1995;80:401–411.
145. Mariathasan S, Newton K, Monack DM, et al. Differential activation of the inflammasome by caspase-1 adaptors ASC and Ipaf. *Nature* 2004;430:213–218.
146. Fahy RJ, Exline MC, Gavrilin MA, et al. Inflammasome mRNA expression in human monocytes during early septic shock. *Am J Respir Crit Care Med* 2008;177:983–988.

147. Kempster SL, Belteki G, Forhead AJ, et al. Developmental control of the Nlrp6 inflammasome and a substrate, IL-18, in mammalian intestine. *Am J Physiol Gastrointest Liver Physiol* 2011;300:G253–G263.
148. Chen H, Li C, Fang M, et al. Understanding *Haemophilus parasuis* infection in porcine spleen through a transcriptomics approach. *BMC Genomics* 2009;10:64.
149. Li R, Zhang A, Chen B, et al. Response of swine spleen to *Streptococcus suis* infection revealed by transcription analysis. *BMC Genomics* 2010;11:556.
150. Swerdlow MP, Kennedy DR, Kennedy JS, et al. Expression and function of TLR2, TLR4, and Nod2 in primary canine colonic epithelial cells. *Vet Immunol Immunopathol* 2006;114:313–319.
151. Asahina Y, Yoshioka N, Kano R, et al. Full-length cDNA cloning of Toll-like receptor 4 in dogs and cats. *Vet Immunol Immunopathol* 2003;96:159–167.
152. Hashimoto M, Asahina Y, Sano J, et al. Cloning of canine Toll-like receptor 9 and its expression in dog tissues. *Vet Immunol Immunopathol* 2005;106:159–163.
153. Ishii M, Hashimoto M, Oguma K, et al. Molecular cloning and tissue expression of canine Toll-like receptor 2 (TLR2). *Vet Immunol Immunopathol* 2006;110:87–95.
154. Okui Y, Kano R, Maruyama H, et al. Cloning of canine Toll-like receptor 7 gene and its expression in dog tissues. *Vet Immunol Immunopathol* 2008;121:156–160.
155. Franchini M, Zini E, Osto M, et al. Feline pancreatic islet-like clusters and insulin producing cells express functional Toll-like receptors (TLRs). *Vet Immunol Immunopathol* 2010;138:70–78.
156. Lehman TL, O'Halloran KP, Fallon SA, et al. Altered bone marrow dendritic cell cytokine production to Toll-like receptor and CD40 ligation during chronic feline immunodeficiency virus infection. *Immunology* 2009;126:405–412.
157. Kwong LS, Parsons R, Patterson R, et al. Characterisation of antibodies to bovine Toll-like receptor (TLR)-2 and cross-reactivity with ovine TLR2. *Vet Immunol Immunopathol* 2011;139:313–318.
158. Dhanasekaran S, Kannan TA, Dhinakar Raj G, et al. Differential expression of Toll-like receptor mRNA in corneal epithelium of ruminants. *Vet Ophthalmol* 2010;13:270–274.
159. Bridger PS, Mohr M, Stamm I, et al. Primary bovine colonic cells: A model to study strain-specific responses to *Escherichia coli*. *Vet Immunol Immunopathol* 2010;137:54–63.
160. Farhat K, Riekenberg S, Jung G, et al. Identification of full length bovine TLR1 and functional characterization of lipopeptide recognition by bovine TLR2/1 heterodimer. *Vet Res* 2010;41:34.
161. Seabury CM, Seabury PM, Decker JE, et al. Diversity and evolution of 11 innate immune genes in *Bos taurus taurus* and *Bos taurus indicus* cattle. *Proc Natl Acad Sci USA* 2010;107:151–156.
162. Arsenaault RJ, Jalal S, Babiuk LA, et al. Kinome analysis of Toll-like receptor signaling in bovine monocytes. *J Recept Signal Transduct Res* 2009;29:299–311.
163. Herath S, Lilly ST, Santos NR, et al. Expression of genes associated with immunity in the endometrium of cattle with disparate postpartum uterine disease and fertility. *Reprod Biol Endocrinol* 2009;7:55.
164. Mucha R, Bhide MR, Chakurkar EB, et al. Toll-like receptors TLR1, TLR2 and TLR4 gene mutations and natural resistance to *Mycobacterium avium* subsp. paratuberculosis infection in cattle. *Vet Immunol Immunopathol* 2009;128:381–388.
165. Zhu J, Brownlie R, Liu Q, et al. Characterization of bovine Toll-like receptor 8: Ligand specificity, signaling essential sites and dimerization. *Mol Immunol* 2009;46:978–990.
166. Jann OC, Werling D, Chang JS, et al. Molecular evolution of bovine Toll-like receptor 2 suggests substitutions of functional relevance. *BMC Evol Biol* 2008;8:288.
167. Goldammer T, Zerbe H, Molenaar A, et al. Mastitis increases mammary mRNA abundance of beta-defensin 5, Toll-like-receptor 2 (TLR2), and TLR4 but not TLR9 in cattle. *Clin Diagn Lab Immunol* 2004;11:174–185.
168. Schneberger D, Lewis D, Caldwell S, et al. Expression of Toll-like receptor 9 in lungs of pigs, dogs and cattle. *Int J Exp Pathol* 2011;92:1–7.
169. Jann OC, King A, Corrales NL, et al. Comparative genomics of Toll-like receptor signalling in five species. *BMC Genomics* 2009;10:216.
170. Mikula I Jr, Mikula I Sr, Characterization of ovine Toll-like receptor 9 protein coding region, comparative analysis, detection of mutations and maedi visna infection. *Dev Comp Immunol* 2011;35:182–192.
171. Plain KM, Purdie AC, Begg DJ, et al. Toll-like receptor (TLR)6 and TLR1 differentiation in gene expression studies of Johne's disease. *Vet Immunol Immunopathol* 2010;137:142–148.
172. Byun SO, Zhou H, Hickford JG. Development of a simple typing method for the ovine Toll-like receptor 4 gene. *Vet Immunol Immunopathol* 2009;130:272–274.
173. Hillman NH, Moss TJ, Nitsos I, et al. Toll-like receptors and agonist responses in the developing fetal sheep lung. *Pediatr Res* 2008;63:388–393.
174. Raja A, Vignesh AR, Mary BA, et al. Sequence analysis of Toll-like receptor genes 1-10 of goat (*Capra hircus*). *Vet Immunol Immunopathol* 2011;140:252–258.
175. Tourais-Esteves I, Bernardet N, Lacroix-Lamande S, et al. Neonatal goats display a stronger TH1-type cytokine response to TLR ligands than adults. *Dev Comp Immunol* 2008;32:1231–1241.
176. Gornik K, Moore P, Figueiredo M, et al. Expression of Toll-like receptors 2, 3, 4, 6, 9, and MD-2 in the normal equine cornea, limbus, and conjunctiva. *Vet Ophthalmol* 2011;14:80–85.
177. Kwon S, Vandenplas ML, Figueiredo MD, et al. Differential induction of Toll-like receptor gene expression in equine monocytes activated by Toll-like receptor ligands or TNF-alpha. *Vet Immunol Immunopathol* 2010;138:213–217.
178. Berndt A, Derksen FJ, Venta PJ, et al. Elevated amount of Toll-like receptor 4 mRNA in bronchial epithelial cells is associated with airway inflammation in horses with recurrent airway obstruction. *Am J Physiol Lung Cell Mol Physiol* 2007;292:L936–943.
179. Singh Suri S, Janardhan KS, Parbhakar O, et al. Expression of Toll-like receptor 4 and 2 in horse lungs. *Vet Res* 2006;37:541–551.
180. Werners AH, Bull S, Vendrig JC, et al. Genotyping of Toll-like receptor 4, myeloid differentiation factor 2 and CD-14 in the horse: An investigation into the influence of genetic polymorphisms on the LPS induced TNF-alpha response in equine whole blood. *Vet Immunol Immunopathol* 2006;111:165–173.
181. Schneberger D, Caldwell S, Suri SS, et al. Expression of Toll-like receptor 9 in horse lungs. *Anat Rec (Hoboken)* 2009;292:1068–1077.
182. Uenishi H, Shinkai H, Morozumi T, et al. Polymorphisms in pattern recognition receptors and their relationship to infectious disease susceptibility in pigs. *BMC Proc* 2011;5(Suppl 4):S27.
183. Shinkai H, Suzuki R, Akiba M, et al. Porcine Toll-like receptors: Recognition of *Salmonella enterica* serovar Choleraesuis and influence of polymorphisms. *Mol Immunol* 2011;48:1114–1120.
184. Evren S, Loai Y, Antoon R, et al. Urinary bladder tissue engineering using natural scaffolds in a porcine model: Role of

Toll-like receptors and impact of biomimetic molecules. *Cells Tissues Organs* 2010;192:250–261.

185. Auray G, Facci MR, van Kessel J, et al. Differential activation and maturation of two porcine DC populations following TLR ligand stimulation. *Mol Immunol* 2010;47:2103–2111.

186. Morozumi T, Uenishi H. Polymorphism distribution and structural conservation in RNA-sensing Toll-like receptors 3, 7, and 8 in pigs. *Biochim Biophys Acta* 2009;1790:267–274.

187. Toka FN, Nfon CK, Dawson H, et al. Accessory-cell-mediated activation of porcine NK cells by Toll-like receptor 7 (TLR7) and TLR8 agonists. *Clin Vaccine Immunol* 2009;16:866–878.

188. Burkey TE, Skjolaas KA, Dritz SS, et al. Expression of porcine Toll-like receptor 2, 4 and 9 gene transcripts in the presence of lipopolysaccharide and *Salmonella enterica* serovars Typhimurium and Choleraesuis. *Vet Immunol Immunopathol* 2009;130:96–101.

189. Liu CH, Chung HC, Chang HL, et al. Expression of Toll-like receptor mRNA and cytokines in pigs infected with porcine reproductive and respiratory syndrome virus. *Vet Microbiol* 2009;136:266–276.

190. Chung HC, Chen CW, Hsieh BL, et al. Toll-like receptor expressions in porcine alveolar macrophages and dendritic cells in responding to poly IC stimulation and porcine reproductive and respiratory syndrome virus (PRRSV) infection. *Comp Immunol Microbiol Infect Dis* 2010;33:197–213.

191. Uenishi H, Shinkai H. Porcine Toll-like receptors: The front line of pathogen monitoring and possible implications for disease resistance. *Dev Comp Immunol* 2009;33:353–361.

192. Ibeagha-Awemu EM, Lee JW, Ibeagha AE, et al. Bacterial lipopolysaccharide induces increased expression of Toll-like receptor (TLR) 4 and downstream TLR signaling molecules in bovine mammary epithelial cells. *Vet Res* 2008;39:11.

193. Worku M, Morris A. Binding of different forms of lipopolysaccharide and gene expression in bovine blood neutrophils. *J Dairy Sci* 2009;92:3185–3193.

194. Cates EA, Connor EE, Mosser DM, et al. Functional characterization of bovine TIRAP and MyD88 in mediating bacterial lipopolysaccharide-induced endothelial NF-kappaB activation and apoptosis. *Comp Immunol Microbiol Infect Dis* 2009;32:477–490.

195. Gil LH, van Olphen AL, Mittal SK, et al. Modulation of PKR activity in cells infected by bovine viral diarrhoea virus. *Virus Res* 2006;116:69–77.

196. Sen A, Pruijssers AJ, Dermody TS, et al. The early interferon response to rotavirus is regulated by PKR and depends on MAVS/IPS-1, RIG-I, MDA-5, and IRF3. *J Virol* 2011;85:3717–3732.

197. Bougarn S, Cunha P, Harmache A, et al. Muramyl dipeptide synergizes with *Staphylococcus aureus* lipoteichoic acid to recruit neutrophils in the mammary gland and to stimulate mammary epithelial cells. *Clin Vaccine Immunol* 2010;17:1797–1809.

198. Whelehan CJ, Meade KG, Eckersall PD, et al. Experimental *Staphylococcus aureus* infection of the mammary gland induces region-specific changes in innate immune gene expression. *Vet Immunol Immunopathol* 2011;140:181–189.

199. Purdie AC, Plain KM, Begg DJ, et al. Candidate gene and genome-wide association studies of *Mycobacterium avium* subsp. paratuberculosis infection in cattle and sheep: A review. *Comp Immunol Microbiol Infect Dis* 2011;34:197–208.

200. Fenhammar J, Rundgren M, Forestier J, et al. Toll-like receptor 4 inhibitor TAK-242 attenuates acute kidney injury in endotoxemic sheep. *Anesthesiology* 2011;114:1130–1137.

201. Alvarez B, Revilla C, Chamorro S, et al. Molecular cloning, characterization and tissue expression of porcine Toll-like receptor 4. *Dev Comp Immunol* 2006;30:345–355.

202. Alves MP, Neuhaus V, Guzylack-Piriou L, et al. Toll-like receptor 7 and MyD88 knockdown by lentivirus-mediated RNA interference to porcine dendritic cell subsets. *Gene Ther* 2007;14:836–844.

203. Moue M, Tohno M, Shimazu T, et al. Toll-like receptor 4 and cytokine expression involved in functional immune response in an originally established porcine intestinal epithelial cell line. *Biochim Biophys Acta* 2008;1780:134–144.

204. Muneta Y, Uenishi H, Kikuma R, et al. Porcine TLR2 and TLR6: Identification and their involvement in *Mycoplasma hyopneumoniae* infection. *J Interferon Cytokine Res* 2003;23:583–590.

205. Shimosato T, Kitazawa H, Katoh S, et al. Swine Toll-like receptor 9(1) recognizes CpG motifs of human cell stimulant. *Biochim Biophys Acta* 2003;1627:56–61.

206. Shinkai H, Tanaka M, Morozumi T, et al. Biased distribution of single nucleotide polymorphisms (SNPs) in porcine Toll-like receptor 1 (TLR1), TLR2, TLR4, TLR5, and TLR6 genes. *Immunogenetics* 2006;58:324–330.

207. Shinkai H, Muneta Y, Suzuki K, et al. Porcine Toll-like receptor 1, 6, and 10 genes: Complete sequencing of genomic region and expression analysis. *Mol Immunol* 2006;43:1474–1480.

208. Thomas AV, Broers AD, Vandegaart HF, et al. Genomic structure, promoter analysis and expression of the porcine (*Sus scrofa*) TLR4 gene. *Mol Immunol* 2006;43:653–659.

209. Wassef A, Janardhan K, Pearce JW, et al. Toll-like receptor 4 in normal and inflamed lungs and other organs of pig, dog and cattle. *Histol Histopathol* 2004;19:1201–1208.

210. Miguel JC, Chen J, Van Alstine WG, et al. Expression of inflammatory cytokines and Toll-like receptors in the brain and respiratory tract of pigs infected with porcine reproductive and respiratory syndrome virus. *Vet Immunol Immunopathol* 2010;135:314–319.

211. Sang Y, Yang J, Ross CR, et al. Molecular identification and functional expression of porcine Toll-like receptor (TLR) 3 and TLR7. *Vet Immunol Immunopathol* 2008;125:162–167.

212. Jozaki K, Shinkai H, Tanaka-Matsuda M, et al. Influence of polymorphisms in porcine NOD2 on ligand recognition. *Mol Immunol* 2009;47:247–252.

213. Kojima-Shibata C, Shinkai H, Morozumi T, et al. Differences in distribution of single nucleotide polymorphisms among intracellular pattern recognition receptors in pigs. *Immunogenetics* 2009;61:153–160.

214. Luo R, Xiao S, Jiang Y, et al. Porcine reproductive and respiratory syndrome virus (PRRSV) suppresses interferon-beta production by interfering with the RIG-I signaling pathway. *Mol Immunol* 2008;45:2839–2846.

215. Tohno M, Shimazu T, Aso H, et al. Molecular cloning and functional characterization of porcine nucleotide-binding oligomerization domain-1 (NOD1) recognizing minimum agonists, meso-diaminopimelic acid and meso-lanthionine. *Mol Immunol* 2008;45:1807–1817.

216. Tohno M, Ueda W, Azuma Y, et al. Molecular cloning and functional characterization of porcine nucleotide-binding oligomerization domain-2 (NOD2). *Mol Immunol* 2008;45:194–203.

217. Tohno M, Shimazu T, Aso H, et al. Molecular cloning and functional characterization of porcine MyD88 essential for TLR signaling. *Cell Mol Immunol* 2007;4:369–376.

218. Kwon S, Gewirtz AT, Hurley DJ, et al. Disparities in TLR5 expression and responsiveness to flagellin in equine neutrophils and mononuclear phagocytes. *J Immunol* 2011;186:6263–6270.

219. Gold JR, Perkins GA, Erb HN, et al. Cytokine profiles of peripheral blood mononuclear cells isolated from septic and healthy neonatal foals. *J Vet Intern Med* 2007;21:482–488.

220. Lopes MA, Salter CE, Vandenplas ML, et al. Expression of inflammation-associated genes in circulating leukocytes

collected from horses with gastrointestinal tract disease. *Am J Vet Res* 2010;71:915–924.

221. Menzies-Gow NJ, Bailey SR, Stevens K, et al. Digital blood flow and plasma endothelin concentration in clinically endotoxemic horses. *Am J Vet Res* 2005;66:630–636.

222. Figueiredo MD, Vandenplas ML, Hurley DJ, et al. Differential induction of MyD88- and TRIF-dependent pathways in equine monocytes by Toll-like receptor agonists. *Vet Immunol Immunopathol* 2009;127:125–134.

223. Allenspach K, House A, Smith K, et al. Evaluation of mucosal bacteria and histopathology, clinical disease activity and expression of Toll-like receptors in German Shepherd Dogs with chronic enteropathies. *Vet Microbiol* 2010;146:326–335.

224. Tian X, Pascal G, Monget P. Evolution and functional divergence of NLRP genes in mammalian reproductive systems. *BMC Evol Biol* 2009;9:202.

225. Burgener IA, Jungi TW. Antibodies specific for human or murine Toll-like receptors detect canine leukocytes by flow cytometry. *Vet Immunol Immunopathol* 2008;124:184–191.

226. Chotimanukul S, Sirivaidyapong S. Differential expression of Toll-like receptor 4 (TLR4) in healthy and infected canine endometrium. *Theriogenology* 2011;76:1152–1161.

227. Abujamra AL, Spanjaard RA, Akinsheye I, et al. Leukemia virus long terminal repeat activates NF-kappaB pathway by a TLR3-dependent mechanism. *Virology* 2006;345:390–403.

228. Burgener IA, Konig A, Allenspach K, et al. Upregulation of Toll-like receptors in chronic enteropathies in dogs. *J Vet Intern Med* 2008;22:553–560.

229. House AK, Binns MM, Gregory SP, et al. Analysis of NOD1, NOD2, TLR1, TLR2, TLR4, TLR5, TLR6 and TLR9 genes in anal furunculosis of German Shepherd Dogs. *Tissue Antigens* 2009;73:250–254.

230. Kathrani A, House A, Catchpole B, et al. Polymorphisms in the TLR4 and TLR5 gene are significantly associated with inflammatory bowel disease in German Shepherd Dogs. *PLoS ONE* 2010;5:e15740.

231. Kathrani A, House A, Catchpole B, et al. Breed-independent Toll-like receptor 5 polymorphisms show association with canine inflammatory bowel disease. *Tissue Antigens* 2011;78:94–101.

232. Mikula I, Bhide M, Pastorekova S. Characterization of ovine TLR7 and TLR8 protein coding regions, detection of mutations and Maedi Visna virus infection. *Vet Immunol Immunopathol* 2010;138:51–59.

233. McCullough KC, Summerfield A. Targeting the porcine immune system—particulate vaccines in the 21st century. *Dev Comp Immunol* 2009;33:394–409.

234. Li J, Birkenheuer AJ, Marr HS, et al. Expression and function of triggering receptor expressed on myeloid cells-1 (TREM-1) on canine neutrophils. *Dev Comp Immunol* 2011;35:872–880.

235. Derive M, Massin F, Gibot S. Triggering receptor expressed on myeloid cells-1 as a new therapeutic target during inflammatory diseases. *Self Nonself* 2010;1:225–230.

236. Zhu J, Mohan C. Toll-like receptor signaling pathways—therapeutic opportunities. *Mediators Inflamm* 2010;2010:781235.

237. Yamamoto M, Takeda K. Current views of Toll-like receptor signaling pathways. *Gastroenterol Res Pract* 2010;2010:240365.

238. Lin Q, Li M, Fang D, et al. The essential roles of Toll-like receptor signaling pathways in sterile inflammatory diseases. *Int Immunopharmacol* 2011;11:1422–1432.

239. Damas P, Reuter A, Gysen P, et al. Tumor necrosis factor and interleukin-1 serum levels during severe sepsis in humans. *Crit Care Med* 1989;17:975–978.

240. Girardin E, Grau GE, Dayer JM, et al. Tumor necrosis factor and interleukin-1 in the serum of children with severe infectious purpura. *N Engl J Med* 1988;319:397–400.

241. Basoglu A, Sen I, Sevinc M, et al. Serum concentrations of tumor necrosis factor-alpha in neonatal calves with presumed septicemia. *J Vet Intern Med* 2004;18:238–241.

242. Otto CM, Drobatz KJ, Soter C. Endotoxemia and tumor necrosis factor activity in dogs with naturally occurring parvoviral enteritis. *J Vet Intern Med* 1997;11:65–70.

243. Bradley JR. TNF-mediated inflammatory disease. *J Pathol* 2008;214:149–160.

244. Naude PJ, den Boer JA, Luiten PG, et al. Tumor necrosis factor receptor cross-talk. *FEBS J* 2011;278:888–898.

245. Vinay DS, Kwon BS. The tumour necrosis factor/TNF receptor superfamily: Therapeutic targets in autoimmune diseases. *Clin Exp Immunol* 2011;164:145–157.

246. Micheau O, Lens S, Gaide O, et al. NF-kappaB signals induce the expression of c-FLIP. *Mol Cell Biol* 2001;21:5299–5305.

247. Tartaglia LA, Pennica D, Goeddel DV. Ligand passing: the 75-kDa tumor necrosis factor (TNF) receptor recruits TNF for signaling by the 55-kDa TNF receptor. *J Biol Chem* 1993;268:18542–18548.

248. Mark KS, Trickler WJ, Miller DW. Tumor necrosis factor-alpha induces cyclooxygenase-2 expression and prostaglandin release in brain microvessel endothelial cells. *J Pharmacol Exp Ther* 2001;297:1051–1058.

249. Pober JS, Bevilacqua MP, Mendrick DL, et al. Two distinct monokines, interleukin 1 and tumor necrosis factor, each independently induce biosynthesis and transient expression of the same antigen on the surface of cultured human vascular endothelial cells. *J Immunol* 1986;136:1680–1687.

250. Munro JM, Pober JS, Cotran RS. Tumor necrosis factor and interferon-gamma induce distinct patterns of endothelial activation and associated leukocyte accumulation in skin of *Papio anubis*. *Am J Pathol* 1989;135:121–133.

251. Sakaue Y, Nezu Y, Yanagisawa S, et al. Effects of continuous low-dose infusion of lipopolysaccharide on expression of E-selectin and intercellular adhesion molecule-1 messenger RNA and neutrophil accumulation in specific organs in dogs. *Am J Vet Res* 2005;66:1259–1266.

252. Miyamoto T, Fujinaga T, Yamashita K, et al. Changes of serum cytokine activities and other parameters in dogs with experimentally induced endotoxic shock. *Jpn J Vet Res* 1996;44:107–118.

253. LeMay DR, LeMay LG, Kluger MJ, et al. Plasma profiles of IL-6 and TNF with fever-inducing doses of lipopolysaccharide in dogs. *Am J Physiol* 1990;259:R126–132.

254. Otto CM, Rawlings CA. Tumor necrosis factor production in cats in response to lipopolysaccharide: An *in vivo* and *in vitro* study. *Vet Immunol Immunopathol* 1995;49:183–188.

255. MacKay RJ, Merritt AM, Zertuche JM, et al. Tumor necrosis factor activity in the circulation of horses given endotoxin. *Am J Vet Res* 1991;52:533–538.

256. Nieto JE, MacDonald MH, Braim AE, et al. Effect of lipopolysaccharide infusion on gene expression of inflammatory cytokines in normal horses *in vivo*. *Equine Vet J* 2009;41:717–719.

257. MacKay RJ, Lester GD. Induction of the acute-phase cytokine, hepatocyte-stimulating factor/interleukin 6, in the circulation of horses treated with endotoxin. *Am J Vet Res* 1992;53:1285–1289.

258. Huang H, Lavoie-Lamoureux A, Moran K, et al. IL-4 stimulates the expression of CXCL-8, E-selectin, VEGF, and inducible nitric oxide synthase mRNA by equine pulmonary artery endothelial cells. *Am J Physiol Lung Cell Mol Physiol* 2007;292:L1147–1154.

259. Barton MH, Collatos C. Tumor necrosis factor and interleukin-6 activity and endotoxin concentration in peritoneal fluid and blood of horses with acute abdominal disease. *J Vet Intern Med* 1999;13:457–464.

260. Stich AN, DeClue AE. Pathogen associated molecular pattern-induced TNF, IL-1beta, IL-6 and CXCL-8 production from feline whole blood culture. *Res Vet Sci* 2011;90:59–63.
261. Osterbur K, Whitehead Z, Sharp CR, et al. Plasma nitrate/nitrite concentrations in dogs with naturally developing sepsis and non-infectious forms of the systemic inflammatory response syndrome. *Vet Rec* 2011;169:554.
262. LeBlanc CJ, Horohov DW, Bauer JE, et al. Effects of dietary supplementation with fish oil on *in vivo* production of inflammatory mediators in clinically normal dogs. *Am J Vet Res* 2008;69:486–493.
263. Bevilacqua MP, Pober JS, Majeau GR, et al. Recombinant tumor necrosis factor induces procoagulant activity in cultured human vascular endothelium: Characterization and comparison with the actions of interleukin 1. *Proc Natl Acad Sci USA* 1986;83:4533–4537.
264. Cartwright N, McMaster SK, Sorrentino R, et al. Elucidation of Toll-like receptor and adapter protein signaling in vascular dysfunction induced by gram-positive *Staphylococcus aureus* or gram-negative *Escherichia coli*. *Shock* 2007;27:40–47.
265. Jimenez R, Belcher E, Sriskandan S, et al. Role of Toll-like receptors 2 and 4 in the induction of cyclooxygenase-2 in vascular smooth muscle. *Proc Natl Acad Sci USA* 2005;102:4637–4642.
266. Schouten M, Wiersinga WJ, Levi M, et al. Inflammation, endothelium, and coagulation in sepsis. *J Leukoc Biol* 2008;83:536–545.
267. Cavaillon JM, Adib-Conquy M, Fitting C, et al. Cytokine cascade in sepsis. *Scand J Infect Dis* 2003;35:535–544.
268. Dinarello CA. Proinflammatory and anti-inflammatory cytokines as mediators in the pathogenesis of septic shock. *Chest* 1997;112:321S–329S.
269. Stylianou E, Saklatvala J. Interleukin-1. *Int J Biochem Cell Biol* 1998;30:1075–1079.
270. Chaitanya GV, Franks SE, Cromer W, et al. Differential cytokine responses in human and mouse lymphatic endothelial cells to cytokines *in vitro*. *Lymphat Res Biol* 2010;8:155–164.
271. Dinarello CA. A clinical perspective of IL-1beta as the gatekeeper of inflammation. *Eur J Immunol* 2011;41:1203–1217.
272. Jawa RS, Anillo S, Huntoon K, et al. Analytic review: Interleukin-6 in surgery, trauma, and critical care: Part I: Basic science. *J Intensive Care Med* 2011;26:3–12.
273. Scheller J, Chalaris A, Schmidt-Arras D, et al. The pro- and anti-inflammatory properties of the cytokine interleukin-6. *Biochim Biophys Acta* 2011;1813:878–888.
274. Borrelli E, Roux-Lombard P, Grau GE, et al. Plasma concentrations of cytokines, their soluble receptors, and antioxidant vitamins can predict the development of multiple organ failure in patients at risk. *Crit Care Med* 1996;24:392–397.
275. Remick DG, Bolgos GR, Siddiqui J, et al. Six at six: Interleukin-6 measured 6 h after the initiation of sepsis predicts mortality over 3 days. *Shock* 2002;17:463–467.
276. Burton AB, Wagner B, Erb HN, et al. Serum interleukin-6 (IL-6) and IL-10 concentrations in normal and septic neonatal foals. *Vet Immunol Immunopathol* 2009;132:122–128.
277. Scumpia PO, Moldawer LL. Biology of interleukin-10 and its regulatory roles in sepsis syndromes. *Crit Care Med* 2005;33:S468–S471.
278. Oberholzer A, Oberholzer C, Moldawer LL. Interleukin-10: A complex role in the pathogenesis of sepsis syndromes and its potential as an anti-inflammatory drug. *Crit Care Med* 2002;30:S58–S63.
279. Seitz M, Loetscher P, Dewald B, et al. Interleukin-10 differentially regulates cytokine inhibitor and chemokine release from blood mononuclear cells and fibroblasts. *Eur J Immunol* 1995;25:1129–1132.
280. Kusske AM, Rongione AJ, Ashley SW, et al. Interleukin-10 prevents death in lethal necrotizing pancreatitis in mice. *Surgery* 1996;120:284–288; discussion 289.
281. Briassoulis G, Venkataraman S, Thompson A. Cytokines and metabolic patterns in pediatric patients with critical illness. *Clin Dev Immunol* 2010;2010:354047.
282. Pusterla N, Magdesian KG, Mapes S, et al. Expression of molecular markers in blood of neonatal foals with sepsis. *Am J Vet Res* 2006;67:1045–1049.
283. Kipar A, Meli ML, Failing K, et al. Natural feline coronavirus infection: Differences in cytokine patterns in association with the outcome of infection. *Vet Immunol Immunopathol* 2006;112:141–155.
284. Gao L, Flores C, Fan-Ma S, et al. Macrophage migration inhibitory factor in acute lung injury: Expression, biomarker, and associations. *Transl Res* 2007;150:18–29.
285. Calandra T, Roger T. Macrophage migration inhibitory factor: A regulator of innate immunity. *Nat Rev Immunol* 2003;3:791–800.
286. Bozza M, Satoskar AR, Lin G, et al. Targeted disruption of migration inhibitory factor gene reveals its critical role in sepsis. *J Exp Med* 1999;189:341–346.
287. Brenner T, Hofer S, Rosenhagen C, et al. Macrophage migration inhibitory factor (MIF) and manganese superoxide dismutase (MnSOD) as early predictors for survival in patients with severe sepsis or septic shock. *J Surg Res* 2010;164:e163–171.
288. Mitchell RA, Liao H, Chesney J, et al. Macrophage migration inhibitory factor (MIF) sustains macrophage proinflammatory function by inhibiting p53: Regulatory role in the innate immune response. *Proc Natl Acad Sci USA* 2002;99:345–350.
289. Stros M. HMGB proteins: Interactions with DNA and chromatin. *Biochim Biophys Acta* 2010;1799:101–113.
290. Yang H, Tracey KJ. Targeting HMGB1 in inflammation. *Biochim Biophys Acta* 2010;1799:149–156.
291. van Zoelen MA, Yang H, Florquin S, et al. Role of Toll-like receptors 2 and 4, and the receptor for advanced glycation end products in high-mobility group box 1-induced inflammation *in vivo*. *Shock* 2009;31:280–284.
292. Huang W, Tang Y, Li L. HMGB1, a potent proinflammatory cytokine in sepsis. *Cytokine* 2010;51:119–126.
293. Mitola S, Belleri M, Urbinati C, et al. Cutting edge: Extracellular high mobility group box-1 protein is a proangiogenic cytokine. *J Immunol* 2006;176:12–15.
294. Yu DH, Nho DH, Song RH, et al. High-mobility group box 1 as a surrogate prognostic marker in dogs with systemic inflammatory response syndrome. *J Vet Emerg Crit Care (San Antonio)* 2010;20:298–302.
295. Karlsson S, Pettila V, Tenhunen J, et al. HMGB1 as a predictor of organ dysfunction and outcome in patients with severe sepsis. *Intensive Care Med* 2008;34:1046–1053.
296. de Jong HK, van der Poll T, Wiersinga WJ. The systemic pro-inflammatory response in sepsis. *J Innate Immun* 2010;2:422–430.
297. Cray C, Zaias J, Altman NH. Acute phase response in animals: A review. *Comp Med* 2009;59:517–526.
298. Moshage H. Cytokines and the hepatic acute phase response. *J Pathol* 1997;181:257–266.
299. Gruys E, Toussaint MJ, Niewold TA, et al. Acute phase reaction and acute phase proteins. *J Zhejiang Univ Sci B* 2005;6:1045–1056.
300. Cecilian F, Giordano A, Spagnolo V. The systemic reaction during inflammation: The acute-phase proteins. *Protein Pept Lett* 2002;9:211–223.
301. Noursadeghi M, Pepys MB, Gallimore R, et al. Relationship of granulocyte colony stimulating factor with other acute phase reactants in man. *Clin Exp Immunol* 2005;140:97–100.

302. Ceron JJ, Eckersall PD, Martynez-Subiela S. Acute phase proteins in dogs and cats: Current knowledge and future perspectives. *Vet Clin Pathol* 2005;34:85–99.
303. Eckersall PD, Bell R. Acute phase proteins: Biomarkers of infection and inflammation in veterinary medicine. *Vet J* 2010;185:23–27.
304. Kjelgaard-Hansen M, Jacobsen S. Assay validation and diagnostic applications of major acute-phase protein testing in companion animals. *Clin Lab Med* 2011;31:51–70.
305. Tvarijonaviciute A, Eralp O, Kocaturk M, et al. Adiponectin and IGF-1 are negative acute phase proteins in a dog model of acute endotoxaemia. *Vet Immunol Immunopathol* 2011;140:147–151.
306. Castelli GP, Pognani C, Meisner M, et al. Procalcitonin and C-reactive protein during systemic inflammatory response syndrome, sepsis and organ dysfunction. *Crit Care* 2004;8:R234–R242.
307. Harbarth S, Holeckova K, Froidevaux C, et al. Diagnostic value of procalcitonin, interleukin-6, and interleukin-8 in critically ill patients admitted with suspected sepsis. *Am J Respir Crit Care Med* 2001;164:396–402.
308. Kuzi S, Aroch I, Peleg K, et al. Canine procalcitonin messenger RNA expression. *J Vet Diagn Invest* 2008;20:629–633.
309. Giunti M, Peli A, Battilani M, et al. Evaluation of CALC-I gene (CALCA) expression in tissues of dogs with signs of the systemic inflammatory response syndrome. *J Vet Emerg Crit Care (San Antonio)* 2010;20:523–527.
310. Mukorera V, Dvir E, van der Merwe LL, et al. Serum C-reactive protein concentration in benign and malignant canine spirocerosis. *J Vet Intern Med* 2011;25:963–966.
311. Koster LS, Van Schoor M, Goddard A, et al. C-reactive protein in canine babesiosis caused by *Babesia rossi* and its association with outcome. *J S Afr Vet Assoc* 2009;80:87–91.
312. Sheahan D, Bell R, Mellanby RJ, et al. Acute phase protein concentrations in dogs with nasal disease. *Vet Rec* 2010;167:895–899.
313. Lowrie M, Penderis J, Eckersall PD, et al. The role of acute phase proteins in diagnosis and management of steroid-responsive meningitis arteritis in dogs. *Vet J* 2009;182:125–130.
314. Griebisch C, Arndt G, Raila J, et al. C-reactive protein concentration in dogs with primary immune-mediated hemolytic anemia. *Vet Clin Pathol* 2009;38:421–425.
315. Mitchell KD, Kruth SA, Wood RD, et al. Serum acute phase protein concentrations in dogs with autoimmune hemolytic anemia. *J Vet Intern Med* 2009;23:585–591.
316. Bathen-Noethen A, Carlson R, Menzel D, et al. Concentrations of acute-phase proteins in dogs with steroid responsive meningitis-arteritis. *J Vet Intern Med* 2008;22:1149–1156.
317. Nakamura M, Takahashi M, Ohno K, et al. C-reactive protein concentration in dogs with various diseases. *J Vet Med Sci* 2008;70:127–131.
318. Ohno K, Yokoyama Y, Nakashima K, et al. C-reactive protein concentration in canine idiopathic polyarthritis. *J Vet Med Sci* 2006;68:1275–1279.
319. Dabrowski R, Wawron W, Kostro K. Changes in CRP, SAA and haptoglobin produced in response to ovariectomy in healthy bitches and those with pyometra. *Theriogenology* 2007;67:321–327.
320. Planellas M, Bassols A, Siracusa C, et al. Evaluation of serum haptoglobin and C-reactive protein in dogs with mammary tumors. *Vet Clin Pathol* 2009;38:348–352.
321. Nielsen L, Toft N, Eckersall PD, et al. Serum C-reactive protein concentration as an indicator of remission status in dogs with multicentric lymphoma. *J Vet Intern Med* 2007;21:1231–1236.
322. Caldin M, Tasca S, Carli E, et al. Serum acute phase protein concentrations in dogs with hyperadrenocorticism with and without concurrent inflammatory conditions. *Vet Clin Pathol* 2009;38:63–68.
323. Chan DL, Rozanski EA, Freeman LM. Relationship among plasma amino acids, C-reactive protein, illness severity, and outcome in critically ill dogs. *J Vet Intern Med* 2009;23:559–563.
324. Gebhardt C, Hirschberger J, Rau S, et al. Use of C-reactive protein to predict outcome in dogs with systemic inflammatory response syndrome or sepsis. *J Vet Emerg Crit Care (San Antonio)* 2009;19:450–458.
325. van der Poll T, de Boer JD, Levi M. The effect of inflammation on coagulation and *vice versa*. *Curr Opin Infect Dis* 2011;24:273–278.
326. Machado FR, Cesar MS. Sepsis, coagulation and anticoagulants. *Endocr Metab Immune Disord Drug Targets* 2010;10:204–213.
327. Sun H. The interaction between pathogens and the host coagulation system. *Physiology (Bethesda)* 2006;21:281–288.
328. Huisse MG, Pease S, Hurtado-Nedelec M, et al. Leukocyte activation: The link between inflammation and coagulation during heatstroke. A study of patients during the 2003 heat wave in Paris. *Crit Care Med* 2008;36:2288–2295.
329. de Laforcade AM, Freeman LM, Shaw SP, et al. Hemostatic changes in dogs with naturally occurring sepsis. *J Vet Intern Med* 2003;17:674–679.
330. Bentz AI, Palmer JE, Dallap BL, et al. Prospective evaluation of coagulation in critically ill neonatal foals. *J Vet Intern Med* 2009;23:161–167.
331. Estrin MA, Wehausen CE, Jessen CR, et al. Disseminated intravascular coagulation in cats. *J Vet Intern Med* 2006;20:1334–1339.
332. Norris CR, Griffey SM, Samii VF. Pulmonary thromboembolism in cats: 29 cases (1987–1997). *J Am Vet Med Assoc* 1999;215:1650–1654.
333. Lucas WE, Kitzmiller JL. The role of intravascular coagulation in feline endotoxin shock. *Surg Gynecol Obstet* 1972;134:73–77.
334. DeClue AE, Williams KJ, Sharp C, et al. Systemic response to low-dose endotoxin infusion in cats. *Vet Immunol Immunopathol* 2009;132:167–174.
335. DelGiudice LA, White GA. The role of tissue factor and tissue factor pathway inhibitor in health and disease states. *J Vet Emerg Crit Care (San Antonio)* 2009;19:23–29.
336. Dallap Schaer BL, Epstein K. Coagulopathy of the critically ill equine patient. *J Vet Emerg Crit Care (San Antonio)* 2009;19:53–65.
337. Hopper K, Bateman S. An updated view of hemostasis: Mechanisms of hemostatic dysfunction associated with sepsis. *J Vet Emerg Crit Car* 2005;15:83–91.
338. Pawlinski R, Mackman N. Cellular sources of tissue factor in endotoxemia and sepsis. *Thromb Res* 2010;125(Suppl 1):S70–S73.
339. Stokol T, Daddona JL, Choi B. Evaluation of tissue factor procoagulant activity on the surface of feline leukocytes in response to treatment with lipopolysaccharide and heat-inactivated fetal bovine serum. *Am J Vet Res* 2010;71:623–629.
340. Grunig G, Hulliger C, Winder C, et al. Spontaneous and lipopolysaccharide-induced expression of procoagulant activity by equine lung macrophages in comparison with blood monocytes and blood neutrophils. *Vet Immunol Immunopathol* 1991;29:295–312.
341. Ouellette AL, Evans RJ, Heath MF. Platelets enhance endotoxin-induced monocyte tissue factor (TF) activity in the horse. *Res Vet Sci* 2004;76:31–35.
342. Henry MM, Moore JN. Clinical relevance of monocyte procoagulant activity in horses with colic. *J Am Vet Med Assoc* 1991;198:843–848.

343. Aras O, Shet A, Bach RR, et al. Induction of microparticle- and cell-associated intravascular tissue factor in human endotoxemia. *Blood* 2004;103:4545–4553.
344. Soriano AO, Jy W, Chirinos JA, et al. Levels of endothelial and platelet microparticles and their interactions with leukocytes negatively correlate with organ dysfunction and predict mortality in severe sepsis. *Crit Care Med* 2005;33:2540–2546.
345. Meziani F, Delabranche X, Asfar P, et al. Bench-to-bedside review: Circulating microparticles—a new player in sepsis? *Crit Care* 2010;14:236.
346. Miyoshi T, Yamashita K, Arai T, et al. The role of endothelial interleukin-8/NADPH oxidase 1 axis in sepsis. *Immunology* 2010;131:331–339.
347. Pawlinski R, Pedersen B, Schabbauer G, et al. Role of tissue factor and protease-activated receptors in a mouse model of endotoxemia. *Blood* 2004;103:1342–1347.
348. Rogers CL, Rozanski EA. Von Willebrand factor antigen concentration in dogs with sepsis. *J Vet Intern Med* 2010;24:229–230.
349. Pantaleon LG, Furr MO, McKenzie HC, et al. Effects of small- and large-volume resuscitation on coagulation and electrolytes during experimental endotoxemia in anesthetized horses. *J Vet Intern Med* 2007;21:1374–1379.
350. Bernard GR, Vincent JL, Laterre PF, et al. Efficacy and safety of recombinant human activated protein C for severe sepsis. *N Engl J Med* 2001;344:699–709.
351. Kerschen EJ, Fernandez JA, Cooley BC, et al. Endotoxemia and sepsis mortality reduction by non-anticoagulant activated protein C. *J Exp Med* 2007;204:2439–2448.
352. Dellinger RP, Carlet JM, Masur H, et al. Surviving Sepsis Campaign guidelines for management of severe sepsis and septic shock. *Intensive Care Med* 2004;30:536–555.
353. Dellinger RP, Levy MM, Carlet JM, et al. Surviving Sepsis Campaign: International guidelines for management of severe sepsis and septic shock: 2008. *Crit Care Med* 2008;36:296–327.
354. Marti-Carvajal AJ, Sola I, Lathyris D, et al. Human recombinant activated protein C for severe sepsis. *Cochrane Database Syst Rev* 2011;4:CD004388.
355. Jackson CV, Bailey BD, Shetler TJ. Pharmacological profile of recombinant, human activated protein C (LY203638) in a canine model of coronary artery thrombosis. *J Pharmacol Exp Ther* 2000;295:967–971.
356. Burdick MD, Schaub RG. Human protein C induces anticoagulation and increased fibrinolytic activity in the cat. *Thromb Res* 1987;45:413–419.
357. Oehmcke S, Herwald H. Contact system activation in severe infectious diseases. *J Mol Med (Berl)* 2010;88:121–126.
358. Frick IM, Bjorck L, Herwald H. The dual role of the contact system in bacterial infectious disease. *Thromb Haemost* 2007;98:497–502.
359. Kaplan AP, Ghebrehiwet B. The plasma bradykinin-forming pathways and its interrelationships with complement. *Mol Immunol* 2010;47:2161–2169.
360. Yarovaya GA, Blokhina TB, Neshkova EA. Contact system. New concepts on activation mechanisms and bioregulatory functions. *Biochemistry (Moscow)* 2002;67:13–24.
361. Rojkaer R, Schmaier AH. Activation of the plasma kallikrein/kinin system on endothelial cell membranes. *Immunopharmacology* 1999;43:109–114.
362. Dong JF. Cleavage of ultra-large von Willebrand factor by ADAMTS-13 under flow conditions. *J Thromb Haemost* 2005;3:1710–1716.
363. Claus RA, Bockmeyer CL, Sossdorf M, et al. The balance between von Willebrand factor and its cleaving protease ADAMTS13: Biomarker in systemic inflammation and development of organ failure? *Curr Mol Med* 2010;10:236–248.
364. Kremer Hovinga JA, Zeerleder S, Kessler P, et al. ADAMTS-13, von Willebrand factor and related parameters in severe sepsis and septic shock. *J Thromb Haemost* 2007;5:2284–2290.
365. Martin K, Borgel D, Lerolle N, et al. Decreased ADAMTS-13 (A disintegrin-like and metalloprotease with thrombospondin type 1 repeats) is associated with a poor prognosis in sepsis-induced organ failure. *Crit Care Med* 2007;35:2375–2382.
366. Ono T, Mimuro J, Madoiwa S, et al. Severe secondary deficiency of von Willebrand factor-cleaving protease (ADAMTS13) in patients with sepsis-induced disseminated intravascular coagulation: Its correlation with development of renal failure. *Blood* 2006;107:528–534.
367. Adib-Conquy M, Cavaillon JM. Compensatory anti-inflammatory response syndrome. *Thromb Haemost* 2009;101:36–47.
368. Bone RC. Sir Isaac Newton, sepsis, SIRS, and CARS. *Crit Care Med* 1996;24:1125–1128.
369. Ward NS, Casserly B, Ayala A. The compensatory anti-inflammatory response syndrome (CARS) in critically ill patients. *Clin Chest Med* 2008;29:617–625, viii.
370. Osuchowski MF, Welch K, Siddiqui J, et al. Circulating cytokine/inhibitor profiles reshape the understanding of the SIRS/CARS continuum in sepsis and predict mortality. *J Immunol* 2006;177:1967–1974.
371. Mokart D, Capo C, Blache JL, et al. Early postoperative compensatory anti-inflammatory response syndrome is associated with septic complications after major surgical trauma in patients with cancer. *Br J Surg* 2002;89:1450–1456.
372. Mokart D, Merlin M, Sannini A, et al. Procalcitonin, interleukin 6 and systemic inflammatory response syndrome (SIRS): Early markers of postoperative sepsis after major surgery. *Br J Anaesth* 2005;94:767–773.
373. Neunaber C, Zeckey C, Andruszkow H, et al. Immunomodulation in polytrauma and polymicrobial sepsis: Where do we stand? *Recent Pat Inflamm Allergy Drug Discov* 2011;5:17–25.
374. Power C, Fanning N, Redmond HP. Cellular apoptosis and organ injury in sepsis: A review. *Shock* 2002;18:197–211.
375. Ward PA. Sepsis, apoptosis and complement. *Biochem Pharmacol* 2008;76:1383–1388.
376. Paula Neto HA, Kubes P. Platelets, endothelium and shear join forces to mislead neutrophils in sepsis. *Crit Care* 2011;15:103.
377. Alves-Filho JC, Spiller F, Cunha FQ. Neutrophil paralysis in sepsis. *Shock* 2010;34(Suppl 1):15–21.
378. Delano MJ, Thayer T, Gabrilovich S, et al. Sepsis induces early alterations in innate immunity that impact mortality to secondary infection. *J Immunol* 2011;186:195–202.
379. Ocui LM, Bamboat ZM, Balachandran VP, et al. Neutrophil IL-10 suppresses peritoneal inflammatory monocytes during polymicrobial sepsis. *J Leukoc Biol* 2011;89:423–432.
380. Escoll P, del Fresno C, Garcia L, et al. Rapid up-regulation of IRAK-M expression following a second endotoxin challenge in human monocytes and in monocytes isolated from septic patients. *Biochem Biophys Res Commun* 2003;311:465–472.
381. Wullaert A, Verstrepen L, Van Huffel S, et al. LIND/ABIN-3 is a novel lipopolysaccharide-inducible inhibitor of NF-kappaB activation. *J Biol Chem* 2007;282:81–90.
382. Wolk K, Docke WD, von Baehr V, et al. Impaired antigen presentation by human monocytes during endotoxin tolerance. *Blood* 2000;96:218–223.
383. Docke WD, Randow F, Syrbe U, et al. Monocyte deactivation in septic patients: Restoration by IFN-gamma treatment. *Nat Med* 1997;3:678–681.
384. Tschaikowsky K, Hedwig-Geissing M, Schiele A, et al. Coincidence of pro- and anti-inflammatory responses in the early

phase of severe sepsis: Longitudinal study of mononuclear histocompatibility leukocyte antigen-DR expression, procalcitonin, C-reactive protein, and changes in T-cell subsets in septic and post-operative patients. *Crit Care Med* 2002;30:1015–1023.

385. Wesche DE, Lomas-Neira JL, Perl M, et al. Leukocyte apoptosis and its significance in sepsis and shock. *J Leukoc Biol* 2005;78:325–337.

386. Hsieh YC, Athar M, Chaudry IH. When apoptosis meets autophagy: Deciding cell fate after trauma and sepsis. *Trends Mol Med* 2009;15:129–138.

387. Pinheiro da Silva F, Nizet V. Cell death during sepsis: Integration of disintegration in the inflammatory response to overwhelming infection. *Apoptosis* 2009;14:509–521.

388. Levine B, Mizushima N, Virgin HW. Autophagy in immunity and inflammation. *Nature* 2011;469:323–335.

389. Giansanti V, Torriglia A, Scovassi AI. Conversation between apoptosis and autophagy: “Is it your turn or mine?”. *Apoptosis* 2011;16:321–333.

390. Watanabe E, Muenzer JT, Hawkins WG, et al. Sepsis induces extensive autophagic vacuolization in hepatocytes: A clinical and laboratory-based study. *Lab Invest* 2009;89:549–561.

391. Bergsbaken T, Fink SL, Cookson BT. Pyroptosis: Host cell death and inflammation. *Nat Rev Microbiol* 2009;7:99–109.

392. Kepp O, Galluzzi L, Zitvogel L, et al. Pyroptosis: A cell death modality of its kind? *Eur J Immunol* 2010;40:627–630.

393. Fink SL, Cookson BT. Apoptosis, pyroptosis, and necrosis: mechanistic description of dead and dying eukaryotic cells. *Infect Immun* 2005;73:1907–1916.

394. Shao W, Yeretssian G, Doiron K, et al. The caspase-1 digestome identifies the glycolysis pathway as a target during infection and septic shock. *J Biol Chem* 2007;282:36321–36329.

395. Nakajima Y, Mikami O, Yoshioka M, et al. Involvement of apoptosis in the endotoxemic lesions of the liver and kidneys of piglets. *J Vet Med Sci* 2000;62:621–626.

396. Sato M, Mikami O, Kobayashi M, et al. Apoptosis in the lymphatic organs of piglets inoculated with classical swine fever virus. *Vet Microbiol* 2000;75:1–9.

397. Messmer UK, Briner VA, Pfeilschifter J. Tumor necrosis factor-alpha and lipopolysaccharide induce apoptotic cell death in bovine glomerular endothelial cells. *Kidney Int* 1999;55:2322–2337.

398. Messmer UK, Winkel G, Briner VA, et al. Glucocorticoids potently block tumour necrosis factor-alpha- and lipopolysaccharide-induced apoptotic cell death in bovine glomerular endothelial cells upstream of caspase 3 activation. *Br J Pharmacol* 1999;127:1633–1640.

399. Sylte MJ, Leite FP, Kuckleburg CJ, et al. Caspase activation during *Haemophilus somnus* lipooligosaccharide-mediated apoptosis of bovine endothelial cells. *Microb Pathog* 2003;35:285–291.

400. Kuckleburg CJ, Tiwari R, Czuprynski CJ. Endothelial cell apoptosis induced by bacteria-activated platelets requires caspase-8 and -9 and generation of reactive oxygen species. *Thromb Haemost* 2008;99:363–372.

401. Wohlschlaeger J, Stubbe HD, Schmitz KJ, et al. Roles of MMP-2/-9 in cardiac dysfunction during early multiple organ failure in an ovine animal model. *Pathol Res Pract* 2005;201:809–817.

402. Umeshappa CS, Singh KP, Nanjundappa RH, et al. Apoptosis and immuno-suppression in sheep infected with bluetongue virus serotype-23. *Vet Microbiol* 2010;144:310–318.

403. Crouser ED, Julian MW, Weinstein DM, et al. Endotoxin-induced ileal mucosal injury and nitric oxide dysregulation are temporally dissociated. *Am J Respir Crit Care Med* 2000;161:1705–1712.

404. Paillet R, Richard S, Bloas F, et al. Toward a detailed characterization of feline immunodeficiency virus-specific T cell immune responses and mediated immune disorders. *Vet Immunol Immunopathol* 2005;106:1–14.

405. Rahal L, Garrido AG, Cruz RJ Jr, et al. Fluid replacement with hypertonic or isotonic solutions guided by mixed venous oxygen saturation in experimental hypodynamic sepsis. *J Trauma* 2009;67:1205–1212.

406. Collins AM, Sewell WA, Edwards MR. Immunoglobulin gene rearrangement, repertoire diversity, and the allergic response. *Pharmacol Ther* 2003;100:157–170.

407. Nikolich-Zugich J, Slifka MK, Messaoudi I. The many important facets of T-cell repertoire diversity. *Nat Rev Immunol* 2004;4:123–132.

408. Hollander GA, Peterson P. Learning to be tolerant: How T cells keep out of trouble. *J Intern Med* 2009;265:541–561.

409. McCaughy TM, Hogquist KA. Central tolerance: What have we learned from mice? *Semin Immunopathol* 2008;30:399–409.

410. Parish IA, Heath WR. Too dangerous to ignore: Self-tolerance and the control of ignorant autoreactive T cells. *Immunol Cell Biol* 2008;86:146–152.

411. Waldmann H. Tolerance: An overview and perspectives. *Nat Rev Nephrol* 2010;6:569–576.

412. Mueller DL. Mechanisms maintaining peripheral tolerance. *Nat Immunol* 2010;11:21–27.

413. Miyara M, Sakaguchi S. Human FoxP3⁺CD4⁺ regulatory T cells: Their knowns and unknowns. *Immunol Cell Biol* 2011;89:346–351.

414. Sakaguchi S. Regulatory T cells: History and perspective. *Methods Mol Biol* 2011;707:3–17.

415. Shevach EM. The resurrection of T cell-mediated suppression. *J Immunol* 2011;186:3805–3807.

416. Garden OA, Pinheiro D, Cunningham F. All creatures great and small: Regulatory T cells in mice, humans, dogs and other domestic animal species. *Int Immunopharmacol* 2011;11:576–588.

417. Lankford S, Petty C, LaVoy A, et al. Cloning of feline FOXP3 and detection of expression in CD4⁺CD25⁺ regulatory T cells. *Vet Immunol Immunopathol* 2008;122:159–166.

418. Petty CS, Tompkins MB, Tompkins WA. Transforming growth factor-beta/transforming growth factor-betaRII signaling may regulate CD4⁺CD25⁺ T-regulatory cell homeostasis and suppressor function in feline AIDS lentivirus infection. *J Acquir Immune Defic Syndr* 2008;47:148–160.

419. Smithberg SR, Fogle JE, Mexas AM, et al. *In vivo* depletion of CD4⁺CD25⁺ regulatory T cells in cats. *J Immunol Methods* 2008;329:81–91.

420. Vahlenkamp TW, Tompkins MB, Tompkins WA. Feline immunodeficiency virus infection phenotypically and functionally activates immunosuppressive CD4⁺CD25⁺ T regulatory cells. *J Immunol* 2004;172:4752–4761.

421. Fogle JE, Tompkins WA, Tompkins MB. CD4⁺CD25⁺ T regulatory cells from FIV⁺ cats induce a unique anergic profile in CD8⁺ lymphocyte targets. *Retrovirology* 2010;7:97.

422. Fogle JE, Mexas AM, Tompkins WA, et al. CD4⁺CD25⁺ T regulatory cells inhibit CD8⁺ IFN-gamma production during acute and chronic FIV infection utilizing a membrane TGF-beta-dependent mechanism. *AIDS Res Hum Retroviruses* 2010;26:201–216.

423. Achleitner A, Clark ME, Bienzle D. T-regulatory cells infected with feline immunodeficiency virus up-regulate programmed death-1 (PD-1). *Vet Immunol Immunopathol* 2011;143:307–313.

424. Biller BJ, Elmslie RE, Burnett RC, et al. Use of FoxP3 expression to identify regulatory T cells in healthy dogs and dogs with cancer. *Vet Immunol Immunopathol* 2007;116:69–78.

425. Keppel KE, Campbell KL, Zuckermann FA, et al. Quantitation of canine regulatory T cell populations, serum interleukin-10 and allergen-specific IgE concentrations in healthy control dogs and canine atopic dermatitis patients receiving allergen-specific immunotherapy. *Vet Immunol Immunopathol* 2008;123:337–344.

426. O'Neill K, Guth A, Biller B, et al. Changes in regulatory T cells in dogs with cancer and associations with tumor type. *J Vet Intern Med* 2009;23:875–881.
427. Horiuchi Y, Tominaga M, Ichikawa M, et al. Increase of regulatory T cells in the peripheral blood of dogs with metastatic tumors. *Microbiol Immunol* 2009;53:468–474.
428. Mizuno T, Suzuki R, Umeki S, et al. Crossreactivity of antibodies to canine CD25 and Foxp3 and identification of canine CD4⁺CD25⁺Foxp3⁺ cells in canine peripheral blood. *J Vet Med Sci* 2009;71:1561–1568.
429. Abrams VK, Hwang B, Lesnikova M, et al. A novel monoclonal antibody specific for canine CD25 (P4A10): Selection and evaluation of canine Tregs. *Vet Immunol Immunopathol* 2010;135:257–265.
430. Risetto KC, Rindt H, Selting KA, et al. Cloning and expression of canine CD25 for validation of an anti-human CD25 antibody to compare T regulatory lymphocytes in healthy dogs and dogs with osteosarcoma. *Vet Immunol Immunopathol* 2010;135:137–145.
431. Horiuchi Y, Tominaga M, Ichikawa M, et al. Relationship between regulatory and type 1 T cells in dogs with oral malignant melanoma. *Microbiol Immunol* 2010;54:152–159.
432. Pinheiro D, Singh Y, Grant CR, et al. Phenotypic and functional characterization of a CD4⁺ CD25^{high} FOXP3^{high} regulatory T-cell population in the dog. *Immunology* 2011;132:111–122.
433. Burton JH, Mitchell L, Thamm DH, et al. Low-dose cyclophosphamide selectively decreases regulatory T cells and inhibits angiogenesis in dogs with soft tissue sarcoma. *J Vet Intern Med* 2011;25:920–926.
434. Kaser T, Gerner W, Hammer SE, et al. Detection of Foxp3 protein expression in porcine T lymphocytes. *Vet Immunol Immunopathol* 2008;125:92–101.
435. Kaser T, Gerner W, Hammer SE, et al. Phenotypic and functional characterisation of porcine CD4⁺CD25^{high} regulatory T cells. *Vet Immunol Immunopathol* 2008;122:153–158.
436. Bolzer K, Kaser T, Saalmuller A, et al. Molecular characterisation of porcine Forkhead-box p3 (Foxp3). *Vet Immunol Immunopathol* 2009;132:275–281.
437. Wongyanin P, Buranapraditkun S, Chokeshai-Usaha K, et al. Induction of inducible CD4⁺CD25⁺Foxp3⁺ regulatory T lymphocytes by porcine reproductive and respiratory syndrome virus (PRRSV). *Vet Immunol Immunopathol* 2010;133:170–182.
438. Silva-Campa E, Flores-Mendoza L, Resendiz M, et al. Induction of T helper 3 regulatory cells by dendritic cells infected with porcine reproductive and respiratory syndrome virus. *Virology* 2009;387:373–379.
439. Dawson H, Solano-Aguilar G, Beal M, et al. Localized Th1-, Th2-, T regulatory cell-, and inflammation-associated hepatic and pulmonary immune responses in *Ascaris suum*-infected swine are increased by retinoic acid. *Infect Immun* 2009;77:2576–2587.
440. Pilon C, Meurens F, Dauba A, et al. Induction of porcine regulatory cells by mycophenolic acid-treated dendritic cells. *Transplant Proc* 2009;41:700–702.
441. Kaser T, Gerner W, Mair K, et al. Current knowledge on porcine regulatory T cells. *Vet Immunol Immunopathol* 2011; <http://doi.org/10.1016/j.vetimm.2011.05.035> (in press).
442. Kaser T, Gerner W, Saalmuller A. Porcine regulatory T cells: Mechanisms and T-cell targets of suppression. *Dev Comp Immunol* 2011;35:1166–1172.
443. Gerner W, Stadler M, Hammer SE, et al. Sensitive detection of Foxp3 expression in bovine lymphocytes by flow cytometry. *Vet Immunol Immunopathol* 2010;138:154–158.
444. Hoek A, Rutten VP, Kool J, et al. Subpopulations of bovine WC1⁺ gammadelta T cells rather than CD4⁺CD25^{high} Foxp3⁺ T cells act as immune regulatory cells ex vivo. *Vet Res* 2009;40:6.
445. Seo KS, Davis WC, Hamilton MJ, et al. Development of monoclonal antibodies to detect bovine FOXP3 in PBMCs exposed to a staphylococcal superantigen. *Vet Immunol Immunopathol* 2009;128:30–36.
446. de Almeida DE, Colvin CJ, Coussens PM. Antigen-specific regulatory T cells in bovine paratuberculosis. *Vet Immunol Immunopathol* 2008;125:234–245.
447. McNeilly TN, McIntyre J, Frew D, et al. Infestation of sheep with *Psoroptes ovis*, the sheep scab mite, results in recruitment of Foxp3(+) T cells into the dermis. *Parasite Immunol* 2010;32:361–369.
448. Rocchi MS, Wattedegera SR, Frew D, et al. Identification of CD4(+)CD25(high) Foxp3(+) T cells in ovine peripheral blood. *Vet Immunol Immunopathol* 2011;144:172–177.
449. Wagner B, Hillegas JM, Brinker DR, et al. Characterization of monoclonal antibodies to equine interleukin-10 and detection of T regulatory 1 cells in horses. *Vet Immunol Immunopathol* 2008;122:57–64.
450. Hamza E, Torsteinsdottir S, Eydal M, et al. Increased IL-4 and decreased regulatory cytokine production following relocation of Icelandic horses from a high to low endoparasite environment. *Vet Immunol Immunopathol* 2010;133:40–50.
451. Hamza E, Steinbach F, Marti E. CD4⁺CD25⁺ T cells expressing FoxP3 in Icelandic horses affected with insect bite hypersensitivity. *Vet Immunol Immunopathol* 2011; <http://doi.org/10.1016/j.vetimm.2011.05.033> (in press).
452. Wagner B, Burton A, Ainsworth D. Interferon-gamma, interleukin-4 and interleukin-10 production by T helper cells reveals intact Th1 and regulatory TR1 cell activation and a delay of the Th2 cell response in equine neonates and foals. *Vet Res* 2010;41:47.
453. Heimann M, Janda J, Sigurdardottir OG, et al. Skin-infiltrating T cells and cytokine expression in Icelandic horses affected with insect bite hypersensitivity: A possible role for regulatory T cells. *Vet Immunol Immunopathol* 2011;140:63–74.
454. Robbin MG, Wagner B, Noronha LE, et al. Subpopulations of equine blood lymphocytes expressing regulatory T cell markers. *Vet Immunol Immunopathol* 2011;140:90–101.
455. Langier S, Sade K, Kivity S. Regulatory T cells: The suppressor arm of the immune system. *Autoimmun Rev* 2010;10:112–115.
456. Sakaguchi S, Wing K, Onishi Y, et al. Regulatory T cells: How do they suppress immune responses? *Int Immunol* 2009;21:1105–1111.
457. Workman CJ, Szymczak-Workman AL, Collison LW, et al. The development and function of regulatory T cells. *Cell Mol Life Sci* 2009;66:2603–2622.
458. Rudensky AY. Regulatory T cells and Foxp3. *Immunol Rev* 2011;241:260–268.
459. Campbell DJ, Ziegler SF. FOXP3 modifies the phenotypic and functional properties of regulatory T cells. *Nat Rev Immunol* 2007;7:305–310.
460. Shalev I, Schmelzle M, Robson SC, et al. Making sense of regulatory T cell suppressive function. *Semin Immunol* 2011;23:282–292.
461. Wan YY. Regulatory T cells: Immune suppression and beyond. *Cell Mol Immunol* 2010;7:204–210.
462. Caramalho I, Lopes-Carvalho T, Ostler D, et al. Regulatory T cells selectively express Toll-like receptors and are activated by lipopolysaccharide. *J Exp Med* 2003;197:403–411.
463. Dai J, Liu B, Li Z. Regulatory T cells and Toll-like receptors: What is the missing link? *Int Immunopharmacol* 2009;9:528–533.
464. Liu H, Komai-Koma M, Xu D, et al. Toll-like receptor 2 signaling modulates the functions of CD4⁺CD25⁺ regulatory T cells. *Proc Natl Acad Sci USA* 2006;103:7048–7053.

465. Nyirenda MH, O'Brien K, Sanvito L, et al. Modulation of regulatory T cells in health and disease: Role of toll-like receptors. *Inflamm Allergy Drug Targets* 2009;8:124–129.
466. Monneret G, Debard AL, Venet F, et al. Marked elevation of human circulating CD4⁺ CD25⁺ regulatory T cells in sepsis-induced immunoparalysis. *Crit Care Med* 2003;31:2068–2071.
467. Saito K, Wagatsuma T, Toyama H, et al. Sepsis is characterized by the increases in percentages of circulating CD4⁺ CD25⁺ regulatory T cells and plasma levels of soluble CD25. *Tohoku J Exp Med* 2008;216:61–68.
468. Venet F, Chung CS, Kherouf H, et al. Increased circulating regulatory T cells (CD4⁺CD25⁺CD127⁻) contribute to lymphocyte anergy in septic shock patients. *Intensive Care Med* 2009;35:678–686.
469. Venet F, Pachot A, Debard AL, et al. Increased percentage of CD4⁺ CD25⁺ regulatory T cells during septic shock is due to the decrease of CD4⁺CD25⁻ lymphocytes. *Crit Care Med* 2004;32:2329–2331.
470. Heuer JG, Zhang T, Zhao J, et al. Adoptive transfer of *in vitro*-stimulated CD4⁺ CD25⁺ regulatory T cells increases bacterial clearance and improves survival in polymicrobial sepsis. *J Immunol* 2005;174:7141–7146.
471. Scumpia PO, Delano MJ, Kelly KM, et al. Increased natural CD4⁺ CD25⁺ regulatory T cells and their suppressor activity do not contribute to mortality in murine polymicrobial sepsis. *J Immunol* 2006;177:7943–7949.
472. Wisnoski N, Chung CS, Chen Y, et al. The contribution of CD4⁺ CD25⁺ T-regulatory-cells to immune suppression in sepsis. *Shock* 2007;27:251–257.
473. Tiemessen MM, Jagger AL, Evans HG, et al. CD4⁺ CD25⁺ Foxp3⁺ regulatory T cells induce alternative activation of human monocytes/macrophages. *Proc Natl Acad Sci USA* 2007;104:19446–19451.
474. Venet F, Pachot A, Debard AL, et al. Human CD4⁺ CD25⁺ regulatory T lymphocytes inhibit lipopolysaccharide-induced monocyte survival through a Fas/Fas ligand-dependent mechanism. *J Immunol* 2006;177:6540–6547.
475. Hodge G, Scott J, Osborn M, et al. Increased T regulatory cells and decreased Th1 pro-inflammatory cytokines correlate with culture-positive infection in febrile neutropenia childhood oncology patients. *Cytokine* 2011;53:286–288.
476. Huang LF, Yao YM, Dong N, et al. Association between regulatory T cell activity and sepsis and outcome of severely burned patients: a prospective, observational study. *Crit Care* 2010;14:R3.
477. Hein F, Massin F, Cravoisy-Popovic A, et al. The relationship between CD4⁺ CD25⁺CD127⁻ regulatory T cells and inflammatory response and outcome during shock states. *Crit Care* 2010;14:R19.
478. van Maren WW, Jacobs JF, de Vries IJ, et al. Toll-like receptor signalling on Tregs: To suppress or not to suppress? *Immunology* 2008;124:445–452.
479. Walker LS. Regulatory T cells overturned: The effectors fight back. *Immunology* 2009;126:466–474.
480. Marshall JC. Sepsis: Rethinking the approach to clinical research. *J Leukoc Biol* 2008;83:471–482.
481. Girbes AR, Beishuizen A, Strack van Schijndel RJ. Pharmacological treatment of sepsis. *Fundam Clin Pharmacol* 2008;22:355–361.
482. Wittebole X, Collienne C, Castanares-Zapatero D, et al. Adjunctive therapies for severe sepsis. *Int J Antimicrob Agents* 2008;32(Suppl 1):S34–S38.