

Supplemental Table II. Different Types of Infection in LVAD.

	Types of Infection
6 month ± 2 weeks before intracranial hemorrhage	Bloodstream infection (N=1) Driveline infection (N=1) Soft tissue infection (N=1)
3 month ± 2 weeks before intracranial hemorrhage	Bloodstream infection (N=2) Driveline infection (N=1) Pneumonia (N=1) Gastrointestinal infection (N=1)
1 month ± 2 weeks before intracranial hemorrhage	Bloodstream infection (N=1) Driveline infection (N=1) Urinary tract infection (N=3) Pleural empyema (N=1) Soft tissue infection (N=1)
Acute intracranial hemorrhage ± 2 weeks	Bloodstream infection (N=16) Driveline infection (N=2) Pump pocket infection (N=1) Urinary tract infection (N=7) Pneumonia (N=4) Gastrointestinal infection (N=1) Meningitis (N=1) Eye (N=1)
1 month ± 2 weeks after intracranial hemorrhage	Bloodstream infection (N=3) Driveline infection (N=2) Pneumonia (N=1)
3 month ± 2 weeks after intracranial hemorrhage	Bloodstream infection (N=2) Driveline infection (N=2) Pump pocket infection (N=1) Urinary tract infection (N=1)
6 month ± 2 weeks after intracranial hemorrhage	Bloodstream infection (N=4) Driveline infection (N=1)

Bloodstream infection = 29, Driveline infection N=10, Urinary tract infection = 8