

SCIENTIFIC REPORTS



OPEN

The impact of long-term organic farming on soil-derived greenhouse gas emissions

Colin Skinner¹, Andreas Gattinger^{1,2}, Maike Krauss¹, Hans-Martin Krause¹, Jochen Mayer³, Marcel G. A. van der Heijden^{4,5,6} & Paul Mäder¹

Agricultural practices contribute considerably to emissions of greenhouse gases. So far, knowledge on the impact of organic compared to non-organic farming on soil-derived nitrous oxide (N₂O) and methane (CH₄) emissions is limited. We investigated N₂O and CH₄ fluxes with manual chambers during 571 days in a grass-clover–silage maize – green manure cropping sequence in the long-term field trial “DOK” in Switzerland. We compared two organic farming systems – biodynamic (BIODYN) and bioorganic (BIOORG) – with two non-organic systems – solely mineral fertilisation (CONMIN) and mixed farming including farmyard manure (CONFYM) – all reflecting Swiss farming practices—together with an unfertilised control (NOFERT). We observed a 40.2% reduction of N₂O emissions per hectare for organic compared to non-organic systems. In contrast to current knowledge, yield-scaled cumulated N₂O emissions under silage maize were similar between organic and non-organic systems. Cumulated on area scale we recorded under silage maize a modest CH₄ uptake for BIODYN and CONMIN and high CH₄ emissions for CONFYM. We found that, in addition to N input, quality properties such as pH, soil organic carbon and microbial biomass significantly affected N₂O emissions. This study showed that organic farming systems can be a viable measure contributing to greenhouse gas mitigation in the agricultural sector.

With a share of 10–12% in carbon-dioxide equivalents (CO₂-eq.), agriculture contributes substantially to global greenhouse gas (GHG) emissions¹. Considering indirect emissions from agriculture-related activities such as fertiliser production and land use change, this share can be up to 30%². In 2005, the agricultural sector contributed 56% to the emissions of the anthropogenic non-CO₂ GHG, i.e. nitrous oxide and methane, with an annual growth of 0.9%¹. An estimated 38% of agriculture’s direct emissions originate from soils, 15% from N₂O from manure on pasture and 12% from N₂O of synthetic fertilisers – the latter at an annual growth rate of 3.9% from 1961 to 2010. In comparison, paddy rice cultivation contributes 11% to the sector’s annual emissions, mainly in the form of CH₄¹. Due to their high global warming potential (GWP) the reduction of N₂O and CH₄ emissions thus gives agriculture’s mitigation potential regarding climate change a high leverage in favour of limiting Global Warming³.

Fostered by the Green Revolution and the availability of cheap synthetic fertilisers from the 1950s onward, world grain harvests doubled to 2.5 billion tons between 1970 and 2010 with an average yield increase from 1600 to 3030 kg ha⁻¹. However, this agricultural intensification was realized mainly through the increase in global fertiliser usage from 32 to 106 Mt yr⁻¹ (+331%). Moreover, these achievements have been accomplished at the expense of environmental damages such as biodiversity loss, accelerated soil erosion and degradation, eutrophication including algal blooms and oceanic dead zones, or pesticide effects on humans and wildlife⁴. There is an increased understanding that the challenges of producing enough food and biomass while maintaining ecosystem services cannot be met by modern intensive “non-organic” agricultural practices that rely heavily on synthetic

¹Department of Soil Sciences, Research Institute of Organic Agriculture (FiBL), Ackerstrasse, CH-5070, Frick, Switzerland. ²Organic Farming with focus on Sustainable Soil Use, Institute of Crop Science and Breeding II, Justus-Liebig University Giessen, 35394, Giessen, Germany. ³Water Protection and Substance Flows, Research Division Agroecology and Environment, Agroscope, CH 8046, Zürich, Switzerland. ⁴Plant Soil Interactions, Research Division Agroecology and Environment, Agroscope, CH 8046, Zürich, Switzerland. ⁵Department of Evolutionary Biology and Environmental Studies, University of Zürich, CH 8057, Zürich, Switzerland. ⁶Institute of Environmental Biology, Faculty of Science, Utrecht University, 3508, TC, Utrecht, The Netherlands. Correspondence and requests for materials should be addressed to A.G. (email: andreas.gattinger@agrar.uni-giessen.de)

fertiliser input and pesticide application⁴. Thus, agro-ecological approaches towards an ecological intensification are fundamental for future food production⁵. Certified “organic” farming is considered as one option towards ecological intensification⁵. Reganold and Wachter recently examined the performance of organic farming in the light of the four key sustainability metrics: productivity, environmental impact, economic viability and social wellbeing. They concluded that, despite lower yields, organic farming delivers greater ecosystem services and social benefits. Thus, organic farming practices need to be considered for the development of sustainable farming systems, which presumably will encompass a blend of organic together with non-organic approaches⁶.

However, there are still knowledge gaps regarding the effects of organic farming on soil carbon (C) and nitrogen (N) fluxes, and thus on N₂O and CH₄ greenhouse gas emissions from the soil⁷. Organic fertilisation strategies through farmyard manure and cropping of N-fixing legumes, grass-clover leys and catch crops influence soil C and N fluxes in complex processes. These can run over years and decouple fertiliser inputs and availability of nutrients⁸. Investigating these processes requires long-term measurements, at best covering complete crop rotations. Skinner *et al.* performed a literature search on measured soil-derived nitrous-oxide and methane fluxes under organic and non-organic management from comparison trials of farming systems, followed by a meta-analysis of the aggregated global dataset. Only 19 studies based on field measurements could be retrieved, of which 13 covered annual measurements. Based on 12 studies comprising annual measurements within the land-use types arable and grassland, lower area-scaled N₂O emissions and (three studies) higher CH₄ uptake in organic compared to non-organic farming systems were determined⁸. However, related to crop yield (eight studies), organically managed soils emitted more N₂O than soils in non-organic systems due to the yield gap of 26% between organic and non-organic farming. Reviews of global datasets resulted with a gap of around 20%^{8–10}. Yet, due to the small number of studies and the average duration of contrasting management of only nine years, further evidence originating from long-term experiments is needed. Along this line, Van Kessel *et al.* found in their global meta-study that management-induced impacts on the N₂O emission pattern become effective only after more than 10 years since adoption of reduced tillage practices¹¹.

There are different kinds of organic farming systems unified under the umbrella of the International Federation of Organic Agricultural Movements IFOAM (www.ifoam.bio)¹². Among these, biodynamic agriculture as proposed by Steiner in 1924 is the oldest certified organic farming type^{13,14}. Even though considerable parts of Steiner’s biodynamic philosophy and practices lie beyond scientific judgement, a fair share of the available peer-reviewed research results from controlled field experiments as well as case studies show effects of biodynamic farming on yield, soil quality and soil biodiversity^{15–19}. However, so far it has not been tested whether the effects of biodynamic farming on GHG emissions differ from other organic farming systems.

The DOK long-term field trial established in 1978 compares organic and non-organic farming systems¹⁵. The differentiated management of the farming systems compared since the start has led to system-specific changes of soil physical, chemical and biological properties^{15,18,19}. By measuring soil gas fluxes in the farming systems of the DOK trial, we aim to extend the knowledge base concerning the impact of organic farming on direct GHG emissions. Two organic farming systems – bioorganic (BIOORG; with slurry and rotted manure) and biodynamic (BIODYN; with slurry and composted manure) – and two non-organic – conventional with livestock (CONFYM; with slurry, fresh stacked manure and synthetic fertilisers) and without (CONMIN; only synthetic fertilisers), all fertilised to an intensity typically practiced in Switzerland were chosen for the experiment. We included the unfertilised control, NOFERT, as a reference. All farming systems are subject to the same crop rotation, but plant protection in the organic and the unfertilised system is based on mechanical weeding, indirect disease control measures and plant extracts together with bio-controls against insects, while in the non-organic systems herbicides, fungicides and pesticides are applied. The monitoring covered a cropping sequence consisting of grass-clover in the second year followed by silage maize (cv. “Colisee”) and green manure (*Brassica chinensis* x *Brassica rapa*) covering 571 days with 86 samplings.

Based on the findings of aforesaid study⁸, which serves as a topical review to present field research, we hypothesised that (i) area-scaled N₂O emissions are lower, (ii) yield-scaled N₂O emissions are higher and (iii) CH₄ uptake is higher in organic compared to non-organic farming systems. Furthermore, we aimed at investigating possible effects of biodynamic farming and identifying major drivers for the observed findings to suggest mitigation options.

Results

N₂O and CH₄ flux dynamics. For all investigated farming systems N₂O fluxes were lowest during the grass-clover phase and highest during the silage maize period with two major emission events (Fig. 1). The first one consisted of distinctly raised fluxes in CONFYM after solid manure application and ploughing, together with peak fluxes in CONMIN that immediately followed the initial mineral fertilisation and sowing of silage maize. The second N₂O emission pulse, pronounced in all systems, developed after the second fertiliser application during silage maize cultivation (Fig. 1; Supplementary Table S1). Enhanced N₂O fluxes at the beginning of the green manure phase that followed silage maize harvest amount to between 20 and 34% of total cumulated area scale N₂O emissions for the entire crop sequence, although no additional N was applied (Fig. 1; Table 1).

Overall, N₂O flux dynamics were best explained by soil moisture expressed as water-filled pore space (WFPS) ($p < 0.0001$), followed by soil nitrate concentrations ($p = 0.021$). Freeze-thaw induced N₂O fluxes were not observed during winter periods 2012/2013 and 2013/2014. The prerequisite condition for frost/thaw events, frozen soil, did not occur (Fig. 1, Supplementary Fig. S1).

CH₄ emissions differed between farming systems with slight CH₄ uptake in BIODYN and CONMIN and no or low CH₄ net emissions in BIOORG and NOFERT (Table 2). CONFYM was a strong source for CH₄, with peaking fluxes starting immediately after solid manure application and ploughing and lasting until the second fertilisation of the silage maize. Weather conditions during this period were rather cool with excessive rainfall (Fig. 1; Supplementary Fig. S1; Table 2).




Figure 1. Temporal dynamics of $\text{N}_2\text{O-N}$ (a) and $\text{CH}_4\text{-C}$ (b) fluxes during the grass-clover – silage maize – green manure cropping sequence (24 Aug 2012 to 18 Mar 2014) in the different farming systems of the DOK long-term field trial. Numbers on top of the panels indicate management operations; detailed information is given in supplementary table S1. The horizontal coloured lines in panel (a) indicate cropping seasons, gaps are falls. Panel (c) depicts the temporal dynamics of N_{min} contents (0–20 cm) per system. Panel (d) shows the temporal dynamics of water filled pore space (WFPS) with the scale on the right side (line colours indicate the systems) and precipitation, air- and soil temperature (10 cm depth) with the scale on the left-hand side.

Farming System	Grass-clover (24 Aug 2012 to 7 May 2013, 256 days)		Silage Maize (7 May 2013 to 24 Sep 2013, 140 days)		Green Manure (24 Sep 2013 to 18 Mar 2014, 175 days)		Overall (24 Aug 2012 to 18 Mar 2014, 571 days)		Annual data (18 Mar 2013 to 18 Mar 2014)	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
BIODYN	0.30	0.04bc	2.07	0.38b	1.15	0.29bc	3.67	0.40c	3.38	0.38b
BIOORG	0.40	0.03ab	3.55	0.23ab	0.89	0.06bc	5.04	0.34bc	4.66	0.33b
CONFYM	0.49	0.02a	4.75	0.49a	2.81	0.51a	8.16	0.47a	7.68	0.48a
CONMIN	0.26	0.04c	4.22	0.85a	1.59	0.36ab	6.16	0.94ab	5.91	0.93ab
NOFERT	— ^a		4.17	1.03a	0.73	0.17c	— ^a		5.03	1.08ab
ANCOVA	F-value	p-value	F-value	p-value	F-value	p-value	F-value	p-value	F-value	p-value
(Intercept)	204.44	<0.0001	155.84	<0.0001	4.37	0.006	248.13	<0.0001	732.44	<0.0001
Clay content	16.04	0.003	1.10	0.316	9.98	0.009	1.62	0.238	4.00	0.070
Farming System	9.28 ^a	0.005	3.38	0.049	7.70	0.003	10.68 ^a	0.003	4.56	0.020
Contrasts	t-value	p-value	t-value	p-value	t-value	p-value	t-value	p-value	t-value	p-value
org vs. non-org	−0.96	0.333	−2.61	0.008	−4.31	<0.0001	−4.94	<0.0001	−3.95	<0.0001

Table 1. Means and standard errors (SE) of cumulative area-scaled N_2O emissions ($\text{kg N}_2\text{O-N ha}^{-1}$) of a grass-clover – silage maize – green manure sequence. ^aMeasurements in NOFERT started at 14 March 2013 and thus were excluded from statistical analysis. ANCOVA on log scaled N_2O emissions per farming system ($n = 4$). Letters = Post-hoc Tukey test ($p < 0.05$). Contrasts: Post-hoc t-test on pairwise comparisons of org (BIODYN + BIOORG) and non-org (CONFYM + CONMIN) systems ($n = 8$).

Area-scaled GHG emissions. Cumulated over the whole observation period of 571 days mean N_2O emissions ranged from 3.67 (BIODYN) to 8.16 $\text{kg N}_2\text{O-N ha}^{-1}$ (CONFYM) (Table 1). Related to a one-year period, with GHG flux data being recorded from all five farming system treatments (including NOFERT), annual N_2O emissions varied between 3.38 (BIODYN) to 7.68 $\text{kg N}_2\text{O-N ha}^{-1}$ (CONFYM). This annual dataset encompasses samplings in silage maize and green manure. For all observation periods a significant impact of farming systems on N_2O emissions was observed ($p = 0.005$; 0.049; 0.003; 0.003 and 0.020) (Table 1). Except for the grass-clover phase, we detected significantly lower area-scaled N_2O emissions for the cropping periods “silage maize” and “green manure” between organic (BIODYN, BIOORG) compared to the non-organic (CONFYM, CONMIN) systems, as revealed by contrast analysis ($p = 0.008$; <0.0001). For the silage-maize cropping period lasting 140 days we found lowest N_2O emissions with 2.07 $\text{kg N}_2\text{O-N ha}^{-1}$ in BIODYN and highest in CONFYM (4.75 $\text{kg N}_2\text{O-N ha}^{-1}$).

Farming System	Grass-clover (24 Aug 2012 to 7 May 2013, 256 days)		Silage Maize (7 May 2013 to 24 Sep 2013, 140 days)		Green Manure (24 Sep 2013 to 18 Mar 2014, 175 days)		Overall (24 Aug 2012 to 18 Mar 2014, 571 days)		Annual data (18 Mar 2013 to 18 Mar 2014)	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
BIODYN	-0.27	0.02b	-0.17	0.04b	0.00	0.03a	-0.48	0.04b	-0.26	0.05b
BIOORG	-0.13	0.06ab	0.02	0.07ab	0.03	0.06a	-0.09	0.15ab	0.00	0.12ab
CONFYM	0.05	0.14a	3.31	1.26a	-0.03	0.04a	3.29	1.21a	3.21	1.31a
CONMIN	-0.29	0.14ab	-0.13	0.03b	-0.06	0.06a	-0.49	0.20ab	-0.26	0.11b
NOFERT	— ^a		0.19	0.14ab	0.15	0.04a	— ^a		0.33	0.15ab
Kruskal Wallis test	χ^2	p-value	χ^2	p-value	χ^2	p-value	χ^2	p-value	χ^2	p-value
Farming System	7.70 ^a	0.053	14.56	0.006	8.31	0.081	10.48 ^a	0.015	14.64	0.006
org vs. non-org	1.86	0.172	2.16	0.141	0.89	0.345	2.48	0.115	1.59	0.208

Table 2. Means and standard errors (SE) of cumulative CH₄ emissions (kg CH₄-C ha⁻¹) of a grass-clover – silage maize – green manure sequence. ^aMeasurements in NOFERT started at 14 March 2013 and thus were excluded from statistical analysis. Kruskal Wallis non-parametric test on CH₄ emissions per farming system (n = 4) and pairwise comparisons of org (BIODYN + BIOORG) and non-org (CONFYM + CONMIN) systems (n = 8). Letters = Post-hoc Kruskal Nemeny test (p < 0.05).

	Silage maize DM ^a yields		Yield-scaled N ₂ O emissions	
	Mg DM ha ⁻¹	% of CONFYM	g N ₂ O-N Mg DM ⁻¹	
Farming system effect				
BIODYN	13.57 ± 0.70b	73	154.8 ± 31.0b	
BIOORG	13.21 ± 0.23b	71	269.0 ± 17.5ab	
CONFYM	18.62 ± 1.22a	100	259.1 ± 32.7ab	
CONMIN	18.00 ± 0.77a	97	238.3 ± 52.5ab	
NOFERT	8.65 ± 0.35c	47	478.8 ± 113.9a	
ANCOVA	F-value	p-value	F-value	p-value
(Intercept)	10577.4	<0.0001	3489.5	<0.0001
Clay content	7.06	0.022	0.21	0.652
Farming System	33.53	<0.0001	3.94	0.031
Contrasts	t-value	p-value	t-value	p-value
org vs. non-org	-7.17	<0.0001	-0.59	0.550

Table 3. Silage maize yields and yield-scaled N₂O emissions (mean ± SE). ^aDry matter (DM) 80% ANCOVA per farming system (n = 4). Letters = Post-hoc Tukey test (p < 0.05). Contrasts: Post-hoc t-test on pairwise comparisons of org (BIODYN + BIOORG) and non-org (CONFYM + CONMIN) systems (n = 8).

N₂O-N ha⁻¹). BIODYN showed significantly (p < 0.05) lowest N₂O emissions for the overall and silage maize observation periods. N₂O emissions for the unfertilised treatment NOFERT under silage maize nearly matched those from fertilised non-organic treatments.

Cumulated over the whole observation period of 571 days, CH₄ emissions ranged from -0.49 (CONMIN) to +3.29 kg CH₄-C ha⁻¹ (CONFYM) (Table 2). Statistical analyses revealed a farming systems effect with significantly (p = 0.015) highest CH₄ emissions in CONFYM but no differences between both organic and non-organic systems. The effect of farming systems was most pronounced (p = 0.006) during silage maize cropping (Table 2).

Yield-scaled N₂O emissions. Yield-scaled N₂O emissions were determined only for silage maize. We recorded by far the lowest yield-scaled N₂O emissions in BIODYN (Table 3). The highest yield-scaled N₂O emissions for silage-maize were observed in NOFERT with 478.8 g N₂O-N Mg⁻¹ dry matter (Table 3). Yield-scaled N₂O emissions in BIOORG, CONFYM and CONMIN were on a medium level in between. Maize yields in 2013 were highest in the non-organic systems CONFYM/CONMIN and significantly (p < 0.0001) lower in the organic systems BIODYN/BIOORG. NOFERT achieved about half of the yield of CONFYM (Table 3). The organic yield gap, calculated on the averages of both organic vs. both non-organic farming systems, amounted to 27% in 2013.

Drivers of cumulative N₂O emissions. To identify the impact of N inputs and soil biogeochemical parameters on N₂O emissions, we compared N₂O emissions of silage maize in the 2013 cropping period with N input and the soil parameters, pH, soil organic carbon (SOC) and microbial biomass carbon (C_{mic}) as determined 2012 in the grass-clover ley (Supplementary Table S2). Since the unfertilised control (NOFERT) showed distinct differences in soil biochemical quality and received no N input, we calculated the correlations including (attributing it zero N-input) or excluding NOFERT (Table 4). If NOFERT was included, soil pH, C_{mic} and SOC were significantly (p < 0.05) correlated with N₂O emissions. These parameters also showed a high level of correlation in between each other. Excluding NOFERT, total N input (N_t input) was most pronounced, the input of mineral N (N_{min} input) revealed a high impact on N₂O emissions together with soil pH.

	Including NOFERT					Excluding NOFERT				
	N ₂ O	Nt input	N _{min} input ¹	pH	SOC	N ₂ O	Nt input	N _{min} input ¹	pH	SOC
N _i input	0.18					0.55*				
N _{min} input ¹	0.37	0.80***				0.63**	0.72**			
pH	-0.54*	0.38	0.25			-0.63**	-0.57*	-0.33		
SOC	-0.46*	0.18	-0.04	0.67**		-0.37	-0.36	-0.44	0.54*	
C _{mic}	-0.50*	0.35	0.02	0.77***	0.88***	-0.49	-0.30	-0.52*	0.57*	0.86***

Table 4. Pearson correlation coefficients between cumulative area-scaled N₂O emissions in silage maize, total N (Nt) and mineralised N (N_{min}) input to maize and soil parameters sampled in the DOK trial in 2012. Correlations were calculated across farming systems by including and excluding NOFERT. ¹N_{min}: nitrate and ammonium. SOC – soil organic carbon, C_{mic} – microbial biomass C. Significant levels for correlations: *p < 0.05, **p < 0.01, ***p < 0.001.

Discussion

With this study we obtained a unique GHG flux dataset from the oldest and longest running farming system comparison trial “DOK”. This GHG monitoring campaign was started 34 years after implementation of contrasting organic and non-organic farming systems. This dataset should thus reflect system-specific physical, chemical and biological soil properties (Supplementary Table S2) resulting from long-term differentiated management of the farming systems compared.

We observed that area-scale cumulated overall N₂O emissions were on average 40.2% lower in organic compared to non-organic farming systems. On an annual basis covering GHG sampling in silage maize and a non-leguminous green manure, organic farming emitted on average 2.78 kg less N₂O-N ha⁻¹ than non-organic farming (Table 1). These results are in accordance with the first hypothesis. The topical review demonstrated based on 70 comparisons, that organic farming emitted annually 1.05 kg less N₂O-N ha⁻¹ than non-organic farming⁸. The results from our field measurements further support the anticipated pronounced time-dependent long-term decrease of nitrous oxide emission rates⁸ as in 2013 the DOK was in the 35th year while the average duration of the trials evaluated in the review was nine years. Soil quality develops on long term towards a balanced steady state of its ecosystem. The time duration since adoption of an alternative farming practice is an important issue because nitrous oxide fluxes have been shown to decrease with a pronounced time dependency¹¹. Longer duration also leads to improvement in soil quality indicated e.g. by raising SOC concentration, microbial biomass and -diversity^{20,21}. Thus, the finding points to an increased N₂O mitigation potential of organic farming over time. More measurements, preferably covering complete crop rotations, are needed to confirm this observation on long-term dynamics within the context of soil quality, soil nutrient fluxes and pools.

During silage maize cropping, organic farming emitted, cumulated on area scale, on average 1.68 kg less N₂O-N ha⁻¹ compared to non-organic farming. N₂O emissions recorded for silage maize cropping were in accordance with previous studies under similar pedoclimatic conditions: Kaiser and Ruser reported annual emissions of 2.1 kg N ha⁻¹ for synthetic fertilisers and 5.3 kg N ha⁻¹ for farmyard manure under a fertilisation intensity of 200 kg N ha⁻¹²².

We observed pronounced N₂O emissions after grass-clover incorporation in all farming systems as has been shown elsewhere²³. Even in the NOFERT system pronounced N₂O emissions were observed following grass-clover incorporation and soil preparation for maize (Fig. 1). Cumulated on area scale, N₂O emissions for the unfertilised treatment NOFERT under silage maize were, with only 0.32 kg N₂O-N ha⁻¹, insignificantly lower than the average emissions in the fertilised non-organic treatments (Table 1). Mineral N derived from decomposition of the incorporated grass-clover ley and from soil organic matter may not have been efficiently taken up by maize plants compared with fertilised systems, as the soil in NOFERT lacks important macro nutrients such as P and K, limiting plant growth and N uptake¹⁹. The extreme measure of excluding fertilisation and relying primarily on soil building during grass-clover leys and N fixation of soya, as done in the NOFERT treatment, not only results in a poor yield performance and soil organic carbon losses^{19,24} resulting to soil degradation, but also leads to adverse effects in terms of GHG emissions.

Despite the pronounced difference in maize yields, yield-scaled N₂O emissions did not differ between organic and non-organic systems in the DOK field trial (Table 2). This contrasts with the findings of Skinner *et al.*, who found higher yield-scaled N₂O emissions in organically managed soils⁶. The silage maize 2013 resulted with an average organic to non-organic yield gap of 27%. The long-standing average in the DOK, however, is only 9% for silage maize and 19% across all crops²⁵. Seufert *et al.* calculated in their global meta-analysis based on 316 observations an average organic to non-organic yield ratio of 0.75 for all crops and a ratio of 0.85 for maize (74 observations)⁹, corresponding to a yield gap of 15% which is in accordance with DOK averages. The dry weather conditions during summer 2013 in Switzerland (Supplementary Fig. S1) led to generally low yields for most crops. The yield performance of organically cultivated silage maize in the DOK was apparently more impaired (-26%) than in the non-organic systems (CONMIN: -13%; CONFYM: -10%) compared to the long-term yielding average. This impairment difference might be related to the cultivar “Colisee” which was originally bred for non-organic farming and judged to be suitable for organic farming under favourable conditions. The 56% lower emissions for BIODYN compared to CONFYM can be related to the 52% lower input, but the resulting yield gap of only 27% for BIODYN (Tables 1, 2, Supplementary Table S4) indicates that this system’s N efficiency is superior and thus contributes to GHG mitigation. Our results show that organic farming management does not per se lead to higher yield-scaled N₂O emissions.

In grass-clover, emissions were comparatively low with lowest emissions in BIODYN and CONFYM. Similar N_2O emissions under organic and non-organic in grass-clover were found elsewhere^{23,26,27}. As the GHG monitoring campaign started in the second year of the grass-clover ley, the calculation of the corresponding yield-scaled N_2O emissions was impossible. Furthermore, yields of the 5th and final cut in 2013 were rather low compared to the long-term average. The ley had to grow under very dry conditions in 2011 and could therefore never establish well. We decided against the calculation of emission factors also because mineralisation of the grass-clover ley introduced an additional source of N in all farming systems, thus questioning the explanatory power of the emission factor approach for single crops in systems with complex crop rotations which also has been concluded elsewhere²³.

Correlation analysis (Table 4) revealed that fertiliser N input and especially the addition of mineralised N to silage maize best explained cumulative N_2O emissions among the fertilised treatments. This is commonly observed, and a reduction of fertilising intensity is therefore suggested to be a good mitigation option³. The soil characteristics C_{mic} , SOC and pH (Supplementary Table S2) appeared to be further explaining variables, at least when NOFERT was included in the correlation analysis. The pH was the most robust variable explaining N_2O emissions, as its impact on N_2O holds true with or without the inclusion of the NOFERT dataset (Table 4). Rather high N_2O emission rates were recorded for NOFERT treatment which showed the lowest soil pH particularly during the maize and the green manure phase. The N_2O reduction process to N_2 may have been restricted in NOFERT soil in comparison to the other treatments. The pH dependence of the N_2O reduction process in soil has been investigated elsewhere²⁸ and has been determined recently in an accompanying laboratory experiment with soil from the DOK trial²⁹. The low pH in NOFERT seems to impede functionality of the relevant N_2O reductase enzyme in soil microorganisms³⁰. In line with this, BIODYN, the farming system with the highest soil pH, showed the lowest area- and yield-scaled N_2O emissions. We found no correlation between the pH from the starting phase in 1977 and the current N_2O emissions. This indicates that rather management-induced changes in soil pH over time contribute to the observed changes in N_2O emissions.

Assessing all systems including NOFERT we have further found good correlation of N_2O emissions with SOC and microbial biomass (Table 4). It has been shown in previous studies, that the contrasting farming management induced a gradient in SOC and microbial biomass in the DOK trial, with the lowest contents in NOFERT and the highest in BIODYN^{18,31}.

Studying the influence of the farming systems on microbial community structure, Esperschütz *et al.* detected distinct microbial communities in BIOORG and BIODYN compared to CONMIN and NOFERT, while samples originating from CONFYM formed a separate cluster¹⁶. Using a high-throughput pyrosequencing approach of bacterial and fungal ribosomal markers Hartmann *et al.* further revealed that manure addition was the most prominent factor influencing the formation of distinct bacterial and fungal communities in the different farming systems of the DOK trial¹⁷. However, functional gene quantification of soil microbes involved in N cycling at the end of the gas sampling campaign did not reveal substantial differences between farming systems. At best, N_2O reducers showed effects between single systems (Supplementary Table S5). Yet it remains unclear how the shift in community composition of soil microbes in the different farming systems affects soil functionality regarding N_2O production and reduction during cropping periods.

Cultivating soil as a habitat for soil organisms and not just as a substrate for plant growth addresses the goal of organic farming's strive for organic matter build-up stimulating soil fertility and nutrient cycling¹⁵ which apparently also influences the N_2O emission properties. Although organically managed soils rich in SOC and microbial biomass may show a higher potential for N_2O production than those from non-organic management under denitrifying conditions²⁹, this does not necessarily translate into the field situation with structured soils and system-specific fertilisation. In turn, this implies that adapted fertilisation strategy in organically managed soils prevents excessive N_2O emissions despite higher N_2O emission potential.

In contrast to the three comparative studies included in the meta-study of Skinner *et al.*⁸, we found no difference in area-scaled CH_4 emissions between organic and non-organic farming. The CONFYM soils and to a lesser extent also NOFERT soils, appeared to be CH_4 sources (Table 2). The systems BIODYN, BIOORG and CONMIN showed a pattern of CH_4 uptake as reported for upland soils elsewhere^{32–34}. It was shown that regular application of stacked cattle manure increases the biomass of methanogenic archaea³⁵. This can lead to considerable CH_4 emissions in combination with high soil water contents^{35,36}. In our study, this may explain high CH_4 emissions in CONFYM at the beginning of the silage maize vegetation phase in 2013 (Fig. 1). Manure application, ploughing and seedbed preparation during May occurred under cold and moist weather conditions (Supplementary Table S1; Supplementary Fig. S1). In contrast, application of composted organic materials were shown in an incubation experiment to transiently foster CH_4 oxidation activity in agriculturally managed soils³⁷, which could explain CH_4 uptake in BIODYN during silage maize cropping (Table 2). Anoxic conditions due to high soil water content disable CH_4 oxidation, which brings uptake to a standstill but allows methane from deeper layers to pass through upper soil horizons to the atmosphere³⁸. This may explain the methane release in NOFERT and BIOORG during silage maize cropping (Table 2). Events with more than 30 mm rainfall or consecutive days with rain in early June increased WFPS in those systems and BIODYN to around 80%, while WFPS in CONMIN and CONFYM was distinctively lower, most probably due to the higher water uptake of their crop (Fig. 1).

This is, according to our knowledge, the first study where soil GHG flux data resulting from bioorganic and biodynamic farming systems were compared. It turned out that BIODYN showed distinctly lower area-scaled N_2O emissions than BIOORG over the whole observation period. Even the yield-scaled N_2O emissions were in tendency lower in BIODYN. The main management difference between BIODYN and BIOORG consists of the type of solid manure applied. This is composted farmyard manure in BIODYN and rotted farmyard manure in BIOORG, although the amount of plant available N was nearly the same for both systems (Supplementary Table S4). Composting of farmyard manure as practiced in BIODYN may raise soil pH over stacked and rotted

manure application³⁹. BIODYN is also the farming system with highest SOC and microbial biomass values over the years^{16,18}.

Furthermore, van Groeningen *et al.* (2010) found that up to a threshold of 187 kg ha⁻¹ fertiliser-N applied, corresponding to the amount applied to BIOORG, emissions remained rather stable at low flux rates to increase sharply for higher application rates⁴⁰. They propose to focus on maximising crop N uptake together with balancing N application to crop N needs. These results are consistent to those of Shcherbak *et al.* who demonstrated an exponential increase of N₂O emissions for fertiliser applications exceeding crop requirements⁴¹.

We are aware, that for a comprehensive GHG assessment of a given farming system, not only the soil-derived emissions but also all other emissions caused by the production of different (synthetic/organic) fertilisers, energy use from farm machinery and emissions caused by livestock and manure management, need to be accounted for⁴². This, however, is beyond the scope of the current study. A previous LCA study, using experimental data of the DOK trial (1985–1998), found the area-scaled (26–35%) and product-based (2–17%) energy use for farm operations such as machinery, synthetic/organic fertilisation and plant protection to be lower in the two organic systems compared to CONFYM in dependence of the fertilisation intensity⁴³.

In our case, ecological intensification by the means of organic farming leads to the same yield-scaled N₂O emissions as found for non-organic farming systems. As, compared to non-organic systems, organic farming systems were reported to deliver greater ecosystem services and social benefits in a wider context, ecological intensification would not just address climate change mitigation but, at the same time, also the other main categories of agricultural sustainability, such as socio-economic aspects⁶.

Crowder and Reganold analysed the economic performance of organic compared to conventional farming with a meta-analysis encompassing data from 55 crops on five continents over 40 years⁴⁴. Taking current price premiums for organic products into account organic farming resulted to be with 22% to 35% more profitable and had a better benefit/cost ratio of 20% to 24% compared to non-organic. However, without price premiums organic farming resulted to be less profitable. Accounting for negative external environmental costs of agriculture, however, is likely to stimulate the expansion of organic agriculture⁴⁴ which currently covers less than 1% of global agricultural land⁴⁵.

Methods

Study site. The DOK long-term farming systems comparison trial, established 1978, is located (47°30'N; 7°32'E; 306 m asl) in Therwil, Switzerland. The soil is a Haplic Luvisol (15% sand, 70% silt and 15% clay) on deep deposits of alluvial Loess.

The trial is designed as a randomized split-block (fertilisation and crop shift), with four replicates arranged in a standard Latin square. It encompasses four farming systems – two organic ones, bio-dynamic, BIODYN (composted manure and slurry) and bioorganic, BIOORG (rotted manure and slurry); a conventional one (with manure, slurry and synthetic fertiliser) CONFYM - all of which are cultivated in a full (1.4 livestock units) as well as a half-fertilised treatment. The second conventional system (exclusively synthetic fertilisation) CONMIN – has only one treatment at normal fertilisation level adjusted to soil P and K contents. An unfertilised control, NOFERT, complements the comparison. Plant protection in the organic and the unfertilised system is based on mechanical weeding, indirect disease control measures and plant extracts together with bio-controls against insects, while in the non-organic systems herbicides, fungicides and pesticides are applied¹⁵.

Preparation of the different manure types is performed on identical amounts of fresh manure. Stacked manure is being compacted and thus undergoes an anaerobic fermentation at about 25 to 30 °C. Rotted manure is being shifted once after four weeks, while compost is turned on a weekly basis. Losses of dry matter, organic matter and nitrogen during preparation account to 22, 29 and 19% for stacked manure, 32, 47 and 30% for rotted manure and 42, 57 and 33% for composted manure⁴⁶.

The seven-year crop rotation is identical for all farming systems and is cultivated in three rotation units with a time shift of one year. The 6th crop rotation since the onset of the trial started 2013. The rotation encompasses silage maize followed by green manure of a *Brassica chinensis* x *Brassica rapa* breed over winter (first year), soybean (second year), winter wheat-1 and green manure (third year), potatoes (fourth year), winter wheat-2 (fourth and fifth year) and two years of a grass-clover ley (years 6 and 7)¹⁵.

The local climate is favourable for agriculture being rather mild and dry with annual (1981–2010) means of 10.5 °C and 842 mm rainfall (1961–1990: 9.6 °C, 778 mm) recorded at Basel/Binningen (BAS; WMO: 06601; 7°35'E; 47°32'N; 316 m asl) by the Federal Office for Meteorology and Climatology.

Monitoring period. CONMIN and the fully fertilised treatments of the farming systems BIODYN, BIOORG and CONFYM were chosen for investigation as they are representative for common agricultural practices in Switzerland. NOFERT was included as a control in the greenhouse gas measurement campaign at a later stage before ley termination in March 2013. Starting on 24 Aug 2012, GHG flux recordings, concomitant soil and agronomic analyses covered a sequence of grass-clover (256 days) – silage maize (140 days) – green manure (175 days) ending on 18 March 2014. Further details and farming operations are displayed in Supplementary Table S1, amount of fertiliser applied to grass-clover in Supplementary Table S3, and fertiliser rates for silage maize in Supplementary Table S4. Yields for silage maize are displayed in Table 3 together with yield-scaled N₂O emissions.

Supplementary Fig. S1 displays monthly means for temperature, rainfall and sunshine hours related to the norm period of 1980–2010. In 2012, the mean annual temperature was 10.9 °C and rainfall amounted to 1048 mm in the Basel area. During the first half of November 2012, extreme rainfall events led to an annual surplus of over a third of normal. In 2013, the mean annual temperature was 10.3 °C and rainfall amounted to 908 mm. The winter was unusually cold and prolonged until end of February. Spring was extremely cloudy, rainy and cool. With only half of the mean hours of sunshine it turned out to be the least sunny spring since 1833. Rainfall amounted to roughly 140% of normal. At the end of May 2013, the weather turned nearly to the other extreme with much

sunshine, occasionally hot, so that a lasting monthly temperature surplus compared to normal resulted until the end of October. Winter 2013/2014 was rather warm, nearly frost-free.

Greenhouse gas sampling. Soil gas fluxes of N₂O, CH₄ and CO₂ were sampled with manual closed chambers of 30 cm inner diameter and 0.7 L volume as described elsewhere^{47,48}. Soil collars were removed for tillage operations only. Chamber height was adjusted to the growing crop with tubular interfaces. Gas flux samplings were performed on a weekly base with additional samplings covering management operations. This extended manual sampling scheme has been shown to provide less than 10% deviation compared to near-continuous automated gas measurements⁴⁹. Samplings were restricted to between nine in the morning and noon (apparent solar time) when actual trace gas fluxes best represent the daily mean of the highly variable flux rates⁵⁰. One gas sampling consisted of four gas samples taken within 50 to 90 minutes with 20 ml plastic syringes. Gas samples were transferred into evacuated 12 ml vials creating an overpressure in vials which was needed to allow analysis via gas chromatography with an autosampler. Gas analysis involved HayeSep Q 80/100 columns at 100 °C oven temperature and N₂O detection with an electron capture detector (μECD), CO₂ reduction with a methanizer and subsequent detection of CO₂ and CH₄ with a flame ionisation detector (FID). Three calibration gases with concentrations from ambient to 10 times ambient were used. Fluxes were calculated with robust linear and non-linear models of the HMR function (package HMR) in R (R Core team) which was slightly adjusted as described elsewhere⁵¹. Cumulative gas emissions were integrated over time using the trapezoidal rule.

Soil sampling. Every gas sampling included a soil sampling (0–20 cm) bulked per treatment to determine soil water content and mineral soil nitrogen (N_{min}). Soil water content (WC) was determined gravimetrically with 24 h drying at 105 °C. Water filled pore space (WFPS) was calculated based on corresponding bulk densities. Soil extractions with 0.01 M CaCl₂-solution (1:4 w/v) were subject to 45 minutes of shaking and filtration before spectrophotometric analysis was employed to determine nitrite, nitrate and ammonium concentrations via continuous flow analysis. A basic sampling for microbial biomass, soil organic carbon and further variables was performed every year during spring time, shortly before starting the farming operations. Determination of soil pH was performed with a soil-water suspension, soil organic carbon concentration by wet oxidation (Walkley Black) and microbial biomass by chloroform fumigation extraction using 0.5 M K₂SO₄ (1:4 w/v) as described elsewhere¹⁸.

Statistical analysis. All calculations were accomplished in R software⁵². Field replicates of gas samples were pooled per plot for statistical analysis. Cumulated N₂O emissions and soil data of 2012 were assessed with a linear mixed effect model with field replication as random effect and clay content as covariate. Relations of cumulated N₂O emissions in maize with N input and soil data were calculated with the Pearson rank correlation (Hmisc package). Time series correlation of N₂O fluxes that were pooled per treatment, with ancillary soil data (n = 1 per treatment) was calculated with a linear mixed effect model with farming system treatments as random effect and sampling date per treatment as correlation term (repeated measurements, nlme package). Due to non-normal distribution, cumulated CH₄ fluxes were assessed with a Kruskal Wallis test for treatment and farming system differences (PCMCRA package).

Data Availability

The datasets generated during and/or analysed during the current study are available from the corresponding author on request.

References

- Smith, P. *et al.* Chapter 11 - Agriculture, forestry and other land use (AFOLU) In *Climate Change 2014: Mitigation of Climate Change, IPCC Working Group III Contribution to AR5 (Cambridge University Press, 2014)*.
- Smith, P. *et al.* Greenhouse gas mitigation in agriculture. *Philos Trans R Soc Lond B Biol Sci.* **363**, 789–813 (2008).
- Venterea, R. T. *et al.* Challenges and opportunities for mitigating nitrous oxide emissions from fertilized cropping systems. *Front Ecol Environ.* **10**, 562–570 (2012).
- Lorenz, K. & Lal, R. Chapter three - environmental impact of organic agriculture In *Advances in Agronomy* (ed. Donald, L. S.) **139**, 99–152 (Academic Press, 2016).
- Halberg, N., Panneerselvam, P. & Treyer, S. Eco-functional intensification and food security: Synergy or compromise? *Sustainable Agriculture Research.* **4**, 126–139 (2015).
- Reganold, J. P. & Wachter, J. M. Organic agriculture in the twenty-first century. *Nature Plants.* **2**, 15221–15229 (2016).
- Van Groenigen, J. *et al.* The soil N cycle: new insights and key challenges. *Soil.* **1**, 235–256 (2015).
- Skinner, C. *et al.* Greenhouse gas fluxes from agricultural soils under organic and non-organic management — a global meta-analysis. *Science of The Total Environment.* **468–469**, 553–563 (2014).
- Seufert, V., Ramankutty, N. & Foley, J. A. Comparing the yields of organic and conventional agriculture. *Nature.* **485**, 229–232 (2012).
- de Ponti, T., Rijk, B. & van Ittersum, M. K. The crop yield gap between organic and conventional agriculture. *Agricultural Systems.* **108**, 1–9 (2012).
- Van Kessel, C. *et al.* Climate, duration, and N placement determine N₂O emissions in reduced tillage systems: a meta-analysis. *Glob Change Biol.* **19**, 33–44 (2013).
- Lampkin, N. *Organic Farming* pp 701 (Farming Press, 1990).
- Turinek, M., Grobelnik-Mlakar, S., Bavec, M. & Bavec, F. Biodynamic agriculture research progress and priorities. *Renewable Agriculture and Food Systems.* **24**, 146–154 (2009).
- Chalker-Scott, L. The science behind biodynamic preparations: A literature review. *Horttech.* **23**, 814–819 (2013).
- Mäder, P. *et al.* Soil fertility and biodiversity in organic farming. *Science.* **296**, 1694–1697 (2002).
- Esperschütz, J., Gattinger, A., Mäder, P., Schloter, M. & Fließbach, A. Response of soil microbial biomass and community structures to conventional and organic farming systems under identical crop rotations. *FEMS Microbiology Ecology.* **61**, 26–37 (2007).
- Hartmann, M., Frey, B., Mayer, J., Mäder, P. & Widmer, F. Distinct soil microbial diversity under long-term organic and conventional farming. *ISME J.* **9**, 1177–1194 (2015).

18. Fließbach, A., Oberholzer, H. R., Gunst, L. & Mäder, P. Soil organic matter and biological soil quality indicators after 21 years of organic and conventional farming. *Agriculture, Ecosystems and Environment*. **118**, 273–284 (2007).
19. Mayer, J. *et al.* Productivity, quality and sustainability of winter wheat under long-term conventional and organic management in Switzerland. *European Journal of Agronomy*. **65**, 27–39 (2015).
20. Gomiero, T., Pimentel, D. & Paoletti, M.-G. Environmental impact of different agricultural management practices: Conventional vs. organic agriculture. *Critical Reviews in Plant Sciences*. **30**, 95–124 (2011).
21. De Gryze, S., Albarracín, M., Catalá-Luque, R., Howitt, R. & Six, J. Modeling shows that alternative soil management can decrease greenhouse gases. *Calif Agr*. **63**, 84–9 (2009).
22. Kaiser, E. & Ruser, R. Nitrous oxide emissions from arable soils in Germany - An evaluation of six long-term field experiments. *Journal of Plant Nutrition and Soil Science*. **163**, 249–260 (2000).
23. Brożyna, M. A., Petersen, S. O., Chirinda, N. & Olesen, J. E. Effects of grass-clover management and cover crops on nitrogen cycling and nitrous oxide emissions in a stockless organic crop rotation. *Agriculture, Ecosystems & Environment*. **181**, 115–126 (2013).
24. Leifeld, J., Reiser, R. & Oberholzer, H.-R. Consequences of conventional versus organic farming on soil carbon: Results from a 27-year field experiment. *Agronomy Journal*. **101**, 1204–1218 (2009).
25. Mäder, P., Fließbach, A., Niggli, U., Gunst, L. & Mayer, J. Ertrag von Mais und Sojabohnen im biologischen und konventionellen Anbausystem des DOK-Versuchs In *Beiträge zur 12. Wissenschaftstagung Ökologischer Landbau* (eds Neuhoff, D. *et al.*) 52–53 (Verlag Dr. Köster, 2013).
26. Syvaesalo, E., Regina, K., Turtola, E., Lemola, R. & Esala, M. Fluxes of nitrous oxide and methane, and nitrogen leaching from organically and conventionally cultivated sandy soil in western Finland. *Agriculture, Ecosystems & Environment*. **113**, 342–348 (2006).
27. Flessa, H. *et al.* Integrated evaluation of greenhouse gas emissions (CO₂, CH₄, N₂O) from two farming systems in southern Germany. *Agriculture, Ecosystems & Environment*. **91**, 175–189 (2002).
28. Čuhel, J. *et al.* Insights into the effect of soil pH on N₂O and N₂ emissions and denitrifier community size and activity. *Appl. Environ. Microbiol.* **76**, 1870–1878 (2010).
29. Krause, H.-M. *et al.* Long term farming systems affect soils potential for N₂O production and reduction processes under denitrifying conditions. *Soil Biol. Biochem.* **114**, 31–41 (2017).
30. Bakken, L. R., Bergaust, L., Liu, B. & Frostegard, A. Regulation of denitrification at the cellular level: a clue to the understanding of N₂O emissions from soils. *Philos Trans R Soc Lond B Biol Sci*. **367**, 1226–1234 (2012).
31. Gattinger, A. *et al.* Enhanced top soil carbon stocks under organic farming. *Proceedings of the National Academy of Sciences*. **109**, 18226–18231 (2012).
32. Conrad, R. Soil microorganisms as controllers of atmospheric trace gases (H₂, CO, CH₄, OCS, N₂O, and NO). *Microbiology and Molecular Biology Reviews*. **60**, 609–640 (1996).
33. Hütsch, B. W. Methane oxidation in non-flooded soils as affected by crop production — invited paper. *European Journal of Agronomy*. **14**, 237–260 (2001).
34. Hiller, R. V. *et al.* Anthropogenic and natural methane fluxes in Switzerland synthesized within a spatially explicit inventory. *Biogeosciences*. **11**, 1941–1959 (2014).
35. Gattinger, A. *et al.* Traditional cattle manure application determines abundance, diversity and activity of methanogenic Archaea in arable European soil. *Environmental Microbiology*. **9**, 612–624 (2007).
36. Radl, V. *et al.* Effects of cattle husbandry on abundance and activity of methanogenic archaea in upland soils. *ISME J*. **1**, 443–452 (2007).
37. Ho, A. *et al.* Unexpected stimulation of soil methane uptake as emergent property of agricultural soils following bio-based residue application. *Glob Chang Biol*. **21**, 3864–3879 (2015).
38. Blagodatsky, S. & Smith, P. Soil physics meets soil biology: Towards better mechanistic prediction of greenhouse gas emissions from soil. *Soil Biol. Biochem.* **47**, 78–92 (2012).
39. Amon, B., Amon, T., Boxberger, J. & Alt, C. Emissions of NH₃, N₂O and CH₄ from dairy cows housed in a farmyard manure tying stall (housing, manure storage, manure spreading). *Nutrient Cycling in Agroecosystems*. **60**, 103–113 (2001).
40. Van Groenigen, J. *et al.* Towards an agronomic assessment of N₂O emissions: a case study for arable crops. *European Journal of Soil Science* **61**, 903–913 (2010).
41. Shcherbak, I., Millar, N. & Robertson, G. P. Global meta-analysis of the nonlinear response of soil nitrous oxide (N₂O) emissions to fertilizer nitrogen. *Proceedings of the National Academy of Sciences*. **111**, 9199–9204 (2014).
42. Meier, M. S. *et al.* Environmental impacts of organic and conventional agricultural products – Are the differences captured by life cycle assessment? *Journal of Environmental Management*. **149**, 193–208 (2015).
43. Nemeček, T., Dubois, D., Huguenin-Elie, O. & Gaillard, G. Life cycle assessment of Swiss farming systems: I. Integrated and organic farming. *Agricultural Systems* **104**, 217–232 (2011).
44. Crowder, D. W. & Reganold, J. P. Financial competitiveness of organic agriculture on a global scale. *PNAS*. **112**, 7611–7616 (2015).
45. Willer, H. & Lernoud, J. *The World of Organic Agriculture. Statistics and Emerging Trends 2018.* Research Institute of Organic Agriculture (FiBL), Frick, and IFOAM – Organics International, Bonn (2018).
46. Berner, A., Scherrer, D. & Alföldi, T. Stickstoffeffizienz von unterschiedlich aufbereiteten Misten in einer Ackerfruchtfolge auf Lösslehm. *Beiträge zur 4. Wissenschaftstagung Ökologischer Landbau* (eds Köpke, U. & Eisele, J. -A.) (Verlag Dr. Köster, 1997).
47. Flessa, H., Dörsch, P. & Beese, F. Seasonal variation of N₂O and CH₄ fluxes in differently managed arable soils in southern Germany. *Journal of Geophysical Research*. **100**, 23115–23124 (1995).
48. Krauss, M. *et al.* Impact of reduced tillage on greenhouse gas emissions and soil carbon stocks in an organic grass-clover ley - winter wheat cropping sequence. *Agriculture, Ecosystems & Environment*. **239**, 324–333 (2017).
49. Flessa, H. *et al.* N₂O and CH₄ fluxes in potato fields: automated measurement, management effects and temporal variation. *Geoderma*. **105**, 307–325 (2002).
50. Alves, B. J. R. *et al.* Selection of the most suitable sampling time for static chambers for the estimation of daily mean N₂O flux from soils. *Soil Biol. Biochem.* **46**, 129–135 (2012).
51. Leiber-Sauheitl, K., Fuss, R., Voigt, C. & Freibauer, A. High greenhouse gas fluxes from grassland on histic gleysol along soil carbon and drainage gradients. *Biogeoscience Discussions*. **10**, 11283–11317 (2007).
52. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, <http://www.R-project.org/> (2013).

Acknowledgements

We gratefully acknowledge the financial support for this project provided by the Swiss Federal Offices for the Environment (FOEN) and Agriculture (FOAG). A. G. received additional funding from the Mercator Foundation Switzerland (Grant No. 2011-0294) and Swiss National Science Foundation within the frame of the National Research Program “Soil as a Resource” (Grant No. 406840_143137). We particularly thank the DOK field teams and the farmer advisory board of the DOK, the lab staff, practitioners and researchers engaged at FiBL, University of Basel and Agroscope for excellent collaboration.

Author Contributions

A.G., P.M., C.S., M.K., J.M. and M.G.A.v.d.H. designed research; C.S., A.G., H.M.K. and M.K. performed research; C.S., H.M.K. and M.K. analysed data; and C.S., A.G., H.M.K. M.K. and P.M. wrote the paper. A.G. is the lead scientist for this project.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-018-38207-w>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019