

# UC Santa Barbara

## UC Santa Barbara Previously Published Works

### Title

The impacts of increasing drought on forest dynamics, structure, and biodiversity in the United States.

### Permalink

<https://escholarship.org/uc/item/0dg4t07p>

### Journal

Global change biology, 22(7)

### ISSN

1354-1013

### Authors

Clark, James S  
Iverson, Louis  
Woodall, Christopher W  
et al.

### Publication Date

2016-07-01

### DOI

10.1111/gcb.13160

Peer reviewed

# **The impacts of increasing drought on forest dynamics, structure, and biodiversity in the United States**

James S. Clark, Louis Iverson, Christopher W. Woodall, Craig D. Allen, David M. Bell, Don C. Bragg, Anthony W. D'Amato, Frank W. Davis, Michelle H. Hersh, Ines Ibanez, Stephen T. Jackson, Stephen Matthews, Neil Pederson, Matthew Peters, Mark Schwartz, Kristen M. Waring, and Niklaus E. Zimmermann

## Affiliations:

Craig D. Allen, U.S. Geological Survey, Fort Collins Science Center, Jemez Mountains Field Station, Los Alamos, New Mexico, USA.

David M. Bell, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Corvallis, OR.

Don C. Bragg, U.S. Department of Agriculture, Forest Service, Southern Research Station, Monticello, AR.

Anthony W. D'Amato, Rubenstein School of Environment and Natural Resources, University of Vermont.

Frank W. Davis, Bren School of Environmental Science and Management, University of California, Santa Barbara.

Michelle H. Hersh, Department of Biology, Sarah Lawrence College, New York.

Ines Ibanez, School of Natural Resources and Environment, University of Michigan.

Stephen T. Jackson, U.S. Geological Survey, Southwest Climate Science Center and Department of Geosciences, University of Arizona.

Stephen Matthews, School of Environment and Natural Resources, Ohio State University.

Neil Pederson, Harvard Forest, Petersham, MA.

Matthew Peters, U.S. Department of Agriculture, Forest Service, Northern Research Station, Delaware, OH.

Mark W. Schwartz, Department of Environmental Science and Policy at UC Davis, California.

Kristen M. Waring, School of Forestry at Northern Arizona University, Arizona.

Niklaus E. Zimmermann, Swiss Federal Research Institute WSL, Birmensdorf, Switzerland.

## **Abstract**

We synthesize insights from current understanding of drought impacts at stand to biogeographic scales, including management options, and we identify challenges to be addressed with new research. Large stand-level shifts underway in western forests already are showing the importance of interactions involving drought, insects, and fire. Diebacks, changes in composition and structure, and shifting range limits are widely observed. In the eastern US, the effects of increasing drought are becoming better understood at the level of individual trees, but this knowledge cannot yet be confidently translated to predictions of changing structure and diversity of forest stands. While eastern forests have not experienced the types of changes seen in western forests in recent decades, they too are vulnerable to drought and could experience significant changes with increased severity, frequency, or duration in drought. Throughout the continental United States, the combination of projected large climate-induced shifts in suitable habitat from modeling studies and limited potential for the rapid migration of tree populations suggests that changing tree and forest biogeography could substantially lag habitat shifts already underway.

Forest management practices can partially ameliorate drought impacts through reductions in stand density, selection of drought-tolerant species and genotypes, artificial regeneration, and the development of multi-structured stands. However, silvicultural treatments also could exacerbate drought impacts unless implemented with careful attention to site and stand characteristics. Gaps in our understanding should motivate new research on the effects of interactions involving climate and other species at the stand scale and how interactions and multiple responses are represented in models. This assessment indicates that, without a stronger empirical basis for drought impacts at the stand scale, more complex models may provide limited guidance.

## **Introduction**

Drought is a departure from the mean climate for a region and represents moisture limitation resulting from below-average precipitation, high temperatures, or both. At the time of this writing, drought conditions have continued over much of the continental United States (US) for up to four years. Combined warming and variable precipitation have increased forest drought severity in the last two decades, not only in the West, but also the South and the Lake States (Box 1). Prolonged drought affects the distributions of species, the biodiversity of landscapes, wildfire, net primary production, and virtually all goods and services provided by forests. Understanding how climatic changes already in progress will affect forests can help us anticipate some of these broader impacts. The synthesis that follows finds that vulnerabilities extend beyond the recent well-publicized forest diebacks in western states to include perhaps all US forests.

Our summary of drought effects emphasizes the fundamental scale both for management and community ecology, the forest stand (O'Hara and Nagel 2013). We build from what can be

learned about climate effects on individual trees, but our principal goal is to anticipate consequences for forest structure and composition, the *size-species distribution* (SSD; Box 2). The SSD is the distribution of trees across species and size classes. The SSD results from interactions of individuals, as each tree responds to local conditions and weather. Competition and climate affect the species and size classes that make up stands in different ways. There is feedback—the structure itself determines how the SSD will respond to drought, through shading and competition for soil moisture. Biogeographic patterns in SSD emerge from these individual responses and interactions with others. Management aims to modify SSDs (e.g., targeted thinning and regeneration) to meet specific resource objectives. However, because SSD responds to climate change as a joint distribution of individuals of many species and size classes, our ability to anticipate impacts and offer solutions to forest managers has been challenged.

This synthesis of current understanding begins with a summary of the extensive but mostly indirect evidence, from studies of individual trees to forest stands, across landscapes and regions, from short-term observations to the paleoecological record. We consider both the responses to individual drought events and the effects of conditions that could be more arid on average than today. Then we compare and contrast evidence available at the individual tree and stand scales, including why the latter is more critical, but harder to obtain. This is followed by a summary of what has been learned from that evidence for forest stands and for biogeography and how management practices might adapt to more frequent drought. Finally, we address critical research gaps between our growing knowledge on individual tree responses (in contrast to the stand scale) and where the relevant forecasts are needed. Recommendations include the assembly and parameterization of models based on SSD data capable of predicting at the SSD scale.

## **Consequences for forest stands: individual responses translate to abundance and size structure**

Not surprisingly, the effects of drought on forest stands are difficult to anticipate due to the novelty of projected new climates and the complexity of interactions across the SSD, including migration. Furthermore, the additional complexity and nonlinear responses associated with forestry practices and how they influence micro climate are also poorly understood (e.g., Bright et al. 2015). Changing temperatures and precipitation patterns will produce novel combinations of drought frequency, intensity, and seasonality (Wehner et al. 2011, Dai 2012). Tree populations can disperse and adapt to local climates, including drought stress (Savolainen et al., 2007; Aitken et al., 2008; Montwé et al., 2015). As tree populations arrive in new locations, they interact with existing populations and form new communities. Some will outrun mutualists, competitors, and natural enemies, and encounter new ones—processes that are too slow, too small, or too large to observe directly and are therefore difficult to study with experiments. Much of what is known relies on observational data or is inferred from model simulations, both of which provide valuable insights and have inherent limitations.

The most striking result from this review was the large gap between knowledge of drought impacts on individual trees (much) versus responses of forest stands (almost none). To see why individual responses do not readily extrapolate to the SSDs of stands (Box 2), consider how the SSD mediates a climate response. For co-dominant trees in crowded stands (trees with crowns in the main canopy), growth and mortality are dominated by competition. Canopy individuals that otherwise might respond positively to a moist growing season are constrained by competing neighbors that also benefit. At the stand level, mortality can increase as a result of favorable conditions—climate and site conditions that benefit individual tree health can increase stand mortality rates, depending on the SSD (e.g., Clark et al. 2014b). Mechanistically, this positive relationship arises because self-thinning is driven by growth—the faster the growth, the sooner resource limitations are reached, and the higher the mortality rate (Assmann 1970). Conversely, drought that depresses growth of individual trees can also decrease crowding pressure by slowing the rate of resource consumption. Unfavorable climate effects could be mitigated by stand characteristics through the active manipulation of stocking (e.g., thinning) or supplementation of limiting resources (e.g., irrigation) (D’Amato et al. 2013, Grant et al. 2013, Erickson and Waring 2014, Dobrowski et al. 2015).

The knowledge gap between individual trees versus stands is important because ecologists and foresters more often need to understand and predict responses of stands than individual trees. The gap comes from the challenge of observing and estimating whole stand responses, and it helps to explain why models for stands rely on parameters from individual trees. Using individuals to predict stands is an example of ‘Simpson’s Paradox’ or the ‘ecological fallacy’. This approach does not permit probabilistic prediction, because individuals within an SSD are interacting with one another. Their joint distribution of responses can only be predictive if it is observed and estimated as a joint distribution. For example, in contrast to co-dominant trees that experience high competition for light, the tallest (dominant) trees with emergent crowns may respond more directly to climate. The rare individuals that make up the right-most diameter extremes (largest trees) in Figure B2.1 are the focus of many tree-ring studies, but they rarely appear in small (0.0672-ha) Forest Inventory and Analysis (FIA) plots (note that trees are sampled on the larger 0.4-ha plots in Western states). Best represented in plot-based studies are the smallest size classes, which, in crowded stands, can have growth rates that are limited both by light and moisture. The large number of positive interactions between light and drought result from the fact that individuals not severely light-limited can respond most to climate variation (Clark et al. 2014b).

### ***Tree growth and mortality patterns in the eastern US***

Despite recent attention to large diebacks in the West, eastern forests are also vulnerable, not only in upland habitats (Abrams 1990, Graumlich 1993, Pederson et al. 2012a) but also in bottomlands and coastal wetlands (Stahle and Cleaveland 1992, Cook et al. 1999). Even where drought does not directly kill trees, the effect of reduced vigor on competitive ability affects

forest composition and structure. The question is, which effects will be most severe, how, and on which parts of the landscape? After all, the growth-related drought responses of tree species are diverse (Fig. 1). For example, the drought sensitivity of some pine species is high in the southeastern U.S. region (Schumacher and Day 1939, Cook et al. 2001, Henderson and Grissino-Mayer 2009, Clark et al. 2014b), while growth of many non-oak hardwoods shows intermediate drought sensitivity (Klos et al. 2009, Clark et al. 2013, Pederson et al. 2013). Combined high temperatures and low moisture could benefit oaks (*Quercus* spp.), as drier than normal conditions tend to have less impact on oak growth rates (Elliot and Swank 1994, Klos et al. 2009, Clark et al. 2011, 2014a, Brzostek et al. 2014), perhaps related to physiology and rooting (Abrams 1990, Abrams and Kubiske 1990, Iverson et al. 2008b). Hence, with increasing drought in the Upper Midwest and Lake States, drought-tolerant pines and oaks may replace drought-intolerant quaking aspen (*Populus tremuloides*), bigtooth aspen (*Populus grandidentata*), paper birch (*Betula papyrifera*), and some boreal and lowland conifers (Scheller and Mladenoff 2008, Handler et al. 2014).

Opportunistic reports of mortality following drought are common (Hough and Forbes 1943, Parshall 1995), but connections between drought and tree death are more difficult to quantify than those for tree growth. Extended morbidity can precede death, a legacy of low vigor spanning decades (Wyckoff and Clark 2002, Anderegg et al. 2013a, b, Berdannier and Clark 2015), potentially related not only to repeated drought (Pedersen 1998, Voelker et al. 2008, Pederson et al. 2014), but also to other risk factors that occur during sample intervals, which might be from one to ten years in many studies. Attribution of death to drought is thus challenging. A synthesis of plot data spanning 50 years, four Midwest states, and 48,000 stems did not find a link between precipitation and mortality, highlighting instead the importance of competition (Yaussy et al. 2013). A number of large studies using FIA data suggest geographic relationships between drought and mortality. Climate variables emerge as weak predictors of mortality at best (Dietze and Moorcroft 2011, Lines et al. 2010, Gustafson and Sturtevant 2013), and patterns may be hard to interpret. For example, a tendency toward higher mortality rates in warm climates is expected on the basis of higher productivity in warm climates—partly explained by the fact that high growth is attended by rapid thinning (Assmann 1970, Clark 1990). This relationship between temperature and mortality does not necessarily constitute a threat of climate change—geographic variation in mortality rate with average temperature need not indicate vulnerability to high temperature (Zhu et al. 2014).

The interactions involving competition and drought could contribute to habitat shifts. A positive interaction between a drought index such as Palmer Drought Severity Index (PDSI) (low PDSI indicates drought) and local moisture status or light availability means that the largest response to PDSI occurs on moist sites and high light (low competition), respectively. Klos et al.'s (2009) suggestion that dense stands may experience the most severe impacts agrees with the positive interaction between drought and competition found at the stand level in the upper Midwest (Fig. 2) and at the tree scale for many species in the eastern US (Clark et al. 2011, 2014b). The latter

study further found positive interactions between drought and local moisture status for many species (e.g., Fig. 3). The possibility that moist sites will provide refuges if climate becomes more xeric (e.g., Frelich and Reich 2010) represents a negative interaction, with the largest response to drought on dry sites. The alternative positive interaction could result from competition—the water-demanding species on wet sites fully utilize abundant moisture supply and thus are especially vulnerable when moisture availability declines. Large growth and fecundity responses to drought in southeastern forests could occur initially for trees at high moisture levels, where leaf area, and thus, moisture demand is greatest (Fig. 4). Mesic sites might see large transitions due to the fact that they also support sensitive species dependent on abundant moisture (Elliot and Swank 1994, Clark et al. 2014b). Still another possibility is that sensitivity could be highest on sites of intermediate moisture (e.g., Dormana et al. 2013). Moreover, the sign of the interaction between drought index and local drainage might shift with time, from short-term positive (loss of moisture demanding species on mesic sites) to negative (eventually the moist sites provide refuges for some species). Sequential drought will have impacts that differ from individual droughts as stands progressively respond (Miao et al. 2009).

Other soil properties can strongly influence the severity of drought. For example, fragipan soils in some pine flatwoods of the West Gulf Coastal Plain restrict root depth and access to deep moisture (Wackerman 1929, Rahman et al. 2006). Drought may also operate differently in stands of different density and age (Esper et al. 2008). If leaf area decreases during drought then understory irradiance increases. For trees beyond the seedling stage Luo and Chen (2013) argue that warming has the greatest impact on mortality of young trees, but there are also reports that old trees show the strongest responses to climate for *Picea glauca* (Wang et al. 2006) and *Quercus robur* (Rozas 2005). Klos et al. (2009) likewise found that the effects of drought on growth and survival might increase with stand age in the Southeast. Due to the large sample interval in many climate-mortality studies, evidence is equivocal (see above). The disparate results could also indicate the importance of unobserved variables that covary with density and stand age (D'Amato et al. 2013).

Even in the eastern US, drought can interact with fire to shape forest dynamics. Over the last century, much of the eastern US has experienced pluvials rather than megadroughts (Stahle et al. 1988, Stahle and Cleaveland 1992, Booth et al. 2006, 2012, Cook et al. 2010, McEwan et al. 2011, Pederson et al. 2013) but fires can occur even during brief periods of low precipitation, high temperatures, or both (Clark 1989, Lynch and Hessl 2010, Lafon and Quiring 2012). Excluding fire has long been a management priority of many landowners and agencies. If this history of fire suppression is responsible for reduced oak regeneration in the East, then climate trends otherwise favorable for oak may be offset by fire suppression. However, evidence that temperate forest stands may see a long-term increase in oaks (Bachelet et al. 2003, Clark et al. 2014b) presents an apparent paradox, in light of the fact that oak recruitment has declined in many regions (Abrams 2003, Fei et al. 2011). Fire suppression can lead to a 'mesophication' as forest canopies close (Nowacki and Abrams 2008) and may explain why oak regeneration

appears to decline relative to that of red (*Acer rubrum*) and sugar (*Acer saccharum*) maples in recent decades (Abrams 1994, 1998, Hutchinson et al. 2008, Iverson et al. 2008a, Fei et al. 2011, Brose et al. 2013). A decrease in flammability may have followed the loss of American chestnut (*Castanea dentata*) from eastern forests (Engber and Varner 2012, Kreye et al. 2013), although human increases in ignition, alteration of fuels, and active suppression make it difficult to characterize pre-settlement fire regimes (Clark and Royall 1996, Parshall and Foster 2003, Guyette et al. 2006).

Taken together, many species are vulnerable to drought in eastern forests. How this vulnerability at the individual scale translates into future forest composition and structure remains uncertain. For instance, the combination of climate, land-use, plant-animal interactions, and fire suppression may have contributed to recent maple recruitment, but this could be reversed by increasing drought (McEwan et al. 2011, Belden and Pallardy 2009, Woodall et al. 2009).

### ***Tree growth and mortality patterns in the western US***

Unlike the East, where drought effects on forest stands are less well documented than the physiological responses of individual trees, the West provides alarming examples of widespread stand replacement, directly or indirectly related to the recent combination of drought and warmer temperatures – “hotter drought” (Allen et al. 2015). Stand- to region-level consequences of hotter drought and forest dieback in the West are now well-documented (e.g, Breshears et al. 2005, van Mantgem et al. 2009, Worrall et al. 2013) (Box 1). Extensive drought across much of the western US and adjoining Canada coincides with declining tree growth, often followed by mortality (Allen et al. 2010, Meddens et al. 2012, Williams et al. 2013, Hicke et al. 2013, Joyce et al. 2014, O’Connor 2013, Peters et al. 2014, 2015).

Increased vulnerability of trees and forests to water stress and mortality risk from warmer droughts is a global phenomenon, well-illustrated in the western U.S. (Allen et al. 2015). High temperatures can increase drought-induced mortality in piñon (Adams et al. 2009) and are especially challenging for seedlings (Kolb and Robberecht 1996, Chmura et al. 2011). In Arizona and New Mexico, high temperatures combined with droughts coincide with widespread mortality of mesic montane tree species (Mueller et al. 2005, Gitlin et al. 2006, Williams et al. 2010, Ganey and Vojta 2011) and patchy die-offs in *Pinus edulis* (Breshears et al. 2005, 2009). Warming is considered most important for seasonal soil water balance due to changes in snowpack dynamics or evapotranspiration (Williams et al. 2013). It contributes to the growth and geographic expansion of insect pest populations (Bentz et al. 2010, Williams et al. 2013, Fig. 5). Varying water deficits appear to be primary drivers of variation in tree recruitment and mortality (Rapacciuolo et al. 2014). A combination of high temperatures during the growing season and low winter-spring precipitation of the previous year can explain much of the variation in conifer growth rates in the Southwest (primarily *Pinus edulis*, *Pinus ponderosa*, and *Pseudotsuga menziesii*) (Williams et al. 2013) and northern California (*Abies concolor*, *Abies magnifica*, *Pinus lambertiana*, *Pinus ponderosa*, and *Pseudotsuga menziesii*) (Yeh and Wensel 2000). Similar relationships between moisture, heat, and growth variation are observed for *Picea*



*glauca* in interior Alaska (Barber et al. 2000), for *Pseudotsuga menziesii* in the central and southwest Rocky Mountains and Mexico (Chen et al. 2010), and for *Populus tremuloides* in western Canada (Hogg et al. 2005). Responses suggest declining growth rates with increasing drought conditions for the western U.S. during the 21<sup>st</sup> century, particularly for the Southwestern U.S. (Williams et al. 2010, 2013).

Interactions between drought, fire, climate change, and human fire suppression have altered SSDs in forests throughout the western US (Westerling et al. 2006, Allen 2007, Littell et al. 2009, Joyce et al. 2014). Fire suppression has led to increased fuel accumulation and a shift to high-density/small diameter stands, from the foothills to the subalpine zone of the Sierra Nevada (Parsons and deBenedetti 1979, Dolanc et al. 2013, Lutz et al. 2009) and extensive semi-arid woodlands and ponderosa pine in the Southwest (e.g., Covington and Moore 1994, Brown and Wu 2005, Mast and Wolf 2006, Fulé et al. 2009). Historically low-density forest conditions resulted from the direct and indirect effects of low moisture (McDowell et al. 2006). The modern shift to high-density forests exacerbates moisture stress during drought. High-severity fires now occur in stands that historically supported frequent, low-severity fires (Barton 2002, Savage and Mast 2005, Goforth and Minnich 2008, Savage et al. 2013), and recent high-severity fires are strongly correlated with forest drought stress (e.g., Fig. 5). Projected warming in the Yellowstone region could increase fire frequency to the point where lodgepole pine (*Pinus contorta*) stands, historically characterized by infrequent fires, are replaced by woodland or non-forest vegetation (Westerling et al. 2011).

Because drought weakens tree defenses, it also makes trees vulnerable to insect attacks and pathogens (Raffa et al. 2008, Weed et al. 2013). The interaction between recent drought and bark beetle outbreaks is clear in the Southwest, where severe drought conditions from ca. 2000-2013 have resulted in widespread tree mortality from bark beetles (Negrón et al. 2009, Ganey and Vojta 2011, Williams et al. 2010, 2013). Defoliator populations may also benefit from drought-stressed trees, particularly on xeric sites (Weed et al. 2013). However, insect-drought interactions may not be intuitive and resistance to insect herbivory may yield lower drought tolerance and higher mortality rates during subsequent drought events (Sthultz et al. 2009). Early successional species that colonize after bark beetle infestations or wildfire may become more frequent in some areas (Pelz and Smith 2013, Shinneman et al. 2013).

Large diebacks have the potential to change species distributions more rapidly than has occurred in the past (Swetnam and Betancourt 1998, Gray et al. 2006, Millar et al. 2012, Fellows and Goulden 2012). Drought-induced mortality in the 1950s is probably responsible for extensive upslope retreat of ponderosa pine in New Mexico (Allen and Breshears 1998) and *Juniperus deppeana* in southeastern Arizona (Brusca 2013). Recent episodes of drought-induced mortality in California extend from high-elevation *Pinus albicaulis* in the Sierra Nevada (Millar et al. 2012) to southern California conifers (Minnich 2007), and a rapid redistribution of coniferous and broadleaf species occurred in southern California mountains during droughts of the early 2000s (Fellows and Goulden 2012). The extent and severity of drought impacts on western

forests raises concern for biodiversity and carbon storage (Gonzalez et al. 2015). The widespread nature of recent drought and its impacts suggest transformations that will have far-reaching consequences.

## **Critical impacts of drought on tree recruitment**

The preceding summaries of eastern and western forests focused on growth and mortality responses to drought rather than on recruitment, mostly because the relationship between drought and recruitment is poorly understood. An example of this limited understanding is the recent decline in oak regeneration (Fig. 6) at a time when the eastern US has experienced higher moisture deficits than the two preceding decades (Fig. B2.2). Recruitment warrants special consideration, both for its central role in decadal-scale responses to drought and because it has been especially difficult to study and predict across multiple scales (Dobowski et al. 2015). Most empirical research on climate effects on seed production are limited to a few years (or less) and a few small study plots (Clark et al. 1999). Some of the longer studies focus on interannual variation, but few provide evidence for decade-scale effects of increasing drought.

Drought influences tree recruitment (and therefore future forest composition) through numerous mechanisms. For some species, drought severely curtails fecundity, limits seed germination, and increases the mortality of shallow-rooted seedlings. Drought effects on fecundity are further complicated by feedbacks with other factors that drive masting cycles, seed predation, and disturbance regimes that disproportionately impact new germinants. Germination, establishment, and early survival are especially susceptible to environmental variation (Harper 1977, Grubb 1977, Silvertown 1987, Ibáñez et al. 2007). Susceptibility of juvenile trees may be particularly acute in dry regions where recruitment is already episodic (Brown and Wu 2005, Jackson et al. 2009). High mortality of seedlings suggests a bottleneck on population growth rate, but direct evidence for its effects on fitness of many interacting species is lacking.

The development of moisture limitation over successive years appears especially important for fecundity. In general, female function in trees is often stimulated by resources, including moisture (Perez-Ramos et al. 2010), CO<sub>2</sub> (LaDeau and Clark 2001), and light availability (Clark et al. 2014b). Seed production of many species shows positive interactions between moisture and light, with trees at high light levels showing the greatest response to moisture availability (Clark et al. 2014b). Warm, dry weather can be beneficial during flower induction the year before seeds ripen (Pucek et al. 1993, Houle 1999), a situation imposed artificially by water restriction in some fruit crops (Owens 1995). This effect may be enhanced if dry conditions follow a wet year (Piovesan and Adams 2001). Drought-induced increases in fecundity may be followed by reduced seed production up to several years thereafter (Innes et al. 1994, Bréda et al. 2006). Increases in late summer temperatures may negatively affect seed cone initiation, which for piñon pine led to a 40% reduction in seed cone production over the past 30 years (Redmond et al. 2012). Furthermore, year-to-year volatility and high spatial variation that comes with the many

feedbacks involving weather, competitors, fungal symbionts, cone and seed insects, and pathogens (e.g. Bell et al. 2014) make this response difficult to quantify.

Interactions involving drought and the biotic environment contribute to recruitment variation following disturbance, e.g., canopy gaps, fires, landslides, ice storms, timber harvesting, and pest outbreaks (e.g., Savage et al. 1996, Brown and Wu 2005, Pederson et al. 2008), and they affect composition, structure, and function for many years (Cooper-Ellis et al. 1999, Dietze and Clark 2008, Kayes and Tinker 2012). Examples of the interactions that can occur between disturbance and moisture availability include the increased recruitment near the prairie-forest ecotone in Minnesota during the 1930s drought (Shuman et al. 2009).

Interactions involving moisture availability and pathogen attack are especially important at the recruitment stage. Seedling mortality during the first year can be high due to damping off, often most severe in shaded understories (Hood et al. 2004, Ichihara and Yamaji 2009). Moist conditions that promote fungal infection (Desprez-Loustau et al. 2006) can also benefit the host plant (Hersh et al. 2012). Combined effects may depend on the pathogen's mode of attack and on the degree of host stress (Desprez-Loustau et al. 2006, Jactel et al. 2012, Oliva et al. 2014). Many pathogens can tolerate a wider range of water stress than the plants they infect, and the combination of pathogen infection and moisture stress on host trees can increase disease severity (Desprez-Loustau et al. 2006). Drought conditions can increase damage from secondary pathogens (those infecting tissue in poor physiological condition), while reducing damage from primary pathogens (those infecting healthy tissue) (Jactel et al. 2012). Long-term the SSD may also be impacted by nurse-plant availability to aid persistence of some species. Piñon pine recruitment in the Southwest may benefit from high canopy cover following disturbance in areas otherwise predicted to become juniper dominated woodlands (Redmond and Barger 2013, Kane et al. 2015). Sugar and Jeffrey pine recruitment in western Nevada also benefits from nurse plants and soil water availability (Legras et al. 2010).

Post-fire recruitment may be particularly susceptible to drought conditions and lead to recruitment failures or unacceptable reductions in regeneration densities. For example, Feddema et al. (2013) and Savage et al. (2013) predict that ponderosa pine regeneration following high-severity fire will decline on dry sites when fires coincide with drought. Recruitment failures and conversion to shrub- or grasslands are common following recent high-severity fires in the Southwest (Roccaforte et al. 2012, Savage et al. 2013). The use of artificial regeneration (planting or direct seeding) offers a potential solution to some of the recruitment failures that have arisen following catastrophic fires and the loss of local seed sources (e.g., Haire and McGarigal 2008, Zhang et al. 2008, Feddema et al. 2013, Ouzts et al. 2015). However, planting must be done to match seedling genotypes and species for given locations (e.g., Blazier et al. 2004, Will et al. 2010), especially if done in the midst of a long-term or deepening drought. For example, the use of more expensive containerized nursery stock (Grossnickle 2005, Nilsson and Örlander 1995) or tree shelters for established seedlings (Taylor et al. 2009) on drought-prone sites may be necessary. If bare-root seedlings are to be planted, those with large root systems

(e.g., shoot:root ratios below 2:1) should be used (Haase and Rose 1993, Pinto et al. 2012) to help reduce drought losses.

## **Scale-dependent consequences of drought**

Understanding the dynamics of drought and forests at one scale does not mean the results can be directly scaled up or down. Gene flow and conservation efforts that span landscapes or regions can influence species composition over scales that are hard to detect at in stand or individual responses. Likewise, interactions that occur within stands mean that stand-level responses to drought will not necessarily agree with studies of individual growth and survival. Said another way, responses of individual trees at low moisture availability do not tell us how the abundances of different species will change as a result of drought. As an example, the behavior of SSD under drought conditions depends on how individual trees responses translate to population growth rates, each population being an aggregate across individuals of all size classes, competitive environments, and microhabitats (Box 2) and subject to population constraints (e.g., ability to disperse across landscapes) and attributes such as genetic diversity. Species that can tolerate xeric conditions might progressively increase in abundance within stands subjected to frequent or persistent drought and, in turn, make lower transpiration demands. Thus, the moisture for which trees compete depends in part on competition feedback (D'Amato et al. 2013).

Attempts to anticipate the effects of increased drought on local or regional species diversity highlight the challenges of extrapolating from tree-level studies. For example, Klos et al.'s (2009) analysis of FIA data specifically targeting the 1999-2001 drought in the Southeast found a weak relationship between stand diversity and drought impacts, suggesting that the partitioning of resources that occurs in diverse stands provides some buffering of drought effects. Relationships between diversity and drought impacts may vary among ecosystems (Grossirod et al. 2014). In western forests increasing drought could result in loss of some species, especially from warm and dry climates at low elevations, potentially accelerated by dieback (Kelly and Goulden 2008, Bell et al. 2014).

### ***Drought-related biogeographic and biome shifts***

Taken together, the evidence for drought effects on forest composition remains mostly indirect. Forests respond to drought not only due to changes in the SSD of trees present in the stand, but also due to immigration and local extinction (Parmesan and Yohe 2003, Chen et al. 2011). A species' geographic range changes when regeneration is successful beyond the current population frontier or when regeneration fails in a portion of the current range. In some cases drought will result in relocation of suitable habitats within the geographic region—for instance, at higher or lower elevations or adjacent to wetlands or bodies of water. Migration is more difficult to evaluate, because it occurs at and beyond range limits, where a species is rare and difficult to study. Local heterogeneity in recruitment success (Pitelka et al. 1997, Morin et al.

2007, Ibanez et al. 2007), low population density, and the potential importance of rare events over broad regions make migration difficult to detect and to quantify (Clark et al. 2003).

Recent reports that some plant species may already be migrating rapidly in response to changing climates make it important to recognize that the term *migration* is not applied consistently. For plants the term most often refers to accumulated gains and losses in the area occupied by a species, typically at a regional scale. Poleward or upslope expansions in response to a warming climate are examples. A second use of the term refers to latitude- or elevation-weighted change in abundance or performance (Feeley et al. 2011, 2013, Gottfried et al. 2012). Such weighted averages can be calculated for samples where observations are individual organisms, abundances of species on plots, or performance (e.g., growth rate) (Lenoir et al. 2008, Woodall et al. 2009). For example, growth rates of trees can serve as weights to calculate a performance-weighted mean latitude for the species. The mean latitude calculated by this approach can change from one survey to the next, regardless of whether or not the population actually moves—even if the range is static, the mean will change if individuals in different parts of the range grow faster/slower than before. Such metrics can provide valuable insight into geographic patterns, though they do not represent a change in a species' geographical distribution. Migration is also hard to assess because most studies inform us more about the centers of population ranges than about range limits. Like weighted averages, models fitted to occurrence, abundance, or demographic rates (e.g., Canham and Thomas 2010, Mok et al. 2012, Clark et al. 2014a, Zhu et al. 2014, Vanderwel et al. 2013) can be dominated by samples where the species is abundant and insensitive to margins. The smooth declines in performance near margins assumed in many models are not widely observed in demographic data (Fig. 7).

Future range shifts are difficult to anticipate, because there is only limited evidence for the combinations of variables that control current range limits. Experimental warming in northern Minnesota showed photosynthesis and growth increases near cold range limits and reductions near warm range limits in planted seedlings (Reich et al. 2015); however, seed germination and establishment were not studied. As documented for decades through forestry provenance trials, trees manifest local adaptation and home site advantage to temperature and moisture conditions (Rehfeldt *et al.*, 1999, Aitken *et al.*, 2008), so that climate change impacts would be expected throughout the range of the species, not just at leading or trailing edges of species ranges (Davis and Shaw, 2001). Potential for rapid adaptation to current changes in climate is not well understood but will depend on the interaction of spatial patterns of genetic variation and modern gene flow (Aitken et al., 2008).

Limited evidence of migration over the last century, a time when the velocity of climate change has been substantial in the northern US (Zhu et al. 2012), is not in agreement with models that suggest that suitable habitats of many species are shifting faster than are the populations themselves (McKenney et al. 2007, 2011). The combination of large projected habitat shifts with limited evidence for the rapid migration that would be needed to track these, suggests that biogeographic patterns could substantially lag behind climate change. Fundamental differences

in migration potential for eastern and western forests result because of the importance of topographic relief in the latter. In the eastern US, with substantial areas of low relief, modest changes in climate can translate to large shifts in locations of suitable habitat (Loarie et al. 2009, IPCC 2014). For the Southeast, the rate of recent climate change produces a climate velocity ranging from 0.2 km per yr in the Appalachians to > 2 km per yr on the Piedmont and coastal plain (Schleip et al. 2015). By 2100 mean isotherms could move northeastward from 400 km for a less CO<sub>2</sub>-sensitive model (PCM) with high energy-resource efficiency (B1) to 800 km for a more sensitive model (HadleyCM3) with a ‘business as usual’ scenario (A1F1) (Iverson et al. 2008a). In western forests, increasing drought could result in loss of some species at low elevations, potentially accelerated by dieback (Kelly and Goulden 2008, Bell et al. 2014), as shifts in climate may be too fast for many populations to adjust by adaptation or migration.

While there is mounting evidence that certain terrestrial and aquatic invertebrates, birds, and herbaceous plants have changed in distribution with warming (Parmesan and Yohe 2003, Chen et al. 2011), there are only a few examples of rapid contemporary natural tree migrations (e.g., Fastie 1995, Pitelka et al. 1997). The paleo record provides some examples of rapid spread in response to climate change, such as *Corylus* expansion into western Europe in the early Holocene (Huntley and Birks 1983). Late Holocene range expansion of three western conifers (*Juniperus osteosperma*, *Pinus edulis*, *P. ponderosa*) may have depended on long-distance dispersal events ranging from 25 to 100 km (Lyford et al. 2003, Gray et al. 2006, Lesser and Jackson 2012, 2013, Norris et al. 2015). Expansion of colonizing populations of *P. ponderosa* may have been slowed by Allee effects (Lesser et al. 2013). However, traditional interpretations of the paleo record suggesting that rapid tree migrations were common in the past are hard to reconcile with known dispersal rates and other life history observations (McLachlan et al. 2005). Paleo evidence can also prove quite ambiguous—for instance, the sporadic occurrence of fossils in lake sediments can mean that a few trees are nearby or that many trees are far away, making it difficult to infer when a population arrives or disappears from a region. Interpretation of Holocene tree migration remains a subject of considerable research.

With their compact moisture and temperature gradients, some of the most effective migrations could be expected in mountainous regions (Jump et al. 2009, Bell et al. 2014). Coops and Waring (2011) predict a distribution shift and reduction in range extent for lodgepole pine in the Pacific Northwest due to late summer drought. Western redcedar (*Thuja plicata*) and western hemlock (*Tsuga heterophylla*) may expand, whereas ponderosa pine, lodgepole pine, grand fir (*Abies grandis*) and noble fir (*Abies procera*) may contract (Coops et al. 2011). In the Green Mountains of Vermont, some work has indicated northern hardwoods have invaded the lower boundary of boreal forest in several locations over the last half century (Beckage et al. 2008), whereas broader patterns for this region suggest downslope migration of boreal species (Foster and D’Amato *in press*). In this location the ecotone is sharp, concentrated within 200 m of elevation. Still, even in such topography where dispersal is probably not limiting, tree upslope

shifts appear to lag climate change in the Alps (Gehrig-Fasel et al. 2007) and Andes (Feeley et al. 2011).

Although latitudinal migration in response to warming and drought stress could be occurring for some species, evidence of poleward movement of trees is even less obvious than upslope migration. Warming over the last century in the continental US has been most rapid in the upper Midwest and Northeast, due to the combination of regional climate change and low relief. Poleward migration would be identified by establishment of new recruitment out ahead of established range boundaries, especially in these areas of rapid change. This pattern is not detected in FIA data from the eastern US (Zhu et al. 2012), but could be occurring at northern limits of several species in Quebec (Boisvert-Marsh et al. 2014). Latitudinal changes might explain some changes in composition at Blackrock Forest of New York (Schuster et al. 2008) and along certain powerline corridors (Treyger and Nowak 2011). Additional evidence of poleward migration could include studies from Woodall et al. (2009), Monleon and Lintz (2015), and Delzon et al. (2013). Clearly, the rapid spread ( $> 10^3 \text{ m yr}^{-1}$ ) required to match the pace of shifting habitats is not occurring.

Changes in fire regime, land cover, and diebacks resulting from combinations of drought, disease, and human action can all contribute to expanding or contracting ranges (Cornwell et al. 2012, Franklin et al. 2013, Jiang et al. 2013), including forest conversion to shrub- and grassland (Lenihan et al. (2008, Man 2013). Increased fire frequency and/or intensity can rapidly shift composition, structure and function. The extent to which large diebacks could promote (Linares et al. 2009, Kane et al. 2011), or that forest fragmentation could reduce (Meier et al. 2012), migration capacity may vary widely. Disturbance could accelerate migration for species that would otherwise fail to invade competitive understories (Dukes et al. 2009, Weed et al. 2013). The capacity for drought-induced dieback to accelerate changes to the SSD, including interactions involving fire and insects, suggests that such change could occur at variable rates through time, with periods of slow change punctuated by episodic rapid transitions.

## **Drought and forest management**

Drought directly and indirectly affects most of the ecosystem services provided by forests, including timber (Woodall et al. 2013b), carbon storage (Gonzalez et al. 2015), recreational value (e.g., Hollenhorst et al. 1993, Aukema et al. 2011, Kovacs et al. 2011), and water yield and quality (Brown et al. 2008). Management practices modify the SSD (Box 2) through the manipulation of species, size, and density. While typically done to achieve productivity goals, management can also mitigate or exacerbate effects of drought at tree- and stand-levels through its influence on local site and climate. For example, in dry western forests density reduction and prescribed burns to promote timber yield or reduce fire risk also can reduce drought vulnerability (Grant et al. 2013, Thomas and Waring 2014).

Management of established forests for a diversity of species can reduce stand vulnerability to drought. Thinning practices may move from simple reductions in density to stand structural attributes that reduce vulnerability to drought (e.g., Guldin 2014, Thomas and Waring 2014). For example, the maintenance of uneven-aged stands may spread risks across ages/sizes of different vulnerabilities (e.g., Carter et al. 1984). Forest restoration practices may allow for greater persistence of large, old trees under drought conditions (e.g., Erickson and Waring 2014). Uncertainty in future climate can motivate a mix of drought-tolerant species and genotypes. Species composition can be altered directly through selective removal of moisture-demanding species and release of suppressed individuals of more drought tolerant species. Such replacement occurs naturally following drought-induced dieback in the pinyon-juniper ecosystems (e.g., Floyd et al. 2009). Thinning to reduce crown competition (Aussenac and Granier 1988, McDowell et al. 2006, Gyenge et al. 2011) also reduces canopy interception of precipitation, thus increasing moisture that reaches the forest floor (Stogsdili Jr et al. 1992, Aussenac 2000) where expanded root systems due to thinning can improve moisture access for individual trees (Dawson 1996). However, the reduced vulnerability of remaining trees in the short term can increase future vulnerability through changes in tree architecture and physiology. Long-term increases in leaf to sapwood area ratios in stands thinned to low densities can increase individual tree water demand (McDowell et al. 2006, Kolb et al. 2007) despite possibly reduced water use by the stand as a whole. Indirect effects can include promoting regeneration (Covington et al. 1997, Moore et al. 1999), also vulnerability to drought (Aussenac 2000) and competition from non-tree species that can increase beneath open canopies (Nilsen et al. 2001). In dry forests of the western US, such negative effects may be offset by the fire hazard reduction that comes with most management options currently being implemented (e.g., Martinson and Omi 2013, Waltz et al. 2014).

Successful regeneration during drought depends on microsite conditions, including competition from non-preferred species. However, current practices and guidelines for seed transfer may need to be re-considered given the potential for locally maladapted genotypes, as well as the possibility of planting more heat- and drought-tolerant genotypes (Aitken et al., 2008, Joyce and Rehfeldt, 2013, Montwé et al., 2015). After all, decades of horticultural practice have clearly shown that growth and reproduction of many species well outside their native ranges is possible, suggesting “assisted migration” is a viable diversity conservation option (e.g., Schwartz et al 2012). Management for drought through active involvement in the regeneration process can also prove costly (Nyland 2007). For species that are especially vulnerable as seedlings (Cavender-Bares and Bazzaz 2000) steps can be taken to maximize belowground development prior to and immediately after planting (e.g., Burdett 1990) or to shelter future crop trees (e.g., Aussenac 2000). Drought may increase reliance on artificial regeneration (i.e., plantings), protection of planted seedlings, and/or seedbed amelioration, such as the manipulation of harvest residues to provide a mulching effect (Roberts et al. 2005, Trottier-Picard et al. 2014). Artificial regeneration may become especially important for conifers that fail to regenerate or are out-competed by sprouting hardwood species (Haire and McGarigal 2008, Zhang et al. 2008,



Feddema et al. 2013, Ouzts et al. 2015). Because recruitment depends on local site conditions, knowledge of how different species and genotypes respond on different sites (Blazier et al. 2004, Will et al. 2010, Erickson et al. 2012) should guide management rather than regional climate projections.

## **Knowledge gaps and future directions**

### ***Challenges/limitations of available data***

Efforts to anticipate future forests rely heavily on observational data, which are often unavailable or difficult to extrapolate. For example, droughts are expected to increase in the Northeast (Melillo et al. 2014), a region that has not experienced severe drought since the 1960s, before the regular collection of forest inventory data. Furthermore, many of the variables that affect forests are changing simultaneously, making it difficult to attribute observed changes to rising CO<sub>2</sub>, N deposition, invasive species, or increasing average age of forest stands (McMahon et al. 2010). The interactions that control stand responses to drought remain poorly understood. Our best understanding comes from the study of individual trees (e.g., Fig. 1). As individuals respond, they interact with one another and with natural enemies. Some interactions occur *within* individuals, such as allocation of carbon resources, with the result that growth, maturation rates, fecundity, and survival can react to drought in different ways (Fig. 3). Other interactions occur *between* individuals, such as competition in crowded stands. Soil moisture depends not only on climate, but also on redistribution by local drainage, and uptake by competing trees (Fig. 4) (e.g., Loik et al. 2004). Interactions with fungal pathogens and herbivores can impact host individuals differently, depending on species, size, resource availability, and host resistance.

In addition to evidence of climate-competition interactions at the scale of individual trees (Cescatti and Piutti 1998, Martin-Benito et al. 2011), evidence also can be found in stands (D'Amato et al. 2013, Thomas and Waring 2015) and across plot networks (Clark et al. 2011, 2014b). Drought effects on SSDs depend on all of these interactions (Box 2). For example, rising CO<sub>2</sub> interacts with SSD, because increase in water-use efficiency of individual leaves or drought-tolerance of whole plants varies widely between species and is expected to depend on the light environment and soil moisture (Battipaglia et al. 2013). Likewise, spatial variation of forest response to moisture and temperature gradients can be confounded by land use, management history, soils, and complex hydrology. For example, private landowners in the Pacific Northwest manage some productive lands for timber production, whereas state and federal agencies mostly manage low productivity and high elevation forests for diverse objectives (Ohmann and Spies 1998). In the Southeast Piedmont, moisture gradients are confounded by land use and stand age. Typical stands of intermediate moisture status established on former cultivated lands a century ago, whereas xeric sites were grazed, and wet bottomlands were not cultivated and thus tend to support older trees (Oosting 1942, Quarterman and Keever 1962). As a consequence, observational data may not yield unambiguous relationships between forest structure and moisture.

Data coverage is also uneven. For example, paleo studies of forest response to past climate come from either tree-ring records or fossil evidence from lake/bog sediments, which are dispersed unevenly in humid regions. Tree-ring data come primarily from mature trees expected to be most sensitive to climate (Fritts 1976) and might respond to climate differently from seedlings and saplings. Moreover, open, low-density stand conditions are often preferentially sampled for tree-ring studies to reduce the growth signal resulting from density-dependent interactions.

Data sets that span sufficient temporal variation in climate are limited. Demographic responses to climate change can be estimated from plot data when there are three or more consistent censuses. Thus far, FIA data provide two consistent censuses for most of the eastern US, but only one census for most of the West. Two censuses provide estimates of mortality rates from numbers of trees that die during the interval (Dietze and Moorcroft 2011, Lines et al. 2010), of recruitment rates from individuals appearing in a census not present previously (Zhu et al. 2014), and of growth rates from changes in size (Vanderwel et al. 2013). However, the geographic relationships between demography and climate may not represent how demography responds to climate change. Understanding forest change through inventory analysis is further complicated when different designs were implemented between the first and second censuses (Goeking in press). Data sets containing long intervals between censuses can be hard to interpret because they integrate many years of climate variation. Many forest plots are re-sampled at intervals of 4 to 10 or more years. Intervals this long can include both exceptionally warm, cold, dry, and wet years (e.g., Williams et al. 2013).

Experiments address some of the limitations of observational data by manipulating the environment in a controlled fashion. However, relatively few experiments are available at a scale that provides general insight for climate changes that affect diverse habitats. Because individual trees can be manipulated more readily than forest stands, there is more evidence of drought effects on trees than on stand-level size-species structure. Experiments sufficiently large and long-term to determine effects on stand composition and structure are costly; not surprisingly, there are still only a few rainfall exclusion and redistribution experiments on mature temperate forests (e.g., McDowell et al. 2013, Hanson and Weltzin 2000), and few experiments at any scale include extreme drought and heat events (Jentsch et al. 2007) or tree-killing levels of drought stress (cf. Allen et al. 2015).

### ***Contributions from models of forest change***

Models for forest response to climate change rely heavily on parameters fitted independently to recruitment, growth, and mortality, and primarily from observations on individual trees, rather than stands. Furthermore, interactions complicate prediction efforts (Tinner et al. 2013). To date, much of the research on climate impacts on stand dynamics relies on simulations of several types, three of which are summarized here:

*Species distribution models* (SDMs) are used to map potential future species habitats under climate scenarios (e.g., Guisan and Thuiller 2005, Franklin 2010, Matthews et al. 2011, Prasad et

al. 2013). Species distributions are calibrated to climate and other environmental variables. The fitted models are then used with climate scenarios generated by GCMs to identify regions of future suitable habitat.

*Dynamic global vegetation models* (DGVMs) are dynamic and non-spatial (Daly et al. 2000, Sitch et al. 2003, Jiang et al. 2013). Species are aggregated as functional types, such as coniferous, deciduous, and mixed forests, savannas and woodlands, or grasslands and shrublands (Bachelet et al. 2003). Some incorporate fire, atmospheric CO<sub>2</sub> (Lenihan et al. 2008, King et al. 2013), establishment mechanisms (Song and Zeng 2014), and patch age structure (Medvigy and Moorcroft 2012). DGVMs are used to predict change in functional types.

*Forest landscape models* (FLMs) simulate forest demography on landscapes that may include drought, fire, land use, and pathogens. Some FLMs explicitly focus on climate change impacts (Scheller and Mladenoff 2008, Loehman et al. 2011), including migration (Lischke et al. 2006, Scheller and Mladenoff 2008, Gustafson and Sturtevant 2013, Nabel et al. 2013, Snell 2014). FLMs are used to predict dynamics of forest stands.

Models of climate effects (including drought) continue to improve but are subject to caveats. First is the uncertainty in climate projections—as an example, three GCMs project climates in 2100 differ by up to 4°C for mean annual temperature and 60% for precipitation over North America (McKenney et al. 2011). This difference suggests mean latitudes for species' habitats could move northeastward from 400 km for a less CO<sub>2</sub>-sensitive model (PCM) with high energy-resource efficiency (B1) to 800 km for a more sensitive model (HadleyCM3) with a 'business as usual' scenario (A1F1) (Iverson et al. 2008b). Second is the uncertainty from heterogeneity not captured in GCM output, from redistribution of precipitation within local drainages—wet and dry sites occupy the same grid cell for regional climate prediction, and from variation in temperature with local topography and vegetation cover. Models of future forest response to future climate begin with this uncertainty in regional and local climate.

Third, all calibration-prediction and simulation approaches incorporate parameters relating drought to recruitment, growth, and survival from separate studies and typically from individual trees, while the interactions that determine drought response, depend on the SSD—the interdependence between individuals within the SSD requires that they be considered together (Box 2). Regional or population-level differences in climate relationships are rarely incorporated into these models (e.g., Sork et al., 2010, Joyce and Rehfeldt 2013, Rehfeldt et al., 2014).

Related to the third point, there remains a need to develop better model representations of climate-mediated mortality (McDowell et al. 2011, Allen et al. 2015) and species interactions (Ibanez et al. 2006). Whether or not populations can move to regions of future suitable climate depends on migration, which is poorly understood. For example, potential distributions predicted from SDMs are sometimes bracketed by two extremes – no migration (species lose but do not gain habitat) and unlimited migration (species occupy all suitable habitat) (Thuiller et al.

2005, Iverson et al. 2008c, Meier et al. 2012). A better understanding of how droughts affect seed production, seed banks, and seedling establishment near range limits, particularly their role in local extinctions and re-colonization (Zimmermann et al. 2009, Jackson et al. 2009) might improve characterization of extremes. The limited studies show large variation in fecundity (Clark et al. 2004, Koenig and Knops 2013) and recruitment (Ibanez et al. 2007). Models that incorporate such estimates predict migration rates that are highly uncertain (Clark et al. 2003). Land cover adds an additional layer of variability, both limiting habitat but often providing recruitment opportunities following disturbance (Iverson et al. 2004, Prasad et al. 2013, Clark et al. 2003).

## Conclusions

There is broad consensus from modeling studies, increasingly supported by observation, that combinations of heat and moisture limitation, and their corresponding indirect effects, will change the health, dynamics, abundance, and distribution of tree species. These changes may accelerate in coming decades. In the eastern US, drought effects are still primarily observed in responses of individual trees. How species differences, well studied at the individual scale, translate to future stand structure and composition is uncertain. Observations of stand-level responses to evaluate how climate changes interact with changing effects of competitors, mutualists, and natural enemies, which are also responding to climate change, are limited. In the western US, stand-level forest transformations are in progress now, already highlighting interactions among warming temperatures, drought, insect attacks, and fire. A proactive management strategy for anticipating change can include promoting drought-tolerant species, managed in lower density stands, and potentially drawing on species or genotypes outside their current geographic ranges.

Despite many important insights from observational evidence, the foregoing knowledge gaps and future climate change highlight the challenge posed by connecting abundant research on individual tree responses to the scale where predictions are needed—the forest stand. Research priorities should include more attention to effects of drought beyond the individual, for example, to focus on the combined size-species interactions that control diversity and productivity of stands. After the uncertainty in climate itself, the greatest obstacle to understanding impacts of future drought is the limited understanding of drought consequences at stand-to-landscape scales. Models will continue to play an important role, one that depends on improved understanding of stand-level responses and the acquisition of suitable long-term data for detection, parameterization, calibration, and validation. This challenge is related to the need for models that accommodate environmental change and forest response as a coherent joint distribution of species and sizes (the SSD), that responds to drought with adequate feedbacks and interactions. The problem persists despite proliferation of bigger and more complex models, faster processing, and increased computer memory. Without the empirical basis for translating fine-scale to aggregate behavior – in the form of allocation constraints, species interactions, and feedbacks –

complex models can provide only limited guidance. These constraints are needed in models when they are fitted to field and experimental data.

Much could be gained from increased efforts focused on the connections from individual to stand, both empirical and modeling. For example, how does decline in individual tree health translate to population structure and abundance of a species, when individuals of all species are responding to climate, often in similar ways? Again, consider the well-known relationship in traditional forestry that the highest mortality rates occur in the most productive stands. Climate changes that place individuals at risk can have unpredictable effects on stands as the individuals within stands respond. At the individual scale, long-term data with regional coverage are needed to infer demographic processes under a range of climates and to detect early signs of change (Breshears et al. 2009). However, predicting changes in stands also requires stand-level inference. The observable physiological responses to temperature and moisture stress must be linked to demographic potential of individuals and to stand attributes, such as size-species distributions. Predicting effects of novel climate on biogeographic patterns would likewise benefit from better understanding of how current biogeography emerges from tree responses to climate. Additional insights might be gained from natural gradients in regions expected to differ in sensitivity to moisture and temperature, with emphasis on connections from individuals to stands. Finally, opportunistic or designed experiments to better understand geographic variation of drought effects still are needed.

## References

- Abrams, M. D. 2003. Where has all the white oak gone? *Bioscience* 53: 927-939.
- Abrams, M. D. 1990. Adaptations and responses to drought in *Quercus* species of North America. *Tree Physiology* 7: 227–238.
- Abrams, M. D. 1994. Fire and the development of oak forests. *Bioscience* 42:346-353.
- Abrams, M. D. 1998. The Red Maple Paradox. *BioScience* 48: 355-364.
- Abrams, M. D. and M. E. Kubiske. 1990. Leaf structural characteristics of 31 hardwood and conifer tree species in central Wisconsin: influence of light regime and shade tolerance rank. *Forest Ecology and Management* 31: 245–253.
- Adams, H. D., M. Guardiola-Claramonte, G. A. Barron-Gafford, J. C. Villegas, D. D. Breshears, C. B. Zou, P. A. Troch, and T. E. Huxman. 2009. Temperature sensitivity of drought-induced tree mortality portends increased regional die-off under global-change-type drought. *Proceedings of the National Academy of Sciences of the United States of America* 106: 7063-7066.
- Aitken SN, Yeaman S, Holliday JA, Wang T, Curtis-McLane S (2008) Adaptation, migration or extirpation: climate change outcomes for tree populations: Climate change outcomes for tree populations. *Evolutionary Applications*, 1, 95–111.
- Allen, C. D. 2007. Interactions across spatial scales among forest dieback, fire, and erosion in northern New Mexico landscapes. *Ecosystems* 10:797-808.
- Allen, C. D., A. K. Macalady, H. Chenchouni, D. Bachelet, N. McDowell, M. Vennetier, T. Kitzberger, A. Rigling, D. D. Breshears, E. H. T. Hogg, P. Gonzalez, R. Fensham, Z. Zhang,

- J. Castro, N. Demidova, J.-H. Lim, G. Allard, S. W. Running, A. Semerci, and N. Cobb. 2010. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management* 259: 660-684.
- Allen, C. D., and D. D. Breshears. 1998. Drought-induced shift of a forest-woodland ecotone: Rapid landscape response to climate variation. *Proceedings of the National Academy of Sciences, U.S.A.* 95:14839-14842.
- Allen, C.D., D.D. Breshears, and N.G. McDowell. 2015. On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* 6(8):129. doi.org/10.1890/ES15-00203.1.
- Anderegg, W. R. L., L. Plavcová, L. D. L. Anderegg, U. G. Hacke, J. A. Berry, and C. B. Fields. 2013a. Drought's legacy: multiyear hydraulic deterioration underlies widespread aspen forest die-off and portends increased future risk. *Global Change Biology* 19: 118-1196.
- Anderegg, W. R. L., J. M. Kane, and L. D. L. Anderegg. 2013b. Consequences of widespread tree mortality triggered by drought and temperature stress. *Nature Climate Change* 3: 30-36.
- Assmann, E. 1970. *The Principles of Forest Yield Study*. Pergamon.
- Anderegg, W. R., Flint, A., Huang, C. Y., Flint, L., Berry, J. A., Davis, F. W., ... & Field, C. B. (2015). Tree mortality predicted from drought-induced vascular damage. *Nature Geoscience*.
- Aussenac, G. 2000. Interactions between forest stands and microclimate: Ecophysiological aspects and consequences for silviculture. *Annals of Forest Science* 57: 287-301.
- Bachelet, D., R. P. Neilson, T. Hickler, R. J. Drapek, J. M. Lenihan, M. T. Sykes, B. Smith, S. Sitch, and K. Thonicke. 2003. Simulating past and future dynamics of natural ecosystems in the United States. *Global Biogeochemical Cycles* 17(2): 14-1.
- Barber, V. A., B. P. Juday, and B. P. Finney. 2000. Reduced growth of Alaskan white spruce in the twentieth century from temperature-induced drought stress. *Nature* 405: 668-673.
- Barton, A.M. 2002. Intense wildfire in southeastern Arizona: transformation of a Madrean pine-oak forest to oak woodland. *Forest Ecology and Management* 165: 205-212.
- Battipaglia, G., M. Saurer, P. Cherubini, C. Calfapietra, H. R. McCarthy, R. J. Norby, and M. F. Cotrufo. 2013. Elevated CO<sub>2</sub> increases tree-level intrinsic water use efficiency: insights from carbon and oxygen isotope analyses in tree rings across three forest FACE sites. *New Phytologist* 197: 544-554.
- Beckage B., B. Osborne, C. Pucko, D. G. Gavin, T. Siccama, and T. Perkins. 2008. An upward shift of a forest ecotone during 40 years of warming in the Green Mountains of Vermont, USA. *Proceedings of the National Academy of Sciences* 105(11): 4197-4202.
- Belden, A. C. and S. G. Pallardy. 2009. Successional trends and apparent *Acer saccharum* regeneration failure in an oak-hickory forest in central Missouri, USA. *Plant Ecology* 204: 305-322.
- Bell, D. M., J. B. Bradford, and W. K. Lauenroth. 2014. Early indicators of change: divergent climate envelopes between tree life stages imply range shifts in the western United States. *Global Ecology and Biogeography* 23: 168-180.
- Bentz, B. J., J. Régnière, C. J. Fettig, E. M. Hansen, J. L. Hayes, J. A. Hicke, R. G. Kelsey, J. F. Negrón, and S. J. Seybold. 2010. Climate change and bark beetles of the western United States and Canada: direct and indirect effects. *BioScience* 60: 602-613.
- Berdanier, A. and J.S. Clark. 2015. Multi-year drought-induced morbidity preceding tree death in Southeastern US forests. *Ecological Applications*, in press.
- Blazier, M.A., T.C. Hennessey, T.B. Lynch, R.F. Wittwer, and M.E. Payton. 2004. Productivity, crown architecture, and gas exchange of North Carolina and

- Oklahoma/Arkansas loblolly pine families growing on a droughty site in southeastern Oklahoma. *Forest Ecology and Management* 194: 83-94.
- Boisvert-Marsh, L., C. Périé, and S. de Blois. 2014. Shifting with climate? Evidence for recent changes in tree species distribution at high latitudes. *Ecosphere* 5:art83.
- Booth, R.K., M. Notaro, S.T. Jackson, & J.E. Kutzbach. 2006. Widespread drought episodes in the western Great Lakes region during the past 2000 years: geographic extent and potential mechanisms. *Earth and Planetary Science Letters* 242:415-427.
- Booth, R.K., S.T. Jackson, V.A. Sousa, M.E. Sullivan, T.A. Minckley & M. Clifford. 2012. Multidecadal drought and amplified moisture variability drove rapid forest community change in a humid region. *Ecology* 93:219-226.
- Bréda N., R. Huc, A. Granier, E. Dreyer. 2006. Temperate forest trees and stands under severe drought: a review of ecophysiological responses, adaption processes and long-term consequences. *Annals of Forest Science* 63: 625–644
- Breshears, D. D., N. S. Cobb, P. M. Rich, K. P. Price, C. D. Allen, R. G. Balice, W. H. Romme, J. H. Kastens, M. L. Floyd, J. Belnap, J. J. Anderson, O. B. Myers, and C. W. Meyer. 2005. Regional vegetation die-off in response to global-change-type drought. *Proceedings of the National Academy of Sciences, USA*. 102(42): 15144-48.
- Breshears, D. D., Myers, O. B., Meyer, C. W., Barnes, F. J., Zou, C. B., Allen, C. D., McDowell, N. G., and Pockman, W. T. 2009. Tree die-off in response to global change-type drought: mortality insights from a decade of plant water potential measurements. *Frontiers in Ecology and the Environment* 7: 185–189.
- Bright, R.M., K. Zhao, R.B. Jackson, and F. Cherubini. 2015. Quantifying surface albedo and other direct biogeophysical climate forcings of forestry activities. *Global Change Biology* 21:3246-3266.
- Brose, P. H., D. C. Dey, R. J. Phillips, and T. A. Waldrop. 2013. A Meta-Analysis of the Fire-Oak Hypothesis: Does Prescribed Burning Promote Oak Reproduction in Eastern North America? *Forest Science* 59: 322-334.
- Brown, P. M., and R. Wu. 2005. Climate and disturbance forcings of episodic tree recruitment in a southwestern ponderosa pine landscape. *Ecology* 86: 3030-3038.
- Brown, T. C., M. T. Hobbins, and J. A. Ramirez. 2008. Spatial Distribution of Water Supply in the Conterminous United States. *Journal of the American Water Resources Association* 44: 1474-1487.
- Brzostek, E. R., D. Dragoni, H. P. Schmid, A. F. Rahman, D. Sims, C. A. Wayson, D. J. Johnson, and R. P. Phillips. 2014. Chronic water stress reduces tree growth and the carbon sink of deciduous hardwood forests. *Global Change Biology*.
- Burdett, A.N. 1990. Physiological processes in plantation establishment and the development of specifications for forest planting stock. *Canadian Journal of Forest Research* 20: 415-427.
- Cabral, J. S. and F. M. Schurr. 2010. Estimating demographic models for the range dynamics of plant species. *Global Ecology and Biogeography* 19: 85-97.
- Canham, C. D. and R. Q. Thomas. 2010. Frequency, not relative abundance, of temperate tree species varies along climate gradients in eastern North America. *Ecology* 91: 3433-3440.
- Carter, G.A., J.H. Miller, D.E. Davis, and R.M. Patterson. 1984. Effectiveness of vegetative competition on the moisture and nutrient status of loblolly pine. *Canadian Journal of Forest Research* 14: 1-9.

- Cavender-Bares, J. and F. A. Bazzaz. 2000. Changes in drought response strategies with ontogeny in *Quercus rubra*: implications for scaling from seedlings to mature trees. *Oecologia* 124: 8-18.
- Cescatti, A., and E. Piutti. 1998. Silvicultural alternatives, competition regime and sensitivity to climate in a European beech forest. *Forest Ecology and Management* 102: 213–223.
- Chen, I. C., J. K. Hill, R. Ohlemüller, D. B. Roy, and C. D. Thomas. 2011. Rapid range shifts of species associated with high levels of climate warming. *Science* 333: 1024-1026.
- Chen, P. Y., C. Welsh, and A. Hamann. 2010. Geographic variation in growth response of Douglas-fir to interannual climate variability and projected climate change. *Global Change Biology* 16:3374-3385.
- Chmura, D. J., P. D. Anderson, G. T. Howe, C. A. Harrington, J. E. Halofsky, D. L. Peterson, D. C. Shaw, and J. Brad St Clair. 2011. Forest responses to climate change in the northwestern United States: ecophysiological foundations for adaptive management. *Forest Ecology and Management* 261: 1121-1142.
- Clark, J., and P. Royall. 1996. Local and regional sediment charcoal evidence for fire regimes in presettlement north-eastern North America. *Journal of Ecology* 84: 35–382.
- Clark, J.S. 1989. Water balance and fire occurrence during the last 160 years in northwestern Minnesota. *Journal of Ecology* 77: 989-1004.
- Clark, J.S. 1990. Integration of ecological levels: individual plant growth, population mortality, and ecosystem dynamics. *Journal of Ecology* 78: 275-299.
- Clark, J. S., A. E. Gelfand, C. W. Woodall, and K. Zhu. 2014a. More than the sum of the parts: forest climate response from joint species distribution models, *Ecological Applications*, in press.
- Clark, J.S., B. Beckage, P. Camill, B. Cleveland, J. Hille Ris Lambers, J. Lichter, J. MacLachlan, J. Mohan, and P. Wyckoff. 1999. Interpreting recruitment limitation in forests. *American Journal of Botany* 86: 1-16.
- Clark, J. S., D. M Bell, M. Kwit, A. Powell, And K. Zhu. 2013. Dynamic inverse prediction and sensitivity analysis with high-dimensional responses: application to climate-change vulnerability of biodiversity. *Journal of Biological, Environmental, and Agricultural Statistics* 18: 376-404.
- Clark, J. S., D. M. Bell, M. C. Kwit, and K. Zhu. 2014b. Competition-interaction landscapes for the joint response of forests to climate change. *Global Change Biology*, in press.
- Clark, J.S., D.M. Bell, M.H. Hersh, and L. Nichols. 2011. Climate change vulnerability of forest biodiversity: climate and resource tracking of demographic rates. *Global Change Biology*, 17, 1834–1849.
- Clark, J. S., M. Lewis, and L. Horvath. 2001. Invasion by extremes: variation in dispersal and reproduction retards population spread. *American Naturalist* 157: 537-554.
- Clark, J. S., M. Lewis, J. S. McLachlan, and J. Hille Ris Lambers. 2003. Estimating population spread: what can we forecast and how well? *Ecology* 84:1979-1988.
- Clark, J. S., S. LaDeau, and I. Ibanez. 2004. Fecundity of trees and the colonization-competition hypothesis, *Ecological Monographs* 74: 415-442.
- Cook, E. R., D. M. Meko, D. W. Stahle, and M. K. Cleaveland. 1999. Drought reconstructions for the continental United States. *Journal of Climate* 12:1145–1162.
- Cook, E. R., J. S. Glitzenstein, P. J. Krusic, and P. A. Harcombe. 2001. Identifying functions groups of trees in west Gulf Coast forests (USA): a tree-ring approach. *Ecological applications* 11:883–903.



- Cook, E., R. Seager, R. R. J. Heim, R. S. Vose, C. Herweijer, and C. A. Woodhouse. 2010. Megadroughts in North America: placing IPCC projections of hydroclimatic change in a long-term palaeoclimate context. *Journal of Quaternary Science* 25:48–61.
- Cooper-Ellis, S., D. R. Foster, G. Carlton and A. Lezberg. 1999. Forest Response to Catastrophic Wind: Results from an Experimental Hurricane. *Ecology* 80: 2683-2696
- Coops, N. C. and R. H. Waring. 2011. A process-based approach to estimate lodgepole pine (*Pinus contorta* Dougl.) distribution in the Pacific Northwest under climate change. *Climatic Change* 105: 313-328.
- Coops, N. C., R. H. Waring, C. Beier, R. Roy-Jauvin, and T. Wang. 2011. Modeling the occurrence of 15 coniferous tree species throughout the Pacific Northwest of North America using a hybrid approach of a generic process-based growth model and decision tree analysis. *Applied Vegetation Science* 14: 402-414.
- Cornwell, W. K., S. A. Stuart, A. Ramirez, C. R. Dolanc, J. H. Thorne, and D. A. Ackerly. 2012. Climate Change Impacts on California Vegetation: Physiology, Life History, and Ecosystem Change. California Energy Commission, Sacramento, California.
- Covington, W. W., and M. M. Moore. 1994. Southwestern ponderosa pine forest structure: Changes since Euro-American settlement. *Journal of Forestry* 92: 39-47.
- D'Amato, A. W., J. B. Bradford, S. Fraver, and B. J. Palik. 2013. Effects of thinning on drought vulnerability and climate response in north temperate forest ecosystems. *Ecological Applications* 23: 1735-1742.
- Dai, A. 2012. Increasing drought under global warming in observations and models. *Nature Climate Change* 3(1): 52-58.
- Daly, C., D. Bachelet, J. M. Lenihan, R. P. Neilson, W. Parton, and D. Ojima. 2000. Dynamic simulation of tree-grass interactions for global change studies. *Ecological Applications* 10: 449-469.
- Davis M, Shaw R (2001) Range shifts and adaptive responses to Quaternary climate change. *Science*, **292**, 673–679.
- Dawson, T. E. 1996. Determining water use by trees and forests from isotopic, energy balance and transpiration analyses: the roles of tree size and hydraulic lift. *Tree Physiology* 16: 263-272.
- Delzon S, Urli M, Samalens J-C, Lamy J-B, Lischke H, Sin F, Zimmermann NE & Porté A, **2013**. Field evidence of colonisation by Holm oak, at the northern margin of its distribution range, during the Anthropocene period. *PLoS ONE* 8(11): e80443.
- Desprez-Loustau, M. L., B. Marçais, L. M. Nageleisen, D. Piou, and A. Vannini. 2006. Interactive effects of drought and pathogens in forest trees. *Annals of Forest Science* 63: 597-612.
- Dietze, M., and J. S. Clark. 2008. Rethinking gap dynamics: the impact of damaged trees and sprouts. *Ecological Monographs* 78: 331-347.
- Dietze, M. C., and P. R. Moorcroft. 2011. Tree mortality in the eastern and central United States: patterns and drivers. *Global Change Biology* 17: 3312–3326.
- Dobrowski SZ, Swanson AK, Abatzoglou JT, Holden ZA, Safford HD, Schwartz MK & Gavin DG, 2015. Forest structure and species traits mediate projected recruitment declines in western US tree species. *Global Ecology and Biogeography*, in press. doi: 10.1111/geb.12302.
- Dolanc, C. R., J. H. Thorne, and H. D. Safford. 2013. Widespread shifts in the demographic structure of subalpine forests in the Sierra Nevada, California, 1934 to 2007. *Global Ecology and Biogeography* 22: 264–276.

- Dormana, M., T. Svoray, A. Perevolotsky, D. Sarris. 2013. Forest performance during two consecutive drought periods: Diverging long-term trends and short-term responses along a climatic gradient. *Forest Ecology and Management* 310: 1-9.
- Dukes, J. S., J. Pontius, D. Orwig, J. R. Garnas, V. L. Rodgers, N. Brazee, B. Cooke, K. A. Theoharides, E. E. Stange, R. Harrington, J. Ehrenfeld, J. Gurevitch, M. Lerda, K. Stinson, R. Wick, and M. Ayres. 2009. Responses of insect pests, pathogens, and invasive plant species to climate change in the forests of northeastern North America: What can we predict? *Canadian Journal of Forest Research* 39: 231-248.
- Elliott, K. J. and W. T. Swank. 1994. Impact of drought on tree mortality and growth in a mixed hardwood forest. *Journal of Vegetation Science* 5: 229–236.
- Engber, E. A., and J. M. Varner. 2012. Patterns of flammability of the California oaks: the role of leaf traits. *Canadian Journal of Forest Research* 42: 1965–1975.
- Erickson, V., C. Aubry, P. Berrang, T. Blush, A. Bower, B. Crane, T. DeSpain, D. Gwaze, J. Hamlin, M. Horning, R. Johnson, M. Mahalovich, M. Maldonado, R. Sniezko, and B. St. Clair. 2012. *Genetic Resource Management and Climate Change: Genetic Options for Adapting National Forests to Climate Change*. Washington, DC: USDA Forest Service, Forest Management.
- Erickson, C. C. and K. M. Waring. 2014. Old ponderosa pine growth and mortality responses to restoration treatments at Mt. Trumbull, AZ. *Applied Vegetation Science*. 17(1):97-108.
- Esper, J., R. Niederer, P. Bebi, and D. Frank. 2008. Climate signal age effects—Evidence from young and old trees in the Swiss Engadin. *Forest Ecology and Management* 255: 3783–3789.
- Fastie, C. L., 1995. Causes and ecosystem consequences of multiple pathways of primary succession at Glacier Bay, Alaska. *Ecology* 76: 1899 – 1916.
- Feddema, J. J., J. N. Mast, and M. Savage. 2013. Modeling high-severity fire, drought and climate change impacts on ponderosa pine regeneration. *Ecological Modelling* 253: 56-69.
- Feeley K. J., M. R. Silman, M. Bush, W. Farfan, K. Garcia Cabrera, Y. Malhi, P. Meir, N. Salinas Revilla, M. N. Raurau Quisiyupanqui, and S. Saatchi. 2011. Upslope migration of Andean trees. *Journal of Biogeography* 38: 783-791.
- Feeley K. J., J. Hurtado, S. Saatchi, M. Silman, and D. Clark. 2013. Compositional shifts in Costa Rican forests due to climate-driven species migrations. *Global Change Biology* 19: 3472-2480.
- Fei S., N. Kong, K. C. Steiner, W. K. Moser, and E. B. Steiner. 2011. Change in oak abundance in the eastern United States from 1980 to 2008. *Forest Ecology and Management* 262: 1370–1377
- Fellows, A. W. and M. L. Goulden. 2012. Rapid vegetation redistribution in Southern California
- Floyd, M.L., M. Clifford, N.S. Cobb, D. Hanna, R. Delph, P. Ford, and D. Turner. 2009. Relationship of stand characteristics to drought-induced mortality in three Southwestern piñon-juniper woodlands. *Ecological Applications* 19(5): 1223-1230.
- Foster, J. R., and A. W. D'Amato. In press. Montane forest ecotones moved downslope in northeastern US in spite of warming between 1984 and 2011. *Global Change Biology*.
- Franklin, J. 2010. Moving beyond static species distribution models in support of conservation biogeography. *Diversity and Distributions* 16: 321-330.
- Franklin, J., F. W. Davis, M. Ikagami, A. D. Syphard, A. Flint, L. Flint, and L. Hannah. 2013. Modeling plant species distributions under future climates: how fine-scale do climate models need to be? *Global Change Biology* 19: 473-483.

- Frelich, L. E., and P. B. Reich. 2010. Will environmental changes reinforce the impact of global warming on the prairie-forest border of central North America? *Frontiers in Ecology and the Environment* 8: 371–378.
- Fritts, H. C. 1976. *Tree Rings and Climate*. Page 567. Academic Press, New York.
- Fule, P. Z., J. E. Korb, and R. Wu. 2009. Changes in forest structure of a mixed conifer forest, southwestern Colorado, USA. *Forest Ecology and Management* 258 (7): 1200-1210.
- Ganey, J. L., and S. C. Vojta. 2011. Tree mortality in drought-stressed mixed-conifer and ponderosa pine forests, Arizona, USA. *Forest Ecology and Management* 261(1): 162–168. .
- Gehrig-Fasel J., A. Guisan, and N. E. Zimmermann. 2007. Treeline shifts in the Swiss Alps: Climate Change or Land Abandonment? *Journal of Vegetation Science* 18: 571-582.
- Gitlin, A. R., C. M. Sthultz, M. A. Bowker, S. Stumpf, K. L. Paxton, K. Kennedy, A. Munoz, J. K. Bailey, and T. G. Whitham. 2006. Mortality gradients within and among dominant plant populations as barometers of ecosystem change during extreme drought. *Conservation Biology* 20: 1477-1486.
- Goeking, S.A. In press. Disentangling forest change from forest inventory change: a case study from the US Interior West. *Journal of Forestry*.
- Goforth, B. R. and R. A. Minnich. 2008. Densification, stand-replacement wildfire, and extirpation of mixed conifer forest in Cuyamaca Rancho State Park, southern California. *Forest Ecology and Management* 256: 36-45.
- Grant, G.E., C.L. Tague, and C.D. Allen. 2013. Watering the forest for the trees: an emerging priority for managing water in forest landscapes. *Frontiers in Ecology and the Environment* 11:314-321.
- Gonzalez, J., J. P. , B. M. Collins, T. Robards, and D. S. Saah. 2015. Aboveground live carbon stock changes of California wildland ecosystems, 2001-2010. *Forest Ecology and Management*, 348, 68-77.
- Gottfried, M., H. Pauli, A. Futschik, M. Akhalkatsi, P. Barancok, J. L. B. Alonso, G. Coldea, J. Dick, B. Erschbamer, M. R. F. Calzado, G. Kazakis, J. Krajci, P. Larsson, M. Mallaun, O. Michelsen, D. Moiseev, P. Moiseev, U. Molau, A. Merzouki, L. Nagy, G. Nakhutsrishvili, B. Pederson, G. Pelino, M. Puscas, G. Rossi, A. Stanisci, J. P. Theurillat, M. Tomaselli, L. Villar, P. Vittoz, I. Vogiatzakis, and G. Grabherr. 2012. Continent-wide response of mountain vegetation to climate change. *Nature Climate Change* 2: 111–115.
- Graumlich, L. 1993. Response of tree growth to climatic variation in the mixed conifer and deciduous forests of the upper Great Lakes region. *Canadian Journal of Forest Research* 23: 133–143.
- Gray, S.T., J.L. Betancourt, S.T. Jackson, & R.G. Eddy. 2006. Role of multidecadal climatic variability in a range extension of pinyon pine. *Ecology* 87:1124-1130.
- Grossiord C., A. Granier, S. Ratcliffe, O. Bouriaud, H. Bruelheide, E. Checko, D. I. Forrester, S. M. Dawud, L. Finer, M. Pollastrini, M. Scherer-Lorenzen, F. Valladares, D. Bonal, and A. Gessler. 2014. Tree diversity does not always improve resistance of forest ecosystems to drought. *Proceedings of the National Academy of Sciences* 111: 14812-14815.
- Grossnickle, S. 2005. Importance of root growth in overcoming planting stress. *New Forests* 30: 273-294.
- Grubb, P. J. 1977. The maintenance of species-richness in plant communities: the importance of the regeneration niche. *Biological Review, Cambridge Philosophical Society* 52: 102-145.
- Guisan, A. and W. Thuiller. 2005. Predicting species distribution: offering more than simple habitat models. *Ecology letters* 8: 993-1009.

- Guldin, J. M. 2014. Adapting silviculture to a changing climate in the southern United States. Pages 173-192 in Vose, J.M. and K.D. Klepzig (eds.), *Climate change for adaptation and mitigation management options: a guide for natural resource managers in southern forest ecosystems*. CRC Press, Boca Raton, FL.
- Gustafson E. J., and B. R. Sturtevant. 2013. Modeling forest mortality caused by drought stress: implications for climate change. *Ecosystems* 16: 60–74.
- Guyette, R. P., M. A. Spetich, and M. C. Stambaugh. 2006. Historic fire regime dynamics and forcing factors in the Boston Mountains, Arkansas, USA. *Forest Ecology and Management* 234(1): 293-304.
- Gyenge, J., M. Fernández, M. Sarasola, and T. Schlichter. 2011. Stand density and drought interaction on water relations of *Nothofagus antarctica*: contribution of forest management to climate change adaptability. *Trees - Structure and Function* 25: 1111-1120.
- Haase, D. L. and R. Rose. 1993. Soil moisture stress induces transplant shock in stored and unstored 2+0 Douglas-fir seedlings of varying root volumes. *Forest Science* 39: 275-294.
- Haire, S.I. and K. McGarigal. 2008. Inhabitants of landscape scars: succession of woody plants after large, severe forest fires in Arizona and New Mexico. *The Southwestern Naturalist* 53(2): 146-161.
- Handler, S. D., J. Matthew, L. Iverson, E. Peters, R. Scheller, K. Wythers, L. Brandt, P. Butler, M. Janowiak, P. D. Shannon, C. Swanston, K. Barrett, R. Kolka, C. McQuiston, B. Palik, P. Reich, C. Turner, M. White, C. Adams, A. D'Amato, S. Hagell, P. Johnson, R. Johnson, M. Larson, S. Matthews, R. Montgomery, S. Olson, M. Peters, A. Prasad, J. Rajala, J. Daley, M. Davenport, M. Emery, D. Fehringer, C. Hoving, G. Johnson, L. Johnson, D. Neitzel, A. Rissman, C. Rittenhouse, and R. Ziel. 2014. *Minnesota Forest Ecosystem Vulnerability Assessment and Synthesis: A report from the Northwoods Climate Change Response Framework*. Gen. Tech. Rep. NRS-133. U.S. Department of Agriculture, Forest Service, Northern Research Station., Newtown Square, PA.
- Hanson P. J., and J. F. Weltzin. 2000. Drought disturbance from climate change: response of United States forests. *Science of the Total Environment* 262: 205-220.
- Harper, J. L. 1977. *Population biology of plants*. Academic Press, London.
- Henderson, J. P., and H. D. Grissino-Mayer. 2009. Climate–tree growth relationships of longleaf pine (*Pinus palustris* Mill.) in the Southeastern Coastal Plain, USA. *Dendrochronologia* 27: 31–43.
- Hersh, M. H., R. Vilgalys, and J. S. Clark. 2012. Evaluating the impacts of multiple generalist fungal pathogens on temperate tree seedling survival. *Ecology* 93: 511–520.
- Hicke, J. A., A. J. H. Meddens, C. D. Allen, C. A. Kolden. 2013. Carbon stocks of trees killed by bark beetles and wildfire in the western United States. *Environ. Res. Lett.* 8 035032 doi:10.1088/1748-9326/8/3/035032 .
- Hogg, E. T., J. P. Brandt, and B. Kochtubajda. 2005. Factors affecting interannual variation in growth of western Canadian aspen forests during 1951-2000. *Canadian Journal of Forest Research* 35: 610-622.
- Hood, L. A., M. D. Swaine, and P. A. Mason. 2004. The influence of spatial patterns of damping-off disease and arbuscular mycorrhizal colonization on tree seedling establishment in Ghanaian tropical forest soil. *Journal of Ecology* 92: 816–823.
- Hough, A. F., and R. D. Forbes. 1943. The ecology and silvics of forests in the High Plateau of Pennsylvania. *Ecological Monographs* 13: 299–320.

- Huntley, B. and H. J. B. Birks. 1983. An atlas of Past and Present Pollen Maps for Europe: 0 - 13000 years ago. Cambridge University Press, Cambridge, UK.
- Hutchinson, T., R. Long, R. Ford, and E. K. Sutherland. 2008. Fire history and the establishment of oaks and maples in second-growth forests. *Canadian Journal of Forestry Research* 38: 1184-1196.
- Ibáñez, I., J. S. Clark, S. LaDeau, and J. Hille Ris Lambers. 2007 Exploiting temporal variability to understand tree recruitment response to climate change. *Ecological Monographs* 77(2): 163-177.
- Ibanez, I., J. S. Clark, M. C. Dietze, K. Feeley, M. Hersh, S. LaDeau, A. McBride, N. E. Welch, and M. S. Wolosin. 2006. Predicting biodiversity change: Outside the climate envelope, beyond the species-area curve. *Ecology* 87: 1896-1906.
- Ichihara, Y. and K. Yamaji. 2009. Effect of light conditions on the resistance of current-year *Fagus crenata* seedlings against fungal pathogens causing damping-off in a natural beech forest: fungus isolation and histological and chemical resistance. *Journal of Chemical Ecology* 35: 1077–1085.
- Innes J. L. 1994. The occurrence of flowering and fruiting on individual trees over 3 years and their effects on subsequent crown condition, *Trees Structure and Function* 8: 139–150.
- Intergovernmental Panel on Climate Change. 1995. IPCC Second Assessment: Climate Change 1995. <http://ipcc.ch/pdf/climate-changes-1995/ipcc-2nd-assessment/2nd-assessment-en.pdf>.
- Intergovernmental Panel on Climate Change. 2014. Climate Change 2014: Impacts, Adaptation, and Vulnerability. Working Group II Contribution to the IPCC 5th Assessment Report Intergovernmental Panel on Climate Change, Stanford, CA.
- Iverson, L. R., A. M. Prasad, S. N. Matthews, and M. Peters. 2008b. Estimating potential habitat for 134 eastern US tree species under six climate scenarios. *Forest Ecology and Management* 254: 390-406.
- Iverson, L. R., A. M. Prasad, S. N. Matthews, and M. Peters. 2008c. Estimating potential habitat for 134 eastern US tree species under six climate scenarios. *Forest Ecology and Management* 254: 390-406.
- Iverson, L. R., M. W. Schwartz, and A. M. Prasad. 2004. How fast and far might tree species migrate in the eastern United States due to climate change? *Global Ecology and Biogeography* 13: 209-219.
- Iverson, L. R., T. F. Hutchinson, A. M. Prasad, and M. P. Peters. 2008a. Thinning, fire, and oak regeneration across a heterogeneous landscape in the eastern U.S.: 7-year results. *Forest Ecology and Management* 255: 3035-3050.
- Jackson, S. T., J. L. Betancourt, R. K. Booth, and S. T. Gray. 2009. Ecology and the ratchet of events: climate variability, niche dimensions, and species distributions. *Proceedings of the National Academy of Science* 106 (2): 19685-19692.
- Jactel, H., J. Petit, M. L. Desprez-Loustau, S. Delzon, D. Piou, A. Battisti, and J. Koricheva. 2012. Drought effects on damage by forest insects and pathogens: a meta-analysis. *Global Change Biology* 18: 1077-1091.
- Jiang, X., S. A. Rauscher, T. D. Ringler, D. M. Lawrence, A. P. Williams, C. D. Allen, A. L. Steiner, D. M. Cai, and N. G. McDowell. 2013. Projected Future Changes in Vegetation in Western North America in the Twenty-First Century. *Journal of Climate* 26.
- Joyce DG, Rehfeldt GE (2013) Climatic niche, ecological genetics, and impact of climate change on eastern white pine (*Pinus strobus* L.): Guidelines for land managers. *Forest Ecology and Management*, **295**, 173–192.

- Joyce, L. A., S. W. Running, D. D. Breshears, V. H. Dale, R. W. Malmshemer, R. N. Sampson, B. Sohngen, and C. W. Woodall. 2014. Ch. 7: Forests. *Climate Change Impacts in the United States: The Third National Climate Assessment*, J. M. Melillo, Terese (T.C.) Richmond, and G. W. Yohe, Eds., U.S. Global Change Research Program, 175-194. doi:10.7930/J0Z60KZC.
- Jump, A. S., C. Matyas, and J. Penuelas. 2009. The altitude-for-latitude disparity in the range retractions of woody species. *Trends in Ecology & Evolution* 24: 694-701.
- Kane, J. M., K. A. Meinhardt, T. Chang, B. L. Cardall, R. Michalet, and T. G. Whitman. 2011. Drought-induced mortality of a foundation species (*Juniperus monosperma*) promotes positive afterlife effects in understory vegetation. *Plant Ecology* 212: 733–741.
- Kane, J. M., F. L. Dugi, and T. E. Kolb. 2015. Establishment and growth of piñon pine regeneration vary by nurse type along a soil substrate age gradient in northern Arizona. *Journal of Arid Environments* 115: 113-119.
- Kayes, L. J., and D. B. Tinker. 2012. Forest structure and regeneration following a mountain pine beetle epidemic in southeastern Wyoming. *Forest Ecology and Management* 263: 57-66.
- Kelly, A. E. and M. L. Goulden. 2008. Rapid shifts in plant distribution with recent climate change. *Proceedings of the National Academy of Science* 105: 11823-11826.
- King, D. A., D. M. Bachelet, and A. J. Symstad. 2013. Climate change and fire effects on a prairie–woodland ecotone: projecting species range shifts with a dynamic global vegetation model. *Ecology and Evolution* 3(15):5076-5097.
- Klos, R. J., G. G. Wang, W. L. Bauerle, and J. R. Rieck. 2009. Drought impact on forest growth and mortality in the southeast USA: an analysis using Forest Health and Monitoring data. *Ecological Applications* 19: 699–708.
- Koenig, W.D. and J. M. H. Knops 2013. Large-scale spatial synchrony and cross-synchrony in acorn production by two California oaks. *Ecology* 94: 83–93.
- Kolb, P. F. and R. Robberecht. 1996. High temperature and drought stress effects on survival of *Pinus ponderosa* seedlings. *Tree Physiology* 16: 665-672.
- Kolb, T. E., J. K. Agee, P. Z. Fule, N. G. McDowell, K. Pearson, A. Sala, and R. H. Waring. 2007. Perpetuating old ponderosa pine. *Forest Ecology and Management* 249: 141-157.
- Kreye, J. K., J. M. Varner, J. K. Hiers, and J. Mola. 2013. Toward a mechanism for eastern North American forest mesophication: differential litter drying across 17 species. *Ecological applications* : a publication of the Ecological Society of America 23:1976–86.
- LaDeau, S. L. and J. S. Clark. 2001. Rising CO<sub>2</sub> levels and the fecundity of forest trees. *Science* 292: 95-98.
- Lafon, C. W. and S. M. Quiring. 2012. Relationships of fire and precipitation regimes in temperate forests of the Eastern United States. *Earth Interactions* 16:1–15.
- Legras, E. C., S. B. Vander Wall, and D. I. Board. The role of germination microsite in the establishment of sugar pine and Jeffrey pine seedlings. *Forest Ecology and Management* 260: 806-813.
- Lenihan, J. M., D. Bachelet, R. P. Neilson, and R. Drapek. 2008. Response of vegetation distribution, ecosystem productivity, and fire to climate change scenarios for California. *Climatic Change* 87: 215-230.
- Lenoir, J., J. C. Gégout, P. A. Marquet, P. de Ruffray, and H. Brisse. 2008. A significant upward shift in plant species optimum elevation during the 20th century. *Science* 320(5884): 1768-1771.
- Lesser, M.R. & S.T. Jackson. 2012. Making a stand: five centuries of population growth in colonizing stands of *Pinus ponderosa*. *Ecology* 93:1071-1081.

- Lesser, M.R. & S.T. Jackson. 2013. Contributions of long-distance dispersal to population growth in colonizing *Pinus ponderosa* populations. *Ecology Letters* 16:380-389.
- Lesser, M.R., T.L. Parchman & S.T. Jackson. 2013. Development of genetic diversity, differentiation and structure over 500 years in four ponderosa pine populations. *Molecular Ecology* 22:2640-2652.
- Linares, C. J., J. Camarero, A. Carreira. 2009. Interacting effects of changes in climate and forest cover on mortality and growth of the southernmost European fir forests. *Global Ecology and Biogeography* 18: 485–497.
- Lines, E. R., D. A. Coomes, and D. W. Purves. 2010. Influences of forest structure, climate and species composition on tree mortality across the eastern US. *PLoS ONE*, 5, e13212.
- Lischke, H., N. E. Zimmermann, J. Bolliger, S. Rickebusch, and T. J. Löffler. 2006. TreeMig: A forest-landscape model for simulating spatio-temporal patterns from stand to landscape scale. *Ecological Modelling* 199: 409-420.
- Littell, J. S., D. McKenzie, D. L. Peterson, and A. L. Westerling. 2009. Climate and wildfire area burned in western US ecoprovinces, 1916-2003. *Ecological Applications* 19: 1003-1021.
- Loarie SR, P. B. Duffy, H. Hamilton, G. P. Asner, C. B. Field, and D. D. Ackerly. 2009. The velocity of climate change. *Nature* 462: 1052-1055.
- Loehman, R. A., A. Corrow, and R. E. Keane. 2011. Modeling climate changes and wildfire interactions: Effects on whitebark pine (*Pinus albicaulis*) and implications for restoration, Glacier National Park, Montana, USA. *Forests*. 2: 832-860.
- Loik, M. E., D. D. Breshears, W. K. Lauenroth, and J. Belnap. 2004. A multi-scale perspective of water pulses in dryland ecosystems: climatology and ecohydrology of the western USA. *Oecologia* 14: 269-281
- Luo, Y. and H. Y. H. Chen. 2013. Observations from old forests underestimate climate change effects on tree mortality. *Nature Communications* 4:1655 doi: 10.1038/ncomms2681
- Lutz, J. A., J. W. van Wagtendonk, and J. F. Franklin. 2009. Twentieth-century decline of large-diameter trees in Yosemite National Park, California, USA. *Forest Ecology and Management* 257: 2296–2307.
- Lyford, M.E., S.T. Jackson, J.L. Betancourt, & S.T. Gray. 2003. Influence of landscape structure and climate variability on a late Holocene plant migration. *Ecological Monographs* 73:567-583.
- Lynch, C., and A. Hessler. 2010. Climatic controls on historical wildfires in West Virginia, 1939–2008. *Physical Geography* 31: 254–269.
- Man, G. 2013. Major Forest Insect and Disease Conditions in the United States: 2012. US Forest Service, SF-1023, Washington, D.C.
- Martin-Benito, D., V. Kint, M. del Río, B. Muys, and I. Cañellas. 2011. Growth responses of West-Mediterranean *Pinus nigra* to climate change are modulated by competition and productivity: Past trends and future perspectives. *Forest Ecology and Management* 262: 1030–1040.
- Martinson, E. J., Omi, P. N. 2013. Fuel treatments and fire severity: A metaanalysis. Res. Pap. RMRS-RP-103WWW. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 38 p.
- Mast, J.N. and J. Wolf. 2006. Spatial patch patterns and altered forest structure in middle elevation versus upper ecotonal mixed-conifer forests, Grand Canyon National Park, Arizona, USA. *Forest Ecology and Management* 236: 241-250.

- Matthews, S. N., L. R. Iverson, A. M. Prasad, M. P. Peters, and P. G. Rodewald. 2011. Modifying climate change habitat models using tree species-specific assessments of model uncertainty and life history-factors. *Forest Ecology and Management* 262: 1460-1472.
- McDowell, N. G., D. J. Beerling, D. D. Breshears, R. A. Fisher, K. F. Raffa, and M. Stitt. 2011. The interdependence of mechanisms underlying climate-driven vegetation mortality. *Trends in Ecology and Evolution* 26: 523-532.
- McDowell, N. G., H. D. Adams, J. D. Bailey, M. Hess, and T. E. Kolb. 2006. Homeostatic maintenance of ponderosa pine gas exchange in response to stand density changes. *Ecological Applications* 16: 1164-1182.
- McDowell, N.G., R.A. Fisher, C. Xu, J. C. Domec, T. Hölttä, D. S. Mackay, J. S. Sperry, A. Boutz, L. Dickman, N. Gehres, J. M. Limousin, A. Macalady, J. Martínez-Vilalta, M. Mencuccini, J. A. Plaut, J. Ogée, R. E. Pangle, D. P. Rasse, M. G. Ryan, S. Sevanto, R. H. Waring, A. P. Williams, E. A. Yezpez, and W. T. Pockman. 2013. Evaluating theories of drought-induced vegetation mortality using a multimodel-experiment framework. *New Phytologist* 200: 304-321.
- McEwan, R. W., J. M. Dyer, and N. Pederson. 2011. Multiple interacting ecosystem drivers: toward an encompassing hypothesis of oak forest dynamics across eastern North America. *Ecography* 34: 244–256.
- McKenney, D. W., J. H. Pedlar, K. Lawrence, K. Campbell, and M. F. Hutchison. 2007. Potential impacts of climate change on distribution of North American trees. *BioScience* 57: 939-948.
- McKenney, D. W., J. H. Pedlar, R. B. Rood, and D. Price. 2011. Revisiting projected shifts in the climate envelopes of North American trees using updated general circulation models. *Global Change Biology* 17: 2720-2730.
- McLachlan, J., J. S. Clark, and P. S. Manos. 2005. Molecular indicators of tree migration capacity under rapid climate change. *Ecology* 86: 2088-2098.
- McMahon, S. M., G. C. Parker, D. R. Miller. 2010. Evidence for a recent increase in forest growth. *Proceedings of the National Academy of Sciences* 107: 3611–3615.
- Medvigy, D. and P. R. Moorcroft. 2012. Predicting ecosystem dynamics at regional scales: an evaluation of a terrestrial biosphere model for the forests of northeastern North America. *Philosophical Transactions of the Royal Society B-Biological Sciences* 367: 222-235.
- Meier, E. S., H. Lischke, D. R. Schmatz, and N. E. Zimmermann. 2012. Climate, competition and connectivity affect future migration and ranges of European trees. *Global Ecology and Biogeography* 21:164-178.
- Melillo, J. M., Terese, T.C., Richmond, and G. W. Yohe, Eds., 2014: *Climate Change Impacts in the United States: The Third National Climate Assessment*. U.S. Global Change Research Program, 841 pp. doi:10.7930/J0Z31WJ2.
- Miao, S., C. B. Zou, and D. D. Breshears. 2009. Vegetation Responses to Extreme Hydrological Events: Sequence Matters. *The American Naturalist* 173:113-118
- Millar, C. I., R. D. Westfall, D. L. Delany, M. J. Bokach, A. L. Flint, and L. E. Flint. 2012. Forest mortality in high-elevation whitebark pine (*Pinus albicaulis*) forests of eastern California, USA; influence of environmental context, bark beetles, climatic water deficit, and warming. *Canadian Journal of Forest Research* 42(4): 749–765.
- Minnich, R. A. 2007. Southern California coniferous forest. In, *Terrestrial vegetation of California*, 3rd edition (M.G. Barbour, T. Keeler-Wolf, and A.S. Schoenherr, eds.). University of California Press , Chapter 18.



- Mok, H. F., S. K. Arndt, and C. R. Nitschke. 2012. Modelling the potential impact of climate variability on species regeneration potential in SE Australia. *Global Change Biology* 18: 1053–1072.
- Monleon, V. J. and H. E. Lintz. 2015. Evidence of Tree Species' Range Shifts in a Complex Landscape. *Plos One* 10:e0118069.
- Montwé D, Spiecker H, Hamann A (2015) Five decades of growth in a genetic field trial of Douglas-fir reveal trade-offs between productivity and drought tolerance. *Tree Genetics & Genomes*, **11**.
- Moore, M. M., W. W. Covington, P. Z. Fulé, D. J. Parsons, T. W. Swetnam, and N. L. Christensen. 1999. Reference conditions and ecological restoration: A southwestern ponderosa pine perspective. *Ecological Applications* 9: 1266-1277.
- Morin, X., C. Augspurger, and I. Chuine. 2007. Process-based modeling of species' distributions: what limits temperate tree species' range boundaries. *Ecology* 88: 2280-2291.
- Mueller, R. C., C. M. Scudder, M. E. Porter, R. Talbot Trotter, C. A. Gehring, and T. G. Whitham. 2005. Differential tree mortality in response to severe drought: evidence for long-term vegetation shifts. *Journal of Ecology* 93: 1085–1093.
- Nabel, J., N. Zurbriggen, and H. Lischke. 2013. Interannual climate variability and population density thresholds can have a substantial impact on simulated tree species' migration. *Ecological Modelling* 257: 88-100.
- National Climatic Data Center, 2014. North American Drought Monitor Indices. <http://www.ncdc.noaa.gov/temp-and-precip/drought/nadm/indices.php>
- Negron, Jose F.; McMillin, Joel D.; Anhold, John A.; Coulson, Dave 2009. Bark beetle-caused mortality in a drought-affected ponderosa pine landscape in Arizona, USA. *Forest Ecology and Management*. 257(4): 1353-1362.
- Nilsen, E. T., B. D. Clinton, T. T. Lei, O. K. Miller, S. W. Semones, and J. F. Walker. 2001. Does *Rhododendron maximum* L. (Ericaceae) Reduce the Availability of Resources above and Belowground for Canopy Tree Seedlings? *American Midland Naturalist* 145: 325-343.
- Nilsson, U. and G. Örlander. 1995. Effects of regeneration methods on drought damage to newly planted Norway spruce seedlings. *Canadian Journal of Forest Research* 25: 790-802.
- Norris, J., J.L. Betancourt, & S.T. Jackson. 2015. Late Holocene expansion of Ponderosa pine in the central Rockies. *Journal of Biogeography*. (in press).
- Nowacki, G. J. and M. D. Abrams. 2008. The Demise of Fire and “Mesophication” of Forests in the Eastern United States. *Bioscience* 58: 123-138.
- Nyland, R. D. 2007. *Silviculture : concepts and applications*. Long Grove, Ill. : Waveland Press, Long Grove, Ill.
- O'Connor, CD. 2013. Spatial and temporal dynamics of disturbance interactions along an ecological gradient. Ph.D. Dissertation, University of Arizona, Tucson, AZ.
- O'Hara, K.L. and L.M. Nagel. 2013. The Stand: Revisiting a central concept in forestry. *J. For.* 111(5):335-340.
- Ohmann, J. L., and T. A. Spies. 1998. Regional gradient analysis and spatial pattern of woody plant communities of Oregon forests. *Ecological Monographs* 68: 151-182.
- Oliva, J., J. Stenlid, and J. Martínez-Vilalta. 2014. The effect of fungal pathogens on the water and carbon economy of trees: implications for drought-induced mortality. *New Phytologist* 203:1028–1035.
- Oosting, H. J., 1942. An Ecological analysis of the plant communities of Piedmont, North Carolina. *American Midland Naturalist* 28: 1-126.

- Ouzts, J., T. Kolb, D. Huffman, and A. Sánchez-Meador. 2015. Post-fire ponderosa pine regeneration with and without planting in Arizona and New Mexico. *Forest Ecology and Management* 354: 281-290.
- Parmesan, C., and G. Yohe. 2003. A globally coherent fingerprint of climate change impacts across natural systems. *Nature* 421: 37-42.
- Parshall T. 1995. Canopy mortality and stand-scale change in a northern hemlock–hardwood forest. *Canadian Journal of Forestry Research* 25: 1466-1478 ISI.
- Parshall, T., and D. Foster. 2003. Fire on the New England landscape: regional and temporal variation, cultural and environmental controls. *Journal of Biogeography* 29: 1305–1317.
- Parsons, D. J. and S. H. DeBenedetti. 1979. Impact of fire suppression on a mixed-conifer forest. *Forest Ecology and Management* 2: 21–33.
- Pedersen, B. 1998. The role of stress in the mortality of midwestern oaks as indicated by growth prior to death. *Ecology* 79: 79–93.
- Pederson, N., A. R. Bell, E. R. Cook, U. Lall, N. Devineni, R. Seager, K. Eggleston, and K. P. Vranes. 2013. Is an Epic Pluvial Masking the Water Insecurity of the Greater New York City Region? *Journal of Climate* 26: 1339–1354.
- Pederson, N., A. R. Bell, T. A. Knight, C. Leland, N. Malcomb, K. J. Anchukaitis, K. Tackett, J. Scheff, A. Brice, B. Catron, W. Blozan, and J. Riddle. 2012. Long-term perspective on a modern drought in the American Southeast. *Environmental Research Letters* 7.
- Pederson, N., J. M. Dyer, R. W. McEwan, A. E. Hessel, C. Mock, D. A. Orwig, H. Rieder, and B. I. Cook. 2014. The legacy of episodic climatic events in shaping broadleaf-dominated forests. *Ecological Monographs*.
- Pederson, N., Tackett, K., McEwan, R.W., Clark, S., Cooper, A., Brosi, G., Eaton, R. & Stockwell, R.D. 2012. Long-term drought sensitivity of trees in second-growth forests in a humid region. *Canadian Journal of Forest Research* 42: 1837–1850.
- Pederson, N., J. M. Varner III, and B. J. Palik. 2008. Canopy disturbance and tree recruitment over two centuries in a managed longleaf pine landscape. *Forest Ecology and Management* 254: 85–95.
- Pelz, K. A., and F. W. Smith. 2013. How will aspen respond to mountain pine beetle? A review of literature and discussion of knowledge gaps. *Forest Ecology and Management* 299: 60-69.
- Perez-Ramos I. M., J. M. Ourcival, J. M. Limousin, and S. Rambal. 2010. Mast seeding under increasing drought: results from a long-term data set and from a rainfall exclusion experiment. *Ecology* 91: 3057–3068.
- Peters, M.P., Iverson, L.R., Matthews, S.N., 2014. Spatio-temporal trends of drought by forest type in the conterminous United States, 1960–2013. Res. Map NRS-7. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. [scale 1:12,000,000].
- Peters, M. P., L. R. Iverson, and S. N. Matthews. 2015. Long-term droughtiness and drought tolerance of eastern US forests over five decades. *Forest Ecology and Management* 345: 56-64.
- Pinto, J. R., J. D. Marshall, R. K. Dumroese, A. S. Davis, and D. R. Cobos. 2012. Photosynthetic response, carbon isotopic composition, survival, and growth of three stock types under water stress enhanced by vegetative competition. *Canadian Journal of Forest Research* 42: 333-344.
- Piovesan G. and J. M. Adams. 2001. Masting behaviour in beech: linking reproduction and climatic variation *Canadian Journal of Botany* 79:1039-1047.

- Pitelka, L. F., R. H. Gardner, J. Ash, S. Berry, H. Gitay, I. R. Noble, A. Saunders, R. H. W. Bradshaw, L. Brubaker, J. S. Clark, M. B. Davis, S. Sugita, J. M. Dyer, R. Hengeveld, G. Hope, B. Huntley, G. A. King, S. Lavorel, R. N. Mack, G. P. Malanson, M. McGlone, I. C. Prentice, and M. Rejmanek. 1997. Plant migration and climate change. *American Scientist* 85: 464-473.
- Prasad, A. M., J. Gardiner, L. Iverson, S. Matthews, and M. Peters. 2013. Exploring tree species colonization potentials using a spatially explicit simulation model: implications for four oaks under climate change. *Global Change Biology* 19: 2196–2208.
- Pucek, Z., W. Jedrzejewski, B. Jedrzejewska, and M. Pucek. 1993. Rodent population dynamics in a primeval deciduous forest (Bialowieza National Park) in relation to weather, seed crop, and predation. *Acta Theriologica* 38: 199–232.
- Quarterman, E. and C. Keever. 1962. Southern mixed hardwood forest: Climax in the Southeastern Coastal Plain, U.S.A.. *Ecological Monographs* 32: 167-185.
- Raffa K. F., B. H. Aukema, B. J. Bentz, A. L. Carroll, J. A. Hicke, M. G. Turner, and W. H. Romme. 2008. Cross-scale drivers of natural disturbances prone to anthropogenic amplification: the dynamics of bark beetle eruptions. *BioScience* 58: 501–517.
- Rahman, M.S., M.G. Messina, and R.F. Fisher. 2006. Intensive forest management affects loblolly pine (*Pinus taeda* L.) growth and survival on poorly drained sites in southern Arkansas. *Southern Journal of Applied Forestry* 30(2): 79-85.
- Rapacciuolo, G., S. P. Maher, A. C. Schneider, T. T. Hammond, M. D. Jabis, R. E. Walsh, K. J. Iknayan, G. K. Walden, M. F. Oldfather, and D. D. Ackerly. 2014. Beyond a warming fingerprint: individualistic biogeographic responses to heterogeneous climate change in California. *Global change biology* 20: 2841-2855.
- Redmond, M. D. and N. N. Barger. 2013. Tree regeneration following drought- and insect-induced mortality in piñon-juniper woodlands. *New Phytologist* 200: 402-412.
- Redmond, M. D., F. Forcella, and N. N. Barger. 2012. Declines in pinyon pine cone production associated with regional warming. *Ecosphere* 3(12): 120.
- Rehfeldt GE, Ying CC, Spittlehouse DL, Hamilton DA (1999) Genetic responses to climate in *Pinus contorta*: Niche breadth, climate change, and reforestation. *Ecological Monographs*, **69**, 375–407.
- Rehfeldt GE, Leites LP, Bradley St Clair J, Jaquish BC, Sáenz-Romero C, López-Upton J, Joyce DG (2014) Comparative genetic responses to climate in the varieties of *Pinus ponderosa* and *Pseudotsuga menziesii*: Clines in growth potential. *Forest Ecology and Management*, **324**, 138–146.
- Reich, P. B., K. M. Sendall, K. Rice, R. L. Rich, A. Stefanski, S. E. Hobbie, and R. A. Montgomery. 2015. Geographic range predicts photosynthetic and growth response to warming in co-occurring tree species. *Nature Clim. Change* 5:148-152.
- Roberts, S. D., C. A. Harrington, and T. A. Terry. 2005. Harvest residue and competing vegetation affect soil moisture, soil temperature, N availability, and Douglas-fir seedling growth. *Forest Ecology and Management* 205: 333-350.
- Roccaforte, J.P., P.Z. Fulé, W.W. Chancellor, and D.C. Laughlin. 2012. Woody debris and tree regeneration dynamics following severe wildfires in Arizona ponderosa pine forests. *Canadian Journal of Forest Research* 42:593-604.
- Rozas V. 2005. Dendrochronology of pedunculate oak (*Quercus robur* L.) in an old-growth pollarded woodland in northern Spain: tree-ring growth responses to climate. *Annals of Forest Science* 62: 209–218.

- Savage, M. and J. N. Mast. 2005. How resilient are southwestern ponderosa pine forests after crown fires? *Canadian Journal of Forest Research* 35: 967-977.
- Savage, M., J. N. Mast, and J. J. Feddema. 2013. Double whammy: high-severity fire and drought in ponderosa pine forests of the Southwest. *Canadian Journal of Forest Research* 43: 570-583.
- Savage, M., P. M. Brown, and J. Feddema. 1996. The role of climate in a pine forest regeneration pulse in the southwestern United States. *Ecoscience* 3:310–318.
- Savolainen O, Pyhäjärvi T, Knürr T (2007) Gene Flow and Local Adaptation in Trees. *Annual Review of Ecology, Evolution, and Systematics*, **38**, 595–619.
- Scheller, R. M. and D. J. Mladenoff. 2008. Simulated effects of climate change, fragmentation, and inter-specific competition on tree species migration in northern Wisconsin, USA. *Climate Research* 36:191-202.
- Schliep, E.M., A.E. Gelfand, and J.S. Clark. 2015. Stochastic modeling for velocity of climate change. *Journal of Agricultural, Biological, and Environmental Statistics*, 20, 323-342.
- Schumacher, F. X. and B. B. Day. 1939. The influence of precipitation upon the width of annual rings of certain timber trees. *Ecological Monographs* 9: 387–429.
- Schuster, W. S. L., K. L. Griffin, H. Roth, M. H. Turnbull, D. Whitehead, and D. T. Tissue. 2008. Changes in composition, structure and aboveground biomass over seventy-six years (1930-2006) in the Black Rock Forest, Hudson Highlands, southeastern New York State. *Tree Physiology* 28: 537-549.
- Schwartz, M. W., J. J. Hellmann, M. M. Jason, D. Sax, J. O. Borevitz, J. Brennan, A. E. Camacho, G. Ceballos, J. R. Clark, H. Doremus, R. Early, J. R. Ettlerson, D. Fielder, J. L. Gill, P. Gonzalez, N. Green, L. Hannah, D. Jamieson, D. Javeline, B. A. Minter, J. Odenbaugh, S. Polasky, D. M. Richardson, T. Root, and H. D. Safford. 2012. Managed relocation: integrating the scientific, regulatory, and ethical challenges. *Bioscience* 62: 732–743.
- Shinneman, D J, Baker, W L, Rogers, P C, Kulakowski, D. 2013. Fire regimes of quaking aspen in the Mountain West. *Forest Ecology and Management* 299: 22-34
- Shuman, B., A. K. Henderson, C. Plank, I. Stefanova, and S. S. Ziegler. 2009. Woodland-to-forest transition during prolonged drought in Minnesota after ca. AD 1300. *Ecology* 90: 2792–807.
- Silvertown, J. W. 1987. *Introduction to Plant Population Ecology*. Longman, White Plains, NY.
- Sitch, S., B. Smith, I. C. Prentice, A. Arneth, A. Bondeau, W. Cramer, J. O. Kaplan, S. Levis, W. Lucht, M. T. Sykes, K. Thonicke, and S. Venevsky. 2003. Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Global Change Biology* 9(2): 161-185.
- Snell, R. S. 2014. Simulating long-distance seed dispersal in a dynamic vegetation model. *Global Ecology and Biogeography* 23: 89-98.
- Song, X. and X. Zeng. 2014. Investigation of uncertainties of establishment schemes in dynamic global vegetation models. *Advances in Atmospheric Sciences* 31: 85-94.
- Sork VL, Davis FW, Westfall R, Flint A, Ikegami M, Wang H, Grivet D (2010) Gene movement and genetic association with regional climate gradients in California valley oak (*Quercus lobata* Nee) in the face of climate change. *Molecular Ecology*, **19**, 3806–3823.
- Stahle, D. W. and M. K. Cleaveland. 1992. Reconstruction and analysis of spring rainfall over the southeastern US for the past 1000 years. *Bulletin of the American Meteorological Society* 73: 1947–1961.

- Stahle, D. W., M. K. Cleaveland, and J. Hehr. 1988. North Carolina climate changes reconstructed from tree rings: AD 372 to 1985. *Science* 240: 1517–1519.
- Sthultz, C. M., C. A. Gehring, and T. G. Whitham. 2009. Deadly combination of genes and drought: Increased mortality of herbivore-resistant trees in a foundation species. *Global Change Biology* 15: 949-961.
- Stogsdili Jr, W. R., R. F. Wittwer, T. C. Hennessey, and P. M. Dougherty. 1992. Water use in thinned loblolly pine plantations. *Forest Ecology and Management* 50: 233-245.
- Swetnam, T. W. and J. L. Betancourt. 1998. Mesoscale disturbance and ecological response to decadal climatic variability in the American Southwest. *Journal of Climate* 11: 3128-3147.
- Taylor, M., D. L. Haase, and R. L. Rose. 2009. Fall Planting and Tree Shelters for Reforestation in the East Washington Cascades. *Western Journal of Applied Forestry* 24:173-179.
- Thomas, Z. and K.M. Waring. 2015. Enhancing resiliency and restoring ecological attributes in second-growth ponderosa pine stands in northern New Mexico, USA. *Forest Science* 61(1):93-104.
- Thuiller, W., S. Lavorel, M. B. Araujo, M. T. Sykes, and I. C. Prentice. 2005. Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences of the United States of America* 102: 8245-8250.
- Tinner, W., D. Colombaroli, O. Heiri, P. D. Henne, M. Steinacher, J. Untenecker, E. Vescovi, J. R. M. Allen, G. Carraro, M. Conedera, F. Joos, A. F. Lotter, J. Luterbacher, S. Samartin, and V. Valsecchi 2013. The past ecology of *Abies alba* provides new perspectives on future responses of silver fir forests to global warming. *Ecological Monographs* 83:419–439.
- Treyger, A. L. and C. A. Nowak. 2011. Changes in tree sapling composition within powerline corridors appear to be consistent with climatic changes in New York State. *Global Change Biology* 17:3439-3452.
- Trottier-Picard, A., E. Thiffault, A. DesRochers, D. Paré, N. Thiffault, and C. Messier. 2014. Amounts of logging residues affect planting microsites: A manipulative study across northern forest ecosystems. *Forest Ecology and Management* 312: 203-215.
- USDA Forest Service and US Geological Survey. 2000. Forest Cover Types. <http://www.fia.fs.fed.us/library/maps/>
- van Mantgem, P. J., N. L. Stephenson, J. C. Byrne, L. D. Daniels, J. F. Franklin, P. Z. Fulé, M. E. Harmon, A. J. Larson, J. M. Smith, A. H. Taylor, and T. T. Veblen. 2009. Widespread increase of tree mortality rates in the western United States. *Proceedings of the National Academy of Science* 323: 521-524.
- Vanderwel M. C., V. S. Lyutsarev, and D. W. Purves. 2013. Climate-related variation in mortality and recruitment determine regional forest-type distributions. *Global Ecology and Biogeography* 22: 1192–1203.
- Voelker, S., R. Muzika, and R. Guyette. 2008. Individual tree and stand level influences on the growth, vigor, and decline of red oaks in the Ozarks. *Forest Science* 54: 8–20.
- Wackerman, A.E. 1929. Why prairies in Arkansas and Louisiana? *Journal of Forestry* 27(6): 726-734.
- Waltz, A. E. M., Stoddard, M. T., Kalies, E. L., Springer, J. D., Huffman, D. W., Sánchez-Meador, A. 2014. Effectiveness of fuel reduction treatments: Assessing metrics of forest resiliency and wildfire severity after the Wallow Fire, AZ. *Forest Ecology and Management* 334: 43-52.

- Wang G. G., S. Chhin, and W. L. Bauerle. 2006. Effect of natural atmospheric CO<sub>2</sub> fertilization suggested by open-grown white spruce in a dry environment. *Global Change Biology* 12: 601–610.
- Weed, A. S., M. P. Ayres, and J. A. Hicke. 2013. Consequences of climate change for biotic disturbances in North American forests. *Ecological Monographs* 83: 441-470.
- Wehner, M., D. R. Easterling, J. H. Lawrimore, R. R. Heim, R. S. Vose, and B. D. Santer. 2011. Projections of Future Drought in the Continental United States and Mexico. *Journal of Hydrometeorology* 12(6): 1359-1377.
- Westerling, A. L., H. G. Hidalgo, D. R. Cayan, and T. W. Swetnam. 2006. Warming and earlier spring increase western US forest wildfire activity. *Science* 313: 940-943.
- Westerling, A. L., M. G. Turner, E. A. H. Smithwick, W. H. Romme, and M. G. Ryan. 2011. Continued warming could transform Greater Yellowstone fire regimes by mid-21st century. *Proceedings of the National Academy of Sciences* 108: 13165-13170.
- Will, R., T. Hennessey, T. Lynch, R. Holeman, and R. Heinemann. 2010. Effects of planting density and seed source on loblolly pine stands in southeastern Oklahoma. *Forest Science* 56(5): 437-443.
- Williams, A.P., C.D. Allen, T.W. Swetnam, C.I. Millar, J. Michaelsen, C.J. Still, and S.W. Leavitt. 2010. Forest responses to increasing aridity and warmth in southwestern North America. *Proceedings of the National Academy of Sciences, U.S.A.* 107:21289-21294.
- Williams, A. P., C. D. Allen, A. K. Macalady, D. Griffin, C. A. Woodhouse, M. M. Meko, T. W. Swetnam, S. A. Rauscher, R. Seager, H. D. Grissino-Mayer, J. S. Dean, E. R. Cook, C. Gangodagamage, M. Cai, and N. G. McDowell. 2013. Temperature as a potent driver of regional forest drought stress and tree mortality. *Nature Climate Change* 3: 292-297.
- Woodall, C., C. M. Oswalt, J. A. Westfall, C. H. Perry, M. D. Nelson, and A. O. Finley. 2009. An indicator of tree migration in forests of the eastern United States. *Forest Ecology and Management* 257: 1434-1444.
- Woodall, C.W., G. M. Domke, K. Riley, C. M. Oswalt, S. J. Crocker, and G. W. Yohe. 2013b. Developing a framework for assessing global change risks to forest carbon stocks. *PLOS One*. 8: e73222.
- Worrall, J. J., G. E. Rehfeldt, A. Hamann, E. H. Hogg, S. B. Marchetti, M. Michaelian, and L. K. Gray. 2013. Recent declines of *Populus tremuloides* in North America linked to climate. *Forest Ecology and Management* 299: 35-51.
- Wyckoff, P. H. and J. S. Clark. 2002. The relationship between growth and mortality for seven co-occurring tree species in the southern Appalachian Mountains. *Journal of Ecology* 90: 604-615.
- Yaussy, D. A., L. R. Iverson, and S. N. Matthews. 2013. Competition and Climate Affects US Hardwood-Forest Tree Mortality. *Forest Science* 59: 416-430.
- Yeh, H. Y., and L. C. Wensel. 2000. The relationship between tree diameter growth and climate for coniferous species in northern California. *Canadian Journal of Forest Research* 30: 1463-1471.
- Zhang, J., J. Webster, R. F. Powers, and J. Mills. 2008. Reforestation after the Fountain Fire in northern California: An untold success story. *Journal of Forestry* 106(8): 425-430.
- Zhu, K, C. W. Woodall, S. Ghosh, A. E. Gelfand, and J. S. Clark. 2014. Dual impacts of climate change: forest migration and turnover through life history. *Global Change Biology* 20: 251-264.

- Zhu, K., C. W. Woodall, and J. S. Clark. 2012. Failure to migrate: lack of tree range expansion in response to climate change. *Global Change Biology* 18: 1042-1052.
- Zimmermann N.E., N. G. Yoccoz, T. C. Edwards, E. S. Meier, W. Thuiller, A. Guisan, D. R. Schmatz, and P. B. Pearman. 2009. Climatic extremes improve predictions of spatial patterns of tree species. *Proceedings of the National Academy of Science USA* 106: 19723-19728.