



## LJMU Research Online

**Byrne, PA, Wood, PJ and Reid, I**

**The impairment of river systems by metal mine contamination: A review including remediation options**

<http://researchonline.ljmu.ac.uk/id/eprint/679/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Byrne, PA, Wood, PJ and Reid, I (2011) The impairment of river systems by metal mine contamination: A review including remediation options. Critical Reviews in Environmental Science and Technology, 42 (19). pp. 2017-2077. ISSN 1064-3389**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

1 **The impairment of river systems by metal mine contamination: A review**  
2 **including remediation options**

3

4 **Byrne, P.,<sup>1</sup> Wood, P.J.,<sup>2</sup> and Reid, I.<sup>2</sup>**

5 1. Centre for Earth and Ecosystem Responses to Environmental Change,  
6 School of Science and Environment, Manchester Metropolitan  
7 University, Manchester, M1 5GD, U.K.

8 2. Centre for Hydrological and Ecosystem Science, Department of  
9 Geography, Loughborough University, Loughborough, Leicestershire,  
10 LE11 3TU, UK.

11

12 **Author for Correspondence**

13 Patrick Byrne

14 Centre for Earth and Ecosystem Responses to Environmental Change,  
15 School of Science and Environment, Manchester Metropolitan University,  
16 Manchester, M1 5GD, U.K.

17 Tel: 44-161-2473660

18 Fax: 44-161-2476318

19 E-mail: [p.byrne@mmu.ac.uk](mailto:p.byrne@mmu.ac.uk)

20

21

22

23

24 **Abstract**

25 Contamination of aquatic environments as a consequence of deep metal  
26 mining for Pb, Zn, Cu, Cd and Fe is of widespread international concern.  
27 Pollution resulting from metal mining activities can result in significant  
28 environmental and ecological degradation and can pose serious risks to  
29 human health through contamination of food and drinking water. This paper  
30 provides a review of the impacts of deep metal mine water discharges on  
31 riverine sedimentology, hydrology and ecology and explores strategies for the  
32 restoration of rivers draining historically abandoned metal mines.

33

34 Physical processes of mine waste dispersal are relatively well understood.  
35 Chemical processes are more complex and much research is now focussed  
36 on understanding geochemical and mineralogical controls on metal  
37 attenuation and release. Recent advances in numerical modelling and  
38 geochemical tracing techniques offer the possibility of identifying present and  
39 predicted future patterns of contamination at the catchment scale.

40

41 The character of mine water has been extensively studied. However,  
42 documented impacts on aquatic ecosystems can vary widely depending on a  
43 range of hydroclimatological and geochemical factors. Numerous studies  
44 have shown that the majority of the annual metal flux in rivers draining mining-  
45 impacted regions occurs during the summer and autumn months as a result of  
46 water table drawdown, sulphide oxidation and dissolution and flushing of  
47 metal salts during subsequent storm periods. There have been few high-

48 resolution studies of stormflow hydrochemistry, despite the importance of high  
49 flows in the translocation of mine wastes.

50

51 A growing number of studies have documented chronic and acute toxic  
52 effects of mine water contaminants, based on both field and laboratory  
53 research, with specific reference to riverine macroinvertebrates. Common  
54 bioindices have been used to examine the impacts of mine water  
55 contaminants on macroinvertebrate ecology, although the success of these  
56 indices has been mixed. Sublethal biomonitoring techniques, as distinct from  
57 traditional laboratory bioassays with lethal endpoints, have gained  
58 prominence as a means of detecting behavioural and physiological responses  
59 of an organism to pulses of contaminants. The development of Biotic Ligand  
60 Models (BLMs) has allowed organism physiology and important  
61 environmental parameters to be factored into assessments of metal toxicity.

62

63 The strategies and technologies available for mine water remediation are  
64 considered and key knowledge gaps are highlighted. Passive remediation  
65 technologies offer a low cost and sustainable alternative to chemical  
66 treatment of deep metal mine discharges. However, at present, these systems  
67 generally fail to remove toxic metals associated with metal mine drainage to  
68 an acceptable standard. New phytoremediation techniques offer the possibility  
69 of immobilisation and extraction of toxic metals in mine spoil and  
70 contaminated soils.

71

72 We conclude by identifying key recommendations for future research:

- 73 (1) Researchers and regulators should consider bioavailable metal fractions in  
74 contaminated sediments, as opposed to total metal concentrations, if  
75 sediment ecotoxicity is to be accurately measured. In addition, more  
76 studies should make use of new spectroscopic techniques (e.g., XANES)  
77 capable of providing more detailed information on metal speciation and,  
78 therefore, sediment ecotoxicity.
- 79 (2) There is a need for better sampling and monitoring of toxic metal  
80 concentrations and fluxes during stormflows in mining-impacted river  
81 systems, especially given future predicted increases in stormflow  
82 occurrence. In addition, further research is required to help understand the  
83 potential toxicological impacts of stormflows in mining-impacted  
84 catchments.
- 85 (3) Further research is required to develop biological indices to identify the  
86 impacts of mine water contamination on macroinvertebrate communities.
- 87 (4) New substrates and techniques for remediation of metal-rich mine waters  
88 are currently being investigated and pilot studies undertaken in the  
89 laboratory and field. Many show promising results at the laboratory scale  
90 but large-scale pilot treatment plants are required to test the efficiency and  
91 long-term sustainability under field conditions.
- 92 (5) An interdisciplinary approach, incorporating the collaborative expertise and  
93 knowledge regarding sedimentological / geological, hydrological, chemical  
94 and ecological consequences of active and historic deep metal mining, is  
95 advocated and should be utilised for effective river basin management and  
96 the remediation and restoration of impacted sites.

97

98 **Keywords:** metal mine, acid mine drainage, river sediment, flood

99 hydrochemistry, benthic macroinvertebrate, mine remediation

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

## 122 **1. Introduction**

123 Environmental impacts of mining on aquatic ecosystems have received  
124 increasing attention in recent years (Gray, 1998; Smolders *et al.*, 2003; Olias  
125 *et al.*, 2004; Batty *et al.*, 2010). Acidic drainage associated with the  
126 abandonment of coal mining activity has been a particular focus of research  
127 (Banks and Banks, 2001). Contaminated discharge from abandoned metal  
128 mines and their spoil heaps has received less attention, reflecting the highly  
129 variable responses associated with the complex and frequently site-specific  
130 hydrogeological context of each, and the highly variable hydrogeochemical  
131 characteristics of the discharge (Environment Agency, 2008a). However,  
132 metal mine discharges have resulted in the severe degradation of many rivers  
133 across the globe (Gray, 1998; Gundersen and Stienes, 2001; Olias *et al.*,  
134 2004; Sola *et al.*, 2004; Poulton *et al.*, 2010).

135

136 Metal mining regions occur on all continents except Antarctica and even  
137 extend to the continental shelf in certain areas where former floodplains have  
138 been submerged by sea-level rise resulting from global warming (Aleva,  
139 1985). As a consequence, significant contamination of the landscape,  
140 including riverine and riparian habitats, has been reported internationally  
141 (Smolders *et al.*, 2003; Asta *et al.*, 2007; Edraki *et al.*, 2005; Gilchrist *et al.*,  
142 2009; Brumbaugh *et al.*, 2010). The most severely contaminated discharges  
143 typically occur shortly after abandonment of a site, when artificial dewatering  
144 has ceased and groundwater levels recover (Robb, 1994). Rising oxygenated  
145 groundwater within deep mines interacts with metal sulphides in exposed  
146 rockfaces, generating a leachate, typically characterised by low pH and high

147 concentrations of dissolved toxic metals and sulphates (Braungardt *et al.*,  
148 2003; Gilchrist *et al.*, 2009). Where the water table reaches the surface,  
149 leachate may enter rivers and lakes as drainage from mine shafts and mine  
150 drainage levels (adits), whilst rainwater may infiltrate through surface spoil  
151 heaps and tailings to enter streams and other surface water bodies.

152

153 Within riverine systems receiving metalliferous drainage, the composition and  
154 health of plant and animal communities can be severely impaired through the  
155 combined toxicity of reactive metals in both the water column and sediments,  
156 sulphates and acidity (Sola *et al.*, 2004; Schmitt *et al.*, 2007; Batty *et al.*,  
157 2010; Chapa-Vargas *et al.*, 2010). Aqueous metal concentrations generally  
158 decline downstream of contaminated sources due to the precipitation of oxide,  
159 hydroxide and sulphate phases, and co-precipitation or sorption of metals  
160 onto these phases (Hudson-Edwards *et al.*, 1999b). However, iron hydroxide  
161 'ochre' and other metal precipitates can cover the entire river bed in extreme  
162 instances and degrade habitat quality and important breeding and feeding  
163 areas for instream organisms (Batty, 2005; Mayes *et al.*, 2008). Chronic  
164 contamination of riverine systems can be exacerbated by episodic flood  
165 events (Bradley 1984; Hudson-Edwards *et al.* 1999a; Dennis *et al.* 2009) or  
166 by the failure of tailing dams (Hudson-Edwards *et al.*, 2003; Macklin *et al.*,  
167 2003; Sola *et al.*, 2004). Such events have led to significant ecological  
168 degradation in many regions of the world and have severely impacted  
169 communities dependent on local rivers and their floodplains for food and  
170 livelihood (Macklin *et al.*, 2006; Taylor *et al.*, 2010).

171



172 Environmental degradation resulting from metal mining is not restricted to  
173 regions of the world where recent or active mineral exploitation is occurring. In  
174 the UK, metal mining reached its peak in the mid-nineteenth century when, for  
175 a time, the UK was the largest lead, tin and copper producer in the world  
176 (Lewin and Macklin, 1987). Following a global reduction in metal prices  
177 associated with the discovery of large deposits of lead and copper in the  
178 Iberian Peninsula, South America and Australia during the late 19<sup>th</sup> and early  
179 20<sup>th</sup> centuries, a decline of metal mining occurred throughout the UK. Today,  
180 the number of abandoned metal mines in England and Wales is estimated at  
181 over 3,000 (Jarvis *et al.*, 2007). The historical legacy of these mines is still  
182 present in the landscape in the form of spoil heaps, abandoned adits and  
183 shafts, and derelict structures. The historical metal mining industry, long  
184 forgotten and often far removed from manufacturing centre's, has left a  
185 significant legacy of environmental contamination which will persist for  
186 centuries to millennia (Environment Agency, 2002; Macklin *et al.*, 2006).  
187 Approximately 20% of all water quality objective failures in England and  
188 Wales are due to drainage from abandoned metal mines (Environment  
189 Agency, 2006). The severity of the problem is underscored by the view of the  
190 Environment Agency of England and Wales that metal mine drainage poses  
191 the most serious threat to water quality objectives after diffuse agricultural  
192 pollution (Environment Agency, 2006).

193

194 Since the 1960s, concerns over the environmental impacts of historic metal  
195 mining activities have gained increasing significance and this is reflected in  
196 the growing body of literature on the topic (e.g., Macklin *et al.*, 2006; Batty *et*

197 *al.*, 2010). However, due to the highly variable nature of environmental  
198 degradation of surface waters draining metal mines and the site-specific  
199 nature of many impacts, the literature is scattered through a wide range of  
200 published sources (Wolkersdorfer, 2004). Unlike most review papers to date,  
201 which largely focus on specific environmental compartments in relative  
202 isolation to the wider aquatic ecosystem, this review paper aims to use an  
203 interdisciplinary perspective to critically review: (1) the sedimentological,  
204 hydrological and ecological impacts of metal mining activities; and (2) the  
205 potential for remediation of metal mine sites and the existing remediation  
206 technologies available.

207

208 The review is organised into 5 main sections. Mine water chemistry has been  
209 studied extensively (e.g., Younger *et al.*, 2002) and is generally well  
210 understood. Therefore, the purpose of section 2 is to provide a brief overview  
211 of the primary variables influencing the generation and character of metal  
212 mine drainage. There have been several systematic reviews of the  
213 sedimentological impacts of mining on the fluvial environment which have  
214 documented the physical and chemical factors controlling metal dispersal and  
215 storage in mining affected rivers systems (Lewin and Macklin, 1987; Macklin,  
216 1996; Miller, 1997). In addition, new technologies and approaches to help  
217 control and remediate sediment contamination have been widely considered  
218 (e.g., Macklin *et al.*, 2006). Section 3 provides a review of the recent  
219 developments centred on new spectroscopic methods for the measurement of  
220 metal mobility and speciation, and evaluate the performance of sediment  
221 environmental quality standards. Section 4 of this review examines the

222 catchment hydrological factors which influence the character of metal mine  
223 drainage in fluvial systems and discusses the important role of stormflows in  
224 transporting mine wastes from mine sites. In section 5, the ecological impacts  
225 associated with metal mines are examined with specific reference to benthic  
226 macroinvertebrate communities. While a significant body of research has  
227 been devoted to examining impacts on fish communities (Hallare *et al.*, 2010),  
228 the benthic lifestyle of macroinvertebrates makes them more representative of  
229 local environmental conditions, and, therefore, more reliable indicators of  
230 biological stress. Previous reviews by Gerhardt (1993) and Batty *et al.* (2010)  
231 have considered the impact of toxic metals and acidity on macroinvertebrates.  
232 The present review builds on previous reviews by considering new  
233 developments in biomonitoring techniques and sublethal measurements of  
234 toxicity assessment. In the final section, remediation practices and  
235 technologies to treat metal mine discharges are evaluated. In each of the four  
236 key review sections (sedimentology, hydrology, ecology, remediation), we  
237 highlight the key research gaps that remain and identify opportunities for  
238 future research.

239

240 Given that previous reviews have considered the environmental impacts  
241 associated with deep and surficial coal mining (Robb, 1994; Banks and  
242 Banks, 2001; Younger, 2002), and in particular acid mine drainage (Robb and  
243 Robinson, 1995; Banks *et al.*, 1997; Gray, 1997), this review focuses on the  
244 impact of deep metal mines on riverine ecosystems with a particular emphasis  
245 on the following widely exploited metals: lead (Pb), zinc (Zn), copper (Cu),  
246 cadmium (Cd) and iron (Fe). All of these metals frequently occur at high

247 concentrations within waters draining metal mines (Novotny, 1995; Younger *et*  
248 *al.*, 2002). The review has broad geographical significance, but highlights  
249 several case studies from the UK to illustrate some of the historic impacts of  
250 metal mining activities on riverine ecosystems. Two search strategies were  
251 used to identify relevant empirical papers. First, key word and title searches of  
252 electronic databases were undertaken independently by the authors before  
253 comparing results. The databases searched were: ASFA Aquatic Sciences,  
254 Biological Sciences, Science Direct, SCOPUS, Toxline, Web of Science and  
255 Zetoc. The key search words were: metal mine, heavy metals, toxic metals,  
256 acid mine drainage, river sediment, flood hydrochemistry, benthic  
257 macroinvertebrate, mine remediation and environmental quality. Databases  
258 were searched from inception to December 2010. Second, relevant  
259 references within any identified papers were followed up. Searches were  
260 limited to papers published in English.

261

## 262 **2. Mine water chemistry**

263 Sulphidic minerals such as galena (lead sulphide - PbS), sphalerite (zinc  
264 sulphide – ZnS) and pyrite (iron disulphide – FeS<sub>2</sub>) are amongst the most  
265 commonly mined metal ores (Novotny, 1995). These minerals are formed  
266 under reducing conditions in the absence of oxygen and remain chemically  
267 stable in dry, anoxic and high pressure environments deep underground.  
268 However, these solid phases become chemically unstable when they are  
269 exposed to the atmosphere (oxygen and water) through natural weathering  
270 processes and long-term landform evolution or anthropogenic activities such  
271 as mining (Johnson, 2003). A series of complex biogeochemical reactions

272 occurs in sulphide weathering processes, leading to the generation of a  
273 potentially toxic leachate and its release into the environment (**Figure 1**;  
274 Younger *et al.*, 2002; Johnson, 2003; Evangelou and Zhang, 1995). The  
275 leachate generated during the sulphide weathering process is complex and is  
276 often referred to as acid mine drainage (AMD) or acid rock drainage (ARD). It  
277 is most commonly characterised by high levels of dissolved toxic metals and  
278 sulphates and low pH (Robb and Robinson, 1995; Braungardt *et al.*, 2003).  
279 However, it should be noted that metal mine discharges are not always acidic  
280 (Banks *et al.*, 2002). In general, an increase in pyrite content of the country  
281 rock results in greater acidity; an increase in base-metal sulphides results in  
282 greater toxic metal concentrations; while an increase in carbonate and silicate  
283 content can result in highly alkaline waters (Oyarzun *et al.*, 2003; Alderton *et*  
284 *al.*, 2005; Cidu and Mereu, 2007). In the UK, much of central and north Wales  
285 is underlain by Lower Palaeozoic shales and mudstones with low  
286 concentrations of base materials (Evans and Adams, 1975). As a result, many  
287 of the headwater streams of the region have low acid-buffering capability,  
288 resulting in extremely acidic discharges containing high levels of dissolved  
289 toxic metals (Abdullah and Royle, 1972; Fuge *et al.*, 1991; Boulton *et al.*, 1994;  
290 Neal *et al.*, 2005). In contrast, in those parts of the English Peak District  
291 where carbonate lithology predominates (Carboniferous Limestone), neutral to  
292 basic mine discharges are common and these have significantly lower  
293 concentrations of dissolved toxic metals (Smith *et al.*, 2003). Aside from  
294 lithology and mineralogy, the character of mine water pollution can vary  
295 considerably between regions as a result of the grain size distribution of  
296 tailings and spoil (Hawkins, 2004), the exposed mineral surface area

297 (Younger *et al.*, 2002), the concentration of reactants such as dissolved  
298 oxygen (Wilkin, 2008), and microbial activity (Hallberg and Johnson, 2005;  
299 Natarajan *et al.*, 2006; Balci, 2008). The highly variable nature of water  
300 chemistry associated with metal mine discharges is outlined in **Table 1**.

301

### 302 **3. Sedimentological impacts**

303 During the lifetime of a deep metal mine, ore extraction and processing can  
304 release vast quantities of solid waste into the riverine environment (Bird *et al.*,  
305 2010). Even after mine abandonment, erosion of material from mine spoil and  
306 tailings can continue to introduce contaminated solid wastes into river  
307 channels and floodplains for many decades (Macklin *et al.*, 2003; Walling *et*  
308 *al.*, 2003; Miller *et al.*, 2004; Dennis *et al.*, 2009). These solid wastes can  
309 have a significant impact on the geochemistry of channel and floodplain  
310 sediments (e.g., Aleksander-Kwaterczak and Helios-Rybicka, 2009; Byrne *et*  
311 *al.*, 2010) and also physical and chemical dispersion patterns of toxic metals  
312 (e.g., Hudson-Edwards *et al.*, 1999b; Dennis *et al.*, 2009).

313

#### 314 3.1 Sediment geochemistry

315 Gross contamination of fluvial sediments both within the channel and on the  
316 floodplain has been reported in most metal mining regions of the world (**Table**  
317 **2**), with metal concentrations in sediments usually being several orders of  
318 magnitude greater than that in the water column (Macklin *et al.*, 2006). Metal  
319 concentrations are greatest in the fine sediment fraction and, in particular, in  
320 the clay-silt fraction (< 63 µm; Lewin and Macklin, 1987; Foster and  
321 Charlesworth, 1996; Stone and Droppo, 1994; Dennis *et al.*, 2003; Förstner,

322 2004). This reflects the higher surface area per unit mass of smaller particles,  
323 and the ion-exchange capacity of silt and clay-sized fractions (which include  
324 clay minerals, iron hydroxides, manganese oxides, and organic matter in  
325 various states of humification).

326

327 Metal speciation is essential to assess geochemical phases and the mobility  
328 of potentially toxic elements in contaminated sediments (Tokalioglu *et al.*,  
329 2003). Until fairly recently, most investigations of sediment metal  
330 concentrations have used strong chemicals (e.g., HNO<sub>3</sub>, HCl, HClO<sub>4</sub>, HF) to  
331 extract the total amount of metals in the sediment, often leading to  
332 oversimplified interpretations that do not take sediment complexity into  
333 account (Linge, 2008). Metals in sediments exist in various geochemical  
334 phases which reflect the degree to which they can be re-mobilised from the  
335 sediment. For this reason, chemical sequential extraction procedures (SEPs)  
336 capable of identifying contaminant partitioning have become increasingly  
337 popular over total dissolution of the sediment achieved by single extractions  
338 (e.g., Tessier *et al.* 1979; Rauret *et al.*, 1999). Various extraction media have  
339 been used to target specific geochemical phases, including electrolytes  
340 (CaCl<sub>2</sub>, MgCl<sub>2</sub>), pH buffers of weak acids (acetic, oxalic acid), chelating  
341 agents (EDTA, DTPA) and reducing agents (NH<sub>2</sub>OH). In many metal mining  
342 regions, the impact on sediment geochemistry has been to increase the  
343 proportion of toxic metals in the more mobile (bioavailable) geochemical  
344 phases. Studies have identified cadmium (Licheng and Guijiu, 1996; Morillo *et*  
345 *al.*, 2002; Vasile and Vladescu, 2010), copper (Jain, 2004), zinc (Morillo *et al.*,  
346 2002; Galan *et al.*, 2003; Aleksander-Kwaterczak and Helios-Rybicka, 2009;

347 Naji *et al.*, 2010) and lead (Byrne *et al.*, 2010) to be highly elevated in the  
348 acid-soluble phases. The largest proportion of metals is usually found in the  
349 reducible phase bound to Fe and Mn oxides (Macklin and Dowsett, 1989).  
350 Copper has been found to associate largely with organic matter in the  
351 oxidisable phase (Licheng and Guijiu, 1996).

352

353 Sequential extraction procedures have allowed the chemical mobility and  
354 toxicological risk posed by contaminated sediments to be established allowing  
355 resource managers to prioritise areas for remediation. However, a number of  
356 doubts concerning the accuracy of selective chemical extraction schemes  
357 have been expressed (Linge, 2008). Particular concerns are whether the  
358 chemical extractant may attack phases other than those expected; and  
359 whether liberated metals may become associated with another sediment  
360 phase rather than staying in solution (Burton, 2010). The multitude of  
361 extraction techniques and media used can also lead to great variability in  
362 results and, in some instances, limits the ability to make direct comparisons  
363 between studies. Since the early 1990s, molecular scale techniques to study  
364 elemental binding have become more popular. X-ray absorption spectroscopy  
365 (XAS) techniques such as X-ray absorption near edge structure (XANES) and  
366 extended X-ray absorption fine structure (EXAFS) have allowed analysis at  
367 the molecular level and direct evidence of surface composition and bonding  
368 characteristics of mining-derived sediments (Esbrí *et al.*, 2010; Van Damme *et al.*,  
369 *et al.*, 2010). By investigating metal speciation at the atomic level, it is possible  
370 to establish metal toxicity, mobility and bioavailability with far greater accuracy  
371 than can be achieved using chemical extraction methods. This allows



372 scientists and environmental managers to more accurately gauge the impact  
373 of toxic metals on ecosystems and human health.

374

### 375 3.2 Physical dispersion and downstream attenuation processes

376 The influx of large volumes of contaminated material into river systems can  
377 significantly alter local sediment transport and deposition and affect chemical  
378 processes that operate at and beneath the river-bed surface (Gilbert, 1917;  
379 Lewin *et al.*, 1977; Wood and Armitage, 1997). A number of reviews have  
380 historically considered the hydrogeomorphic response of riverine systems to  
381 mining activities (e.g., Miller, 1997; Macklin *et al.*, 2006) and as a result only  
382 limited coverage is provided here. Based on research in the UK, Lewin and  
383 Macklin (1987) suggested that disturbances of the river channel due to mining  
384 can be categorized as involving processes of '*passive dispersal*' and '*active*  
385 *transformation*'.

386

#### 387 Passive Dispersal

388 During *passive dispersal*, mine waste is transported from the mine site with no  
389 significant alteration of the prevailing sediment load of the river. Changes can  
390 occur in depositional environments, with slow flowing and deep pools being  
391 preferential sites for the deposition of contaminant-enriched fine sediment (<  
392 2000  $\mu\text{m}$ ). Transport of coarse sediment (> 2000  $\mu\text{m}$ ) may be limited to  
393 modest and high flow events. However, fine sediments may be transported  
394 under a range of different flows, including extended periods of base-flow. In-  
395 channel sediment contamination generally decreases downstream from the  
396 contaminant source at rates that vary between systems but which, in many

397 cases, are negatively exponential (Lewin and Macklin, 1987). This pattern is  
398 functionally related to the hydraulic sorting of sediment based on density and  
399 size of ore particles (e.g., galena is more dense than sphalerite and smaller  
400 grains travel less fast than coarser grains - Wolfenden and Lewin 1978);  
401 dilution by uncontaminated sediments (Marcus, 1987); hydrogeochemical  
402 reactions (Hudson-Edwards *et al.*, 1996); and biological uptake (Lewin and  
403 Macklin, 1987). In many cases, good fits between metal concentration and  
404 distance downstream of mining input can be achieved using regression  
405 analysis (Wolfenden and Lewin, 1977; Lewin and Macklin, 1987) or non-linear  
406 mixing models which incorporate clean and contaminated sediment sources  
407 within a river catchment (Marcus, 1987). However, these models are often  
408 specific to both the individual metal and the catchment for which they were  
409 developed (Dawson and Macklin, 1998; Miller, 1997). Movement of sediment  
410 can also occur in large-scale bed forms or 'slugs', which have been identified  
411 as associated with highs and lows in an otherwise downward trending metal  
412 concentration with distance downstream (Miller, 1997).

413

#### 414 Active Transformation

415 *Active transformation* occurs in association with a significant increase in the  
416 input of mining debris to the fluvial system. This may cause intrinsic  
417 thresholds to be exceeded and, consequently, lead to a local loss of hydraulic  
418 or geomorphological equilibrium that manifests itself in changes in channel  
419 character (Lewin and Macklin, 1987). The type, rate and magnitude of  
420 erosional and depositional processes can change (Miller, 1997). Channel  
421 aggradation may be associated with sediment inputs from active mining and

422 channel degradation may occur after mining has ceased (Gilbert, 1917;  
423 Knighton, 1991). Meandering channels may be transformed into braided  
424 forms (Warburton *et al.*, 2002). Other depositional features can include scroll  
425 bars that arise from rapid accretion of sequentially developing point bars as a  
426 response to high sediment loads and channel migration, and substantial  
427 overbank floodplain deposits, particularly where overbank splays lead to  
428 avulsion channels that cross the flood plain (Miller, 1997; Walling *et al.*, 2003;  
429 Dennis *et al.*, 2009).

430

431 Toxic metal contaminants can be extremely persistent within the environment  
432 and can remain stored within floodplain deposits for decades to millennia  
433 (Miller, 1997). Since the 1970's, a significant amount of research has  
434 focussed on the role of historical metal mining in the contamination of  
435 floodplains (**Table 2**). Analyses of floodplain overbank sediments in the River  
436 Ouse catchment in northeast England revealed contaminated sedimentary  
437 successions reflecting over 2000 years of lead and zinc mining (Hudson-  
438 Edwards *et al.*, 1999a). It has been estimated that over 55% of the  
439 agriculturally important River Swale floodplain, a tributary of the Ouse, is  
440 significantly contaminated by toxic metals (Brewer *et al.*, 2005). It has been  
441 estimated that approximately 28% of the lead produced in the Swale  
442 catchment remains within channel and floodplain sediments. At present rates  
443 of valley-bottom reworking through channel migration and erosion, it may take  
444 in excess of 5,000 years for all of the metal-rich sediment to be exported from  
445 the catchment (Dennis *et al.*, 2009). These studies indicate that large areas of  
446 agricultural land are potentially contaminated and that there may be long-term

447 health concerns for those ingesting contaminants via crops produced on this  
448 land (Albering *et al.*, 1999; Conesa *et al.*, 2010).

449

450 Recent advances in geochemical tracing techniques and numerical modelling  
451 have led to improved understanding and predictability of dispersal rates and  
452 patterns of sediment-associated toxic metal contamination. Owens *et al.*  
453 (1999) used geochemical fingerprinting to identify the proportion of sediment  
454 from mining areas in the River Ouse catchment, UK. Using isotope  
455 signatures, several studies have differentiated specific geographical sources  
456 in mining-affected catchments (Hudson-Edwards *et al.*, 1999a; Bird *et al.*,  
457 2010). Bird *et al.* (2010) were able to discriminate between sediments derived  
458 from mine waste and river sediments using lead isotope signatures. They  
459 surmised that approximately 30% of the sediment load of the lower River  
460 Danube was derived from mining. Numerical modelling techniques now allow  
461 the prediction of contamination patterns in river catchments now and in the  
462 future. For example, the catchment sediment model TRACER has been  
463 applied to identify sediment contamination 'hot spots' in the River Swale  
464 catchment, UK (Coulthard and Macklin, 2003). The model also revealed that  
465 over 200 years after the cessation of mining activities, over 70% of the  
466 deposited contaminants remain in the Swale catchment.

467

### 468 3.3 Chemical dispersion and attenuation processes

469 Chemical transportation processes in sediments of metal mining-affected  
470 rivers become increasingly important after the closure and abandonment of  
471 deep mines (Lewin and Macklin, 1987; Bradley *et al.*, 1995). Toxic metals can

472 be attenuated downstream of a mining input through pH buffering, acid  
473 neutralisation, and precipitation and adsorption reactions (Routh and  
474 Ikramuddin, 1996; Ford *et al.*, 1997; Lee *et al.*, 2002; Ren and Packman,  
475 2004). The often termed 'master variable' for determining metal speciation in  
476 aquatic systems is pH (Kelly, 1988; Younger *et al.*, 2002). As pH increases,  
477 aqueous metal species generally display an increasing tendency to precipitate  
478 as carbonate, oxide, hydroxide, phosphate, silicate or hydroxysulphate  
479 minerals (Salomons, 1993). The effects of increasing pH below mine  
480 discharges can be seen in some rivers by changes in precipitate mineralogy,  
481 with proximal capture by iron hydroxides and distal capture by aluminium  
482 oxides (e.g., Munk *et al.*, 2002). Therefore, a major control on metal  
483 attenuation, acid production and stream pH at abandoned mine sites is the  
484 amount of carbonate minerals present in the surrounding geology. Carbonate  
485 minerals such as calcite, dolomite and siderite weather quickly and can buffer  
486 pH and act as adsorption sites for dissolved toxic metals. Non-carbonate  
487 minerals weather slowly and, where they predominate, can be extremely slow  
488 to react to changes in pH (Wilkin, 2008). The precipitation of solid-form metals  
489 limits the concentration of metals which are transported through the aquatic  
490 system as free ion species (Enid Martinez and McBride, 1998). These  
491 secondary minerals can also act as sorbents for dissolved metals (Enid-  
492 Martinez and McBride, 1998; Asta *et al.*, 2007; Wilkin, 2008). Adsorption of  
493 metals usually increases at higher pH so that substantial changes in dissolved  
494 metal concentrations can occur with small changes in pH, typically over 1 –  
495 1.5 pH units (Salomons, 1993). Aside from pH, several other water quality  
496 parameters can influence metal speciation, including the concentration of the

497 metal, presence of ligands, redox conditions, salinity, hardness, and the  
498 presence of other metals (Novotny, 2003). High levels of salinity, hardness  
499 and organic matter content are known to increase metal attenuation by  
500 providing binding sites for metal sorption (Salomons, 1980; Dojlido and  
501 Taboryska, 1991; Achterberg *et al.*, 2003).

502

503 Under invariant environmental conditions, sediment geochemical phases are  
504 stable, chemical attenuation of metals will proceed at regular rates and, thus,  
505 metals remain immobile in river bed sediments (Morillo *et al.*, 2002). However,  
506 sediments are not a permanent sink for metals and they may be released into  
507 the water column when suitable conditions for dissolution occur. Several  
508 studies have reported the mobilisation of reduced sediment-bound metals to  
509 the water column under oxidising conditions, for example, during floods and  
510 dredging activities (Calmano *et al.*, 1993; Petersen *et al.*, 1997; Kuwabara *et al.*,  
511 *et al.*, 2000; Zoumis *et al.*, 2001; Butler, 2009; Knott *et al.*, 2009). In sediments  
512 from Hamburg harbour, Calmano *et al.* (1993) observed oxidation episodes to  
513 decrease pH in the suspended sediments from 7 to 3.4, leading to the  
514 mobilisation of zinc and cadmium. Similarly, oxidation of anoxic sediments  
515 from Mulde reservoir, Germany, resulted in the mobilisation of zinc and  
516 cadmium and redistribution of toxic metals to more bioavailable geochemical  
517 phases (Zoumis *et al.*, 2001). Mullinger (2004) reported diffuse discharges of  
518 metals from bed sediments accounted for up to 40% of zinc, cadmium and  
519 copper entering surface waters of the Cwm Rheidol mine, Wales. Bioturbation  
520 (Zoumis *et al.*, 2001) and changes in pH (Hermann and Neumann-Mahlkau,  
521 1985), dissolved organic carbon (Butler, 2009), ionic concentration (Dojlido

522 and Taboryska, 1991), and the concentration of complexing agents  
523 (Fergusson, 1990; Morillo *et al.*, 2002) have also been reported to lead to the  
524 release of 'stored' toxic metals into the wider environment.

525

526 The contamination risk posed by toxic metals stored in aquatic sediments of  
527 former and current industrial centres (including metal mining regions), and the  
528 potential for these toxic metals to contaminate areas beyond the source of  
529 contamination, has prompted many national regulatory authorities to introduce  
530 sediment environmental quality standards (SEQS) (e.g., Environment Agency,  
531 2008b) based on total metal concentrations in the sediment. The practical  
532 application of SEQS is made difficult by a number of factors relating to the  
533 nature of heavy metal pollutants, including variation in natural background  
534 concentrations, the existence of chemical species, the concentrations of  
535 physico-chemical parameters, variations in organism sensitivity, and the fact  
536 that some heavy metals are essential elements for organisms (Comber *et al.*,  
537 2008). In order to classify accurately the ecological status of rivers impacted  
538 by metal mining, sediment assessments may need to be unique to each river  
539 catchment and incorporate: background metal concentrations, an assessment  
540 of bioavailable fractions, and concurrent water quality measurements  
541 (including major ions) (Netzband *et al.* 2007; Brils 2008; Förstner 2009). As  
542 far as is known by the authors, most national monitoring and assessment  
543 programmes for freshwater systems measure total metal concentrations in  
544 sediments rather than the concentration of metals in different geochemical  
545 phases. Measurement of total quantities of metals in sediment provides little  
546 information regarding their ecotoxicity and their potential mobility. With the

547 achievement of Good Ecological Status (GES) at the centre of many  
548 environmental improvement programmes (e.g., to comply with the European  
549 Water framework Directive), it is argued that measurement of bioavailable  
550 metals in the sediment, which can interact relatively easily with aquatic  
551 organisms, would provide a more comprehensive and robust assessment of  
552 ecological risk. In this respect, there is a real risk that such programmes are  
553 failing to meet their own objectives.

554

#### 555 **4. Hydrological impacts**

556 The generation of mine water pollution is a product of many factors including  
557 local mineralogy, lithology, contaminant source area, and biogeochemical  
558 reactions (Younger *et al.*, 2002). The character of mine water pollution in  
559 surface waters is strongly influenced by a wide range of hydroclimatological  
560 factors (including rainfall characteristics), land use (both catchment-wide and  
561 any changes associated with spoil heaps), seasonality, antecedent conditions  
562 to rainfall or snow-melt (particularly soil and spoil moisture content but also  
563 temperature), dominant hydrological transport pathways, and stream  
564 discharge (Gammons *et al.*, 2005; Canovas *et al.*, 2008). Once released to  
565 the water column, metals can move through the aquatic environment,  
566 resulting in impaired water quality in reaches of a river or estuary that were  
567 unaffected directly by deep mine drainage. Released metals can also interact  
568 with aquatic animals, resulting in the deterioration of aquatic ecosystem health  
569 (Farag *et al.*, 1998).

570



571 Traditionally, discharge has been seen as a master variable driving river  
572 hydrochemistry (Bradley and Lewin, 1982). Heavy metal ion concentrations in  
573 rivers are generally thought to be greatest during low flows and lowest  
574 coinciding with high flows, when uncontaminated runoff dilutes solute  
575 concentrations (Webb and Walling, 1983). Since the 1970s, many  
576 researchers have documented the effects of seasonal variability in stream  
577 discharge on toxic metal concentrations (e.g., Grimshaw *et al.*, 1976; Keith *et*  
578 *al.*, 2001; Sullivan and Drewer, 2001; Nagorski *et al.*, 2003; Desbarats and  
579 Dirom, 2005; Hammarstrom *et al.*, 2005). Annual patterns (hysteresis  
580 patterns) of dissolved metal concentrations are apparent in many rivers,  
581 reflecting the flushing of oxidised sulphides accumulated over dry summer  
582 (low flow) months (Canovas *et al.*, 2008). Many researchers have noted  
583 maximum toxic metal concentrations as occurring during the first heavy rains  
584 of the hydrological year, during the autumn (Bradley and Lewin, 1982; Bird,  
585 1987; Boulton *et al.*, 1994; Braungardt *et al.*, 2003; Desbarats and Dirom, 2005;  
586 Olias *et al.*, 2004; Mighanetara *et al.*, 2009). Contaminant concentrations  
587 typically decrease in winter and increase gradually through spring and  
588 summer as a result of increased sulphide oxidation and evaporation.  
589 Therefore, the timing of maximum contaminant flux will be largely a function of  
590 hydroclimatology, catchment characteristics and the minerals present at a  
591 mine site.

592

593 It is understood that a major part of element transfer in rivers takes place  
594 during short episodes of high river flow, i.e. floods (Sanden *et al.*, 1997).

595 However, to date, very little research has been directed towards detailing toxic

596 metal fluxes and hydrochemical variability during individual high flow events in  
597 former mining regions (Grimshaw *et al.*, 1976; Bradley and Lewin, 1982;  
598 Sanden *et al.*, 1997; Dawson and Macklin, 1998; Lambing *et al.*, 1999; Wirt *et*  
599 *al.*, 1999; Keith *et al.*, 2001; Gammons *et al.*, 2005; Canovas *et al.*, 2008).  
600 One of the earliest studies by Grimshaw *et al.* (1976), on the River Ystwyth,  
601 Wales, observed hysteretic behaviour in the relation between metal  
602 concentrations and discharge, whereby metal concentrations increased on the  
603 rising limb of the flood hydrograph and decreased on the falling limb,  
604 associated with flushing and exhaustion (or dilution), respectively. This  
605 general pattern has also been reported in a number of more recent studies  
606 (e.g., Keith *et al.*, 2001; Canovas *et al.*, 2008; Byrne *et al.*, 2009). In some  
607 instances, the source of metals in the initial flush was metal sulphates  
608 accumulated on the surface of mine waste (Keith *et al.*, 2001) or  
609 contaminated groundwater efflux from mine portals (Canovas *et al.*, 2008).  
610 Metal attenuation on the falling limb is principally due to rain-water dilution  
611 and the fact that the available contaminant are scavenged in the first flush  
612 (Canovas *et al.*, 2008). The frequent occurrence of peak iron, manganese and  
613 aluminium concentrations on the falling limb of the hydrograph indicates that  
614 adsorption onto, or precipitation with, iron solids may be an important toxic  
615 metal attenuation mechanism during stormflow events in some rivers (Lee *et*  
616 *al.*, 2002; Asta *et al.*, 2007; Byrne *et al.*, 2009).  
617  
618 The mobilisation and transport of mine wastes during stormflows and the  
619 consequent contamination of agricultural lands is an important issue for  
620 environmental managers of former metal mining regions (Dennis *et al.*, 2003;

621 Connelly, 2009). During the 1990s, there was a marked increased interest in  
622 toxic metal contamination in floodplains in the UK following a number of  
623 devastating floods and an increased focus on the potential effects of climate  
624 change on hydrological regimes and sediment transport dynamics (**Table 2**).  
625 The autumn and winter floods of 2000-2001 across a substantial part of  
626 Europe caused large-scale remobilisation and deposition of contaminated  
627 sediments in floodplains and farm-land (Dennis *et al.*, 2003; Macklin *et al.*,  
628 2003; Macklin *et al.*, 2006). In future, predicted increases in the frequency and  
629 magnitude of floods as a function of climate change may result in increased  
630 mobilisation and deposition of toxic metals in floodplains across Europe  
631 (Macklin *et al.*, 2006; Environment Agency, 2008b; Förstner and Salomons,  
632 2008). Therefore, there is a need to monitor and assess stormflow events and  
633 river hydrochemistry in detail in order to quantify metal fluxes with reasonable  
634 levels of accuracy in order to allow environmental managers to prioritise areas  
635 for remediation.

636

637 Aside from contamination of floodplains, the large-scale movement of mine  
638 waste during stormflow events has significance for aquatic ecosystem health.  
639 The highly elevated toxic metal concentrations during stormflows undoubtedly  
640 cause harm to aquatic communities and degrade biological quality (Wolz *et*  
641 *al.*, 2009). The long-term effects of these transient conditions can be  
642 established through investigations of aquatic ecosystem health. However, the  
643 added or individual impact of stormflow events is still largely unknown due to  
644 the difficulty of measuring it. Predicted increases in the frequency and  
645 magnitude of floods across Europe due to climate change (Wilby *et al.*, 2006)

646 have put an emphasis on bridging the knowledge gap between the physical  
647 remobilisation of contaminants during stormflows and the potential  
648 toxicological impacts (Wolz *et al.*, 2009). Understanding the toxicological  
649 impacts of stormflows will be important in the achievement of environmental  
650 quality standards in mining-affected river catchments.

651

652 Most metal mines are associated with significant volumes of waste material  
653 deposited as surface spoil heaps and tailings. The hydrological behaviour of  
654 these waste deposits can be significantly different to the wider catchment due  
655 to the alteration of local surface and sub-surface flow pathways (Younger *et*  
656 *al.*, 2002). Considering the important role of spoil material in the production of  
657 metal contaminants, comparatively little research has been undertaken into  
658 flow pathways and contaminant generating processes within mine spoil. Due  
659 to the artificial stratification and the discontinuities in permeability that occur  
660 within spoil heaps, they often have 'perched aquifers' that lie well above the  
661 underlying bedrock, producing unique flow paths (Younger *et al.*, 2002). The  
662 development of a water table in mine spoil depends on the predominant  
663 lithology of the spoil. For example, sandstone generally forms highly  
664 permeable spoil whereas mudstone produces spoil of low permeability. Highly  
665 permeable spoil can contain as much as 25% or more of ore as fines or  
666 solutes (Davies and Thornton, 1983). Where rainfall infiltration-excess is  
667 typical, because, for instance, fine-grained material produces a surface seal,  
668 surface runoff will be the predominant flow path (Younger *et al.*, 2002). This  
669 will, through gully erosion, transfer large quantities of contaminated solids into  
670 the local water course.

671

672 Changes in flow paths and direction within mine spoil can occur slowly  
673 through the seasons or more rapidly during rainfall events as different flow  
674 paths become active with the fluctuation of perched water tables (Walling and  
675 Webb, 1980). Differential hydrology can induce variability in toxic metal  
676 speciation in mine spoils and tailings (Kovacs *et al.*, 2006). Generally,  
677 oxidation of sulphide minerals occurs in a shallow oxidation zone near the  
678 surface of the spoil (Jurjovec *et al.*, 2002). Dissolution and flushing of these  
679 oxidised metals can then occur during wet periods (Navarro *et al.*, 2008).  
680 Several studies of metal flushing during storms have reported the importance  
681 of weathered metal salts on and near the surface of mine spoil as responsible  
682 for increasing metal concentrations during runoff (Canovas *et al.*, 2008; Byrne  
683 *et al.*, 2009). Below the oxidation zone, a zone of transition from saturated to  
684 unsaturated sediments typically occurs, often characterised by a 'hard pan' of  
685 metal precipitates (Romero *et al.*, 2007). Toxic metals can be attenuated in  
686 the mine spoil through a series of precipitation, co-precipitation and  
687 adsorption reactions. Reducing conditions in saturated sediments can lead to  
688 the formation of insoluble metal sulphides. pH buffering can occur in the  
689 shallow oxidizing zone with secondary-phase precipitation occurring near the  
690 deeper saturated zone (McGregor *et al.*, 1998). In order to effectively plan for  
691 mine site remediation, it is essential that mine spoils and tailings are  
692 characterised in terms of mineralogy, metal speciation and hydrology,  
693 especially where contamination of groundwater is an issue. Such information  
694 is necessary to understand the mechanisms controlling the release and  
695 attenuation of metals at these sites.

696

697 **5. Ecological impacts of metal mine contamination on macroinvertebrate**  
698 **communities**

699 As early as the 1960s, the adverse impacts of mining activities on  
700 macroinvertebrates were being acknowledged (Reish and Gerlinger, 1964).  
701 Metal mine drainage can severely impact aquatic ecosystems by affecting  
702 primary and secondary production, nutrient cycling, energy flow and  
703 decomposition (Stoertz *et al.*, 2002; Knott *et al.*, 2009; Younger and  
704 Wolkersdorfer, 2004; Batty *et al.*, 2010). Freshwater macroinvertebrates fulfil  
705 important roles in the river ecosystem, being vital food sources for many  
706 aquatic and terrestrial predators and playing a significant part in the cycling of  
707 organic matter and nutrients (Gerhardt, 1993). The pivotal position of benthic  
708 macroinvertebrates in aquatic food webs means that negative impacts on  
709 them can have widespread consequences within aquatic and terrestrial food-  
710 webs for primary producers, predators and the wider ecosystem. As a result,  
711 macroinvertebrates have increasingly been used as indicators of stream  
712 ecosystem health associated with metal mining (e.g., Batty *et al.*, 2010;  
713 Poulton *et al.*, 2010).

714

715 **5.1 Changes in community composition**

716 A wide range of changes to macroinvertebrate community structure and  
717 composition have been reported in the scientific literature associated with  
718 metal mining activities. Reductions in abundance, number of taxa and  
719 biodiversity are common impacts reported in association with metal mining-  
720 activities internationally (e.g., Willis, 1985; Gray, 1998; Amisah and Cowx,

721 2000; Watanabe *et al.*, 2000; Hirst *et al.*, 2002; Kiffney and Clements, 2003)  
722 (**Table 3**). Investigations have generally revealed that some  
723 macroinvertebrate taxa display a tolerance or sensitivity to contamination  
724 (**Table 3**). Whilst investigating contaminated stretches of two rivers in Ohio,  
725 USA, Winner *et al.* (1980) hypothesised that habitats heavily polluted with  
726 toxic metals may be dominated by Chironomidae (Diptera – true fly larvae);  
727 moderately polluted habitats by Chironomidae and Trichoptera (caddisfly);  
728 and minimally or unpolluted habitats by caddisflies and Ephemeroptera  
729 (mayfly). Armitage *et al.* (1980; 2007) examined macroinvertebrate species  
730 composition of the mining impacted River Nent. Diptera and Plecoptera  
731 (stonefly) were the dominant orders observed in the river system. Trichoptera  
732 and mayfly (Ephemeroptera) were not abundant and seemed particularly  
733 sensitive to the mine water pollution. In contaminated reaches of the River  
734 Vasco, Portugal, the number of predators increased and the number of EPT  
735 taxa (Ephemeroptera – Plecoptera - Trichoptera) decreased, probably  
736 reflecting the presence of thick layers of metal hydroxides on the river  
737 substrate (Gerhardt *et al.*, 2004). Sites subject to severe AMD contamination  
738 showed high levels of biodiversity due to high species richness of the tolerant  
739 species. In general, the order of toxicity of metal mine contamination to the  
740 most common macroinvertebrate orders is: Ephemeroptera > Trichoptera >  
741 Plecoptera > Diptera. However, there can be considerable variability in metal  
742 tolerance between macroinvertebrate taxa and species. For example,  
743 Ephemeroptera are generally considered to be highly sensitive to metal  
744 contamination despite some species (e.g., *Baetis rhodani* and *Caenis cf.*  
745 *luctuosa*) being reported to display some tolerance to metal contaminants

746 (Roline, 1988; Beltman *et al.*, 1999; Gower *et al.*, 1994; Gerhardt *et al.*, 2004;  
747 Gerhardt *et al.*, 2005b). Several authors have reported impacts of mine water  
748 contamination on ecosystem function (**Table 3**), including reduced secondary  
749 production (Carlisle and Clements, 2005; Woodcock and Huryn, 2007), and a  
750 reduction in leaf matter (detritus) decomposition rates and microbial  
751 respiration (Kiffney and Clements, 2003; Carlisle and Clements, 2005).

752

753 Relatively predictable changes in macroinvertebrate community structure as a  
754 result of pollution (e.g., decreased abundance and biodiversity, elimination of  
755 sensitive taxa) have led to the development of a number of biotic and diversity  
756 indices (e.g., Shannon, 1948; Berger and Parker, 1970). However, the  
757 performance of biological indices / metrics appear to vary widely when applied  
758 to mine water contaminated sites (Smolders *et al.*, 2003; Van Damme *et al.*,  
759 2008; Chadwick and Canton, 1984; Willis, 1985; Chadwick *et al.*, 1986; Rhea  
760 *et al.*, 2006). Variability in success is likely to be a function of the complicated  
761 interplay between the mine water components, other water quality  
762 parameters, and natural tolerances and sensitivities of organisms. Gray and  
763 Delaney (2008) suggest a modification of the Acid Waters Indicator  
764 Community (AWIC) index (Davy-Bowker *et al.*, 2005) to incorporate metal  
765 toxicity may be required. However, such a revision would also need to  
766 address the pH bias in the calibration data and the (possibly) inaccurate  
767 grouping of macroinvertebrates in sensitivity groups. A revision of the  
768 Biological Monitoring Working Party (BMWP) system (Biological Monitoring  
769 Working Party, 1978), based on species' tolerance to acidity and metal  
770 contamination, has also been suggested (Gray and Delaney, 2008) and some



771 success has been achieved using a multi-metric approach by considering  
772 multiple biological metrics simultaneously (e.g., Clews and Ormerod, 2009).  
773 Clearly, there is scope for a biological index designed specifically for detecting  
774 the impacts of mine water contamination on aquatic communities. However,  
775 such an index would need to incorporate the effects on a community of  
776 multiple environmental stressors, the most important of which are probably  
777 dissolved metals and acidity.

778

## 779 5.2 Changes in macroinvertebrate physiology and behaviour

780 More subtle community alterations as a result of physiological or behavioural  
781 changes are less easy to diagnose (Younger and Wolkersdorfer, 2004) (**Table**  
782 **3**). For example, Petersen and Petersen (1983) reported anomalies in the  
783 construction of filter feeding nets of Hydropsychidae (Trichoptera) in rivers  
784 affected by a gradient of toxic metal pollution. Disruption of silk-spinning by  
785 contamination caused the caddisfly to spend more time in open habitats  
786 repairing the structure and thus more vulnerable to potential predators. Vuori  
787 (1994) observed metal exposure to affect the territorial behaviour of  
788 Hydropsychidae, relaxing levels of interspecific competition and increasing  
789 susceptibility to predation. Brinkman and Johnston (2008) reported decreased  
790 moulting rates (*Rhithrogena hageni*: Ephemeroptera) after exposure to high  
791 levels of copper, cadmium and zinc. In an experimental stream study,  
792 Clements *et al.* (1989) reported that high copper doses increased predation  
793 pressure, so much that the numbers of caddisfly, mayfly and chironomids  
794 were dramatically reduced. Maltby and Naylor (1990) found high zinc  
795 concentrations significantly impacted *Gammarus pulex* reproduction by  
796 causing a reduction in energy absorption and an increase in the number of

797 broods aborted. Other behavioural responses reported associated with metal  
798 mine contamination include increased drift rates, physical avoidance of  
799 contaminated sediments, reduced burrowing / burial rates (Leland *et al.*, 1989;  
800 Roper *et al.*, 1995) and reduced leaf litter processing rates and microbial  
801 respiration (Kiffney and Clements, 2003; Carlisle and Clements, 2005). Many  
802 of the species specific differences reported within the literature have been  
803 attributed to trophic status with herbivores and detritivores typically being  
804 more sensitive to contamination than predators (Leland *et al.*, 1989;  
805 Schultheis *et al.*, 1997; Gerhardt *et al.*, 2004; Poulton *et al.*, 2010). Acute  
806 metal contamination can induce deformities and mutations of head and  
807 feeding structure in macroinvertebrate fauna (e.g., Groenendijk *et al.*, 1998;  
808 Vermeulen *et al.*, 2000; Groenendijk *et al.*, 2002; De Bisthoven *et al.*, 2005).  
809 Both zinc and lead have been implicated as teratogens (inducing deformities  
810 as a result of chronic exposure during the lifetime of the organism) and as a  
811 mutagen (inducing deformities in offspring due to DNA damage in parents  
812 from chronic exposure) in *Chironomus riparius* (Chironomidae) (Martinez *et*  
813 *al.*, 2004).

814

815 More recent studies have made use of biomonitoring techniques which are  
816 capable of detecting sublethal behavioural and physiological responses in an  
817 organism when exposed to a contaminant (e.g., De Bisthoven *et al.*, 2004;  
818 Gerhardt *et al.*, 2004; Gerhardt *et al.*, 2005a; De Bisthoven *et al.*, 2006;  
819 Gerhardt, 2007; Macedo-Sousa *et al.*, 2007) (**Table 3**). A conceptual  
820 Stepwise Stress Model (SSM), proposed by Gerhardt *et al.* (2005a),  
821 postulates that an organism will display a time-dependent sequence of

822 different regulatory and behavioural responses during exposure to  
823 contaminants over a certain threshold. Several species have been found to  
824 show a pH-dependent response to AMD involving, first, an increase in  
825 locomotion, followed by an increase in ventilation (e.g., Gerhardt *et al.*, 2005a;  
826 De Bisthoven *et al.*, 2006). An increased ventilation rate reflects changes in  
827 the organism's respiratory and physiological system, and may be due to  
828 damage to gill membranes or nerve tissues. Locomotory activity probably  
829 represents an avoidance strategy from potentially toxic conditions.  
830 Importantly, biomonitoring methods integrate biochemical and physiological  
831 processes and so are a more comprehensive method than single biochemical  
832 or physiological parameters. In combination with the Stepwise Stress Model,  
833 online biomonitoring offers the possibility of a graduated 'early warning'  
834 system for the detection of pollution waves (Gerhardt *et al.*, 2005a).

835

### 836 5.3 Metal bioaccumulation in macroinvertebrates

837 A significant body of research has concentrated on evaluating the  
838 bioaccumulation of toxic metals in macroinvertebrates as a measure of the  
839 bioavailability of contaminants (e.g., Farag *et al.*, 1998; Smolders *et al.*, 2003;  
840 Yi *et al.*, 2008). Metals which are bioaccumulated by organisms and plants  
841 can be concentrated or magnified in the food chain (Sola *et al.*, 2004) (**Table**  
842 **3**). Benthic primary producers and decomposers are known to accumulate  
843 significant amounts of metals with little or no deleterious effects (Farag *et al.*,  
844 1998; Sanchez *et al.*, 1998). These metals can be transferred to herbivorous  
845 and detritivorous macroinvertebrates which in turn can transfer the metals to  
846 higher trophic levels (Younger and Wolkersdorfer, 2004). Metal accumulation

847 can vary between species, depending on a great number of physiological (e.g.  
848 cuticle type, the presence or absence of external plate gills, the processes  
849 which control metal distribution in the cell) and behavioural factors such as an  
850 organisms feeding strategy, contact with benthic sediments, larval stage and  
851 size (Dressing *et al.*, 1982; Farag *et al.*, 1998; Goodyear and McNeill, 1999;  
852 Sola and Prat, 2006; Cid *et al.*, 2010). Metal intake can take place through  
853 direct exposure to metals in surface and pore waters or indirectly via food  
854 supply. Those metals which, through their chemistry, are almost completely  
855 sediment-bound (Fe, Mn, Pb, Al), will usually be most important for particle  
856 feeders. Metal intake in the tissue takes place at a cell membrane, typically in  
857 the gill or gut, depending on whether the metal is in solution in the  
858 surrounding water body or if it was ingested with food. A range of  
859 environmental factors determine the potential for metal bioaccumulation  
860 including metal concentration in the surrounding water, water hardness,  
861 presence of organic matter, feeding group and the ionic state of the metal  
862 (Gower and Darlington, 1990; Farag *et al.*, 1998; Sola and Prat, 2006). The  
863 accumulation of metals in different organisms can also vary greatly as a result  
864 of natural or evolved tolerance mechanisms (Spehar *et al.*, 1978; Gower and  
865 Darlington, 1990; Bahrndorff *et al.*, 2006). For example, *Plectrocnemia*  
866 *conspersa* (Trichoptera), common in streams in south-west England affected  
867 by metal mine drainage were found to be tolerant of copper pollution (Gower  
868 and Darlington 1990). Some controlled microcosm experiments have reported  
869 tolerance to metal polluted sediments by *Chironomus februaris*  
870 (Chironomidae) (Bahrndorff *et al.* 2006). Mechanisms of tolerance might be  
871 methylation, increased metal excretion or decreased metallothionein

872 production. Metallothionein is a metal-binding protein with the principal  
873 function of accumulating essential metals for normal metabolic processes  
874 (Howard, 1998). Its presence facilitates the accumulation of toxic metals,  
875 however decreased production of this protein may allow certain organisms to  
876 accumulate lower amounts of toxic metals. Despite the great range of factors  
877 which can affect metal bioaccumulation in organisms, bioaccumulation factors  
878 (BAFs) which consider tissue metal concentration in relation to the  
879 surrounding abiotic medium, are possibly a more robust biodiagnostic method  
880 than measurement of metal concentrations in the water column and benthic  
881 sediments. If water quality guidelines are to continue to be used, then  
882 additional research will need to be undertaken to determine appropriate  
883 guidelines (possibly above existing guidelines) to support aquatic  
884 communities. In the future, metal bioaccumulation will need to be studied in a  
885 greater range of macroinvertebrates in order to fully understand metal-  
886 organism interactions in aquatic systems. A review of metal bioaccumulation  
887 studies by Goodyear and McNeill (1999) found that most studies primarily  
888 considered Ephemeropteran and Dipteran taxa and especially collector-  
889 gatherer and predatory functional feeding groups / traits.

890

#### 891 5.4 Effects of environmental parameters on the toxicity of mine discharges

892 Changes in some environmental parameters can affect the chemistry and,  
893 therefore, the toxicity of metals to organisms. The effects of salinity, water  
894 hardness and alkalinity on metal toxicity have been studied extensively (Stiff,  
895 1971; Brkovic-Popovic and Popovic, 1977; Gauss *et al.*, 1985; Gower *et al.*,  
896 1994; Yim *et al.*, 2006, Riba *et al.*, 2010 – **Table 3**). All of these studies

897 reported metal toxicity increases for macroinvertebrate and fish species under  
898 low salinity, alkalinity and water hardness conditions. Increased metal toxicity  
899 has also been reported at low turbidity (Garcia-Garcia and Nandini, 2006) and  
900 DOM (dissolved organic material) levels (Gower *et al.*, 1994). In river  
901 systems, carbonate minerals, clay minerals and DOM act as sorption sites for  
902 toxic metals and, therefore, high levels of these parameters help to reduce the  
903 concentration of dissolved toxic metals in bioavailable forms. However,  
904 bottom-dwelling organisms will take up sediment-bound metals through  
905 ingestion.

906

907 While bioassay and microcosm studies have revealed much information on  
908 metal ecotoxicity, a possible criticism of them could be that they are too  
909 simplistic in seeking to evaluate the response of macroinvertebrate species or  
910 communities to a single metal contaminant. In reality, most contaminated  
911 mine waters will contain mixtures of different metals in solution (**Table 3**). The  
912 simplest solution has been to assume the toxic effects of the metals present in  
913 the mixture are additive (Vermeulen, 1995). However, the interaction between  
914 metals can result in synergistic effects. For example, Hickey and Golding  
915 (2002) reported total abundance of heptageniid mayflies, community  
916 respiration and macroinvertebrate drift were most sensitive to solutions with a  
917 mixture of zinc and copper. Clements (2004), in stream mesocosms, found  
918 negative responses were generally greatest with zinc alone or with zinc and  
919 cadmium. A possible explanation for this synergism is the physiological  
920 inhibition of metal excretion by one of the metals, allowing the other metal(s)  
921 to have greater toxic effects (Berninger and Pennanen, 1995). Mixtures of

922 metals have also been shown to have antagonistic effects. Morley *et al.*  
923 (2002) found zinc and cadmium to have an antagonistic effect leading to  
924 increased survival of the cercarial stage of the parasitic fluke *Diplostomum*  
925 *spathaceum*. In some cases, antagonistic effects of metal mixtures are  
926 probably related to competition between metal ions for common sites of  
927 uptake (Younger and Wolkersdorfer, 2004). A study by Vermeulen (1995)  
928 illustrated the difficulty in predicting how metal mixtures will affect metal  
929 toxicity to organisms. Out of the 26 studies analysed, thirteen reported  
930 synergistic effects, six reported antagonistic effects, and seven reported  
931 additive effects. The problem of metal mixture toxicity is further compounded  
932 by other water quality parameters such as hardness, salinity and organic  
933 matter content. These parameters can increase or decrease metal toxicity and  
934 comparable mixtures of metals can also show contrasting toxicity effects  
935 between different groups, species and populations of organisms (Younger  
936 and Wolkersdorfer, 2004).

937

938 The task of evaluating metal toxicity is made even more difficult when acidity  
939 is considered. Most commonly, a decrease in pH will increase the amount of  
940 toxic free metal ions due to changes in metal speciation, mobility and  
941 bioavailability (Campbell and Stokes, 1985). However, at low pH, metals tend  
942 to desorb from organisms due to competition with hydrogen ions for binding  
943 sites (Gerhardt, 1993). The effects of low pH on stream biota in the absence  
944 of dissolved metals can be lethal or sublethal, inducing a range of  
945 physiological changes including an upset of the ionic balance across  
946 organism membranes and hydrolysing of cellular components (Kelly, 1988).

947 Campbell and Stokes (1985) suggested acidity can affect metal-organism  
948 interactions in two key ways. First, if a decrease in pH causes little change in  
949 metal speciation and there is only weak binding of metals at biological  
950 surfaces, the decrease in pH will decrease the toxicity of the metal due to  
951 competition with hydrogen ions for binding sites. Second, if a decrease in pH  
952 causes changes in speciation and there is strong binding at biological  
953 surfaces, then acidification will increase metal availability and toxicity. In the  
954 first instance, acidity will be the primary threat to ecosystems. In the second  
955 scenario, low pH and high dissolved metals may both influence toxicity.

956

957 The multi-factor nature of contaminated mine discharges (acidity, dissolved  
958 metals, metal precipitates, sulphates) and the natural variability in water  
959 chemistry between regions means that metal toxicity can be highly variable.  
960 Historically, ambient water quality criteria have specified permissible total or  
961 dissolved metal concentrations even though metal toxicity is heavily  
962 dependent on water chemistry (e.g., hardness, pH, DOM). The Biotic Ligand  
963 Model (BLM) (Di Toro *et al.*, 2001) was developed to predict metal toxicity by  
964 incorporating basic principles of physiology and toxicology, and the effects of  
965 water chemistry on metal speciation and bioavailability. The model has gained  
966 widespread use amongst the scientific / academic and water industry  
967 communities due to its potential for identifying water quality criteria and in  
968 facilitating risk assessment of aquatic environments (Paquin *et al.*, 2002). In  
969 order to gain wider applicability and relevance, BLMs will need to be applied  
970 to a wider range of organisms and pollutants in the future, and to be able to  
971 incorporate metal mixtures into toxicity predictions (Niyogi and Wood, 2004).



972

973 **6. Remediation of mining-impacted river systems**

974 The prevention of contaminated discharge from mine sites is now required by  
975 law in many countries (Macklin *et al.*, 2006). In the USA, the Clean Water Act  
976 (1972) was established to minimise the impact of anthropogenic pressures  
977 (including mining) on surface waters. In Europe, the adoption of the Water  
978 Framework Directive (2000/60/EC), and subsequent Mining Waste Directive  
979 (2006/21/EC), has necessitated the development of inventories of  
980 contaminant impacts at active and abandoned mine sites (Hering *et al.*, 2010).  
981 New legislation, based on a greater understanding of water quality and  
982 ecological integrity issues arising from mine discharges, have prompted  
983 research into remediation technologies aimed at reducing the environmental  
984 impact of metal mines (PIRAMID Consortium, 2003).

985

986 Mine water remediation technologies can be broadly categorised into active  
987 and passive treatment. Active treatment technologies are well established and  
988 involve the utilisation of electrical energy and mechanised procedures (Jarvis  
989 *et al.*, 2006) and are dependent on continuous monitoring and maintenance  
990 (Robb and Robinson, 1995). Traditional active treatment processes involve a  
991 sequence of oxidation by physical or chemical means, the addition of alkaline  
992 chemicals to raise pH and accelerate oxidation and precipitation of metals  
993 (Robb and Robinson, 1995; Lund and McCullough, 2009), and settlement and  
994 filtration (PIRAMID Consortium, 2003). However, active treatment incurs  
995 substantial set-up, material and maintenance costs (PIRAMID Consortium,  
996 2003). In response, passive remediation utilising natural physical, chemical

997 and biological processes and materials has found increasing favour over the  
998 past 30 years (Geroni *et al.*, 2009). Passive remediation systems use  
999 naturally available energy (e.g., topographical gradient, metabolic energy,  
1000 photosynthesis) to drive the remediative processes and have the principal  
1001 advantages over active remediation of reduced set up and maintenance costs  
1002 (Pulles and Heath, 2009). Some passive systems (e.g., wetlands) require  
1003 significantly greater land area than active treatment systems; although they do  
1004 not require costly reagents and incur less operational maintenance (Norton,  
1005 1992; Hedin *et al.*, 1994). Detailed characterisation of contaminant loading  
1006 over a sufficiently long time period is required prior to implementation of  
1007 treatment systems, including measurements of seasonal variation and the  
1008 impact of episodic contaminant flushing events, e.g., associated with spate  
1009 flows (Younger *et al.*, 2005; Byrne *et al.*, 2009). Equally important is the  
1010 linking of all mine water sources with a treatment system. Many abandoned  
1011 mine sites have substantial diffuse sources (Pirrie *et al.*, 2003; Mayes *et al.*,  
1012 2008; Mighanetara *et al.*, 2009; Byrne *et al.*, 2010), including mine spoil and  
1013 mobile metal fractions in the river bed. As a result it may be difficult to collect  
1014 and route contaminated runoff to treatment areas.

1015

1016 Mine water treatment technologies have been extensively reviewed elsewhere  
1017 (e.g., Brown *et al.*, 2002; Younger *et al.*, 2002; PIRAMID Consortium, 2003;  
1018 Lottermoser, 2007) and so a brief overview is provided (**Table 4**). Both  
1019 wetlands and Reducing and Alkalinity Producing Systems (RAPS) are now  
1020 well established remediation technologies throughout North America (e.g.,  
1021 Hedin *et al.*, 1994) and Europe (e.g., Whitehead and Prior, 2005) as passive

1022 treatment options for sulphate and Fe-rich, net-alkaline and net-acidic coal  
1023 mine discharges (Batty and Younger, 2004). In anoxic systems, removal of  
1024 toxic metals (e.g., zinc, lead, copper, cadmium) is hypothesised to occur  
1025 through the formation of insoluble metal sulphides and carbonates (Younger  
1026 *et al.*, 2002 – See **Table 4**). In aerobic systems, some toxic metals can be  
1027 removed either by direct precipitation as oxides and hydroxides or carbonate  
1028 phases or by co-precipitation with iron, manganese and aluminium  
1029 hydroxides. However, rates of toxic metal removal in these systems  
1030 (particularly zinc) have, in general, proved insufficient in circum-neutral and  
1031 net-alkaline mine waters, where chalcophile metals are the principal  
1032 contaminants (Robb and Robinson, 1995; Nuttall and Younger, 2000). Some  
1033 success has been achieved using variations of conventional calcite and  
1034 organic-based treatment systems in laboratory-scale experiments (Nuttall and  
1035 Younger, 2000; Rotting *et al.*, 2007; Mayes *et al.*, 2009). A large number of  
1036 researchers have also demonstrated the potential for organic and inorganic  
1037 sorbent media to remove toxic metals (Cui *et al.*, 2006; Perkins *et al.*, 2006;  
1038 Madzivire *et al.*, 2009; Mayes *et al.*, 2009; Rieuwertz *et al.*, 2009; Koukouzas  
1039 *et al.*, 2010; Vinod *et al.*, 2010). However, many of these technologies are still  
1040 at the experimental stage and will require further refinement and large-scale  
1041 field pilot studies before their full potential is realised. Frequent blocking of  
1042 filtering media with metal precipitates and rapid consumption of reactive  
1043 surfaces limit the metal removal efficiency of many of these systems to very  
1044 short time scales – hours to days in some instances (Younger *et al.*, 2002).  
1045

1046 Even with mine water treatment, the legacy of contamination in river  
1047 sediments and floodplains will represent a significant secondary diffuse  
1048 source of pollution long after other water quality parameters have improved to  
1049 acceptable levels. Therefore, contaminated sediments of mining-affected  
1050 rivers will continue to pose a serious threat to ecological integrity and the  
1051 achievement of Good Chemical Status (GCS) and Good Ecological Status  
1052 (GES) under the EU Water Framework Directive. The historical, preferred  
1053 method of dealing with contaminated sediment is removal by dredging (Nayar  
1054 *et al.*, 2004). This is an expensive and destructive process which may  
1055 mobilise vast reservoirs of bioavailable metals as part of the process (Nayar  
1056 *et al.*, 2004; Knott *et al.*, 2009). Furthermore, the sediment removed still  
1057 requires treatment and safe disposal. Recently, geochemical engineering  
1058 approaches involving in-situ and ex-situ biological and chemical treatment of  
1059 contaminated soils and sediments have gained attention as alternatives  
1060 (Förstner, 2004), and some success has been achieved in the stabilisation  
1061 and removal of toxic metals (Guangwei *et al.*, 2009; Luoping *et al.*, 2009;  
1062 Scanferla *et al.*, 2009). However, the principal necessity for the protection of  
1063 sediment and aquatic systems is considered to be the development of  
1064 guidelines concerning sediment quality (Burton, 2010; Byrne *et al.*, 2010).

1065

1066 Some efforts have focussed on the prevention of the generation of  
1067 contaminated mine water, so-called source control techniques. Conventional  
1068 techniques have focussed on physical and chemical stabilisation (Mendez  
1069 and Maier, 2008). Physical stabilisation involves covering mine waste with  
1070 inert material (e.g., clay, gravel) to reduce oxygen inflow and water ingress

1071 into the contaminated material (Gandy and Younger, 2003; Waygood and  
1072 Ferriera, 2009). However, clay caps in arid and semi-arid regions have tended  
1073 to crack from wetting and drying cycles resulting in the failure of the air-tight  
1074 cap (Newson and Fahey, 2003). Chemical stabilisation is achieved by adding  
1075 a resinous adhesive to form a crust over the mine waste, however, these also  
1076 are prone to cracking and failure (Tordoff *et al.*, 2000). More recently,  
1077 phytoremediation (phytoextraction and phytostabilisation) techniques have  
1078 developed as less costly alternatives (Marques *et al.*, 2009).

1079 Phytostabilisation creates a vegetative cap on the mine waste which  
1080 immobilises metals by adsorption and accumulation in the rhizosphere  
1081 (Mendez and Maier, 2008). Some success has been achieved in laboratory  
1082 trials investigating reforestation of mine tailings using endemic tree species  
1083 (Pollmann *et al.*, 2009). Phytoextraction offers the possibility of recovery of  
1084 metals through the hyperaccumulation of metals in plant tissues (Ernst, 2005).  
1085 However, the long-term performance of these new strategies needs to be  
1086 evaluated, as does the bioavailability of metals to wildlife which may feed on  
1087 the vegetative covers.

1088

## 1089 **7. Synthesis and conclusions**

1090 This paper provides a critical synthesis of scientific literature related to the  
1091 sedimentological, hydrological and ecological impacts of metal mining on  
1092 aquatic ecosystems. It has also highlighted the potential for remediation of  
1093 mine sites and provided an overview of current research and technological  
1094 developments in this area.

1095

1096 The important role of sediments in the dispersal, storage and recycling of  
1097 metal contaminants within the fluvial environment has been highlighted.  
1098 Significant quantities of contaminated sediment are eroded and transported  
1099 into aquatic systems from abandoned metal mines and both physical and  
1100 chemical processes influence the distribution of toxic metals within riverine  
1101 ecosystems. Physical dispersal processes are generally well understood and  
1102 can be classified as passive or active (Lewin and Macklin, 1987), the latter  
1103 prevailing when the addition of mine wastes to a river system results in a  
1104 threshold crossing event and the collapse of geomorphological equilibrium.  
1105 Under these circumstances, significant contamination of floodplains by toxic  
1106 metals can occur, with long-term potential consequences for the environment,  
1107 society and human health. However, recent advances in geochemical tracing  
1108 techniques and numerical modelling have led to improved understanding and  
1109 predictability of dispersal rates and patterns of sediment-associated toxic  
1110 metal contamination (Coulthard and Macklin, 2003). Chemical dispersal of  
1111 mine wastes tends to predominate after mine closure and four principal  
1112 processes result in toxic metal attenuation downstream of inputs – pH  
1113 buffering, acid neutralisation, precipitation and adsorption. However, river  
1114 sediments are not a permanent store for toxic metals and they may be  
1115 released into the water column if there are fluctuations in some important  
1116 environmental parameters (i.e. pH and redox potential). As a result,  
1117 establishing metal speciation, bioavailability and potential mobility is essential  
1118 in order to prioritise sites for remediation. Recently, molecular scale  
1119 techniques to study elemental binding have become more accessible to  
1120 researchers. A greater number of geochemical studies should make use of

1121 these techniques to provide more accurate information on bonding  
1122 characteristics of metals in sediments. Environmental regulators are  
1123 beginning to acknowledge the central role of sediments in maintaining  
1124 ecological quality in river systems. We have argued that the measurement /  
1125 quantification of total metal concentrations, as is practiced by many  
1126 regulators, provides limited information on the potential toxicity of sediments.  
1127 Measurement of the bioavailable metal fraction within benthic sediments is  
1128 considered a more accurate gauge of potential metal toxicity.

1129

1130 The character of metal mine drainage after it enters surface waters is affected  
1131 by many factors including stream discharge, rainfall characteristics, conditions  
1132 antecedent to rainfall-runoff events and season, and the interaction of a large  
1133 permutation of processes which must be understood and quantified in order to  
1134 mitigate effectively. Seasonal variability in metal concentrations is linked to  
1135 oxidation and dissolution of metal sulphates, leading to elevated metal  
1136 concentrations in summer and autumn months. At many mine sites, the  
1137 transport of significant amounts of mine waste is limited to stormflows.

1138 Typically, hysteresis is evident in the relationship between metal  
1139 concentrations and discharge. Peak metal concentrations are achieved before  
1140 peak discharge, associated with the dissolution of surface oxidised material.

1141 Despite the importance of stormflows for the transport of mine wastes, little  
1142 research has concentrated on investigating toxic metal fluxes and  
1143 hydrochemical variability under these conditions. Predicted increases in the  
1144 frequency and magnitude of floods as a function of climate change may result  
1145 in increased mobilisation and deposition of toxic metals in floodplains across

1146 Europe. Stormflow hydrochemistry in rivers draining mine sites should be  
1147 studied in more detail in order to quantify metal fluxes more accurately and  
1148 allow environmental managers to prioritise areas for remediation. Toxic metal  
1149 flushing during stormflows potentially impacts stream ecosystems by  
1150 significantly increasing the toxicity of the river water, even if only for short time  
1151 periods. More research is needed to help understand the potential  
1152 toxicological impacts of stormflows in mining-affected river catchments.  
1153 Relatively few studies have investigated mine spoil hydrology and metal  
1154 attenuation and release processes. Environmental investigations at  
1155 abandoned metal mine sites should include assessments of mine spoil in  
1156 terms of mineralogy, metal speciation and hydrology, especially where  
1157 contamination of groundwater is an issue.

1158

1159 Metal mine contaminants in river systems can have a variety of negative  
1160 impacts on macroinvertebrate ecology and biology, including changes to  
1161 community structure, physiological and behavioural impacts as well as direct  
1162 mortality. Typically, rivers heavily impacted by metal mine drainage have  
1163 reduced species diversity and abundance, and tend to be dominated by  
1164 Dipteran species. The order of toxicity in mining-impacted streams generally  
1165 proceeds in the order Ephemeroptera > Trichoptera > Plecoptera > Diptera.  
1166 Bioindices are used widely to quantify contaminant impacts on  
1167 macroinvertebrate communities. However, their effectiveness in discerning  
1168 the impacts of metal mine contamination is questionable, with widely varying  
1169 performance reported in the literature. The problem appears to be related to  
1170 the multi-factor nature of mine discharges. Further research is required to



1171 develop a biological index specifically for the detection of the impacts of mine  
1172 water contamination on macroinvertebrate communities and the wider  
1173 ecosystem. Traditionally, laboratory bioassay experiments have been used to  
1174 investigate metal and AMD toxicity, with organism mortality being the test  
1175 endpoint. Recently, biomonitoring techniques capable of detecting sublethal  
1176 behavioural and physiological responses in an organism have become  
1177 popular (e.g., Gerhardt *et al.*, 2004). They have the principal advantage over  
1178 bioassays of integrating both biochemical and physiological processes. A  
1179 major criticism of bioassay and microcosm studies is that they generally do  
1180 not consider metal mixtures or the influence of other environmental  
1181 parameters on metal toxicity. The development of the Biotic Ligand Model has  
1182 allowed organism physiology and important environmental parameters to be  
1183 factored into assessments of metal toxicity (Di Toro *et al.*, 2001). However, to  
1184 reach their full potential, BLMs will need to be applied to a wider range of  
1185 organisms and pollutants, and they will need to be able to incorporate metal  
1186 mixtures into toxicity predictions.

1187

1188 An increasing range of remediation technologies have been developed for the  
1189 treatment of contaminated mine water which can be applied in a variety of  
1190 topographical settings. Chemical treatment of mine waters is expensive and  
1191 unsustainable over the substantial time periods treatment will be required.

1192 Passive remediation technologies offer a low cost and sustainable alternative.

1193 Passive systems for the treatment of coal mine discharges, where iron,  
1194 sulphates and acidity are the principal contaminants, are considered proven  
1195 technology. However, these systems generally fail to remove toxic metals

1196 (e.g., Zn, Pb, Cd), associated with metal mine discharges, to an acceptable  
1197 standard. New substrates and techniques aimed at removing high  
1198 concentrations of these toxic metals are being trialled and many show  
1199 promise at the laboratory scale. However, large-scale pilot treatment plants  
1200 are needed in order to develop these new systems and to test them in field-  
1201 relevant conditions. Even with mine water treatment, mine spoil and  
1202 contaminated soils in mining regions will continue to pose a threat to water  
1203 and ecological quality for many years into the future. New bio-based source  
1204 control techniques such as phytoremediation offer the possibility of stabilising,  
1205 immobilising and extracting toxic metals from soils at low cost, by using plants  
1206 which hyper-accumulate toxic metals in their tissue. However, the long-term  
1207 functioning and ecological impact of these new strategies needs to be  
1208 evaluated.

1209

1210 A management approach which can draw on the expertise of separate but  
1211 related and relevant disciplines such as hydrology, hydrochemistry, sediment  
1212 geochemistry, fluvial geomorphology and aquatic ecology affords the  
1213 opportunity for a more complete understanding of processes and impacts in  
1214 mining-impacted river catchments. It is hoped that this review will help to  
1215 contribute to our knowledge and understanding of the impacts of metal mining  
1216 on aquatic ecosystems and highlight the usefulness of approaching such  
1217 problems from a multi-disciplinary geographical point of view.

1218

1219 **Acknowledgements**

1220 PB gratefully acknowledges the support of a Loughborough University

1221 Department of Geography Scholarship which allowed this research to be

1222 undertaken.

1223

1224 **References**

1225 Abdullah MI, Royle LG. Heavy metal content of some rivers and lakes in  
1226 Wales. *Nature* 1972; 238: 329-330.

1227 Aisemberg J, Nahabedian DE, Wider EA, Verrengia Guerrero NR.  
1228 Comparative study on two freshwater invertebrates for monitoring  
1229 environmental lead exposure. *Toxicology* 2005; 210: 45-53.

1230 Albering HJ, van Leusen SM, Moonen EJC, Hoogewerff JA, Kleinjans JCS.  
1231 Human health risk assessment: A case study involving heavy metal soil  
1232 contamination after the flooding of the river Meuse during the winter of  
1233 1993-1994. *Environmental Health Perspectives* 1999; 107: 37-43.

1234 Alderton DHM, Serafimovski T, Mullen B, Fairall K, James S. The chemistry of  
1235 waters associated with metal mining in Macedonia. *Mine Water and the  
1236 Environment* 2005; 24: 139-149.

1237 Aleksander-Kwaterczak U, Helios-Rybicka E. Contaminated sediments as a  
1238 potential source of Zn, Pb, and Cd for a river system in the historical  
1239 metalliferous ore mining and smelting industry area of South Poland.  
1240 *Journal of Soils and Sediments* 2009; 9: 13-22.

1241 Aleva GJJ. Indonesian Fluvial Cassiterite Placers and Their Genetic  
1242 Environment. *Journal of the Geological Society* 1985; 142: 815-836.

1243 Amisah S, Cowx IG. Impacts of abandoned mine and industrial discharges on  
1244 fish abundance and macroinvertebrate diversity of the upper River Don  
1245 in South Yorkshire, UK. *Journal of Freshwater Biology* 2000; 15: 237-  
1246 250.

1247 Armitage PD. The effects of mine drainage and organic enrichment on  
1248 benthos in the River Nent system, Northern Pennines. *Hydrobiologia*  
1249 1980; 74: 119-128.

1250 Armitage PD, Bowes MJ, Vincent HM. Long-term changes in  
1251 macroinvertebrate communities of a heavy metal polluted stream: the  
1252 River Nent (Cumbria, UK) after 28 years. *River Research and  
1253 Applications* 2007; 23: 997-1015.

1254 Asta MP, Cama J, Gault AG, Charnock JM, Queralt I. Characterisation of  
1255 AMD sediments in the discharge of the Tinto Santa Rosa mine (Iberian  
1256 Pyritic Belt, SW Spain). In: Cidu R, Frau F, editors. *International Mine  
1257 Water Association Symposium 2007: Water in Mining Environments,  
1258 Cagliari (Mako Edizioni)*, 2007.

1259 Bahrndorff S, Ward J, Pettigrove V, Hoffmann AA. A microcosm test of  
1260 adaptation and species specific responses to polluted sediments  
1261 applicable to indigenous chironomids (Diptera). *Environmental  
1262 Pollution* 2006; 139: 550-560.

- 1263 Balci NC. Effects of bacterial activity on the release of trace metals from  
1264 sphalerite oxidation. In: Rapantova N, Hrkal Z, editors. Mine Water and  
1265 the Environment. Ostrava (VSB – Technical University of Ostrava),  
1266 2008.
- 1267 Banks D, Parnachev VP, Frengstad B, Holden W, Vedernikov AA, Kannachuk  
1268 OV. Alkaline mine drainage from metal sulphide and coal mines:  
1269 examples from Svalbard and Siberia. In: Younger PL, Robins NS,  
1270 editors. Mine Water Hydrogeology and Geochemistry. The Geological  
1271 Society, London, 2002, pp. 287-296.
- 1272 Banks D, Younger PL, Arnesen RT, Iversen ER, Banks SB. Mine-water  
1273 chemistry: the good, the bad and the ugly. *Environmental Geology*  
1274 1997; 32: 157-174.
- 1275 Banks SB, Banks D. Abandoned mines drainage: impact assessment and  
1276 mitigation of discharges from coal mines in the UK. *Engineering*  
1277 *Geology* 2001; 60: 31-37.
- 1278 Batty LC. The potential importance of mine sites for biodiversity. *Mine Water*  
1279 *and the Environment* 2005; 24: 101-103.
- 1280 Batty LC, Auladell M, Sadler J. The impacts of metalliferous drainage on  
1281 aquatic communities. In: Batty LC, Hallberg KB, editors. *Ecology of*  
1282 *Industrial Pollution*. Cambridge University Press, Cambridge, 2010, pp.  
1283 70-100.
- 1284 Batty LC, Younger PL. The use of waste materials in the passive remediation  
1285 of mine water pollution. *Surveys in Geophysics* 2004; 25: 55-67.
- 1286 Beltman DJ, Clements WH, Lipton J, Cacela D. Benthic invertebrate metals  
1287 exposure, accumulation and community-level effects downstream from  
1288 a hard rock mine site. *Environmental Toxicology and Chemistry* 1999;  
1289 18: 299-307.
- 1290 Benner SG, Blowes DW, Ptacek CJ. A full-scale porous reactive wall for  
1291 prevention of acid mine drainage. *Ground Water Monitoring and*  
1292 *Remediation* 1997; 17: 99-107.
- 1293 Berger WH, Parker FL. Diversity of planktonic foraminifera in deep-sea  
1294 sediments. *Science of the Total Environment* 1970; 168: 1345-1347.
- 1295 Berninger K, Pennanen J. Heavy metals in perch (*Perca fluviatilis* L.) from two  
1296 acidified lakes in the Salpausselkae esker area in Finland. *Water, Air,*  
1297 *and Soil Pollution* 1995; 81: 283-294.
- 1298 Biological Monitoring Working Party. Final Report: Assessment and  
1299 Presentation of Biological Quality of Rivers in Great Britain.  
1300 Unpublished report. Department of the Environment Water Data Unit,  
1301 1978.
- 1302 Bird G, Brewer PA, Macklin MG, Nikolova M, Kotsev T, Mollov M, et al. Pb  
1303 isotope evidence for contaminant-metal dispersal in an international  
1304 river system: The lower Danube catchment, Eastern Europe. *Applied*  
1305 *Geochemistry* 2010; 25: 1070-1084.
- 1306 Bird SC. The effect of hydrological factors on trace metal contamination in the  
1307 River Tawe, South Wales. *Environmental Pollution* 1987; 45: 87-124.
- 1308 Boulton S, Collins DN, White KN, Curtis CD. Metal transport in a stream polluted  
1309 by acid mine drainage - the Afon Goch, Anglesey, UK. *Environmental*  
1310 *Pollution* 1994; 84: 279-284.
- 1311 Bradley JB, Cox JJ. The significance of the floodplain to the cycling of metals  
1312 in the River Derwent catchment, UK. *Science of the Total Environment*

1313 1990; 97/98: 441-454.

1314 Bradley SB. Flood effects of the transport of heavy metals. *International*  
1315 *Journal of Environmental Studies* 1984; 22: 225-230.

1316 Bradley SB, Foster IDL, Gurnell AM, Webb BW. Long-term dispersal of metals  
1317 in mineralised catchments by fluvial processes. In: Foster IDL, Gurnell  
1318 AM, Webb BW, editors. *Sediment and Water Quality in River*  
1319 *Catchments*. John Wiley & Sons Ltd, Chichester, 1995, pp. 161-177.

1320 Bradley SB, Lewin J. Transport of heavy metals on suspended sediments  
1321 under high flow conditions in a mineralised region of Wales.  
1322 *Environmental Pollution (Series B)* 1982; 4: 257-267.

1323 Braungardt CB, Achterberg EP, Elbaz-Poulichet F, Morley NH. Metal  
1324 geochemistry in a mine-polluted estuarine system in Spain. *Applied*  
1325 *Geochemistry* 2003; 18: 1757-1771.

1326 Brewer PA, Dennis IA, Macklin MG. The use of geomorphological mapping  
1327 and modelling for identifying land affected by metal contamination on  
1328 river floodplains: DEFRA, 2005.

1329 Brinkman SF, Johnston WD. Acute toxicity of aqueous copper, cadmium, and  
1330 zinc to the mayfly *Rithrogena hageni*. *Archives of Environmental*  
1331 *Contamination and Toxicology* 2008; 54: 466-472.

1332 Brkovic-Popovic I, Popovic M. Effects of heavy metals on survival and  
1333 respiration rate of tubificid worms: Part 1 - effects on survival.  
1334 *Environmental Pollution* 1977; 13: 65-72.

1335 Brumbaugh WG, Mora MA, May TW, Phalen DN. Metal exposure and effects  
1336 in voles and small birds near a mining haul road in Cape Krusenstern  
1337 National Monument, Alaska. *Environmental Monitoring and*  
1338 *Assessment* 2010; 170: 73-86.

1339 Burrows IG, Whitton BA. Heavy metals in water, sediments and invertebrates  
1340 from a metal-contaminated river free of organic pollution. *Hydrobiologia*  
1341 1983; 106: 263-273.

1342 Burton AG. Metal Bioavailability and Toxicity in Sediments. *Critical Reviews in*  
1343 *Environmental Science and Technology* 2010; 40: 852 - 907.

1344 Butler BA. Effect of pH, ionic strength, dissolved organic carbon, time, and  
1345 particle size on metals release from mine drainage impacted  
1346 streambed sediments. *Water Research* 2009; 43: 1392-1402.

1347 Byrne P, Reid I, Wood PJ. Short-term fluctuations in heavy metal  
1348 concentrations during flood events through abandoned metal mines,  
1349 with implications for aquatic ecology and mine water treatment.  
1350 *International Mine Water Conference*. Water Institute of Southern  
1351 Africa and International Mine Water Association, Pretoria, South Africa,  
1352 2009, pp. 124-129.

1353 Byrne P, Reid I, Wood PJ. Sediment geochemistry of streams draining  
1354 abandoned lead/zinc mines in central Wales: the Afon Twymyn.  
1355 *Journal of Soils and Sediments* 2010; 4: 683-697.

1356 Calmano W, Hong J, Forstner U. Binding and mobilisation of heavy metals in  
1357 contaminated sediments affected by pH and redox potential. *Water*  
1358 *Science and Technology* 1993; 28: 223-235.

1359 Campbell PGC, Stokes PM. Acidification and toxicity of metals to aquatic  
1360 biota. *Canadian Journal of Fisheries and Aquatic Sciences* 1985; 42:  
1361 2034-2049.

1362 Canovas CR, Hubbard CG, Olias M, Nieto JM, Black S, Coleman ML.

1363 Hydrochemical variations and contaminant load in the Rio Tinto (Spain)  
1364 during flood events. *Journal of Hydrology* 2008; 350: 25-40.

1365 Carlisle DM, Clements WH. Leaf litter breakdown and shredder production in  
1366 metal-polluted streams. *Freshwater Biology* 2005; 50: 380-390.

1367 Carpenter J, Odum WE, Mills A. Leaf litter decomposition in a reservoir  
1368 affected by acid mine drainage. *Oikos* 1983; 41: 165-172.

1369 Chadwick JW, Canton SP. Inadequacy of diversity indices in discerning metal  
1370 mine drainage effects on a stream invertebrate community. *Water, Air,  
1371 and Soil Pollution* 1984; 22: 217-223.

1372 Chadwick JW, Canton SP, Dent RL. Recovery of benthic invertebrate  
1373 communities in Silver Bow Creek, Montana, following improved metal  
1374 mine wastewater treatment. *Water, Air and Soil Pollution* 1986; 28:  
1375 427-438.

1376 Chapa-Vargas L, Mejia-Saavedra JJ, Monzalvo-Santos K, Puebla-Olivares F.  
1377 Blood lead concentrations in wild birds from a polluted mining region at  
1378 Villa de La Paz, San Luis Potosi, Mexico. *Journal of Environmental  
1379 Science and Health Part a-Toxic/Hazardous Substances &  
1380 Environmental Engineering* 2010; 45: 90-98.

1381 Cid N, Ibanez C, Palanques A, Prat N. Patterns of metal bioaccumulation in  
1382 two filter-feeding macroinvertebrates: Exposure distribution, inter-  
1383 species differences and variability across developmental stages.  
1384 *Science of the Total Environment* 2010; 408: 2795-2806.

1385 Cidu R, Di Palma M, Medas D. The Fluminese Mining District (SW Sardinia,  
1386 Italy): Impact of the past lead-zinc exploitation on aquatic environment.  
1387 In: Cidu R, Frau F, editors. *International Mine Water Association  
1388 Symposium 2007: Water in the Mining Environment*, Cagliari (Mako  
1389 Edizioni), 2007, pp. 47-51.

1390 Cidu R, Mereu L. The abandoned copper-mine of Funtana Raminosa  
1391 (Sardinia): Preliminary evaluation of its impact on the aquatic system.  
1392 In: Cidu R, Frau F, editors. *International Mine Water Association  
1393 Symposium 2007: Water in Mining Environments*, Cagliari (Mako  
1394 Edizioni), 2007, pp. 53-57.

1395 Clements WH. Small-scale experiments support causal relationships between  
1396 metal contamination and macroinvertebrate community responses.  
1397 *Ecological Applications* 2004; 14: 954-967.

1398 Clements WH, Carlisle DM, Lazorchak JM, Johnson PC. Heavy metals  
1399 structure benthic communities in Colorado mountain streams.  
1400 *Ecological Applications* 2000; 10: 626-638.

1401 Clements WH, Cherry DS, Cairns J. The influence of copper exposure on  
1402 predator-prey interactions in aquatic insect communities. *Freshwater  
1403 Biology* 1989; 21: 483-488.

1404 Clements WH, Cherry DS, Van Hassel JH. Assessment of the impact of  
1405 heavy metals on benthic communities at the Clinch River (Virginia):  
1406 Evaluation of an index of community sensitivity. *Canadian Journal of  
1407 Fisheries and Aquatic Sciences* 1992; 49: 1686-1694.

1408 Clews E, Ormerod SJ. Improving bio-diagnostic monitoring using simple  
1409 combinations of standard biotic indices. *River Research and  
1410 Applications* 2009; 25: 348-361.

1411 Comber SD, Merrington G, Sturdy L, Delbeke K, van Assche F. Copper and  
1412 zinc water quality standards under the EU Water Framework Directive:

- 1413 The use of a tiered approach to estimate the levels of failure. *Science*  
1414 *of the Total Environment* 2008; 403: 12-22.
- 1415 Conesa HM, Perez-Chacon JA, Arnaldos R, Moreno-Caselles J, Faz-Cano A.  
1416 In situ heavy metal accumulation in lettuce growing near a former  
1417 mining waste disposal area: Implications for agricultural management.  
1418 *Water, Air and Soil Pollution* 2010; 208: 377-383.
- 1419 Connelly RJ. Rehabilitation and construction issues for Silvermines  
1420 Abandoned Mine Area, Ireland. *International Mine Water Conference.*  
1421 *Water Institute of South Africa and International Mine Water*  
1422 *Association, Pretoria, South Africa, 2009, pp. 298-307.*
- 1423 Coulthard TJ, Macklin MG. Modelling long-term contamination in river  
1424 systems from historical metal mining. *Geology* 2003; 31: 451-454.
- 1425 Cui H, Li LY, Grace JR. Exploration of remediation of acid rock drainage with  
1426 clinoptilolite as sorbent in a slurry bubble column for both heavy metal  
1427 capture and regeneration. *Water Research* 2006; 40: 3359-3366.
- 1428 Davies BE, Lewin J. Chronosequences in alluvial soils with special reference  
1429 to historic lead pollution in Cardiganshire, Wales. *Environmental*  
1430 *Pollution* 1974; 6: 49-57.
- 1431 Davies BE, Thornton I. Heavy metal contamination from base metal mining  
1432 and smelting: implications for man and his environment. In: Thornton I,  
1433 editor. *Applied Environmental Geochemistry.* Academic Press, London,  
1434 1983, pp. 425-462.
- 1435 Davy-Bowker J, Murphy JF, Rutt GP, Steel JEC, Furse MT. The development  
1436 and testing of a macroinvertebrate biotic index for detecting the impact  
1437 of acidity on streams. *Archiv fuer Hydrobiologie* 2005; 163: 383-403.
- 1438 Dawson EJ, Macklin MG. Speciation of heavy metals on suspended sediment  
1439 under high flow conditions in the River Aire, West Yorkshire, UK.  
1440 *Hydrological Processes* 1998; 12: 1483-1494.
- 1441 De Bisthoven JL, Gerhardt A, Guhr K, Soares AMVM. Behavioural changes  
1442 and acute toxicity to the freshwater shrimp *Atyaephyra desmaresti*  
1443 *Millet (Decapoda: Natantia)* from exposure to acid mine drainage.  
1444 *Ecotoxicology* 2006; 15: 215-227.
- 1445 De Bisthoven JL, Gerhardt A, Soares AMVM. Effects of acid mine drainage on  
1446 larval *Chironomus (Diptera, Chironomidae)* measured with the  
1447 *Multispecies Freshwater Biomonitor.* *Environmental Toxicology and*  
1448 *Chemistry* 2004; 23: 1123-1128.
- 1449 De Bisthoven JL, Gerhardt A, Soares AMVM. Chironomidae larvae as  
1450 bioindicators of an acid mine drainage in Portugal. *Hydrobiologia* 2005;  
1451 532: 181-191.
- 1452 DeNicola DM, Stapleton MG. Impact of acid mine drainage on benthic  
1453 communities in streams: the relative roles of substratum vs. aqueous  
1454 effects. *Environmental Pollution* 2002; 119: 303-315.
- 1455 Dennis IA, Coulthard TJ, Brewer PA, Macklin MG. The role of floodplains in  
1456 attenuating contaminated sediment fluxes in formerly mined drainage  
1457 basins. *Earth Surface Processes and Landforms* 2009; 34: 453-466.
- 1458 Dennis IA, Macklin MG, Coulthard TJ, Brewer PA. The impact of the October-  
1459 November 2000 floods on contaminant metal dispersal in the River  
1460 Swale catchment, North Yorkshire, UK. *Hydrological Processes* 2003;  
1461 17: 1641-1657.
- 1462 Desbarats AJ, Dirom GC. Temporal variation in discharge chemistry and

- 1463 portal flow from the 8-Level adit, Lynx Mine, Myra Falls Operations,  
1464 Vancouver Island, British Columbia. *Environmental Geology* 2005; 47:  
1465 445-456.
- 1466 Di Toro DM, Allen HE, Bergman HL, Meyer JS, Paquin PR, Santore RC.  
1467 Biotic ligand model of the acute toxicity of metals. 1, Technical basis.  
1468 *Environmental Toxicology and Chemistry* 2001; 20: 2383-2396.
- 1469 Dojlido JR, Taboryska B. Exchange of heavy metals between sediment and  
1470 water in the Wloclawek Reservoir on the Vistula River. In: Peters NE  
1471 WD, editor. *Sediment and Stream Water Quality in a Changing*  
1472 *Environment: Trends and Explanations*. IAHS Pub. no. 203, Vienna,  
1473 1991, pp. 315-320.
- 1474 Dressing SA, Mass RP, Weiss CM. Effect of chemical speciation on the  
1475 accumulation of cadmium by the caddisfly, *Hydropsyche* sp. *Bulletin of*  
1476 *Environmental Contamination and Toxicology* 1982; 28: 172-180.
- 1477 Edraki M, Golding SD, Baublys KA, Lawrence MG. Hydrochemistry,  
1478 mineralogy and sulfur isotope geochemistry of acid mine drainage at  
1479 the Mt. Morgan mine environment, Queensland, Australia. *Applied*  
1480 *Geochemistry* 2005; 20: 789-805.
- 1481 Enid Martinez C, McBride MB. Solubility of Cd<sup>2+</sup>, Cu<sup>2+</sup>, Pb<sup>2+</sup>, and Zn<sup>2+</sup>  
1482 in aged coprecipitates with amorphous iron hydroxides. *Environmental*  
1483 *Science and Technology* 1998; 32: 743-748.
- 1484 Environment Agency. *Metal Mine Strategy for Wales*. Environment Agency  
1485 Wales, Cardiff, 2002.
- 1486 Environment Agency. *Attenuation of mine pollutants in the hyporheic zone*.  
1487 Environment Agency, Bristol, 2006.
- 1488 Environment Agency. *Abandoned mines and the water environment*. Bristol.  
1489 Environment Agency, 2008a.
- 1490 Environment Agency. *Assessment of metal mining-contaminated river*  
1491 *sediments in England and Wales*. Environment Agency, Bristol, 2008b.
- 1492 Ernst WHO. Phytoextraction of mine wastes - options and impossibilities.  
1493 *Chemie der Erde* 2005; 65: 29-42.
- 1494 Esbri JM, Bernaus A, Avila M, Kocman D, Garcia-Noquero EM, Gaona X, et  
1495 al. XANES speciation of mercury in three mining districts - Almaden,  
1496 Asturia (Spain), Idria (Slovenia). *Journal of Synchrotron Radiation*  
1497 2010; 17: 179-186 Part 2.
- 1498 Evangelou VP, Zhang YL. A review - pyrite oxidation mechanisms and acid-  
1499 mine drainage prevention. *Critical Reviews in Environmental Science*  
1500 *and Technology* 1995; 25: 141-199.
- 1501 Evans LJ, Adams WA. Chlorite and illite in some lower Palaeozoic mudstones  
1502 of mid-Wales. *Clay Minerals* 1975; 10: 387-397.
- 1503 Farag AM, Woodward DF, Goldstein JN, Brumbaugh W, Meyer JS.  
1504 Concentrations of metals associated with mining waste in sediments,  
1505 biofilm, benthic macroinvertebrates, and fish from the Coeur d'Alene  
1506 River Basin, Idaho. *Archives of Environmental Contamination and*  
1507 *Toxicology* 1998; 34: 119-127.
- 1508 Fergusson JE. *The Heavy Elements. Chemistry, Environmental Impact and*  
1509 *Health Effects*. Oxford: Pergamon Press, 1990.
- 1510 Filipek LH, Nordstrom DK, Ficklin WH. Interaction of acid mine drainage with  
1511 waters and sediments of West Squaw Creek in the West Shasta Mining  
1512 District, California. *Environmental Science and Technology* 1987; 21:



1513 388-396.

1514 Ford RG, Bertsch PM, Farley KJ. Changes in transition and heavy metal  
1515 partitioning during hydrous iron oxide aging. *Environmental Science*  
1516 *and Technology* 1997; 31: 2028-2033.

1517 Forstner U. Sediment dynamics and pollutant mobility in rivers: An  
1518 interdisciplinary approach. *Lakes and Reservoirs. Research and*  
1519 *Management* 2004; 9: 25-40.

1520 Forstner U. Sediments and priority substances in river basins. *Journal of Soils*  
1521 *and Sediments* 2009; 9: 89-93.

1522 Forstner U, Salomons W. Trends and challenges in sediment research 2008:  
1523 the role of sediments in river basin management. *Journal of Soils and*  
1524 *Sediments* 2008; 8: 281-283.

1525 Foster IDL, Charlesworth SM. Heavy metals in the hydrological cycle: trends  
1526 and explanations. *Hydrological Processes* 1996; 10: 227-261.

1527 Fuge R, Laidlaw IMS, Perkins WT, Rogers KP. The influence of acidic mine  
1528 and spoil drainage on water quality in the mid-Wales area.  
1529 *Environmental Geochemistry and Health* 1991; 13: 70-75.

1530 Galan E, Gomez-Ariza JL, Gonzalez I, Fernandez-Caliani JC, Morales E,  
1531 Giraldez I. Heavy metal partitioning in river sediments severely polluted  
1532 by acid mine drainage in the Iberian Pyrite Belt. *Applied Geochemistry*  
1533 2003; 18: 409-421.

1534 Gammons CH, Shope CL, Duaine TE. A 24 h investigation of the  
1535 hydrogeochemistry of baseflow and stormwater in an urban area  
1536 impacted by mining: Butte, Montana. *Hydrological Processes* 2005; 19:  
1537 2737-2753.

1538 Gandy CJ, Younger PL. Effect of a clay cap on oxidation of Pyrite within mine  
1539 spoil. *Quarterly Journal of Engineering Geology and Hydrogeology*  
1540 2003; 36: 207-215.

1541 Garcia-Garcia G, Nandini S. Turbidity mitigates lead toxicity to cladocerans  
1542 (Cladocera). *Ecotoxicology* 2006; 15: 425-436.

1543 Gauss JD, Woods PE, Winner RW, Skillings JH. Acute toxicity of copper to  
1544 three life stages of Chironomus tentans as affected by water  
1545 hardness-alkalinity. *Environmental Pollution (Series A)* 1985; 37: 149-  
1546 157.

1547 Geer R. Reconstructing the geomorphological and sedimentological impacts  
1548 of a catastrophic flood event, Dale Beck Valley, Caldbeck Fells,  
1549 Cumbria. MSc thesis. University of Leeds, Leeds, 2004.

1550 Gerhardt A. Review of impact of heavy metals on stream invertebrates with  
1551 special emphasis on acid conditions. *Water, Air, and Soil Pollution*  
1552 1993; 66: 289-314.

1553 Gerhardt A. Importance of exposure route for behavioural responses in  
1554 *Lumbricus variegatus* Muller (Oligochaeta: Lumbriculida) in short-  
1555 term exposures to Pb. *Environmental Science and Pollution Research*  
1556 2007; 14: 430-434.

1557 Gerhardt A, De Bisthoven JL, Soares AMVM. Effects of acid mine drainage  
1558 and acidity on the activity of *Choroterpes picteti* (Ephemeroptera:  
1559 *Leptophlebiidae*). *Archives of Environmental Contamination and*  
1560 *Toxicology* 2005a; 48: 450-458.

1561 Gerhardt A, De Bisthoven LJ, Soares AMVM. Macroinvertebrate response to  
1562 acid mine drainage: community metrics and on-line behavioural toxicity

- 1563 bioassay. *Environmental Pollution* 2004; 130: 263-274.
- 1564 Gerhardt A, De Bisthoven LJ, Soares AMVM. Evidence for the Stepwise  
1565 Stress Model: *Gambusia holbrooki* and *Daphnia magna* under acid  
1566 mine drainage and acidified reference water stress. *Environmental  
1567 Science and Technology* 2005b; 39: 4150-4158.
- 1568 Geroni JN, Sapsford DJ, Barnes A, Watson IA, Williams KP. Current  
1569 performance of passive treatment systems in south Wales, UK.  
1570 International Mine Water Conference. Water Institute of Southern  
1571 Africa and International Mine Water Association, Pretoria, South Africa,  
1572 2009, pp. 486-496.
- 1573 Giesy JP. Cadmium inhibition of leaf decomposition in an aquatic microcosm.  
1574 *Chemosphere* 1978; 7: 467-475.
- 1575 Gilbert GK. Hydraulic-mining debris in the Sierra Nevada. US Geological  
1576 Survey Paper 105 1917.
- 1577 Gilchrist S, Gates A, Szabo Z, Lamothe PJ. Impact of AMD on water quality in  
1578 critical watershed in the Hudson River drainage basin: Phillips Mine,  
1579 Hudson Highlands, New York. *Environmental Geology* 2009; 57: 397-  
1580 409.
- 1581 Goodyear KL, McNeill S. Bioaccumulation of heavy metals by aquatic  
1582 macroinvertebrates of different feeding guilds: a review. *Science of the  
1583 Total Environment* 1999; 229: 1-19.
- 1584 Goodyear KL, Ramsey MH, Thorton I, Rosenbaum MS. Source identification  
1585 of Pb-Zn contamination in the Allen Basin, Cornwall, S.W. England.  
1586 *Applied Geochemistry* 1996; 11: 61-68.
- 1587 Gower AM, Darlington ST. Relationships between copper concentrations in  
1588 larvae of *Plectrocnemia conspersa* (Curtis) (Trichoptera) and in mine  
1589 drainage streams. *Environmental Pollution* 1990; 65: 155-168.
- 1590 Gower AM, Myers G, Kent M, Foulkes ME. Relationships between  
1591 macroinvertebrate communities and environmental variables in metal-  
1592 contaminated streams in south-west England. *Freshwater Biology*  
1593 1994; 32: 199-221.
- 1594 Gray NF. Environmental impact and remediation of acid mine drainage: a  
1595 management problem. *Environmental Geology* 1997; 30: 62-71.
- 1596 Gray NF. Acid mine drainage composition and the implications for its impact  
1597 on lotic systems. *Water Research* 1998; 32: 2122-2134.
- 1598 Gray NF, Delaney E. Comparison of benthic macroinvertebrate indices for  
1599 the assessment of the impact of acid mine drainage on an Irish river  
1600 below an abandoned Cu-S mine. *Environmental Pollution* 2008; 155:  
1601 31-40.
- 1602 Grimshaw DL, Lewin J, Fuge R. Seasonal and short-term variations in the  
1603 concentration and supply of dissolved zinc to polluted aquatic  
1604 environments. *Environmental Pollution* 1976; 11: 1-7.
- 1605 Groenendijk D, Lucker SMG, Plans M, Kraak MHS, Admiraal W. Dynamics of  
1606 metal adaptation in riverine chironomids. *Environmental Pollution* 2002;  
1607 117: 101-109.
- 1608 Groenendijk D, Zenstra LWM, Postma JF. Fluctuating assymetry and mentum  
1609 gaps in populations of the midge *Chironomous riparius* (Diptera:  
1610 Chironomidae) from a metal-contaminated river. *Environmental  
1611 Toxicology and Chemistry* 1998; 17: 1999-2005.
- 1612 Guangwei Y, Hengyi L, Tao B, Zhong L, Qiang Y, Xianqiang S. In-situ

1613 stabilisation followed by ex-situ composting for treatment and disposal  
1614 of heavy metals polluted sediments. *Journal of Environmental Sciences*  
1615 2009; 21: 877-883.

1616 Gundersen P, Steinnes E. Influence of temporal variations in river discharge,  
1617 pH, alkalinity and Ca on the speciation and concentration of heavy  
1618 metals in some mining polluted rivers. *Aquatic Geochemistry* 2001; 7:  
1619 173-193.

1620 Hallare AV, Seiler T-B, Hollert H. The versatile, changing, and advancing  
1621 roles of fish in sediment toxicity assessment - a review. *Journal of Soils*  
1622 *and Sediments* 2010; 11.

1623 Hallberg KB, Johnson DB. Mine water microbiology. *Mine Water and the*  
1624 *Environment* 2005; 24: 28-32.

1625 Hammarstrom JM, Seal RR, II., Meier AM, Kornfeld JM. Secondary sulfate  
1626 minerals associated with acid drainage in the eastern US: recycling of  
1627 metals and acidity in surficial environments. *Chemical Geology* 2005;  
1628 215: 407-431.

1629 Hawkins JW. Predictability of surface mine spoil hydrologic properties in the  
1630 Appalachian Plateau. *Groundwater* 2004; 42: 119-125.

1631 Hedin RS, Nairn RW, Kleinmann RLP. *Passive Treatment of Coal Mine*  
1632 *Drainage*. US Bureau of Mines, 1994.

1633 Hering D, et al. The European Water Framework Directive at the age of 10: A  
1634 critical review of the achievements with recommendations for the  
1635 future. *Science of the Total Environment* 2010; 408: 4007-4019.

1636 Hermann R, Neumann-Mahlkau P. The mobility of zinc, cadmium, copper,  
1637 lead, iron and arsenic in ground water as a function of redox potential  
1638 and pH. *Science of the Total Environment* 1985; 43: 1-12.

1639 Herr C, Gray NF. Seasonal variation of metal contamination of riverine  
1640 sediments below a copper and sulphur mine in south-east Ireland.  
1641 *Water Science and Technology* 1996; 35: 255-261.

1642 Hickey CW, Golding LA. Response of macroinvertebrates to copper and zinc  
1643 in a stream mesocosm. *Environmental Toxicology and Chemistry* 2002;  
1644 21: 1854-1863.

1645 Hirst H, Juttner I, Ormerod SJ. Comparing the responses of diatoms and  
1646 macroinvertebrates to metals in upland streams of Wales and  
1647 Cornwall. *Freshwater Biology* 2002; 47: 1752-1765.

1648 Howard AG. *Aquatic Environmental Chemistry*. Oxford: Oxford University  
1649 Press, 1998.

1650 Hudson-Edwards KA, Macklin MG, Curtis CD, Vaughn DJ. Processes of  
1651 formation and distribution of Pb, Zn, Cd and Cu bearing minerals in the  
1652 Tyne Basin, northeast England: Implications for metal-contaminated  
1653 river systems. *Environmental Science and Technology* 1996; 30: 72-  
1654 80.

1655 Hudson-Edwards KA, Macklin MG, Jamieson HE, Brewer PA, Coulthard TJ,  
1656 Howard AJ, et al. The impact of tailings dam spills and clean-up  
1657 operations on sediment and water quality in river systems: the Rios  
1658 Agrio-Guadiamar, Aznalcollar, Spain. *Applied Geochemistry* 2003; 18:  
1659 221-239.

1660 Hudson-Edwards KA, Macklin MG, Taylor M. Historic metal mining inputs to  
1661 Tees river sediment. *Science of the Total Environment* 1997; 194/195:  
1662 437-445.

- 1663 Hudson-Edwards KA, Macklin MG, Taylor MP. 2000 years of sediment-borne  
1664 heavy metal storage in the Yorkshire Ouse basin, NE England, UK.  
1665 Hydrological Processes 1999a; 13: 1087-1102.
- 1666 Hudson-Edwards KA, Schell C, Macklin MG. Mineralogy and geochemistry of  
1667 alluvium contaminated by metal mining in the Rio Tinto area, southwest  
1668 Spain. Applied Geochemistry 1999b; 14: 1015-1030.
- 1669 Jage C, Zipper C, Noble R. Factors affecting alkalinity generation by  
1670 successive alkalinity producing systems: regression analysis. Journal  
1671 of Environmental Quality 2001; 30: 1015-1022.
- 1672 Jain CK. Metal fractionation study on bed sediments of River Yamuna, India.  
1673 Water Research 2004; 38: 569-578.
- 1674 Jarvis AP, Fox A, Gozzard E, Hill S, Mayes WM, Potter HAB. Prospects for  
1675 effective national management of abandoned metal mine water  
1676 pollution in the UK. In: Cidu R, Frau F, editors. International Mine  
1677 Water Association Symposium 2007: Water in Mining Environments,  
1678 Cagliari (Mako Edizioni), 2007, pp. 77-81.
- 1679 Jarvis AP, Moustafa M, Orme PHA, Younger PL. Effective remediation of  
1680 grossly polluted acidic, and metal-rich, spoil heap drainage using a  
1681 novel, low-cost, permeable reactive barrier in Northumberland, UK.  
1682 Environmental Pollution 2006; 143: 261-268.
- 1683 Jarvis AP, Younger PL. Passive treatment of ferruginous mine waters using  
1684 high surface area media. Water Research 2001; 35: 3643-3648.
- 1685 Johnson DB. Chemical and microbiological characteristics of mineral spoils  
1686 and drainage waters at abandoned coal and metal mines. Water, Air,  
1687 and Soil Pollution 2003; 3: 47-66.
- 1688 Johnson DB, Hallberg KB. Acid mine drainage remediation options: A review.  
1689 Science of the Total Environment 2005; 338: 3-14.
- 1690 Jop KM. Concentration of metals in various larval stages of four  
1691 Ephemeroptera species. Bulletin of Environmental Contamination and  
1692 Toxicology 1991; 46: 901-905.
- 1693 Jurjovec J, Ptacek CJ, Blowes DW. Acid neutralization mechanisms and  
1694 metal release in mine tailings: A laboratory column experiment.  
1695 Geochimica et Cosmochimica Acta 2002; 66: 1511-1523.
- 1696 Keith DC, Runnells DD, Esposito KJ, Chermak JA, Levy DB, Hannula SR, et  
1697 al. Geochemical models of the impact of acidic groundwater and  
1698 evaporative sulfate salts on Boulder Creek at Iron Mountain, California.  
1699 Applied Geochemistry 2001; 16: 947-961.
- 1700 Kelly M. Mining and the Freshwater Environment. Barking: Elsevier Science  
1701 Publishing, 1988.
- 1702 Kepler D, McCleary E. Successive alkalinity producing systems (SAPS) for  
1703 the treatment of acid mine drainage. Proceedings of the International  
1704 Land Reclamation and Mine Drainage Conference, Pittsburgh, PA,  
1705 USA, 1994.
- 1706 Kiffney PM. Main and interactive effects of invertebrate density, predation,  
1707 and metals on a Rocky Mountain stream macroinvertebrate  
1708 community. Canadian Journal of Fisheries and Aquatic Sciences 1996;  
1709 53: 1595-1601.
- 1710 Kiffney PM, Clements WH. Responses of periphyton and insects to  
1711 experimental manipulation of riparian buffer width along forest streams.  
1712 Journal of Applied Ecology 2003; 40: 1060-1076.

- 1713 Knighton AD. Channel Bed Adjustment Along Mine-Affected Rivers of  
1714 Northeast Tasmania. *Geomorphology* 1991; 4: 205-219.
- 1715 Knott NA, Aulbury JP, Brown TH, Johnston EL. Contemporary ecological  
1716 threats from historical pollution sources: impacts of large-scale  
1717 resuspension of contaminated sediments on sessile invertebrate  
1718 recruitment. *Journal of Applied Ecology* 2009; 46: 770-781.
- 1719 Koukouzas N, Vasilatos C, Itskos G, Mitsis I, Moutsatsou A. Removal of  
1720 heavy metals from wastewater using CFB-coal fly ash zeolitic  
1721 materials. *Journal of Hazardous Materials* 2010; 173: 581-588.
- 1722 Kovacs E, Dubbin WE, Tamas J. Influence of hydrology on heavy metal  
1723 speciation and mobility in a Pb-Zn mine tailing. *Environmental Pollution*  
1724 2006; 141: 310-320.
- 1725 Krantzberg G. Metal accumulation by chironomid larvae: the effects of age  
1726 and body weight on metal body burdens. *Hydrobiologia* 1989; 188/189:  
1727 497-506.
- 1728 Kuwabara J, Berelson W, Balistrieri L, Woods P, Topping B, Steding D, et al.  
1729 Benthic flux of metals and nutrients into the water column of Lake  
1730 Coeur d'Alene, Idaho: report of an August 1999 pilot study. US  
1731 Geological Survey Water Resources Investigation 00-4132 (CD-ROM),  
1732 Menlow Park, California, 2000.
- 1733 Lambing JH, Nimick DA, Cleasby TE. Short-term variation of trace-element  
1734 concentrations during base flow and rainfall runoff in small basins. U.S.  
1735 Geological Survey, 1999.
- 1736 Lee G, Bigham JM, Faure G. Removal of trace metals by coprecipitation with  
1737 Fe, Al and Mn from natural waters contaminated with acid mine  
1738 drainage in the Ducktown Mining District, Tennessee. *Applied*  
1739 *Geochemistry* 2002; 17: 569-581.
- 1740 Leland HV, Fend SV, Dudley TL, Carter JL. Effects of copper on species  
1741 composition of benthic insects in a Sierra Nevada, California, stream.  
1742 *Freshwater Biology* 1989; 21: 163-179.
- 1743 Lewin J, Bradley SB, Macklin MG. Historical valley alluviation in mid-Wales.  
1744 *Geological Journal* 1983; 19: 331-350.
- 1745 Lewin J, Davies BE, Wolfenden PJ. Interactions between channel change and  
1746 historic mining sediment. In: Gregory KJ, editor. *River Channel*  
1747 *Changes*. Wiley, Chichester, 1977, pp. 353-367.
- 1748 Lewin J, Macklin MG. Metal mining and floodplain sedimentation in Britain. In:  
1749 Gardiner V, editor. *International Geomorphology 1986 Part I*. John  
1750 Wiley & Sons Ltd., 1987, pp. 1009-1027.
- 1751 Licheng Z, Guijiu Z. The species and geochemical characteristics of heavy  
1752 metals in the sediments of Kangjixi River in the Shuikoushan Mine  
1753 Area, China. *Applied Geochemistry* 1996; 11: 217-222.
- 1754 Linge KL. Methods for investigating trace element binding in sediments.  
1755 *Critical Reviews in Environmental Science and Technology* 2008; 38:  
1756 165-196.
- 1757 Lord RA, Morgan PA. Metal contamination of active stream sediments in  
1758 Upper Weardale, Northern Pennine Orefield, UK. *Environmental*  
1759 *Geochemistry and Health* 2003; 25: 95-104.
- 1760 Lottermoser BG. *Mine Waste. Characterization, Treatment, Environmental*  
1761 *Impacts*. New York: Springer, 2007.
- 1762 Lund MA, McCullough CD. *Biological Remediation of Low Sulphate Acidic Pit*

- 1763 Lake Waters with Limestone pH Neutralisation and Nutrients.  
1764 International Mine Water Conference. Water Institute of Southern  
1765 Africa and International Mine Water Association, Pretoria, South Africa,  
1766 2009, pp. 519-525.
- 1767 Luoping Z, Huan F, Xiaoxia L, Xin Y, Youhai J, Tong O. Heavy metal  
1768 contaminant remediation study of western Xiamen Bay sediment,  
1769 China: Laboratory bench scale testing results. *Journal of Hazardous*  
1770 *Materials* 2009; 172: 108-116.
- 1771 Macedo-Sousa JA, Gerhardt A, Brett CMA, Noqueira AJA, Soares AMVM.  
1772 Behavioural responses of indigenous benthic invertebrates  
1773 (*Echinogammarus meridionalis*, *Hydropsyche pellucidula* and  
1774 *Choroterpes picteti*) to a pulse of acid mine drainage: A laboratorial  
1775 study. *Environmental Pollution* 2008; 156: 966-973.
- 1776 Macedo-Sousa JA, Pestana JLT, Gerhardt A, Noqueira AJA, Soares AMVM.  
1777 Behavioural and feeding responses of *Echinogammarus meridionalis*  
1778 (Crustacea, Amphipoda) to acid mine drainage. *Chemosphere* 2007;  
1779 67: 1663-1670.
- 1780 Macklin MG. Fluxes and storage of sediment-associated heavy metals in  
1781 floodplain systems: assessment and river basin management issues at  
1782 a time of rapid environmental change. In: Anderson MG, Walling DE,  
1783 Bates PD, editors. *Floodplain Processes*. Wiley, Chichester, 1996, pp.  
1784 441-460.
- 1785 Macklin MG, Brewer PA, Balteanu D, Coulthard TJ, Driga B, Howard AJ, et al.  
1786 The long-term fate and environmental significance of contaminant  
1787 metals released by the January and March 2000 mining tailings dam  
1788 failures in Maramures County, upper Tisa Basin, Romania. *Applied*  
1789 *Geochemistry* 2003; 18: 241-257.
- 1790 Macklin MG, Brewer PA, Hudson-Edwards KA, Bird G, Coulthard TJ, Dennis  
1791 IA. A geomorphological approach to the management of rivers  
1792 contaminated by metal mining. *Geomorphology* 2006; 79: 423-447.
- 1793 Macklin MG, Dowsett RB. The chemical and physical speciation of trace  
1794 elements in fine grained overbank flood sediments in the Tyne Basin,  
1795 North-East England. *Catena* 1989; 16: 135-151.
- 1796 Macklin MG, Johnston EL, Lewin J. Pervasive and long-term forcing of  
1797 Holocene river instability and flooding in Great Britain by centennial-  
1798 scale climate change. *The Holocene* 2005; 15: 937-943.
- 1799 Macklin MG, Ridgway J, Passmore DG, Rumsby BT. The use of overbank  
1800 sediment for geochemical mapping and contamination assessment:  
1801 results from selected English and Welsh floodplains. *Applied*  
1802 *Geochemistry* 1994; 9: 689-700.
- 1803 Macklin MG, Rumsby BT, Newson MD, Billi P, Hey RD, Tacconi P, et al.  
1804 Historic overbank floods and vertical accretion of fine-grained alluvium  
1805 in the lower Tyne valley, north-east England. *Dynamics of Gravel-bed*  
1806 *Rivers*. Proceedings of the Third International Workshop on Gravel-bed  
1807 Rivers. Wiley, Chichester, 1992, pp. 564-580.
- 1808 Madzivire G, Petrik LF, Gitari WM, Balfour G, Vadapalli VRK, Ojumu TV. Role  
1809 of pH in sulphate removal from circumneutral mine water using coal fly  
1810 ash. Proceedings of the International Mine Water Conference. Water  
1811 Institute of Southern Africa's Mine Water Division and International  
1812 Mine Water Association, Pretoria, South Africa, 2009, pp. 462-471.

1813 Malmqvist B, Hoffsten P. Influence of drainage from old mine deposits on  
1814 benthic communities in central Swedish streams. *Water Research*  
1815 1999; 33: 2415-2423.

1816 Maltby L, Naylor C. Preliminary observations on the ecological relevance of  
1817 the Gammarus 'scope for growth' assay: effect of zinc on reproduction.  
1818 *Functional Ecology* 1990; 4: 393-397.

1819 Marcus WA. Copper dispersion in ephemeral stream sediments. *Earth*  
1820 *Surface Processes and Landforms* 1987; 12: 217-228.

1821 Marques MJ, Martinez-Conde E, Rovira JV. Effects of zinc and lead mining on  
1822 the benthic macroinvertebrates of a fluvial ecosystem. *Water, Air, and*  
1823 *Soil Pollution* 2003; 148: 363-388.

1824 Marques MJ, Martinez-Conde E, Rovira JV, Ordonez S. Heavy metals  
1825 pollution of aquatic ecosystems in the vicinity of a recently closed  
1826 underground lead-zinc mine (Basque Country, Spain). *Environmental*  
1827 *Geology* 2001; 40: 1125-1137.

1828 Marques APGC, Rangel AOSS, Castro PML. Remediation of heavy metal  
1829 contaminated soils: Phytoremediation as a potentially promising clean-  
1830 up technology. *Critical Reviews in Environmental Science and*  
1831 *Technology* 2009; 39: 622-654.

1832 Martinez EA, Moore BC, Schaumloffel J, Dasgupta N. Teratogenic versus  
1833 mutagenic abnormalities in Chironomid larvae exposed to zinc and  
1834 lead. *Archives of Environmental Contamination and Toxicology* 2004;  
1835 47: 193-198.

1836 Mayes WM, Gozzard E, Potter HAB, Jarvis AP. Quantifying the importance of  
1837 diffuse minewater pollution in a historically heavily coal mined  
1838 catchment. *Environmental Pollution* 2008; 151: 165-175.

1839 Mayes WM, Potter HAB, Jarvis AP. Novel approach to zinc removal from  
1840 circum-neutral mine waters using pelletised recovered hydrous ferric  
1841 oxide. *Journal of Hazardous Materials* 2009; 162: 512-520.

1842 McGinness S, Johnson BD. Seasonal variation in the microbiology and  
1843 chemistry of an acid mine drainage stream. *Science of the Total*  
1844 *Environment* 1993; 132: 27-41.

1845 McGregor RG, Blowes DW, Jambor JL, Robertson WD. The solid-phase  
1846 controls on the mobility of heavy metals at the Copper Cliff tailings  
1847 area, Sudbury, Ontario, Canada. *Journal of Contaminant Hydrology*  
1848 1998; 33: 247-271.

1849 Mendez MO, Maier RM. Phytostabilisation of mine tailings in arid and semi-  
1850 arid environments - An emerging remediation technology.  
1851 *Environmental Health Perspectives* 2008; 113: 278-283.

1852 Mighanetara K, Braungardt CB, Rieuwerts JS, Azizi F. Contaminant fluxes  
1853 from point and diffuse sources from abandoned mines in the River  
1854 Tamar catchment, UK. *Journal of Geochemical Exploration* 2009; 100:  
1855 116-124.

1856 Miller JR. The role of fluvial geomorphic processes in the dispersal of heavy  
1857 metals from mine sites. *Journal of Geochemical Exploration* 1997; 58:  
1858 101-118.

1859 Miller JR, Hudson-Edwards KA, Lechler PJ, Preston D, Macklin MG. Heavy  
1860 metal contamination of water, soil and produce within riverine  
1861 communities of the Rio Pilcomayo basin, Bolivia. *Science of the Total*  
1862 *Environment* 2004; 320: 189-209.

1863 Morillo J, Usero J, Gracia I. Partitioning of metals in sediments from the Odiel  
1864 River (Spain). *Environment International* 2002; 28: 263-271.

1865 Morley NJ, Crane M, Lewis JW. Toxicity of cadmium and zinc mixtures to  
1866 *Diplostomum spathaceum* (Trematoda: Diplostomidae) cercarial  
1867 survival. *Archives of Environmental Contamination and Toxicology*  
1868 2002; 43: 28-33.

1869 Mullinger N. Review of environmental and ecological impacts of drainage from  
1870 abandoned mines in Wales. Cardiff: Environment Agency, 2004.

1871 Munk L, Faure G, Pride DE, Bigham JM. Sorption of trace metals to an  
1872 aluminium precipitate in a stream receiving acid rock-drainage; Snake  
1873 River, Summit County, Colorado. *Applied Geochemistry* 2002; 17: 421-  
1874 430.

1875 Nagorski SA, McKinnon TE, Moore JN. Seasonal and storm-scale variations  
1876 in heavy metal concentrations of two mining-contaminated streams,  
1877 Montana, USA. *Journal De Physique IV* 2003; 107: 909-912.

1878 Nagorski SA, Moore JN, Smith DB. Distribution of metals in water and bed  
1879 sediment in a mineral-rich watershed, Montana, USA. *Mine Water and*  
1880 *the Environment* 2002; 21: 121-136.

1881 Netzband A, et al. Sediment management: An essential element of River  
1882 Basin Management Plans. *Journal of Soils and Sediments* 2007; 7:  
1883 117-132.

1884 Naji A, Ismail A, Ismail AR. Chemical speciation and contamination  
1885 assessment of Zn and Cd by sequential extraction in surface sediment  
1886 of Klang River, Malaysia. *Microchemical Journal* 2010; 95: 285-292.

1887 Natarajan KA, Subramanian S, Braun JJ. Environmental impact of metal  
1888 mining - biotechnological aspects of water pollution and remediation -  
1889 an Indian experience. *Journal of Geochemical Exploration* 2006; 88:  
1890 45-48.

1891 Navarro A, Cardellach E, Mendoza JL, Corbella M, Domenech LM. Metal  
1892 mobilization from base-metal smelting slag dumps in Sierra Almagrera  
1893 (Almeria, Spain). *Applied Geochemistry* 2008; 23: 895-913.

1894 Nayar S, Goh BPL, Chou LM. Environmental impact of heavy metals from  
1895 dredged and resuspended sediments on phytoplankton and bacteria  
1896 assessed in in situ mesocosms. *Ecotoxicology and Environmental*  
1897 *Safety* 2004; 59: 349-369.

1898 Neal C, Whitehead PG, Jeffrey H, Neal M. The water quality of the River  
1899 Carnon, west Cornwall, November 1992 to March 1994: the impacts of  
1900 Wheal Jane discharges. *Science of the Total Environment* 2005; 338:  
1901 23-39.

1902 Newson TA, Fahey M. Measurement of evaporation from saline tailings  
1903 storages. *Engineering Geology* 2003; 70: 217-233.

1904 Niyogi S, Wood CM. Biotic ligand model, a flexible tool for developing site-  
1905 specific water quality guidelines for metals. *Environmental Science and*  
1906 *Technology* 2004; 38: 6177-6192.

1907 Nordstrom DK, Alpers CN, Ptacek CJ, Blowes DW. Negative pH and  
1908 extremely acidic mine waters from Iron Mountain, California.  
1909 *Environmental Science and Technology* 2000; 34: 254-258.

1910 Norton PJ. The control of acid mine drainage with wetlands. *Mine Water and*  
1911 *the Environment* 1992; 11: 27-34.

1912 Novotny V. *Diffuse Pollution and Watershed Management*. New York: Wiley &



- 1913 Sons, 2003.
- 1914 Novotny V, Salomons W, Forstner V, Mader P. Diffuse sources of pollution by  
1915 toxic metals and impact on receiving waters. *Heavy Metals, Problems*  
1916 *and Solutions*. Springer, New York, 1995, pp. 33-52.
- 1917 Nuttall CA, Younger PL. Zinc removal from hard circum-neutral mine waters  
1918 using a novel closed-bed limestone reactor. *Water Research* 2000; 34:  
1919 1262-1268.
- 1920 Olias M, Nieto JM, Sarmiento AM, Ceron JC, Canovas CR. Seasonal water  
1921 quality variations in a river affected by acid mine drainage: the Odiel  
1922 River (South West Spain). *Science of the Total Environment* 2004; 333:  
1923 267-281.
- 1924 Ouyang Y, Higman J, Thompson J, O'Toole T, Campbell D. Characterisation  
1925 and spatial distribution of heavy metals in sediment from Cedar and  
1926 Ortega rivers subbasin. *Journal of Contaminant Hydrology* 2002; 54:  
1927 19-35.
- 1928 Owens PN, Walling DE, Leeks GJL. Use of floodplain sediment cores to  
1929 investigate recent historical changes in overbank sedimentation rates  
1930 and sediment sources in the catchment of the River Ouse, Yorkshire,  
1931 UK. *Catena* 1999; 36: 21-47.
- 1932 Oyarzun J, Maturana H, Paulo A, Pasiieczna A. Heavy metals in stream  
1933 sediments from the Coquimbo Region (Chile): Effects of sustained  
1934 mining and natural processes in a semi-arid Andean Basin. *Mine Water*  
1935 *and the Environment* 2003; 22: 155-161.
- 1936 Paquin PR, Gorsuch JW, Apte S, Batley GE, Bowles KC, Campbell PGC,  
1937 Delos CG, Di Toro DM, Dwyer RL, Galvez F, Gensemer RW, Goss  
1938 GG, Hogstrand C, Janssen CR, McGeer JC, Naddy RB, Playle RC,  
1939 Santore RC, Schneider U, Stubblefield WA, Wood KB, Wu. The biotic  
1940 ligand model: a historical overview. *Comparative Biochemistry and*  
1941 *Physiology C-Toxicology & Pharmacology* 2002; 133: 3-35.
- 1942 Perkins WT, Hartley S, Pearce NJG, Dinelli E, Edyvean R, Sandlands L.  
1943 Bioadsorption in remediation of metal mine drainage: The use of  
1944 dealginated seaweed in the BIOMAN project. *Geochimica et*  
1945 *Cosmochimica Acta* 2006; 70: A482-A482.
- 1946 Petersen LBM, Petersen RC. Anomalies in hydropsychid capture nets from  
1947 polluted streams. *Freshwater Biology* 1983; 13: 185-191.
- 1948 Petersen W, Willer E, Williamowski C. Remobilization of trace elements from  
1949 polluted anoxic sediments after resuspension in oxic water. *Water, Air*  
1950 *and Soil Pollution* 1997; 99: 515-22.
- 1951 PIRAMID Consortium. Engineering guidelines for the passive remediation of  
1952 acidic and/or metalliferous mine drainage and similar waste waters.  
1953 University of Newcastle Upon Tyne, Newcastle, 2003.
- 1954 Pirrie D, Power MR, Rollinson G, Camm GS, Huges SH, Butcher AR, et al.  
1955 The spatial distribution and source of arsenic, copper, tin and zinc  
1956 within the surface sediments of the Fal Estuary, Cornwall, UK.  
1957 *Sedimentology* 2003; 50: 579-595.
- 1958 Pollmann O, van Rensburg L, Lange C. Reforestation and Landscaping on  
1959 Mine Tailings. International Mine Water Conference. Water Institute of  
1960 Southern Africa and International Mine Water Association, Pretoria,  
1961 South Africa, 2009, pp. 837-842.
- 1962 Poulton BC, Albert AL, Besser JM, Scmitt CJ, Brumbaugh WG. Fairchild, J.F.

- 1963 A macroinvertebrate assessment of Ozark streams located in lead-zinc  
1964 mining areas of Viburnum Trend in southeastern Missouri, USA.  
1965 Environmental Monitoring and Assessment 2010; 163: 619-641.
- 1966 Pulles W, Heath R. The evolution of passive mine water treatment technology  
1967 for sulphate removal. International Mine Water Conference. Water  
1968 Institute of Southern Africa's Mine Water Division and International  
1969 Mine Water Association, Pretoria, South Africa, 2009, pp. 2-14.
- 1970 Rauret G, Lopez-Sanchez JF, Sahuquillo A, Rubio R, Davidson C, Ure A, et  
1971 al. Improvement of the BCR three step sequential extraction procedure  
1972 prior to certification of new sediment and soil reference materials.  
1973 Journal of Environmental Monitoring 1999; 1: 57-61.
- 1974 Reish DJ, Gerlinger TV. The effects of cadmium, lead and zinc on survival  
1975 and reproduction in the polychaetus annelid *Neanthus*  
1976 *arenaceodentata* (F. Neriedidae). In: Hutchings PA, editor.  
1977 Proceedings of the First International Polychaete Conference. Linear  
1978 Society, Sydney, 1964.
- 1979 Ren J, Packman AI. Stream-subsurface exchange of zinc in the presence of  
1980 silica and kaolinite colloids. Environmental Science and Technology  
1981 2004; 38: 6571-6581.
- 1982 Rhea DT, Harper DD, Farag AM, Brumbaugh WG. Biomonitoring in the  
1983 Boulder River watershed, Montana, USA: metal concentrations in  
1984 biofilm and macroinvertebrates, and relations with macroinvertebrate  
1985 assemblage. Environmental Monitoring and Assessment 2006; 115:  
1986 381-393.
- 1987 Riba I, Garcia-Luque E, Maz-Courrau A, Gonzalez de Canales ML, Delvals  
1988 TA. Influence of salinity in the bioavailability of Zn in sediments of the  
1989 Gulf of Cadiz (Spain). Water, Air and Soil Pollution 2010; 212: 329-336.
- 1990 Rieuwerts JS, Austin S, Harris EA. Contamination from historic metal mines  
1991 and the need for non-invasive remediation techniques: a case study  
1992 from Southwest England. Environmental Monitoring and Assessment  
1993 2009; 148: 149-158.
- 1994 Robb GA. Environmental consequences of coal mine closure. The  
1995 Geographical Journal 1994; 160: 33-40.
- 1996 Robb GA, Robinson JDF. Acid drainage from mines. The Geographical  
1997 Journal 1995; 161: 47-54.
- 1998 Roline RA. The effects of heavy metals pollution of the upper Arkansas River  
1999 on the distribution of aquatic macroinvertebrates. Hydrobiologia 1988;  
2000 160: 3-8.
- 2001 Romero FM, Armienta MA, Gonzalez-Hernandez G. Solid-phase control on  
2002 the mobility of potentially toxic elements in an abandoned lead/zinc  
2003 mine tailings impoundment, Taxco, Mexico. Applied Geochemistry  
2004 2007; 22: 109-127.
- 2005 Roper DS, Nipper MG, Hickey CW, Martin ML, Weatherhead MA. Burial,  
2006 crawling and drifting behaviour of the Bivalve *Macomona liliiana* in  
2007 response to common sediment contaminants. Marine Pollution Bulletin  
2008 1995; 31: 471-478.
- 2009 Rotting TS, Ayora C, Carrera J. Chemical and hydraulic performance of  
2010 "dispersed alkaline substrate" (DAS) for passive treatment of acid mine  
2011 drainage with high metal concentrations. In: Cidu R, Frau F, editors.  
2012 International Mine Water Association Symposium 2007: Water in

- 2013 Mining Environments, Cagliari (Mako Edizioni), 2007, pp. 255-259.
- 2014 Routh J, Ikramuddin M. Thrace-element geochemistry of Onion Creek near  
2015 Van Stone lead-zinc mine (Washington, USA) - chemical analysis and  
2016 geochemical modeling. *Chemical Geology* 1996; 133: 211-224.
- 2017 Salomons W. Adsorption processes and hydrodynamic conditions in  
2018 estuaries. *Environmental Technology Letters* 1980; 1: 356-365.
- 2019 Salomons W. Sediment Pollution in the EEC. Office for Official Publications of  
2020 the European Communities, Luxembourg, 1993.
- 2021 Sanchez J, Marino N, Vaquero MC, Ansorena J, Leqorburu I. Metal pollution  
2022 by old lead-zinc mines in Urumea River Valley (Basque Country,  
2023 Spain). *Soil, biota and sediment. Water, Air, and Soil Pollution* 1998;  
2024 107: 303-319.
- 2025 Sanchez Espana J, Lopez Pamo E, Santofimia Pastor E, Reyes Andres J,  
2026 Martin Rubi JA. The impact of acid mine drainage on the water quality  
2027 of the Odiel River (Huelva, Spain): Evolution of precipitate mineralogy  
2028 and aqueous geochemistry along the Concepcion-Tintillo segment.  
2029 *Water, Air and Soil Pollution* 2006; 173: 121-149.
- 2030 Sanden P, Karlsson S, Duker A, Ledin A, Lundman L. Variations in  
2031 hydrochemistry, trace metal concentration and transport during a rain  
2032 storm event in a small catchment. *Journal of Geochemical Exploration*  
2033 1997; 58: 145-155.
- 2034 Sapsford DJ, Williams KP. Sizing criteria for a low footprint passive mine  
2035 water treatment system. *Water Research* 2009; 43: 423-432.
- 2036 Scanferla P, Ferrari G, Pellay R, Ghirardini AV, Zanetto G, Libralato G. An  
2037 innovative stabilisation/solidification treatment for contaminated soil  
2038 remediation: demonstration of project results. *Journal of Soils and*  
2039 *Sediments* 2009; 9: 229-236.
- 2040 Schmitt CJ, Brumbaugh WG, May TW. Accumulation of metals in fish from  
2041 lead-zinc mining areas of southeastern Missouri, USA. *Ecotoxicology*  
2042 *and Environmental Safety* 2007; 67: 14-30.
- 2043 Schultheis AS, Sanchez M, Hendricks AC. Structural and functional  
2044 responses of stream insects to copper pollution. *Hydrobiologia* 1997;  
2045 346: 85-93.
- 2046 Shannon CE. A mathematical theory of communication. *Bell System*  
2047 *Technical Journal* 1948; 27: 379-423.
- 2048 Smith H, Wood PJ, Gunn J. The influence of habitat structure and flow  
2049 permanence on invertebrate communities in karst spring systems.  
2050 *Hydrobiologia* 2003; 510: 53-66.
- 2051 Smolders AJP, Lock RAC, Van der Velde G, Medina Hoyos RI, Roelofs JGM.  
2052 Effects of mining activities on heavy metal concentrations in water,  
2053 sediment, and macroinvertebrates in different reaches of the Pilcomayo  
2054 River, South America. *Archives of Environmental Contamination and*  
2055 *Toxicology* 2003; 44: 314-323.
- 2056 Sola C, Burgos M, Plazuelo A, Toja J, Plans M, Prat N. Heavy metal  
2057 bioaccumulation and macroinvertebrate community changes in a  
2058 Mediterranean stream affected by acid mine drainage and an  
2059 accidental spill (Guadamar River, SW Spain). *Science of the Total*  
2060 *Environment* 2004; 333: 109-126.
- 2061 Sola C, Prat N. Monitoring metal and metalloid bioaccumulation of  
2062 Hydropsyche (Trichoptera, Hydropsychidae) to evaluate metal pollution

2063 in a mining river. Whole body versus tissue content. *Science of the*  
2064 *Total Environment* 2006; 359: 221-231.

2065 Spehar RL, Anderson RL, Fiandt JT. Toxicity and bioaccumulation of  
2066 cadmium and lead in aquatic invertebrates. *Environmental Pollution*  
2067 1978; 15: 195-208.

2068 Stiff MJ. The chemical states of copper in polluted fresh water and a scheme  
2069 of analysis to differentiate them. *Water Research* 1971; 5: 585-599.

2070 Stoertz MW, Bourne H, Knotts C, White MM. The effects of isolation and acid  
2071 mine drainage on fish and macroinvertebrate communities of Monday  
2072 Creek, Ohio, USA. *Mine Water and the Environment* 2002; 21: 60-72.

2073 Stone M, Droppo IG. In-channel surficial fine grained sediment laminae. Part  
2074 II: Chemical characteristics and implications for contaminant transport  
2075 in fluvial systems. *Hydrological Processes* 1994; 8: 113-124.

2076 Sullivan AB, Drever JI. Spatiotemporal variability in stream chemistry in a  
2077 high-elevation catchment affected by mine drainage. *Journal of*  
2078 *Hydrology* 2001; 252: 237-250.

2079 Tordoff GM, Baker AJM, Willis AJ. Current approaches to the revegetation  
2080 and reclamation of metalliferous mine wastes. *Chemosphere* 2000; 41:  
2081 219-228.

2082 Taylor MP. The variability of heavy metals in floodplain sediments: a case  
2083 study from mid Wales. *Catena* 1996; 28: 71-87.

2084 Taylor MP, Mackay AK, Hudson-Edwards KA, Holz E. Soil Cu, Pb and Zn  
2085 contaminants around Mount Isa city, Queensland, Australia: potential  
2086 sources and risks to human health. *Applied Geochemistry* 2010; 25:  
2087 841-855.

2088 Tessier A, Campbell PGC, Bisson M. Sequential extraction procedure for the  
2089 speciation of particulate trace metals. *Analytical Chemistry* 1979; 7: 41-  
2090 54.

2091 Tokalioglu S, Kartal S, Birol G. Application of a three-stage sequential  
2092 extraction procedure for the determination of extractable metal  
2093 contents in highway soils. *Turkish Journal of Chemistry* 2003; 27: 333-  
2094 346.

2095 Van Damme A, Degryse F, Smolders E, Sarret G, Dewit J, Swennen R, et al.  
2096 Zinc speciation in mining and smelter contaminated overbank  
2097 sediments by EXAFS spectroscopy. *Geochimica et Cosmochimica*  
2098 *Acta* 2010; 74: 3707-3720.

2099 Van Damme PA, Harnel C, Ayala A, Bervoets L. Macroinvertebrate  
2100 community response to acid mine drainage in rivers of the High Andes  
2101 (Bolivia). *Environmental Pollution* 2008; 156: 1061-1068.

2102 Vasile GD, Vladescu L. Cadmium partition in river sediments from an area  
2103 affected by mining activities. *Environmental Monitoring and*  
2104 *Assessment* 2010; 167: 349-357.

2105 Vermeulen AC. Elaborating chironomid deformities as bioindicators of toxic  
2106 sediment stress: The potential application of mixture toxicity concepts.  
2107 *Annales Zoologici Fennici* 1995; 32: 265-285.

2108 Vermeulen AC, Liberloo G, Dumont P, Ollevier F, Goddeeris B. Exposure of  
2109 *Chironomus riparius* larvae (diptera) to lead, mercury and beta-  
2110 sitosterol: effects on mouthpart deformation and moulting.  
2111 *Chemosphere* 2000; 41: 1581-1591.

2112 Vuori K-M. Rapid behavioural and morphological responses of hydropsychid

- 2113 larvae (trichoptera, hydropsychidae) to sublethal cadmium exposure.  
2114 Environmental Pollution 1994; 84: 291-299.
- 2115 Vinod VTP, Sashidar RB, Sukumar AA. Competitive adsorption of toxic heavy  
2116 metal contaminants by gum kondagogu (*Cochlospermum gossypium*):  
2117 A natural hydrocolloid. Colloids and Surfaces B: Biointerfaces 2010;  
2118 75: 490-495.
- 2119 Vivian CMG, Massie KS. Trace metals in waters and sediments of the River  
2120 Tawe, South Wales, in relation to local sources. Environmental  
2121 Pollution 1977; 14: 47-61.
- 2122 Walling DE, Owens PN, Carter J, Leeks GJL, Lewis S, Meharg AA, et al.  
2123 Storage of sediment-associated nutrients and contaminants in river  
2124 channel and floodplain systems. Applied Geochemistry 2003; 18: 195-  
2125 220.
- 2126 Walling DE, Webb BW. The spatial dimension in the interpretation of stream  
2127 solute behaviour. Journal of Hydrology 1980; 47: 129-149.
- 2128 Warburton J, Danks M, Wishart D. Stability of an upland gravel-bed stream,  
2129 Swinhope Burn, Northern England. Catena 2002; 49: 309-329.
- 2130 Watanabe NC, Harada S, Komai Y. Long-term recovery from mine drainage  
2131 disturbance of a macroinvertebrate community in the Ichi-kawa River,  
2132 Japan. Hydrobiologia 2000; 429: 171-180.
- 2133 Watzlaf G, Schroeder K, Kairies C. Long-term performance of anoxic  
2134 limestone drains. Mine Water and the Environment 2000; 19: 98-110.
- 2135 Waygood CG, Ferreira S. A review of the current strategy for capping mining  
2136 spoils. International Mine Water Conference. Water Institute of  
2137 Southern Africa and International Mine Water Association, Pretoria,  
2138 South Africa, 2009, pp. 738-745.
- 2139 Webb BW, Walling DE. Stream solute behaviour in the River Exe basin,  
2140 Devon, UK. Dissolved loads of Rivers and Surface Water  
2141 Quality/Quantity Relationships. International Association of  
2142 Hydrological Sciences Publication No. 141, 1983, pp. 153-169.
- 2143 Whitehead PG, Prior H. Bioremediation of acid mine drainage: An introduction  
2144 to the Wheal Jane wetlands project. Science of the Total Environment  
2145 2005; 338: 15-21.
- 2146 Wilby RL, Orr HG, Hedger M, Forrow D, Blackmore M. Risks posed by climate  
2147 change to the delivery of Water Framework Directive objectives in the  
2148 UK. Environment International 2006; 32: 1043-1055.
- 2149 Wilkin RT. Contaminant attenuation processes at mine sites. Mine Water and  
2150 the Environment 2008; 27: 251-258.
- 2151 Willis M. Analysis of the effects of zinc pollution on the macroinvertebrate  
2152 populations of the Afon Crafnant, North Wales. Environmental  
2153 Geochemistry and Health 1985; 7: 98-109.
- 2154 Winner RW, Boesel MW, Farrell MP. Insect community structure as an index  
2155 of heavy-metal pollution in lotic ecosystems. Canadian Journal of  
2156 Fisheries and Aquatic Sciences 1980; 37: 647-655.
- 2157 Wirt L, Leib KJ, Bove DJ, Mast MA, Evans JB, Meeker GP. Determination of  
2158 chemical-constituent loads during base-flow and storm-runoff  
2159 conditions near historical mines in Prospect Gulch, Upper Animas  
2160 River Watershed, Southwestern Colorado. U.S. Geological Survey,  
2161 1999.
- 2162 Wolfenden PJ, Lewin J. Distribution of metal pollutants in floodplain

2163 sediments. *Catena* 1977; 4: 317.

2164 Wolfenden PJ, Lewin J. Distribution of metal pollutants in active stream  
2165 sediments. *Catena* 1978; 5: 67-78.

2166 Wolkersdorfer C. Mine water literature in ISI's Science Citation Index  
2167 expanded. *Mine Water and the Environment* 2004; 23: 96-99.

2168 Wolz JEA, et al. In search for the ecological and toxicological relevance of  
2169 sediment re- mobilisation and transport during flood events. *Journal of*  
2170 *Soils and Sediments* 2009; 9: 1-5.

2171 Wood PJ, Armitage PD. Biological effects of fine sediment in the lotic  
2172 environment. *Environmental Management* 1997; 21: 203-217.

2173 Woodcock TS, Huryn AD. The response of macroinvertebrate production to a  
2174 pollution gradient in a headwater stream. *Freshwater Biology* 2007; 52:  
2175 177-196.

2176 Yi Y, Wang Z, Zhang K, Yu G, Duan X. Sediment pollution and its effect on  
2177 fish through food chain in the Yangtze River. *International Journal of*  
2178 *Sediment Research* 2008; 23: 338-347.

2179 Yim JH, Kim KW, Kim SD. Effect of hardness on acute toxicity of metal  
2180 mixtures using *Daphnia magna*: prediction of acid mine drainage  
2181 toxicity. *Journal of Hazardous Materials* 2006; B138: 16-21.

2182 Yim WWS. Geochemical investigations on fluvial sediments contaminated by  
2183 tin mine tailings, Cornwall, England. *Environmental Geology* 1981; 3:  
2184 245-256.

2185 Younger PL. The adoption and adaptation of passive treatment technologies  
2186 for mine waters in the United Kingdom. *Mine Water and the*  
2187 *Environment* 2000; 19: 84-97.

2188 Younger PL. Coalfield closure and the water environment in Europe.  
2189 *Transactions of the Institution of Mining and Metallurgy (Section A:*  
2190 *Mining Technology)* 2002; 111: A201-A209.

2191 Younger PL, Banwart SA, Hedin RS. *Mine Water. Hydrology, Pollution,*  
2192 *Remediation.* Dordrecht: Kluwer Academic Publishers, 2002.

2193 Younger PL, Coulton RH, Frogatt EC. The contribution of science to risk-  
2194 based decision-making: lessons from the development of full-scale  
2195 treatment measures for acidic mine waters at Wheal Jane, UK. *Science*  
2196 *of the Total Environment* 2005; 338: 138-154.

2197 Younger PL, Wolkersdorfer CH. *Mining Impacts on the Fresh Water*  
2198 *Environment: Technical and Managerial Guidelines for Catchment*  
2199 *Scale Management.* *Mine Water and the Environment* 2004;  
2200 Supplement to Volume 23: S1-S80.

2201 Ziemkiewicz P, Skousen J, Brant D, Sterner P, Lovett R. Acid mine drainage  
2202 treatment with armoured limestone in open limestone channels.  
2203 *Journal of Environmental Quality* 1997; 26: 560-569.


2204 Zoumis T, Schmidt A, Grigorova L, Calmano W. Contaminants in sediments:  
2205 remobilisation and demobilisation. *Science of the Total Environment*  
2206 2001; 226: 195-202.

2207

2208

2209

2210



2211

2212

2213 **Figure 1** The process of pyrite weathering in a deep metal mine. Four general

2214 equations describe the chemistry of pyrite weathering and the production of

2215 AMD – (1) The oxidation of pyrite by oxygen and water in atmospheric

2216 conditions to produce dissolved ferrous iron and sulphuric acid; (2) the

2217 oxidation of dissolved ferrous iron to ferric iron; (3) the hydrolysis of ferric iron

2218 with water to produce iron hydroxide precipitate (ochre) and acidity; (4) the

2219 oxidation of additional pyrite by the ferric iron generated in reaction (2) to

2220 produce dissolved ferrous iron and sulphuric acid. The acidic conditions

2221 generated during these processes can dissolve oxidised trace metals. The

2222 process is accelerated by the presence of sulphide and iron-oxidising

2223 bacteria.

2224

2225 **Table 1** A comparison of dissolved metal (mg/l), sulphate (mg/l) and pH concentrations from waters impacted by historical deep  
 2226 metal mining.

Location	Sample type	Pb	Zn	Cu	Cd	Fe	SO <sub>4</sub>	pH	Author(s)
<i>Europe</i>									
River Carnon, England	Mine drainage	<0.01 - 0.02	0.12 - 23	0.02 - 1.3	<0.01 - 0.02	<0.01 - 49	77 - 789	3.3 - 7.7	Neal et al. (2005)
River Tamar, England	Adit drainage	<0.01 - 0.17	<0.1 - 2.5	<0.01 - 1.4	<0.01 - 0.01	0.05 - 2.6	10 - 89	3.4 - 7.8	Mighanetara et al. (2009)
Funtana Raminosa District, Italy	Mining Tailings drainage	<0.01	0.08 - 34	<0.01 - 0.04	<0.01 - 0.85	0.02 - 0.25	22 - 1680	7.1 - 7.8	Cidu and Mereu (2007)
Buchim Mining district, Macedonia	Mine stream	0.03*	0.03*	0.62*	<0.01*	0.3*	-	5.1*	Alderton et al. (2005)
Zletovo Mining District, Macedonia	Adit drainage	0.06*	21.57*	0.46*	0.14*	98.2*	-	3.4*	Alderton et al. (2005)
River Zletovska, Macedonia	Channel	<0.03 - 0.8	0.04 - 70.07	<0.01 - 1.05	<0.01 - 0.24	0.1 - 103.3	-	3.4 - 7.6	Alderton et al. (2005)
River Bjorgasen, Norway	Channel	-	5.4 <sup>b</sup>	2.7 <sup>b</sup>	0.01 <sup>b</sup>	-	-	3.2 <sup>b</sup>	Gundersen and Stienes (2001)
Rio Tinto, Spain	Channel	0.1 - 2.4	0.3 - 420	0.05 - 240	-	-	2800 - 16000	1.4 - 7.6	Hudson-Edwards et al. (1999b)
Troya Mine, Spain	Tailings pond	0.02 - 0.05	4.99 - 18.95	<0.01 - 0.03	0.01 - 0.03	0.04 - 0.33	-	-	Marques et al. (2001)
River Odiel, Spain	Channel	<0.01 - 1.18	0.17 - 130.23	0.01 - 37.62	<0.01 - 0.38	0.03 - 262.71	50.7 - 3960	2.5 - 6.3	Olias et al. (2004)
Tintillo River, Spain	Mine drainage	0.01 - 0.07	7.3 - 216	3.5 - 115	<0.01 - 0.51	264 - 1973	1300 - 11580	2.3 - 2.8	Sanchez Espana et al. (2006)
Tinto Santa Rosa Mine, Spain	Mine drainage	<0.01 - 0.08	56 - 85	15 - 23	0.09 - 0.15	234 - 881	2704 - 4026	2.6 - 3.4	Asta et al. (2007)



Fluminese Mining District, Spain	Mine water	<0.01 - 0.05	0.88 - 40	-	<0.01 - 0.09	<0.01 - 12	17 - 640	6.3 - 8.2	<i>Cidu et al. (2007)</i>
River Tawe, Wales	Channel	<0.01 - 0.15	0.01 - 8.8	<0.01 - 0.04	<0.01 - 0.16	-	-	-	<i>Vivian and Massie (1977)</i>
River Rheidol, Wales	Channel	<0.01	0.08 - 0.29	-	<0.01	-	5.3 - 7.1	5.5 - 6.4	<i>Fuge et al. (1991)</i>
River Yswyth, Wales	Channel	0.06 - 0.09	0.17 - 0.36	-	<0.01	-	nd - 5.3	4.1 - 4.6	<i>Fuge et al. (1991)</i>
Cwm Rheidol Mine, Wales	Adit drainage	0.02 - 0.04	38 - 72	0.03 - 0.07	0.04 - 0.11	-	441 - 846	2.8 - 3.0	<i>Fuge et al. (1991)</i>
Cwm Ystwyth Mine, Wales	Spoil drainage	0.29 - 3.3	1.5 - 4.6	<0.01	<0.01	-	nd	4.1*	<i>Fuge et al. (1991)</i>
Cae Coch Pyrite Mine, Wales	Mine water	-	-	-	-	2261 <sup>b</sup>	6590 <sup>b</sup>	2.4 <sup>b</sup>	<i>McGinness and Johnson (1993)</i>
River Goch, Wales	Channel	-	<0.01 - 4.19	<0.01 - 5.99	-	<0.01 - 25.98	-	2.3 - 7.7	<i>Boult et al. (1994)</i>
Cwm Rheidol Mine, Wales	Spoil drainage	-	577 - 978	1.2 - 9.35	-	-	-	2.6 - 2.7	<i>Johnson (2003)</i>
<i>North America</i>									
West Squaw Creek, USA	Channel	-	0.01 - 156	<0.01 - 190	-	0.03 - 500	2.6 - 5100	2.4 - 6.9	<i>Filipek et al. (1987)</i>
Richmond Mine, USA	Mine water	1 - 120	0.06 - 23.5 <sup>a</sup>	0.21 - 4.76 <sup>a</sup>	4 - 2110	2.47 - 79.7 <sup>a</sup>	14 - 760 <sup>a</sup>	-3.6 - 1.5	<i>Nordstrom et al. (2000)</i>
Peru Creek, USA	Channel	-	0.55 - 1.89	0.05 - 0.22	-	0.08 - 0.5	29.6 - 73	4.7 - 5.9	<i>Sullivan and Drever (2001)</i>
Boulder Creek, USA	Channel	<0.032*	0.469*	0.246*	<0.01*	2.82*	97.4*	3.3*	<i>Keith et al. (2001)</i>
Black Foot River, USA	Channel	-	<0.2 - 535	<0.8 - 4	<0.5 - 2.6	<5 - 37	5.5 - 88.8	7.3 - 8.8	<i>Nagorski et al. (2002)</i>
Phillips Mine, USA	Channel	<0.01	<0.01 - 0.17	0.02 - 3.13	-	0.16 - 42.4	25 - 368	2.3 - 6.5	<i>Gilchrist et al. (2009)</i>
<i>Australasia</i>									
River Dee, Australia	Channel	<0.01 - 0.6	<0.01 - 10.4	<0.01 - 45.03	-	<0.01 - 74	340 - 5950	2.7 - 7.0	<i>Edraki et al. (2005)</i>

---

Mt. Morgan Mine, Australia	Open pit	1.51*	21.97*	44.54*	-	253*	13600*	2.7*	<i>Edraki et al. (2005)</i>
----------------------------	----------	-------	--------	--------	---	------	--------	------	-----------------------------

---

nd = not detectable. \* single observation. <sup>a</sup> grams per litre. <sup>b</sup> mean value

---

2227

2228

2229

2230

2231

2232

2233

2234

2235

2236

2237

2238

2239

2240 **Table 2** Comparison of metal concentrations (mg/kg) in channel and floodplain sediments from historic deep metal mining impacted  
 2241 rivers.

River location	Geomorphic-type site	Grain size fraction	Metal phase extracted	Pb	Zn	Cu	Cd	Author(s)
<i>Europe</i>								
Red River, England	Channel	<2000 µm	Total	nd - 120	nd - 630	nd - 1320	-	<i>Yim (1981)</i>
River Derwent, England	Channel	<1000 µm	Total	96 - 3120	82 - 2760	-	0.6 - 13.8	<i>Burrows and Whitton (1983)</i>
River Derwent, England	Floodplain	<2000 µm	Total	131 - 1179	<10 - 1696	2.9 - 64	0.08 - 12.5	<i>Bradley and Cox (1990)</i>
River Tyne, England	Floodplain	<2000 µm	Total	615 - 2340	722 - 2340	11 - 42.5	2.6 - 8	<i>Macklin et al. (1992)</i>
River Swale, England	Floodplain	<63 µm	Total	56 - 5507	15 - 3066	-	1 - 18	<i>Macklin et al. (1994)</i>
River Allen, England	Channel	<170 µm	Total	2330*	1410*	-	-	<i>Goodyear et al. (1996)</i>
River Severn, England	Floodplain	<2000 µm	Total	23 - 204	173 - 936	30 - 67	0.35 - 6.4	<i>Taylor (1996)</i>
River Tees, England	Channel	<2000 µm	Total	522 - 6880	404 - 1920	20 - 77	0.95 - 5.95	<i>Hudson-Edwards et al. (1997)</i>
River Aire, England	Channel	<63 µm	Total	90 - 237	274 - 580	118 - 198	-	<i>Walling et al. (2003)</i>
River Swale, England	Floodplain	<63 µm	Total	10000*	14000*	-	7500*	<i>Dennis et al. (2003)</i>
River Calder, England	Channel	<63 µm	Total	199 - 343	397 - 907	141 - 235	-	<i>Walling et al. (2003)</i>
River Wear, England	Channel	<150 µm	Total	20 - 15000	40 - 1500	<10 - 340	-	<i>Lord and Morgan (2003)</i>
Dale Beck, England	Channel	<2000 µm	Total	13693*	442*	206*	-	<i>Geer (2004)</i>
River Avoca, Ireland	Channel	<1000 µm	Total	-	1520 <sup>a</sup>	674 <sup>a</sup>	-	<i>Herr and Gray (1996)</i>
River Mala Panew, Poland	Channel	<63 µm	Total	36 - 3309	126 - 11153	3.97 - 483	0.18 - 559	<i>Aleksander-Kwaterczak and Helios-</i>

---

									<i>Rybicka (2009)</i>
River Somes, Romania	Channel	<2000 µm	Total	28 - 6800	64 - 19600	12 - 8400	0.8 - 110		<i>Macklin et al. (2003)</i>
River Viseu, Romania	Floodplain	<2000 µm	total	17 - 850	110 - 2760	32 - 1000	0.5 - 17		<i>Macklin et al. (2005)</i>
Gezala Creek, Spain	Channel	<177 µm	Total	10.6 - 37630	216 - 25676	2.7 - 1691	0.22 - 45		<i>Marques et al. (2001)</i>
River Tinto, Spain	Channel	<2000 µm	Total	3200 - 16500	600 - 67300	1800 - 26500	<1 - 23		<i>Galan et al. (2003)</i>
River Odiel, Spain	Channel	<2000 µm	Total	1900 - 16600	1000 - 74600	3500 - 20900	1.4 - 10.2		<i>Galan et al. (2003)</i>
River Rheidol, Wales	Floodplain	<2000 µm	Total	291 - 2098	242 - 630	21 - 85	0.08 - 3.5		<i>Davies and Lewin (1974)</i>
River Tawe, Wales	Channel	<2000 µm	Total	63 - 6993	20 - 31199	34 - 2000	2 - 335		<i>Vivian and Massie (1977)</i>
River Rheidol, Wales	Floodplain	<210 µm	Total	813 - 1717	201 - 383	33 - 120	-		<i>Wolfenden and Lewin (1977)</i>
River Towy, Wales	Channel	<2000 µm	Total	36 - 5732	106 - 3722	44 - 259	0.78 - 83		<i>Wolfenden and Lewin (1978)</i>
River Twymyn, Wales	Channel	<2000 µm	Total	593 - 6411	159 - 6955	44 - 2557	1.5 - 44		<i>Wolfenden and Lewin (1978)</i>
River Ystywth, Wales	Floodplain	<2000 µm	Total	73 - 4646	123 - 1543	-	-		<i>Lewin et al. (1983)</i>
River Twymyn, Wales	Channel	<63 µm	Non-residual	1.1 - 2914	0.7 - 148	0.3 - 30	<0.01 - 0.9		<i>Byrne et al. (2010)</i>
<i>North America</i>									
West Squaw Creek, USA	Channel	<177 µm	Total	-	32 - 5940	254 - 4090	-		<i>Filipek et al. (1987)</i>
Black Foot River, USA	Channel	<63 µm	Total	1100 - 8700	1700 - 9600	1400 - 9900	<1 - 115		<i>Nagorski et al. (2002)</i>
River Cedar, USA	Channel	-	Total	4.5 - 420	9.75 - 2050	2.3 - 107	0.07 - 3.8		<i>Ouyang et al. (2002)</i>
Copper Mine Brook, USA	Channel	<1000 µm	Total	9.9 - 30	9 - 67	31 - 398	-		<i>Gilchrist et al. (2009)</i>
<i>Australasia</i>									
River Kangjiaxi, China	Channel	-	Non-residual	1154 - 8034	124 - 2319	23 - 209	2.6 - 41		<i>Licheng and Guiju (1996)</i>

---

---

nd = not detectable. \* maximum value. <sup>b</sup> mean value.

---

2242

2243

2244

2245

2246

2247

2248

2249

2250

2251

2252

2253

2254

2255 **Table 3** Impacts of metal mine drainage on instream macroinvertebrates reported within the scientific literature. Types of studies  
 2256 are - <sup>a</sup> stream survey, <sup>b</sup> microcosm experiment and <sup>c</sup> laboratory bioassay.

Primary impact reported	Additional information	Author(s)
<i>Community composition</i>		
Shift in community structure	Clean sites dominated by Ephemeroptera and Plecoptera; moderately contaminated sites dominated by Plecoptera and Diptera; and heavily contaminated sites dominated by Diptera	Armitage (1980) <sup>a</sup>
	Clean sites dominated by Ephemeroptera; moderately contaminated sites by Tricoptera; and heavily contaminated sites dominated by Diptera	Winner <i>et al.</i> (1980) <sup>a</sup>
	Contaminated sites dominated by Orthocladiinae (Chironomidae) and species of net-spinning Tricoptera	Clements <i>et al.</i> (1992) <sup>a</sup>
	Contaminated sites dominated by Chironomidae	Gray (1998) <sup>a</sup>
	Ephemeroptera reduced by > 75% in moderately contaminated streams	Clements <i>et al.</i> (2000) <sup>a</sup>
	Clean sites dominated by Stenopsychidae (Trichoptera); contaminated sites dominated by Chironomidae and <i>Epeorus latifolium</i> (Ephemeroptera)	Watanabe <i>et al.</i> (2000) <sup>a</sup>
	Contaminated sites dominated by Chironomidae, Tubificidae, Baetidae and Simuliidae	Marques <i>et al.</i> (2003) <sup>a</sup>
	Heavily contaminated sites dominated by Chironomidae	Smolders <i>et al.</i> (2003) <sup>a</sup>
	Dominance of predators in very acidic mining sites	Gerhardt <i>et al.</i> (2004) <sup>a</sup>
	Heavily contaminated sites characterised by high proportion of Chironominae and predatory Tanypodinae	Janssens de Bisthoven <i>et al.</i> (2005) <sup>a</sup>
Decrease in abundance	Reduction in abundance recorded	Willis (1985) <sup>a</sup> , Gray (1998) <sup>a</sup> , Hirst <i>et al.</i> (2002) <sup>a</sup>
	Ephemeroptera comprised less than 5% of individuals at one location	Clements <i>et al.</i> (1992) <sup>a</sup>

---

	Abundance significantly lower in experiments with metal mixtures and high predation pressure	Kiffney (1996) <sup>b</sup>
	Abundance positively related to stream alkalinity and pH	Malmqvist and Hoffsten (1999) <sup>a</sup>
	Ephemeroptera and Plecoptera particularly affected	Clements (2004) <sup>b</sup>
Decrease in number of taxa	Reduced number of taxa recorded	Willis (1985) <sup>a</sup> , Kiffney (1996) <sup>b</sup> , Gray (1998) <sup>a</sup>
	Decrease most pronounced in low flow conditions	Clements <i>et al.</i> (1992) <sup>a</sup>
Decrease in EPT taxa	EPT richness positively related to stream pH	Malmqvist and Hoffsten (1999) <sup>a</sup>
	Near extinction of mayfly species	Hickey and Golding (2002) <sup>a</sup>
	Reduced number of EPT taxa recorded	Gerhardt <i>et al.</i> (2004) <sup>a</sup>
Decrease in species diversity	Reduced species diversity recorded	Amisah and Cowx (2000) <sup>a</sup> , Hirst <i>et al.</i> (2002) <sup>a</sup>
	Dominance of Chironomidae	Smolders <i>et al.</i> (2003) <sup>a</sup>
	Dominance of Chironomidae, Baetidae and Simuliidae	Van Damme <i>et al.</i> (2008) <sup>a</sup>
Impaired ecosystem function	Microbial colonisation of leaf material and leaf decomposition inhibited by high Cd concentrations	Giesy <i>et al.</i> (1978) <sup>b</sup>
	Microbial activity and leaf decomposition rates significantly lower at contaminated sites	Carpenter <i>et al.</i> (1983) <sup>a</sup>
	Secondary production of shredders negatively associated with metal contamination; leaf decomposition rates decreased; microbial respiration decreased	Carlisle and Clements (2005) <sup>a</sup>
	Reduced secondary production and organic matter storage	Woodcock and Huryn (2007) <sup>a</sup>
	Greater vulnerability of net-spinning Tricoptera to predation possibly due to spending more time in the open repairing capture nets	Clements <i>et al.</i> (1989) <sup>b</sup>

Physiological response	Differences in metal sensitivity related to trophic status; herbivores and detritivores more sensitive than predators	Leland <i>et al.</i> (1989) <sup>a</sup>
	Decrease in reproduction rates of <i>Gammurus pulex</i> (Gammaridae)	Maltby and Naylor (1990) <sup>c</sup>
	Differences in sensitivity related to trophic status; reduced leaf decomposition rates suggests shredders sensitive to pollution	Schultheis <i>et al.</i> (1997) <sup>a</sup>
	Increase incident of deformity (mentum structure) in <i>Chironomus riparius</i> (Chironomidae)	Groenendijk <i>et al.</i> (1998) <sup>a</sup>
	Increased incident of deformity (mentum structure) and decreased moulting success in <i>Chironomus riparius</i> (Chironomidae)	Vermeulen <i>et al.</i> (2000) <sup>c</sup>
	pH-dependent decrease in locomotion of <i>Atyaephyra desmaersti</i> (Crustacea) in AMD solutions	Gerhardt <i>et al.</i> (2004) <sup>c</sup>
	Locomotion and ventilation of <i>Choroterpes picteti</i> (Leptophlebiidae) greater in acid only solutions than in AMD solutions	Gerhardt <i>et al.</i> (2005a) <sup>c</sup>
	pH-dependent increase in locomotion and ventilation of <i>Gambusia holbrooki</i> (Crustacea) in AMD solutions	Gerhardt <i>et al.</i> (2005b) <sup>c</sup>
	pH-dependent decrease in locomotion and ventilation of <i>Atyaephyra desmaersti</i> (Crustacea) in AMD solutions	Janssens De Bisthoven <i>et al.</i> (2006) <sup>c</sup>
	Contaminated water causes higher locomotory activity in <i>Lumbriculus variegatus</i> (Oligochaeta) than contaminated sediment	Gerhardt (2007) <sup>c</sup>
	Decrease in pH and increase in dissolved metals caused decrease in locomotion and inhibition of feeding rate in <i>Echinogammarus meridionalis</i> (Crustacea)	Macedo-Sousa <i>et al.</i> (2007) <sup>c</sup>
	Pulse of AMD caused early warning responses in <i>Echinogammarus meridionalis</i> (Crustacea) consisting of increased locomotion and subsequent increase in ventilation	Macedo-Sousa <i>et al.</i> (2008) <sup>c</sup>
	Average daily moulting rate of <i>Rithrogena hageni</i> (Heptageniidae) decreased after exposure to aqueous copper, cadmium and zinc	Brinkman and Johnston (2008) <sup>c</sup>
Behavioural response	Anomalies in capture nets of Hydropsychidae	Petersen and Petersen (1983) <sup>a</sup>



	Decrease in burrowing rates and increase in crawling and drifting rates of <i>Macomona liliiana</i> (Bivalve)	Roper <i>et al.</i> (1995) <sup>c</sup>
Morphological deformities	Cross-breeding of <i>Chironomous riparius</i> (Chironomidae) from contaminated and clean rivers revealed some level of genetic adaptation to metals in offspring	Groenendijk <i>et al.</i> (2002) <sup>b</sup>
	Macroinvertebrate drift and respiration significant correlated with metal concentrations	Clements (2004) <sup>b</sup>
	Increased incident of adult and larval deformities in <i>Chironomous tentans</i> (Chironomidae)	Martinez <i>et al.</i> (2004) <sup>c</sup>
	Decreased locomotory activity of <i>Chironomous</i> sp. (Chironomidae) in AMD solutions	Janssens De Bisthoven <i>et al.</i> (2004) <sup>c</sup>
Metal bioaccumulation	Younger instars had higher metal concentrations than older instars	Krantzberg (1989) <sup>c</sup>
	Concentration of metals in Ephemeropteran species decreased in consecutive larval stages	Jop (1991) <sup>c</sup>
	Metal bioaccumulation dependent on feeding group; shredders and scrapers accumulated the highest metal concentrations (biofilm contained more metals than sediments)	Farag <i>et al.</i> (1998) <sup>a</sup>
	Whole-body metal concentrations of <i>Hydropsyche</i> sp. (Hydropsychidae) greater in species exposed to dissolved metals than in species exposed to AMD precipitates	DeNicola and Stapleton (2002) <sup>b</sup>
	<i>Chironomus februarius</i> (Chironomidae) exhibited adaptation to and tolerance of metal-polluted sediments	Bahrndorff <i>et al.</i> (2006) <sup>b</sup>
	Macroinvertebrate metrics significantly correlated with metals in biofilm, suggesting biofilm is a better index than macroinvertebrates for monitoring metal impacts on aquatic systems	Rhea <i>et al.</i> (2006) <sup>a</sup>
	Whole-body metal concentrations of <i>Hydropsyche</i> sp. (Hydropsychidae) were strongly positively correlated with metal concentrations in water and sediment	Sola and Prat (2006) <sup>a</sup>
<i>Effects of environmental parameters on the toxicity of metal mine discharges</i>		
Water hardness and alkalinity	Increased water hardness and alkalinity reduces metal toxicity in <i>Chironomous tentans</i> (Chironomidae)	Gauss <i>et al.</i> (1985) <sup>c</sup>
	Increasing water hardness reduces community sensitivity to metal contamination	Gower <i>et al.</i> (1994) <sup>a</sup>
	Increased water hardness reduces metal toxicity in <i>Daphnia magna</i> (Dapniidae)	Yim <i>et al.</i> (2006) <sup>c</sup>

---

Metal mixtures	Abundance of heptageniidae, community respiration and macroinvertebrate drift were more sensitive to metal mixtures than single metal solutions	Hickey and Golding (2002) <sup>c</sup>
	Survival of <i>Diplostomum spathaceum</i> (Diplostomatidae) greater in metal mixtures than in single metal solutions	Morley <i>et al.</i> (2002) <sup>c</sup>
	Community sensitivity greatest in combined metal mixtures compared to single metal solutions	Clements (2004) <sup>b</sup>
Other parameters	Increased turbidity reduces metal toxicity to Cladocera by decreasing bioavailability of metals	Garcia-Garcia and Nandini (2006) <sup>c</sup>
	Inverse correlation between salinity and lesion index of gills in <i>Ruditapes philippinarum</i> (Bivalvia)	Riba <i>et al.</i> (2010) <sup>c</sup>

---

2257

2258

2259

2260

2261

2262

2263

2264

2265

2266

2267

2268 **Table 4** Typology of common passive mine water treatment units and source control techniques: indicating the nature of mine  
 2269 water drainage and the principal advantages and limitations of each method.

Name	Mine water type	Brief description	Advantages	Limitations	Example reference(s)
<i>Passive mine water treatment technologies</i>					
Aerobic wetlands	Net alkaline ferruginous	A system of shallow ponds, cascades and vegetated substrate encourage aeration of mine waters and oxidation, hydrolysis and precipitation of some heavy metals (mainly Fe and Al)	Efficient Fe and Al removal; low maintenance requirement; cost-effective; easy integration into landscape and connection with existing ecosystems	Not suitable for highly toxic, sulphate-rich and acidic mine waters; large land surface area requirement; occasional removal of substrate precipitates required	Robb and Robinson (1995); Johnson and Hallberg (2005)
Anaerobic wetlands	Net acidic ferruginous with high sulphate concentrations	A thick anoxic substrate of saturated organic material neutralises acidity and generates alkalinity through processes of bacterial	Often used to neutralise acidity and generate alkalinity prior to discharge to aerobic wetlands; efficient Fe and sulphate removal; some toxic	Not suitable for high toxic metal concentrations (especially Zn and Cd); large land surface area requirement; occasional	Younger <i>et al.</i> (2002); Johnson and Hallberg (2005)

---

sulphate reduction and calcite dissolution; heavy metals (mainly Fe and Al) are removed as precipitates  
 metals are removed through precipitation of sulphides and adsorption to organic matter; low maintenance requirement; cost-effective; easy integration into landscape and connection with existing ecosystems  
 removal of substrate precipitates required; requires high sulphate (>100 mg/l) concentrations; often produce hydrogen sulphide gas

Anoxic Limestone Drains (ALDs) Net acidic, low Al and Fe, low dissolved oxygen concentrations Mine water is routed into a buried limestone trench which neutralises acidity and generates alkalinity Often used to neutralise acidity and generate alkalinity prior to discharge to aerobic wetlands; efficient Fe and Al removal at low concentrations (<2 mg/l) Not suitable for high toxic metal mine waters; vulnerable to precipitation of Al and Fe on limestone; only suitable for mine waters above pH 5 with low ferric Fe, Al (<2 mg/l) and dissolved oxygen content (<1 mg/l) Nuttall and Younger (2000); Watzlaf *et al.* (2000)

Oxidic Limestone Drains Net acidic, low to An open (exposed to the Often used to neutralise Not suitable for high toxic Ziemkiewicz *et al.* (1997)

---

---

(OLDs)	moderate sulphate	atmosphere)	limestone	acidity and generate alkalinity	metal mine waters; high flow	
		trench which	neutralises	prior to discharge to aerobic	velocities required to prevent	
		acidity and	generates	wetlands; good rates of	Fe and Al precipitation on the	
		alkalinity		alkalinity generation with low	limestone	
				water residence times; easy		
				to construct and low cost		
				alternative to more		
				technically challenging and		
				costly systems		
Reducing	and	Net acidic	A layer of limestone beneath	Often used to neutralise	Not suitable for high toxic	Kepler and McCleary (1994);
Alkalinity	Producing		a thick anoxic substrate of	acidity and generate alkalinity	metal mine waters; requires	Jage <i>et al.</i> (2001)
Systems (RAPS)			organic material neutralises	prior to discharge to aerobic	significant hydraulic head	
			acidity and generates	wetlands; efficient Fe and		
			alkalinity through processes	sulphate removal; suitable for		
			of bacterial sulphate reduction	net acidic mine waters with		
			and calcite dissolution; heavy	high ferric Fe, Al and		
			metals (mainly Fe and Al) are	dissolved oxygen content (>1		

---

---

removed as precipitates      mg/l); low footprint

Surface Catalyzed Net      *alkaline*      Containers are packed with      More efficient Fe removal      Not suitable for high toxic      Younger (2000); Jarvis and  
Oxidation Of Ferrous ferruginous      high specific surface area      than aerobic wetlands; low      metal mine waters; requires      Younger (2001); Sapsford  
Iron (SCOOFI)      inorganic media (e.g. plastic      footprint      significant hydraulic head;      and Williams (2009)  
trickle filter, ochre, blast      requires regular cleaning and  
furnace slag) which      replacing of filtering media  
encourage sorption and  
oxidation of ferrous Fe and  
accretion of ferric  
oxyhydroxide

*Source control technologies and techniques*

---

---

Permeable Barriers (PRBs)	reactive Net acidic	<p>PRBs provide a vertical and permeable compost-based medium in the path of and instead travel as polluted mine water which groundwater plumes neutralises acidity and promotes the generation of alkalinity through bacterial sulphate reduction and calcite dissolution</p>	<p>Useful for mine waters which do not emerge at the surface and instead travel as depth of aquifer</p>	<p>Limited evidence for removal of toxic metals; limited by</p>	<p>Benner <i>et al.</i> (1997); Jarvis <i>et al.</i> (2006)</p>
Physical of mine wastes	stabilisation -	<p>Covering of mine waste with inert material (e.g. clay, gravel) to reduce oxygen inflow and water ingress into the contaminated material and, hence, the concentrations of contaminants in drainage</p>	<p>Immobilises contaminants at source and prevents generation of mine drainage</p>	<p>Clay caps tend to crack in arid and semi-arid regions from wetting and drying cycles resulting in failure of air-tight cap</p>	<p>Gandy and Younger (2003); Waygood and Ferriera (2009)</p>

---

---

waters

Chemical stabilisation -  
of mine wastes

Addition of a resinous adhesive to form a crust over the mine waste

Immobilises contaminants at source and prevents generation of mine drainage

Similar to clay caps, crusts are prone to cracking resulting in failure of air-tight cap

Tordoff *et al.* (2000)

Phytostabilisation -

A vegetative cap on the mine waste to immobilise contaminants by adsorption and accumulation in the rhizosphere

Immobilises contaminants at source and prevents generation of mine drainage; creates wildlife habitat

Concerns over bioavailability of contaminants to wildlife; need for metal tolerant plants

Mendez and Maier (2008);  
Pollmann *et al.* (2009)

Phytoextraction -

A vegetative cap on the mine waste to immobilise contaminants through hyperaccumulation in plant tissues

Immobilises contaminants at source and prevents generation of mine drainage; creates wildlife habitat; offers the possibility of recovery of metals from plant tissues; improves land for agriculture

Concerns over bioavailability of contaminants to wildlife

Ernst (2005)

---



---

and forestry use

---

2270