

## The importance of reducing the systematic error due to non-linearity in N<sub>2</sub>O flux measurements by static chambers

P. S. Kroon · A. Hensen · W. C. M. van den Bulk ·  
P. A. C. Jongejan · A. T. Vermeulen

Published online: 31 October 2008  
© Springer Science+Business Media B.V. 2008

**Erratum to: Nutr Cycl Agroecosyst (2008) 82:175–  
186 DOI 10.1007/s10705-008-9179-x**

In the original publication, in Figure 10 (on page 185) the cumulative emissions were incorrect. Correct cumulative emissions should read:

---

The online version of the original article can be found under doi:[10.1007/s10705-008-9179-x](https://doi.org/10.1007/s10705-008-9179-x).

---

P. S. Kroon (✉) · A. Hensen · W. C. M. van den Bulk ·  
P. A. C. Jongejan · A. T. Vermeulen  
Department of Air Quality and Climate Change, Energy  
Research Centre of the Netherlands (ECN),  
Westerduinweg 3, 1755 LE Petten, The Netherlands  
e-mail: p.kroon@ecn.nl

**Fig. 10** Comparison of cumulative  $N_2O$  emission measured continuously and weekly at Cabauw in the Netherlands in September 2005 (a), February 2006 (b), April 2006 (c) and May 2006 (d). The *black arrow* indicates the moment of fertilizing

