



Original Article

The influence of the stomatognathic system on explosive strength: a pilot study

ANTONINO PATTI, MSc^{1, 2)*}, ANTONINO BIANCO¹⁾, GIUSEPPE MESSINA^{1, 2)}, ANTONIO PAOLI³⁾,
MARIANNA BELLAFFIORE¹⁾, GIUSEPPE BATTAGLIA¹⁾, ANGELO IOVANE¹⁾, MARCELLO TRAINA¹⁾,
ANTONIO PALMA¹⁾

¹⁾ Sport and Exercise Sciences Research Unit, University of Palermo: Via Giovanni Pascoli 6, 90144
Palermo, Italy

²⁾ Posturalab Italia, Italy

³⁾ Department of Biomedical Science, University of Padua, Italy

Abstract. [Purpose] Recent findings suggest there is an interesting interaction between the stomatognathic system and the musculoskeletal system. The aim of this study was therefore to examine the influence of the temporomandibular joint on the explosive strength of the lower limbs. [Subjects and Methods] An observational study was carried out. The subjects were 60 male football players who voluntarily participated in the investigation. After a warm-up phase of 10 minutes, each participant performed three Squat Jumps (SJ) with different mandible positions: mouth closed and mouth open. SJ heights were recorded using a Sensor Medica force platform and the FreeMed system. [Results] Sixty participants were enrolled in this study (age: 24 ± 7 yrs; height: 174 ± 4.6 cm; weight 63.7 ± 7.6 kg). The SJ heights with the mouth closed, 38.50 ± 4.0 cm, were shorter than those with the mouth open, 40.4 ± 4.1 cm. Statistical analysis showed there was a statistically significant difference between the performances. [Conclusion] This pilot study highlighted that occlusal factors can influence physical performance and this could have practical applications in sports and exercise science. However, our results have to be confirmed in studies with larger numbers of participants and supported by other investigations.

Key words: Stomatognathic system, Explosive strength, Dental occlusion

(This article was submitted Aug. 25, 2015, and was accepted Oct. 9, 2015)

INTRODUCTION

The literature suggests that there is a correlation between the stomatognathic system and the musculoskeletal system. Indeed, it seems certain that an alteration of the tooth-mandible-tongue complex affects postural attitude¹⁾. It has been reported that proprioception and postural control are of great importance for optimal sports injury prevention²⁻⁵⁾. In this context, Lee et al. confirmed that an exercise program aiming to improve proprioception can positively influence the balance ability of stroke patients⁶⁾. Tecco et al. reported a change in the postural stability of subjects with knee pathologies. These pathologies can generate changes in the function of the masticatory muscles, neck and trunk⁷⁾. Moreover, the mandible position is determined by dental occlusion, the relative position of teeth to each other^{8, 9)}. In addition, some studies have found that the stomatognathic system and mandible disorders can influence the spine^{10, 11)}. Tecco et al. also observed that the loading percentage and the loading surface on the right and the left feet were influenced by an experimentally induced imbalance of occlusion created by a cotton roll positioned in dental arches¹²⁾. Furthermore, it was observed that the temporomandibular joint (TMJ) affects other systems^{13, 14)}. Björne showed that tinnitus and vertigo were common in patients with TMJ dysfunction. Many of their patients also had symptoms of cervical spine disorders and head, neck and shoulder pain¹⁴⁾. It is also interesting to note that craniocervical flexion exercise increased postural stability¹⁵⁾. Yong et al. investigated the postural stability of twenty-four

*Corresponding author. Antonino Patti (E-mail: patti.nino@libero.it)

©2016 The Society of Physical Therapy Science. Published by IPEC Inc.

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <<http://creativecommons.org/licenses/by-nc-nd/3.0/>>.

subjects, randomly assigned to an exercise or control group, who performed a craniocervical flexion exercise four days per week for five weeks and reported their postural stability increased¹⁵). Moreover, Björne in 2007 showed that muscular problems and tinnitus were reduced by injection of lidocaine into the jaw muscle¹⁴). Recently, Martines et al. confirmed that tinnitus influences the postural control¹³). Furthermore, some studies have reported that the stomatognathic system affects physical performance, and that muscle activity is dependent on the position of the TMJ^{1, 12}). Some studies have shown that the stomatognathic system could elicit improvements in sport performances^{16, 17}), but the hypothesis has not yet been confirmed¹⁸). In 2012, Baldini et al. reported there was an increase in the relative strength of the quadriceps muscles when a patient wore an occlusal splint¹⁹). Our interest has focused on the role of the mandible position in physical performance, and the aim of this pilot study was to examine the influence of the temporomandibular joint and dental occlusion on the explosive strength of football players.

SUBJECTS AND METHODS

We performed an observational study. This study was performed in compliance with the Declaration of Helsinki, and the principles of the Italian data protection act (196/2003) were observed. The athletes selected for this study were 60 male football player volunteers. The squat jump (SJ) was the evaluation test adopted and all participants attempts were properly randomized (N° 30 with MO vs. MC; N° 30 MC vs. MO). SJ is well known and has been studied widely²⁰⁻²³). The SJ execution was standardized with a joint angle at departure of 90° degrees. Participants were instructed to jump for maximal height. The importance of avoiding any kind of preparatory countermovement was stressed. The protocol established for the execution of the jumps was as follows. In a warm-up phase of 10 minutes, the athletes performed light running (<65% VO₂max) and stretching exercises (four times 6 static postures for 20 seconds each), then each individual performed three practice SJs with the mouth closed (MC), and mouth open (MO). Then the athletes climbed on the platform for measurement and proceeded to perform SJs under the two different occlusal conditions. The participants were given a 2 minutes rest period between trials, and SJ was performed without shoes. The values were recorded using the FreeMed system and a force platform (Sensor Medica; Guidonia Montecelio, Roma). Statistical analysis was performed by a statistician. All data were input to an Excel file and the best performance of both occlusal conditions was selected for comparison. Statistical analysis was performed using StatSoft's STATISTICA software for Windows, Vers. 8.0 (Tulsa, OK). The paired t-test, one tail (p<0.05), was used to detect significant differences.

RESULTS

Sixty subjects were enrolled in this study (age: 24 ± 7.0 yrs; height: 174 ± 4.6 cm; weight 63.7 ± 7.6 kg). The best performances recorded were 38.50 ± 4.001 cm with MC and 40.41 ± 4.093 cm with MO. The Statistical analysis showed there was a statistically significant difference between the performances (p< 0.0001). The mouth open condition resulted in improvement in the vertical jump height compared to the mouth closed condition.

DISCUSSION

Some studies have found that there is a correlation between the stomatognathic system and the musculoskeletal system, and that an alteration of the tooth-mandible-tongue complex seems to affect postural attitude^{1, 11}). However, the conclusions of the different studies are not unanimous. Many studies are found in the literature, but despite the high number of articles published, the mechanism through which orthodontic treatment influences the etiology of TMJ dysfunction is still unknown²⁴). In the present pilot study, we sought to investigate whether the position of the mandible affects the explosive force. Our results are certainly encouraging but, according to the scientific literature, it is conceivable that the protagonist of the change is the trigeminal nerve²⁵⁻²⁷). Numerous anatomical associations have been described between the trigeminal system and the nervous structures implicated in posture control. The mesencephalic nucleus of the trigeminal nerve is a sensorial nucleus with unique characteristics²⁸). Gangloff and Perrin showed that the trigeminal nerve affected postural stabilization. Their results demonstrated the effects of trigeminal afferences on postural stabilization²⁹): postural deviation under the eyes closed condition was observed after unilateral truncular anaesthesia of the mandibular nerve in static posturography²⁹). Similarly, our results show that the mandible position influences physical performance. Another item of interest is the relationship between the trigeminal nerve and the muscle-fascial chain. The fascial system is important not only because it can passively distribute tension in the body muscles when mechanically stimulated, but also because it contains mechanoreceptors and possesses an autonomous contractile ability that influences the tension of the fasciae. These tensions seem to influence the body posture²⁸). Oh et al. showed that the stomatognathic alignment exercise improved TMJ function³⁰). The stomatognathic alignment exercise program was performed by an experimental group and consisted of mobility exercises for the TMJ and neck, and postural correction. At the end of the study, all the parameters of neck mobility and mouth opening had significantly improved. In conclusion, the result of this pilot investigation suggests that the mandible position influences physical performance, and that this could have practical applications in sports and exercise sciences. These conclusions need to be supported by further studies with different cohorts and larger populations.

REFERENCES

- 1) Fujimoto M, Hayakawa L, Hirano S, et al.: Changes in gait stability induced by alteration of mandibular position. *J Med Dent Sci*, 2001, 48: 131–136. [[Medline](#)]
- 2) Romero-Franco N, Gallego-Izquierdo T, Martínez-López EJ, et al.: Postural stability and subsequent sports injuries during indoor season of athletes. *J Phys Ther Sci*, 2014, 26: 683–687. [[Medline](#)] [[CrossRef](#)]
- 3) Şahin N, Bianco A, Patti A, et al.: Evaluation of knee joint proprioception and balance of young female volleyball players: a pilot study. *J Phys Ther Sci*, 2015, 27: 437–440. [[Medline](#)] [[CrossRef](#)]
- 4) Bianco A, Patti A, Bellafigliore M, et al.: Group fitness activities for the elderly: an innovative approach to reduce falls and injuries. *Aging Clin Exp Res*, 2014, 26: 147–152. [[Medline](#)] [[CrossRef](#)]
- 5) Mir SM, Talebian S, Naseri N, et al.: Assessment of knee proprioception in the anterior cruciate ligament injury risk position in healthy subjects: a cross-sectional study. *J Phys Ther Sci*, 2014, 26: 1515–1518. [[Medline](#)] [[CrossRef](#)]
- 6) Lee H, Kim H, Ahn M, et al.: Effects of proprioception training with exercise imagery on balance ability of stroke patients. *J Phys Ther Sci*, 2015, 27: 1–4. [[Medline](#)] [[CrossRef](#)]
- 7) Tecco S, Salini V, Calvisi V, et al.: Effects of anterior cruciate ligament (ACL) injury on postural control and muscle activity of head, neck and trunk muscles. *J Oral Rehabil*, 2006, 33: 576–587. [[Medline](#)] [[CrossRef](#)]
- 8) Passero PL, Wyman BS, Bell JW, et al.: Temporomandibular joint dysfunction syndrome. A clinical report. *Phys Ther*, 1985, 65: 1203–1207. [[Medline](#)]
- 9) Ohlendorf D, Seebach K, Hoerzer S, et al.: The effects of a temporarily manipulated dental occlusion on the position of the spine: a comparison during standing and walking. *Spine J*, 2014, 14: 2384–2391. [[Medline](#)] [[CrossRef](#)]
- 10) Kim HS, Na HJ, Kim HJ, et al.: Evaluation of proximal contact strength by postural changes. *J Adv Prosthodont*, 2009, 1: 118–123. [[Medline](#)] [[CrossRef](#)]
- 11) Solow B, Sonnesen L: Head posture and malocclusions. *Eur J Orthod*, 1998, 20: 685–693. [[Medline](#)] [[CrossRef](#)]
- 12) Tecco S, Polimeni A, Saccucci M, et al.: Postural loads during walking after an imbalance of occlusion created with unilateral cotton rolls. *BMC Res Notes*, 2010, 3: 141. [[Medline](#)] [[CrossRef](#)]
- 13) Martines F, Messina G, Patti A, et al.: Effects of tinnitus on postural control and stabilization: a pilot study. *Acta Med Mediter*, 2015, 31: 907–912.
- 14) Björne A: Assessment of temporomandibular and cervical spine disorders in tinnitus patients. *Prog Brain Res*, 2007, 166: 215–219. [[Medline](#)] [[CrossRef](#)]
- 15) Yong MS, Lee HY, Ryu YU, et al.: Effects of craniocervical flexion exercise on upper-limb postural stability during a goal-directed pointing task. *J Phys Ther Sci*, 2015, 27: 2005–2007. [[Medline](#)] [[CrossRef](#)]
- 16) Smith SD: Muscular strength correlated to jaw posture and the temporomandibular joint. *N Y State Dent J*, 1978, 44: 278–285. [[Medline](#)]
- 17) Sforza C, Tartaglia GM, Solimene U, et al.: Occlusion, sternocleidomastoid muscle activity, and body sway: a pilot study in male astronauts. *Cranio*, 2006, 24: 43–49. [[Medline](#)]
- 18) Perinetti G: Dental occlusion and body posture: no detectable correlation. *Gait Posture*, 2006, 24: 165–168. [[Medline](#)] [[CrossRef](#)]
- 19) Baldini A, Beraldi A, Nota A, et al.: Gnathological postural treatment in a professional basketball player: a case report and an overview of the role of dental occlusion on performance. *Ann Stomatol (Roma)*, 2012, 3: 51–58. [[Medline](#)]
- 20) Anderson FC, Pandy MG: Storage and utilization of elastic strain energy during jumping. *J Biomech*, 1993, 26: 1413–1427. [[Medline](#)] [[CrossRef](#)]
- 21) Bobbert MF, Gerritsen KG, Litjens MC, et al.: Why is countermovement jump height greater than squat jump height? *Med Sci Sports Exerc*, 1996, 28: 1402–1412. [[Medline](#)] [[CrossRef](#)]
- 22) Bobbert MF, van Zandwijk JP: Dynamics of force and muscle stimulation in human vertical jumping. *Med Sci Sports Exerc*, 1999, 31: 303–310. [[Medline](#)] [[CrossRef](#)]
- 23) van Zandwijk JP, Bobbert MF, Munneke M, et al.: Control of maximal and submaximal vertical jumps. *Med Sci Sports Exerc*, 2000, 32: 477–485. [[Medline](#)] [[CrossRef](#)]
- 24) Fernández-González FJ, Cañigral A, López-Caballo JL, et al.: Influence of orthodontic treatment on temporomandibular disorders. A systematic review. *J Clin Exp Dent*, 2015, 7: e320–e327. [[Medline](#)] [[CrossRef](#)]
- 25) Gangloff P, Louis JP, Perrin PP: Dental occlusion modifies gaze and posture stabilization in human subjects. *Neurosci*

- Lett, 2000, 293: 203–206. [[Medline](#)] [[CrossRef](#)]
- 26) Helling E, McWilliam J, Reigo T, et al.: The relationship between craniofacial morphology, head posture and spinal curvature in 8, 11 and 15-year-old children. *Eur J Orthod*, 1987, 9: 254–264. [[Medline](#)] [[CrossRef](#)]
 - 27) Milani RS, De Perière DD, Lapeyre L, et al.: Relationship between dental occlusion and posture. *Cranio*, 2000, 18: 127–134. [[Medline](#)]
 - 28) Cuccia A, Caradonna C: The relationship between the stomatognathic system and body posture. *Clinics (Sao Paulo)*, 2009, 64: 61–66. [[Medline](#)] [[CrossRef](#)]
 - 29) Gangloff P, Perrin PP: Unilateral trigeminal anaesthesia modifies postural control in human subjects. *Neurosci Lett*, 2002, 330: 179–182. [[Medline](#)] [[CrossRef](#)]
 - 30) Oh DW, Kang TW, Kim SJ: Effect of stomatognathic alignment exercise on temporomandibular joint function and swallowing function of stroke patients with limited mouth opening. *J Phys Ther Sci*, 2013, 25: 1325–1329. [[Medline](#)] [[CrossRef](#)]