

THE INTRINSIC RANDOM FUNCTIONS AND THEIR APPLICATIONS

G. MATHERON, *Centre de Morphologie Mathématique, Fontainebleau*

Abstract

The intrinsic random functions (IRF) are a particular case of the Gelfand generalized processes with stationary increments. They constitute a much wider class than the stationary RF, and are used in practical applications for representing non-stationary phenomena. The most important topics are: existence of a generalized covariance (GC) for which statistical inference is possible from a unique realization; theory of the best linear intrinsic estimator (BLIE) used for contouring and estimating problems; the turning bands method for simulating IRF; and the models with polynomial GC, for which statistical inference may be performed by automatic procedures.

GUELFAND GENERALIZED STOCHASTIC PROCESS; INTRINSIC RANDOM FUNCTION; GENERALIZED COVARIANCES; POLYNOMIAL COVARIANCE; TURNING BANDS METHOD

TABLE OF CONTENTS	Page
0. INTRODUCTION	440
1. DEFINITION AND GENERAL PROPERTIES	442
1.1 The spaces Λ and M_c	442
1.2 Definition of the IRF	443
1.3 Representations of a k -IRF	444
1.4 The translations formula	446
1.5 A decomposition theorem	448
1.6 Drift of an IRF	449
2. THE GENERALIZED COVARIANCES (GC)	450
2.1 The class of the GC of a k -IRF	450
2.2 k -IRF without drift	451
2.3 Examples	452
2.4 Applications of Theorem 2.1	453
3. THE BEST LINEAR INTRINSIC ESTIMATORS (BLIE)	457
Example 3.1 (the finite case)	459
Example 3.2 (estimation of a drift)	460
4. THE TURNING BANDS METHOD	461
5. THE POLYNOMIAL GC	462
6. k -IRF LOCALLY STATIONARY	465

Received in revised form 22 March 1973.

0. INTRODUCTION

The aim of the present study is essentially to act as a theoretical support to the optimum automatic contouring procedures being developed under the name of universal kriging, and which have been presented elsewhere in a more practical context ([7], [8], [9]). The problem we are dealing with is the following. We know the values $z(x_i)$ taken by the physical variable of interest at several experimental locations x_i (in general anywhere in two- or three-dimensional space) and we want to estimate at each point x the value of the function z (or of any other function deduced from it by a linear operation). We then assume that the function z can be considered as a realization of an order-two random function Z , and we compute at each point x the best linear predictor $z^*(x)$ of z (in the sense of Wiener [12]). But for this the covariance function of Z must be known. Statistical inference from a unique realization is in general reasonably possible in the stationary case (although some difficulties still arise when the experimental points are not located on a regular grid). Unfortunately, in numerous cases, this assumption of stationarity is physically inadmissible. Hence, it was necessary to find a wider class than that of order-two random functions, but one that would present the same advantages concerning statistical inference.

This class is that of intrinsic random functions (IRF) which constitutes, in fact, a particular case of the generalized stochastic processes with stationary increments of order k defined by Guelfand and Vilenkin ([3], [4]), namely the case where the generalized processes are random functions (and not only distributions). This circumstance leads to special properties which deserve to be studied, especially because of their interest regarding applications. For instance, here the generalized covariances will be conditionally positive definite functions (and not only distributions). The aim is, moreover, to present IRF's as a generalization of stationary random functions (stationary, in this paper, is always taken in the sense of weak stationarity), and hence without referring to the theory of distributions.

The simplest example of an IRF is that of a RF of the form $Y(x) = Y_0(x) + P(x)$, when $Y_0(x)$ is a SRF and $P(x)$ a polynomial of degree k with random coefficients. Besides, we will see that any IRF is, in a way, a limit of RF's of this type. The established concept of introducing this form of RF's comes naturally to mind when one tries to represent non-stationary phenomena. Indeed, it corresponds to the simplest hypothesis that can be made with a view to making statistical inference possible from a single realization, but IRF's present the same advantages concerning statistical inference, while offering a wider scope of possible models.

Realizations of an IRF of order 0, like Brownian motion, show characteristics that do not evoke the intuitive idea of stationarity (see, for example,

[2], p. 87). This circumstance is amplified when the order k of the IRF increases (see Figures 1, 2, 3) so that the theory becomes applicable to larger and larger ranges of non-stationary phenomena, while keeping, for the greatest part, the advantages linked to stationarity.




Figure 1
 Realization of a 0-IRF with the GC $K(h) = -|h|$. The representation chosen vanishes at the center of the figure (from Orfeuil [10]).




Figure 2
 Realization of a 1-IRF with the GC $K(h) = -|h|^3$. The representation chosen, as well as its first derivatives, vanishes at the center of the figure (from Orfeuil [10]).

After giving the definitions and general properties of IRF's (Sections 1 and 2), we examine in Section 3 their generalized covariances, and the conditions that an IRF must fulfil to be differentiable, or identical to an IRF of lower order. The theory of the best linear intrinsic estimator (BLIE), given




Figure 3
Realization of a 2-IRF with the GC $K(h) = -|h|^5$. The representation chosen, as well as its first and second derivatives, vanishes at the center of the figure (from Orfeuill [10]).

in Section 4, is a straightforward generalization of Wiener's best linear predictor [12], and is fundamental for the applications, as is the turning bands method for simulating IRF's (Section 5). From a practical point of view, statistical inference is particularly easy to carry out by automatic procedures for a generalized covariance whose expression depends linearly on some unknown parameters. This is the justification for models with polynomial covariances (Section 6). These models are isotropic, but this is not so limiting as might be thought at first sight. The structure of an IRF is indeed determined by its generalized covariance only up to a random polynomial and, in practical cases, this implicit polynomial is often adequate to take into account the anisotropies of the real phenomenon. On the other hand, the IRF's with polynomial covariances are locally stationary, i.e., may be locally identified with a stationary RF up to a random polynomial. These circumstances enable us to define a precise and locally significant notion of trend, or drift, and this is also important in the applications (Section 7).

1. DEFINITION AND GENERAL PROPERTIES

1.1 The spaces Λ and M_c

I shall denote by Λ the vector space of real measures in \mathbf{R}^n with *finite supports*. For any function f on \mathbf{R}^n and $\lambda \in \Lambda$, the integral $\int f(x) \lambda(dx)$ is thus a finite linear expression of the form $\sum_i \lambda_i f(x_i)$. If $Z: \mathbf{R}^n \rightarrow L^2(\Omega, \mathcal{A}, P)$ is a real order-two random function, it admits a linear extension $Z: \Lambda \rightarrow L^2(\Omega, \mathcal{A}, P)$ defined by putting $Z(\lambda) = \int \lambda(dx) Z(x)$, $\lambda \in \Lambda$. The function $\lambda \rightarrow \| \lambda \| = \| Z(\lambda) \|$

is a norm on Λ if $\int \lambda(dx)Z(x)$ implies $\lambda = 0$, this condition being equivalent to the strict positivity of the covariance matrix $\langle Z(x_\alpha), Z(x_\beta) \rangle$ for any finite set of distinct points x_α in \mathbf{R}^n . Under this condition, Λ is prehilbertian for the norm $\|\lambda\|$, and the completed space $\tilde{\Lambda}$ may be identified with the real Hilbert space $H \subset L^2(\Omega, \mathcal{A}, P)$ generated by the $Y(x)$, $x \in \mathbf{R}^n$. If the condition of strict positivity is not fulfilled, it is always possible to take a convenient quotient-space instead of Λ itself, and nothing is changed substantially.

It will often be useful in the sequel to consider another topology for Λ . Let $M_c \supset \Lambda$ be the space of measures with compact supports. M_c is the exact dual of the space \mathcal{C} of the continuous functions on \mathbf{R}^n (for the compact convergence topology). We shall consider only the weak topology on M_c , and the corresponding relative topology on Λ . For this weak topology, the convergence $\mu_n \rightarrow \mu$ in M_c is equivalent to the following two conditions:

- (1) the sequence $\{\mu_n\}$ is weakly convergent towards μ ;
- (2) the supports of the measures μ_n are contained in a fixed compact set.

The random function $Z: \mathbf{R}^n \rightarrow L^2(\Omega, \mathcal{A}, P)$ is then *strongly continuous* if and only if the mapping $\lambda \rightarrow Z(\lambda)$ is continuous on Λ for the relative M_c topology. The if part is obvious. Conversely, if Z is strongly continuous on \mathbf{R}^n , the covariance function $\langle Z(x), Z(y) \rangle$ is continuous on $\mathbf{R}^n \times \mathbf{R}^n$. But the weak convergence $\lambda_n \rightarrow 0$ in $\Lambda \subset M_c$ implies $\lambda_n \otimes \lambda_n \rightarrow 0$ in $M_c(\mathbf{R}^n \times \mathbf{R}^n)$, and

$$\|Z(\lambda_n)\|^2 = \int \lambda_n(dx) \langle Z(x), Z(y) \rangle \lambda_n(dy) \rightarrow 0.$$

Thus, the mapping $\lambda \rightarrow Z(\lambda)$ is continuous.

If Z is continuous on Λ , it admits a unique continuous extension on M_c , defined by putting $Z(\mu) = \int \mu(dx) Z(x)$ ($\mu \in M_c$). If $\|Z(x)\| = 0$ implies $\mu = 0$, M_c may be identified with a subspace (generally not closed) of $\tilde{\Lambda}$, but the weak topology on M_c is generally strictly stronger than the prehilbertian relative topology induced by $\tilde{\Lambda}$.

For any $h \in \mathbf{R}^n$ and $\mu \in M_c$, we define the *translated measure* $\tau_h \mu$ by

$$\int \tau_h(dx) f(x) = \int \mu(dx) f(x + h) \quad (f \in \mathcal{C}).$$

If Z is a *stationary random function* (SRF), there exists a *group of unitary operators* U_h , $h \in \mathbf{R}^n$ on $H = Z(\tilde{\Lambda})$ satisfying $U_h Z(\lambda) = Z(\tau_h \lambda)$ ($\lambda \in \Lambda$). Moreover, if Z is a continuous SRF, the group U_h is continuous and we also have $U_h Z(\mu) = Z(\tau_h \mu)$ for $\mu \in M_c$. This property may be used as definition of the SRF and, with a slight modification, will lead us to the notion of IRF.

1.2 Definition of the IRF

Now let Λ' be a *subspace* of Λ closed in Λ for the relative M_c topology. In other words, there exists a family f^l , $l \in L$ of continuous functions on

\mathbf{R}^n such that $\Lambda' = \{\lambda: \lambda \in \Lambda, \int \lambda f^l = 0, l \in L\}$, or $\Lambda' = M' \cap \Lambda$ if M' is the subspace of M_c orthogonal to the family $f^l, l \in L$. Then, we shall say that a linear mapping

$$Z: \Lambda' \rightarrow L^2(\Omega, \mathcal{A}, P)$$

is a *generalized (order-two) random function on Λ'* . We denote $H(Z)$, or simply H , the closure of the range $Z(\Lambda')$ in $L^2(\Omega, \mathcal{A}, P)$. If $\|Z(\lambda)\| = 0$ implies $\lambda = 0$ for $\lambda \in \Lambda'$, the completed space $\tilde{\Lambda}'$ of Λ' for the norm $\|\lambda\| = \|Z(\lambda)\|$ may be identified with H itself. If this condition is not fulfilled, $\tilde{\Lambda}'$ will denote the completion of the convenient quotient space. If Z is continuous on Λ' for the relative M_c topology (*continuous generalized random function*), it admits a unique continuous extension $Z: M' \rightarrow L^2(\Omega, \mathcal{A}, P)$, and M' (or a convenient quotient space) may be identified with a dense (but not closed) subspace of $\tilde{\Lambda}'$. In order to generalize the notion of stationarity, we now suppose Λ' *closed for translations* (i.e., $\tau_h \lambda \in \Lambda'$ for $h \in \mathbf{R}^n$ and $\lambda \in \Lambda'$), and we say that the generalized random function $Z: \Lambda' \rightarrow L^2(\Omega, \mathcal{A}, P)$ is an *intrinsic random function (IRF)* on Λ' if the mapping $h \rightarrow Z(\tau_h \lambda)$ is a SRF for any $\lambda \in \Lambda'$, i.e., if there exists a *group of unitary operators* $U_h, h \in \mathbf{R}^n$ on $H = Z(\tilde{\Lambda}')$ satisfying $U_h Z(\lambda) = Z(\tau_h \lambda)$ for any $\lambda \in \Lambda'$. Obviously, an IRF Z is continuous on Λ' if and only if the mapping $(h, \lambda) \rightarrow Z(\tau_h \lambda)$ is continuous on $\mathbf{R}^n \times \Lambda'$. In this case, the group U_h is continuous on \mathbf{R}^n , and we also have $U_h Z(\mu) = Z(\tau_h \mu)$ for any $\mu \in M'$.

The subspace $M' \subset M_c$ being closed for translations, so also is its orthogonal complement, i.e., the closed subspace of \mathcal{C} generated by the functions $f^l, l \in L$. In particular, if L is *finite*, this implies that the f^l are exponential-polynomial functions. In what follows, we shall examine only the case where the f^l functions *are polynomials*. More precisely: for k , an integer ≥ 0 , we denote M_k the subspace of M_c defined by the condition $\mu \in M_k$ if $\mu \in M_c$ and

$$\int x_1^{i_1} x_2^{i_2} \dots x_n^{i_n} \mu(dx) = 0$$

for i_1, \dots, i_n , integers ≥ 0 , such that $i_1 + \dots + i_n \leq k$. For brevity, we shall write l instead of $(i_1, \dots, i_n), l \leq k$ instead of $i_1 + \dots + i_n \leq k$ and $f^l(x)$ for $x_1^{i_1} x_2^{i_2} \dots x_n^{i_n}$. The subspace $\Lambda \cap M_k$ (closed in Λ) will be denoted Λ_k , and we shall say that an IRF Z on Λ_k is an *IRF of order k , or a k -IRF*. If a k -IRF Z is continuous, its unique continuous extension on M_k will also be denoted by Z . In the sequel, we consider only *continuous k -IRF's*.

1.3 Representations of a k -IRF

If Z is an IRF on a subspace $\Lambda' \subset \Lambda$, we say that an order-two random function $Y: \mathbf{R}^n \rightarrow L^2(\Omega, \mathcal{A}, P)$ is a *representation of Z* if $Z(\lambda) = \int Y(x)\lambda(dx)$

for any $\lambda \in \Lambda'$. In the sequel, we examine only the case of a continuous k -IRF Z . It is easy to find measures $\lambda_l \in \Lambda$ ($l = i_1 + i_2 + \dots + i_n \leq k$) such that

$$(1.1) \quad \int \lambda_l(dx) f^s(x) = \delta_l^s$$

($s = (j_1, \dots, j_n)$, $j_1 + \dots + j_n \leq k$, $\delta_l^s = 0$ for $l \neq s$ and $\delta_l^s = 1$ if $s = l$). If δ_x is the Dirac measure at $x \in \mathbb{R}^n$, clearly $(\delta_x - f_x^l \lambda_l) \in \Lambda_k$. Then, the random function $Y(x)$, $x \in \mathbb{R}^n$, defined by

$$(1.2) \quad Y(x) = Z(\delta_x - f_x^l \lambda_l)$$

is a representation of Z . For, if $\lambda \in \Lambda_k$, i.e., $\int \lambda(dx) f^l(x) = 0$, we also have $\lambda = \int \lambda(dx) [\delta_x - f_x^l \lambda_l]$. Z being a linear mapping of Λ_k into $H = Z(\bar{\Lambda}_k)$, we may write

$$Z(\lambda) = \int \lambda(dx) Z(\delta_x - f_x^l \lambda_l) = \int \lambda(dx) Y(x)$$

and $Y(x)$ is a representation of the k -IRF Z .

Let $X(x)$ be another representation of Z . By the very definition, we may write

$$Z(\lambda) = \int \lambda(dx) Y(x) = \int \lambda(dx) X(x) \text{ for any } \lambda \in \Lambda_k.$$

In particular, for $\lambda = \delta_x - f_x^l \lambda_l$, we find

$$Y(x) = Y(x) - f_x^l \int \lambda_l(dy) Y(y) = X(x) - f_x^l \int \lambda_l(dy) X(y).$$

From this relationship, we conclude:

(a) The representation defined by the relationship (1.2) is *characterized* by

$$(1.3) \quad \int \lambda_l(dx) Y(x) = 0.$$

For, if another representation $X(x)$ satisfies (1.3), we find

$$Y(x) = X(x) - f_x^l \int \lambda_l(dy) X(y) = X(x).$$

(b) Any other representation $X(x)$ of Z is of the form

$$(1.4) \quad X(x) = Y(x) + A_l f^l(x)$$

for random variables $A_l \in L^2(\Omega, \mathcal{A}, P)$ satisfying $A_l = \int \lambda_l(dx) X(x)$. Converse-

ly, for any random variables $A_i \in L^2(\Omega, \mathcal{A}, P)$, the random function $X(x)$ defined by the relationship (2.4) is obviously a representation of Z , because

$$\int \lambda(dx) X(x) = \int \lambda(dx) Y(x) \text{ for any } \lambda \in \Lambda_k.$$

This representation satisfies

$$(1.5) \quad A_i = \int \lambda_i(dx) X(x)$$

by (1.1) and (1.3). Thus, the relationships (1.2) and (1.4) give us *the general form of all the representations* of the k -IRF Z .

The k -IRF Z being continuous on Λ_k (for the relative M_c topology), all its representations are continuous, and, conversely, if Z admits a continuous representation, it is a continuous k -IRF.

1.4 The translations formula

Let Z be a continuous k -IRF. Let us choose measures λ_i satisfying (1.1), and consider the representation $Y(x) = Z(\delta_x - f_x^i \lambda_i)$. From the very definition of an IRF, we have

$$U_h Y(x) = Z(\delta_{x+h} - f_x^i \tau_h \lambda_i)$$

for $h, x \in \mathbb{R}^n$. This may be written as follows

$$U_h Y(x) = Z(\delta_{x+h} - f_{x+h}^i \lambda_i) + Z(f_{x+h}^i \lambda_i - f_x^i \tau_h \lambda_i).$$

But $Z(\delta_{x+h} - f_{x+h}^i \lambda_i) = Y(x+h)$, and the relationship (1.3) implies

$$\begin{aligned} Z(f_{x+h}^i \lambda_i - f_x^i \tau_h \lambda_i) &= f_{x+h}^i \int \lambda_i(dy) Y(y) - f_x^i \int \lambda_i(dy) Y(y+h) \\ &= -f_x^i \int \lambda_i(dy) Y(y+h). \end{aligned}$$

Thus we obtain the following *translations formula*

$$(1.6) \quad \begin{aligned} U_h Y(x) &= Y(x+h) - A_i(h) f^i(x), \\ A_i(h) &= \int \lambda_i(dy) Y(y+h). \end{aligned}$$

From the relationship (1.6) we can obtain an important *inequality*. Because $\|Y(x)\|$ is continuous on \mathbb{R}^n , and the measures $\lambda_i \in \Lambda$ have their supports included in the same compact, there exists a real $B > 0$ such that

$$|h| \leq 1 \Rightarrow \left\| \int \lambda_i(dy) Y(x+h) \right\| \leq B.$$

always exists, and is an infinitely differentiable random function, as can easily be verified.

Besides, we note that for any $\lambda \in \Lambda_k$, the integral $\int \lambda(dx)Y_\phi(x)$ does not depend on the choice of the particular representation $Y(x)$ of the k -IRF we have used, as shown by

$$\begin{aligned} \int \lambda(dx)Y_\phi(x) &= \int \phi(y)dy \int \lambda(dx)Y(x+y) \\ &= \int \phi(y)U_yZ(\lambda)dy. \end{aligned}$$

In other words, the regularized RF $Y_\phi(x)$ is a representation of the k -IRF Z_ϕ defined on Λ_k by putting $Z_\phi(\lambda) = \int \phi(y)U_yZ(\lambda)dy$.

We shall say that the k -IRF Z_ϕ is the regularized function of Z by ϕ . Clearly, Z_ϕ is infinitely differentiable if $\phi \in \mathcal{S}$ (i.e., any representation of Z_ϕ is infinitely differentiable, or $Z_\phi(D)$ exists for any derivative D). If D is a derivative of order p , the corresponding derivative DZ_ϕ , defined by

$$DZ_\phi(\lambda) = (-1)^p \int D\phi(y)U_yZ(\lambda)dy$$

is a $(k-p)$ -IRF for $p \leq k$, and a SRF for $p > k$.

1.5 A decomposition theorem

If $Y(x)$, $x \in \mathbf{R}^n$ is a SRF, the mapping $\lambda \rightarrow \int \lambda(dx)Y(x)$ obviously defines a k -IRF on Λ_k , which may be denoted $\lambda \rightarrow Y(\lambda)$. It is thus possible to add a k -IRF Z_c and the SRF Y considered as a k -IRF. The sum $Z_c + Y$ is the k -IRF Z defined by $Z(\lambda) = Z_c(\lambda) + \int \lambda(dx)Y(x)$, $\lambda \in \Lambda_k$. Then we may state the following theorem.

Theorem 1.5. Any continuous k -IRF Z is the sum of a SRF and of an infinitely differentiable k -IRF.

Proof. Let $p_a \in \mathcal{S}$ be the function defined by its Fourier transform

$$\tilde{p}_a(u) = (1 + \frac{1}{2}a|u|^2 + \dots + \frac{1}{2}k'^{-1}a^{k'}|u|^{2k'}) \exp(-a|u|^2) \text{ for } a > 0$$

and an integer $k' > \frac{1}{2}k$. All the derivatives of $1 - \tilde{p}_a$ up to order $2k' > k$ vanish in $u=0$. Thus, the measure $\delta - p_a(x)dx$ is in M_k . Put $Y_a(x) = Z(\delta_x - \tau_x p_a)$, i.e., $Y_a(x) = U_xZ(\delta - p_a)$. In other words, $Y_a(x)$ is a SRF. From the definition of the regularized Z_{p_a} , we then have $Z = Y_a + Z_{p_a}$, where Y_a is the mapping $\lambda \rightarrow \int Y_a(x)\lambda(dx)$, $\lambda \in \Lambda_k$, and the regularized Z_{p_a} is an infinitely differentiable k -IRF.

For any representation $Y(x)$ of Z , we have

$$Y_a(x) = Z(\delta_x - \tau_x p_a) = Y(x) - \int p_a(y)Y(x + y)dy,$$

because $(\delta_x - \tau_x p_a) \in \Lambda_k$, and the decomposition theorem gives simply $Y(x) = Y_a(x) + Y_{p_a}(x)$, where $Y_a(x) = Y(x) - Y_{p_a}(x)$ is a SRF and $Y_{p_a}(x)$ a representation of the infinitely differentiable k -IRF Z_{p_a} .

Also note the following corollary.

Corollary. For any k -IRF, there exists a sequence $\{Y_n\}$ of SRF such that $\int \lambda(dx) Y_n(x)$ strongly converges toward $Z(\lambda)$ for any $\lambda \in \Lambda_k$.

With the preceding notation, we have to show that, for $\lambda \in \Lambda_k$, $\lim Y_a(\lambda) = Z(\lambda)$ for $a \rightarrow \infty$, i.e., $\lim \int p_a(y)U_y Z(\lambda)dy = 0$. Let $\chi(du)$ be the spectral measure associated with the SRF $U_y Z(\lambda)$. Then, by $|\tilde{p}_a| \leq 1$ and $|\tilde{p}_a(u)|^2 \rightarrow 0$ for any $u \in \mathbb{R}^n$, we obtain

$$\left\| \int p_a(y)U_y Z(\lambda)dy \right\|^2 = \int |\tilde{p}_a(u)|^2 \chi(du) \rightarrow 0,$$

i.e., $Y_a(\lambda) \rightarrow Z(\lambda)$.

1.6 Drift of an IRF

Let Z be a continuous k -IRF, H_0 the subspace of $H = H(\tilde{\Lambda}_k)$ containing the invariant elements of H (i.e., $X \in H_0$ if $X \in H$ and $U_h X = X$ for any $h \in \mathbb{R}^n$) and Π_0 the projector on H_0 (in the ergodic case, Π_0 may be identified with the expected value). By putting $m_0(\lambda) = \Pi_0 Z(\lambda)$ for $\lambda \in \Lambda_k$, we define a continuous k -IRF m_0 obviously invariant for the group U_h . Let us choose $\lambda \in \Lambda$ satisfying (1.1), and write $m_0(x) = m_0(\delta_x - f_x^! \lambda)$. Then, the representation $m_0(x)$ of the k -IRF m_0 is a polynomial of degree $\leq k + 1$ with coefficients in H_0 (constant in the ergodic case).

Proof. If Z is infinitely differentiable, so also are m_0 and its representation $m_0(x)$. By $m_0(x) \in H_0$ and the translations formula (1.6), we obtain $m_0(x) = m_0(x + h) - A_l(h)f^l(x)$. Differentiation of order $k + 1$ with respect to x eliminates the terms $A_l(h)f^l(x)$, and then differentiation of order 1 with respect to h eliminates $m_0(x)$. Hence, all derivatives of order $k + 2$ of $m_0(x)$ vanish identically, and $m_0(x)$ is a polynomial of degree $\leq k + 1$. The decomposition theorem shows that the result remains true when Z is not differentiable.

The polynomial $m_0(x)$ with invariant coefficients depends on the choice of the representation we have used. But it is easy to verify that the terms of degree $k + 1$ do not depend on it, and thus constitute a property of the k -IRF Z itself. The corresponding homogeneous polynomial of degree $k + 1$ with invariant coefficients will be called the *drift* (or *intrinsic drift*) of the k -IRF.

If $m_0 = 0$, the k -IRF Z is said to be *without drift*. Obviously, Z is without drift if and only if $\Pi_0 Y(x)$ is a polynomial of degree $\leq k$ with invariant coefficients for a representation of Z (and then, the same is true for any representation).

2. THE GENERALIZED COVARIANCES (GC)

2.1 The class of the GC of a k -IRF

Let Z be a continuous k -IRF. We say that a continuous and symmetric function K on \mathbf{R}^n is a *generalized covariance* (GC) of Z if

$$\langle Z(\lambda), Z(\mu) \rangle = \int \lambda(dx) K(x - y) \mu(dy) \text{ for } \lambda, \mu \in \Lambda_k.$$

This is equivalent to the condition:

$$(2.1) \quad \int \lambda(dx) K(x - y) \lambda(dy) = \|Z(\lambda)\|^2 \quad (\lambda \in \Lambda_k).$$

The family of the continuous and symmetric functions K satisfying the condition (2.1) will be called *the class of the GC of Z* . We will show that such GC always exist, and, if one of them is known, we obtain all the others by adding arbitrary even polynomials of degree $\leq 2k$. In other words, there is *an existence theorem*, and a *uniqueness theorem*, the latter *up to an equivalence* defined by the relationship $K_1 \equiv K_2$ if $K_1 - K_2$ is an even polynomial of degree $\leq 2k$.

We shall also say that a continuous, symmetric function K on \mathbf{R}^n is *conditionally positive definite* of order k if

$$\int \lambda(dx) K(x - y) \lambda(dy) \geq 0 \text{ for any } \lambda \in \Lambda_k.$$

From the definition, the GC of a k -IRF (if any) are k -conditionally positive definite. Conversely, if K is a continuous k -conditionally positive definite function, there exists a continuous k -IRF Z such that the $Z(\lambda)$, $\lambda \in \Lambda_k$ are Gaussian random variables satisfying (2.1), and these two classes of functions may be identified. The following theorem gives their characterization.

Theorem 2.1. Let K be a continuous and symmetrical function on \mathbf{R}^n . Then, K is k -conditionally positive definite if and only if it admits the representation:

$$(2.2) \quad K(h) = \int \frac{\cos(2\pi(uh)) - 1_B(u)P_k(2\pi(uh))}{(4\pi^2|u|^2)^{k+1}} \chi_0(du) + K_0(h),$$

where K_0 is an even k -conditionally positive definite polynomial of degree

$\leq 2k + 2$, 1_B the indicator of a neighbourhood of the origin, and χ_0 a positive measure, necessarily unique, without atom at the origin, and such that

$$(2.3) \quad \int [1 + 4\pi^2 |u|^2]^{-k-1} \chi_0(du) < \infty$$

(P_k is the polynomial $P_k(x) = \sum_0^k (-1)^p x^{2p} / (2p)!$).

Proof. It follows from Section 1.4 that any representation of a k -IRF is a generalized stochastic process with stationary increments, so that the existence and uniqueness theorem as well as Theorem 2.1 are simple consequences of the Guelfand-Vilenkin theory ([3], Chapter 2, Section 4 and Chapter 3, Section 3). Note that it is also possible to derive them by spectral analysis, i.e., without using the distribution theory.

2.2 k -IRF without drift

Let Z be a k -IRF, K a GC of Z which admits the representation (2.2), $\lambda \in \Lambda_k$ and $\tilde{\lambda}$ its Fourier transform. Then, it follows from (2.2) that the spectral measure χ_λ associated with the SRF $x \rightarrow U_x Z(\lambda)$ is

$$\chi_\lambda = (|\tilde{\lambda}|^2 / 4\pi^2 |u|^{2k+2}) \chi_0 + a\delta,$$

and the atom a is given by

$$a = \int \lambda(dx) K_0(x-y) \lambda(dy).$$

In other words, Z is without drift if and only if the degree of the polynomial K_0 is $\leq 2k$. More precisely, we may state the following theorem.

Theorem 2.2. Let Z be a continuous k -IRF, K a GC and $Y(x)$, $x \in \mathbb{R}^n$ a representation of Z . Then, the following three conditions are equivalent:

- (a) Z is without drift (i.e., $\Pi_0 Z = 0$),
- (b) $K(h) / |h|^{2k+2} \rightarrow 0$ for $|h| \rightarrow \infty$,
- (c) $Y(x) / |x|^{k+1} \rightarrow 0$ strongly for $|x| \rightarrow \infty$.

Proof. By the inequality $|\cos x - P_k(x)| \leq x^{2k+2} / (2k + 2)!$, we have

$$\left| \frac{\cos(2\pi(uh)) - P_k(2\pi(uh))}{(4\pi^2 u^2)^{k+1}} \right| \leq \frac{|h|^{2k+2}}{(2k + 2)!}$$

and, by the dominated convergence theorem

$$\lim_{|h| \rightarrow \infty} \int_B \frac{\cos(2\pi(uh)) - P_k(2\pi(uh))}{|h|^{2k+2} (4\pi^2 u^2)^{k+1}} \chi_0(du) = 0.$$

Thus, the condition (b) is fulfilled if and only if the polynomial K_0 of Theorem 2.3 is of degree $\leq 2k$, i.e., if (a) is satisfied, and thus (a) and (b) are equivalent.

By the relationship (1.4), we may assume that the representation $Y(x)$ of Z is of the form (1.2). Then, the covariance of $Y(x)$ is the function

$$\begin{aligned} \langle Y(x), Y(y) \rangle &= K(x-y) - f^l(x) \int K(z-y)\lambda_l(dz) - f^l(y) \int K(z-x)\lambda_l(dz) \\ &\quad + f^l(x)f^s(y) \int \lambda_l(dz)K(z-z')\lambda_l(dz'). \end{aligned}$$

It follows from this relationship that (b) implies (c).

Finally, the representation $\Pi_0 Y(x)$ of the drift $m_0 = \Pi_0 Z$ is a polynomial of degree $\leq k + 1$ with invariant coefficients. If the condition (c) is fulfilled, this polynomial is of degree $\leq k$, and thus Z is without drift.

2.3 Examples

By Theorem 2.1, a positive measure χ_0 without atom at the origin and satisfying the condition (2.3) characterizes a k -conditionally positive definite function, and thus a possible model of a k -IRF.

For instance, the function r^α on \mathbf{R}^n ($r = |x|$, α real, > 0 and different from an even integer) admits as a Fourier transform (in the sense of the distribution theory) the (pseudo) function $\pi^{-\alpha-\frac{1}{2}n} \Gamma[\frac{1}{2}(\alpha+n)]/\Gamma(-\frac{1}{2}\alpha)\rho^{-\alpha-n}$ (with $\rho = |u|$). If $\alpha < 2k + 2$, the absolutely continuous measure $\rho^{2k+2-\alpha-n} du$ satisfies the conditions of Theorem 2.1. Thus $\Gamma(-\frac{1}{2}\alpha)r^\alpha$ is k -conditionally positive definite. If $\alpha = 2p + 1$ ($p \leq k$) is an odd integer, the function $(-1)^{p+1}r^{2p+1}$ may be used as GC of a k -IRF. More generally, the function K defined on \mathbf{R}^n by

$$(2.4) \quad K(r) = \sum_{p=0}^k (-1)^{p+1} a_p r^{2p+1}$$

is a k -GC if and only if the coefficients a_p satisfy the condition

$$\sum_{p=0}^k \frac{a_p}{\pi^{2p+2+\frac{1}{2}n}} \frac{\Gamma[\frac{1}{2}(2p+1+n)]}{\Gamma[1+\frac{1}{2}(2p+1)]} \rho^{-n-2p+1} \geq 0$$

for any $\rho > 0$. These ‘‘polynomial isotropic’’ GC are interesting from the applications point of view, because their expression depends linearly on the coefficients a_p , an advantageous property for statistical inference. (See succeeding paragraph).

Now let us examine the case $n = 1$. Let $X(x)$ be a SRF on \mathbf{R}^1 , $\sigma(h)$ its stationary covariance and χ its spectral measure, and consider the successive integrals

$$Y_0(x) = \int_0^x X(\xi)d\xi, \dots, Y_k(x) = \int_0^x Y_{k-1}(\xi)d\xi,$$

or explicitly,

$$Y_k(x) = \int_0^x \frac{(x - \xi)^k}{k!} X(\xi)d\xi.$$

Then, Y_k is a representation of a k -IRF Z_k with the GC K_k defined by

$$K_k(h) = \int_0^{|h|} \frac{(|h| - \xi)^{2k+1}}{(2k + 1)!} \sigma(\xi)d\xi.$$

If $\Pi_0 X(x) = 0$, Z_k is without drift, and the spectral measure χ_0 of $X(x)$ is identical with the measure χ_0 occurring in Theorem 2.3.

Now if $W_0(x)$ is a 0-IRF with the GC $-|h|$ (for instance, W_0 may be a Brownian motion), the expression

$$W_k(x) = \int_0^x \frac{(x - \xi)^{k-1}}{(k - 1)!} W_0(\xi)d\xi$$

constitutes a representation of a k -IRF with the GC $(-1)^{k+1}|h|^{2k+1}/(2k + 1)!$. More generally, if we put

$$(2.5) \quad Y(x) = \sum_{p=0}^k b_p W_p(x)$$

with arbitrary real coefficients b_p , we obtain a realization of a k -IRF which admits a polynomial GC of type (2.4). Conversely, it is possible to show that any k -IRF with a polynomial GC of type (2.4) admits a representation of the form (2.5).

2.4 Applications of Theorem 2.1

It follows from Theorem 2.1 that the classical results of the harmonic analysis may be applied to the k -IRF. Let Z be a k -IRF without drift, and χ_0 the spectral measure associated by Theorem 2.1 with its GC. Then, as we have seen, for any $\lambda \in \Lambda_k$, the SRF $x \rightarrow U_x Z(\lambda)$ admits the spectral measure $\chi_\lambda = (|\tilde{\lambda}|^2/(4\pi^2|u|^2)^{k+1})\chi_0$ ($\tilde{\lambda}$ is the Fourier transform of λ). Moreover, if we denote by ζ_λ the orthogonal random measure whose Fourier transform is $U_x Z(\lambda)$, the following relationship holds

$$(2.6) \quad \tilde{\lambda}'\zeta_\lambda = \tilde{\lambda}\zeta_{\lambda'} \quad (\lambda, \lambda' \in \Lambda_k).$$

Furthermore, the mapping $\lambda \rightarrow \tilde{\lambda}/(4\pi^2|u|^2)^{\frac{1}{2}(k+1)}$ may be extended by an isomorphism from the completion $\tilde{\Lambda}_k$ onto $L^2(\mathbb{R}^n/\chi_0)$, so that (2.6) remains valid for any $\lambda, \lambda' \in \tilde{\Lambda}_k$.

As a direct consequence, we see that the k -IRF Z is differentiable up to order p if and only if

$$(2.7) \quad \int [1 + 4\pi^2 |u|^2]^{p-k-1} \chi_0(du) < \infty.$$

Thus, the integral

$$(2.8) \quad K_1(h) = \int \frac{\cos(2\pi(uh)) - P_k(2\pi(uh))}{(4\pi^2 |u|^2)^{k+1}} \chi_0(du)$$

exists, and represents a GC of Z , if and only if Z is differentiable up to order k . If so, K_1 is the unique GC of Z which vanishes at $h = 0$ as also do its derivatives up to order $2k$. In particular, any 0-IRF without drift admits a GC of the form

$$K_1(h) = \int \frac{\cos(2\pi(uh)) - 1}{4\pi^2 |u|^2} \chi_0(du).$$

Any k -GC K satisfies the inequality

$$(2.9) \quad |K(0) - K(h)| \leq a + b |h|^{2k+2}$$

for convenient constants $a, b \geq 0$, as can be shown from (2.2) or (1.7). Then, the following criterion is easy to prove by harmonic analysis: a k -IRF is differentiable if and only if its GC satisfies inequalities of the form

$$(2.10) \quad |K(0) - K(h)| \leq \alpha |h|^2 + \beta |h|^{2k+2}.$$

Let us now examine under which condition a k -IRF Z actually is of order $< k$, i.e., is the restriction to Λ_k of a $(k-1)$ -IRF \bar{Z} .

Theorem 2.4. Let Z be a continuous k -IRF without drift, $k \geq 1$ (respectively, a 0-IRF such that $\Pi_0 Z = 0$). Then, the following three conditions are equivalent.

(a) Z is the restriction to Λ_k of a continuous $(k-1)$ -IRF \bar{Z} (resp. the restriction to Λ_0 of a continuous SRF \bar{Z}), and \bar{Z} is unique up to a drift (resp. to an invariant).

(b) There exists a measure $\chi'_0 \geq 0$ without atom at the origin such that $\int [1 + 4\pi^2 |u|^2]^{-k} \chi'_0(du) < \infty$, and the spectral measure associated with Z by Theorem 2.1 is $\chi_0 = (4\pi^2 |u|^2) \chi'_0$. If so, χ'_0 is the spectral measure associated with the $(k-1)$ -IRF \bar{Z} (with the SRF \bar{Z}).

(c) The GC's of Z satisfy inequalities of the form $|K(h)| \leq a + b |h|^{2k}$ (resp. are bounded on \mathbb{R}^n).

Proof. Let Z be a k -IRF and \bar{Z}, \bar{Z}' $(k-1)$ -IRF such that $\bar{Z}(\lambda) = \bar{Z}'(\lambda) = Z(\lambda)$ for any $\lambda \in \Lambda_k$. Then, for any $\lambda' \in \Lambda_{k-1}$, we find

$$(U_h - I)Z'(\lambda) = Z(\tau_h\lambda - \lambda) = (U_h - I)Z(\lambda).$$

Thus, $Z'(\lambda) = Z(\lambda)$ up to an invariant, and Z is unique up to a drift.

The implications (a) \Rightarrow (b) and (b) \Rightarrow (c) are trivial. Let K be a k -GC such that (c) holds, and χ_0 its spectral measure. If B is the neighbourhood of 0 such that (2.1) holds, and $C = B^c$ the complementary set, the measure $1_C\chi_0$ trivially satisfies (b), so that we may suppose $\chi_0 = 1_B\chi_0$, i.e., Z is infinitely differentiable. Then, the function K_1 defined by (2.8) exists and is a GC of Z . Moreover, by (2.4) and (c), there exists a real $B < \infty$ such that

$$(2.11) \quad |K_1(h)| \leq B|h|^{2k}.$$

It remains to prove that (2.11) implies (b). We start from the obvious relationship

$$\cos x - P_k(x) = (-1)^{k+1} \int_0^x \frac{(x - \xi)^{2k-1}}{(2k-1)!} [1 - \cos \xi] d\xi$$

which implies, for $r = |h|$ and $\alpha = h/r$,

$$\begin{aligned} &\cos 2\pi(uh) - P_k(2\pi(uh)) \\ &= (-1)^{k+1} (2\pi(u\alpha))^{2k} \int_0^r \frac{(r - \rho)^{2k-1}}{(2k-1)!} [1 - \cos(2\pi(u\alpha)\rho)] d\rho. \end{aligned}$$

By substituting in (2.8) we obtain

$$(2.12) \quad (-1)^{k+1} K_1(\alpha r) = \int_0^r \frac{(r - \rho)^{2k-1}}{(2k-1)!} d\rho \int \frac{(u\alpha)^{2k}}{(u^2)^k} \frac{1 - \cos(2\pi(u\alpha)\rho)}{4\pi^2 u^2} \chi_0(du).$$

Let Φ_α be the Laplace transform of $(-1)^{k+1} K_0(\alpha r)$, i.e.,

$$\Phi_\alpha(\lambda) = (-1)^{k+1} \int_0^\infty K_1(\alpha r) e^{-\lambda r} dr \quad (\lambda > 0).$$

By (2.12), the function $(-1)^{k+1} K_1(\alpha r)$ is the convolutive product in \mathbf{R}_+ of two functions, the Laplace transforms of which are respectively λ^{-2k} and

$$\int \frac{(u\alpha)^{2k}}{(u^2)^k} \frac{\chi_0(du)}{\lambda(\lambda^2 + 4\pi^2(u\alpha)^2)}.$$

Thus, we obtain

$$(2.13) \quad \Phi_\alpha(\lambda) = \frac{1}{\lambda^{2k+1}} \int \frac{(u\alpha)^{2k}}{(u^2)^k} \frac{\chi_0(du)}{\lambda^2 + (2\pi(u\alpha))^2}.$$

On the other hand, the function $(-1)^{k+1} K_0(\alpha r)$ is ≥ 0 by (2.8), and thus (2.11) implies the inequality $\Phi_\alpha(\lambda) \leq B(2k)!/\lambda^{2k+1}$. By (2.13) and $(u\alpha)^2 \leq u^2\alpha^2 = u^2$, we obtain

$$\int \frac{(u\alpha)^{2k}}{(u^2)^k} \frac{\chi_0(du)}{\lambda^2 + 4\pi^2 u^2} \leq B(2k)!.$$

If α is the unit vector of the u_i axis, this implies

$$(2.14) \quad \int \frac{u_i^{2k}}{(\sum u_j^2)^k} \frac{\chi_0(du)}{(\lambda^2 + 4\pi^2 u^2)} \leq B(2k)!.$$

By the convexity relationship $\sum_1^n x_i^n \geq n^{1-p}(\sum x_i)^p$, x_i and $p \geq 0$, we have $(\sum u_i^2)^k \leq n^{k-1} \sum u_i^{2k}$. Thus, by substituting in (2.14) and summing from $i = 1$ to $i = k$, we find

$$\int \frac{\chi_0(du)}{\lambda^2 + 4\pi^2 u^2} \leq n^k(2k)! B.$$

The decreasing family $\chi_\lambda = \chi_0/(\lambda^2 + 4\pi^2 u^2)$ being dominated, it follows that there exists a positive bounded measure $\chi'_0 = \lim \chi_\lambda$, for $\lambda \downarrow 0$. By the relationship $4\pi^2 u^2 \chi_\lambda = \chi_0 - \lambda^2 \chi_\lambda$, we get $\chi_0 = \lim 4\pi^2 u^2 \chi_\lambda = 4\pi^2 u^2 \chi'_0$. Thus, (b) is true.

It remains to prove (b) implies (a). Let us at first examine the case $k = 0$. Let Z be a 0-IRF without drift, satisfying (b), i.e., admitting a GC of the form

$$(2.15) \quad K(h) = \int [\cos 2\pi(uh) - 1] \chi'_0(du)$$

for a bounded positive measure χ'_0 without atom at the origin. Let $\chi_\lambda = |\tilde{\lambda}|^2 \chi'_0$ and ζ_λ be the spectral measure and the orthogonal random measure associated with a $\lambda \in \Lambda_k$. The function $1/\tilde{\lambda}$ belongs to $L^2(\mathbf{R}^n, \chi_\lambda)$, and thus the integral

$$Y(x) = \int [(\exp\{-2i\pi(ux)\})/\tilde{\lambda}(u)] \zeta_\lambda(du)$$

exists and defines an SRF $Y(x)$, $x \in \mathbf{R}^n$. For any $\mu \in \Lambda_k$, we may write

$$\int Y(x)\mu(dx) = \int (\tilde{\mu}(u)/\tilde{\lambda}(u))\zeta_\lambda(du) = \int \zeta_\mu(du) = Z(\mu)$$

by $\tilde{\mu}\zeta_\lambda = \tilde{\lambda}\zeta_\mu$. Then (a) is true.

Now let Z be a k -IRF without drift, $k > 0$, suppose (b) is true and prove (a). For $\lambda \in \Lambda_{k-1}$, put $Y_\lambda(x) = Z(\tau_x \lambda - \lambda)(x \in \mathbf{R}^n)$. $Y_\lambda(x)$ is a representation of a 0-IRF and satisfies the relationship

$$\|Y_\lambda(x)\|^2 = \int \frac{|\exp\{-2i\pi(ux)\} - 1|^2 |\tilde{\lambda}|^2}{(4\pi^2 u^2)^k} \chi'_0(du).$$

The function $|\tilde{\lambda}|^2/(4\pi^2 u^2)^k$ being bounded on $\mathbf{R}^n \setminus \{0\}$, and χ'_0 without atom at the origin, it follows that $\|Y_\lambda(x)\|$ is bounded on \mathbf{R}^n . By the result which we have already proved for the 0-IRF, there exists a unique element $Y_\lambda \in H$ such that $Y_\lambda(x) = (U_x - I) Y_\lambda$ and $\Pi_0 Y_\lambda = 0$. The spectral measure associated

with Y_λ is $|\tilde{\lambda}|^2 \chi'_0 / (4\pi^2 u^2)^k$. The continuity of the linear mapping $\lambda \rightarrow Y_\lambda$ from Λ_{k-1} into H is then easy to prove, so that this mapping is a continuous $(k-1)$ -IRF. For $\lambda \in \Lambda_k \subset \Lambda_{k-1}$, obviously

$$(U_x - I)Y_\lambda = Z(\tau_x \lambda) - Z(\lambda) = (U_x - I)Z(\lambda),$$

and thus $Y_\lambda = Z(\lambda)$ up to an invariant. But this invariant is null, for $\Pi_0 Y_\lambda = \Pi_0 Z(\lambda) = 0$, and the equality $Y_\lambda = Z(\lambda)$ holds. This achieves the proof of Theorem 2.3. As an immediate consequence, we may state the following corollary.

Corollary. A $(k+p)$ -IRF (respectively a $(p-1)$ -IRF) with $k \geq 0, p \geq 1$ is the restriction to $\Lambda_{k+p}(\Lambda_{p-1})$ of a k -IRF (a SRF) if and only if one of its GC satisfies an inequality of the form $|K(h)| \leq a + b|h|^{2k+2}$ (resp. is bounded on \mathbf{R}^n).

3. THE BEST LINEAR INTRINSIC ESTIMATORS (BLIE)

In practical applications, we generally have to interpret the concerned phenomenon as a realization of a certain representation $Y(x)$ of a k -IRF Z . Statistical inference is then reasonably possible as far as the GC of Z are concerned. On the contrary, if only one realization of $Y(x)$ is available, it is entirely impossible to specify which particular representation of Z is involved in the experimental data. Thus, only ‘‘authorized’’ integrals, i.e., integrals of the form $\int \lambda(dx) Y(x), \lambda \in \Lambda_k$ or $\lambda \in M_k$, may be assigned a computable variance (because they depend only on Z , and not on the choice of the particular representation $Y(x)$).

Then, if Y_0 is an element of the Hilbert space $H(Y)$ generated by the $Y(x), x \in \mathbf{R}^n$, we shall say that another element $Y^* \in H(Y)$ is an *intrinsic estimator* of Y_0 if the *difference* $(Y^* - Y_0)$ is itself an authorized integral, or a strong limit of authorized integrals. For, in this case only, the variance of the ‘‘error’’ $(Y^* - Y_0)$ uniquely depends on Z and not on the representation—and may be computed at least approximately. In this context, it is natural to develop a theory of the *best linear intrinsic estimators* (BLIE).

The element $Y_0 \in H(Y)$ we have to estimate will be, for instance, the ‘‘value’’ $Y(x_0)$ of $Y(x)$ at a given point $x_0 \in \mathbf{R}^n$, or the integral $\int p(dx) Y(x)$, where the measure p is known, or a derivative of $Y(x)$ at a given point, and so on. Generally speaking, we consider the case $Y_0 = \mathcal{L}(Y)$, where \mathcal{L} is an element of the completed $\tilde{\Lambda}_Y$ of Λ for the norm $\|\lambda\| = \|\int \lambda(dx) Y(x)\|$ (or of the convenient quotient space, if $\|\lambda\|$ is not a norm). But the space $\tilde{\Lambda}_Y$ depends on the choice of the particular representation $Y(x)$, and not only on the k -IRF Z itself, and in practice we do not know which particular representation is involved. For this reason, the operator \mathcal{L} must be taken so that $\mathcal{L}(Y)$ is

defined for any representation Y of Z . In other words, by (1.4), the monomials f^l of degree $\leq k$ necessarily belong to the domain of \mathcal{L} . Or, which is the same, there exists in Λ a sequence $\{\lambda_n\}$ such that $\int \lambda_n(dx) Y(x)$ strongly converges in $H(Y)$ towards the limit $\mathcal{L}Y$ for any representation $Y(x)$ of Z . In particular, this implies the numerical convergence of the sequences $\{\int \lambda_n(dx) f^l(x)\}$ towards limits denoted $\mathcal{L}f^l$. Instead of $\mathcal{L}(Y) = \lim \int \lambda_n(dx) Y(x)$, we shall use the symbolic notation $\mathcal{L}(Y) = \int \mathcal{L}(dx) Y(x)$ for brevity.

In order to estimate $\mathcal{L}(Y)$, we know, say, the elements $Y(x)$, $x \in S$ belonging to a compact set S (the set of the “experimental data”). In other words, the only possible estimators Y^* are in the form $\int \lambda(dx) Y(x)$, for measures $\lambda \in \Lambda(S)$ (i.e., with finite supports included into S) and, more generally, strong limits of such elements, i.e., elements of the form $\int \mathcal{L}^*(dx) Y(x)$ for operators \mathcal{L}^* with “support” in S . For $\lambda \in \Lambda(S)$, the element $\int \lambda(dx) Y(dx)$ is an intrinsic estimator of $\mathcal{L}(Y)$ if and only if λ satisfies the “universality conditions”

$$\int \lambda(dx) f^l(x) = \mathcal{L}f^l,$$

i.e., $\lambda \in \mathcal{L} + \tilde{\Lambda}_k$. Now, if a sequence $\{\lambda_n\}$ in $\Lambda(S) \cap (\mathcal{L} + \tilde{\Lambda}_k)$ is such that $\{\int \lambda_n(dx) Y(x)\}$ strongly converges in $H(Y)$ for a given representation $Y(x)$, it follows from the universality condition that the sequence $\{\int \lambda_n(dx) X(x)\}$ is also convergent for any other representation $X(x)$. In other words, the set of the operators \mathcal{L}^* with support in S and such that $Y^* = \mathcal{L}^*(Y)$ is an intrinsic estimator of $\mathcal{L}(Y)$ (i.e., the closure $\overline{\Lambda(S) \cap (\mathcal{L} + \tilde{\Lambda}_k)}$) does not depend on the choice of the representation $Y(x)$ we have used to define the norm $\|\lambda\| = \|\int \lambda(dx) Y(x)\|$.

This result may be stated in another equivalent manner: if

$$\lambda_n \in \Lambda(S) \cap (\mathcal{L} + \tilde{\Lambda}_k),$$

i.e., $\mathcal{L} - \lambda_n \in \tilde{\Lambda}_k$, the element

$$\int \mathcal{L}(dx) Y(x) - \int \lambda_n(dx) Y(x) = Z(\mathcal{L} - \lambda_n)$$

does not depend on the choice of the representation $Y(x)$. In other words, the strong convergence $\int \lambda_n(dx) Y(x) \rightarrow \mathcal{L}^*(Y)$ for a particular representation $Y(x)$ is equivalent to the convergence $Z(\mathcal{L} - \lambda_n) \rightarrow Z(\mathcal{L} - \mathcal{L}^*)$ and thus implies $\int \lambda_n(dx) X(x) \rightarrow \mathcal{L}^*(X)$ for any other representation $X(x)$.

The variety $\Lambda(S) \cap (\mathcal{L} + \tilde{\Lambda}_k)$ is empty if it is not possible to find $\lambda \in \Lambda(S)$ with $\int \lambda(dx) f^l(x) = \mathcal{L}f^l$, i.e., if there exist coefficients C_l such that $C_l f^l(x) = 0$ for any $x \in S$ and $C_l \mathcal{L}f^l \neq 0$. For this reason, we shall always suppose the monomials f^l linearly independent on S , i.e., $C_l f^l(x) = 0$ for any $x \in S$ implies $C_l = 0$, so that the variety containing the intrinsic estimators on S will never be empty.

In order to determine the BLIE, it remains to write that $Z(\mathcal{L} - \mathcal{L}^*)$ is the projection of 0 into the linear variety $Z(\mathcal{L} - \Lambda(S) \cap (\mathcal{L} + \tilde{\Lambda}_k))$, which is closed and non-empty, by the above considerations. We obtain the condition: $\langle Z(\mathcal{L} - \mathcal{L}^*), Z(\lambda) \rangle = 0$ for any $\lambda \in \Lambda_k(S)$. This condition expresses that the continuous function $y \rightarrow \int (\mathcal{L}(dx) - \mathcal{L}^*(dx))K(x - y)$ is orthogonal to $\Lambda_k(S)$, and thus coincides on S with a linear combination $\mu_l f^l$ (i.e., a polynomial of degree $\leq k$). Finally, the operator \mathcal{L}^* with support in S associated with the BLIE $Y^* = \mathcal{L}^*(Y)$ is characterized by the following conditions:

$$(3.1) \quad \begin{aligned} \mathcal{L}^* f^l &= \mathcal{L} f^l, \\ \int \mathcal{L}^*(dx)K(x - y) &= \int \mathcal{L}(dx)K(x - y) + \mu_l f^l(y) \text{ for all } y \in S; \end{aligned}$$

and the corresponding ‘‘estimation variance’’ is

$$\begin{aligned} &\int (\mathcal{L}(dx) - \mathcal{L}^*(dx))K(x - y)(\mathcal{L}(dy) - \mathcal{L}^*(dy)) \\ &= \int \mathcal{L}(dx)K(x - y)\mathcal{L}(dy) - \int \mathcal{L}(dx)K(x - y)\mathcal{L}^*(dy) + \mu_l \mathcal{L} f^l. \end{aligned}$$

Example 3.1 (The finite case)

If the set S of ‘‘experimental data’’ is finite, the preceding discussion is greatly simplified, because we only have to consider estimators of the form $\int \lambda(dx) Y(x)$ with $\lambda \in \Lambda(S)$. Let $x_\alpha, \alpha = 1, 2, \dots, N$ be the points of S , and write $Y_\alpha, f^l_\alpha, K_{\alpha\beta}$, etc., instead of $Y(x_\alpha), f^l(x_\alpha), K(x_\alpha - x_\beta)$, etc. Then, the BLIE on S for $Y_0 = \mathcal{L}(Y)$ is $Y^* = \lambda^\alpha Y_\alpha$, with coefficients λ^α satisfying the following system:

$$(3.2) \quad \begin{aligned} \lambda^\alpha f^l_\alpha &= \mathcal{L} f^l, \\ \lambda^\alpha K_{\alpha\beta} &= \int \mathcal{L}(dx)K(x - x_\beta) + \mu_l f^l_\beta. \end{aligned}$$

The BLIE $Y^* = \lambda^\alpha Y_\alpha$ always exists, the functions f^l being independent on S , so that the system (3.2) admits solutions. The unicity of Y^* does not imply the regularity of the system (3.2), but only the relationship $\lambda_1^\alpha Y_\alpha = \lambda_2^\alpha Y_\alpha$ a.s. if λ_1 and λ_2 are two distinct solutions. As a matter of fact, the difference $v = \lambda_1 - \lambda_2$ between two solutions will satisfy the system:

$$v^\alpha f^l_\alpha = 0, \quad v^\alpha K_{\alpha\beta} = \mu_l f^l_\beta,$$

from which it follows $\|v^\alpha Y_\alpha\|^2 = v^\alpha K_{\alpha\beta} v^\beta = \mu_l f^l_\beta v^\beta = 0$. It is not difficult to show that the system (3.2) is regular if and only if the following two conditions are fulfilled:

- (a) the matrix $K_{\alpha\beta}$ is strictly conditionally positive definite,
- (b) the functions f^l are linearly independent on S (i.e., $C_l f_\alpha^l = 0 \Rightarrow C_l = 0$).

Example 3.2 (Estimation of a drift)

We suppose now that the k -IRF Z admits as representation a SRF $Z(x) = U_x Z_0$ with $\Pi_0 Z_0 = 0$. By the corollary of Theorem 2.4, it is the case if and only if Z is without drift and admits a bounded GC, say σ . We may suppose that σ is the stationary covariance of the SRF $Z(x) = U_x Z_0$. Any other representation of Z is of the form

$$Y(x) = Z(x) + A_l f^l(x)$$

where the A_l are order-two random variables. In the present context, we shall say that the (random) polynomial $A_l f^l$ is the *drift* of the RF $Y(x)$. We may ask if it is possible to get a BLIE for the drift value $A_l f^l(x)$ at a given point x , and for the coefficients A_l themselves. This new formulation of an old problem (see, for instance, [5]) will remain valid if Z is only *locally* stationary (see Section 7), but not stationary, so that it will be possible to define a locally significant notion of a drift and to get a BLIE for it.

The first question that arises is whether $A_l \in H(Y)$ and $A_l = \mathcal{L}_l Y$ for operators \mathcal{L}_l including the monomials f^s in their domains. For this purpose, take $\lambda_l \in \Lambda$ satisfying the ‘‘universality condition’’ $\int \lambda_l(dx) f^s(x) = \delta_l^s$ ($= 0$ if $l \neq s$, $= 1$ if $l = s$), and denote ϕ_t the density of the Gaussian centered law with variance t . The regularized functions

$$\phi_t(x, t) = \int \lambda_l(dy) \phi_t(x + y)$$

still satisfy the universality conditions

$$\int \phi_t(x, t) f^s(x) dx = \delta_t^s,$$

and we get

$$\int \phi_t(x, t) Y(x) dx = \int \phi_t(x, t) Z(x) dx + A_l.$$

For $t \rightarrow \infty$, the ergodic theorem asserts that $\int \phi_t(x, t) Z(x) dx$ strongly converges towards 0. Thus, the operators \mathcal{L}_l defined by

$$\mathcal{L}_l(Y) = \lim_{t \rightarrow \infty} \int \phi_t(x, t) Y(x) dx$$

satisfy $\mathcal{L}_l(Y) = A_l$ for any representation Y , as required, and in particular $\mathcal{L}_l f^s = \delta_l^s$.

Thus, there exists a BLIE for $A_l = \mathcal{L}_l Y$, say $A_l^* = \mathcal{L}_l^*(Y)$. The operator \mathcal{L}_l^* (with support in the compact set S of experimental data) is characterized by the system (3.1) written with σ instead of K . But, in the second relationship (3.1), the term $\int \mathcal{L}_l(dx)\sigma(x - y)$ vanishes, by the ergodic theorem. Finally, we get the following system:

$$(3.3) \quad \int \mathcal{L}_l^*(dx)f^s(x) = \delta_l^s,$$

$$\int \mathcal{L}_l^*(dx)\sigma(x - y) = \mu_{ls}f^s(y) \text{ for all } y \in S.$$

The matrix (μ_{ls}) of the ‘‘Lagrange parameters’’ admits a very simple interpretation: it is identical with the covariance matrix $\langle A_l - A_l^*, A_s - A_s^* \rangle$, as is easy to verify by direct calculation. It is also easy to show that, for a given $x \in \mathbb{R}^n$, $A_l^* f^l(x)$ is the BLIE for the drift value in x , $A_l f^l(x)$.

Let $H_Y(S)$ be the Hilbert space generated by the $Y(x)$, $x \in S$, and $H_k(S) \subset H_Y(S)$ the Hilbert space generated by the $\int \lambda(dx)Y(x) = Z(\lambda)$, $\lambda \in \Lambda_k(S)$. By the system (3.3), the A_l^* belong to the orthogonal $H_k^\perp(S)$ of $H_k(S)$ in $H_Y(S)$. Conversely, if $Y_0 \in H_k^\perp(S)$, this element is of the form $Y_0 = C^l A_l^*$. For, if $x \in S$, the relationship $Y(x) = Z(\delta_x - f_x^l \mathcal{L}_l^*) + A_l^* f^l(x)$ implies $\langle Y_0, Y(x) \rangle = \langle Y_0, A_l^* \rangle f^l(x)$, and the projection of Y_0 into the orthogonal of the A_l vanishes. Thus, $H_k^\perp(S)$ is spanned by the A_l^* . We shall say that $H_k^\perp(S)$ is the space of the drift and $H_k(S)$ the space of the residuals.

Then, by relationship $Y(x) = Z(\delta_x - f^l(x)\mathcal{L}_l^*) + A_l^* f^l(x)$, we conclude that for $x \in S$ the BLIE $A_l^* f^l(x)$ of the drift value at $x \in S$ is the projection of $Y(x)$ itself into the drift space $H_k^\perp(S)$. But this result does not remain true for $x \notin S$.

4. THE TURNING BANDS METHOD

In practical applications, one must sometimes simulate realizations of a given random function in \mathbb{R}^2 or \mathbb{R}^3 , and we will now describe a procedure which may be used in the isotropic case for SRF as well as for k -IRF.

First, let $t \rightarrow Y(t)$ be a SRF on \mathbb{R}^1 , C_1 its covariance, and s a unit vector in \mathbb{R}^n . Then, $x \rightarrow Z_s(x) = Y(\langle x, s \rangle)$ is a SRF on \mathbb{R}^n and its covariance is defined by $\langle Z_s(x), Z_s(y) \rangle = C_1(\langle (x - y), s \rangle)$. If now s is substituted by the unit random vector, $Z_s(x)$ is changed into a SRF $Z(x)$, the covariance C of which is defined by

$$C(h) = \int C_1(\langle h, s \rangle) \varpi_n(ds)$$

where ϖ_n is the probability concentrated on the unit sphere and invariant

under rotations. Clearly, the covariance C is isotropic, i.e., there exists a function C_n on \mathbf{R}_+ such that $C(h) = C_n(|h|)$, say explicitly

$$(4.1) \quad C_n(r) = 2\Gamma(\frac{1}{2}n)\pi^{-\frac{1}{2}}(\Gamma[\frac{1}{2}(n-1)])^{-1} \int_0^1 C_1(vr)(1-v^2)^{\frac{1}{2}(n-3)}dv.$$

Let G be the positive measure on \mathbf{R}_+ such that

$$C_1(r) = \int_0^\infty \cos 2\pi \rho r G(d\rho).$$

An easy calculation gives

$$(4.2) \quad C_n(r) = 2^{\frac{1}{2}n-1}\Gamma(\frac{1}{2}n) \int_0^\infty (2\pi\rho r)^{1-\frac{1}{2}n} J_{\frac{1}{2}n-1}(2\pi\rho r) G(d\rho)$$

where $J_{\frac{1}{2}n-1}$ is the classical Bessel function. But we know from the Bochner theorem that (4.2) is the general form of an isotropic covariance on \mathbf{R}^n , so that the mapping $C_1 \rightarrow C_n$ is one-to-one from the covariances on \mathbf{R}^1 onto the isotropic covariances on \mathbf{R}^n . In other words, for any isotropic covariance C_n , there exists a unique covariance C_1 on \mathbf{R}^1 such that (4.1) holds. For instance, if $n = 3$, we find $C_3(r) = \int_0^1 C_1(vr)dv$, and, conversely, $C_1(r) = drC_3(r)/dr$.

If an isotropic covariance C_n on \mathbf{R}^n is given, the procedure described above (i.e., the “turning bands” method) will yield a realization of a SRF with covariance C_n , if it is applied to a realization of a SRF Y with the corresponding covariance C_1 on \mathbf{R}^1 .

The turning bands method may also be used in order to construct representations of a k -IRF with a given isotropic GC on \mathbf{R}^n , for it is easy to see from Theorem 2.1 that the relationship (4.1) defines a one-to-one mapping from the k -GC on \mathbf{R}^1 onto the isotropic k -GC on \mathbf{R}^n . Moreover, the monomials r^α , $\alpha \geq 0$ are eigen-functions for the turning bands operator (4.1), so that the polynomial isotropic GC on \mathbf{R}^n (see Section 5 below) are generated by the polynomial GC on \mathbf{R}^1 . This procedure was used by Orfeuill [10], who constructed the originals of the Figures 1, 2 and 3 above. These figures give realizations of IRF order $k = 0, 1, 2$ with GC proportional to $-|h|$, $|h|^3$ and $-|h|^5$ respectively. The representations chosen by Orfeuill vanish at the center of the figures, as well as their derivatives up to order k , and this explains the feature of Figure 3 in which nothing resembles an isotropic and stationary phenomenon any longer.

5. THE POLYNOMIAL GC

Let us denote by T_n the turning bands operator defined in (4.1), so that for any $\alpha > 0$ the function $r \rightarrow r^\alpha$ is an eigen-function for T_n . If $\alpha = 2p + 1$ is an odd integer, we find explicitly

$$T_n r^{2p+1} = B_n p r^{2p+1}, \quad B_{n,p} = p! \Gamma(\frac{1}{2}n) / (\pi^{\frac{1}{2}} \Gamma(p + \frac{1}{2}(1+n))).$$

As noted above, the function K_1 defined by

$$(5.1) \quad K_1(r) = \sum_{p=0}^k (-1)^{p+1} a_p r^{2p+1} / (2p+1)!$$

is a k -GC on R^1 if and only if the function

$$(5.2) \quad K_n(r) = \sum_{p=0}^k (-1)^{p+1} a_p B_{np} r^{2p+1} / (2p+1)!$$

is itself an isotropic K -GC on R^n . With the notations (5.1) and (5.2), the conditions the coefficients a_p have to satisfy do not depend on the dimension n .

From the relationship

$$(-1)^{p+1} \frac{r^{2p+1}}{(2p+1)!} = 4 \int_0^\infty \frac{\cos(2\pi\rho r) - P_p(2\pi\rho r)}{(4\pi^2\rho^2)^{p+1}} d\rho,$$

it follows that the function K_1 defined in (5.1) admits (up to an even polynomial of degree $\leq 2k$) the representation

$$K_1(r) = \int_0^\infty \frac{\cos 2\pi\rho r - 1_B(\rho) P_k(2\pi\rho r)}{(4\pi^2\rho^2)^{k+1}} g(\rho) d\rho,$$

$$g(\rho) = 4 \sum_{p=0}^k a_p (4\pi^2\rho^2)^{k-p}.$$

In other words, K_1 (and K_n) are k -GC if and only if the polynomial $\Pi(x) = \sum a_p x^{2(k-p)}$ is ≥ 0 for x real.

Lemma 5. An even polynomial Π with real coefficients and degree $2k$ satisfies $\Pi(x) \geq 0$ for x real if and only if there exists a polynomial Φ with real coefficients and degree k such that $\Pi(x) = |\Phi(ix)|^2$.

Proof. The if part is obvious. Conversely, let Π be an even polynomial with real coefficients and degree $2k$ such that $\Pi(x) \geq 0$ for x real. Then (up to a positive multiplicative constant) $\Pi(z)$ is the product of terms of the type

$$(z + ib)(z - ib) = (b + iz)(b - iz),$$

with b real, or

$$(z - a)(z + a)(z - \bar{a})(z + \bar{a}) = (\alpha + iz)(\bar{\alpha} + iz)(\alpha - iz)(\bar{\alpha} - iz),$$

with $\alpha = ia$ and $a = a_1 + ia_2$, a_1, a_2 real and $a_1 \neq 0$. Each of these terms is of the form $\phi(iz)\phi(-iz)$ for a polynomial ϕ with real coefficients. The lemma follows.

Let us now give the general form of the k -IRF with polynomial GC on \mathbf{R}^1 .

Theorem 5. A k -IRF Z on \mathbf{R}^1 admits a polynomial GC if and only if it admits a representation Y such that

$$(5.3) \quad Y(x) = b_0W(x) + b_1 \int_0^x W(\xi)d\xi + \dots + b_k \int_0^x \frac{(x - \xi)^{k-1}}{(k - 1)!} W(\xi)d\xi$$

where the b_p are real coefficients and $W(x)$ a representation of a 0-IRF with the GC $K(h) = -|h|$.

Proof. If Z admits the representation (5.3),

$$X(x) = \int_0^x (x - \xi)^{k-1} W(\xi)/(k - 1)!d\xi$$

is a representation of a k -IRF with the GC $(-1)^{k+1}|h|^{2k+1}/(2k + 1)!$. If D^p denotes the derivation order p , we have

$$Y(x) = (\sum b_p D^{k-p})X(x)$$

and the k -IRF Z admits the GC

$$K(h) = (\sum b_p D^{k-p}) (\sum (-1)^{k-p} D^{k-p}) |h|^{2k+1}/(2k + 1)!$$

i.e., a polynomial GC.

Conversely, let Z be a k -IRF on \mathbf{R}^1 admitting a GC $K(h)$ of the form (5.1). From Lemma 5 and the preceding considerations we may write (up to an even polynomial of degree $\leq 2k$)

$$K(h) = 2 \int_{-\infty}^{\infty} \frac{\cos 2\pi u h - 1_B(u)P_k(2\pi u h)}{(4\pi^2 u^2)^{k+1}} |\Phi(2i\pi u)|^2 du,$$

$$\Phi(u) = \sum_{p=0}^k b_p u^{k-p},$$

with convenient real coefficients b_p . Now let $\alpha \in \tilde{\Lambda}_k$ be the function defined by its Fourier transform $\tilde{\alpha}(u) = (2i\pi u)^k \exp(-u^2)$, and ζ_α the random orthogonal measure such that $Z(\tau_h \alpha) = \int \exp(-2i\pi u h) \zeta_\alpha(du)$. The corresponding spectral measure χ_α is

$$\chi_\alpha(du) = 2[|\tilde{\alpha}|^2 |\Phi(2i\pi u)|^2 / (4\pi^2 u^2)^{k+1}] du,$$

and, for any $\lambda \in \Lambda_k$ the Fourier transform of which is $\tilde{\lambda}, \tilde{\lambda}/\tilde{\alpha} \in L^2(\mathbf{R}^1, \chi_\alpha)$ implies

$$(5.4) \quad Z(\lambda) = \int (\tilde{\lambda}(u)/\tilde{\alpha}(u)) \zeta_\alpha(du).$$

The function $(\exp(-2i\pi u x) - 1)(-2i\pi u)^k / [\tilde{\alpha}(u)\Phi(-2i\pi u)]$ is also in $L^2(\mathbf{R}^1, \chi_\alpha)$, and thus there exists a random function $W(x)$ defined by

$$(5.5) \quad W(x) = \int_{-\infty}^{+\infty} \frac{(\exp\{-2i\pi ux\} - 1)(-2i\pi u)^k}{\tilde{\alpha}(u)\Phi(-2i\pi u)} \zeta_{\alpha}(du).$$

If $\lambda \in \Lambda_0$, we have

$$\int \lambda(dx)W(x) = \int_{-\infty}^{+\infty} \tilde{\lambda}(u)(-2i\pi u)^k/(\tilde{\alpha}(u)\Phi(-2i\pi u))\zeta_{\alpha}(du)$$

and thus

$$\begin{aligned} \left\| \int \lambda(dx)W(x) \right\|^2 &= \int \frac{|\tilde{\lambda}|^2(4\pi^2u^2)^k}{|\tilde{\alpha}|^2|\Phi(-2i\pi u)|^2} \chi_{\alpha}(du) \\ &= 2 \int (|\tilde{\lambda}|^2/4\pi^2u^2)du. \end{aligned}$$

From this relationship, it follows that $\lambda \rightarrow \int \lambda(dx)W(x)$ is a 0-IRF with the GC

$$2 \int_{-\infty}^{+\infty} \frac{\cos 2\pi uh - 1}{4\pi^2u^2} = -|h|.$$

Now let $Y(x)$ be defined by (5.3) and (5.5), and $\mu \in \Lambda_k$. By (5.5), we may write

$$\begin{aligned} \int \mu(dx)Y(x) &= \int \frac{\tilde{\mu}(u)(-2i\pi u)^k \sum b_p/(-2i\pi u)^p}{\tilde{\alpha}(u)\Phi(-2i\pi u)} \zeta_{\alpha}(du) \\ &= \int (\tilde{\mu}(u)/\tilde{\alpha}(u))\zeta_{\alpha}(du). \end{aligned}$$

Thus, by (4.4), we have $\int \mu(dx)Y(x) = Z(\mu)$, and $Y(x)$ is a representation of Z . This completes the proof.

It is very easy to construct realizations of a 0-FAI with GC $-|h|$ on \mathbf{R}^1 . For instance, we may choose a Wiener-Lévy process (Brownian motion). By Theorem 5, we are thus able to construct realizations of a k -IRF with a given polynomial GC, and by the turning bands method we may also obtain realizations of a k -IRF with a given polynomial covariance isotropic on \mathbf{R}^n .

6. k -IRF LOCALLY STATIONARY

Any stationary covariance is also a k -GC, and thus there exist k -IRF's which admit stationary representations and may be called *stationary k -IRF's*. Any representation of a stationary k -IRF is then of the form $Y(x) = Y_0(x) + P(x)$, where $Y_0(x)$ is a SRF and $P(x)$ a polynomial of degree $\leq k$ with random coefficients A_i . From Theorem 2.4, we know that a k -IRF is stationary if and only if it admits a GC bounded on \mathbf{R}^n . This condition is never fulfilled by a polynomial GC. But, on the other hand, we shall see that the k -IRF with polynomial GC are always locally stationary in the following sense.

Definition. A k -IRF Z is *locally stationary* if there exist a bounded open set $V \subset \mathbf{R}^n$ and a representation $Y(x)$ of Z such that $Y(x)$ coincides on V with a SRF $Y_V(x)$.

Clearly, if Z is locally stationary on an open set V , it is still so on any translation V' of V , but the SRF $Y_{V'}$ cannot be identical to Y_V if Z itself is not stationary. This property is important for practical applications, because it leads to a locally significant notion of the drift, i.e., the polynomial $\phi(x) = Y(x) - Y_V(x)$, $x \in V$. In particular, the estimation of $P(x)$ in a point $x \in V$, or of its coefficients A_i is then possible by the technique of the best linear intrinsic estimator (BLIE, see Section 3). There exist k -IRF's which cannot be locally stationary. For instance, if an analytic k -IRF were locally stationary, it would be stationary on the whole space \mathbf{R}^n .

Theorem 6. Any k -IRF with polynomial GC is locally stationary on any bounded open set V .

Proof. If the theorem is true in \mathbf{R}^1 , it will be true in \mathbf{R}^n for any $n > 0$ by the turning bands method. Then, suppose $n = 1$, and examine at first the case $k = 0$.

Let Z be a 0-IRF with the GC $-|h|$ on \mathbf{R}^1 , and choose the representation $W(x) = Z(\delta_x - \frac{1}{2}(\delta_L + \delta_0))$, for an arbitrary real $L > 0$. The covariance of $W(x)$ satisfies

$$\langle W(x), W(y) \rangle = \frac{1}{2}L - |x - y|$$

for $x, y \in [0, L]$, and, in particular, $W(0) + W(L) = 0$. Thus, it is possible to define a periodic random function $Y_0(x)$ continuous on \mathbf{R}^1 by putting

$$Y_0(x) = W(x) \text{ if } x \in [0, L]$$

and

$$Y_0(x) = (-1)^k Y_0(x + kL) \text{ for } kL \leq x \leq (k + 1)L, k \text{ integer.}$$

The period of $Y_0(x)$ is $2L$. Let C denote the function with period $2L$ defined by $C(h) = \frac{1}{2}L - |h|$ for $|h| \leq L$. For $x = x_0 + kL$, $y = y_0 + k'L$, $x_0, y_0 \in [0, L]$, we have

$$\begin{aligned} \langle Y_0(x), Y_0(y) \rangle &= (-1)^{k+k'} [\frac{1}{2}L - |x_0 - y_0|] \\ &= (-1)^{k+k'} [\frac{1}{2}L - |x - y - (k - k')L|] \\ &= C(|x - y|) \end{aligned}$$

by the relationship $C(h + kL) = (-1)^k C(h)$.

Thus $Y_0(x)$ is a SRF and its covariance is $C(h)$. By $W(x) = Y_0(x)$ for $0 \leq x \leq L$, Z is then locally stationary on $[0, L]$.

Now let Z be a k -IRF with polynomial covariance on \mathbf{R}^1 , and $W(x)$ the 0-IRF occurring in its representation (5.3) (Theorem 5) which may be chosen in such a way that $W(0) + W(L) = 0$. By the preceding part of the proof, we have $W(x) = Y_0(x)$ if $0 \leq x \leq L$ for a periodic SRF $Y_0(x)$, the covariance of which is $C(h)$. By the Fourier expansion,

$$C(h) = \sum_{q=0}^{\infty} 4L\pi^{-2}(2q + 1)^{-2} \cos(2q + 1)\pi(h/L),$$

there exists a sequence $\{C_l\}$, $l = \pm 1, \pm 3, \dots$ of complex orthogonal variables, with $C_{-l} = \bar{C}_l$ and $\|C_l\|^2 = 4L/(\pi l)^2$ such that

$$Y_0(x) = \sum_l C_l \exp(-i\pi l x/L).$$

For p integer > 0 , the SRF $Y_p(x)$ defined by

$$Y_p(x) = \sum_l (iL/\pi l)^p C_l \exp(-i\pi l x/L)$$

admits the covariance

$$C_p(h) = 4 \sum_{q=0}^{\infty} L^{2p+1} ((2q + 1)\pi)^{-2p} \cos(2q + 1)\pi(h/L),$$

and is equal to $\int_0^x (x - \xi)^{p-1} Y_0(\xi) / (p-1)! d\xi$ up to a random polynomial of degree $\leq p$. By the formula (5.3), Theorem 5, the SRF $\sum b_p Y_p(x)$ is thus equal to $Y(x)$, up to a random polynomial of degree $\leq k$ for any $x \in [0, L]$. Thus, Z is locally stationary on $[0, L]$.

Corollary. Let Z be a k -IRF with the GC

$$K(h) = \sum_0^k (-1)^p a_p |h|^{2p+1} / (2p + 1)!$$

on \mathbf{R}^1 , and $C(h)$ the covariance of the SRF with period $2L$ which coincides on $[0, L]$ with a representation of Z . Then,

$$C(h) = 4 \sum_{q=0}^{\infty} B_q \cos(2q + 1)\pi(h/L),$$

$$B_q = \sum_{p=0}^k a_p L^{2p+1} \pi^{-2p-2} (2q + 1)^{-2p-2}.$$

The corollary follows easily from Lemma 5 and

$$\sum a_p x^{2(k-p)} = |\sum b_p (ix)^{k-p}|^2.$$

Note that $C(h) = K(h)$ for $|h| \leq L$, up to an even polynomial of degree $\leq 2k$.

References

- [1] CRAMÉR, H. AND LEADBETTER, M. R. (1969) *Stationary and Related Stochastic Processes*. Wiley, New York.
- [2] FELLER, W. (1957) *An Introduction to Probability Theory and its Applications*. Vol. 1. Wiley, New York.
- [3] GUELFAND, M. AND VILENKIN, N. Y. (1961) *Nekotorye primeneniya garmonitsheskovo analiza*. Moscow.
- [4] GUELFAND, M. AND VILENKIN, N. Y. (1967) *Les Distributions*. Vol. 4. Dunod, Paris.
- [5] GRENANDER, U. AND ROSENBLATT, M. (1957) *Stationary Time Series*. Wiley, New York.
- [6] GUIBAL, D. (1972) Simulations de schémas intrinsèques. *Internal report*, Centre de Morphologie Mathématique, Fontainebleau.
- [7] HUIJBREGTS, CH. AND MATHERON, G. (1970) Universal Kriging — an optimal method for estimating and contouring in trend surface analysis. *CIMM International Symposium*, Montreal.
- [8] MATHERON, G. (1969) Le Krigeage Universel. *Fasc. No. 1*, Cahiers du Centre de Morphologie Mathématique, Fontainebleau.
- [9] MATHERON, G. (1971) The theory of regionalized variables, and its applications. *Fasc. No. 5*, Cahiers du Centre de Morphologie Mathématique, Fontainebleau.
- [10] ORFEUIL, J. P. (1972) Simulation du Wiener-Lévy et de ses intégrales. *Internal report*, Centre de Morphologie Mathématique, Fontainebleau.
- [11] YAGLOM, A. M. (1962) *Stationary Random Functions*. Prentice Hall, Englewood Cliffs, N. J.
- [12] WIENER, N. AND MASANI, P. (1958) The prediction theory for multivariate stochastic processes. I. *Acta Math.* **98**, 111–150; II. *Acta Math.* **99**, 93–137.