

Open Research Online

The Open University's repository of research publications and other research outputs

The iterated minimum modulus and conjectures of Baker and Eremenko

Journal Item

How to cite:

Osborne, John W.; Rippon, Philip J. and Stallard, Gwyneth M. (2019). The iterated minimum modulus and conjectures of Baker and Eremenko. *Journal d'Analyse Mathematique*, 139(2) pp. 521–558.

For guidance on citations see [FAQs](#).

© 2019 Journal d'Analyse Mathematique



<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1007/s11854-019-0065-z>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

THE ITERATED MINIMUM MODULUS AND CONJECTURES OF BAKER AND EREMENKO

J.W. OSBORNE, P.J. RIPPON AND G.M. STALLARD

ABSTRACT. In transcendental dynamics significant progress has been made by studying points whose iterates escape to infinity at least as fast as iterates of the maximum modulus. Here we take the novel approach of studying points whose iterates escape at least as fast as iterates of the *minimum* modulus, and obtain new results related to Eremenko's conjecture and Baker's conjecture, and the rate of escape in Baker domains. To do this we prove a result of wider interest concerning the existence of points that escape to infinity under the iteration of a positive continuous function.

For Walter Hayman on the occasion of his ninetieth birthday.

1. INTRODUCTION

Denote the n th iterate of a function f by f^n , for $n \in \mathbb{N}$. If f is a transcendental entire function then the *Fatou set* $F(f)$ is the set of points $z \in \mathbb{C}$ such that the family of functions $\{f^n : n \in \mathbb{N}\}$ is normal in some neighbourhood of z and the *Julia set* $J(f)$ is the complement of $F(f)$. We refer to [4, 5, 8, 19] for the fundamental properties of these sets and an introduction to complex dynamics.

The *escaping set* $I(f) = \{z \in \mathbb{C} : f^n(z) \rightarrow \infty \text{ as } n \rightarrow \infty\}$ was first studied for a general transcendental entire function f by Eremenko [9]. In recent years, the *fast escaping set* $A(f)$ has played a significant role in transcendental dynamics, for example, in progress on *Eremenko's conjecture*, that all the components of $I(f)$ are unbounded, and on *Baker's conjecture*, that if f has order at most $1/2$, minimal type, then all the components of $F(f)$ are bounded. Despite many partial results, both conjectures remain open.

The set $A(f)$ was introduced in [6] and consists of those points whose iterates under f eventually grow at least as fast as iterates of the maximum modulus, $M(r) = \max_{|z|=r} |f(z)|$. It can be defined as follows:

$$A(f) = \bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(A_R(f)), \text{ where } A_R(f) = \{z : |f^n(z)| \geq M^n(R), \text{ for } n \in \mathbb{N}\}.$$

Here $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, $M^n(r)$ denotes the n th iterate of the function $r \mapsto M(r)$, and $R > 0$ is such that $M^n(R) \rightarrow \infty$ as $n \rightarrow \infty$. Note that there always exists $R > 0$ such that, for $r \geq R$, we have $M(r) > r$ and hence $M^n(R) \rightarrow \infty$ as $n \rightarrow \infty$, and the definition of $A(f)$ is independent of the choice of such an R .

The set $A(f)$ has many strong properties [29, 32] and was used in [37] to show that the escaping set $I(f)$ is either connected or has infinitely many unbounded components. See [25] and [38] for other partial results on Eremenko's conjecture.

2010 *Mathematics Subject Classification.* Primary 37F10, Secondary 30D05.
The last two authors were supported by the EPSRC grant EP/K031163/1.

In this paper, we study those points whose iterates under f eventually grow at least as fast as iterates of the *minimum modulus*, $m(r) = \min_{|z|=r} |f(z)|$. Replacing $M(r)$ by $m(r)$ in the definition of $A(f)$ does not in general yield a subset of $I(f)$. Indeed, if the function $m(r)$ is bounded, then its iterates tell us nothing about $I(f)$; this is the case, for example, when f is in the Eremenko-Lyubich class \mathcal{B} of transcendental entire functions with a bounded set of singular values (that is, critical values and asymptotic values).

It turns out, however, that iterating the minimum modulus is of significant interest for the many entire functions with the property that

$$(1.1) \quad \text{there exists } r > 0 \text{ with } m^n(r) \rightarrow \infty \text{ as } n \rightarrow \infty.$$

We introduced condition (1.1) in [23] in the context of investigating the connectedness properties of the set $I^+(f)$ of points at which the iterates of f form an unbounded sequence. In this paper, we make a deeper study of the condition (1.1), and in this way obtain new results related to Eremenko's conjecture and Baker's conjecture, and about the rate of escape in Baker domains (defined in Section 7).

In order to work with (1.1) and, in particular, to identify transcendental entire functions for which (1.1) holds, it is useful to introduce the function

$$\tilde{m}(r) := \max_{0 \leq s \leq r} m(s), \quad \text{for } r \in [0, \infty).$$

We show that (1.1) is true if and only if

$$(1.2) \quad \text{there exists } R > 0 \text{ such that } \tilde{m}(r) > r, \text{ for } r \geq R.$$

In fact, in Section 2 we prove a general result about escaping points of positive continuous functions, which is of wider interest.

Using the condition (1.2) and several classical results about the size of the minimum modulus, we can show that condition (1.1) holds for many classes of entire functions. The terminology used in the following result is explained in Section 3.

Theorem 1.1. *Let f be a transcendental entire function. Then (1.1) holds if*

- (a) f is of order less than $1/2$, or
- (b) f has finite order and Fabry gaps, or
- (c) f has Hayman gaps, or
- (d) f exhibits the pits effect, as defined by Littlewood and Offord, or
- (e) f has a multiply connected Fatou component.

In Section 8 we give several further examples of familiar entire functions that satisfy the condition (1.1), such as $f(z) = 2z(1 + e^{-z})$.

We note that if the condition (1.1) holds, then it holds for many different values of r but not all. For example, we shall see in Section 2 that if (1.1) holds, then for different values of r the sequence $(m^n(r))$ may be bounded or it may tend to infinity arbitrarily slowly.

In contrast, it is clear that the *fastest* rate at which $m^n(r)$ can tend to infinity must be related to the growth of $\tilde{m}^n(r)$; indeed, this rate is always attained.

Theorem 1.2. *Let f be a transcendental entire function such that (1.1) holds. Then there exists $R > 0$ such that*

$$(1.3) \quad \tilde{m}^n(R) \rightarrow \infty \text{ as } n \rightarrow \infty,$$

and, for any such R , there exists $r \geq R$ such that

$$(1.4) \quad m^n(r) \geq \tilde{m}^n(R), \text{ for } n \in \mathbb{N}.$$

Moreover, for every r and R satisfying (1.3) and (1.4), the set

$$\bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(\{z : |f^n(z)| \geq \tilde{m}^n(R), \text{ for } n \in \mathbb{N}\})$$

is independent of the choice of R and is equal to the set

$$\bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(\{z : |f^n(z)| \geq m^n(r), \text{ for } n \in \mathbb{N}\}).$$

Whenever (1.1) holds we define the following set, by analogy with $A(f)$:

$$V(f) = \bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(V_R(f)), \text{ where } V_R(f) = \{z : |f^n(z)| \geq \tilde{m}^n(R), \text{ for } n \in \mathbb{N}\}.$$

By Theorem 1.2, the set $V(f)$ is well defined and independent of R , provided R is so large that $\tilde{m}(r) > r$ for $r \geq R$. Moreover, $V(f)$ is completely invariant under f .

Since $M^n(r) \geq \tilde{m}^n(r)$ for $n \in \mathbb{N}$, we always have $A(f) \subset V(f)$ and thus $V(f) \neq \emptyset$. Perhaps surprisingly, there are many classes of functions for which $V(f) = A(f)$. In fact, as we show in Section 5, all the functions listed in Theorem 1.1 have this property provided they satisfy a certain regularity condition.

The following result shows that the conclusions of both Eremenko's conjecture and Baker's conjecture hold for any function for which $V(f) = A(f)$.

Theorem 1.3. *Let f be a transcendental entire function such that (1.1) holds, so that the set $V(f)$ is well defined.*

(a) *We have $V(f) = A(f)$ if and only if there exist $r \geq R > 0$ such that*

$$m^n(r) \geq M^n(R), \text{ for } n \in \mathbb{N}, \quad \text{and} \quad M^n(R) \rightarrow \infty \text{ as } n \rightarrow \infty.$$

(b) *If the equivalent conditions in part (a) hold, then*

- (i) $A_R(f)$, $V(f)$ and $I(f)$ are spiders' webs, and
- (ii) $F(f)$ has no unbounded components.

Here, a set E is a *spider's web* if it is connected and there exists a sequence (G_n) of bounded, simply connected domains such that

$$G_n \subset G_{n+1}, \quad \partial G_n \subset E, \text{ for } n \in \mathbb{N}, \quad \text{and} \quad \bigcup_{n \in \mathbb{N}} G_n = \mathbb{C}.$$

Part (b) of Theorem 1.3 shows that if $V(f) = A(f)$, then the set $A_R(f)$ is a spider's web, a property that has several strong consequences [32]. Showing that $A_R(f)$ is a spider's web is, in essence, the approach used to prove all partial results on Baker's conjecture prior to the recent papers [21, 33]; see [2, 16, 17, 31] and the discussion in [33, Introduction].

In view of Theorem 1.3, it is natural to ask the following.

Question 1.4. *Let f be a transcendental entire function such that (1.1) holds. Do the following conclusions hold under a weaker hypothesis than $V(f) = A(f)$?*

- (1) $V(f)$ and $I(f)$ are spiders' webs;
- (2) $F(f)$ has no unbounded components.

Various recent results suggest that a significant weakening of the hypothesis here that $V(f) = A(f)$ is indeed plausible. For example, by [35] there exist entire functions of order less than $1/2$ (so (1.1) holds) for which $A_R(f)$ is not a spider's web. Thus, by the discussion after Theorem 1.3, we have $V(f) \neq A(f)$, and yet it can be shown using [33, Theorems 1.1 and 2.3] that properties (1) and (2) hold. Also, the most recent work on Baker's conjecture [21, Corollary 1.2] shows that property (2) holds whenever f is a real function of order at most $1/2$, minimal type, with only real zeros. Note, however, that there are transcendental entire functions of order greater than $1/2$ for which (1.1) holds and for which the Fatou set has an unbounded component (see Example 8.1), so some hypothesis is needed in addition to (1.1).

As a step towards making progress on Question 1.4, we prove two main results, the first of which is a refinement of [23, Theorem 1.2]. There we showed that if (1.1) holds, then $I^+(f)$ is connected, where $I^+(f)$ is the set of points at which the iterates of f form an unbounded sequence. We now define $V^+(f)$ to be the set derived from $V(f)$ in the same way that $I^+(f)$ is derived from $I(f)$ – that is, by adding those points for which only a *subsequence* of iterates satisfies its defining property. Thus

$$V^+(f) = \bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(V_R^+(f)),$$

where

$$V_R^+(f) = \{z : \exists (n_j) \text{ such that } |f^{n_j}(z)| \geq \tilde{m}^{n_j}(R), \text{ for } j \in \mathbb{N}\}.$$

Here, (n_j) is a strictly increasing sequence of positive integers that in general depends on z , and $R > 0$ is such that $\tilde{m}(r) > r$ for $r \geq R$.

The following is our refinement of [23, Theorem 1.2].

Theorem 1.5. *Let f be a transcendental entire function such that (1.1) holds. Then the set $V^+(f)$, and also the set $I^+(f)$, is a weak spider's web.*

By a *weak spider's web* we mean a connected set whose complement contains no unbounded closed connected sets. The name arises from the fact that a spider's web has the stronger property that its complement contains no unbounded connected sets; see Figure 1 for an illustration of a set that is a weak spider's web but not a spider's web.

Remark If we could replace $V^+(f)$ by $V(f)$, or $I^+(f)$ by $I(f)$, in Theorem 1.5, then this would show that the conclusion of Eremenko's conjecture holds for all entire functions satisfying (1.1). Indeed, if $V(f)$ is connected, then $I(f)$ is connected; see Theorem 4.3.

In order to prove Theorem 1.5, we show that a major consequence of the condition (1.1) is that certain sufficiently long continua must contain a point of $V^+(f)$; see Theorem 6.2 for a detailed statement. We also use this fact about long continua meeting $V^+(f)$ to obtain our second result related to Question 1.4. This second result strengthens several results of Zheng [43], which relate the dynamical behaviour of a transcendental entire function f in an unbounded component of $F(f)$ to the size of the minimum modulus of f , and hence are related to Baker's conjecture.




FIGURE 1. A weak spider's web that is not a spider's web

First we restate the results of Zheng below, using the notation of this paper. (See Section 7 for the definition of a wandering domain.)

(a) If

$$m(r) > r \text{ for an unbounded sequence of values of } r$$

and U is an unbounded component of $F(f)$, then $U \subset I^+(f)$; see [43, Theorem 2].

(b) If

$$\limsup_{r \rightarrow \infty} \frac{m(r)}{r} = \infty$$

and U is an unbounded component of $F(f)$, then U is a wandering domain and $U \subset I^+(f)$; see [43, Theorem 1].

(c) If f has order less than $1/2$ and U is an unbounded component of $F(f)$, then U is a wandering domain and $U \subset Z^+(f)$; see [43, Theorem 3].

The set $Z^+(f)$ in result (c) is defined as follows:

$$Z^+(f) := \left\{ z : \exists (n_j) \text{ such that } \frac{\log^+ \log^+ |f^{n_j}(z)|}{n_j} \rightarrow \infty \text{ as } j \rightarrow \infty \right\},$$

where (n_j) is a strictly increasing sequence of positive integers. The set $Z^+(f)$ is formed from the set $Z(f)$ of points whose iterates 'zip' to ∞ (for which, see [28]) by adding those points at which only a subsequence of iterates has this property.

It is clear that Zheng's result (c) is related to Baker's conjecture. To see that (a) and (b) are also related to this conjecture, note that the hypotheses about $m(r)$ in (a) and (b) both hold whenever f has order at most $1/2$, minimal type, by a classical theorem of Wiman [42]; see [14, Theorem 6.4], and [15] for a more precise version of Wiman's theorem. Also, note that the condition on $m(r)$ in (a) ensures that f is a strongly polynomial-like function; see [22], where a generalisation of Zheng's result (a) to all strongly polynomial-like functions is given.

We use the property of long continua meeting $V^+(f)$ to prove the following theorem, which gives more detailed information than Zheng's results (a) and (b) whenever (1.1) holds, and strengthens Zheng's result (c).

Theorem 1.6. *Let f be a transcendental entire function and let U be an unbounded component of $F(f)$.*

- (a) *If (1.1) holds, then $U \cap V^+(f) \neq \emptyset$, and hence U is either a Baker domain (or preimage of a Baker domain) or a wandering domain. If, in addition,*

$$\liminf_{r \rightarrow \infty} \frac{\tilde{m}(r)}{r} > 1,$$

then $U \subset V^+(f)$.

- (b) *If*

$$\lim_{r \rightarrow \infty} \frac{\tilde{m}(r)}{r} = \infty,$$

then U is a wandering domain and $U \subset V^+(f)$.

- (c) *If*

$$\lim_{r \rightarrow \infty} \frac{\log \tilde{m}(r)}{\log r} = \infty,$$

then U is a wandering domain and $U \subset V^+(f) \subset Z^+(f)$.

The hypothesis on \tilde{m} in part (c) of this result holds for many classes of entire functions, including those of order less than $1/2$; see Section 3 and, in particular, Lemma 3.1.

Finally, we also provide new information about the rate of escape that occurs in certain Baker domains.

Theorem 1.7. *Let f be a transcendental entire function and let U be an invariant Baker domain of f .*

If (1.1) holds, then there exists $R > 0$ and, for each $z \in U$, a constant $C(z) > 1$ such that

$$|f^n(z)| \geq \tilde{m}^n(R)/C(z), \quad \text{for } n \in \mathbb{N}, \quad \text{and} \quad \tilde{m}^n(R) \rightarrow \infty \text{ as } n \rightarrow \infty.$$

If, in addition,

$$\liminf_{r \rightarrow \infty} \frac{\tilde{m}(r)}{r} > 1,$$

then $U \subset V(f)$.

The organisation of this paper is as follows. In Section 2, we prove our results about the existence of escaping points of positive continuous functions; in particular, Theorem 2.1 gives several equivalent conditions for the existence of such points. In Section 3, we use Theorem 2.1 to deduce Theorem 1.1 on classes of functions for which condition (1.1) holds. In Sections 4 and 5, we prove Theorems 1.2 and 1.3 concerning the set $V(f)$ and show that $V(f) = A(f)$ for the functions listed in Theorem 1.1 provided they satisfy a mild regularity condition. The proofs of Theorem 1.5 and related results are given in Section 6, and the proofs of Theorems 1.6 and 1.7 are given in Section 7. Finally, in Section 8, we give some examples to illustrate our results.

2. ESCAPING POINTS OF POSITIVE CONTINUOUS FUNCTIONS

Let $\varphi : [0, \infty) \rightarrow [0, \infty)$ be a continuous function and define the function $\tilde{\varphi}$ by

$$\tilde{\varphi}(t) := \max_{0 \leq s \leq t} \varphi(s), \quad \text{for } t \in [0, \infty).$$

The following result gives a number of conditions equivalent to the existence of escaping points of φ . In particular, this theorem justifies our assertion in the introduction that

$$\text{there exists } r > 0 \text{ with } m^n(r) \rightarrow \infty \text{ as } n \rightarrow \infty$$

if and only if

$$\text{there exists } R > 0 \text{ such that } \tilde{m}(r) > r, \text{ for } r \geq R.$$

Theorem 2.1. *Let $\varphi : [0, \infty) \rightarrow [0, \infty)$ be continuous. Then the following statements are equivalent.*

- (a) *There exists $t > 0$ such that $\varphi^n(t) \rightarrow \infty$ as $n \rightarrow \infty$.*
- (b) *There exists $t' > 0$ such that the set $\{\varphi^n(t') : n \in \mathbb{N}_0\}$ is unbounded.*
- (c) *There exists $T > 0$ such that $\tilde{\varphi}(t) > t$, for $t \geq T$.*
- (d) *There exist $t \geq T > 0$ such that*

$$\varphi^n(t) \text{ and } \tilde{\varphi}^n(T) \text{ increase strictly with } n \text{ to } \infty,$$

and

$$\varphi^n(t) \in [\tilde{\varphi}^n(T), \tilde{\varphi}^{n+1}(T)], \quad \text{for } n \in \mathbb{N}_0.$$

- (e) *There exists a sequence (t_n) of positive real numbers such that $t_n \rightarrow \infty$ as $n \rightarrow \infty$ and*

$$\varphi(t_n) \geq t_{n+1}, \quad \text{for } n \in \mathbb{N}_0.$$

Remarks 1. It follows from a recent result of Short and Sixsmith [39, Theorem 1.6] that if $t > 0$ satisfies the condition in Theorem 2.1 part (a), then every open interval containing t includes uncountably many escaping points.

2. The condition in part (d) of Theorem 2.1 shows that there always exist points that escape at the fastest possible rate. We use this fact in Section 4 in the proof of Theorem 1.2.

3. In Section 6, we use Theorem 2.1 to show that if a transcendental entire function f satisfies (1.1), then any curve that tends to ∞ and is invariant under f must meet $I(f)$.

In the introduction we also stated that if (1.1) holds, then $m^n(r)$ can tend to ∞ arbitrarily slowly. This follows from our next result because any transcendental entire function either has infinitely many zeros or has asymptotic value 0, by Iversen's theorem, so

$$\liminf_{r \rightarrow \infty} m(r) = 0.$$

Theorem 2.2. *Let $\varphi : [0, \infty) \rightarrow [0, \infty)$ be continuous and suppose there exists $t > 0$ such that $\varphi^n(t) \rightarrow \infty$ as $n \rightarrow \infty$. Suppose also that*

$$(2.1) \quad \liminf_{t \rightarrow \infty} \varphi(t) < \infty.$$

If $a = (a_n)$ is a positive sequence such that $a_n \rightarrow \infty$ as $n \rightarrow \infty$, then there exist $t_a > 0$ and $N_a \in \mathbb{N}$ such that $\varphi^n(t_a) \rightarrow \infty$ as $n \rightarrow \infty$ and

$$(2.2) \quad \varphi^n(t_a) \leq a_n, \quad \text{for } n \geq N_a.$$

The key techniques used in the proofs of Theorems 2.1 and 2.2 are given in the following lemma.

Lemma 2.3. *Let φ and $\tilde{\varphi}$ be as defined before the statement of Theorem 2.1, and suppose there exists $T > 0$ such that $\tilde{\varphi}(t) > t$, for $t \geq T$. Then*

- (a) *the sequence $\tilde{t}_n := \tilde{\varphi}^n(T)$, $n \in \mathbb{N}_0$, is strictly increasing and tends to ∞ ;*
 (b) *if $E_n = [\tilde{t}_n, \tilde{t}_{n+1}]$, for $n \in \mathbb{N}_0$, then*

$$\varphi(E_n) \supset E_{n+1}, \quad \text{for } n \in \mathbb{N}_0;$$

- (c) *if, in addition,*

$$\liminf_{t \rightarrow \infty} \varphi(t) < \infty,$$

then there is a subsequence $(E_{n(k)})$, $k \in \mathbb{N}_0$, such that

$$\varphi(E_{n(k)}) \supset \bigcup_{n=n(0)}^{n(k)} E_n, \quad \text{for } k \in \mathbb{N}_0.$$

Proof. The hypothesis of the lemma clearly implies part (a). It follows from the definitions of $\tilde{\varphi}$ and \tilde{t}_n that

$$(2.3) \quad \varphi(\tilde{t}_n) \leq \tilde{\varphi}(\tilde{t}_n) = \tilde{t}_{n+1}, \quad \text{for } n \in \mathbb{N}_0,$$

and

$$(2.4) \quad \tilde{t}_{n+2} = \tilde{\varphi}(\tilde{t}_{n+1}) = \varphi(t), \quad \text{for some } t \leq \tilde{t}_{n+1}.$$

Now we cannot have $t < \tilde{t}_n$, for otherwise, using part (a),

$$\varphi(t) \leq \tilde{\varphi}(t) \leq \tilde{\varphi}(\tilde{t}_n) = \tilde{t}_{n+1} < \tilde{t}_{n+2},$$

which contradicts (2.4). Thus $t \in [\tilde{t}_n, \tilde{t}_{n+1}] = E_n$ and, together with (2.3), this shows that

$$\varphi(E_n) \supset E_{n+1}, \quad \text{for } n \in \mathbb{N}_0,$$

which proves part (b).

Part (c) follows by choosing $n(0)$ so that $\tilde{t}_{n(0)} > \liminf_{t \rightarrow \infty} \varphi(t)$. □

We now give the proofs of Theorems 2.1 and 2.2. The proof of Theorem 2.1 makes use of a simple topological result, which is widely used, going back at least to [9], and stated explicitly in [30, Lemma 1]. We quote here a version of this result which we also need later in the paper; compare [40, Lemma 3.1].

Lemma 2.4. *Let $(E_j)_{j \in \mathbb{N}_0}$ be a sequence of compact sets in \mathbb{C} , $(m_j)_{j \in \mathbb{N}_0}$ be a sequence of positive integers and $f : \mathbb{C} \rightarrow \hat{\mathbb{C}}$ be a continuous function such that*

$$f^{m_j}(E_j) \supset E_{j+1}, \quad \text{for } j \in \mathbb{N}_0.$$

Then there exists $\zeta \in E_0$ such that

$$f^{m_0+m_1+\dots+m_n}(\zeta) \in E_{n+1}, \quad \text{for } n \in \mathbb{N}_0.$$

Proof of Theorem 2.1. We show that (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (d) \Rightarrow (e), and then that (e) \Rightarrow (c). Since it is clear that (d) \Rightarrow (a), this will prove the theorem.

It is obvious that (a) \Rightarrow (b) and that (d) \Rightarrow (e).

Suppose (b) holds, and that $t' > 0$ is such that the set $\{\varphi^n(t') : n \in \mathbb{N}_0\}$ is unbounded. Then, for $t \geq t'$, there is an integer $N = N(t) \in \mathbb{N}$ such that $\varphi^{N-1}(t') \leq t$ and $\varphi^N(t') > t$. It follows that

$$\tilde{\varphi}(t) \geq \tilde{\varphi}(\varphi^{N-1}(t')) \geq \varphi^N(t') > t,$$

which proves (c) with $T = t'$.

Now suppose (c) holds, and that $T > 0$ is such that $\tilde{\varphi}(t) > t$ for $t \geq T$. Define

$$E_n = [\tilde{\varphi}^n(T), \tilde{\varphi}^{n+1}(T)], \text{ for } n \in \mathbb{N}_0.$$

Then it follows by Lemma 2.4 and Lemma 2.3 part (b) that there is a point $t \in E_0$ such that $\varphi^n(t) \in E_n$, for $n \in \mathbb{N}_0$, and therefore $\varphi^n(t) \geq \tilde{\varphi}^n(T) \rightarrow \infty$, as $n \rightarrow \infty$. Moreover, the sequence $(\varphi^n(t))$, $n \in \mathbb{N}_0$, is strictly increasing, since otherwise it would eventually be constant. This proves (d).

It remains to prove that (e) \Rightarrow (c). Suppose (e) holds, and let (t_n) be a positive sequence that tends to ∞ as $n \rightarrow \infty$, with $\varphi(t_n) \geq t_{n+1}$ for $n \in \mathbb{N}_0$. Set $T = t_0$. If $t \geq T$, then $t \in [t_n, t_{n+1})$ for some $n \in \mathbb{N}_0$, so

$$\tilde{\varphi}(t) \geq \varphi(t_n) \geq t_{n+1} > t,$$

which proves (c) and completes the proof of the theorem. \square

Proof of Theorem 2.2. By Theorem 2.1, there exists $T > 0$ such that $\tilde{\varphi}(t) > t$ for $t \geq T$. Thus it follows from Lemma 2.3 that the sequence $\tilde{t}_n = \tilde{\varphi}^n(T)$ is strictly increasing and that, if $E_n = [\tilde{t}_n, \tilde{t}_{n+1}]$ for $n \in \mathbb{N}_0$, then

$$(2.5) \quad \varphi(E_n) \supset E_{n+1}, \text{ for } n \in \mathbb{N}_0.$$

Furthermore, since (2.1) holds, it follows from Lemma 2.3 part (c) that there is a subsequence $(E_{n(k)})$, $k \in \mathbb{N}_0$, such that

$$(2.6) \quad \varphi(E_{n(k)}) \supset E_{n(k)}, \text{ for } k \in \mathbb{N}_0.$$

We now construct a new sequence of intervals by selecting terms from the sequence (E_n) in such a way that we can apply Lemma 2.4 and hence deduce the existence of $t_a > 0$ and $N_a \in \mathbb{N}$ with the properties in the statement of the theorem. The basic idea is that, in this new sequence, we repeat each of the intervals $E_{n(k)}$ sufficiently often that, by using property (2.6), we ‘slow down’ the rate at which $\varphi^n(t_a) \rightarrow \infty$ to ensure that (2.2) is satisfied.

Without loss of generality, we can assume that $a = (a_n)$ is increasing. Let the new sequence (F_m) , $m \in \mathbb{N}_0$, consist of all the intervals E_n for $n \geq n(0)$, taken in order of increasing n , but with each interval $E_{n(k)}$, $k \in \mathbb{N}_0$, repeated $m(k)$ times, where $m(k)$ is so large that

$$\bigcup_{n=n(0)}^{n(k+1)} E_n \subset [0, a_{m(k)}].$$

Since the interval $E_{n(k)}$ is repeated $m(k)$ times in the new sequence (F_m) , it follows that if $F_m = E_n$, where $n(k) + 1 \leq n \leq n(k+1)$, then $m \geq m(k)$. Thus, for such n , we have

$$F_m = E_n \subset [0, a_{m(k)}] \subset [0, a_m],$$

and it follows that

$$(2.7) \quad F_m \subset [0, a_m], \text{ for } m \geq m(0).$$

Now by (2.5) and (2.6) we have

$$\varphi(F_m) \supset F_{m+1}, \quad \text{for } m \in \mathbb{N}_0,$$

so applying Lemma 2.4 we obtain $t_a \in F_0$ such that $\varphi^m(t_a) \in F_m$, for $m \in \mathbb{N}_0$. Clearly $\varphi^m(t_a) \rightarrow \infty$ as $m \rightarrow \infty$, and it follows from (2.7) that

$$\varphi^m(t_a) \leq a_m, \quad \text{for } m \geq m(0).$$

This completes the proof. \square

Remark We note that the property (2.6) implies, by Lemma 2.4, that each interval of the form $E_{n(k)}$, $k \in \mathbb{N}_0$, contains a value of t such that $\varphi^n(t) \in E_{n(k)}$, for $n \in \mathbb{N}_0$, and so the sequence $(\varphi^n(t))$ is bounded. The combination of (2.5) and (2.6) can also be used to obtain values of t such that $(\varphi^n(t))$ is unbounded but not escaping; see [24] where a similar approach was used for transcendental entire functions.

3. CLASSES OF FUNCTIONS FOR WHICH CONDITION (1.1) HOLDS

In this section we prove Theorem 1.1, which lists a number of large classes of transcendental entire functions for which condition (1.1) holds. A key step in the proof of parts (a)–(d) of Theorem 1.1 is the following simple lemma, which will also be needed in Section 7.

Lemma 3.1. *Suppose f is a transcendental entire function with the property that there exist $C > 1$ and $R_0 > 0$ such that, for $r \geq R_0$,*

$$(3.1) \quad \text{there exists } s \in (r, r^C) \text{ with } m(s) \geq M(r).$$

Then condition (1.1) is satisfied and, more strongly,

$$(3.2) \quad \frac{\log \tilde{m}(r)}{\log r} \rightarrow \infty \quad \text{as } r \rightarrow \infty;$$

in particular, for r sufficiently large, we have

$$(3.3) \quad \frac{\log^+ \log^+ \tilde{m}^n(r)}{n} \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

Proof. By (3.1), we have

$$(3.4) \quad \tilde{m}(r^C) \geq m(s) \geq M(r), \quad \text{for } r \geq R_0.$$

Then (3.2) follows from (3.4) together with the well known fact that if f is a transcendental entire function, then

$$(3.5) \quad \frac{\log M(r)}{\log r} \rightarrow \infty \quad \text{as } r \rightarrow \infty,$$

and (1.1) follows from (3.2) and Theorem 2.1. Finally, (3.3) follows easily from (3.2). \square

To prove Theorem 1.1, we show that each of the classes (a)–(d) in the theorem meets the condition in Lemma 3.1 and then deal with class (e) separately.

We first define some terms needed here and later in the paper. The *order* ρ , *lower order* λ and *type* τ of an entire function f are defined by

$$\rho = \limsup_{r \rightarrow \infty} \frac{\log \log M(r, f)}{\log r},$$

$$\lambda = \liminf_{r \rightarrow \infty} \frac{\log \log M(r, f)}{\log r}$$

and

$$\tau = \limsup_{r \rightarrow \infty} \frac{\log M(r, f)}{r^\rho}.$$

If $\tau = 0$, then f is said to be of *minimal type*.

We note that the arguments used in the proof of Lemma 3.2 were also used in the proof of [32, Corollary 8.3(a)], but we include a summary of them here for completeness.

Lemma 3.2. *The functions in each of the classes (a)–(d) in Theorem 1.1 satisfy condition (3.1) and hence condition (1.1).*

Proof. We consider each of the classes (a)–(d) in turn.

(a) It was shown by Baker [1, Satz 1] that a transcendental entire function f of order less than $1/2$ satisfies (3.1) for sufficiently large positive values of C ; this also follows from the version of the $\cos \pi \rho$ theorem proved by Barry [3].

(b) A transcendental entire function f is said to have *Fabry gaps* if

$$(3.6) \quad f(z) = \sum_{k=0}^{\infty} a_k z^{n_k},$$

where $n_k/k \rightarrow \infty$ as $k \rightarrow \infty$. It follows from a result of Fuchs [12, Theorem 1] that, if f has finite order and Fabry gaps, then for each $\varepsilon > 0$,

$$\log m(r) > (1 - \varepsilon) \log M(r),$$

for values of r outside a set of zero logarithmic density. It is easy to check that this implies that functions of finite order with Fabry gaps satisfy (3.1) for $C > 1$.

(c) Hayman [13, Theorem 3] showed that the conclusion of Fuchs' result above holds for transcendental entire functions of *any* order provided that a stronger gap condition is satisfied, which we call *Hayman gaps*. The condition is that, in the expansion (3.6), we have

$$n_k > k \log k (\log \log k)^\alpha,$$

for some $\alpha > 2$ and sufficiently large values of $k \in \mathbb{N}$. As before, it follows that such functions satisfy (3.1) for $C > 1$.

(d) Loosely speaking, a function exhibits the *pits effect* if it has very large modulus except in small regions (pits) around its zeros. Littlewood and Offord [18] showed that, if $\sum_{n=0}^{\infty} a_n z^n$ is a transcendental entire function of order $\rho \in (0, \infty)$ and lower order $\lambda > 0$, and if

$$S = \left\{ f : f(z) = \sum_{n=0}^{\infty} \varepsilon_n a_n z^n \right\}$$

where the ε_n take the values ± 1 with equal probability then, in some precise sense, almost all functions in the set S exhibit the pits effect. For such functions, it is shown in [32, Proof of Example 2] that, if $|z| = r$, then

$$\log |f(z)| > \frac{1}{4} \log M(r),$$

outside a set of values of r of finite logarithmic measure. Again, this is sufficient to show that (3.1) holds for large positive values of C . \square

Finally, we consider functions with multiply connected Fatou components, that is, multiply connected components of the Fatou set. The papers [7] and [36] give a very detailed analysis of dynamical behaviour in such components, including the following result about the existence of large annuli whose union is forward invariant [36, Lemma 3.3]. Here we use the notation $\delta(r) := 1/\sqrt{\log r}$ and also

$$A(r, R) := \{z : r < |z| < R\}, \quad 0 < r < R.$$

Lemma 3.3. *Let f be a transcendental entire function with a multiply connected Fatou component. Then there exist sequences (r_n) and (k_n) , with $r_n > 1$ and $k_n > 1$, for $n \in \mathbb{N}_0$, such that the annuli*

$$A'_n = A(r_n^{1+6\pi\delta_n}, r_n^{k_n(1-6\pi\delta_n)}), \quad \text{where } \delta_n = \delta(r_n), \quad n \in \mathbb{N}_0,$$

have the properties that, for $n \in \mathbb{N}_0$,

$$f(A'_n) \subset A'_{n+1},$$

and

$$r_{n+1} = M(r_n) > r_n^{16}.$$

It follows from Lemma 3.3 that if $r > 0$ and $r \in A'_0$, then $m^n(r) \in A'_n$, so

$$(3.7) \quad m^n(r) > r_n^{1+6\pi\delta_n} \geq r_n = M^n(r_0), \quad \text{for } n \in \mathbb{N}_0.$$

Hence $m^n(r) \rightarrow \infty$ as $n \rightarrow \infty$ whenever f has a multiply connected Fatou component. This completes the proof of Theorem 1.1.

4. THE SETS $V(f)$ AND $V^+(f)$: PROOF OF THEOREM 1.2

In this section we prove a number of basic properties of the sets $V(f)$ and $V^+(f)$, starting with Theorem 1.2. Recall that, for a transcendental entire function such that (1.1) holds, we define

$$V(f) = \bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(V_R(f)), \quad \text{where } V_R(f) = \{z : |f^n(z)| \geq \tilde{m}^n(R), \text{ for } n \in \mathbb{N}\},$$

and $R > 0$ is such that $\tilde{m}(r) > r$ for $r \geq R$.

We first show that this definition is unambiguous. To do this, we return to the question of the rate at which $m^n(r)$ tends to infinity for a transcendental entire function satisfying (1.1). We showed in Theorem 2.2 that $m^n(r)$ can tend to infinity arbitrarily slowly. By contrast, the *fastest* rate at which $m^n(r)$ can tend to infinity must be limited by the growth of $\tilde{m}^n(r)$, and we now prove Theorem 1.2 which shows that this fastest rate is always attained.

Proof of Theorem 1.2. If f is a transcendental entire function such that (1.1) holds, then it follows from Theorem 2.1 that there exist $R > 0$ such that

$$(4.1) \quad \tilde{m}^n(R) \rightarrow \infty \text{ as } n \rightarrow \infty,$$

and $r \geq R$ such that

$$(4.2) \quad m^n(r) \geq \tilde{m}^n(R), \text{ for } n \in \mathbb{N}.$$

We now show that the set

$$V_1(R) = \bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(\{z : |f^n(z)| \geq \tilde{m}^n(R), \text{ for } n \in \mathbb{N}\})$$

is independent of the choice of R satisfying (4.1) and, for $r \geq R$ satisfying (4.2), is equal to the set

$$V_2(r) = \bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(\{z : |f^n(z)| \geq m^n(r), \text{ for } n \in \mathbb{N}\}).$$

Suppose $R' > R$, where R satisfies (4.1). Then since $\tilde{m}^n(R') \geq \tilde{m}^n(R)$ for $n \in \mathbb{N}$, it is clear that $V_1(R') \subset V_1(R)$. On the other hand, by (4.1) there exists $k \in \mathbb{N}$ such that $\tilde{m}^k(R) \geq R'$, so for $z \in V_1(R)$ there exists $\ell \in \mathbb{N}_0$ such that

$$|f^{n+\ell}(z)| \geq \tilde{m}^n(R) \geq \tilde{m}^{n-k}(R'), \text{ for } n \geq k.$$

Hence $|f^{n+\ell+k}(z)| \geq \tilde{m}^n(R')$ for $n \in \mathbb{N}_0$, and therefore $V_1(R) \subset V_1(R')$. It follows that $V_1(R) = V_1(R')$ and thus that $V_1(R)$ is independent of the choice of R .

The proof that $V_1(R) = V_2(r)$ is similar. Suppose r and R satisfy (4.1) and (4.2). Then (4.2) clearly implies that $V_2(r) \subset V_1(R)$ and, since by (4.1) there exists $j \in \mathbb{N}$ such that $\tilde{m}^j(R) \geq r$, we deduce as above that $V_1(R) \subset V_2(r)$. \square

It follows from Theorem 1.2 that, if (1.1) holds, then the set $V(f)$ is well defined and independent of R , provided $R > 0$ is such that (4.1) holds, or equivalently that $\tilde{m}(r) > r$ for $r \geq R$.

Now recall that

$$V^+(f) = \bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(V_R^+(f)),$$

where

$$V_R^+(f) = \{z : \exists (n_j) \text{ such that } |f^{n_j}(z)| \geq \tilde{m}^{n_j}(R), \text{ for } j \in \mathbb{N}\}.$$

In this definition, $R > 0$ satisfies $\tilde{m}(r) > r$ for $r \geq R$ and (n_j) is a strictly increasing sequence of positive integers, which in general depends on z . Using an argument similar to the proof of Theorem 1.2, it is easy to see that if f is a transcendental entire function such that condition (1.1) holds, and $r \geq R > 0$ satisfy (4.1) and (4.2), then

$$(4.3) \quad V^+(f) = \bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(\{z : \exists (n_j) \text{ such that } |f^{n_j}(z)| \geq m^{n_j}(r), \text{ for } j \in \mathbb{N}\}).$$

Thus, the set $V^+(f)$ is also well defined and independent of R , provided $R > 0$ is such that $\tilde{m}(r) > r$ for $r \geq R$.

We now show that $V(f)$ and $V^+(f)$ have some of the basic properties of $I(f)$ proved in [9]. There, Eremenko showed that

$$(4.4) \quad I(f) \neq \emptyset, \quad I(f) \cap J(f) \neq \emptyset, \quad J(f) = \partial I(f),$$

and that $\overline{I(f)}$ has no bounded components. (Recall that the fast escaping set $A(f)$ also has the properties listed in (4.4), and in addition $A(f)$ has no bounded components; see [6], [29] and [32].) For $V(f)$ and $V^+(f)$, we have the following.

Theorem 4.1. (a) *Let f be a transcendental entire function such that (1.1) holds. Then*

$$(4.5) \quad V(f) \neq \emptyset, \quad V(f) \cap J(f) \neq \emptyset, \quad J(f) = \overline{V(f) \cap J(f)}, \quad J(f) \subset \partial V(f),$$

and $\overline{V(f)}$ has no bounded components.

(b) *If, in addition,*

$$(4.6) \quad \liminf_{r \rightarrow \infty} \frac{\tilde{m}(r)}{r} > 1,$$

then every Fatou component of f that meets $V(f)$ must lie entirely in $V(f)$, and $J(f) = \partial V(f)$.

(c) *The properties in (a) and (b) hold if $V(f)$ is replaced by $V^+(f)$.*

We make use of the following result, which is part of [28, Theorem 3].

Lemma 4.2. *Let f be a transcendental entire function. If U is a simply connected component of $F(f)$ and K is a compact subset of U , then there exists $C = C(K) \in (1, \infty)$ such that*

$$\frac{|f^n(z_2)|}{|f^n(z_1)| + 1} \leq C, \quad \text{for } z_1, z_2 \in K \text{ and } n \in \mathbb{N}.$$

Proof of Theorem 4.1. (a) The first three properties of $V(f)$ in (4.5) are immediate since these properties hold for $A(f)$ (see [32]) and $A(f) \subset V(f)$.

Because $V(f)$ is infinite and completely invariant under f , we have $J(f) \subset \overline{V(f)}$, and this implies that $J(f) \subset \partial V(f)$ since any open subset of $V(f)$ is contained in $F(f)$.

Finally, if $\overline{V(f)}$ has a bounded component, E say, then there is an open topological annulus A which surrounds E and lies in the complement of $\overline{V(f)}$. Since $\overline{V(f)}$ is completely invariant under f we deduce that $A \subset F(f)$, and since $J(f) \subset \partial V(f)$ it follows that A is contained in a multiply connected Fatou component. But any multiply connected Fatou component of f is contained in $A(f)$ (see [29]) and hence in $V(f)$, so we obtain a contradiction.

(b) The first statement of part (b) is immediate if the Fatou component is multiply connected, since in that case it lies in $A(f)$; see [29]. Otherwise this statement follows from Lemma 4.2 and the hypothesis (4.6). Indeed, if U is a simply connected Fatou component of f and $z \in U \cap V(f)$, then there exists $\ell \in \mathbb{N}$ such that

$$|f^{n+\ell}(z)| \geq \tilde{m}^n(R), \quad \text{for } n \in \mathbb{N},$$

where $R > 0$ is such that $\tilde{m}(r) > r$ for $r \geq R$. We deduce from Lemma 4.2 that for any $z' \in U$ there exists $C(z') > 1$ such that, for each $n \in \mathbb{N}$,

$$|f^n(z')| \geq |f^n(z)|/C(z').$$

The hypothesis (4.6) implies that there exists $k = k(C(z')) \in \mathbb{N}$ such that

$$\tilde{m}^{n+k}(R) = \tilde{m}^k(\tilde{m}^n(R)) > C(z')\tilde{m}^n(R), \quad \text{for } n \in \mathbb{N}.$$

Thus

$$|f^{n+\ell}(z')| \geq |f^{n+\ell}(z)|/C(z') \geq \tilde{m}^n(R)/C(z') \geq \tilde{m}^{n-k}(R), \quad \text{for } n > k,$$

so $z' \in U \cap V(f)$, as required.

This property of Fatou components implies immediately that $J(f) \supset \partial V(f)$. Since, by part (a), we also have $J(f) \subset \partial V(f)$, it follows that $J(f) = \partial V(f)$ whenever (4.6) holds.

(c) Similar arguments show that the properties in parts (a) and (b) also hold for $V^+(f)$; we omit the details. \square

Remarks. 1. Note that, if f is a transcendental entire function such that (1.1) holds, then $V^+(f)$ is connected by Theorem 1.5, from which it is immediate that $V^+(f)$ and $\overline{V^+(f)}$ have no bounded components.

2. It is natural to ask if the statement that $J(f) \subset \partial V(f)$ in Theorem 4.1 part (a) can be strengthened to $J(f) = \partial V(f)$ for all transcendental entire functions that satisfy (1.1), and similarly for $V^+(f)$.

Finally in this section, we record various relationships involving the connectedness properties of $V(f)$, $V^+(f)$, $I(f)$ and $I^+(f)$. Recall that a connected set E is a spider's web if there exists a sequence (G_n) of bounded, simply connected domains such that

$$G_n \subset G_{n+1}, \quad \partial G_n \subset E, \quad \text{for } n \in \mathbb{N}, \quad \text{and} \quad \bigcup_{n \in \mathbb{N}} G_n = \mathbb{C},$$

and E is a weak spider's web if its complement contains no unbounded closed connected sets.

Theorem 4.3. *Let f be a transcendental entire function such that (1.1) holds. We have the following implications:*

- (a) *if $V(f)$ is connected, then $I(f)$ is connected;*
- (b) *if $V(f)$ is a (weak) spider's web, then $I(f)$ is a (weak) spider's web;*
- (c) *if $V^+(f)$ is a spider's web, then $I^+(f)$ is a spider's web.*

Note that if (1.1) holds, then both $V^+(f)$ and $I^+(f)$ are weak spiders' webs; see Theorem 1.5.

To prove part (a) of Theorem 4.3 we use the following result [37, Theorem 1.2].

Lemma 4.4. *Let f be a transcendental entire function and let E be a set such that $E \subset I(f)$ and $J(f) \subset \overline{E}$. Either $I(f)$ is connected or it has infinitely many components that meet E ; in particular, if E is connected, then $I(f)$ is connected.*

Proof of Theorem 4.3. Part (a) follows from Lemma 4.4 by taking $E = V(f)$, since $J(f) \subset \overline{V(f)}$ by Theorem 4.1.

To prove parts (b) and (c) we note that, by the definitions, if a connected set contains a (weak) spider's web, then it is a (weak) spider's web. The fact that $I(f)$ is connected follows from part (a), and the proof that $I^+(f)$ is connected was given in [23, Theorem 1.2]. \square

5. FUNCTIONS FOR WHICH $V(f) = A(f)$: PROOF OF THEOREM 1.3

In this section we consider functions satisfying (1.1) for which $V(f) = A(f)$. We show that there are many classes of functions with this property. The main focus of the section is the proof of Theorem 1.3.

Part (b) of Theorem 1.3 says that, for a transcendental entire function f satisfying (1.1), if $V(f) = A(f)$ then $A_R(f)$, $V(f)$ and $I(f)$ are spiders' webs, and $F(f)$ has no unbounded components. This proves that Eremenko's conjecture holds for such functions, and that Baker's conjecture holds for such functions of order less than $1/2$, minimal type.

Part (a) of the theorem gives a useful equivalent condition that we use in the proof of part (b), namely that $V(f) = A(f)$ if and only if

$$(5.1) \quad \begin{aligned} & \text{there exist } r \geq R > 0 \text{ such that } m^n(r) \geq M^n(R), \text{ for } n \in \mathbb{N}, \\ & \text{and } M^n(R) \rightarrow \infty \text{ as } n \rightarrow \infty. \end{aligned}$$

We need three further results for the proof. The first is part of [36, Theorem 1.4].

Lemma 5.1. *Let f be a transcendental entire function. Then there exists $R_0 > 0$ with the property that, whenever (a_n) is a positive sequence such that*

$$a_n \geq R_0 \quad \text{and} \quad a_{n+1} \leq M(a_n), \quad \text{for } n \in \mathbb{N}_0,$$

there exists $z \in \mathbb{C}$ and a sequence (n_j) with $n_j \rightarrow \infty$ as $j \rightarrow \infty$ such that

$$|f^n(z)| \geq a_n, \quad \text{for } n \in \mathbb{N}_0,$$

and

$$|f^{n_j}(z)| \leq M^2(a_{n_j}), \quad \text{for } j \in \mathbb{N}.$$

Next, we need the following result from plane topology, which will also be used several times later in the paper.

Lemma 5.2. [20, page 84] *If E_0 is a continuum in $\hat{\mathbb{C}}$, E_1 is a closed subset of E_0 and C is a component of $E_0 \setminus E_1$, then \bar{C} meets E_1 .*

Finally, we need the following sufficient condition for the set $A_R(f)$ to be a spider's web. Recall that

$$A_R(f) = \{z : |f^n(z)| \geq M^n(R), \text{ for } n \in \mathbb{N}\},$$

where $R > 0$ is such that $M^n(R) \rightarrow \infty$ as $n \rightarrow \infty$.

Lemma 5.3. [32, Corollary 8.2] *Let f be a transcendental entire function and let $R > 0$ be such that $M^n(R) \rightarrow \infty$ as $n \rightarrow \infty$. Then $A_R(f)$ is a spider's web if there exists a sequence $(s_n)_{n \in \mathbb{N}_0}$ such that*

$$s_n \geq M^n(R) \quad \text{and} \quad m(s_n) \geq s_{n+1}, \quad \text{for } n \in \mathbb{N}_0.$$

We are now in a position to prove Theorem 1.3.

Proof of Theorem 1.3. We first prove part (a). Suppose that f is a transcendental entire function satisfying condition (1.1), and that (5.1) also holds. If $z \in V(f)$, then it follows from Theorem 1.2 and (5.1) that, for some $\ell \in \mathbb{N}_0$,

$$|f^{n+\ell}(z)| \geq \tilde{m}^n(r) \geq m^n(r) \geq M^n(R), \quad \text{for } n \in \mathbb{N},$$

and so $z \in A(f)$. Thus $V(f) \subset A(f)$. As it is always the case that $A(f) \subset V(f)$, we have shown that (5.1) implies $V(f) = A(f)$.

To establish the converse, we prove the contrapositive. First note that, since (1.1) holds, it follows from Theorem 1.2 that there exist $r \geq R > 0$ such that

$$(5.2) \quad m^n(r) \geq \tilde{m}^n(R) \text{ for } n \in \mathbb{N}, \quad \text{and} \quad \tilde{m}^n(R) \rightarrow \infty \text{ as } n \rightarrow \infty.$$

It then follows from Lemma 5.1 that there exists $z \in \mathbb{C}$ and a sequence (n_j) with $n_j \rightarrow \infty$ as $j \rightarrow \infty$ such that

$$(5.3) \quad |f^n(z)| \geq m^n(r), \quad \text{for } n \in \mathbb{N}_0,$$

and

$$(5.4) \quad |f^{n_j}(z)| \leq M^2(m^{n_j}(r)), \quad \text{for } j \in \mathbb{N}.$$

Suppose now that (5.1) does *not* hold. Then for every $\ell \in \mathbb{N}_0$, there exists $N_\ell \in \mathbb{N}$ such that

$$m^{N_\ell+\ell}(r) < M^{N_\ell}(r),$$

and therefore

$$(5.5) \quad m^{n+\ell}(r) < M^n(r), \quad \text{for } n \geq N_\ell.$$

Now it follows from (5.2) and (5.3) that $z \in V(f)$. However, by (5.5), we deduce that, for each $\ell \in \mathbb{N}$, there exists $j = j(\ell)$ such that

$$m^{n_j}(r) < M^{n_j-\ell}(r).$$

Therefore, by (5.4),

$$\begin{aligned} |f^{n_j}(z)| &\leq M^2(m^{n_j}(r)) \\ &\leq M^{n_j-\ell+2}(r), \end{aligned}$$

so $z \notin A(f)$. Thus $V(f) \neq A(f)$, and this completes the proof of part (a).

To prove part (b), observe that condition (5.1) implies that the set $A_R(f)$ is a spider's web, by Lemma 5.3 with $s_n = m^n(r)$. It now follows from [32, Theorems 1.4 and 1.5] that $A(f)$ and $I(f)$ are also spiders' webs, and that f has no unbounded Fatou components. Since (5.1) is equivalent to the condition $V(f) = A(f)$ by part (a), this completes the proof. \square

We conclude this section by discussing when the functions included in Theorem 1.1 have the property that $V(f) = A(f)$. First it is clear that functions with a multiply connected Fatou component have this property, by Theorem 1.3 part (a) and (3.7). The following result shows that the other functions covered by Theorem 1.1 have the property that $V(f) = A(f)$ provided they also satisfy the weak regularity condition given in part (b) below. (Recall that, by Lemma 3.2, the functions in parts (a)–(d) of Theorem 1.1 all satisfy the condition (3.1), which is the same as the condition in part (a) below.)

Theorem 5.4. *Let f be a transcendental entire function and let $R > 0$ be such that $M(r) > r$ for $r \geq R$. Then $V(f) = A(f)$ if, for some $C > 1$,*

(a) *there exists $R_0 > 0$ such that, for $r \geq R_0$,*

there exists $s \in (r, r^C)$ with $m(s) \geq M(r)$, and

(b) *f has regular growth in the sense that there exists a sequence $(r_n)_{n \in \mathbb{N}_0}$ with*

$$(5.6) \quad r_n \geq M^n(R) \text{ and } M(r_n) \geq r_{n+1}^C, \text{ for } n \in \mathbb{N}_0.$$

Proof. Let $R_0 > 0$ be as in part (a). Then, by part (b), there exists a sequence $(r_n)_{n \in \mathbb{N}_0}$ satisfying (5.6) with $r_n > R_0$, for $n \in \mathbb{N}_0$. So, by part (a), for each $n \in \mathbb{N}_0$, there exists $s_n \in (r_n, r_n^C)$ with

$$m(s_n) \geq M(r_n) \geq r_{n+1}^C > s_{n+1}.$$

Therefore

$$\tilde{m}^n(s_0) \geq s_n \geq r_n \geq M^n(R), \text{ for } n \in \mathbb{N}_0,$$

so $\tilde{m}^n(s_0) \rightarrow \infty$ as $n \rightarrow \infty$. Thus, by Theorem 1.2, there exists $r' \geq s_0$ such that $m^n(r') \geq M^n(R)$, for $n \in \mathbb{N}$. It now follows from Theorem 1.3 part (a) that $V(f) = A(f)$. \square

Remark Theorem 5.4 shows that certain hypotheses imply that $V(f) = A(f)$, whereas we showed in [32, Corollary 8.3] that the same hypotheses imply that $A_R(f)$ is a spider's web. Since the property that $V(f) = A(f)$ implies that $A_R(f)$ is a spider's web, by Theorem 1.3 part (b), it appears that Theorem 5.4 is a stronger result than [32, Corollary 8.3]. However, we do not have an example of a function for which $A_R(f)$ is a spider's web but $V(f) \neq A(f)$.

6. THE ITERATED MINIMUM MODULUS AND LONG CONTINUA

In [23] we showed that, for a transcendental entire function f , the set of points $I^+(f)$ at which the iterates of f are unbounded is connected whenever (1.1) holds. The proof of [23, Theorem 1.2] also shows that the complement of $I^+(f)$ contains no unbounded closed connected sets, and thus that $I^+(f)$ is a weak spider's web. This is one part of Theorem 1.5.

In this section we prove the other part of Theorem 1.5, which states that, if condition (1.1) holds, then $V^+(f)$ is also a weak spider's web. In fact, we show that if (1.1) holds, then very many subsets of $I^+(f)$ are weak spider's webs.

For a transcendental entire function f such that $m^n(r) \rightarrow \infty$ as $n \rightarrow \infty$, for some $r > 0$, we put

$$I^+(f, (m^n(r))) := \{z : \text{there exists } (n_j) \text{ such that } |f^{n_j}(z)| \geq m^{n_j}(r), \text{ for } j \in \mathbb{N}\},$$

where (n_j) is a strictly increasing sequence of positive integers, which in general depends on z . We prove the following result.

Theorem 6.1. *Let f be a transcendental entire function satisfying (1.1) and let $r > 0$ be such that $m^n(r) \rightarrow \infty$ as $n \rightarrow \infty$. Then each of the sets*

$$I^+(f, (m^n(r))) \quad \text{and} \quad \bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(I^+(f, (m^n(r))))$$

is a weak spider's web, and hence $V^+(f)$ is a weak spider's webs.

The fact that $V^+(f)$ is a weak spider's web follows, by property (4.3), from the fact that $\bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(I^+(f, (m^n(r))))$ is a weak spider's web.

As noted above, the result that $I^+(f)$ is a weak spider's web whenever (1.1) holds was proved in [23]. The proofs given here for the sets $I^+(f, (m^n(r)))$ and $\bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(I^+(f, (m^n(r))))$ use a similar approach, but they are significantly more complicated and they yield more information about the structure of the set $I^+(f)$. Indeed, recall from Theorem 2.2 that if (1.1) holds, then there exist values of $r > 0$ such that $m^n(r) \rightarrow \infty$ more slowly than any given rate. For such r

the set $\bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(I^+(f, (m^n(r))))$ is correspondingly larger than $V^+(f)$, which is therefore in this sense the smallest subset of $I^+(f)$ having this form.

We deduce Theorem 6.1 from a new fundamental result which states that if $m^n(r) \rightarrow \infty$ as $n \rightarrow \infty$, where $r > 0$, then in a precise sense certain long continua must meet $I^+(f, (m^n(r)))$. Here, and in what follows, we denote the complement of $I^+(f, (m^n(r)))$ by

$$K(f, (m^n(r))) = \{z : |f^n(z)| < m^n(r) \text{ for sufficiently large } n\}.$$

Theorem 6.2. *Let f be a transcendental entire function satisfying (1.1), let $r > 0$ be such that $m^n(r) \rightarrow \infty$ as $n \rightarrow \infty$, and put*

$$(6.1) \quad D_n = \{z \in \mathbb{C} : |z| < m^n(r)\}, \text{ for } n \in \mathbb{N}_0.$$

Suppose that $\alpha \in K(f, (m^n(r)))$ and let $N_0 = N_0(\alpha) \in \mathbb{N}$ be such that

$$f^n(\alpha) \in D_n, \text{ for } n \geq N_0.$$

If $K \subset K(f, (m^n(r)))$ is a continuum, and $\alpha \in K$, then

$$f^n(K) \subset D_n, \text{ for } n \geq N_0.$$

Moreover, there exists $N_1 = N_1(\alpha)$ such that $K \subset D_{N_1}$.

The following lemma contains the induction step used in the proof of Theorem 6.2. The idea of the proof of the lemma is similar to that of [23, Lemma 3.2], though the details are different.

Lemma 6.3. *Let f be a transcendental entire function such that (1.1) holds and let D_n , $n \in \mathbb{N}_0$, be defined as in (6.1). Suppose that, for some $j \in \mathbb{N}_0$, there exists $n_j \in \mathbb{N}_0$ and a continuum Γ_{n_j} with the following properties:*

- (i) $\Gamma_{n_j} \subset f^{n_j}(K(f, (m^n(r)))) \cap (\mathbb{C} \setminus D_{n_j})$;
- (ii) *there is a point $z_{n_j} \in \Gamma_{n_j} \cap \partial D_{n_j}$;*
- (iii) *there is a point $z'_{n_j} \in \Gamma_{n_j}$ such that $f^n(z'_{n_j}) \in D_{n_j+n}$, for $n \in \mathbb{N}$.*

Then there exists $n_{j+1} > n_j$ and a continuum $\Gamma_{n_{j+1}} \subset f^{n_{j+1}-n_j}(\Gamma_{n_j})$ such that properties (i), (ii) and (iii) hold with n_j replaced by n_{j+1} throughout.

Proof. Since $z_{n_j} \in \Gamma_{n_j}$, it follows from property (i) that there exists a minimal integer $N = N(z_{n_j}) \in \mathbb{N}_0$ such that

$$(6.2) \quad f^n(z_{n_j}) \in D_{n_j+n}, \text{ for } n > N.$$

On the other hand, the properties of the minimum modulus function imply that

$$(6.3) \quad f(\partial D_n) \subset \mathbb{C} \setminus D_{n+1}, \text{ for } n \in \mathbb{N}_0,$$

so by property (ii) we have

$$f(z_{n_j}) \in \mathbb{C} \setminus D_{n_{j+1}}.$$

Hence $N \geq 1$. Now define $n_{j+1} = n_j + N$. Then, by (6.2) and the minimality of N ,

$$(6.4) \quad f^n(z_{n_j}) \in D_{n_j+n}, \text{ for } n > n_{j+1} - n_j,$$

and

$$f^{n_{j+1}-n_j}(z_{n_j}) \in \mathbb{C} \setminus D_{n_{j+1}}.$$

Moreover, $f^{n_{j+1}-n_j}(z_{n_j}) \notin \partial D_{n_{j+1}}$, by (6.3) and (6.4), so

$$(6.5) \quad f^{n_{j+1}-n_j}(z_{n_j}) \in \mathbb{C} \setminus \overline{D}_{n_{j+1}}.$$

Also, by property (iii),

$$(6.6) \quad f^{n_{j+1}-n_j}(z'_{n_j}) \in D_{n_{j+1}}.$$

It follows from (6.5) and (6.6) that the continuum $f^{n_{j+1}-n_j}(\Gamma_{n_j})$ includes points from both $D_{n_{j+1}}$ and $\mathbb{C} \setminus \overline{D}_{n_{j+1}}$ (see Figure 2).




FIGURE 2. Proof of Lemma 6.3

Now let $\Gamma_{n_{j+1}}$ be the component of the closed set

$$f^{n_{j+1}-n_j}(\Gamma_{n_j}) \cap (\mathbb{C} \setminus D_{n_{j+1}})$$

that contains the point

$$(6.7) \quad z'_{n_{j+1}} := f^{n_{j+1}-n_j}(z_{n_j}).$$

Then

$$\Gamma_{n_{j+1}} \subset f^{n_{j+1}}(K(f, (m^n(r)))) \cap (\mathbb{C} \setminus D_{n_{j+1}}),$$

and we deduce that $\Gamma_{n_{j+1}}$ meets $\partial D_{n_{j+1}}$ by applying Lemma 5.2 with

$$E_0 = f^{n_{j+1}-n_j}(\Gamma_{n_j}) \cup \overline{D}_{n_{j+1}} \quad \text{and} \quad E_1 = \overline{D}_{n_{j+1}}.$$

Thus there exists $z_{n_{j+1}} \in \Gamma_{n_{j+1}} \cap \partial D_{n_{j+1}}$. Therefore, properties (i) and (ii) hold with n_j replaced by n_{j+1} , and property (iii) also holds, since

$$f^n(z'_{n_{j+1}}) = f^{n+n_{j+1}-n_j}(z_{n_j}) \in D_{n_{j+1}+n}, \quad \text{for } n \in \mathbb{N},$$

by (6.4) and (6.7). □

Next, we use Lemma 6.3 to prove Theorem 6.2.

Proof of Theorem 6.2. Let $\alpha \in K(f, (m^n(r)))$ and let $N_0 \in \mathbb{N}$ be such that $f^n(\alpha) \in D_n$ for $n \geq N_0$. Also, let $K \subset K(f, (m^n(r)))$ be a continuum such that $\alpha \in K$.

We assume for a contradiction that the conclusion of the theorem does not hold; that is, there exists $N \geq N_0$ such that $f^N(K) \cap (\mathbb{C} \setminus D_N) \neq \emptyset$. We show that this assumption implies that we can select a certain continuum Γ to act as the starting

point for the construction of a sequence (Γ_{n_j}) of continua with the properties stated in Lemma 6.3. This then enables us to obtain the required contradiction.

Step 1: Selection of the continuum Γ . Let $\alpha' := f^N(\alpha)$ and $\Gamma := f^N(K)$. Then Γ is a continuum containing α' , and $\alpha' \in D_N$, by hypothesis. Also, $\Gamma \cap (\mathbb{C} \setminus D_N) \neq \emptyset$, by assumption. Hence, we have

$$\Gamma \subset f^N(K(f, (m^n(r))))), \quad \Gamma \cap \partial D_N \neq \emptyset, \quad \alpha' \in \Gamma \quad \text{and} \quad f^n(\alpha') \in D_{N+n}, \quad \text{for } n \in \mathbb{N}_0.$$

Step 2: Construction of a sequence of continua Γ_{n_j} . We now relabel D_n as D_{n-N} for $n \geq N$, and put $r' = m^N(r)$, giving

$$\Gamma \subset K(f, (m^n(r'))), \quad \Gamma \cap \partial D_0 \neq \emptyset, \quad \alpha' \in \Gamma \quad \text{and} \quad f^n(\alpha') \in D_n, \quad \text{for } n \in \mathbb{N}_0.$$

Let $z_0 \in \Gamma \cap \partial D_0$. Then, since $\Gamma \subset K(f, (m^n(r')))$, there exists $n_0 \in \mathbb{N}$ such that

$$(6.8) \quad f^{n_0}(z_0) \in \mathbb{C} \setminus D_{n_0} \quad \text{and} \quad f^n(z_0) \in D_n, \quad \text{for } n > n_0.$$

Note that $f^{n_0}(\Gamma)$ meets $\mathbb{C} \setminus \overline{D_{n_0}}$, since if $z'_{n_0} := f^{n_0}(z_0)$ lay in ∂D_{n_0} we would have $f^{n_0+1}(z_0) \in \mathbb{C} \setminus D_{n_0+1}$, contradicting (6.8). Since $f^{n_0}(\alpha') \in D_{n_0}$, it follows that $f^{n_0}(\Gamma)$ meets ∂D_{n_0} .

Now let Γ_{n_0} be the component of $f^{n_0}(\Gamma) \setminus D_{n_0}$ that contains z'_{n_0} . Then

$$(6.9) \quad \Gamma_{n_0} \subset f^{n_0}(\Gamma),$$

and Γ_{n_0} meets ∂D_{n_0} , by Lemma 5.2, applied with

$$E_0 = f^{n_0}(\Gamma) \cup \overline{D_{n_0}} \quad \text{and} \quad E_1 = \overline{D_{n_0}}.$$

Hence Γ_{n_0} satisfies

- (i) $\Gamma_{n_0} \subset f^{n_0}(K(f, (m^n(r')))) \cap (\mathbb{C} \setminus D_{n_0})$;
- (ii) there is a point $z_{n_0} \in \Gamma_{n_0} \cap \partial D_{n_0}$;
- (iii) there is a point $z'_{n_0} \in \Gamma_{n_0}$ such that $f^n(z'_{n_0}) \in D_{n_0+n}$, for $n \in \mathbb{N}$.

Thus by Lemma 6.3 with r replaced by r' , there exist a strictly increasing sequence $(n_j)_{j \in \mathbb{N}_0}$ and a sequence of continua $(\Gamma_{n_j})_{j \in \mathbb{N}_0}$ such that, for each $j \in \mathbb{N}_0$,

- (i) $\Gamma_{n_j} \subset f^{n_j}(K(f, (m^n(r')))) \cap (\mathbb{C} \setminus D_{n_j})$;
- (ii) there is a point $z_{n_j} \in \Gamma_{n_j} \cap \partial D_{n_j}$;
- (iii) there is a point $z'_{n_j} \in \Gamma_{n_j}$ such that $f^n(z'_{n_j}) \in D_{n_j+n}$, for $n \in \mathbb{N}$;
- (iv) $f^{n_{j+1}-n_j}(\Gamma_{n_j}) \supset \Gamma_{n_{j+1}}$.

Step 3: Construction of a point in $\Gamma \cap I^+(f, (m^n(r')))$. We now apply Lemma 2.4 with

$$E_j = \Gamma_{n_j} \quad \text{and} \quad m_j = n_{j+1} - n_j, \quad \text{for } j \in \mathbb{N}_0.$$

By property (iv) above,

$$f^{m_j}(E_j) \supset E_{j+1}, \quad \text{for } j \in \mathbb{N}_0,$$

and we deduce from Lemma 2.4 that there exists $\zeta \in E_0 = \Gamma_{n_0}$ such that

$$f^{m_0+\dots+m_k}(\zeta) \in E_{k+1}, \quad \text{for } k \in \mathbb{N}_0;$$

that is,

$$f^{n_{k+1}-n_0}(\zeta) \in \Gamma_{n_{k+1}}, \quad \text{for } k \in \mathbb{N}_0.$$

Thus, by property (i) of the sequence of continua (Γ_{n_j}) ,

$$f^{n_{k+1}-n_0}(\zeta) \in \mathbb{C} \setminus D_{n_{k+1}}, \quad \text{for } k \in \mathbb{N}_0.$$

It follows from (6.9) that there exists $\zeta' \in \Gamma$ such that $f^{n_0}(\zeta') = \zeta$, and hence

$$f^{n_{k+1}}(\zeta') \in \mathbb{C} \setminus D_{n_{k+1}}, \quad \text{for } k \in \mathbb{N}_0.$$

Therefore $\zeta' \in I^+(f, (m^n(r)))$, which contradicts the fact that $\zeta' \in \Gamma \subset K(f, (m^n(r)))$.

This completes the proof that $f^n(K) \subset D_n$, for $n \geq N_0$.

Finally, if we choose $N_1 = N_1(\alpha)$ such that

$$\{f^k(\alpha) : 0 \leq k < N_0\} \subset D_{N_1} \quad \text{and} \quad D_{N_0} \subset D_{N_1} \subset D_{N_1+1},$$

then $K \subset D_{N_1}$, for otherwise we can deduce, by repeatedly applying the minimum modulus property (1.1) to the continua $f^k(K)$, $k = 0, \dots, N_0 - 1$, that $f^{N_0}(K)$ meets ∂D_{N_1} , and hence meets ∂D_{N_0} . \square

We are now in a position to complete the proof of Theorem 6.1. To do this, we make use of the following characterisation of a disconnected subset of the plane.

Lemma 6.4. [26, Lemma 3.1] *A subset S of \mathbb{C} is disconnected if and only if there exists a closed, connected set $\Omega \subset \mathbb{C}$ such that $S \cap \Omega = \emptyset$ and at least two different components of Ω^c intersect S .*

Proof of Theorem 6.1. It follows from Theorem 6.2 and another application of Lemma 5.2 that $K(f, (m^n(r)))$ contains no unbounded closed connected set. Therefore, to complete the proof that $I^+(f, (m^n(r)))$ is a weak spider's web, we must show that this set is connected.

Suppose that $I^+(f, (m^n(r)))$ is disconnected. Then, by Lemma 6.4, there is a closed connected set $E \subset K(f, (m^n(r)))$ such that at least two different components of E^c intersect $I^+(f, (m^n(r)))$. Now E is bounded by Theorem 6.2, so at least one such component of E^c , say G , is bounded. Clearly, G is simply connected, so ∂G is a continuum.

We now show that, as in the proof of Theorem 6.2, we can select a certain continuum Γ to act as the starting point for the construction of a sequence (Γ_{n_j}) of continua with the properties stated in Lemma 6.3, which leads to a contradiction. The argument is identical to the proof of Theorem 6.2 except for the selection of this initial continuum Γ , which we now describe.

By the choice of G we have

$$\partial G \subset K(f, (m^n(r))) \quad \text{and} \quad G \cap I^+(f, (m^n(r))) \neq \emptyset,$$

so there exist $\alpha \in \partial G$, $\beta \in G$ and $N = N(\alpha, \beta) \in \mathbb{N}$ such that

$$f^n(\alpha) \in D_n, \quad \text{for } n \geq N,$$

and

$$f^N(\beta) \in \mathbb{C} \setminus D_N.$$

Since $f^N(\alpha) \in f^N(\partial G)$, whereas $f^N(\beta)$ lies in a bounded complementary component of $f^N(\partial G)$, it follows that $f^N(\partial G)$ meets ∂D_N . We deduce, by applying Lemma 5.2 and using the fact that $\partial G \subset K(f, (m^n(r)))$, that the component Γ of $f^N(\partial G) \cap \overline{D}_N$ that contains $\alpha' := f^N(\alpha)$ is a continuum satisfying

$$\Gamma \subset f^N(K(f, (m^n(r)))) \quad \Gamma \cap \partial D_N \neq \emptyset, \quad \alpha' \in \Gamma \quad \text{and} \quad f^n(\alpha') \in D_{N+n}, \quad \text{for } n \in \mathbb{N}_0.$$

The proof now proceeds as from Step 2 in the proof of Theorem 6.2, and leads to the conclusion that $\Gamma \cap I^+(f, (m^n(r))) \neq \emptyset$, which is a contradiction. Thus we

have shown that $I^+(f, (m^n(r)))$ is connected, and hence that it is a weak spider's web.

The proof that $\bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(I^+(f, (m^n(r))))$ is also a weak spider's web now follows fairly easily. First, this set contains $I^+(f, (m^n(r)))$, so there is no unbounded closed connected set in its complement. Thus we just need to show that $\bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(I^+(f, (m^n(r))))$ is connected. However, for each $\ell \in \mathbb{N}_0$, the set $f^{-\ell}(I^+(f, (m^n(r))))$ can be shown to be connected by minor modifications of the above arguments that $I^+(f, (m^n(r)))$ is connected, and so the connectedness of the nested union $\bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(I^+(f, (m^n(r))))$ follows.

This completes the proof of Theorem 6.1. \square

In [23, Theorem 3.1], we showed that $I^+(f)$ is a weak spider's web for functions satisfying more general conditions than (1.1). We remark that Theorem 6.1 can be generalised in a similar way. Suppose that f is a transcendental entire function and that there exists a sequence of bounded, simply connected domains $(D_n)_{n \in \mathbb{N}_0}$ such that

$$(6.10) \quad f(\partial D_n) \text{ surrounds } D_{n+1}, \text{ for } n \in \mathbb{N}_0,$$

and

$$(6.11) \quad \text{every disc centred at } 0 \text{ is contained in } D_n \text{ for sufficiently large } n.$$

These domains D_n generalise the discs given by (6.1), which were used in the proof of Theorem 6.1. Using these domains, we define the set

$$I^+(f, (D_n)) = \{z : \text{there exists } (n_j) \text{ such that } f^{n_j}(z) \in \mathbb{C} \setminus D_{n_j}, \text{ for } j \in \mathbb{N}\},$$

where (n_j) is a strictly increasing sequence of positive integers that, in general, depends on z . Then the following result can be proved by making only slight changes to the proof of Theorem 6.1; we omit the details.

Theorem 6.5. *Suppose the transcendental entire function f and the sequence $(D_n)_{n \in \mathbb{N}_0}$ of bounded, simply connected domains satisfy (6.10) and (6.11). Then each of the sets*

$$I^+(f, (D_n)) \quad \text{and} \quad \bigcup_{\ell \in \mathbb{N}_0} f^{-\ell}(I^+(f, (D_n)))$$

is a weak spider's web.

We conclude this section by using Theorem 2.1 to show that if (1.1) holds and there exists an invariant curve under f , which tends to ∞ , then this curve must contain a point of $I(f)$.

Theorem 6.6. *Let f be a transcendental entire function such that condition (1.1) holds and let Γ be a simple curve tending to ∞ and invariant under f . Then $\Gamma \cap I(f) \neq \emptyset$.*

Proof. Let $\psi : \Gamma \rightarrow [0, \infty)$ be a homeomorphism and define

$$\varphi(t) = \psi \circ f \circ \psi^{-1}(t) \quad \text{and} \quad \tilde{\varphi}(t) := \max_{0 \leq s \leq t} \varphi(s), \quad \text{for } t \in [0, \infty).$$

We claim that there exists t_0 such that $\varphi^n(t_0) \rightarrow \infty$ as $n \rightarrow \infty$. The theorem follows from the claim since, if $z_0 = \psi^{-1}(t_0)$, then

$$f^n(z_0) = f^n(\psi^{-1}(t_0)) = \psi^{-1}(\varphi^n(t_0)) \rightarrow \infty \quad \text{as } n \rightarrow \infty,$$

so $z_0 \in \Gamma \cap I(f)$.

To prove the claim we use Theorem 2.1 to choose $r \geq R > 0$ such that

$$m^n(r) \text{ and } \tilde{m}^n(R) \text{ increase strictly with } n \text{ to } \infty,$$

and

$$m^n(r) \in [\tilde{m}^n(R), \tilde{m}^{n+1}(R)], \text{ for } n \in \mathbb{N}_0.$$

Without loss of generality, we can assume that Γ meets each of the circles $\{z : |z| = m^n(r)\}$, for $n \in \mathbb{N}_0$. Then there exists $T \geq 0$ such that, for each $t \geq T$,

$$m^n(r) \leq |\psi^{-1}(t)| < m^{n+1}(r), \text{ for some } n \in \mathbb{N}_0.$$

Moreover, for $t \geq T$, there exists $z_t \in \Gamma$ and a maximal $N \in \mathbb{N}_0$ such that

$$(6.12) \quad |z_t| = m^N(r) \quad \text{and} \quad \psi(z_t) \leq t.$$

Then

$$|f(z_t)| \geq m^{N+1}(r)$$

and also, because Γ is invariant under f and N is maximal for (6.12),

$$\tilde{\varphi}(t) \geq \varphi(\psi(z_t)) = \psi(f(z_t)) > t.$$

Thus we have shown that $\tilde{\varphi}(t) > t$ for $t \geq T$. Hence, by Theorem 2.1 there exists $t_0 > 0$ such that $\varphi^n(t_0) \rightarrow \infty$ as $n \rightarrow \infty$. This completes the proof. \square

Remark. It follows from the result of Short and Sixsmith [39, Theorem 1.6], mentioned in Section 2, that the conclusion of Theorem 6.6 can be strengthened to state that Γ contains uncountably many points in $I(f)$.

7. UNBOUNDED FATOU COMPONENTS

In this section we prove Theorems 1.6 and 1.7 which, for a transcendental entire function f for which the condition (1.1) holds, concern the relationship between *unbounded* Fatou components and the sets $V^+(f)$ and $V(f)$. We begin by noting that if a point z belongs to a Fatou component of f that lies in an attracting or parabolic basin, or a Siegel disc (or a preimage of a Siegel disc), then the orbit of z must be bounded. Thus any Fatou component of f that meets $V^+(f)$ must be a Baker domain (or a preimage of a Baker domain) or a wandering domain.

Recall that, if $U = U_0$ is a Fatou component, then $f^n(U) \subset U_n$ for some Fatou component U_n , for each $n \in \mathbb{N}$. A Fatou component of a transcendental entire function is called a *wandering domain* if it is not eventually periodic, that is, if $U_m \neq U_n$ for $m \neq n$, and it is called a *Baker domain* if it is periodic and $f^n(z) \rightarrow \infty$ as $n \rightarrow \infty$ for $z \in U$. Note that Baker domains of transcendental entire functions are always unbounded whereas wandering domains can be either bounded or unbounded. (For a full description of the possible types of Fatou components of a transcendental entire function see, for example, [5].)

Proof of Theorem 1.6. (a) Let f be a transcendental entire function satisfying the condition (1.1) and let U be an unbounded Fatou component of f . By Theorem 1.2 we can take $r \geq R > 0$ such that

$$m^n(r) \geq \tilde{m}^n(R), \text{ for } n \in \mathbb{N}_0, \quad \text{and} \quad \tilde{m}^n(R) \rightarrow \infty \text{ as } n \rightarrow \infty.$$

Then we can use Theorem 6.2 to show that $U \cap V^+(f) \neq \emptyset$. Indeed, if this is false, then $U \subset K(f, (m^n(r)))$ and, by Theorem 6.2, for any $\alpha \in U$ there exists

$N_1 = N_1(\alpha)$ such that any continuum K in U that contains α must lie in D_{N_1} , which is a contradiction. Therefore, we must have $U \cap V^+(f) \neq \emptyset$. As noted above, this implies that U is either a Baker domain (or a preimage of a Baker domain) or a wandering domain.

The second statement of part (a) follows from Theorem 4.1 part (c).

(b) Zheng's result (b) before the statement of Theorem 1.6 states that if

$$\limsup_{r \rightarrow \infty} \frac{m(r)}{r} = \infty,$$

then any unbounded component U of $F(f)$ must be a wandering domain. This condition is clearly satisfied if $\lim_{r \rightarrow \infty} \tilde{m}(r)/r = \infty$. The fact that such a wandering domain must be in $V^+(f)$ follows from part (a).

(c) The fact that U is a wandering domain in $V^+(f)$ follows from part (b), and $V^+(f) \subset Z^+(f)$ because the hypothesis about $\tilde{m}(r)$ implies that, for sufficiently large r ,

$$\frac{\log^+ \log^+ \tilde{m}^n(r)}{n} \rightarrow \infty \text{ as } n \rightarrow \infty,$$

by Lemma 3.1. □

Finally in this section, we prove our result about the rate of escape in certain Baker domains.

Proof of Theorem 1.7. Suppose that (1.1) holds and that U is an invariant Baker domain of the transcendental entire function f . By Theorem 2.1, we can take $r \geq R > 0$ such that

$$(7.1) \quad m^n(r) \geq \tilde{m}^n(R), \text{ for } n \in \mathbb{N}_0, \text{ and } \tilde{m}^n(R) \rightarrow \infty \text{ as } n \rightarrow \infty.$$

and

$$m^n(r) \text{ increases strictly with } n \text{ (to } \infty).$$

We now take $\alpha \in U$ and let Γ be a compact curve in U joining α to $f(\alpha)$. Moreover, we assume without loss of generality (by choosing α suitably and replacing r by some iterate $m^n(r)$ if necessary) that

$$|\alpha| \leq r < |f(\alpha)|.$$

Now let n_0 denote the largest value of $n \in \mathbb{N}_0$ such that

$$(7.2) \quad \Gamma \cap \{z : |z| = m^n(r)\} \neq \emptyset.$$

Then

$$(7.3) \quad f(\Gamma) \cap \{z : |z| \geq m^{n_0+1}(r)\} \neq \emptyset.$$

Also, $f(\alpha) \in \Gamma \cap f(\Gamma)$, so by the definition of n_0 and the fact that $(m^n(r))$ is strictly increasing, we have

$$(7.4) \quad f(\Gamma) \cap \{z : |z| < m^{n_0+1}(r)\} \neq \emptyset.$$

It follows from (7.3) and (7.4) that, if we let n_1 denote the largest value of $n \in \mathbb{N}_0$ such that

$$(7.5) \quad f(\Gamma) \cap \{z : |z| = m^n(r)\} \neq \emptyset,$$

then $n_1 \geq n_0 + 1 \geq 1$.

By repeating this process we find that, for each $k \in \mathbb{N}_0$, there exists $n_k \geq k$ such that

$$f^k(\Gamma) \cap \{z : |z| = m^{n_k}(r)\} \neq \emptyset.$$

It now follows from Lemma 4.2 that there exists $C = C_\Gamma > 1$ such that, for each $z \in \Gamma$ and each $k \in \mathbb{N}_0$,

$$|f^k(z)| \geq m^{n_k}(r)/C \geq m^k(r)/C.$$

Finally, by applying Lemma 4.2 again, we deduce from (7.1) that for any $z \in U$ there exists $C(z) > 1$ such that, for each $n \in \mathbb{N}$,

$$(7.6) \quad |f^n(z)| \geq m^n(r)/C(z) \geq \tilde{m}^n(R)/C(z),$$

as required.

If we also have

$$\liminf_{r \rightarrow \infty} \frac{\tilde{m}(r)}{r} > 1,$$

then, for any $z \in U$ there exists $k = k(C(z)) \in \mathbb{N}$ such that

$$\tilde{m}^{n+k}(R) \geq C(z)\tilde{m}^n(R), \quad \text{for } n \in \mathbb{N}_0,$$

where R was defined above, and hence, by (7.6),

$$|f^{n+k}(z)| \geq \tilde{m}^n(R), \quad \text{for } n \in \mathbb{N}_0;$$

that is, $z \in V(f)$. □

8. EXAMPLES

In this section we illustrate our results with examples of fairly simple entire functions, some of which satisfy (1.1) and some of which do not.

Our first example is a function f of order 1 which does not belong to the classes of functions covered by Theorem 1.1. We show that this function satisfies condition (1.1) and has an invariant Baker domain that lies in $V(f)$.

Example 8.1. *Let f be the function*

$$f(z) = 2z(1 + e^{-z}).$$

Then

- (a) *there exists $r > 0$ such that $m^n(r) \rightarrow \infty$ as $n \rightarrow \infty$;*
- (b) *f has an invariant Baker domain that lies in $V(f)$.*

Proof. Put $z = re^{i\theta}$, so

$$f(re^{i\theta}) = 2re^{i\theta}(1 + e^{-r \cos \theta} e^{-ir \sin \theta}),$$

and therefore

$$\begin{aligned} \mu(r, \theta) &:= |f(re^{i\theta})|^2 = 4r^2 \left((1 + e^{-r \cos \theta} \cos(r \sin \theta))^2 + (e^{-r \cos \theta} \sin(r \sin \theta))^2 \right) \\ &= 4r^2 \left(1 + 2e^{-r \cos \theta} \cos(r \sin \theta) + e^{-2r \cos \theta} \right) \\ &= 4r^2(1 - e^{-r \cos \theta})^2 + 8r^2 e^{-r \cos \theta} (1 + \cos(r \sin \theta)). \end{aligned}$$

Note that both terms in the last line are non-negative for all values of r and θ .

Now let $r_n = 2n\pi$, for $n \geq 2$, so $r_n \rightarrow \infty$ as $n \rightarrow \infty$, and $r_{n+1} \leq \frac{3r_n}{2}$. We claim that

$$\mu(r_n, \theta) \geq \left(\frac{3r_n}{2}\right)^2 \geq r_{n+1}^2, \quad \text{for } n \geq 2,$$

from which it follows that $m(r_n) \geq 3r_n/2 \geq r_{n+1}$ for $n \geq 2$, and this proves part (a) by Theorem 2.1.

To prove the claim, first observe that, if $e^{-r \cos \theta} \leq \frac{1}{4}$ or $e^{-r \cos \theta} \geq \frac{7}{4}$, then

$$\mu(r, \theta) \geq 4r^2(1 - e^{-r \cos \theta})^2 \geq 4r^2 \left(\frac{3}{4}\right)^2 = \left(\frac{3r}{2}\right)^2.$$

Suppose therefore that $\frac{1}{4} < e^{-r \cos \theta} < \frac{7}{4}$. Then

$$-\frac{1}{r} \log \frac{7}{4} < \cos \theta < -\frac{1}{r} \log \frac{1}{4},$$

so we can put $\theta = \frac{1}{2}\pi + \varepsilon(r)$, where $|\varepsilon(r)| \leq C/r$ for some positive absolute constant C . Thus, for $n \geq 2$ we have

$$r_n \sin \theta = r_n \left(1 - \frac{O(1)}{r_n^2}\right) = 2n\pi - \frac{O(1)}{2n\pi},$$

and hence

$$\cos(r_n \sin \theta) = 1 - \frac{O(1)}{(2n\pi)^2}.$$

It follows that, for large n ,

$$\mu(r_n, \theta) \geq 8r_n^2 e^{-r_n \cos \theta} (1 + \cos(r_n \sin \theta)) \geq 2r_n^2 \left(2 - \frac{O(1)}{(2n\pi)^2}\right) \geq \left(\frac{3r_n}{2}\right)^2,$$

and this completes the proof of the claim, and of part (a).

That f has an invariant Baker domain, U say, follows from [27, Theorem 2], which describes a large family of entire functions with Baker domains, including this function. Since we have just shown that for the sequence $r_n = 2n\pi$ we have $m(r_n) \geq 3r_n/2$, it follows that $\liminf_{r \rightarrow \infty} \tilde{m}(r)/r > 1$ and thus that $U \subset V(f)$ by Theorem 1.7 part (b). \square

Our next example concerns the transcendental entire function $f(z) = z + 1 + e^{-z}$, first investigated by Fatou [11] and often named after him. For this function, it is known that $F(f)$ is a completely invariant Baker domain, that $I(f)$ is a spider's web but $A(f)$ is not, and that f is strongly polynomial-like; see [10, Theorem 1.1] and [22, Example 5.4] for these results and an explanation of the terminology. We use Theorem 2.1 to show that, nevertheless, condition (1.1) does not hold for this function f .

Example 8.2. *Let f be the Fatou function,*

$$f(z) = z + 1 + e^{-z}.$$

Then there does not exist $r > 0$ such that $m^n(r) \rightarrow \infty$ as $n \rightarrow \infty$.

Proof. We will show that $\tilde{m}(r) < r$ for arbitrarily large r , from which the result follows by Theorem 2.1.

Consider the images under f of points ir , $r > 0$. As r increases, the image points travel clockwise around a circle of radius 1, whose centre is at the same time moving up the line $\operatorname{Re} z = 1$. Clearly

$$|f(ir)| = r, \quad \text{for } r = (2k+1)\pi, \quad k \in \mathbb{N},$$

and it is easy to see that there exists $\varepsilon_k > 0$, with $\varepsilon_k \rightarrow 0$ as $r \rightarrow \infty$, such that

$$(8.1) \quad m(r) \leq |f(ir)| \leq r, \quad \text{for } 2k\pi + \varepsilon_k \leq r \leq (2k+1)\pi, \quad k \in \mathbb{N}.$$

Moreover, for $r > 0$, we have

$$(8.2) \quad |f(ir)| \leq r + 1 + |e^{-ir}| \leq r + 2.$$

Now let $r_k = (2k+1)\pi - \delta_k$, where $\delta_k > 0$. Then for δ_k sufficiently small we have

$$m(s) \leq r_k, \quad \text{for } 2k\pi + \varepsilon_k \leq s \leq r_k,$$

by (8.1), and

$$\max \{m(s) : 0 \leq s \leq 2k\pi + \varepsilon_k\} \leq 2k\pi + \varepsilon_k + 2 < r_k,$$

by (8.2). Thus we have shown that

$$\tilde{m}(r_k) = \max \{m(s) : 0 \leq s \leq r_k\} < r_k, \quad \text{for } k \in \mathbb{N},$$

and this completes the proof. \square

We remark that a similar argument shows that (1.1) fails for every function of the form $f(z) = z + a + be^{-z}$, where $a, b > 0$.

The following example shows that the condition (1.1) can hold even if we have

$$(8.3) \quad \lim_{r \rightarrow \infty} \frac{\tilde{m}(r)}{r} = 1.$$

Example 8.3. Condition (1.1) holds for any function of the form

$$f(z) = z + b \sin z, \quad \text{where } b > 2\pi.$$

Proof. We claim that such a function f has the property that if $r_n = 2n\pi + \pi/2$, where $n \in \mathbb{N}$, then

$$m(r_n) \geq r_n + 2\pi = r_{n+1}, \quad \text{for sufficiently large } n,$$

from which it follows that condition (1.1) holds, by Theorem 2.1. It is also clear that (8.3) holds for functions of this form.

To prove the claim, suppose first that $z = x + iy$, where $x, y \geq 0$, $|z| = r_n$, for some $n \in \mathbb{N}$, and $y \geq \log(3r_n)$. Then clearly

$$\sin^2 x + \sinh^2 y \geq \sinh^2 y \geq r_n^2,$$

and so

$$|\sin z| \geq |z|.$$

Hence, for such z we have

$$(8.4) \quad |f(z)| \geq b|\sin z| - |z| \geq 3|z| - |z| \geq 2|z| \geq |z| + 2\pi.$$

Next suppose that $z = x + iy$, where $x, y \geq 0$ and $|z| = r_n$, for some $n \in \mathbb{N}$, and $y \leq \log(3r_n)$. Then

$$x^2 = r_n^2 - y^2 \geq r_n^2 - (\log(3r_n))^2 = r_n^2 \left(1 - \left(\frac{\log(3r_n)}{r_n} \right)^2 \right),$$

so

$$r_n \geq x \geq r_n \left(1 - \left(\frac{\log(3r_n)}{r_n} \right)^2 \right)^{1/2} \geq r_n - \frac{(\log(3r_n))^2}{r_n}.$$

Thus, for such z , we have (since $r_n = 2n\pi + \pi/2$)

$$\operatorname{Re}(\sin z) = \sin x \cosh y \geq \sin x \geq 1 - \frac{(\log(3r_n))^2}{r_n},$$

from which it follows that

$$(8.5) \quad |f(z)| = |z + b \sin z| \geq r_n + b - o(1) \quad \text{as } n \rightarrow \infty,$$

uniformly for such z .

The claim about $m(r_n)$ now follows from (8.4) and (8.5), together with the symmetry properties of the sine function. \square

It follows from Theorem 1.1 that condition (1.1) always holds for transcendental entire functions of order less than $1/2$. Our final example shows that this condition may or may not hold for transcendental entire functions of order $1/2$.

Example 8.4. (a) Condition (1.1) does not hold for the function $f(z) = \cos \sqrt{z}$.
 (b) Condition (1.1) holds for the function $g(z) = 2z \cos \sqrt{z}$.

Proof. (a) Since f is bounded on the positive real axis, it follows that $m(r)$ is bounded and thus that there is no $r > 0$ such that $m^n(r) \rightarrow \infty$ as $n \rightarrow \infty$.

(b) First note that the minimum modulus of g is attained on the positive real axis for every $r > 0$. This can be deduced, for example, from the fact that g is an entire function of order $1/2$ with its zeros on the positive real axis, and so (see [41]) can be written in the form

$$g(z) = 2z \prod_{n=0}^{\infty} \left(1 - \frac{z}{(n + 1/2)^2 \pi^2} \right).$$

Now suppose that $r \geq 9\pi^2$. Then $r \in [n^2\pi^2, (n+1)^2\pi^2)$ for some integer $n \geq 3$, and it follows that

$$\tilde{m}(r) \geq m(n^2\pi^2) = 2n^2\pi^2 > (n+1)^2\pi^2 > r.$$

Thus we have shown that $\tilde{m}(r) > r$ for $r \geq 9\pi^2$, so it follows from Theorem 2.1 that condition (1.1) is satisfied. \square

Remarks. 1. Example 8.4 can be modified to apply to transcendental entire functions of any positive integer order p , giving that condition (1.1) is not satisfied for the function $f(z) = \cos z^p$, but is satisfied for the function $g(z) = 2z \cos z^p$.

2. The case of functions of order $1/2$, *minimal type*, mentioned in the statement of Baker's conjecture, is rather delicate. In forthcoming work we will show, using a method pioneered by Kjellberg [14, page 821], that there are examples of such functions that do satisfy condition (1.1) and other examples that do not.

REFERENCES

- [1] I.N. Baker, Zusammensetzungen ganzer Funktionen, *Math. Z.*, 69 (1958), 121–163.
- [2] I. N. Baker, The iteration of polynomials and transcendental entire functions, *J. Austral. Math. Soc. (Series A)*, 30 (1981), 483–495.
- [3] P. D. Barry, On a theorem of Besicovitch, *Quart. J. Math. Oxford (2)*, 14 (1963), 293–302.
- [4] A. F. Beardon, *Iteration of rational functions*, Graduate Texts in Mathematics 132, Springer-Verlag, 1991.
- [5] W. Bergweiler, Iteration of meromorphic functions, *Bull. Amer. Math. Soc.*, 29 (1993), 151–188.
- [6] W. Bergweiler and A. Hinkkanen, On semiconjugation of entire functions, *Math. Proc. Camb. Phil. Soc.*, 126 (1999), 565–574.
- [7] W. Bergweiler, P. J. Rippon and G. M. Stallard, Multiply connected wandering domains of entire functions, *Proc. London Math. Soc.*, 107 (2013), 1261–1301.
- [8] L. Carleson and T. W. Gamelin, *Complex dynamics*, Springer-Verlag, 1993.
- [9] A. E. Eremenko, On the iteration of entire functions, *Dynamical systems and ergodic theory*, Banach Center Publications 23, Polish Scientific Publishers, Warsaw, 1989, 339–345.
- [10] V. Evdoridou, Fatou’s web, *Proc. Amer. Math. Soc.*, 144 (2016), 5227–5240.
- [11] P. Fatou, Sur l’itération des fonctions transcendentes entières, *Acta Math.*, 47 (1926), 337–360.
- [12] W. H. J. Fuchs, Proof of a conjecture of G. Pólya concerning gap series, *Illinois J. Math.*, 7 (1963), 661–667.
- [13] W. K. Hayman, Angular value distribution of power series with gaps, *Proc. London Math. Soc. (3)*, 24 (1972), 590–624.
- [14] W. K. Hayman, *Subharmonic functions, Volume 2*, Academic Press, 1989.
- [15] M. Heins, Entire functions with bounded minimum modulus; subharmonic function analogues, *Annals of Mathematics*, 49 (1948), 200–213.
- [16] A. Hinkkanen, Entire functions with bounded Fatou components, in *Transcendental dynamics and complex analysis*, P.J. Rippon and G.M. Stallard (eds.), Cambridge University Press, 2008, 187–216.
- [17] A. Hinkkanen and J. Miles, Growth conditions for entire functions with only bounded Fatou components, *J. Anal. Math.*, 108 (2009), 87–118.
- [18] J.E. Littlewood and A.C. Offord, On the distribution of zeros and a -values of a random integral function (II), *Annals of Mathematics, Second Series*, 49 (4) (1948), 885–952.
- [19] J. Milnor, *Dynamics in one complex variable*, Third Edition, Princeton University Press, 2006.
- [20] M. H. A. Newman, *Elements of the topology of plane sets of points*, Cambridge University Press, Reprinted 2008.
- [21] D. A. Nicks, P. J. Rippon and G. M. Stallard, Baker’s conjecture for functions with real zeros. Preprint, arXiv: 1502.02526.
- [22] J. W. Osborne, Connectedness properties of the set where the iterates of an entire function are bounded, *Math. Proc. Camb. Phil. Soc.*, 155 (3) (2013), 391–410.
- [23] J. W. Osborne, P. J. Rippon and G. M. Stallard, Connectedness properties of the set where the iterates of an entire function are unbounded, *Ergodic Theory and Dynamical Systems*, First view, doi: 10.1017/etds.2015.85, arXiv: 1504.05611.
- [24] J. W. Osborne and D. J. Sixsmith, On the set where the iterates of an entire function are neither escaping nor bounded, *Ann. Acad. Sci. Fenn. Math.*, 41 (2016), 561–578.
- [25] L. Rempe, On a question of Eremenko concerning escaping components of entire functions, *Bull. London Math. Soc.*, 39 (4) (2007), 661–666.
- [26] L. Rempe, Connected escaping sets of exponential maps, *Ann. Acad. Sci. Fenn. Math.*, 36 (2011), 71–80.
- [27] P. J. Rippon and G. M. Stallard, Families of Baker domains II, *Conformal geometry and dynamics*, 3 (1999), 67–78.
- [28] P. J. Rippon and G. M. Stallard, On sets where iterates of a meromorphic function zip towards infinity, *Bull. London Math. Soc.*, 32 (2000), 528–536.

- [29] P. J. Rippon and G. M. Stallard, On questions of Fatou and Eremenko, *Proc. Amer. Math. Soc.*, 133 (2005), 1119–1126.
- [30] P. J. Rippon and G. M. Stallard, Slow escaping points of meromorphic functions, *Trans. Amer. Math. Soc.*, 363 (8) (2011), 4171–4201.
- [31] P. J. Rippon and G. M. Stallard, Functions of small growth with no unbounded Fatou components, *J. Anal. Math.*, 108 (2009), 61–86.
- [32] P. J. Rippon and G. M. Stallard, Fast escaping points of entire functions, *Proc. London Math. Soc.*, 105 (4) (2012), 787–820.
- [33] P. J. Rippon and G. M. Stallard, Baker’s conjecture and Eremenko’s conjecture for functions with negative zeros, *J. Anal. Math.*, 120 (2013), 291–309.
- [34] P. J. Rippon and G. M. Stallard, Regularity and fast escaping points of entire functions, *International Mathematics Research Notices*, Oct 2014, Vol. 2014 Issue 19, 5203–5229.
- [35] P. J. Rippon and G. M. Stallard, A sharp growth condition for a fast escaping spider’s web, *Advances in Mathematics*, 244 (2013), 337–353.
- [36] P. J. Rippon and G. M. Stallard, Annular itineraries for entire functions, *Trans. Amer. Math. Soc.*, 367 (2015), 377–399.
- [37] P. J. Rippon and G. M. Stallard, Boundaries of univalent Baker domains. To appear in *J. Anal. Math.*, arXiv: 1411.6999.
- [38] G. Rottenfusser, J. Rückert, L. Rempe and D. Schleicher, Dynamic rays of bounded-type entire functions, *Annals of Mathematics*, 173 (1) (2011), 77–125.
- [39] I. Short and D. J. Sixsmith, Escaping sets of continuous functions. To appear in *J. Anal. Math.*, arXiv:1601.04010.
- [40] D. J. Sixsmith, Maximally and non-maximally fast escaping points of transcendental entire functions, *Math. Proc. Camb. Phil. Soc.*, 158 (2) (2015), 365–383.
- [41] E. C. Titchmarsh, *The theory of functions*, Oxford Univ. Press, 2nd edition, 1939.
- [42] A. Wiman, Sur une extension d’un théorème de M. Hadamard, *Ark. Mat. Astr. Fys.* (1905), No. 14.
- [43] J. H. Zheng, Unbounded domains of normality of entire functions of small growth, *Math. Proc. Camb. Phil. Soc.*, 128 (2000), 355–361.

SCHOOL OF MATHEMATICS AND STATISTICS, THE OPEN UNIVERSITY, WALTON HALL, MILTON KEYNES MK7 6AA, UK

E-mail address: john.osborne@open.ac.uk, phil.rippon@open.ac.uk, gwyneth.stallard@open.ac.uk