

THE JOINT UNIVERSALITY AND THE FUNCTIONAL INDEPENDENCE FOR LERCH ZETA-FUNCTIONS

ANTANAS LAURINĆIKAS¹ AND KOHJI MATSUMOTO

Abstract. The joint universality theorem for Lerch zeta-functions $L(\lambda_l, \alpha_l, s)$ ($1 \leq l \leq n$) is proved, in the case when λ_l s are rational numbers and α_l s are transcendental numbers. The case $n = 1$ was known before ([12]); the rationality of λ_l s is used to establish the theorem for the "joint" case $n \geq 2$. As a corollary, the joint functional independence for those functions is shown.

1. Introduction

Let $s = \sigma + it$ be a complex variable, and let $\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}$ and \mathbb{C} denote the set of all natural numbers, integers, rational numbers, real numbers and complex numbers, respectively. The Lerch zeta-function $L(\lambda, \alpha, s)$, for $\sigma > 1$, is defined by

$$L(\lambda, \alpha, s) = \sum_{m=0}^{\infty} \frac{e^{2\pi i \lambda m}}{(m + \alpha)^s}.$$

Here $\alpha, \lambda \in \mathbb{R}$, $0 < \alpha \leq 1$, are fixed parameters. When $\lambda \in \mathbb{Z}$ the Lerch zeta-function $L(\lambda, \alpha, s)$ reduces to the Hurwitz zeta-function $\zeta(s, \alpha)$. If $\lambda \notin \mathbb{Z}$, then the function $L(\lambda, \alpha, s)$ is analytically continuable to an entire function. Clearly, in this case we may suppose that $0 < \lambda < 1$. In what follows we will deal with this case only.

The Lerch zeta-function is one of the classical objects in number theory, introduced by M. Lerch [16] in 1887.

In recent years the value-distribution of the Lerch zeta-function was studied by D. Klusch, R. Garunkštis, M. Katsurada, W. Zhang, by the authors and other mathematicians. In [12] the universality theorem for the function $L(\lambda, \alpha, s)$ was proved. In order to state it we need some notation.

Received February 18, 1999.

¹Partially supported by Grant from Lithuanian Foundation of Studies and Science.

By $\text{meas}\{A\}$ we denote the Lebesgue measure of the set A , and, for $T > 0$, we use the notation

$$\nu_T(\dots) = \frac{1}{T} \text{meas}\{\tau \in [0, T], \dots\},$$

where in place of dots some condition satisfied by τ is to be written. Let $D = \{s \in \mathbb{C} : 1/2 < \sigma < 1\}$. Then the result of [12] is as follows.

Let α be a transcendental number. Let K be a compact subset of the strip D with the connected complement, $f(s)$ be a continuous function on K which is analytic in the interior of K . Then for any $\varepsilon > 0$ it holds that

$$\liminf_{T \rightarrow \infty} \nu_T \left(\sup_{s \in K} |L(\lambda, \alpha, s + i\tau) - f(s)| < \varepsilon \right) > 0.$$

The universality for $L(\lambda, \alpha, s)$ was also studied in [5], [13].

It is the purpose of the present paper to obtain a joint universality theorem for Lerch zeta-functions. Suppose $n \geq 2$.

THEOREM 1. *Let $\alpha_1, \dots, \alpha_n$ be transcendental numbers, $\lambda_1 = a_1/q_1, \dots, \lambda_n = a_n/q_n$, $(a_1, q_1) = 1, \dots, (a_n, q_n) = 1$, where q_1, \dots, q_n are distinct positive integers and a_1, \dots, a_n are positive integers with $a_1 < q_1, \dots, a_n < q_n$. Let K_1, \dots, K_n be compact subsets of the strip D with connected complements, and, for $1 \leq l \leq n$, let $f_l(s)$ be a continuous function on K_l which is analytic in the interior of K_l . Then for every $\varepsilon > 0$ it holds that*

$$\liminf_{T \rightarrow \infty} \nu_T \left(\sup_{1 \leq l \leq n} \sup_{s \in K_l} |L(\lambda_l, \alpha_l, s + i\tau) - f_l(s)| < \varepsilon \right) > 0.$$

Joint universality theorems for Dirichlet L -functions were obtained by B. Bagchi [2], [3], S.M. Gonek [7], and S.M. Voronin [18], [19]. For more general Dirichlet series such theorems were proved in [8], [9], [14].

The proof of Theorem 1 is based on Bagchi’s method [2], [3], but some new ideas are necessary for the proof of Lemmas 5 and 6 below.

In the case of the aforementioned universality theorem [12] for a single zeta-function, the arithmetic nature of λ is irrelevant. However, in the proof of Theorem 1, the fact that $\lambda_l \in \mathbb{Q}$ ($1 \leq l \leq n$) is used essentially. In Section 4 we will discuss briefly the case when $\lambda_l \notin \mathbb{Q}$.

As an application of Theorem 1, we will show the joint functional independence.

THEOREM 2. *Let $\alpha_l, \lambda_l = a_l/q_l$ be as in Theorem 1, and F_j ($0 \leq j \leq k$) be continuous functions on \mathbb{C}^{N^n} . Suppose*

$$\sum_{j=0}^k s^j F_j(L(\lambda_1, \alpha_1, s), \dots, L(\lambda_n, \alpha_n, s), L'(\lambda_1, \alpha_1, s), \dots, L'(\lambda_n, \alpha_n, s), \dots, L^{(N-1)}(\lambda_1, \alpha_1, s), \dots, L^{(N-1)}(\lambda_n, \alpha_n, s)) = 0$$

identically for all $s \in \mathbb{C}$. Then $F_j \equiv 0$ ($0 \leq j \leq k$).

This theorem gives a generalization of the result proved in Garunkštis-Laurinčikas [6]. A quite different approach to this type of problems has recently been developed by Amou-Katsurada [1].

2. A joint limit theorem for Lerch zeta-functions

For the proof of Theorem 1 we will apply a joint limit theorem in the sense of weak convergence of probability measures for the Lerch zeta-functions $L(\lambda_1, \alpha_1, s), \dots, L(\lambda_n, \alpha_n, s)$ in the space of analytic functions. Denote by $H(D)$ the space of analytic on D functions equipped with the topology of uniform convergence on compacta. Let $\mathcal{B}(S)$ stand for the class of Borel sets of the space S . Define on $(H^n(D), \mathcal{B}(H^n(D)))$ the probability measure

$$P_T(A) = \nu_T\left(\left(L(\lambda_1, \alpha_1, s+i\tau), \dots, L(\lambda_n, \alpha_n, s+i\tau)\right) \in A\right), \quad A \in \mathcal{B}(H^n(D)).$$

What we need is a limit theorem in the sense of weak convergence of probability measures for P_T as $T \rightarrow \infty$, with an explicit form of the limit measure. Denote by γ the unit circle on \mathbb{C} , i.e. $\gamma = \{s \in \mathbb{C} : |s| = 1\}$, and let

$$\Omega = \prod_{m=0}^{\infty} \gamma_m,$$

where $\gamma_m = \gamma$ for all $m = 0, 1, 2, \dots$. With the product topology and pointwise multiplication the infinite dimensional torus Ω is a compact topological Abelian group. Denoting by m_H the probability Haar measure on $(\Omega, \mathcal{B}(\Omega))$, we obtain the probability space $(\Omega, \mathcal{B}(\Omega), m_H)$. Let $\omega(m)$ be the projection of $\omega \in \Omega$ to the coordinate space γ_m , and define on the probability space $(\Omega, \mathcal{B}(\Omega), m_H)$ the $H^n(D)$ -valued random element $L(s, \omega)$ by

$$L(s, \omega) = (L(\lambda_1, \alpha_1, s, \omega), \dots, L(\lambda_n, \alpha_n, s, \omega)),$$

where

$$L(\lambda_l, \alpha_l, s, \omega) = \sum_{m=0}^{\infty} \frac{e^{2\pi i \lambda_l m \omega(m)}}{(m + \alpha_l)^s}, \quad s \in D, \omega \in \Omega, l = 1, \dots, n.$$

The proof that $L(\lambda_l, \alpha_l, s, \omega)$ is an $H(D)$ -valued random element can be found in [11]. Let P_L stand for the distribution of the random element $L(s, \omega)$, i.e.

$$P_L(A) = m_H(\omega \in \Omega : L(s, \omega) \in A), \quad A \in \mathcal{B}(H^n(D)).$$

LEMMA 1. *The probability measure P_T converges weakly to P_L as $T \rightarrow \infty$.*

Proof. Let $D_0 = \{s \in \mathbb{C} : \sigma > 1/2\}$. Then in [15] the result of the lemma was proved in the case of the space $H^n(D_0)$. Obviously, from this the lemma follows.

3. The support of the random element L

In this section we will consider the support of the measure P_L . We recall that the minimal closed set $S_{P_L} \subseteq H^n(D)$ such that $P_L(S_{P_L}) = 1$ is called the support of P_L . The set S_{P_L} consists of all $\underline{f} \in H^n(D)$ such that for every neighbourhood \mathcal{G} of \underline{f} the inequality $P_L(\mathcal{G}) > 0$ is satisfied.

The support of the distribution of the random element X is called the support of X and is denoted by S_X .

LEMMA 2. *Let $\{X_m\}$ be a sequence of independent $H^n(D)$ -valued random elements, and suppose that the series*

$$\sum_{m=1}^{\infty} X_m$$

converges almost surely. Then the support of the sum of this series is the closure of the set of all $\underline{f} \in H^n(D)$ which may be written as a convergent series

$$\underline{f} = \sum_{m=1}^{\infty} \underline{f}_m, \quad \underline{f}_m \in S_{X_m}.$$

Proof of the lemma in the case $n = 1$ is given in [10], Theorem 1.7.10. The proof when $n > 1$ is similar to that of the case $n = 1$.

Let $\underline{f}(s) = (f_1(s), \dots, f_n(s)) \in H^n(D)$. Then we write

$$|\underline{f}(s)|^2 = \sum_{l=1}^n |f_l(s)|^2.$$

LEMMA 3. Let $\{\underline{f}_m = (f_{1m}, \dots, f_{nm}), m \geq 1\}$ be a sequence in $H^n(D)$ which satisfies:

a) If μ_1, \dots, μ_n are complex measures on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ with compact supports contained in D such that

$$\sum_{m=1}^{\infty} \left| \sum_{l=1}^n \int_{\mathbb{C}} f_{lm} d\mu_l \right| < \infty,$$

then

$$\int_{\mathbb{C}} s^r d\mu_l(s) = 0$$

for all $l = 1, \dots, n, r = 0, 1, 2, \dots$

b) The series

$$\sum_{m=1}^{\infty} \underline{f}_m$$

converges in $H^n(D)$.

c) For any compact $K \subseteq D$

$$\sum_{m=1}^{\infty} \sup_{s \in K} |\underline{f}_m(s)|^2 < \infty.$$

Then the set of all convergent series

$$\sum_{m=1}^{\infty} a_m \underline{f}_m$$

with $a_m \in \gamma$ is dense in $H^n(D)$.

Proof. This lemma is Lemma 5.2.9 of [2], see also [3]. In [10] the proof in the case $n = 1$ is given, see Theorem 6.3.10. The proof of the general case is obtained in a similar way.

Now we state two lemmas on entire functions of exponential type. Recall that an entire function $f(s)$ is of exponential type if

$$\limsup_{r \rightarrow \infty} \frac{\log |f(re^{i\theta})|}{r} < \infty$$

uniformly in θ , $|\theta| \leq \pi$.

LEMMA 4. *Let μ be a complex measure on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ with the compact support contained in the half-plane $\sigma > \sigma_0$, and let*

$$f(z) = \int_{\mathbb{C}} e^{sz} d\mu(s), \quad z \in \mathbb{C}.$$

If $f(z) \not\equiv 0$, then

$$\limsup_{x \rightarrow \infty} \frac{\log |f(x)|}{x} > \sigma_0.$$

This lemma is due to B. Bagchi [2]. For the proof see Lemma 6.4.10 of [10].

Let \mathcal{M} be a set of natural numbers having a positive density, i.e.

$$(1) \quad \lim_{x \rightarrow \infty} \frac{1}{x} \#\{m \in \mathcal{M} : m \leq x\} = d > 0.$$

LEMMA 5. *Let $f(s)$ be an entire function of exponential type, and let*

$$\limsup_{r \rightarrow \infty} \frac{\log |f(r)|}{r} > -1.$$

Then

$$\sum_{m \in \mathcal{M}} |f(\log m)| = \infty.$$

Proof. Let $\alpha > 0$ be such that

$$(2) \quad \limsup_{y \rightarrow \infty} \frac{\log |f(\pm iy)|}{y} \leq \alpha.$$

Let us fix a positive number β such that $\alpha\beta < \pi$, and suppose, on the contrary, that

$$(3) \quad \sum_{m \in \mathcal{M}} |f(\log m)| < \infty.$$

Consider the set $A = \{m \in \mathbb{N} : \exists r \in ((m - 1/4)\beta, (m + 1/4)\beta) \text{ and } |f(r)| \leq e^{-r}\}$. Let, for brevity,

$$m_{\mathcal{M}}(x) = \sum_{\substack{m \leq x \\ m \in \mathcal{M}}} 1.$$

Clearly, we have

$$(4) \quad \sum_{m \in \mathcal{M}} |f(\log m)| \geq \sum_{m \notin A} \sum'_m |f(\log k)| \geq \sum_{m \notin A} \sum'_m \frac{1}{k},$$

where \sum'_m denotes the sum extended over all natural numbers $k \in \mathcal{M}$ satisfying $(m - 1/4)\beta < \log k \leq (m + 1/4)\beta$. If we denote

$$a = \exp \left\{ \left(m - \frac{1}{4} \right) \beta \right\}, \quad b = \exp \left\{ \left(m + \frac{1}{4} \right) \beta \right\},$$

then we have that

$$\sum'_m \frac{1}{k} = \sum_{\substack{k \in \mathcal{M} \\ a < k \leq b}} \frac{1}{k}.$$

Summing by parts, we find

$$(5) \quad \sum_{\substack{k \in \mathcal{M} \\ a < k \leq b}} \frac{1}{k} = \frac{1}{b} \sum_{\substack{k \in \mathcal{M} \\ a < k \leq b}} 1 + \int_a^b \left(\sum_{\substack{k \in \mathcal{M} \\ a < k \leq u}} 1 \right) \frac{du}{u^2}.$$

Obviously,

$$\sum_{\substack{k \in \mathcal{M} \\ a < k \leq u}} 1 = m_{\mathcal{M}}(u) - m_{\mathcal{M}}(a).$$

The assumption (1) implies

$$m_{\mathcal{M}}(x) = dx(1 + o(1)), \quad x \rightarrow \infty,$$

hence, for any $\varepsilon > 0$, there exists a number $x_0 = x_0(\varepsilon)$ such that

$$\begin{aligned} m_{\mathcal{M}}(u) &\geq du(1 - \varepsilon), \\ m_{\mathcal{M}}(a) &\leq da(1 + \varepsilon) \end{aligned}$$

if $a \geq x_0$. Therefore

$$(6) \quad \sum_{\substack{k \in \mathcal{M} \\ a < k \leq u}} 1 \geq d((u - a) - \varepsilon(a + u)).$$

Let η satisfy the inequality $1 < \eta < \exp\{\beta/2\}$, and consider the case $u \geq \eta a$. Then we have

$$\begin{aligned} \frac{1}{2}(u - a) - \varepsilon(a + u) &\geq \frac{1}{2}\left(u - \frac{u}{\eta}\right) - \varepsilon\left(\frac{u}{\eta} + u\right) \\ &= u\left(\left(\frac{1}{2} - \varepsilon\right) - \frac{1}{\eta}\left(\frac{1}{2} + \varepsilon\right)\right) > 0 \end{aligned}$$

if we choose ε sufficiently small. Hence and from (6) we obtain

$$\sum_{\substack{k \in \mathcal{M} \\ a < k \leq u}} 1 \geq \frac{d}{2}(u - a), \quad u \geq \eta a.$$

Combining this with (5), and using partial summation again, we have

$$\begin{aligned} (7) \quad \sum_{\substack{k \in \mathcal{M} \\ a < k \leq b}} \frac{1}{k} &\geq \frac{d}{2b}(b - a) + \frac{d}{2} \int_{\eta a}^b (u - a) \frac{du}{u^2} \\ &\geq \frac{d}{2} \left\{ \frac{1}{b}([b] - [\eta a]) + \int_{\eta a}^b ([u] - [\eta a]) \frac{du}{u^2} + \frac{B}{a} \right\} = \frac{d}{2} \sum_{\eta a < k \leq b} \frac{1}{k} + \frac{B}{a}, \end{aligned}$$

where $[x]$ denotes the integer part of x , and B is a number (not always the same) bounded by a constant. Clearly,

$$\begin{aligned} \sum_{\eta a < k \leq b} \frac{1}{k} &= \log b - \log(\eta a) + \frac{B}{\eta a} \\ &= \left(m + \frac{1}{4}\right)\beta - \log \eta - \left(m - \frac{1}{4}\right)\beta + Be^{-m\beta} \\ &= \frac{\beta}{2} - \log \eta + Be^{-m\beta}. \end{aligned}$$

From the choice of η it follows that

$$\frac{\beta}{2} - \log \eta > 0.$$

Now (7) shows

$$\sum_{\substack{k \in \mathcal{M} \\ a < k \leq b}} \frac{1}{k} \geq \frac{d}{2} \left(\frac{\beta}{2} - \log \eta\right) + Be^{-m\beta}.$$

This together with (3) and (4) implies

$$\sum_{m \notin A} \left(\frac{d}{2} \left(\frac{\beta}{2} - \log \eta \right) + B e^{-m\beta} \right) \leq \sum_{m \in \mathcal{M}} |f(\log m)| < \infty,$$

hence

$$(8) \quad \sum_{m \notin A} 1 < \infty.$$

Let $A = \{a_m : a_1 < a_2 < \dots\}$. Then (8) gives that

$$(9) \quad \lim_{m \rightarrow \infty} \frac{a_m}{m} = 1.$$

By the definition of the set A , there exists a sequence $\{\lambda_m\}$ such that

$$\left(a_m - \frac{1}{4} \right) \beta < \lambda_m \leq \left(a_m + \frac{1}{4} \right) \beta,$$

and $|f(\lambda_m)| \leq e^{-\lambda_m}$. Hence, in view of (9),

$$(10) \quad \lim_{m \rightarrow \infty} \frac{\lambda_m}{m} = \beta,$$

and

$$\limsup_{m \rightarrow \infty} \frac{\log |f(\lambda_m)|}{\lambda_m} \leq -1.$$

Now we apply Theorem 6.4.12 of [10]. The assumptions of that theorem are satisfied by (10), (2), and the condition $\alpha\beta < \pi$. Hence by that theorem it follows that

$$\limsup_{r \rightarrow \infty} \frac{\log |f(r)|}{r} \leq -1.$$

This contradicts the assumption of the lemma, and Lemma 5 is proved.

LEMMA 6. *The support of the measure P_L is the whole of $H^n(D)$.*

Proof. It follows from the definition of Ω that $\{\omega(m)\}$ is a sequence of independent random variables with respect to the measure m_H . Hence $\{\underline{f}_m(s, \omega(m)), m \in \mathbb{N} \cup \{0\}\}$ is a sequence of independent $H^n(D)$ -valued random elements, where

$$\underline{f}_m(s, \omega(m)) = \left(\frac{e^{2\pi i \lambda_1 m \omega(m)}}{(m + \alpha_1)^s}, \dots, \frac{e^{2\pi i \lambda_n m \omega(m)}}{(m + \alpha_n)^s} \right).$$

The support of each $\omega(m)$ is the unit circle γ . Therefore the set $\{f_m(s, a) : a \in \gamma\}$ is the support of the random element $f_m(s, \omega(m))$. Consequently, by Lemma 2 the closure of the set of all convergent series

$$\sum_{m=0}^{\infty} f_m(s, a_m), \quad a_m \in \gamma,$$

is the support of the random element $L(s, \omega)$. It remains to check that the latter set is dense in $H^n(D)$.

Let μ_1, \dots, μ_n be complex measures on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ with compact supports contained in D such that

$$(11) \quad \sum_{m=0}^{\infty} \left| \sum_{l=1}^n \int_{\mathbb{C}} \frac{e^{2\pi i \lambda_l m}}{(m + \alpha_l)^s} d\mu_l(s) \right| < \infty.$$

It is well known that for all $s \in \mathbb{C}$

$$e^s = 1 + B|s|e^{|s|}.$$

Therefore, for $m \geq 2$,

$$\begin{aligned} (m + \alpha_l)^{-s} &= m^{-s} \left(1 + \frac{\alpha_l}{m}\right)^{-s} = m^{-s} \exp \left\{ -s \log \left(1 + \frac{\alpha_l}{m}\right) \right\} \\ &= m^{-s} \exp \left\{ \frac{B|s|}{m} \right\} = m^{-s} \left(1 + \frac{B|s|}{m} e^{B|s|}\right) \\ &= m^{-s} + Bm^{-1-\sigma} |s| e^{B|s|}. \end{aligned}$$

Hence, taking into account the properties of the measures μ_1, \dots, μ_n , we deduce from (11) that

$$\sum_{m=0}^{\infty} \left| \sum_{l=1}^n \int_{\mathbb{C}} \frac{e^{2\pi i \lambda_l m}}{m^s} d\mu_l(s) \right| < \infty,$$

which can be rewritten in the form

$$(12) \quad \sum_{\substack{m=0 \\ m \equiv r \pmod{q}}}^{\infty} \left| \sum_{l=1}^n \int_{\mathbb{C}} \frac{e^{2\pi i \lambda_l r}}{m^s} d\mu_l(s) \right| < \infty, \quad 1 \leq r \leq q,$$

where $q = [q_1, \dots, q_n]$. Now let

$$\nu_r(A) = \sum_{l=1}^n e^{2\pi i \lambda_l r} \mu_l(A), \quad A \in \mathcal{B}(\mathbb{C}), \quad 1 \leq r \leq q.$$

Note that the measures ν_1, \dots, ν_q have the same properties as μ_1, \dots, μ_n . Using this notation, we may write the relation (12) as follows:

$$(13) \quad \sum_{\substack{m=0 \\ m \equiv r \pmod{q}}}^{\infty} \left| \int_{\mathbb{C}} m^{-s} d\nu_r(s) \right| < \infty, \quad 1 \leq r \leq q.$$

Let

$$\tilde{\varrho}_r(z) = \int_{\mathbb{C}} e^{-sz} d\nu_r(s), \quad z \in \mathbb{C}.$$

Then (13) becomes the following condition

$$(14) \quad \sum_{\substack{m=0 \\ m \equiv r \pmod{q}}}^{\infty} |\tilde{\varrho}_r(\log m)| < \infty, \quad 1 \leq r \leq q.$$

By Lemma 4 we obtain that $\tilde{\varrho}_r(z) \equiv 0$, or

$$\limsup_{x \rightarrow \infty} \frac{\log |\tilde{\varrho}_r(x)|}{x} > -1, \quad 1 \leq r \leq q.$$

Lemma 5 shows that the last inequality contradicts (14). Hence

$$(15) \quad \tilde{\varrho}_r(z) \equiv 0$$

for $1 \leq r \leq q$. Let

$$\varrho_l(z) = \int_{\mathbb{C}} e^{-sz} d\mu_l(s), \quad z \in \mathbb{C}, \quad l = 1, \dots, n.$$

Then by the definitions of ν_r and $\tilde{\varrho}_r$ we have

$$\begin{aligned} \tilde{\varrho}_r(z) &= \int_{\mathbb{C}} e^{-sz} \sum_{l=1}^n e^{2\pi i \lambda_l r} d\mu_l(s) = \sum_{l=1}^n e^{2\pi i \lambda_l r} \int_{\mathbb{C}} e^{-sz} d\mu_l(s) \\ &= \sum_{l=1}^n e^{2\pi i \lambda_l r} \varrho_l(z), \end{aligned}$$

which is identically equal to zero by (15). Multiplying by $e^{-2\pi i \lambda_j}$, we have

$$(16) \quad \sum_{l=1}^n e^{2\pi i (\lambda_l - \lambda_j) r} \varrho_l(z) \equiv 0, \quad 1 \leq r \leq q.$$

Taking into account that

$$\sum_{r=1}^q e^{2\pi i(\lambda_l - \lambda_j)r} = \begin{cases} q & \text{if } (\lambda_l - \lambda_j) \in \mathbb{Z}, \\ 0 & \text{otherwise,} \end{cases}$$

and the fact that $\lambda_l - \lambda_j \in \mathbb{Z}$ only if $l = j$, and summing (16) over $r = 1, \dots, q$, we find that

$$\varrho_j(z) = \int_{\mathbb{C}} e^{-sz} d\mu_j(s) \equiv 0, \quad j = 1, 2, \dots, n.$$

Differentiating this equality r times and then putting $z = 0$, we find that

$$\int_{\mathbb{C}} s^r \mu_j(s) = 0$$

for all $j = 1, \dots, n, r = 0, 1, 2, \dots$. Thus the condition a) of Lemma 3 for the sequence $\{f_m(s, 1), m \geq 1\}$ is satisfied.

Let, for a natural number N ,

$$S(\lambda, N) = \sum_{m=0}^N e^{2\pi i\lambda m}.$$

If $\lambda \notin \mathbb{Z}$, then we have

$$(17) \quad S(\lambda, N) = \frac{1 - e^{2\pi i\lambda(N+1)}}{1 - e^{2\pi i\lambda}}$$

which is uniformly bounded for all $N \geq 1$. Summing by parts, we find

$$\sum_{m=0}^N \frac{e^{2\pi i\lambda m}}{(m + \alpha)^s} = S(\lambda, N)(N + \alpha)^{-s} + s \int_0^N S(\lambda, u) \frac{du}{(u + \alpha)^{s+1}}.$$

Taking $N \rightarrow \infty$ we obtain

$$\sum_{m=0}^{\infty} \frac{e^{2\pi i\lambda m}}{(m + \alpha)^s} = s \int_0^{\infty} S(\lambda, u) \frac{du}{(u + \alpha)^{s+1}},$$

which converges for $\sigma > 0$ in view of (17). Consequently, the series

$$(18) \quad \sum_{m=0}^{\infty} \frac{e^{2\pi i\lambda m}}{(m + \alpha)^s}$$

with $\lambda \notin \mathbb{Z}$ converges (Corollary 2.1.3 of [10]) uniformly on compacta in the half-plane $\sigma > \sigma_0$ for any $\sigma_0 > 0$. This shows that the series

$$\sum_{m=0}^{\infty} \underline{f}_m(s, 1)$$

converges in $H^n(D)$, i.e. the condition b) of Lemma 3 holds for the sequence $\{\underline{f}_m(s, 1), m \geq 1\}$. The condition c) of Lemma 3 is also satisfied clearly, since for $s \in K$ we have that $\sigma > 1/2$.

Now, applying Lemma 3, we have that the set of all convergent series

$$\sum_{m=0}^{\infty} a_m \underline{f}_m(s, 1) = \sum_{m=0}^{\infty} \underline{f}_m(s, a_m)$$

with $a_m \in \gamma$ is dense in $H^n(D)$. This completes the proof of the lemma.

4. Proof of Theorem 1

The following deduction of Theorem 1 from the above lemmas is standard (cf. Section 6.5 of [10]), but we present it for the convenience of readers.

We begin with the Mergelyan theorem.

LEMMA 7. *Let K be a compact subset of \mathbb{C} whose complement is connected. Then any continuous function $f(s)$ on K which is analytic in the interior of K is approximable uniformly on K by polynomials of s .*

Proof is given, for example, in [20].

Proof of Theorem 1. First suppose that functions $f_l(s), l = 1, \dots, n$, can be continued analytically to the whole of D . Denote by \mathcal{G} the set of all $(g_1, \dots, g_n) \in H^n(D)$ such that

$$\sup_{1 \leq l \leq n} \sup_{s \in K_l} |g_l(s) - f_l(s)| < \frac{\varepsilon}{4}.$$

Let P_n and P be probability measures defined on $(S, \mathcal{B}(S))$. It is well known (see [4], Theorem 2.1) that P_n converges weakly to P as $n \rightarrow \infty$ if and only if

$$\liminf_{n \rightarrow \infty} P_n(G) \geq P(G)$$

for all open sets G .

The set \mathcal{G} is open, and, by Lemma 1, the measure P_T converges weakly to P_L as $T \rightarrow \infty$. Therefore, using the above property of the weak convergence of probability measures and Lemma 6, we obtain

$$(19) \liminf_{T \rightarrow \infty} \nu_T \left(\sup_{1 \leq l \leq n} \sup_{s \in K_l} |L(\lambda_l, \alpha_l, s + i\tau) - f_l(s)| < \frac{\varepsilon}{4} \right) = P_L(\mathcal{G}) > 0.$$

Now let the functions $f_l(s), l = 1, \dots, n$, be the same as in the statement of Theorem 1. By Lemma 7 there exist polynomials $p_l(s), l = 1, \dots, n$, such that

$$(20) \sup_{1 \leq l \leq n} \sup_{s \in K_l} |p_l(s) - f_l(s)| < \frac{\varepsilon}{2}.$$

By the first part of the proof we have that

$$(21) \liminf_{T \rightarrow \infty} \nu_T \left(\sup_{1 \leq l \leq n} \sup_{s \in K_l} |L(\lambda_l, \alpha_l, s + i\tau) - p_l(s)| < \frac{\varepsilon}{2} \right) > 0.$$

Obviously, for $l = 1, \dots, n$

$$\sup_{s \in K_l} |L(\lambda_l, \alpha_l, s + i\tau)| \leq \sup_{s \in K_l} |L(\lambda_l, \alpha_l, s + i\tau) - p_l(s)| + \sup_{s \in K_l} |f_l(s) - p_l(s)|.$$

Therefore by (20) it is easily seen that

$$\begin{aligned} & \left\{ \tau : \sup_{1 \leq l \leq n} \sup_{s \in K_l} |L(\lambda_l, \alpha_l, s + i\tau) - f_l(s)| < \varepsilon \right\} \\ & \supseteq \left\{ \tau : \sup_{1 \leq l \leq n} \sup_{s \in K_l} |L(\lambda_l, \alpha_l, s + i\tau) - p_l(s)| < \frac{\varepsilon}{2} \right\}. \end{aligned}$$

This and (21) yield the assertion of Theorem 1.

Now we discuss briefly the case that $1, \lambda_1, \dots, \lambda_n$ are linearly independent over \mathbb{Q} . Then the sequence

$$\{(\lambda_1 m, \dots, \lambda_n m), m \in \mathbb{N}\}$$

is uniformly distributed mod 1 in \mathbb{R}^n (see Kuipers-Niederreiter [17], Section 1.6, Example 6.1), hence the set

$$N_\varepsilon = \{m \in \mathbb{N} : (\lambda_1 m, \dots, \lambda_n m) \in (-\varepsilon, \varepsilon)^n \text{ mod } 1\}$$

has the positive density $(2\varepsilon)^n$. From (11) we have

$$(22) \quad \sum_{m \in N_\varepsilon} \left| \sum_{l=1}^n e^{2\pi i \lambda_l m} \varrho_l(\log m) \right| < \infty,$$

which suggests that

$$(23) \quad \sum_{m \in N_\varepsilon} \left| \sum_{l=1}^n \varrho_l(\log m) \right| < \infty$$

might be also true. If (23) would be true, then using Lemmas 4 and 5 we obtain

$$\sum_{l=1}^n \varrho_l(z) \equiv 0$$

for any $z \in \mathbb{C}$. We could prove

$$-\varrho_1(z) + \sum_{l=2}^n \varrho_l(z) \equiv 0$$

in the same way, hence $\varrho_1(z) \equiv 0$, and similarly $\varrho_l(z) \equiv 0$, $l = 2, 3, \dots, n$. From this fact we could deduce the joint universality theorem in this case. If we could prove the above conclusion $\varrho_l(z) \equiv 0$ not only from (23), but also from (22), then this argument would be complete.

5. Proof of Theorem 2

It is sufficient to give a sketch, because the proof is a direct generalization of that in [6]. Define the mapping $h : \mathbb{R} \rightarrow \mathbb{C}^{Nn}$ by

$$h(t) = (L(\lambda_1, \alpha_1, \sigma + it), \dots, L(\lambda_n, \alpha_n, \sigma + it), \\ L'(\lambda_1, \alpha_1, \sigma + it), \dots, L'(\lambda_n, \alpha_n, \sigma + it), \dots, \\ L^{(N-1)}(\lambda_1, \alpha_1, \sigma + it), \dots, L^{(N-1)}(\lambda_n, \alpha_n, \sigma + it)).$$

For any $\varepsilon > 0$ and any $s_{\nu l} \in \mathbb{C}$ ($0 \leq \nu \leq N - 1$, $1 \leq l \leq n$), we can find $\tau \in \mathbb{R}$ such that

$$|L^{(\nu)}(\lambda_l, \alpha_l, \sigma + i\tau) - s_{\nu l}| < \varepsilon \quad (0 \leq \nu \leq N - 1, 1 \leq l \leq n).$$

This can be shown by the same way as in Lemma 3 of [6], by taking the polynomial

$$p_{lN}(s) = \sum_{\nu=0}^{N-1} \frac{s_{\nu l} s^\nu}{\nu!} \quad (1 \leq l \leq n)$$

and applying Theorem 1. Hence the image of \mathbb{R} by the mapping h is dense in \mathbb{C}^{Nn} . From this, similarly to [6] (or Section 6.6 of [10]), we can deduce the conclusion of Theorem 2.

REFERENCES

- [1] M. Amou and M. Katsurada, *Differential transcendence of a class of generalized Dirichlet series*, in preparation.
- [2] B. Bagchi, *The statistical behaviour and universality properties of the Riemann zeta-function and other allied Dirichlet series*, Ph. D. Thesis, Calcutta, Indian Statistical Institute, 1981.
- [3] ———, *Joint universality theorem for Dirichlet L-functions*, Math. Zeitschrift, **181** (1982), 319–334.
- [4] P. Billingsley, *Convergence of Probability Measures*, John Wiley, New York, 1968.
- [5] R. Garunkštis, *The universality theorem with weight for the Lerch zeta-function*, In: Analytic and Probabilistic Methods in Number Theory, New Trends in Probability and Statistics, Vol.4, A. Laurinčikas et al. (Eds.), VSP/TEV, 1997, pp. 59–67.
- [6] R. Garunkštis and A. Laurinčikas, *On one Hilbert's problem for the Lerch zeta-function*, Publ. Inst. Math. (Beograd) (N.S.), **65** (79) (1999), 63–68.
- [7] S.M. Gonek, *Analytic properties of zeta and L-functions*, Ph. D. Thesis, University of Michigan, 1979.
- [8] A. Laurinčikas, *On the zeros of linear forms of Dirichlet series*, Liet. Matem. Rink., **26**(3) (1986), 468–477 (in Russian) = Lith. Math. J., **26**(3) (1986), 244–251.
- [9] ———, *Application of probability methods in the theory of Dirichlet series*, Thesis Doctor of Sciences, Vilnius, 1989 (in Russian).
- [10] ———, *Limit Theorems for the Riemann Zeta-Function*, Kluwer Academic Publishers, Dordrecht/ Boston/ London, 1996.
- [11] ———, *On limit distribution of the Lerch zeta-function*, In: Analytic and Probabilistic Methods in Number Theory, New Trends in Probability and Statistics, Vol.4, A. Laurinčikas et al.(Eds.), VSP/TEV, 1997, pp.135–148.
- [12] ———, *The universality of the Lerch zeta-function*, Liet. Matem. Rink., **37**(3) (1997), 367–375 (in Russian) = Lith. Math. J., **37**(3) (1997), 275–280.
- [13] ———, *On the Lerch zeta-function with rational parameters*, Liet. Matem. Rink., **38**(1) (1998), 113–124 (in Russian) = Lith. Math. J., **38**(1) (1998), 89–97.
- [14] ———, *On the zeros of linear combinations of the Matsumoto zeta-functions*, Liet. Matem. Rink., **38**(2) (1998), 185–204 (in Russian) = Lith. Math. J., **38**(2) (1998), 144–159.
- [15] A. Laurinčikas and K. Matsumoto, *Joint value-distribution theorems on Lerch zeta-functions*, Liet. Matem. Rink., **38**(3) (1998), 312–326 = Lith. Math. J., **38**(3) (1998), 238–249.
- [16] M. Lerch, *Note sur la fonction $K(w, x, s) = \sum_{n \geq 0} \exp\{2\pi inx\}(n+w)^{-s}$* , Acta Math., **11** (1887), 19–24.

- [17] L. Kuipers and H. Niederreiter, *Uniform Distribution of Sequences*, John Wiley & Sons, New York/London/Sydney/Toronto, 1974.
- [18] S.M. Voronin, *On the functional independence of Dirichlet L-functions*, *Acta Arith.*, **27** (1975), 493–503 (in Russian).
- [19] S.M. Voronin, *Analytic properties of generating Dirichlet functions of arithmetical objects*, Thesis Doctor phys.-matem. nauk, Moscow, 1977 (in Russian).
- [20] J.L. Walsh, *Interpolation and Approximation by Rational Functions in the Complex Domain*, Amer. Math. Soc. Coll. Publ., **20** (1960).

Antanas Laurinčikas
Department of Mathematics
Vilnius University
Naugarduko, 24
2006 Vilnius
Lithuania
`antanas.laurincikas@maf.vu.lt`

Kohji Matsumoto
Graduate School of Mathematics
Nagoya University
Chikusa-ku, Nagoya, 464-8602
Japan
`kohjimat@math.nagoya-u.ac.jp`