



This is a repository copy of *The Largest Left Quotient Ring of a Ring*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/124646/>

Version: Accepted Version

Article:

Bavula, V.V. (2016) The Largest Left Quotient Ring of a Ring. *Communications in Algebra*, 44 (8). pp. 3219-3261. ISSN 0092-7872

<https://doi.org/10.1080/00927872.2015.1082577>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

The largest strong left quotient ring of a ring

V. V. Bavula

Abstract

For an arbitrary ring R , the *largest strong left quotient ring* $Q_i^s(R)$ of R and the *strong left localization radical* \mathfrak{l}_R^s are introduced and their properties are studied in detail. In particular, it is proved that $Q_i^s(Q_i^s(R)) \simeq Q_i^s(R)$, $\mathfrak{l}_{R/\mathfrak{l}_R^s}^s = 0$ and a criterion is given for the ring $Q_i^s(R)$ to be a semisimple ring. There is a canonical homomorphism from the classical left quotient ring $Q_{l,cl}(R)$ to $Q_i^s(R)$ which is not an isomorphism, in general. The objects $Q_i^s(R)$ and \mathfrak{l}_R^s are explicitly described for several large classes of rings (semiprime left Goldie ring, left Artinian rings, rings with left Artinian left quotient ring, etc).

Key Words: the (largest) strong left quotient ring of a ring, Goldie's Theorem, the strong left localization radical, the left quotient ring of a ring, the largest left quotient ring of a ring, a maximal left denominator set, the left localization radical of a ring.

Mathematics subject classification 2010: 16S85, 16U20, 16P50, 16P60, 16P20.

Contents

1. Introduction.
2. Preliminaries, the largest strong left denominator set $T_l(R)$ of R and its characterizations.
3. The largest strong left quotient ring of a ring and its properties.
4. The largest strong quotient ring of a ring.
5. The largest strong quotient ring of $Q_i(R)$.
6. Examples.

1 Introduction

The aim of the paper is, for an arbitrary ring R , to introduce new concepts: *the largest strong left denominator set* $T_l(R)$ of R , *the largest strong left quotient ring* $Q_i^s(R) := T_l(R)^{-1}R$ of R and *the strong left localization radical* \mathfrak{l}_R^s of R , and to study their properties.

In this paper, the following notation is fixed:

- R is a ring with 1 and R^* is its group of units;
- $\mathcal{C} = \mathcal{C}_R$ is the set of *regular* elements of the ring R (i.e. \mathcal{C} is the set of non-zero-divisors of the ring R);
- ${}'\mathcal{C}_R$ is the set of *left regular* elements of the ring R , i.e. ${}'\mathcal{C}_R := \{c \in R \mid \ker(\cdot c) = 0\}$ where $\cdot c : R \rightarrow R, r \mapsto rc$;
- $Q = Q_{l,cl}(R) := \mathcal{C}^{-1}R$ is the *left quotient ring* (the *classical left ring of fractions*) of the ring R (if it exists, i.e. if \mathcal{C} is a left Ore set) and Q^* is the group of units of Q ;
- $\text{Ore}_l(R) := \{S \mid S \text{ is a left Ore set in } R\}$;
- $\text{Den}_l(R) := \{S \mid S \text{ is a left denominator set in } R\}$;
- $\text{Ass}_l(R) := \{\text{ass}(S) \mid S \in \text{Den}_l(R)\}$ where $\text{ass}(S) := \{r \in R \mid sr = 0 \text{ for some } s = s(r) \in S\}$;

- $\text{Den}_l(R, \mathfrak{a})$ is the set of left denominator sets S of R with $\text{ass}(S) = \mathfrak{a}$ where \mathfrak{a} is an ideal of R ;
- $S_{\mathfrak{a}} = S_{\mathfrak{a}}(R) = S_{l,\mathfrak{a}}(R)$ is the *largest element* of the poset $(\text{Den}_l(R, \mathfrak{a}), \subseteq)$ and $Q_{\mathfrak{a}}(R) := Q_{l,\mathfrak{a}}(R) := S_{\mathfrak{a}}^{-1}R$ is the *largest left quotient ring associated with \mathfrak{a}* . The fact that $S_{\mathfrak{a}}$ exists is proven in [3, Theorem 2.1] (but also see Lemma 2.5 below for the easy proof in other contexts);
- In particular, $S_0 = S_0(R) = S_{l,0}(R)$ is the largest element of the poset $(\text{Den}_l(R, 0), \subseteq)$, i.e. the *largest regular left Ore set* of R , and $Q_l(R) := S_0^{-1}R$ is the *largest left quotient ring* of R [3];
- $\text{max.Den}_l(R)$ is the set of maximal left denominator sets of R (it is always a *non-empty* set, see [3], or Lemma 2.5 below for the proof).

The largest strong left quotient ring of a ring. Consider the following subsets of a ring R : The sets

$$\begin{aligned} \mathcal{L}_l^s(R) &:= \bigcap_{S \in \text{max.Den}_l(R)} S \stackrel{\text{Prop. 2.3.(1)}}{=} \{c \in R \mid \frac{c}{1} \in (S^{-1}R)^* \text{ for all } S \in \text{max.Den}_l(R)\}, \\ \mathcal{C}_R^w &:= \{c \in R \mid \frac{c}{1} \in \mathcal{C}_{S^{-1}R} \text{ for all } S \in \text{max.Den}_l(R)\}, \\ {}'\mathcal{C}_R^w &:= \{c \in R \mid \frac{c}{1} \in {}'\mathcal{C}_{S^{-1}R} \text{ for all } S \in \text{max.Den}_l(R)\}, \end{aligned}$$

are called respectively *the set of strongly left localizable elements*, *the set of weak regular elements* and *the set of weak left regular elements* of R .

- (Proposition 2.8 and Proposition 2.15) *Each of the sets $\mathcal{L}_l^s(R)$, \mathcal{C}_R^w and ${}'\mathcal{C}_R^w$ contains a unique largest left denominator set, and all three largest left denominator sets coincide and are denoted by $T_l(R)$.*

The set $T_l(R)$ is called the *largest strong left denominator set* of R and the ring $Q_l^s(R) := T_l(R)^{-1}R$ is called the *largest strong left quotient ring* of R . The ideal of R given by $\mathfrak{l}_R^s := \text{ass}(T_l(R)) := \{r \in R \mid tr = 0 \text{ for some } t \in T_l(R)\}$ is called the *strong left localization radical* of R . In the above definitions, the adjective ‘strong’ reflects their connections with the set $\mathcal{L}_l^s(R)$ of strongly left localizable elements of R . The set $T_l(R)$ is the largest left denominator set of R that consists of elements that are invertible in *all* maximal left localizations of the ring R .

In general, for a ring R , its left (right; two sided) localizations, especially maximal ones, are unrelated. The intuition behind the construction of the largest strong left quotient ring of R is to have the *largest* possible left localization of R that is *related* to all maximal left localizations of the ring R , i.e. there exists a ring R -homomorphism (necessarily, unique) from $Q_l^s(R)$ to $S^{-1}R$ for every $S \in \text{max.Den}_l(R)$.

In Section 5, the triple $T_l(R)$, \mathfrak{l}_R^s , $Q_l^s(R)$ is found explicitly for the following four classes of rings: semiprime left Goldie rings (Theorem 5.1); rings of $n \times n$ lower/upper triangular matrices with coefficients in a left Goldie domain (Theorem 5.2 and Theorem 5.3); left Artinian rings (Theorem 5.6), and rings with left Artinian left quotient ring (Theorem 5.7). In particular, for semiprime left Goldie rings R : $T_l(R) = \mathcal{C}_R$, $\mathfrak{l}_R^s = 0$ and $Q_l^s(R) = Q_{l,cl}(R)$ (Theorem 5.1). In general, none of the three equalities holds for the remaining three (just mentioned) classes of rings but the results are natural and beautiful (very symmetrical), eg, for a ring A such that $Q_{l,cl}(R)$ is a left Artinian ring (Theorem 5.7):

$$T_l(A) = \bigcap_{S' \in \text{max.Den}_l(A)} S' \text{ and } Q_l^s(A) \simeq \prod_{S' \in \text{max.Den}_l(A)} S'^{-1}A.$$

In particular, the ring $Q_l^s(A)$ has a more regular/simple structure than the ring $Q_{l,cl}(A)$.

It would be interesting to find the ring $Q_l^s(R)$ for other classes of rings. Theorem 1.6.(3), which states that $Q_l^s(R) \simeq Q_l^s(Q_l(R))$ for an arbitrary ring R , opens a way for tackling more challenging types of rings.

The main results of the paper are the following six theorems. The first one describes $T_l(R)$, $Q_l^s(R)$ and $Q_l^s(R)^*$.

Theorem 1.1 *Let R be a ring, $\pi : R \rightarrow R/\mathfrak{I}_R^s$, $r \mapsto \bar{r} = r + \mathfrak{I}_R^s$; $\sigma : R \rightarrow Q_l^s(R)$, $r \mapsto \frac{r}{1}$, and $Q_l^s(R)^*$ be the group of units of the ring $Q_l^s(R)$. Then*

1. $T_l(R) = S_{l, \mathfrak{I}_R^s}(R)$.
2. $Q_l^s(R) = Q_{l, \mathfrak{I}_R^s}(R) \simeq Q_l(R/\mathfrak{I}_R^s)$.
3. $T_l(R) = \sigma^{-1}(Q_l^s(R)^*)$.
4. $T_l(R) = \pi^{-1}(S_{l, 0}(R/\mathfrak{I}_R^s))$.
5. $Q_l^s(R)^* = \{s^{-1}t \mid s, t \in T_l(R)\}$.

The second one describes the objects $Q_l^s(Q_l^s(R))$, $\mathfrak{I}_{R/\mathfrak{I}_R^s}^s$, $T_l(R/\mathfrak{I}_R^s)$, $Q_l^s(R/\mathfrak{I}_R^s)$ and their connections with their counterparts for the ring R .

Theorem 1.2 *We keep the notation of Theorem 1.1. Then*

1. $Q_l^s(Q_l^s(R)) = Q_l^s(R)$.
2. $T_l(R/\mathfrak{I}_R^s) = \pi(T_l(R))$ and $T_l(R) = \pi^{-1}(T_l(R/\mathfrak{I}_R^s))$.
3. $T_l(R/\mathfrak{I}_R^s) = S_{l, 0}(R/\mathfrak{I}_R^s)$.
4. $\mathfrak{I}_{R/\mathfrak{I}_R^s}^s = 0$.
5. $T_l(Q_l^s(R)) = Q_l^s(R)^*$ and $\mathfrak{I}_{Q_l^s(R)}^s = 0$.
6. $\pi(\mathcal{L}_l^s(R)) = \mathcal{L}_l^s(R/\mathfrak{I}_R^s)$ and $\mathcal{L}_l^s(R) = \pi^{-1}(\mathcal{L}_l^s(R/\mathfrak{I}_R^s))$.
7. $Q_l^s(R/\mathfrak{I}_R^s) = Q_l(R/\mathfrak{I}_R^s)$.

Semisimplicity criterion for the ring $Q_l^s(R)$. A ring is called a *left Goldie ring* if it does not contain infinite direct sums of nonzero left ideals and satisfies the ascending chain condition on left annihilators.

Theorem 1.3 *Let R be a ring. The following statements are equivalent.*

1. $Q_l^s(R)$ is a semisimple ring.
2. R/\mathfrak{I}_R^s is a semiprime left Goldie ring.
3. $Q_l(R/\mathfrak{I}_R^s)$ is a semisimple ring.
4. $Q_{l, cl}(R/\mathfrak{I}_R^s)$ is a semisimple ring.

If one of the equivalent conditions holds then

$$Q_l^s(R) \simeq Q_l(R/\mathfrak{I}_R^s) \simeq Q_{l, cl}(R/\mathfrak{I}_R^s),$$

$T_l(R) = \pi^{-1}(\mathcal{C}_{R/\mathfrak{I}_R^s})$ and $T_l(R/\mathfrak{I}_R^s) = \mathcal{C}_{R/\mathfrak{I}_R^s}$ where $\pi : R \rightarrow R/\mathfrak{I}_R^s$, $r \mapsto \bar{r} = r + \mathfrak{I}_R^s$, and $\mathcal{C}_{R/\mathfrak{I}_R^s}$ is the set of regular elements of the ring R/\mathfrak{I}_R^s .

Goldie's Theorem [7] is a criterion for a ring to have semisimple left quotient ring (earlier, criteria were given, by Goldie [6] and Lesieur and Croisot [10], for a ring to have a simple Artinian left quotient ring). Recently, the author [4] has given several more new criteria. For a left Noetherian ring which has a left quotient ring, Talintyre [17] has established necessary and sufficient conditions for the left quotient ring to be left Artinian. Small [13, 14], Robson [12], and later Tachikawa [16] and Hajarnavis [9], and recently the author [2] have given different criteria for a ring to have a left Artinian left quotient ring.

Semisimplicity criterion for the ring $Q_{l,cl}(R)$. The statement of Goldie's Theorem is a semisimplicity criterion for the ring $Q_{l,cl}(R)$ which states that the ring $Q_{l,cl}(R)$ is a semisimple ring iff R is a semiprime left Goldie ring. Recently, four new criteria for semisimplicity of $Q_{l,cl}(R)$ are given in [4] using completely different ideas and approach. Below, another semisimplicity criterion for $Q_{l,cl}(R)$ is given via $Q_l^s(R)$ and \mathfrak{I}_R^s .

Theorem 1.4 *Let R be a ring. The following statements are equivalent.*

1. $Q_{l,cl}(R)$ is a semisimple ring.
2. $Q_l(R)$ is a semisimple ring.
3. (a) $Q_l^s(R)$ is a semisimple ring.
(b) $\mathfrak{I}_R^s = 0$.

If one of the equivalent conditions 1–3 holds then $Q_{l,cl}(R) \simeq Q_l(R) \simeq Q_l^s(R)$ and $\mathcal{C}_R = T_l(R) = \mathcal{L}_l^s(R)$.

Theorem 1.5 *Let R be a ring. Then, for all $\mathfrak{a} \in \text{Ass}_l(R)$ with $\mathfrak{a} \subseteq \mathfrak{I}_R^s$, $S_{l,\mathfrak{a}}(R) \subseteq T_l(R)$, and so there is a ring R -homomorphism $Q_{l,\mathfrak{a}}(R) \rightarrow Q_l^s(R)$, $s^{-1}r \mapsto s^{-1}r$.*

For an arbitrary ring R , Theorem 1.6 reveals natural and tight connections between triples $T_l(R)$, \mathfrak{I}_R^s , $Q_l^s(R)$ and $T_l(Q_l(R))$, $\mathfrak{I}_{Q_l(R)}^s$, $Q_l^s(Q_l(R))$.

Theorem 1.6 *Let R be a ring. Then*

1. $T_l(Q_l(R)) = Q_l(R)^* T_l(R) = \{s^{-1}t \mid s \in S_l(R), t \in T_l(R)\}$ and $T_l(R) = R \cap T_l(Q_l(R))$.
2. $\mathfrak{I}_{Q_l(R)}^s = S_l(R)^{-1} \mathfrak{I}_R^s$ and $\mathfrak{I}_R^s = R \cap \mathfrak{I}_{Q_l(R)}^s$.
3. $Q_l^s(R) \simeq Q_l^s(Q_l(R))$.

The paper is organized as follows. In Section 2, we prove Proposition 2.8 and Proposition 2.15 (mentioned above). We show that that $S_{l,0}(R) \subseteq T_l(R)$ (Lemma 2.10.(3)) and as a result there is a canonical homomorphism

$$\theta : Q_l(R) \rightarrow Q_l^s(R), \quad s^{-1}r \mapsto s^{-1}r, \quad (s \in S_l(R), r \in R).$$

The lemma below is a criterion for the homomorphism θ to be an isomorphism.

- (Lemma 2.13) $S_{l,0}(R) = T_l(R)$ iff θ is an isomorphism iff $\mathfrak{I}_R^s = 0$.

In Section 3, proofs of Theorems 1.1–1.5 are given. In Section 4, the two-sided theory (i.e. about left and right denominators sets) is developed and analogous results to the five theorems above are proved. In Section 3, Theorem 1.6 is proved.

2 Preliminaries, the largest strong left denominator set $T_l(R)$ of R and its characterizations

In this section, for reader's convenience we collect necessary results that are used in the proofs of this paper. Several characterizations (Proposition 2.8 and Proposition 2.15) of $T_l(R)$ are given. A criterion is given for the inclusion $S_{l,0}(R) \subseteq T_l(R)$ (which always holds by Lemma 2.10.(3)) to be an equality, and for the canonical ring homomorphism $Q_{l,cl}(R) \rightarrow Q_l^s(R)$ to be an isomorphism (Lemma 2.13).

More results on localizations of rings (and some of the missed standard definitions) the reader can find in [8], [11] and [15].

The largest regular left Ore set and the largest left quotient ring of a ring. Let R be a ring. A *multiplicatively closed subset* S of R or a *multiplicative subset* of R (i.e. a multiplicative sub-semigroup of (R, \cdot) such that $1 \in S$ and $0 \notin S$) is said to be a *left Ore set* if it satisfies the *left Ore condition*: for each $r \in R$ and $s \in S$, $Sr \cap Rs \neq \emptyset$. Let $\text{Ore}_l(R)$ be the set of all left Ore sets of R . For each $S \in \text{Ore}_l(R)$ the set $\text{ass}(S) := \{r \in R \mid sr = 0 \text{ for some } s \in S\}$ is an ideal of the ring R .

A left Ore set S is called a *left denominator set* of the ring R if $rs = 0$ for some elements $r \in R$ and $s \in S$ implies $tr = 0$ for some element $t \in S$, i.e. $r \in \text{ass}(S)$. Let $\text{Den}_l(R)$ be the set of all left denominator sets of R . For $S \in \text{Den}_l(R)$, let $S^{-1}R = \{s^{-1}r \mid s \in S, r \in R\}$ be the *left localization* of the ring R at S (the *left quotient ring* of R at S). Let us stress that in Ore's method of localization one can localize *precisely* at left denominator sets.

In general, the set \mathcal{C} of regular elements of a ring R is neither a left nor right Ore set of the ring R and as a result neither the left nor right classical quotient ring ($Q_{l,cl}(R) := \mathcal{C}^{-1}R$ and $Q_{r,cl}(R) := R\mathcal{C}^{-1}$) exists. Remarkably, there exists a largest regular left Ore set $S_0 = S_{l,0} = S_{l,0}(R)$, [3]. This means that the set $S_{l,0}(R)$ is an Ore set of the ring R that consists of regular elements (i.e. $S_{l,0}(R) \subseteq \mathcal{C}$) and contains all the left Ore sets in R that consist of regular elements. Also, there exists a largest regular (left and right) Ore set $S_{l,r,0}(R)$ of any ring R . In general, all the sets \mathcal{C} , $S_{l,0}(R)$, $S_{r,0}(R)$ and $S_{l,r,0}(R)$ are distinct. For example, these sets are different for the ring $\mathbb{I}_1 = K\langle x, \partial, f \rangle$ of polynomial integro-differential operators over a field K of characteristic zero, [1]. In [1], these four sets are found explicitly for $R = \mathbb{I}_1$.

Definition: Following the terminology of [1], [3], we call the ring

$$Q_l(R) := S_{l,0}(R)^{-1}R$$

(respectively, $Q_r(R) := RS_{r,0}(R)^{-1}$ and $Q(R) := S_{l,r,0}(R)^{-1}R \simeq RS_{l,r,0}(R)^{-1}$) the *largest left* (respectively, *right and two-sided*) *quotient ring* of the ring R .

In general, the rings $Q_l(R)$, $Q_r(R)$ and $Q(R)$ are not isomorphic, for example, for $R = \mathbb{I}_1$ as shown in Section 8 of [1]. The next theorem gives various properties of the ring $Q_l(R)$. In particular, it describes its group of units.

Theorem 2.1 [3]

1. $S_0(Q_l(R)) = Q_l(R)^*$ and $S_0(Q_l(R)) \cap R = S_0(R)$.
2. $Q_l(R)^* = \langle S_0(R), S_0(R)^{-1} \rangle$, i.e. the group of units of the ring $Q_l(R)$ is generated by the sets $S_0(R)$ and $S_0(R)^{-1} := \{s^{-1} \mid s \in S_0(R)\}$.
3. $Q_l(R)^* = \{s^{-1}t \mid s, t \in S_0(R)\}$.
4. $Q_l(Q_l(R)) = Q_l(R)$.

The maximal left denominator sets and the maximal left localizations of a ring. The set $(\text{Den}_l(R), \subseteq)$ is a poset (partially ordered set). In [3], it is proved that the set $\text{max.Den}_l(R)$ of its maximal elements is a *non-empty* set.

Definition, [3]. An element S of the set $\text{max.Den}_l(R)$ is called a *maximal left denominator set* of the ring R and the ring $S^{-1}R$ is called a *maximal left quotient ring* of the ring R or a *maximal left localization ring* of the ring R . The intersection

$$\mathfrak{l}_R := \text{l.rad}(R) := \bigcap_{S \in \text{max.Den}_l(R)} \text{ass}(S) \quad (1)$$

is called the *left localization radical* of the ring R , [3].

Properties of the maximal left quotient rings of a ring. The next theorem describes various properties of the maximal left quotient rings of a ring. In particular, their groups of units and their largest left quotient rings. It is the key fact in the proof of the characterization of the set \mathcal{L}_l^s (Proposition 2.3).

Theorem 2.2 [3] *Let $S \in \text{max.Den}_l(R)$, $A = S^{-1}R$, A^* be the group of units of the ring A ; $\mathfrak{a} := \text{ass}(S)$, $\pi_{\mathfrak{a}} : R \rightarrow R/\mathfrak{a}$, $a \mapsto a + \mathfrak{a}$, and $\sigma_{\mathfrak{a}} : R \rightarrow A$, $r \mapsto \frac{r}{1}$. Then*

1. $S = S_{\mathfrak{a}}(R)$, $S = \pi_{\mathfrak{a}}^{-1}(S_0(R/\mathfrak{a}))$, $\pi_{\mathfrak{a}}(S) = S_0(R/\mathfrak{a})$ and $A = S_0(R/\mathfrak{a})^{-1}R/\mathfrak{a} = Q_l(R/\mathfrak{a})$.
2. $S_0(A) = A^*$ and $S_0(A) \cap (R/\mathfrak{a}) = S_0(R/\mathfrak{a})$.
3. $S = \sigma_{\mathfrak{a}}^{-1}(A^*)$.
4. $A^* = \langle \pi_{\mathfrak{a}}(S), \pi_{\mathfrak{a}}(S)^{-1} \rangle$, i.e. the group of units of the ring A is generated by the sets $\pi_{\mathfrak{a}}(S)$ and $\pi_{\mathfrak{a}}(S)^{-1} := \{\pi_{\mathfrak{a}}(s)^{-1} \mid s \in S\}$.
5. $A^* = \{\pi_{\mathfrak{a}}(s)^{-1}\pi_{\mathfrak{a}}(t) \mid s, t \in S\}$.
6. $Q_l(A) = A$ and $\text{Ass}_l(A) = \{0\}$. In particular, if $T \in \text{Den}_l(A)$ then $T \subseteq A^*$.

Theorems 2.1 and 2.2 are used in many proofs in this paper.

Definition, [4]. The sets

$$\mathcal{L}_l(R) := \bigcup_{S \in \text{max.Den}_l(R)} S \text{ and } \mathcal{NL}_l(R) := R \setminus \mathcal{L}_l(R)$$

are called the sets of *left localizable* and *left non-localizable elements* of R , respectively, and the intersection

$$\mathcal{L}_l^s(R) := \bigcap_{S \in \text{max.Den}_l(R)} S$$

is called the *set of strongly (or completely) left localizable elements* of R . Clearly, $\mathcal{L}_l^s(R)$ is a multiplicative set and

$$R^* \subseteq \mathcal{L}_l^s(R) \quad (2)$$

since $R^* \subseteq S$ for all $S \in \text{max.Den}_l(R)$, by Theorem 2.2.(3). By Proposition 3.3.(1),

$$S_{l,0}(R) \subseteq \mathcal{L}_l^s(R). \quad (3)$$

In particular, if the set \mathcal{C}_R of regular elements of the ring R is a left Ore set then $\mathcal{C}_R = S_{l,0}(R)$ and so

$$\mathcal{C}_R \subseteq \mathcal{L}_l^s(R). \quad (4)$$

The next proposition is a characterization of the set $\mathcal{L}_l^s(R)$ which says that the set $\mathcal{L}_l^s(R)$ contains *precisely* the elements of the ring R that are units in all maximal left localizations of R .

Proposition 2.3 *Let R be a ring. Then*

1. $\mathcal{L}_l^s(R) = \{s \in R \mid \frac{s}{1} \in (S^{-1}R)^* \text{ for all } S \in \max.\text{Den}_l(R)\}$ where $(S^{-1}R)^*$ is the group of units of the ring $S^{-1}R$.
2. For all automorphisms $\sigma \in \text{Aut}(R)$, $\sigma(\mathcal{L}_l^s(R)) = \mathcal{L}_l^s(R)$.

Proof. 1. Let \mathcal{R} be the RHS of the claimed equality. By the very definition of the set $\mathcal{L}_l^s(R)$, we have the inclusion $\mathcal{L}_l^s(R) \subseteq \mathcal{R}$. Conversely, let $s \in \mathcal{R}$ and $\sigma_S : R \rightarrow S^{-1}R$, $r \mapsto \frac{r}{1}$, where $S \in \max.\text{Den}_l(R)$. Then $s \in \sigma_S^{-1}((S^{-1}R)^*) = S$ for all $S \in \max.\text{Den}_l(R)$ (Theorem 2.2 parts 2 and 3), hence $s \in \mathcal{L}_l^s(R)$.

2. Obvious. \square

Let R be a ring. Let S, T be submonoids of the multiplicative monoid (R, \cdot) . We denote by ST the *submonoid* of (R, \cdot) generated by S and T . This notation should not be confused with the product of two sets which is *not* used in this paper. The next result is a criterion for the set ST to be a left Ore (denominator) set.

Lemma 2.4 1. *Let $S, T \in \text{Ore}_l(R)$. If $0 \notin ST$ then $ST \in \text{Ore}_l(R)$.*

2. *Let $S, T \in \text{Den}_l(R)$. If $0 \notin ST$ then $ST \in \text{Den}_l(R)$.*

3. *Statements 1 and 2 hold also for Ore sets and denominator sets, respectively.*

Proof. 1. Since $0 \notin ST$, the set $P := ST$ is multiplicative. It remains to show that the left Ore condition holds for P . Given an element $p = s_1 t_1 \cdots s_n t_n \in P$ and $r \in R$ (where $s_i \in S$ and $t_i \in T$) we have to find elements $p' \in P$ and $r' \in R$ such that $p'r = r'p$. There are elements $t'_n \in T$ and $r'_n \in R$ such that $t'_n r = r'_n t_n$. Similarly, $s'_n r'_n = r''_n s_n$ for some $s'_n \in S$ and $r''_n \in R$. Hence, $s'_n t'_n r = r''_n s_n t_n$. Then repeating these two steps $n - 1$ more times we find elements $s'_i \in S$, $t'_i \in T$ and $r' \in R$ such that

$$s'_1 t'_1 \cdots s'_n t'_n r = r' s_1 t_1 \cdots s_n t_n.$$

So, it suffices to take $p' = s'_1 t'_1 \cdots s'_n t'_n$.

2. By statement 1, it remains to show that if $rp = 0$ for some elements $r \in R$ and $p = s_1 t_1 \cdots s_n t_n \in P$ then $p'r = 0$ for some $p' \in P$. $0 = rp = (rs_1 t_1 \cdots s_n) t_n \Rightarrow t'_n r s_1 t_1 \cdots s_n = 0$ for some element $t'_n \in T$. Similarly, $s'_n t'_n r s_1 t_1 \cdots s_{n-1} t_{n-1} = 0$ for some element $s'_n \in S$. Repeating the same two steps $n - 1$ more times we have $s'_1 t'_1 \cdots s'_n t'_n r = 0$ for some elements $s'_i \in S$ and $t'_i \in T$. It suffices to take $p' = s'_1 t'_1 \cdots s'_n t'_n$.

3. Statement 3 follows from statements 1 and 2. \square

Criterion for a left Ore/denominator set to be maximal. There are posets $(\text{Ore}_l(R), \subseteq)$, $(\text{Den}_l(R), \subseteq)$, $(\text{Ore}(R), \subseteq)$ and $(\text{Den}(R), \subseteq)$. The next lemma states that the sets of maximal elements of these posets are non-empty sets.

Lemma 2.5 *Let R be a ring.*

1. *The set $\max.\text{Ore}_l(R)$ of maximal left Ore sets in R is a non-empty set.*
2. *The set $\max.\text{Den}_l(R)$ of maximal left denominator sets in R is a non-empty set.*
3. *The set $\max.\text{Ore}(R)$ of maximal (left and right) Ore sets in R is a non-empty set.*
4. *The set $\max.\text{Den}(R)$ of maximal (left and right) denominator sets in R is a non-empty set.*

Proof. All statements follow at once from Zorn's Lemma and the fact that given a linearly ordered chain of left (resp. left and right) Ore sets [resp. denominator sets] then their union is a left (resp. left and right) Ore set [resp. a denominator set]. \square

The next proposition is a criterion for a left Ore/denominator set to be a *maximal* left Ore/denominator set.

Proposition 2.6 *Let R be a ring.*

1. *Let $S \in \text{Ore}_l(R)$ (resp. $S \in \text{Ore}(R)$). Then $S \in \text{max.Ore}_l(R)$ (resp. $S \in \text{max.Ore}(R)$) iff $0 \in ST$ for all $T \in \text{Ore}_l(R)$ such that $T \not\subseteq S$ (resp. $T \in \text{Ore}(R)$).*
2. *Let $S \in \text{Den}_l(R)$ (resp. $S \in \text{Den}(R)$). Then $S \in \text{max.Den}_l(R)$ (resp. $S \in \text{max.Den}(R)$) iff $0 \in ST$ for all $T \in \text{Den}_l(R)$ such that $T \not\subseteq S$ (resp. $T \in \text{Den}(R)$).*

Proof. 1. Statement 1 follows from Lemma 2.4.(1,3) and the inclusion $S \subseteq ST$.
2. Statement 1 follows from Lemma 2.4.(2,3) and the inclusion $S \subseteq ST$. \square

Let $\{S_i \mid i \in I\} \subseteq \text{Ore}_l(R)$, $I \neq \emptyset$, $\mathcal{F} := \{J \subseteq I \mid 1 \leq |J| < \infty\}$ and

$$\bigvee_{i \in I} S_i := \bigcup_{F \in \mathcal{F}} \prod_{i \in F} S_i. \quad (5)$$

- Lemma 2.7**
1. *Let $\{S_i \mid i \in I\} \subseteq \text{Ore}_l(R)$. Suppose that $0 \notin \prod_{i \in F} S_i$ for all non-empty finite subsets $F \subseteq I$. Then $\bigvee_{i \in I} S_i$ is the least upper bound of $\{S_i \mid i \in I\}$ in $\text{Ore}_l(R)$.*
 2. *Let $\{S_i \mid i \in I\} \subseteq \text{Den}_l(R)$. Suppose that $0 \notin \prod_{i \in F} S_i$ for all non-empty finite subsets $F \subseteq I$. Then $\bigvee_{i \in I} S_i$ is the least upper bound of $\{S_i \mid i \in I\}$ in $\text{Den}_l(R)$.*
 3. *Statements 1 and 2 hold also for Ore sets and denominator sets, respectively.*

Proof. 1. By Lemma 2.4.(1), $\bigvee_{i \in I} S_i \in \text{Ore}_l(R)$. Now, statement 1 is obvious.
2. By Lemma 2.4.(2), $\bigvee_{i \in I} S_i \in \text{Den}_l(R)$. Now, statement 2 is obvious.
3. Statement 3 follows from statements 1 and 2. \square

The largest strong left denominator set $T_l(R)$. The set $\text{Den}_l^s(R) := \{T \in \text{Den}_l(R) \mid T \subseteq \mathcal{L}_l^s(R)\}$ is a non-empty set since $R^* \in \text{Den}_l^s(R)$. The elements of $\text{Den}_l^s(R)$ are called the *strong left denominator sets* of R and the rings $T^{-1}R$ where $T \in \text{Den}_l^s(R)$ are called the *strong left quotient rings* or the *strong left localizations* of R . The next proposition shows that the set of maximal elements (w.r.t. to inclusion) $\text{max.Den}_l^s(R)$ is a non-empty set. Moreover, it contains a single element. Namely,

$$T_l(R) := \bigcup_{S \in \text{max.Den}_l^s(R)} S. \quad (6)$$

Proposition 2.8 *Let R be a ring. Then $\text{max.Den}_l^s(R) = \{T_l(R)\}$.*

Proof. (i) For all $S, T \in \text{Den}_l^s(R)$, $0 \notin ST$ where ST is the semigroup of (R, \cdot) generated by the sets S and T : Suppose that $0 \in ST$ for some $S, T \in \text{Den}_l^s(R)$, i.e. $s_1 t_1 s_2 t_2 \cdots s_n t_n = 0$ for some elements $s_i \in S$ and $t_i \in T$. Take $\mathcal{S} \in \text{max.Den}_l(R)$. Then $S, T \subseteq \mathcal{S}$ and so $0 \neq s_1 t_1 s_2 t_2 \cdots s_n t_n / 1 \in \mathcal{S}^{-1}R$, a contradiction.

(ii) $\bigvee_{S \in \text{max.Den}_l^s(R)} S = T_l(R)$ (see (5)): This follows from the fact that $\text{Den}_l^s(R)$ is a monoid, by (i).

(iii) $\text{max.Den}_l^s(R) = \{T_l(R)\}$: By (ii) and Lemma 2.7.(2), $T_l(R)$ is the least upper bound of $\text{Den}_l^s(R)$ in the set $\text{Den}_l(R)$. Since $T_l(R) \in \text{Den}_l^s(R)$, $T_l(R)$ is the largest element in $\text{Den}_l^s(R)$. \square

So, $T_l(R)$ is the *largest strong denominator set* of R .

Definition. The ideal of the ring R ,

$$\mathfrak{I}_R^s := \text{ass}(T_l(R)) = \bigcup_{S \in \text{max.Den}_l^s(R)} \text{ass}(S)$$

is called the *strong left localization radical* of the ring R .

The set $\text{Den}_l^s(R)$ is invariant under the action of the group of automorphisms $\text{Aut}(R)$ of the ring R .

Lemma 2.9 1. For all automorphisms $\sigma \in \text{Aut}(R)$, $\sigma(T_l(R)) = T_l(R)$ and $\sigma(\mathfrak{l}_R^s) = \mathfrak{l}_R^s$.

2. $\mathfrak{l}_R^s \subseteq \mathfrak{l}_R$ where $\mathfrak{l}_R := \bigcap_{S \in \max.\text{Den}_l(R)} \text{ass}(S)$.

Proof. 1. Obvious.

2. For all $S \in \max.\text{Den}_l(R)$, $T_l(R) \subseteq S$, and so $\text{ass}(T_l(R)) \subseteq \text{ass}(S)$. Therefore, $\mathfrak{l}_R^s \subseteq \mathfrak{l}_R$. \square

The next result shows that the sets $T_l(R)$ and $\mathcal{L}_l^s(R)$ are closed under addition of elements of \mathfrak{l}_R^s and \mathfrak{l}_R , respectively.

Lemma 2.10 Let R be a ring. Then

1. $T_l(R) + \mathfrak{l}_R^s = T_l(R)$.

2. $\mathcal{L}_l^s(R) + \mathfrak{l}_R \subseteq \mathcal{L}_l^s(R)$. In particular, $T_l(R) + \mathfrak{l}_R \subseteq \mathcal{L}_l^s(R)$.

3. $R^* \subseteq S_{l,0}(R) \subseteq T_l(R)$.

Proof. 1. Let $T_l := T_l(R)$. By Corollary 2.12.(1) (or Lemma 2.11) (see below), $T := T_l + \mathfrak{l}_R^s \in \text{Den}_l(R)$. To finish the proof of statement 1 it suffices to show that $T \subseteq \mathcal{L}_l^s(R)$ (since then $T \subseteq T_l$, as T_l is the largest element in $\mathcal{L}_l^s(R)$, Proposition 2.8). Let $t \in T_l$ and $a \in \mathfrak{l}_R^s$. We have to show that $t + a \in \mathcal{L}_l^s(R)$. For all $S \in \max.\text{Den}_l(R)$, $T_l \subseteq S$ and so $\mathfrak{l}_R^s \subseteq \text{ass}(S)$. Hence,

$$S^{-1}R \ni \frac{t+a}{1} = \frac{t}{1} \in (S^{-1}R)^*.$$

By Proposition 2.3.(1), $t + a \in \mathcal{L}_l^s(R)$.

2. Let $t \in \mathcal{L}_l^s(R)$ and $l \in \mathfrak{l}_R$. For all $S \in \max.\text{Den}_l(R)$, $S^{-1}R \ni \frac{t+l}{1} = \frac{t}{1} \in (S^{-1}R)^*$. By Proposition 2.3, $t + l \in \mathcal{L}_l^s(R)$.

3. By the inclusion (3), $S_{l,0}(R) \in \text{Den}_l^s(R)$. Hence, $S_{l,0}(R) \subseteq T_l(R)$, by Proposition 2.8. By Theorem 2.1.(1), $R^* \subseteq S_{l,0}(R)$. \square

The next lemma shows that under natural conditions pre-images of left denominator sets are also left denominator sets.

Lemma 2.11 Let R be a ring, $S \in \text{Den}_l(R, \mathfrak{a})$, $\pi : R \rightarrow R/\mathfrak{a}$, $r \mapsto \bar{r} := r + \mathfrak{a}$, $T \in \text{Den}_l(R/\mathfrak{a}, \mathfrak{b}/\mathfrak{a})$ where \mathfrak{b} is an ideal of R such $\mathfrak{a} \subseteq \mathfrak{b}$, and $T' := \pi^{-1}(T)$. If $S \subseteq T'$ then $T' \in \text{Den}_l(R, \mathfrak{b})$.

Proof. (i) $T' \in \text{Ore}_l(R)$: Given elements $t' \in T'$ and $r \in R$. Since $T \in \text{Ore}_l(R/\mathfrak{a})$, $\bar{t}\bar{r} = \bar{r}_1\bar{t}$ for some elements $t \in T'$ and $r_1 \in R$. Then $s(tr - r_1t') = 0$ for some element $s \in S$, and so $st \cdot r = sr_1 \cdot t'$ where $st \in T'$, i.e. $T' \in \text{Ore}_l(R)$.

(ii) $T' \in \text{Den}_l(R)$: If $rt' = 0$ for some elements $r \in R$ and $t' \in T'$ then $\bar{r}\bar{t}' = 0$, and so $\bar{t}'_1\bar{r} = 0$ for some element $t'_1 \in T'$. Now, $t'_1r \in \mathfrak{a}$. There exists an element $s \in S$ such that $st'_1 \cdot r = 0$. Notice that $st'_1 \in T'$ (since $s \in S \subseteq T'$). Therefore, $T' \in \text{Den}_l(R)$, by (i).

(iii) $\text{ass}(T') = \mathfrak{b}$: If $t'r = 0$ for some elements $t' \in T'$ and $r \in R$ then $\bar{t}'\bar{r} = 0$, and so $\bar{r} \in \mathfrak{b}/\mathfrak{a}$ and $r \in \mathfrak{b}$. Then $\text{ass}(T') \subseteq \mathfrak{b}$. Let $b \in \mathfrak{b}$. Then $\bar{t}'\bar{b} = 0$ for some element $t' \in T'$ (since $\pi(T') = T \in \text{Den}_l(R/\mathfrak{a}, \mathfrak{b}/\mathfrak{a})$). Now, $t'b \in \mathfrak{a}$. Hence, $st' \cdot b = 0$ for some $s \in S$. Since $st' \in T'$, $b \in \text{ass}(T')$. Then, $\text{ass}(T') = \mathfrak{b}$. \square

Corollary 2.12 Let R be a ring and $S \in \text{Den}_l(R, \mathfrak{a})$. Then

1. $S + \mathfrak{a} \in \text{Den}_l(R, \mathfrak{a})$.

2. $S + \text{ass}(S) = S$ for all $S \in \max.\text{Den}_l(R)$.

3. $S + \text{ass}(S) = S$ for all $S \in \max.\text{Den}(R)$.

Proof. 1. Let $\pi : R \rightarrow R/\mathfrak{a}$, $r \mapsto \bar{r} := r + \mathfrak{a}$. Since $T := \pi(S) \in \text{Den}_l(R/\mathfrak{a}, 0)$ and $S + \mathfrak{a} = \pi^{-1}(T) \supseteq S$, we see that $S + \mathfrak{a} \in \text{Den}_l(R, \mathfrak{a})$, by Lemma 2.11.

2 and 3. Statements 2 and 3 follow from statement 1 and the inclusion $S \subseteq S + \text{ass}(S)$. \square

The largest strong left quotient ring $Q_l^s(R)$ of a ring R .

Definition. The ring $Q_l^s(R) := T_l(R)^{-1}R$ is called the *largest strong left quotient ring* of the ring R .

There are exact sequences:

$$0 \rightarrow \mathfrak{I}_R^s \rightarrow R \rightarrow Q_l^s(R), \quad r \mapsto \frac{r}{1}, \quad (7)$$

$$0 \rightarrow S_{l,0}(R)^{-1}\mathfrak{I}_R^s \rightarrow Q_l(R) \xrightarrow{\theta} Q_l^s(R), \quad \theta(s^{-1}r) = s^{-1}r, \quad (8)$$

where $s \in S_{l,0}(R)$ and $r \in R$ (if $\theta(s^{-1}r) = 0$ then $\frac{r}{1} = 0$ in $Q_l^s(R)$, hence $r \in \mathfrak{I}_R^s$, and so $\ker(\theta) = S_{l,0}^{-1}(R)^{-1}\mathfrak{I}_R^s$).

For each $S \in \text{max.Den}_l(R)$, there is a commutative diagram of ring homomorphisms:

$$\begin{array}{ccc} R & \xrightarrow{\sigma} & Q_l^s(R) \\ & \searrow \sigma_s & \downarrow \sigma_s^s \\ & & S^{-1}R \end{array} \quad (9)$$

where, for $r \in R$ and $s \in T_l(R)$, $\sigma(r) = \frac{r}{1}$, $\sigma_s(r) = \frac{r}{1}$ and $\sigma_s^s(s^{-1}r) = s^{-1}r$. Clearly,

$$\ker(\sigma_s^s) = T_l(R)^{-1}\text{ass}(S). \quad (10)$$

In more detail, if $\sigma_s^s(s^{-1}r) = 0$ then $0 = \frac{r}{1} \in S^{-1}R$, hence $r \in \text{ass}(S)$, and so $\ker(\sigma_s^s) = T_l(R)^{-1}\text{ass}(S)$. Moreover, there is a canonical exact sequence

$$0 \rightarrow T_l(R)^{-1}\mathfrak{I}_R \rightarrow Q_l^s(R) \xrightarrow{\sigma^s} \prod_{S \in \text{max.Den}_l(R)} S^{-1}R, \quad \text{where } \sigma^s := \prod_{S \in \text{max.Den}_l(R)} \sigma_s^s. \quad (11)$$

In more detail, if $\sigma^s(s^{-1}r) = 0$ where $s \in T_l(R)$ and $r \in R$ then $S^{-1}R \ni \sigma_s^s(s^{-1}r) = s^{-1}r = 0$ for all $S \in \text{max.Den}_l(R)$, and so $r \in \text{ass}(S)$ for all $S \in \text{max.Den}_l(R)$, i.e. $r \in \bigcap_{S \in \text{max.Den}_l(R)} \text{ass}(S) = \mathfrak{I}_R$. Therefore, $\ker(\sigma^s) = T_l(R)^{-1}\mathfrak{I}_R$.

Criterion for $S_{l,0}(R) = T_l(R)$. Recall that $S_{l,0}(R) \subseteq T_l(R)$ (Lemma 2.10.(3)). The next lemma is a criterion for a ring R to have the property that $S_{l,0}(R) = T_l(R)$ (or/and that the homomorphism $\theta : Q_l(R) \rightarrow Q_l^s(R)$ is an isomorphism).

Lemma 2.13 *Let R be a ring. The following statements are equivalent.*

1. $S_{l,0}(R) = T_l(R)$.
2. $\mathfrak{I}_R^s = 0$.
3. θ is an isomorphism.
4. θ is a monomorphism.
5. θ is a epimorphism and $S_{l,0}(R) + \mathfrak{I}_R^s \subseteq S_{l,0}(R)$.

Proof. The following implications are obvious (see the exact sequence (8)): $1 \Leftrightarrow 2 \Leftrightarrow 4$, $1 \Rightarrow 3 \Rightarrow 4$ and $3 \Rightarrow 5$.

(5 \Rightarrow 2) It suffices to show that $T_l := T_l(R) \subseteq S_l := S_{l,0}(R)$. Let $t \in T_l$. Then $Q_l^s(R) \ni t^{-1} = \theta(s^{-1}r)$ for some elements $s \in S_l$ and $r \in R$. Hence, $rt = s + a =: s'$ for some element $a \in \mathfrak{I}_R^s$.

Since, by the assumption, $S_l + \mathfrak{l}_R^s \subseteq S_l$, we have $s' \in S_l$. Hence, $\ker(t \cdot) = 0$ (where $t \cdot : R \rightarrow R$, $x \mapsto tx$) since $\ker(t \cdot) \subseteq \ker(rt \cdot) = \ker(s' \cdot) = 0$ as $s' \in S_l$. Thus $\mathfrak{l}_R^s = 0$. \square

Two more characterizations of the set $T_l(R)$. For a ring R , let $'\mathcal{C}_R := \{c \in R \mid \ker(\cdot c) = 0\}$ be the set of left regular elements of R where $\cdot c : R \rightarrow R$, $r \mapsto rc$.

Definition. The sets

$$\begin{aligned}\mathcal{C}_R^w &:= \{c \in R \mid \frac{c}{1} \in \mathcal{C}_{S^{-1}R} \text{ for all } S \in \max.\text{Den}_l(R)\}, \\ '\mathcal{C}_R^w &:= \{c \in R \mid \frac{c}{1} \in '\mathcal{C}_{S^{-1}R} \text{ for all } S \in \max.\text{Den}_l(R)\},\end{aligned}$$

are called the sets of weak regular and weak left regular elements of R .

The sets \mathcal{C}_R^w and $'\mathcal{C}_R^w$ are multiplicative sets such that $R^* \subseteq \mathcal{C}_R^w \subseteq '\mathcal{C}_R^w$.

Lemma 2.14 *Let R be a ring. Then*

1. $\mathcal{C}_R \subseteq '\mathcal{C}_R \subseteq '\mathcal{C}_R^w$.
2. $\mathcal{L}_l^s(R) \subseteq \mathcal{C}_R^w$.

Proof. 1. We have to show that $'\mathcal{C}_R \subseteq '\mathcal{C}_R^w$. Given $c \in '\mathcal{C}_R$. Suppose that $c \notin '\mathcal{C}_R^w$, we seek a contradiction. Then there exist $S \in \max.\text{Den}_l(R)$ and $r \in R$ such that $\frac{r}{1} \frac{c}{1} = 0$ and $\frac{r}{1} \neq 0$. Then $rc \in \text{ass}(S)$, and so $src = 0$ for some $s \in S$. Now, $sr = 0$ (since $c \in '\mathcal{C}_R$). Therefore, $\frac{r}{1} = 0$, a contradiction.

2. Statement 2 follows from Proposition 2.3.(1). \square

By Lemma 2.14.(2),

$$T_l(R) \subseteq \mathcal{L}_l^s(R) \subseteq \mathcal{C}_R^w \subseteq '\mathcal{C}_R^w. \quad (12)$$

Two more characterizations of the set $T_l(R)$ are given below.

Proposition 2.15 1. *The set $T_l(R)$ is the largest left denominator set in the set $'\mathcal{C}_R^w$.*

2. *The set $T_l(R)$ is the largest left denominator set in the set \mathcal{C}_R^w .*

Proof. 1. Given $T \in \text{Den}_l(R)$ such that $T \subseteq '\mathcal{C}_R^w$. We have to show that $T \subseteq T_l(R)$.

(i) For all $S \in \max.\text{Den}_l(R)$, $ST \in \text{Den}_l(R)$: By Lemma 2.4.(2), we have to show that $0 \notin ST$. Suppose that $0 \in ST$ for some $S \in \max.\text{Den}_l(R)$, we seek a contradiction. Then $s_1 t_1 \cdots s_n t_n = 0$ for some elements $s_i \in S$ and $t_i \in T$, and in the ring $S^{-1}R$, $s_1 t_1 \cdots s_n t_n / 1 = 0$. Now, $s_1 t_1 \cdots s_n t_n / 1 = 0$ implies $s_1 t_1 \cdots s_n / 1 = 0$ (since $t_n \in '\mathcal{C}_R^w$) implies $s_1 t_1 \cdots s_{n-1} t_{n-1} / 1 = 0$ (since s_n is a unit in $S^{-1}R$). Continue in this way we obtain that $s_1 / 1 = 0$, a contradiction.

(ii) $T \subseteq \mathcal{L}_l^s(R)$. By (i), for all $S \in \max.\text{Den}_l(R)$, $S \subseteq ST$, hence $S = ST$ (by the maximality of S), and so $T \subseteq ST = S$. Therefore, $T \subseteq \bigcap_{S \in \max.\text{Den}_l(R)} S = \mathcal{L}_l^s(R)$.

(iii) $T \subseteq T_l(R)$, by Proposition 2.8.

2. By (12), $'\mathcal{C}_R^w \supseteq \mathcal{L}_l^s(R) \supseteq T_l(R)$. Now, statement 2 follows from statement 1. \square

3 The largest strong left quotient ring of a ring and its properties

The aim of this section is to prove Theorem 1.1 and Theorem 1.2, and to give a criterion for a ring R to have a semisimple strong left quotient ring (Theorem 1.3).

Let us collect/prove the results (Lemma 3.1, Lemma 3.2 and Proposition 3.3) that are used in the proofs of these theorems. Let $S, T \in \text{Den}_l(R)$. The denominator set T is called *S-saturated* if $sr \in T$, for some $s \in S$ and $r \in R$, then $r \in T$, and if $r's' \in T$, for some $s' \in S$ and $r' \in R$, then $r' \in T$.

Lemma 3.1 [3] Let $S \in \text{Den}_l(R, \mathfrak{a})$, $\pi : R \rightarrow R/\mathfrak{a}$, $a \mapsto a + \mathfrak{a}$, and $\sigma : R \rightarrow S^{-1}R$, $r \mapsto r/1$.

1. Let $T \in \text{Den}_l(S^{-1}R, 0)$ be such that $\pi(S), \pi(S)^{-1} \subseteq T$. Then $T' := \sigma^{-1}(T) \in \text{Den}_l(R, \mathfrak{a})$, T' is S -saturated, $T = \{s^{-1}t' \mid s \in S, t' \in T'\}$, and $S^{-1}R \subseteq T'^{-1}R = T^{-1}R$.
2. $\pi^{-1}(S_0(R/\mathfrak{a})) = S_{\mathfrak{a}}(R)$, $\pi(S_{\mathfrak{a}}(R)) = S_0(R/\mathfrak{a})$ and $Q_{\mathfrak{a}}(R) = S_{\mathfrak{a}}(R)^{-1}R = Q_l(R/\mathfrak{a})$.

The next lemma shows that there is a canonical bijection between the sets of maximal left denominator sets of the rings R and R/\mathfrak{l}_R^s .

Lemma 3.2 Let R be a ring, $\pi : R \rightarrow R/\mathfrak{l}_R^s$, $r \mapsto \bar{r} := r + \mathfrak{l}_R^s$. Then the map

$$\max.\text{Den}_l(R) \rightarrow \max.\text{Den}_l(R/\mathfrak{l}_R^s), \quad S \mapsto \pi(S),$$

is a bijection with the inverse $T \mapsto \pi^{-1}(T)$.

Proof. (i) For all $S \in \max.\text{Den}_l(R)$, $\pi(S) \in \max.\text{Den}_l(R/\mathfrak{l}_R^s)$: Since $\mathfrak{l}_R^s \subseteq \mathfrak{l}_R \subseteq \text{ass}(S)$, $\pi(S) \in \text{Den}_l(R/\mathfrak{l}_R^s, \text{ass}(S)/\mathfrak{l}_R^s)$. Suppose that $T \in \text{Den}_l(R/\mathfrak{l}_R^s)$ with $\pi(S) \subseteq T$, we have to show that $\pi(S) = T$. Let $\pi' : R/\mathfrak{l}_R^s \rightarrow R/\text{ass}(S)$, $r + \mathfrak{l}_R^s \mapsto r + \text{ass}(S)$. Since $\text{ass}(\pi(S)) \subseteq \text{ass}(T)$,

$$\pi'(T) \in \text{Den}_l(R/\text{ass}(S), \text{ass}(T)/\text{ass}(\pi(S))).$$

Using Lemma 2.11 in the situation when $S \in \text{Den}_l(R, \text{ass}(S))$, $\pi'' : R \rightarrow R/\text{ass}(S)$, $r \mapsto r + \text{ass}(S)$, and $\pi'(T) \in \text{Den}_l(R/\text{ass}(S), \text{ass}(T)/\text{ass}(\pi(S))) \simeq \mathfrak{b}/\text{ass}(S)$ where $\mathfrak{b} = \pi^{-1}(\text{ass}(T))$, we conclude that

$$S \subseteq \pi^{-1}\pi(S) \subseteq \pi^{-1}(T) = \pi''^{-1}(\pi'(T)) \in \text{Den}_l(R).$$

Therefore, $S = \pi^{-1}(T)$, by the maximality of S . Hence, $\pi(S) = \pi\pi^{-1}(T) = T$, as required.

(ii) For all $T \in \max.\text{Den}_l(R/\mathfrak{l}_R^s)$, $\pi^{-1}(T) \in \max.\text{Den}_l(R)$: Since $T_l := T_l(R) \in \text{Den}_l(R, \mathfrak{l}_R^s)$, we have $\pi(T_l) \in \text{Den}_l(R/\mathfrak{l}_R^s, 0)$. We claim that

$$0 \notin T\pi(T_l)$$

where $T\pi(T_l)$ is the submonoid of $(R/\mathfrak{l}_R^s, \cdot)$ generated by T and $\pi(T_l)$. Suppose that $0 \in T\pi(T_l)$, i.e. $t_1s_1 \cdots t_ns_n = 0$ for some elements $t_i \in T$ and $s_i \in \pi(T_l)$, we seek a contradiction. Then $t_1s_1 \cdots t_ns_n = 0$ (since $s_n \in \pi(T_l)$ and $\text{ass}(\pi(T_l)) = 0$) and so

$$t'_n t_1 s_1 \cdot t_2 s_2 \cdots t_{n-1} s_{n-1} = 0$$

for some element $t'_n \in T$ (since $t_n \in T$ and $T \in \text{Den}_l(R/\mathfrak{l}_R^s)$). Repeating the same argument $n-1$ more times we obtain elements $t'_2, \dots, t'_{n-1} \in T$ such that $T \ni t'_2 t'_3 \cdots t'_n t_1 = 0$, a contradiction.

Since $T \in \max.\text{Den}_l(R/\mathfrak{l}_R^s)$ and $0 \notin T\pi(T_l)$, we must have $T \subseteq T\pi(T_l) \in \text{Den}_l(R/\mathfrak{l}_R^s)$, by Lemma 2.4.(2). Therefore, $T = T\pi(T_l)$ (by the maximality of T) and so

$$\pi(T_l) \subseteq T.$$

Using Lemma 2.11 in the situation when $S := T_l(R) \in \text{Den}_l(R, \mathfrak{a} := \mathfrak{l}_R^s)$, $\pi : R \rightarrow R/\mathfrak{l}_R^s$ and $T \in \text{Den}_l(R/\mathfrak{l}_R^s, \text{ass}(T))$, we conclude that $T_l(R) \subseteq \pi^{-1}(T) \in \text{Den}_l(R)$. To finish the proof of (ii) we have to show that if $\pi^{-1}(T) \subseteq S'$ for some $S' \in \max.\text{Den}_l(R)$ then $\pi^{-1}(T) = S'$. The inclusion $\pi^{-1}(T) \subseteq S'$ implies the inclusion $T = \pi\pi^{-1}(T) \subseteq \pi(S')$. By (i) and $T \in \max.\text{Den}_l(R/\mathfrak{l}_R^s)$, $T = \pi(S')$. Therefore,

$$\pi^{-1}(T) = \pi^{-1}\pi(S') = S' + \mathfrak{l}_R^s = S'$$

since $\mathfrak{l}_R^s \subseteq \mathfrak{l}_R \subseteq \text{ass}(S')$ and $S' + \text{ass}(S') = S'$, by Corollary 2.12.(2).

(iii) For all $S \in \max.\text{Den}_l(R)$, $\pi^{-1}\pi(S) = S$: $\pi^{-1}\pi(S) = S + \text{ass}(S) = S$, by Corollary 2.12.(2).

(iv) For all $T \in \max.\text{Den}_l(R/\mathfrak{l}_R^s)$, $\pi\pi^{-1}(T) = T$: Trivial.

The proof of the lemma is complete. \square

A bijection between $\max.\text{Den}_l(R)$ and $\max.\text{Den}_l(Q_l(R))$.

Proposition 3.3 [4, Proposition 2.10] *Let R be a ring, S_l be the largest regular left Ore set of the ring R , $Q_l := S_l^{-1}R$ be the largest left quotient ring of the ring R , and \mathcal{C} be the set of regular elements of the ring R . Then*

1. $S_l \subseteq S$ for all $S \in \max.\text{Den}_l(R)$. In particular, $\mathcal{C} \subseteq S$ for all $S \in \max.\text{Den}_l(R)$ provided \mathcal{C} is a left Ore set.
2. Either $\max.\text{Den}_l(R) = \{\mathcal{C}\}$ or, otherwise, $\mathcal{C} \notin \max.\text{Den}_l(R)$.
3. The map

$$\max.\text{Den}_l(R) \rightarrow \max.\text{Den}_l(Q_l), \quad S \mapsto SQ_l^* = \{c^{-1}s \mid c \in S_l, s \in S\},$$

is a bijection with the inverse $\mathcal{T} \mapsto \sigma^{-1}(\mathcal{T})$ where $\sigma : R \rightarrow Q_l$, $r \mapsto \frac{r}{1}$, and SQ_l^* is the sub-semigroup of (Q_l, \cdot) generated by the set S and the group Q_l^* of units of the ring Q_l , and $S^{-1}R = (SQ_l^*)^{-1}Q_l$.

4. If \mathcal{C} is a left Ore set then the map (where $Q = Q_{l,cl}(R)$)

$$\max.\text{Den}_l(R) \rightarrow \max.\text{Den}_l(Q), \quad S \mapsto SQ^* = \{c^{-1}s \mid c \in \mathcal{C}, s \in S\},$$

is a bijection with the inverse $\mathcal{T} \mapsto \sigma^{-1}(\mathcal{T})$ where $\sigma : R \rightarrow Q$, $r \mapsto \frac{r}{1}$, and SQ^* is the sub-semigroup of (Q, \cdot) generated by the set S and the group Q^* of units of the ring Q , and $S^{-1}R = (SQ^*)^{-1}Q$.

Proof of Theorem 1.1. 1. Let $T_l := T_l(R)$ and $\mathcal{S} := S_{l, \mathfrak{I}_R^s}(R)$. Since $T_l \in \text{Den}_l(R, \mathfrak{I}_R^s)$, we have the inclusion $T_l \subseteq \mathcal{S}$. It remains to show that $T_l \supseteq \mathcal{S}$. By Proposition 3.3.(1), and Lemma 3.2, $S_{l,0}(R/\mathfrak{I}_R^s) \subseteq \pi(S)$ for all $S \in \max.\text{Den}_l(R)$. Therefore,

$$\begin{aligned} \mathcal{S} &= \pi^{-1}(S_{l,0}(R/\mathfrak{I}_R^s)) \quad (\text{Lemma 3.1.(2)}) \\ &\subseteq \pi^{-1}\left(\bigcap_{T \in \max.\text{Den}_l(R/\mathfrak{I}_R^s)} T\right) \quad (\text{Proposition 3.3.(1)}) \\ &= \pi^{-1}\left(\bigcap_{S \in \max.\text{Den}_l(R)} \pi(S)\right) \quad (\text{Lemma 3.2}) \\ &= \bigcap_{S \in \max.\text{Den}_l(R)} \pi^{-1}\pi(S) = \bigcap_{S \in \max.\text{Den}_l(R)} (S + \mathfrak{I}_R^s) \\ &= \bigcap_{S \in \max.\text{Den}_l(R)} S \quad (\text{by Corollary 2.12.(2) and } \mathfrak{I}_R^s \subseteq \text{ass}(S)) \\ &= \mathcal{L}_l(R). \end{aligned}$$

Hence, $\mathcal{S} \subseteq T_l$, by the maximality of T_l .

2. By statement 1, $Q_l^s(R) = T_l(R)^{-1}R = S_{l, \mathfrak{I}_R^s}(R)^{-1}R = Q_{l, \mathfrak{I}_R^s}(R)$. By Lemma 3.1.(2), $Q_{l, \mathfrak{I}_R^s}(R) \simeq Q_l(R/\mathfrak{I}_R^s)$.

3 and 4. By Theorem 2.1.(1), $S_{l,0}(R/\mathfrak{I}_R^s) = R/\mathfrak{I}_R^s \cap Q_l(R/\mathfrak{I}_R^s)^*$. Now,

$$\begin{aligned} T_l(R) &\stackrel{\text{st.1}}{=} S_{l, \mathfrak{I}_R^s}(R) = \pi^{-1}(S_{l,0}(R/\mathfrak{I}_R^s)) \quad (\text{Lemma 3.1.(2)}) \\ &= \pi^{-1}(R/\mathfrak{I}_R^s \cap Q_l(R/\mathfrak{I}_R^s)^*) \\ &= \pi^{-1}(R/\mathfrak{I}_R^s \cap Q_l^s(R/\mathfrak{I}_R^s)^*) \quad (\text{by statement 2}) \\ &= \sigma^{-1}(Q_l^s(R/\mathfrak{I}_R^s)^*). \end{aligned}$$

5. By Theorem 2.1.(3),

$$Q_l(R/\mathfrak{I}_R^s)^* = \{s^{-1}t \mid s, t \in S_{l,0}(R/\mathfrak{I}_R^s)\}.$$

By statements 1, 2 and 4, $Q_i^s(R)^* = \{s^{-1}t \mid s, t \in T_l(R)\}$. \square

Proof of Theorem 1.2. 6. Statement 6 follows at once from Lemma 3.2:

$$\begin{aligned} \pi^{-1}(\mathcal{L}_l^s(R/\mathfrak{t}_R^s)) &= \pi^{-1}\left(\bigcap_{T \in \max.\text{Den}_l(R/\mathfrak{t}_R^s)} T\right) = \bigcap_{T \in \max.\text{Den}_l(R/\mathfrak{t}_R^s)} \pi^{-1}(T) \\ &= \bigcap_{S \in \max.\text{Den}_l(R)} S = \mathcal{L}_l^s(R). \end{aligned}$$

Hence, $\pi(\mathcal{L}_l^s(R)) = \pi\pi^{-1}(\mathcal{L}_l^s(R/\mathfrak{t}_R^s)) = \mathcal{L}_l^s(R/\mathfrak{t}_R^s)$.

2 and 4. Since $\pi(T_l(R)) \in \text{Den}_l(R/\mathfrak{t}_R^s, 0)$ and $\pi(T_l(R)) \subseteq \pi(\mathcal{L}_l^s(R)) = \mathcal{L}_l^s(R/\mathfrak{t}_R^s)$, by statement 6, we see that $\pi(T_l(R)) \subseteq T_l(R/\mathfrak{t}_R^s)$, by the maximality of $T_l(R/\mathfrak{t}_R^s)$.

Conversely, by Lemma 3.2, $T_l(R/\mathfrak{t}_R^s) \subseteq \pi(S)$ for all $S \in \max.\text{Den}_l(R)$. Therefore,

$$\pi^{-1}(T_l(R/\mathfrak{t}_R^s)) \subseteq \pi^{-1}\pi(S) = S \text{ for all } S \in \max.\text{Den}_l(R) \text{ (Lemma 3.2),}$$

hence $\pi^{-1}(T_l(R/\mathfrak{t}_R^s)) \subseteq \bigcap_{S \in \max.\text{Den}_l(R)} S = \mathcal{L}_l^s(R)$. Since $T_l(R/\mathfrak{t}_R^s) \in \text{Den}_l(R/\mathfrak{t}_R^s)$, $T_l(R) \in \text{Den}_l(R, \mathfrak{t}_R^s)$ and $\pi(T_l(R)) \subseteq T_l(R/\mathfrak{t}_R^s)$, we obtain that $T_l(R) \subseteq \pi^{-1}(T_l(R/\mathfrak{t}_R^s)) \in \text{Den}_l(R)$, by Lemma 2.11. Hence,

$$\pi^{-1}(T_l(R/\mathfrak{t}_R^s)) = T_l(R),$$

by the maximality of $T_l(R)$. Hence,

$$T_l(R/\mathfrak{t}_R^s) = \pi\pi^{-1}(T_l(R/\mathfrak{t}_R^s)) = \pi(T_l(R)) \in \text{Den}_l(R/\mathfrak{t}_R^s, 0).$$

This means that statements 2 and 4 hold.

3. Statement 3 follows from statement 4 and Theorem 1.1.(1).

5. Let $\bar{\sigma} : R/\mathfrak{t}_R^s \rightarrow Q_l(R/\mathfrak{t}_R^s)$, $\bar{\sigma} \mapsto \frac{\bar{\sigma}}{\bar{\sigma}}$. By Proposition 3.3.(3), the map

$$\max.\text{Den}_l(R/\mathfrak{t}_R^s) \rightarrow \max.\text{Den}_l(Q_l(R/\mathfrak{t}_R^s)), \quad S \mapsto SQ_l(R/\mathfrak{t}_R^s)^* = \{c^{-1}s \mid c \in S_{l,0}(R/\mathfrak{t}_R^s), s \in S\}, \quad (13)$$

is a bijection with the inverse $T \mapsto \bar{\sigma}^{-1}(T) = T \cap R/\mathfrak{t}_R^s$. Let $\mathcal{T} := T_l(Q_l(R/\mathfrak{t}_R^s))$. Then $Q_l(R/\mathfrak{t}_R^s)^* \subseteq \mathcal{T}$, by Lemma 2.10.(3). By (13), $\mathcal{T} \subseteq SQ_l(R/\mathfrak{t}_R^s)^*$ for all $S \in \max.\text{Den}(R/\mathfrak{t}_R^s)$. Hence,

$$\begin{aligned} \mathcal{T} \cap R/\mathfrak{t}_R^s &\subseteq \bigcap_{S \in \max.\text{Den}_l(R/\mathfrak{t}_R^s)} R/\mathfrak{t}_R^s \cap SQ_l(R/\mathfrak{t}_R^s)^* \\ &= \bigcap_{S \in \max.\text{Den}_l(R/\mathfrak{t}_R^s)} S \quad (\text{since } S = R/\mathfrak{t}_R^s \cap SQ_l(R/\mathfrak{t}_R^s)^*, \text{ by (13)}) \\ &= \mathcal{L}_l^s(R/\mathfrak{t}_R^s). \end{aligned}$$

Also,

$$\begin{aligned} \mathcal{T} \cap R/\mathfrak{t}_R^s &\supseteq R/\mathfrak{t}_R^s \cap Q_l(R/\mathfrak{t}_R^s)^* \quad (\text{since } \mathcal{T} \supseteq Q_l(R/\mathfrak{t}_R^s)^*) \\ &= S_{l,0}(R/\mathfrak{t}_R^s) \quad (\text{Theorem 2.1.(1)}) \\ &= T_l(R/\mathfrak{t}_R^s), \quad (\text{by statement 3}). \end{aligned}$$

Applying Lemma 3.1 to the case where $S = S_{l,0}(R/\mathfrak{t}_R^s) \in \text{Den}_l(R/\mathfrak{t}_R^s, \mathfrak{a} := 0)$ and $T = \mathcal{T}$ (notice that $Q_l(R/\mathfrak{t}_R^s)^* \subseteq \mathcal{T}$) we see that

$$\mathcal{T} \cap R/\mathfrak{t}_R^s = \bar{\sigma}^{-1}(\mathcal{T}) \in \text{Den}_l(R/\mathfrak{t}_R^s).$$

This fact together with the inclusions (see above) $T_l(R/\mathfrak{t}_R^s) \subseteq \mathcal{T} \cap R/\mathfrak{t}_R^s \subseteq \mathcal{L}_l^s(R/\mathfrak{t}_R^s)$ implies the equality

$$T_l(R/\mathfrak{t}_R^s) = \mathcal{T} \cap R/\mathfrak{t}_R^s, \quad (14)$$

by the maximality of $T_l(R/\mathfrak{l}_R^s)$. The inclusion $Q_l(R/\mathfrak{l}_R^s)^* \subseteq \mathcal{T}$ and the fact that $Q_l(R/\mathfrak{l}_R^s)^* = \{s^{-1}t \mid s, t \in S_{l,0}(R/\mathfrak{l}_R^s)\}$ (Theorem 2.1.(3)) imply that

$$\mathcal{T} = \{s^{-1}r \mid s \in S_{l,0}(R/\mathfrak{l}_R^s); r \in \mathcal{T} \cap R/\mathfrak{l}_R^s\}.$$

In more detail, if $s^{-1}r \in \mathcal{T}$ for some $s \in S_{l,0}(R/\mathfrak{l}_R^s) \subseteq \mathcal{T}$ and $\frac{r}{1} \in R/\mathfrak{l}_R^s$ then $\frac{r}{1} \in s\mathcal{T} \subseteq \mathcal{T}\mathcal{T} = \mathcal{T}$, and so $r \in \mathcal{T} \cap R/\mathfrak{l}_R^s$.

By (14) and $T_l(R/\mathfrak{l}_R^s) = S_{l,0}(R/\mathfrak{l}_R^s)$ (statement 3),

$$\mathcal{T} = \{s^{-1}r \mid s \in T_l(R/\mathfrak{l}_R^s), r \in T_l(R/\mathfrak{l}_R^s)\} = Q_l(R/\mathfrak{l}_R^s)^*.$$

Hence, $\text{ass}(\mathcal{T}) = \text{ass}(Q_l(R/\mathfrak{l}_R^s)^*) = 0$.

1. Statement 1 follows from statement 5.

7. By statement 4 and Theorem 1.1.(2), $Q_l^s(R/\mathfrak{l}_R^s) \simeq Q_l((R/\mathfrak{l}_R^s)/\mathfrak{l}_{R/\mathfrak{l}_R^s}^s) = Q_l(R/\mathfrak{l}_R^s)$. \square

Necessary and sufficient conditions for $Q_l(R)$ to be a semi-simple ring. A ring Q is called a *ring of quotients* if every element $c \in \mathcal{C}_Q$ is invertible. A subring R of a ring of quotients Q is called a *left order* in Q if \mathcal{C}_R is a left Ore set and $\mathcal{C}_R^{-1}R = Q$. A ring R has *finite left rank* (i.e. *finite left uniform dimension*) if there are no infinite direct sums of nonzero left ideals in R .

The next theorem gives an answer to the question of when $Q_l(R)$ is a semi-simple ring. Theorem 3.4 is the key result in the proof of Theorem 1.3.

Theorem 3.4 [3] *The following properties of a ring R are equivalent.*

1. $Q_l(R)$ is a semi-simple ring.
2. $Q_{cl}(R)$ exists and is a semi-simple ring.
3. R is a left order in a semi-simple ring.
4. R has finite left rank, satisfies the ascending chain condition on left annihilators and is a semi-prime ring.
5. A left ideal of R is essential iff it contains a regular element.

If one of the equivalent conditions hold then $S_0(R) = \mathcal{C}_R$ and $Q_l(R) = Q_{cl}(R)$.

Proof of Theorem 1.3. (2 \Leftrightarrow 4) This is the Goldie's Theorem for the ring R/\mathfrak{l}_R^s .

(1 \Leftrightarrow 3 \Leftrightarrow 4) By Theorem 1.1.(2), $Q_l^s(R) \simeq Q_l(R/\mathfrak{l}_R^s)$. Theorem 3.4 implies that (1 \Leftrightarrow 3 \Leftrightarrow 4) and that $Q_l(R/\mathfrak{l}_R^s) \simeq Q_{l,cl}(R/\mathfrak{l}_R^s)$ and $S_{l,0}(R/\mathfrak{l}_R^s) = \mathcal{C}_{R/\mathfrak{l}_R^s}$. By Theorem 1.2.(3), $S_{l,0}(R/\mathfrak{l}_R^s) = T_l(R/\mathfrak{l}_R^s)$. Then, by Theorem 1.2.(2), $T_l(R) = \pi^{-1}(T_l(R/\mathfrak{l}_R^s)) = \pi^{-1}(S_{l,0}(R/\mathfrak{l}_R^s))$. By Theorem 1.2.(2), $T_l(R/\mathfrak{l}_R^s) = \pi(T_l) = \pi\pi^{-1}(S_{l,0}(R/\mathfrak{l}_R^s)) = S_{l,0}(R/\mathfrak{l}_R^s) = \mathcal{C}_{R/\mathfrak{l}_R^s}$. \square

The maximal left denominator sets of a finite direct product of rings.

Theorem 3.5 [4, Theorem 2.9] *Let $R = \prod_{i=1}^n R_i$ be a direct product of rings R_i . Then for each $i = 1, \dots, n$, the map*

$$\text{max.Den}_l(R_i) \rightarrow \text{max.Den}_l(R), \quad S_i \mapsto R_1 \times \cdots \times S_i \times \cdots \times R_n, \quad (15)$$

is an injection. Moreover, $\text{max.Den}_l(R) = \coprod_{i=1}^n \text{max.Den}_l(R_i)$ in the sense of (15), i.e.

$$\text{max.Den}_l(R) = \{S_i \mid S_i \in \text{max.Den}_l(R_i), i = 1, \dots, n\},$$

$$S_i^{-1}R \simeq S_i^{-1}R_i, \quad \text{ass}_R(S_i) = R_1 \times \cdots \times \text{ass}_{R_i}(S_i) \times \cdots \times R_n.$$

Proof of Theorem 1.4. (1 \Leftrightarrow 2) These implications are Theorem 3.4.(1,2).

(3 \Rightarrow 1) The implication follows from Theorem 1.3. In particular, $Q_{l,cl}(R) \simeq Q_l^s(R)$.

(1 \Rightarrow 3) (i) $\mathcal{L}_l^s(R) = \mathcal{C}_R$: Since $Q_{l,cl}(R)$ is a semisimple ring, $\mathcal{C}_R = S_{l,0}(R) \subseteq \mathcal{L}_l^s(R)$, by (3). It remains to show that $\mathcal{L}_l^s(R) \subseteq \mathcal{C}_R$. The ring $Q_{l,cl}(R) = \prod_{i=1}^n R_i$ is a semisimple ring where R_i are simple Artinian rings (i.e. matrix rings over division rings). Clearly, $\max.\text{Den}_l(R_i) = \{R_i^*\}$ where R_i^* is the group of units of R_i . By Theorem 3.5, $\max.\text{Den}_l(R) = \{\mathcal{S}_1, \dots, \mathcal{S}_n\}$ where $\mathcal{S}_i = R_1 \times \dots \times R_{i-1} \times R_i^* \times R_{i+1} \times \dots \times R_n$ for $i = 1, \dots, n$. The map

$$\sigma : R \rightarrow Q_{l,cl}(R) = \prod_{i=1}^n R_i, \quad r \mapsto \frac{r}{1} = (r_1, \dots, r_n),$$

is a monomorphism. Then an element $r \in R$ is regular iff the element $\frac{r}{1} = (r_1, \dots, r_n) \in Q_{l,cl}(R) = \prod_{i=1}^n R_i$ is regular iff $r_1 \in \mathcal{C}_{R_1} = R_1^*, \dots, r_n \in \mathcal{C}_{R_n} = R_n^*$ iff $\frac{r}{1} \in \bigcap_{i=1}^n \mathcal{S}_i$ iff

$$r = \sigma^{-1}(r) \in \sigma^{-1}\left(\bigcap_{i=1}^n \mathcal{S}_i\right) = \bigcap_{i=1}^n \sigma^{-1}(\mathcal{S}_i) = \bigcap_{S \in \max.\text{Den}_l(R)} S = \mathcal{L}_l^s(R),$$

by Proposition 3.3.(3) and Theorem 3.5.

(ii) $T_l(R) = \mathcal{C}_R$: By (3), $\mathcal{C}_R = S_{l,0}(R) \subseteq T_l(R) \subseteq \mathcal{L}_l^s(R) = \mathcal{C}_R$.

(iii) $Q_{l,cl}(R) = Q_l^s(R)$ is a semisimple ring, by (ii) and $\mathfrak{I}_R^s = \text{ass}(\mathcal{C}_R) = 0$. \square

The operations $\mathcal{L}_l^s(\cdot) : R \mapsto \mathcal{L}_l^s(R)$, $T_l(\cdot) : R \mapsto T_l(R)$ and $Q_l^s(\cdot) : R \mapsto Q_l^s(R)$ commute with finite direct products as the next theorem shows.

Theorem 3.6 *Let $R = \prod_{i=1}^n R_i$ be a direct product of rings. Then*

1. $\mathcal{L}_l^s(R) = \prod_{i=1}^n \mathcal{L}_l^s(R_i)$
2. $T_l(R) = \prod_{i=1}^n T_l(R_i)$ and $Q_l^s(R) \simeq \prod_{i=1}^n Q_l^s(R_i)$.

Proof. 1. Statement 1 follows from Theorem 3.5:

$$\begin{aligned} \mathcal{L}_l^s(R) &= \bigcap_{S \in \max.\text{Den}_l(R)} S = \prod_{i=1}^n \bigcap_{S_i \in \max.\text{Den}_l(R_i)} S_i \quad (\text{Theorem 3.5}) \\ &= \prod_{i=1}^n \mathcal{L}_l^s(R_i). \end{aligned}$$

2. Let $T_l := T_l(R)$ and $T = \prod_{i=1}^n T_l(R_i)$. We have to show that $T_l = T$. Clearly, $T \in \text{Den}_l(R)$ and $T \subseteq \prod_{i=1}^n \mathcal{L}_l^s(R_i) = \mathcal{L}_l^s(R)$, by statement 1. Therefore, $T \subseteq T_l$, by the maximality of T_l . It remains to show that $T_l \subseteq T$. Since $T_l = \prod_{i=1}^n T'_i$ for some $T'_i \in \text{Den}_l(R_i)$ such that $T'_i \subseteq \mathcal{L}_l^s(R_i)$ (by statement 1). Therefore, $T'_i \subseteq T_l(R_i)$ for $i = 1, \dots, n$, by the maximality of $T_l(R_i)$, and so $T_l \subseteq T$. \square

Lemma 3.7 [3] *Let $S \in \text{Den}_l(R, \mathfrak{a})$ and $T \in \text{Den}_l(R, \mathfrak{b})$ such that $\mathfrak{a} \subseteq \mathfrak{b}$. Then*

1. $r.\text{ass}(ST) \subseteq \mathfrak{b}$ where $r.\text{ass}(ST) := \{r \in R \mid rc = 0 \text{ for some } c \in ST\}$.
2. $ST \in \text{Den}_l(R, \mathfrak{c})$ and $\mathfrak{b} \subseteq \mathfrak{c}$.

Proof of Theorem 1.5. (i) $S_{l,\mathfrak{a}}(R) \subseteq S$ for all $S \in \max.\text{Den}_l(R)$: By Lemma 3.7, $S_{l,\mathfrak{a}}(R)S \in \text{Den}_l(R)$ since $\mathfrak{a} \subseteq \mathfrak{I}_R^s \subseteq \text{ass}(S)$. Therefore, $S_{l,\mathfrak{a}}(R) \subseteq S_{l,\mathfrak{a}}(R)S = S$, by the maximality of S .

(ii) $S_{l,\mathfrak{a}}(R) \subseteq T_l(R)$: By (i), $S_{l,\mathfrak{a}}(R) \subseteq \mathcal{L}_l^s(R)$, hence $S_{l,\mathfrak{a}}(R) \subseteq T_l(R)$, by the maximality of $T_l(R)$, Proposition 2.8. Hence, there is a ring R -homomorphism $Q_{l,\mathfrak{a}}(R) \rightarrow Q_l^s(R)$, $s^{-1}r \mapsto s^{-1}r$. \square

The largest strong quotient ring of $Q_l(R)$

For an arbitrary ring R , Theorem 1.6 establishes natural and tight connections between triples $T_l(R)$, \mathfrak{l}_R^s , $Q_l^s(R)$ and $T_l(Q_l(R))$, $\mathfrak{l}_{Q_l(R)}^s$, $Q_l^s(Q_l(R))$. The applications of this theorem are given in Section 5 where it is used in giving explicit descriptions of the triple $T_l(R)$, \mathfrak{l}_R^s , $Q_l^s(R)$ for every ring R such that its classical left quotient ring $Q_{l,cl}(R)$ is a left Artinian ring, see Theorem 5.7 (notice that in this case $Q_{l,cl}(R) = Q_l(R)$ ([3, Corollary 2.10]).

Proof of Theorem 1.6. 1 and 2. Let $S_l = S_l(R)$ and $Q = Q_l(R)$. Since $S_l(R) \subseteq T_l(R)$ (Lemma 2.10.(3)), the multiplicative submonoid $Q^*T_l(R)$ of Q generated by Q^* and $T_l(R)$ belongs to $\text{Den}_l(Q, S_l(R)^{-1}\mathfrak{l}_R^s)$. In view of the bijection between the sets $\text{max.Den}_l(R)$ and $\text{max.Den}_l(Q)$ given by Proposition 3.3.(3), we have the inclusion $Q^*T_l(R) \subseteq \mathcal{L}_l^s(Q)$ which immediately implies the inclusion $Q^*T_l(R) \subseteq T_l(Q)$. We identify the ring R with its image in Q via $\sigma : R \rightarrow Q, r \mapsto \frac{r}{1}$. By Lemma 3.1.(1), $T_l(Q) \cap R \in \text{Den}_l(R, R \cap \mathfrak{l}_Q^s)$ and $T_l(Q) = \{s^{-1}t \mid s \in S_l(R), t \in T_l(Q) \cap R\}$. By Proposition 3.3.(3), $T_l(Q) \cap R \subseteq \mathcal{L}_l^s(R)$, and so $T_l(Q) \cap R \subseteq T_l(R)$. Now,

$$T_l(R) \subseteq Q^*T_l(R) \cap R \subseteq T_l(Q) \cap R \subseteq T_l(R),$$

i.e. $T_l(R) = Q^*T_l(R) \cap R = T_l(Q) \cap R \in \text{Den}_l(R, R \cap \mathfrak{l}_R^s)$. In particular, $\mathfrak{l}_R^s = R \cap \mathfrak{l}_Q^s$. By Lemma 3.1.(1),

$$Q^*T_l(R) = \{s^{-1}t \mid s \in S_l(R), t \in T_l(R) = T_l(Q) \cap R\} = T_l(Q).$$

So, statement 1 is proven. Since $T_l(Q) = Q^*T_l(R) \in \text{Den}_l(Q, S_l(R)^{-1}\text{ass}(T_l(R)))$, we must have $\mathfrak{l}_Q^s = \text{ass}(T_l(Q)) = S_l(R)^{-1}\mathfrak{l}_R^s$. So, statement 2 is proven.

3. Let $Q = Q_l(R)$. Then

$$Q_l^s(R) = T_l(R)^{-1}R \simeq (Q^*T_l(R))^{-1}S_l(R)^{-1}R \stackrel{\text{st.1}}{\simeq} T_l(Q)^{-1}Q \simeq Q_l^s(Q). \quad \square$$

4 The largest strong quotient ring of a ring

In this section, the two-sided versions of the concepts appeared in Sections 2 and 3 are introduced: the *largest strong denominator set* $T(R)$, the *largest strong quotient ring* $Q^s(R) = T(R)^{-1}R$ and the *strong localization radical* $\mathfrak{l}_{R,t}^s$ (the subscript t stands for ‘two-sided’, i.e. ‘left and right’). All the results of the previous sections are true (with obvious adjustments) for *left and right* Ore/denominator sets. For the analogous versions we state the corresponding results and the proofs are left for the reader as an exercise in the case when they are literally the same (with obvious modifications). The following notation is fixed:

- $\text{Den}(R, \mathfrak{a})$ is the set of (left and right) denominator sets S of R with $\text{ass}(S) = \mathfrak{a}$;
- $S_{\mathfrak{a}} = S_{\mathfrak{a}}(R) = S_{l,r,\mathfrak{a}}(R)$ is the largest element of the poset $(\text{Den}(R, \mathfrak{a}), \subseteq)$, i.e. the *largest denominator set in R associated with \mathfrak{a}* , and $Q_{\mathfrak{a}}(R) := Q_{l,r,\mathfrak{a}}(R) := S_{l,r,\mathfrak{a}}^{-1}R$ is the *largest (left and right) quotient ring associated with \mathfrak{a}* , $S_{l,r,\mathfrak{a}}$ exists, [3];
- $\text{max.Den}(R)$ is the set of maximal denominator sets of R (it is a *non-empty* set, Lemma 2.5.(3));
- $\text{Ass}(R) := \{\text{ass}(S) \mid S \in \text{Den}(R)\}$.

The sets

$$\mathcal{L}(R) := \bigcup_{S \in \text{max.Den}(R)} S \quad \text{and} \quad \mathcal{NL}(R) := R \setminus \mathcal{L}(R)$$

are called the sets of *localizable* and *non-localizable elements* of R , respectively, and the intersection

$$\mathcal{L}^s(R) := \bigcap_{S \in \text{max.Den}(R)} S$$

is called the *set of strongly (or completely) localizable elements* of R . Clearly, $\mathcal{L}^s(R)$ is a multiplicative set and

$$R^* \subseteq \mathcal{L}^s(R) \quad (16)$$

since $R^* \subseteq S$ for all $S \in \max.\text{Den}(R)$, by Lemma 3.7.(2). Similarly, by Lemma 3.7.(2),

$$S_{l,r,0}(R) \subseteq \mathcal{L}^s(R). \quad (17)$$

The next proposition is a characterization of the set $\mathcal{L}^s(R)$ which says that the set $\mathcal{L}^s(R)$ contains *precisely* the elements of the ring R that are units in all maximal localizations of R .

Proposition 4.1 *Let R be a ring. Then*

1. $\mathcal{L}^s(R) = \{s \in R \mid \frac{s}{1} \in (S^{-1}R)^* \text{ for all } S \in \max.\text{Den}(R)\}$ where $(S^{-1}R)^*$ is the group of units of the ring $S^{-1}R$.
2. For all automorphisms $\sigma \in \text{Aut}(R)$, $\sigma(\mathcal{L}^s(R)) = \mathcal{L}^s(R)$.

Proof. 1. Let \mathcal{R} be the RHS of the equality. By the very definition of the set $\mathcal{L}^s(R)$, $\mathcal{L}^s(R) \subseteq \mathcal{R}$. Conversely, let $s \in \mathcal{R}$ and $\sigma_S : R \rightarrow S^{-1}R$, $r \mapsto \frac{r}{1}$, where $S \in \max.\text{Den}(R)$. Then $s \in \sigma_S^{-1}((S^{-1}R)^*) = S$ for all $S \in \max.\text{Den}(R)$, [3, Theorem 4.11.(2,3)], hence $s \in \mathcal{L}^s(R)$.

2. Obvious. \square

The largest strong denominator set $T(R)$. The set $\text{Den}^s(R) := \{T \in \text{Den}(R) \mid T \subseteq \mathcal{L}_l^s(R)\}$ is a *non-empty* set since $R^* \in \text{Den}^s(R)$. The elements of $\text{Den}^s(R)$ are called the *strong denominator sets* of R and the rings $T^{-1}R$ where $T \in \text{Den}^s(R)$ are called the *strong quotient rings* or the *strong localizations* of R . Proposition 4.2 shows that the set of maximal elements $\max.\text{Den}^s(R)$ of the poset $(\text{Den}^s(R), \subseteq)$ is a non-empty set. Moreover, it contains a single element. Namely,

$$T(R) := \bigcup_{S \in \text{Den}^s(R)} S. \quad (18)$$

Proposition 4.2 *Let R be a ring. Then $\max.\text{Den}^s(R) = \{T(R)\}$.*

So, $T(R)$ is the *largest strong denominator set* of R . The ideal of the ring R ,

$$\mathfrak{l}_{R,t}^s := \text{ass}(T(R)) = \bigcup_{S \in \text{Den}^s(R)} \text{ass}(S)$$

is called the *strong localization radical* of the ring R .

The ideal

$$\mathfrak{l}_{R,t} := \bigcap_{S \in \max.\text{Den}(R)} \text{ass}(S)$$

is called the *(two-sided) localization radical* of R . There is an exact sequence

$$0 \rightarrow \mathfrak{l}_{R,t} \rightarrow R \xrightarrow{\sigma} \prod_{S \in \max.\text{Den}(R)} S^{-1}R, \quad \sigma = \prod_{S \in \max.\text{Den}(R)} \sigma_S,$$

where $\sigma_S : R \rightarrow S^{-1}R$, $r \mapsto \frac{r}{1}$. The set $\text{Den}^s(R)$ is invariant under the action of the group of automorphisms $\text{Aut}(R)$ of the ring R .

Lemma 4.3 1. For all automorphisms $\sigma \in \text{Aut}(R)$, $\sigma(T(R)) = T(R)$, $\sigma(\mathfrak{l}_{R,t}) = \mathfrak{l}_{R,t}$ and $\sigma(\mathfrak{l}_{R,t}^s) = \mathfrak{l}_{R,t}^s$.

2. $\mathfrak{l}_{R,t}^s \subseteq \mathfrak{l}_{R,t}$.

Lemma 4.4 *Let R be a ring. Then*

1. $T(R) + \mathfrak{I}_{R,t}^s \subseteq T(R)$.
2. $\mathcal{L}^s(R) + \mathfrak{I}_{R,t} \subseteq \mathcal{L}^s(R)$. In particular, $T(R) + \mathfrak{I}_{R,t} \subseteq \mathcal{L}^s(R)$.
3. $R^* \subseteq S_{l,r,0}(R) \subseteq T(R)$.

The largest strong quotient ring $Q^s(R)$ of a ring R . The ring $Q_l(R) := T_l(R)^{-1}R$ is called the *largest strong quotient ring* of the ring R . There are exact sequences:

$$0 \rightarrow \mathfrak{I}_{R,t}^s \rightarrow R \rightarrow Q^s(R), \quad r \mapsto \frac{r}{1}, \quad (19)$$

$$0 \rightarrow S_{l,r,0}(R)^{-1}\mathfrak{I}_{R,t}^s \rightarrow Q(R) := S_{l,r,0}(R)^{-1}R \xrightarrow{\theta} Q^s(R), \quad \theta(s^{-1}r) = s^{-1}r, \quad (20)$$

where $s \in S_{l,r,0}(R)$ and $r \in R$ (if $\theta(s^{-1}r) = 0$ then $\frac{r}{1} = 0$ in $Q^s(R)$, hence $r \in \mathfrak{I}_{R,t}^s$, and so $\ker(\theta) = S_{l,r,0}^{-1}(R)^{-1}\mathfrak{I}_{R,t}^s$) and $Q(R)$ is the *largest (two-sided) quotient ring* of R , [3].

For each $S \in \max.\text{Den}(R)$, there is a commutative diagram of ring homomorphisms:

$$\begin{array}{ccc} R & \xrightarrow{\sigma} & Q^s(R) \\ & \searrow \sigma_s & \downarrow \sigma_s^s \\ & & S^{-1}R \end{array} \quad (21)$$

where, for $r \in R$ and $s \in T(R)$, $\sigma(r) = \frac{r}{1}$, $\sigma_s(r) = \frac{r}{1}$ and $\sigma_s^s(s^{-1}r) = s^{-1}r$. Clearly,

$$\ker(\sigma_s^s) = T(R)^{-1}\text{ass}(S). \quad (22)$$

In more detail, if $\sigma_s^s(s^{-1}r) = 0$ then $0 = \frac{r}{1} \in S^{-1}R$, hence $r \in \text{ass}(S)$, and so $\ker(\sigma_s^s) = T(R)^{-1}\text{ass}(S)$. Moreover, there is a canonical exact sequence

$$0 \rightarrow T(R)^{-1}\mathfrak{I}_{R,t} \rightarrow Q^s(R) \xrightarrow{\sigma^s} \prod_{S \in \max.\text{Den}(R)} S^{-1}R, \quad \text{where } \sigma^s := \prod_{S \in \max.\text{Den}(R)} \sigma_s^s. \quad (23)$$

In more detail, if $\sigma^s(s^{-1}r) = 0$ where $s \in T(R)$ and $r \in R$ then $S^{-1}R \ni \sigma_s^s(s^{-1}r) = s^{-1}r = 0$ for all $S \in \max.\text{Den}(R)$, and so $r \in \text{ass}(S)$ for all $S \in \max.\text{Den}(R)$, i.e. $r \in \bigcap_{S \in \max.\text{Den}(R)} \text{ass}(S) = \mathfrak{I}_{R,t}$. Therefore, $\ker(\sigma^s) = T(R)^{-1}\mathfrak{I}_{R,t}$.

Criterion for $S_{l,r,0}(R) = T(R)$. Recall that $S_{l,r,0}(R) \subseteq T_l(R)$ (Lemma 4.4.(3)). The next lemma is a criterion for a ring R to have the property that $S_{l,r,0}(R) = T(R)$ (or/and that the homomorphism $\theta : Q_l(R) \rightarrow Q^s(R)$ is an isomorphism).

Lemma 4.5 *Let R be a ring. The following statements are equivalent.*

1. $S_{l,r,0}(R) = T(R)$.
2. $\mathfrak{I}_{R,t}^s = 0$.
3. θ is an isomorphism.
4. θ is a monomorphism.
5. θ is an epimorphism and $S_{l,r,0}(R) + \mathfrak{I}_{R,t}^s \subseteq S_{l,r,0}(R)$.

Two more characterizations of the set $T(R)$. The sets

$$\begin{aligned}\mathcal{C}_{R,t}^w &:= \{c \in R \mid \frac{c}{1} \in \mathcal{C}_{S^{-1}R} \text{ for all } S \in \max.\text{Den}(R)\}, \\ {}'\mathcal{C}_{R,t}^w &:= \{c \in R \mid \frac{c}{1} \in {}'\mathcal{C}_{S^{-1}R} \text{ for all } S \in \max.\text{Den}(R)\},\end{aligned}$$

are called the *sets of two-sided weak regular* and *weak left regular elements* of R , respectively.

The sets $\mathcal{C}_{R,t}^w$ and ${}'\mathcal{C}_{R,t}^w$ are multiplicative sets such that $R^* \subseteq \mathcal{C}_{R,t}^w \subseteq {}'\mathcal{C}_{R,t}^w$.

Lemma 4.6 *Let R be a ring. Then*

1. $\mathcal{C}_R \subseteq {}'\mathcal{C}_R \subseteq {}'\mathcal{C}_{R,t}^w$.
2. $\mathcal{L}^s(R) \subseteq \mathcal{C}_{R,t}^w$.

Proof. 1. We have to show that ${}'\mathcal{C}_R \subseteq {}'\mathcal{C}_{R,t}^w$. Given $c \in {}'\mathcal{C}_R$. Suppose that $c \notin {}'\mathcal{C}_{R,t}^w$, we seek a contradiction. Then there exist $S \in \max.\text{Den}(R)$ and $r \in R$ such that $\frac{r}{1} \frac{c}{1} = 0$ and $\frac{r}{1} \neq 0$. Then $rc \in \text{ass}(S)$, and so $src = 0$ for some $s \in S$. Now, $sr = 0$ (since $c \in {}'\mathcal{C}_R$). Therefore, $\frac{r}{1} = 0$, a contradiction.

2. Statement 2 follows from Proposition 4.1.(1). \square

By Lemma 4.6.(2),

$$T(R) \subseteq \mathcal{L}^s(R) \subseteq \mathcal{C}_{R,t}^w \subseteq {}'\mathcal{C}_{R,t}^w. \quad (24)$$

The next proposition gives two more characterizations of the set $T(R)$.

Proposition 4.7 1. *The set $T(R)$ is the largest left denominator set in the set ${}'\mathcal{C}_{R,t}^w$.*

2. *The set $T(R)$ is the largest left denominator set in the set $\mathcal{C}_{R,t}^w$.*

The next lemma shows that there is a bijection between maximal denominator sets of the rings R and $R/\mathfrak{I}_{R,t}^s$.

Lemma 4.8 *Let R be a ring, $\pi : R \rightarrow R/\mathfrak{I}_{R,t}^s$, $R \mapsto \bar{r} := r + \mathfrak{I}_{R,t}^s$. Then the map*

$$\max.\text{Den}(R) \rightarrow \max.\text{Den}(R/\mathfrak{I}_{R,t}^s), \quad S \mapsto \pi(S),$$

is a bijection with the inverse $T \mapsto \pi^{-1}(T)$.

Theorem 4.9 *Let R be a ring, $\pi : R \rightarrow R/\mathfrak{I}_{R,t}^s$, $r \mapsto \bar{r} = r + \mathfrak{I}_{R,t}^s$; $\sigma : R \rightarrow Q^s(R)$, $r \mapsto \frac{r}{1}$, and $Q^s(R)^*$ be the group of units of the ring $Q^s(R)$. Then*

1. $T(R) = S_{l,r,\mathfrak{I}_{R,t}^s}(R)$.
2. $Q^s(R) = Q_{l,r,\mathfrak{I}_{R,t}^s}(R) \simeq Q_l(R/\mathfrak{I}_{R,t}^s)$.
3. $T(R) = \sigma^{-1}(Q^s(R)^*)$.
4. $T(R) = \pi^{-1}(S_{l,r,0}(R/\mathfrak{I}_{R,t}^s))$.
5. $Q^s(R)^* = \{s^{-1}t \mid s, t \in T(R)\}$.

Theorem 4.10 *We keep the notation of Theorem 4.9. Then*

1. $Q^s(Q^s(R)) = Q^s(R)$.
2. $T(R/\mathfrak{I}_{R,t}^s) = \pi(T(R))$ and $T(R) = \pi^{-1}(T(R/\mathfrak{I}_{R,t}^s))$.
3. $T(R/\mathfrak{I}_{R,t}^s) = S_{l,r,0}(R/\mathfrak{I}_{R,t}^s)$.

4. $\mathfrak{l}_{R/\mathfrak{l}_{R,t}^s}^s = 0$.
5. $T(Q^s(R)) = Q^s(R)^*$ and $\mathfrak{l}_{Q^s(R),t}^s = 0$.
6. $\pi(\mathcal{L}^s(R)) = \mathcal{L}^s(R/\mathfrak{l}_{R,t}^s)$ and $\mathcal{L}^s(R) = \pi^{-1}(\mathcal{L}^s(R/\mathfrak{l}_{R,t}^s))$.
7. $Q^s(R/\mathfrak{l}_{R,t}^s) = Q(R/\mathfrak{l}_{R,t}^s)$.

Semisimplicity criterion for the ring $Q^s(R)$. A ring is called a *Goldie ring* if it is a left and right Goldie ring.

Theorem 4.11 *Let R be a ring. The following statements are equivalent.*

1. $Q^s(R)$ is a semisimple ring.
2. $R/\mathfrak{l}_{R,t}^s$ is a semiprime Goldie ring.
3. $Q(R/\mathfrak{l}_{R,t}^s)$ is a semisimple ring.
4. $Q_{l,cl}(R/\mathfrak{l}_R^s) \simeq Q_{r,cl}(R/\mathfrak{l}_R^s)$ is a semisimple ring.

If one of the equivalent conditions holds then

$$Q^s(R) \simeq Q_l(R/\mathfrak{l}_{R,t}^s) \simeq Q_{l,cl}(R/\mathfrak{l}_{R,t}^s) \simeq Q_{r,cl}(R/\mathfrak{l}_{R,t}^s),$$

$T(R) = \pi^{-1}(\mathcal{C}_{R/\mathfrak{l}_{R,t}^s})$ and $T(R/\mathfrak{l}_{R,t}^s) = \mathcal{C}_{R/\mathfrak{l}_{R,t}^s}$ where $\pi : R \rightarrow R/\mathfrak{l}_{R,t}^s$, $r \mapsto \bar{r} = r + \mathfrak{l}_{R,t}^s$, and $\mathcal{C}_{R/\mathfrak{l}_{R,t}^s}$ is the set of regular elements of the ring $R/\mathfrak{l}_{R,t}^s$.

Theorem 4.12 *Let $R = \prod_{i=1}^n R_i$ be a direct product of rings. Then*

1. $\mathcal{L}^s(R) = \prod_{i=1}^n \mathcal{L}^s(R_i)$.
2. $T(R) = \prod_{i=1}^n T(R_i)$ and $Q^s(R) \simeq \prod_{i=1}^n Q^s(R_i)$.

Theorem 4.13 *Let R be a ring. Then $S_{l,r,\mathfrak{a}}(R) \subseteq T(R)$ for all $\mathfrak{a} \in \text{Ass}(R)$ with $\mathfrak{a} \subseteq \mathfrak{l}_{R,t}^s$.*

5 Examples

In this section, the strong left quotient ring $Q_l^s(R)$, the strong left localization radical \mathfrak{l}_R^s and the largest strong left denominator set $T_l(R)$ are explicitly found for the following classes of rings: semiprime left Goldie rings (Theorem 5.1); rings of $n \times n$ lower/upper triangular matrices with coefficients in a left Goldie domain (Theorem 5.2 and Theorem 5.3); left Artinian rings (Theorem 5.6), and rings with left Artinian left quotient ring (Theorem 5.7).

Semiprime left Goldie rings. For each semiprime left Goldie ring R , the theorem below describes its largest strong left quotient ring $Q_l^s(R)$, \mathfrak{l}_R^s and $T_l(R)$.

Theorem 5.1 *Let R be a semiprime left Goldie ring. Then*

1. $Q_l^s(R) = Q_{l,cl}(R)$.
2. $\mathfrak{l}_R^s = 0$.
3. $T_l(R) = \mathcal{C}_R$.

Proof. The statements follow from Goldie's Theorem and Theorem 1.4. \square

Rings of lower/upper triangular matrices with coefficients in a left Goldie domain.

Let R be a left Goldie domain, $D := Q_{l,cl}(R)$ be its left quotient ring (it is a division ring), $L_n(R)$ and $U_n(D)$ be respectively the ring of lower and upper triangular matrices with coefficients in R and D . There are natural inclusions $R \subseteq L_n(R) \subseteq L_n(D)$ (each element $r \in R$ is identified with the diagonal matrix where all the diagonal elements are equal to r). Then $\mathcal{C}_R = R \setminus \{0\} \in \text{Den}_l(L_n(R), 0)$ with $\mathcal{C}_R^{-1}L_n(R) \simeq L_n(D)$. Hence, $\mathcal{C}_R \subseteq S_l(L_n(R))$ and $Q_{l,cl}(L_n(R)) \simeq L_n(D)$ since $Q_{l,cl}(L_n(D)) = L_n(D)$. Let E_{ij} ($i, j = 1, \dots, n$) be the matrix units. Every element $a = (a_{ij}) \in L_n(R)$ is a unique sum $a = \sum_{1 \leq i, j \leq n} a_{ij} E_{ij}$ where $a_{ij} \in R$.

Theorem 5.2 *Let R be a left Goldie domain. Then*

1. $\max.\text{Den}_l(L_n(R)) = \{T_l(R)\}$ and $T_l(R) = \{a = (a_{ij}) \in L_n(R) \mid a_{11} \neq 0\}$.
2. $\mathfrak{l}_{L_n(R)}^s = \{a = (a_{ij}) \in L_n(R) \mid a_{11} = 0\}$.
3. $Q_l^s(L_n(R)) = Q_{l,cl}(R)$.

Proof. Briefly, the statements follow from Proposition 3.3 and the following three facts

(i) $\max.\text{Den}_l(L_n(D)) = \{T_{E_{11}}\}$ where $T_{E_{11}} = \{a = (a_{ij}) \in L_n(D) \mid a_{11} \neq 0\}$, [5, Lemma 7.11.(2)],

(ii) $T_{E_{11}}^{-1}L_n(D) \simeq D$ [5, Lemma 7.11.(2)],

(iii) $\text{ass}(T_{E_{11}}) = (1 - E_{11})L_n(D) = \{a = (a_{ij}) \in L_n(D) \mid a_{11} = 0\}$, [5, Lemma 7.11.(3)].

In more detail, statement 1 follows from Proposition 3.3.(3) and the statement (i).

The inclusion $T_l(R) \subseteq T_{E_{11}}$ implies the inclusion

$$\mathfrak{l}_R^s = \text{ass}(T_l(R)) \subseteq \text{ass}(T_{E_{11}}) \cap L_n(R) = \mathfrak{a} := \{a = (a_{ij}) \in L_n(R) \mid a_{11} = 0\},$$

by the statement (iii). Since $E_{11} \in T_l(R)$ and $E_{11}\mathfrak{a} = 0$, we have the opposite inclusion $\mathfrak{l}_R^s \supseteq \mathfrak{a}$, i.e. $\mathfrak{l}_R^s = \mathfrak{a}$. This finishes the proof of statement 2.

Statement 3 follows from the statement (ii) and Proposition 3.3.(4):

$$Q_l^s(R) = T_l(R)^{-1}L_n(R) \simeq T_{E_{11}}^{-1}L_n(Q) \simeq D \simeq Q_{l,cl}(R). \quad \square$$

Let $U_n(R)$ be the ring of $n \times n$ upper triangular matrices with coefficients in R .

Theorem 5.3 *Let R be a left Goldie domain. Then*

1. $\max.\text{Den}_l(U_n(R)) = \{T_l(R)\}$ and $T_l(R) = \{a = (a_{ij}) \in U_n(R) \mid a_{nn} \neq 0\}$.
2. $\mathfrak{l}_{U_n(R)}^s = \{a = (a_{ij}) \in U_n(R) \mid a_{nn} = 0\}$.
3. $Q_l^s(U_n(R)) = Q_{l,cl}(R)$.

Proof. The theorem follows at once from Theorem 5.2 and the fact that the R -homomorphism

$$U_n(R) \rightarrow L_n(R), \quad E_{ij} \mapsto E_{n+1-i, n+1-j},$$

is a ring isomorphism. \square

Left Artinian rings. Before giving a proof of Theorem 5.6, let us introduce notation and cite two results from [5]. Let R be a left Artinian ring, $\text{rad}(R)$ be its radical, $\overline{R} := R/\text{rad}(R) = \prod_{i=1}^s \overline{R}_i$ is a direct product of simple Artinian rings \overline{R}_i , $\overline{1}_i$ be the identity element of the ring \overline{R}_i . So, $1 = \sum_{i=1}^s \overline{1}_i$ is the sum of orthogonal central idempotents of \overline{R} , $1 = \sum_{i=1}^s 1_i$ is a sum of orthogonal idempotents of R such that 1_i is a lifting of $\overline{1}_i$. For each non-empty set I of $\{1, \dots, s\}$, let $e_I := \sum_{i \in I} 1_i$,

$$\mathcal{I}'_I := \mathcal{I}'_I(R) := \{e_I \mid e_I R(1 - e_I) = 0\}.$$

The finite set \mathcal{I}'_I is a partially ordered set where $e_I \leq e_J$ if $I \subseteq J$.

Proposition 5.4 [5, Corollary 4.14] *Let R be a left Artinian ring and $e := \sum_{e' \in \min \mathcal{I}'_I(R)} e'$. Then*

1. $S_e := \{1, e\} \in \text{Den}_l(R, (1 - e)R)$.
2. $\text{ass}(S_e) = \mathfrak{l}_R$.
3. e is the least upper bound of the set $\min \mathcal{I}'_I(R)$ in $\mathcal{I}'_I(R)$.

The next theorem provides a description of the maximal left denominator sets of a left Artinian ring.

Theorem 5.5 [5, Theorem 4.10] *Let R be a left Artinian ring. Then*

1. $\max.\text{Den}_l(R) = \{T_e \mid e \in \min \mathcal{I}'_l(R)\}$ where $T_e = \{u \in R \mid u + (1 - e)R \in (R/(1 - e)R)^*\}$.
2. $|\max.\text{Den}_l(R)| \leq s$ (s is the number of isomorphism classes of left simple R -modules).
3. $|\max.\text{Den}_l(R)| = s$ iff R is a semisimple ring.

The next theorem explicitly describes the triple $T_l(R)$, \mathfrak{I}_R^s , $Q_l^s(R)$ for all left Artinian rings R .

Theorem 5.6 *Let R be a left Artinian ring and $e = \sum_{e' \in \min \mathcal{I}'_l(R)} e'$. Then*

1. $T_l(R) = \bigcap_{S \in \max.\text{Den}_l(R)} S = \{u \in R \mid u + (1 - e')R \in (R/(1 - e')R)^* \text{ for all } e' \in \min \mathcal{I}'_l(R)\}$.
2. $\mathfrak{I}_R^s = \mathfrak{I}_R = (1 - e)R$.
3. $Q_l^s(R) = R/\mathfrak{I}_R^s \simeq R/(1 - e)R \simeq \prod_{e' \in \min \mathcal{I}'_l(R)} R/(1 - e')R \simeq \prod_{S \in \max.\text{Den}_l(R)} S^{-1}R$.

Proof. 2. By Theorem 5.5.(1), $e \in \mathcal{L}_l^s(R)$. By Proposition 5.4.(1), $S_e = \{1, e\} \in \text{Den}_l(R)$, and so $S_e \subseteq T_l(R)$, by the maximality of $T_l(R)$. Notice that $\mathfrak{I}_R^s \subseteq \mathfrak{I}_R$ (Lemma 2.9.(2)) and $\text{ass}(S_e) = \mathfrak{I}_R$ (Proposition 5.4.(2)). Now,

$$\mathfrak{I}_R \supseteq \mathfrak{I}_R^s = \text{ass } T_l(R) \supseteq \text{ass}(S_e) = (1 - e)R \stackrel{\text{Pr. 5.4.(2)}}{=} \mathfrak{I}_R,$$

i.e. $\mathfrak{I}_R^s = \mathfrak{I}_R = (1 - e)R$.

3. By Theorem 1.1.(2) and statement 2,

$$Q_l^s(R) = Q_l(R/\mathfrak{I}_R^s) = R/\mathfrak{I}_R^s = R/(1 - e)R.$$

The isomorphism $R/(1 - e)R \simeq \prod_{e' \in \min \mathcal{I}'_l(R)} R/(1 - e')R$ follows from the decomposition [5, Equality (21)]. It remains to notice that $R/(1 - e')R \simeq T_{e'}^{-1}R$ and $\max.\text{Den}_l(R) = \{T_{e'} \mid e' \in \min \mathcal{I}'_l(R)\}$ (Theorem 5.5.(1)).

1. Let $\pi : R \rightarrow R/\mathfrak{I}_R^s$, $r \mapsto r + \mathfrak{I}_R^s$. Since the group of units U of the ring R/\mathfrak{I}_R^s is a left denominator set of $R/\mathfrak{I}_R^s \simeq T_l(R)^{-1}R$ and $\text{ass}(T_l(R)) = \mathfrak{I}_R^s$, the pre-image

$$\pi^{-1}(U) = \{u \in R \mid u \in (1 - e')R \in (R/(1 - e')R)^* \text{ for all } e' \in \min \mathcal{I}'_l(R)\} = \bigcap_{S \in \max.\text{Den}_l(R)} S$$

(Theorem 5.5.(1)) belongs to $\text{Den}_l(R, \mathfrak{I}_R^s)$ (Lemma 2.11), hence $T_l(R) = \pi^{-1}(U)$. \square

Rings with left Artinian left quotient ring. Let A be a ring such that $R := Q_{l,cl}(A)$ is a left Artinian ring. We keep the notation of the previous subsection.

Theorem 5.7 *Let A be a ring such that $R = Q_{l,cl}(A)$ is a left Artinian ring and $e = \sum_{e' \in \min \mathcal{I}'_l(R)} e'$. Then*

1. $T_l(A) = \bigcap_{S' \in \max.\text{Den}_l(R)} S'$.
2. $\mathfrak{I}_A^s = A \cap \mathfrak{I}_R^s = A \cap \mathfrak{I}_R = A \cap (1 - e)R$.
3. $Q_l^s(A) \simeq Q_l^s(R) = R/\mathfrak{I}_R^s \simeq R/(1 - e)R \simeq \prod_{e' \in \min \mathcal{I}'_l(R)} R/(1 - e')R \simeq \prod_{S \in \max.\text{Den}_l(R)} S^{-1}R \simeq \prod_{S' \in \max.\text{Den}_l(A)} S'^{-1}A$.

Proof. The theorem follows from Theorem 1.6, Theorem 5.6 and Proposition 3.3.(4). \square

Acknowledgements

The work is partly supported by the Royal Society and EPSRC.

References

- [1] V. V. Bavula, The algebra of integro-differential operators on an affine line and its modules, *J. Pure Appl. Algebra*, **217** (2013) 495-529. (Arxiv:math.RA: 1011.2997).
- [2] V. V. Bavula, Characterizations of left orders in left Artinian rings. *Journal of Alg. and its Appl.*, **13** (2014) no 6, 1-21. (Arxiv:math.RA:1212.3529).
- [3] V. V. Bavula, The largest left quotient ring of a ring, *Comm. in Algebra*, to appear. Arxiv:math.RA:1101.5107.
- [4] V. V. Bavula, New criteria for a ring to have a semisimple left quotient ring, *Journal of Alg. and its Appl.*, **6** (2015) no. 6, [28 pages] DOI: 10.1142/S0219498815500905. Arxiv:math.RA:1303.0859.
- [5] V. V. Bavula, Left localizations of left Artinian rings, Arxiv:math.RA:1405.0214.
- [6] A. W. Goldie, The structure of prime rings under ascending chain conditions. *Proc. London Math. Soc.* (3) **8** (1958) 589–608.
- [7] A. W. Goldie, Semi-prime rings with maximum condition. *Proc. London Math. Soc.* (3) **10** (1960) 201–220.
- [8] A. V. Jategaonkar, Localization in Noetherian Rings, Lndon Math. Soc. LMS 98, Cambridge Univ. Press, 1986.
- [9] C. R. Hajarnavis, On Small's Theorem, *J. London Math. Soc.* (2) **5** (1972) 596–600.
- [10] L. Lesieur and R. Croisot, Sur les anneaux premiers noethériens à gauche. *Ann. Sci. École Norm. Sup.* (3) **76** (1959) 161-183.
- [11] J. C. McConnell and J. C. Robson, Noncommutative Noetherian rings. With the cooperation of L. W. Small. Revised edition. Graduate Studies in Mathematics, 30. American Mathematical Society, Providence, RI, 2001. 636 pp.
- [12] J. C. Robson, Artinian quotient rings. *Proc. London Math. Soc.* (3) **17** (1967) 600–616.
- [13] L. W. Small, Orders in Artinian rings. *J. Algebra* **4** (1966) 13–41.
- [14] L. W. Small, Correction and addendum: Orders in Artinian rings. *J. Algebra* **4** (1966) 505–507.
- [15] B. Stenström, Rings of Quotients, Springer-Verlag, Berlin, Heidelberg, New York, 1975.
- [16] H. Tachikawa, Localization and Artinian Quotient Rings, *Math. Z.* **119** (1971) 239–253.
- [17] T. D. Talintyre, Quotient rings with minimum condition on right ideals, *J. London Math. Soc.* **41** (1966) 141–144.

Department of Pure Mathematics
University of Sheffield
Hicks Building
Sheffield S3 7RH
UK
email: v.bavula@sheffield.ac.uk