



# The LDL1/2-HDA6 Histone Modification Complex Interacts With TOC1 and Regulates the Core Circadian Clock Components in *Arabidopsis*

Fu-Yu Hung<sup>1,2</sup>, Fang-Fang Chen<sup>1</sup>, Chenlong Li<sup>2,3,4</sup>, Chen Chen<sup>2,3</sup>, Jian-Hao Chen<sup>1</sup>, Yuhai Cui<sup>2,3\*</sup> and Keqiang Wu<sup>1\*</sup>

<sup>1</sup> Institute of Plant Biology, National Taiwan University, Taipei, Taiwan, <sup>2</sup> Agriculture and Agri-Food Canada, London Research and Development Centre, London, ON, Canada, <sup>3</sup> Department of Biology, Western University, London, ON, Canada, <sup>4</sup> State Key Laboratory of Biocontrol and Guangdong Key Laboratory of Plant Resource, School of Life Sciences, Sun Yat-sen University, Guangzhou, China

## OPEN ACCESS

### Edited by:

Jean-Benoit Charron,  
McGill University, Canada

### Reviewed by:

R. Glen Uhrig,  
University of Alberta, Canada  
Jun Xiao,  
Institute of Genetics  
and Developmental Biology (CAS),  
China

### \*Correspondence:

Yuhai Cui  
Yuhai.cui@canada.ca  
Keqiang Wu  
kewu@ntu.edu.tw

### Specialty section:

This article was submitted to  
Plant Cell Biology,  
a section of the journal  
Frontiers in Plant Science

Received: 25 October 2018

Accepted: 12 February 2019

Published: 26 February 2019

### Citation:

Hung F-Y, Chen F-F, Li C, Chen C, Chen J-H, Cui Y and Wu K (2019) The LDL1/2-HDA6 Histone Modification Complex Interacts With TOC1 and Regulates the Core Circadian Clock Components in *Arabidopsis*. *Front. Plant Sci.* 10:233. doi: 10.3389/fpls.2019.00233

In *Arabidopsis*, the circadian rhythm is associated with multiple important biological processes and maintained by multiple interconnected loops that generate robust rhythms. The circadian clock central loop is a negative feedback loop composed of the core circadian clock components. *TOC1* (*TIMING OF CAB EXPRESSION 1*) is highly expressed in the evening and negatively regulates the expression of *CCA1* (*CIRCADIAN CLOCK ASSOCIATED 1*)/*LHY* (*LATE ELONGATED HYPOCOTYL*). *CCA1/LHY* also binds to the promoter of *TOC1* and represses the *TOC1* expression. Our recent research revealed that the histone modification complex comprising of *LYSINE-SPECIFIC DEMETHYLASE 1* (*LSD1*)-LIKE 1/2 (*LDL1/2*) and *HISTONE DEACETYLASE 6* (*HDA6*) can be recruited by *CCA1/LHY* to repress *TOC1* expression. In this study, we found that *HDA6*, *LDL1*, and *LDL2* can interact with *TOC1*, and the *LDL1/2-HDA6* complex is associate with *TOC1* to repress the *CCA1/LHY* expression. Furthermore, *LDL1/2-HDA6* and *TOC1* co-target a subset of genes involved in the circadian rhythm. Collectively, our results indicate that the *LDL1/2-HDA6* histone modification complex is important for the regulation of the core circadian clock components.

**Keywords:** H3K4 demethylases, HDA6, circadian clock, CCA1/LHY, TOC1, *Arabidopsis*

## INTRODUCTION

The circadian rhythm is an endogenous oscillation widely observed in plants, animals, fungi, and cyanobacteria (Edgar et al., 2012). The plant circadian rhythm is highly associated with multiple important biological processes, and maintained by multiple interconnected loops that generate robust rhythms. The circadian clock central loop is a negative feedback loop composed of the core circadian clock components such as *TOC1* (*TIMING OF CAB EXPRESSION 1*) and *CCA1* (*CIRCADIAN CLOCK ASSOCIATED 1*)/*LHY* (*LATE ELONGATED HYPOCOTYL*). *TOC1* is highly expressed in the evening, but low expressed at dawn (Alabadi et al., 2001). Furthermore, *TOC1*

was identified as a repressor of *CCA1* and *LHY* by binding to their promoters in the evening (Gendron et al., 2012; Huang et al., 2012). In contrast, *CCA1* and *LHY* are highly expressed in the morning, but low expressed at nightfall (Schaffer et al., 1998; Wang and Tobin, 1998; Alabadi et al., 2001). *CCA1* and *LHY* bind to the evening element (EE) on the promoter of *TOC1* to inhibit its expression (Schaffer et al., 1998; Wang and Tobin, 1998; Alabadi et al., 2001; Nagel et al., 2015). *CHE* (*CCA1* HIKING EXPEDITION) is an evening-expressed TCP-family transcription factor, which also targets the *CCA1* promoter to repress its expression. Furthermore, *CCA1* and *LHY* were shown to repress the *CHE* expression by targeting the *CHE* promoter (Pruneda-Paz et al., 2009).

Histone modifications play important roles in the regulation of gene expression. Histone methyltransferases and demethylases determine the methylation levels, whereas histone acetylation levels are regulated by histone acetyltransferases (HATs) and histone deacetylases (HDACs or HDAs). HDACs and the H3K4 demethylase LSD1 (Lysine-Specific Demethylase 1) are the core components of the Mi2/NuRD and CoREST protein complexes in yeast and animal cells (Khochbin et al., 2001; Lee et al., 2005; Wang et al., 2009). They act co-operatively to repress gene expression in mammals (Huang et al., 2011). The interactions among the core protein components of the HDAC complexes are relatively stable and the HDAC complexes can also interact with various transcription factors under different environmental conditions (Joshi et al., 2013; Liu et al., 2014). FLD (FLOWERING LOCUS D), LDL1 (Lysine-Specific Demethylase-LIKE 1), LDL2, and LDL3 are the LSD1 homologs in *Arabidopsis* (Jiang et al., 2007). LDL1 and LDL2 act redundantly to regulate *FLC* (FLOWERING LOCUS C) by H3K4 demethylation (Jiang et al., 2007). Furthermore, *Arabidopsis* HISTONE DEACETYLASE 6 (HDA6) directly interacts with FLD to repress *FLC*, *MAF4*, and *MAF5* by reducing H3K4 methylation (H3K4me) and H3 acetylation (H3Ac) to regulate flowering time (Yu et al., 2011). In addition, HDA6 can also interact with LDL1 and LDL2 to regulate gene expression (Hung et al., 2018).

The HDAC inhibitor TSA treated plants show delayed phases and higher amplitudes of *TOC1* expression (Perales and Más, 2007). In addition, the expression of *Arabidopsis* *CCA1*, *LHY*, and *TOC1* is specifically associated with H3Ac and H3K4me changes (Hemmes et al., 2012; Malapeira et al., 2012), indicating that the expression of the core circadian clock components is associated with H3Ac and H3K4me level changes. Our recent study indicated that *CCA1* and *LHY* can interact with the HDAC complex containing LDL1, LDL2, and HDA6. Furthermore, the LDL1/2-HDA6 complex can be recruited by the transcription repressors *CCA1* and *LHY* to their target genes including *TOC1*. Since *CCA1* and *LHY* are low expressed at nightfall, the expression of *TOC1* is increased due to the release of LDL1/2-HDA6 from the *TOC1* promoter (Hung et al., 2018). In this study, we demonstrated that LDL1/2-HDA6 can also interact with *TOC1* to regulate the expression of *CCA1* and *LHY*. Furthermore, LDL1/2-HDA6 and *TOC1* co-target a subset of genes involved in the circadian rhythm.

## MATERIALS AND METHODS

### Plant Materials and Growth Conditions

The *Arabidopsis thaliana* Columbia (Col-0) ecotype was used. Plants were grown at 22°C under 12/12 h light/dark conditions in growth chambers. The mutants used in this study were previously described, including *ldl1/ldl2* (Jiang et al., 2007), *hda6* (*axe1-5*) (Yu et al., 2011), *hda6/ldl1/2* (Hung et al., 2018), *toc1*, and *cca1/lhy* (Wang et al., 2011). *35Spro::LDL1:GFP*, *35Spro::GFP:HDA6*, *LDL1pro::LDL1:GFP* and *HDA6pro::HDA6:GFP* transgenic plants were previously described (Yu et al., 2011; Hung et al., 2018).

The full-length coding sequence (CDS) fragment of *TOC1* was PCR-amplified and cloned into the *pCR8/GW/TOPO* vector (Invitrogen), and then recombined into the *PK7WGF2* binary vector or *3xFLAG* Gateway vector (Invitrogen<sup>1</sup>). The *35S::TOC1:GFP* vector was transformed into Col-0 WT or *hda6/ldl1/2* by the floral dip method.

### Bimolecular Fluorescence Complementation (BiFC) Assays

To generate the constructs for BiFC assays, the full-length coding sequence (CDS) fragment of *TOC1* was amplified by PCR and cloned into the *pCR8/GW/TOPO* vector, and then recombined into the *pEarleyGate201-YN* (Lu et al., 2010). *LDL1-YC* and *HDA6-YC* were described in the previous studies (Yu et al., 2011; Hung et al., 2018). Constructed vectors were transformed into *Arabidopsis* protoplasts or tobacco (*Nicotiana benthamiana*) leaves for transient assays. Transformed protoplasts and tobacco leaves were then examined by confocal spectral microscope imaging system (NTU-TCS SP5, Leica<sup>2</sup>).



### Yeast Two-Hybrid (Y2H) Assays and Co-immunoprecipitation (Co-IP) Assays

Yeast two-hybrid assays were performed based on the instruction for the Matchmaker GAL4-based two-hybrid system 3 (Clontech). The *LDL1*, *LDL2*, and *TOC1* full length cDNA fragments were sub-cloned into *pGADT7* and *pGBKT7* vectors. All constructs were transformed into the yeast (*Saccharomyces cerevisiae*) strain AH109 by the lithium acetate method, and yeast cells were grown on a minimal medium/-Leu-Trp according to the manufacturer's instructions (Clontech). Transformed colonies were grown on the medium containing X- $\alpha$ -gal for the  $\alpha$ -galactosidase activity assay or minimal medium/-Leu-Trp-His (3DO) with 0.25 mM 3-amino-1,2,4-triazole (3AT).

Co-immunoprecipitation assays were performed as previously described (Yu et al., 2011). The *35S::TOC1:3xFLAG* plasmid was transformed into *Arabidopsis* protoplasts extracted from *LDL1pro::LDL1:GFP* or *35Spro::GFP* transgenic plants. Total proteins were then extracted from the transformed protoplasts. Anti-GFP (Santa Cruz Biotechnologies, catalog no. SC-9996; 1:3000 dilution) and anti-FLAG (SIGMA catalog no. M2; 1:3000

<sup>1</sup><https://www.psb.ugent.be/core-facilities/380-gateway-vectors>

<sup>2</sup><https://www.leica-microsystems.com/products/confocal-microscopes/p/leica-tcs-sp5/>



dilution) antibodies were used as primary antibodies for Western blot. The resulting signals were detected by using a Pierce ECL Western blotting kit (Pierce<sup>3</sup>).

## Quantitative Real-Time PCR (qRT-PCR) Analysis

The TRIZOL reagent (Invitrogen, 15596026) was used for total RNA isolation according to the manufacturer's instructions. Total RNA treated with 2 µg of DNase (Promega, RQ1 #M6101) were then used for cDNA synthesis (Promega, #1012891). The iQ SYBR Green Supermix solution (Bio-Rad, #170-8880) was used for real-time quantitative PCR assays with the CFX96 real-time PCR Detection System (Bio-Rad Laboratories, Inc.). Cycling conditions were started with 95°C/10 min, followed by 45 cycles of 95°C/15 s, 60°C/30 s, and then fluorescent detection, and melting curve detection (65–95°C, incrementing 0.5°C for 5 s, and plate reading). Each sample was normalized by calculating delta quantification cycle (Cq) to the expression of the *UBQ10* (*Ubiquitin10*) internal control and quantified at least in triplicate. The Cq and relative expression level are calculated by the Biorad CFX Manager 3.1 based on the MIQE guidelines (Bustin et al., 2009). **Supplementary Table S1** listed the gene specific primers used for qRT-PCR. Standard deviations (SD) represent at least three technical and three biological replicates. The variance in average data is represented by standard error of the mean (SEM). The SD, SEM determination and *P*-value were calculated using Student's paired *t*-test.

## Protoplast Transient Assays

The *CCA1pro::LUC* plasmid construct was previously described (Wang et al., 2011). For transcriptional activity assays, the *35Spro::TOC1*, *35Spro::LDL1*, *35Spro::HDA6*, or *35Spro::GFP* effector constructs were co-transformed into protoplasts with *CCA1pro::LUC*, and the plant samples were collected at ZT0 after 12 h. The relative activities of LUC (luciferase) reporter were standardized by activities of co-expressed Renilla LUC. Experiments were repeated at least three times for each reporter-effector combination. The dual luciferase assay reagent (Promega) was used for Firefly LUC and Renilla LUC detection.

## Chromatin Immunoprecipitation (ChIP) Assays and ChIP-seq Data Analyses

Chromatin immunoprecipitation assays were accomplished as previously described (Yu et al., 2011; Hung et al., 2018). Plant seedlings were treated with 1% formaldehyde for chromatin extraction. The extracted DNA was sheared to the mean length near 500 bp by sonication, proteins, and DNA fragments were then immunoprecipitated by the H3K9K14 (Millipore, catalog no. 06-599), H3K4me3 (Millipore, catalog no. 04-745), or GFP (Abcam, catalog no. ab290) antibodies. The cross-link between DNA with immunoprecipitated proteins were reversed, and then analyzed by real-time PCR using specific primers (**Supplementary Table S1**). The quantification cycle (Cq) was

calculated by Biorad CFX Manager 3.1 based on the MIQE guideline (Bustin et al., 2009). Percent input was calculated as  $2^{[Cq(IN)-Cq(IP)]} \times 100$ . Each sample was quantified at least in triplicate, and normalized by calculating delta Cq to the expression of the internal control. Standard deviations (SD) represent at least three technical and three biological replicates. The variance in average data is represented by standard error of the mean (SEM). The SD, SEM determination and *P*-value were calculated using Student's paired *t*-test.

ChIP-seq assays were performed based on previous research (Li et al., 2015, 2016; Hung et al., 2018). The LDL1 ChIP-seq data were deposited to NCBI-Gene Expression Omnibus (GEO) database (GSE118025) (Hung et al., 2018). The ChIP-Seq files from other research groups, GSE35952 (Huang et al., 2012) and (Kamioka et al., 2016), were downloaded from the NCBI-GEO database.

## RESULTS

### LDL1 and HDA6 Interact With TOC1 and Directly Target on *CCA1* and *LHY*

Our recent study indicated that *CCA1/LHY* can interact with the LDL1/2-HDA6 complex to repress *TOC1* (Hung et al., 2018). In addition, the expression of *TOC1*, *CCA1* and *LHY* is also associated with H3K4me and H3 acetylation changes (Hemmes et al., 2012; Malapeira et al., 2012). We further analyzed the functional correlation between TOC1 and the LDL1/2-HDA6 complex. TOC1 directly interacted with both LDL1 and LDL2 in BiFC assays by using *Arabidopsis* protoplasts and *Agrobacterium*-infiltrated tobacco leaves. The YFP fluorescence signal was detected in nucleus of the transformed cells (**Figure 1A** and **Supplementary Figure S1**). The interaction between LDL1, LDL2, and TOC1 was further confirmed by yeast two-hybrid assays (**Figures 1B,C**) and Co-IP assays using *Arabidopsis* protoplasts (**Figure 1D** and **Supplementary Figure S1**). Furthermore, TOC1 can also interact with HDA6 in BiFC assays (**Supplementary Figures S2A,B**). These results suggested that TOC1 may recruit the LDL1/2-HDA6 histone modification complex to its target genes such as *CCA1* and *LHY*.

We further analyzed the binding of LDL1 and HDA6 to *CCA1* and *LHY* by ChIP assays. The *LDL1:GFP* and *HDA6:GFP* transgenic plants were previously described (Yu et al., 2011; Hung et al., 2018). 14 days old plants grown under 12 h light/12 h dark condition were collected on Zeitgeber time 0 (ZT0) and ZT12. An anti-GFP antibody was used for ChIP assays, and the binding of LDL1 and HDA6 was analyzed by qPCR. We identified that both LDL1 and HDA6 can bind to the promoters of *CCA1* and *LHY*. Furthermore, the binding of LDL1 and HDA6 to the promoters of *CCA1* and *LHY* were significantly decreased on ZT0 compared to ZT12 (**Figure 2** and **Supplementary Figure S2C**). The binding of LDL1 and HDA6 to the *CCA1* and *LHY* promoters is correlated to TOC1 accumulation, since *TOC1* is highly expressed at nightfall but low expressed in the morning (Alabadi et al., 2001).

<sup>3</sup><https://www.lifetechnologies.com/>

## TOC1 and LDL1 Co-target Genes Involved in the Circadian Rhythm


Previously, we identified the global binding sites of LDL1 by ChIP-Seq assays (Hung et al., 2018). The GO-BP (Gene Ontology\_Biological Process) analysis of LDL1-targeted genes revealed that LDL1 targets on a subset of circadian rhythm genes. Furthermore, LDL1 also binds to a cluster of circadian rhythm genes regulated by CCA1 (Hung et al., 2018). In this study, we further analyzed whether the LDL1 and TOC1 also co-target genes involved in the circadian rhythm.

We compared the previously published TOC1 ChIP-Seq data (Huang et al., 2012) with the LDL1 ChIP-Seq data (Hung et al., 2018). The genome browser views by Integrative Genomics Viewer (IGV) indicated that LDL1 bound to *CCA1* and *LHY*, and the binding peaks of LDL1 are highly correlated with the TOC1 binding regions on *CCA1* and *LHY* promoters (Figure 3A). Among 772 genes occupied by TOC1 (Huang et al., 2012), 195 of them are also co-occupied by LDL1 ( $P = 1.14 \times 10^{-16}$ ) (Figure 3B). Furthermore, the genomic binding regions of TOC1 are closed to the LDL1 binding regions (Figure 3C), indicating that TOC1 and LDL1 tend to bind to the similar genome

sites. GO-BP analysis also indicated that LDL1 and TOC1 co-target on a subgroup of genes involved in circadian rhythm and response to cold (Figure 3D). In GO-BP analysis, the ratio of the circadian genes of LDL1/TOC1 co-targeted genes is increased when compared to the LDL1-targeted genes or the TOC1-targeted genes alone (Supplementary Figure S3). Interestingly, the ratio of the circadian rhythm genes is further increased in the LDL1/CCA1/TOC1 co-targeted genes (Supplementary Figure S3). Previous studies indicated that several *cis*-elements are enriched in the promoters of TOC1 regulated genes, including the (AG/CT)<sub>n</sub> repeat, G-box (CACGTG), Evening Element (EE)-like and TCP binding site (TBS, GGCCCA) (Gendron et al., 2012; Huang et al., 2012). Similar *cis*-elements are also enriched in the LDL1-targeted promoter regions (Hung et al., 2018).

## LDL1/2-HDA6 Is Involved in the Regulation of CCA1/LHY

TOC1 is a repressor and targets on the promoters of *CCA1* and *LHY*. The expression of *CCA1* and *LHY* is decreased in *TOC1* over-expressing (*TOC1-OE*) plants (Gendron et al., 2012; Huang et al., 2012). Furthermore, additional *TOC1*




**FIGURE 3** | LDL1-occupied sites in the genome identified by ChIP-seq analysis. **(A)** Integrated genome view of LDL1 binding peaks on *CCA1* and *LHY*. green BARS indicate the TOC1-binding regions from previous published data (Huang et al., 2012). **(B)** Overlap between TOC1 target genes (Huang et al., 2012) and LDL1 targeted genes (Hung et al., 2018) (hypergeometric distribution of TOC1 and LDL1 co-targeted genes:  $p = 1.14 \times 10^{-16}$ ). **(C)** Distribution of distances between the total binding sites of LDL1 and TOC1. **(D)** GO-BP annotation of LDL1/TOC1 co-occupied genes. Annotation terms with  $p$ -value  $< 0.01$  were listed.


expression causes increased period length of *CCA1* (Mas et al., 2003a). To investigate the functional relationship between TOC1 and LDL1/2-HDA6, we generated *TOC1* over-expressing plants in WT (*TOC1-OE*) and the *hda6/ldl1/2* background (*TOC1-OE/hda6/ldl1/2*). The binary vector containing *CaMV 35S promoter* driven *GFP:TOC1* (*35S::GFP:TOC1*) was transformed into WT or *hda6/ldl1/2*. The expression patterns of *CCA1* and *LHY* were compared by qRT-PCR in wild-type (WT), *TOC1-OE* and *ldl1/2/hda6* plants grown under 12 h light/12 h dark for 14 days. As reported previously (Gendron et al., 2012; Huang et al., 2012), the expression of *CCA1* and *LHY* was decreased in *TOC1-OE* plants. However, the expression of *CCA1* and *LHY* was not significantly decreased in *hda6/ldl1/2* compared to WT (Figure 4A and Supplementary Figures S4A,B). Furthermore, the decrease of *CCA1* and *LHY* expression was recovered when *TOC1* was over-expressed in *hda6/ldl1/2* (Figure 4A and Supplementary Figures S4B,C). We also compared the daily expression patterns of *CCA1*, *LHY*, and *TOC1* in *ldl1/ldl2*, *hda6*, *hda6/ldl1/2*, and WT grown under 12 h light/12 h dark conditions. The expression of *CCA1* and *LHY* was not significantly decreased or shifted in *ldl1/ldl2*, *hda6*, and *hda6/ldl1/2* compared to WT (Supplementary Figure S4A). The expression patterns of other TOC1 targets such as *GI*, *PRR7* and *PRR9* in *ldl1/ldl2*, *hda6*, and *hda6/ldl1/2* were analyzed in

our previous study (Hung et al., 2018). *XTH27* and *AT1G10020* were previously identified to be the target genes regulated by TOC1 (Gendron et al., 2012; Huang et al., 2012), which are also targeted by LDL1 (Hung et al., 2018). The expression of *XTH27* and *AT1G10020* was increased in *ldl1/ldl2*, *hda6*, and *hda6/ldl1/2* compared to WT (Supplementary Figure S4C).

We further analyzed the functional correlation between LDL1, HDA6, and TOC1. *CCA1pro::CCA1:LUC* (*pCCA1:LUC*) was co-expressed with *35Spro::TOC1*, *35Spro::LDL1*, *35Spro::HDA6*, or *35Spro::GFP* in *Arabidopsis* protoplasts. Although the activity of *CCA1:LUC* was only slightly reduced when co-expressed with LDL1, and activity was further decreased when TOC1 was co-expressed with LDL1 (Figure 4B). Similar results were also observed when TOC1 was co-expressed with HDA6 (Figure 4C).

We also analyzed H3K4me and H3Ac levels of *CCA1* and *LHY* in WT, *TOC1-OE* plants and *hda6/ldl1/2*. For ChIP-qPCR assays, 14-days old plants grown under 12 h light/12 h dark conditions were collected on ZT0. H3K4me and H3Ac of *CCA1* and *LHY* were decreased in *TOC1-OE* plants (Figure 5), indicating that TOC1 affects the levels of H3K4me and H3Ac on *CCA1* and *LHY*. We further analyzed H3Ac and H3K4me levels of *CCA1* and *LHY* in 14 days old *hda6*, *ldl1/ldl2*, *hda6/ldl1/2*, and WT on ZT0 and ZT12. The H3Ac and H3K4me levels of *CCA1* and *LHY* were not decreased in *hda6*, *ldl1/ldl2*, *hda6/ldl1/2*





**FIGURE 5 |** LDL1/2-HDA6 is involved in regulation of H3Ac and H3K4me of *CCA1/LHY*. (A) Schematic diagram of *CCA1* and *LHY*. P: promoter region, E: coding region, U: 3' UTR. (B) ChIP analysis of H3ac and H3K4me levels of *CCA1* and *LHY* in *TOC1-OE* plants on ZT0. The amounts of DNA after ChIP were quantified by qRT-PCR and normalized to *ACT2*. Plants were grown under 12/12 : light/dark for 14 days. Data points represent the average of three technical replicates. Error bars correspond to SD from three biological replicates. \* $P < 0.05$ , \*\* $P < 0.005$  (Student's *t*-test).


(Supplementary Figure S4D). Interestingly, decreased H3K4me and H3Ac in *TOC1-OE* were recovered in *TOC1-OE/hda6/ldl1/2*, since the H3Ac and H3K4me levels of *CCA1* and *LHY* were significant higher in *TOC1-OE/hda6/ldl1/2* compared to the *TOC1-OE* plants (Figure 5). These results suggested that *TOC1* is involved in regulation of H3K4me and H3Ac on *CCA1* and *LHY*, and *TOC1* repressed *CCA1* and *LHY* expression is dependent on the function of LDL1/2-HDA6 complex.

## DISCUSSION

*Arabidopsis* HDA6 is a class I RPD3-like histone deacetylase associated with regulation of rRNA and transcription repression (Murfett et al., 2001; Probst et al., 2004; Earley et al., 2006; Liu et al., 2012; Yu et al., 2017). Different transcription factors can recruit HDA6 to regulate the gene expression involved in flowering, leaf development, abiotic stress response, and senescence (Wu et al., 2008; Chen et al., 2010; Yu et al., 2011; Luo et al., 2012; Liu et al., 2014). In animal and yeast cells, HDACs and LSD1 regulate gene expression cooperatively and they are both identified as the core components of Mi2/NuRD and CoREST

complexes (Khochbin et al., 2001; Lee et al., 2005; Wang et al., 2009). Our recent study demonstrated that the *Arabidopsis* H3K4 demethylases LDL1 and LDL2 can interact with HDA6 to repress gene expression (Hung et al., 2018). The LDL1/2-HDA6 complex can also interact with *CCA1/LHY* and reduce H3Ac and H3K4me levels of the circadian core component *TOC1* (Hung et al., 2018). Furthermore, a subset of genes involved in the circadian clock are co-targeted by LDL1 and *CCA1* (Hung et al., 2018).

*Arabidopsis* circadian clock genes are regulated by a complicate feedback regulation network forming multiple interconnected loops. The central loop is comprised of the core clock components, such as *TOC1* and *CCA1/LHY* (Schaffer et al., 1998; Wang and Tobin, 1998; Alabadi et al., 2001; Gendron et al., 2012; Huang et al., 2012; Nagel et al., 2015). The central loop is interlocked with the evening loop and morning loop. *PRR5*, *PRR7*, *PRR9*, and *CCA1/LHY* constitute the morning loop (Nakamichi et al., 2010; Salomé et al., 2010; Pokhilko et al., 2012), whereas *PRR3*, *GI*, *ZTL* (*ZEITLUPE*), and *TOC1* comprise the evening loop (Kim et al., 2003; Mas et al., 2003b; Para et al., 2007; McClung and Gutiérrez, 2010). We found that LDL1 and *CCA1* co-target to a subset of circadian genes, which are repressed by *CCA1* in the morning. However, LDL1 also targets to the morning expressed circadian genes, which may not be repressed by *CCA1* and *LHY* (Nagel et al., 2015; Kamioka et al., 2016; Hung et al., 2018). Although the binding of LDL1 on the LDL1/*CCA1* co-targeted genes are reduced in the *cca1/lhy* mutant, their binding is not completely abolished (Hung et al., 2018). These results suggested that in addition to *CCA1* and *LHY*, the LDL1/2-HDA6 complex may also functionally associate with other circadian clock genes. EC (Evening Complex) is



**FIGURE 6 |** A model for LDL1/2 and HDA6 functions in the regulation of core circadian clock components. Both morning accumulated *CCA1/LHY* and evening accumulated *TOC1* interact with the same histone modification complex containing LDL1/2 and HDA6. *CCA1/LHY* act as transcription repressors and recruit the histone modification complex to their target loci such as *TOC1* in the morning. Meanwhile, *TOC1* also recruits the histone modification complex to its targets such as *CCA1* and *LHY* in the evening.

also associated with regulation of the circadian genes, which is comprised of LUX (LUX ARRHYTHMO), ELF3 (EARLY FLOWERING3), and ELF4 (EARLY FLOWERING4) (Hazen et al., 2005; Nusinow et al., 2011). A previous study indicated that *Arabidopsis* HDACs are associated with PRR9 through direct interacting with TPL/TPR (TOPLESS/TOPLESS-RELATED) to regulate the expression of *CCA1* (Wang et al., 2013). Further research is required to investigate the functional correlation among LDL1/2-HDA6, PRR9, and EC.

The central loop of *Arabidopsis* circadian clock is consisted of the core clock components including *CCA1*, *LHY*, and *TOC1* (Schaffer et al., 1998; Wang and Tobin, 1998; Alabadi et al., 2001; Gendron et al., 2012; Huang et al., 2012; Nagel et al., 2015). Although *CCA1* and *LHY* are low expressed at nightfall, they are highly induced at dawn (Schaffer et al., 1998; Wang and Tobin, 1998; Alabadi et al., 2001). Previously, we found that *CCA1* interacts with *LDL1* in the morning (Hung et al., 2018). The binding of *LDL1* and *HDA6* on promoter of *TOC1* is higher in the morning but decreased in the evening (Hung et al., 2018). Furthermore, *HDA6*, *LDL1*, and *LDL2* are constitutively expressed at different time periods. *CCA1/LHY* can therefore recruit the *LDL1/2-HDA6* complex to suppress *TOC1* expression at dawn (Hung et al., 2018). In this study, we found that *LDL1/2* and *HDA6* also interact with *TOC1* to regulate the expression of *CCA1* and *LHY*. In consistent with the fact that *TOC1* is highly accumulated at nightfall (Alabadi et al., 2001), we found that the binding of *LDL1* and *HDA6* on the *CCA1* and *LHY* promoters is higher in the evening but decreased in the morning. Since *TOC1* is a repressor of *CCA1* and *LHY*, the expression of *CCA1* and *LHY* is decreased with increased *TOC1* expression (Gendron et al., 2012; Huang et al., 2012). We found that histone acetylation and H3K4 methylation levels of *CCA1* and *LHY* are decreased in *TOC1-OE* plants. However, the H3Ac, H3K4me and expression levels of *CCA1* and *LHY* are significantly increased in *TOC1-OE/hda6/ldl1/2* compared to the *TOC1-OE* plants, indicating that the *LDL1/2-HDA6* complex is functionally associated with the regulation of *CCA1* and *LHY* expression. Although the expression of *TOC1* is highly increased in *hda6/ldl1/2* compared to wild type, the expression of *CCA1* and *LHY* is not decreased. It is possible that in addition to *LDL1/2-HDA6*, other unknown proteins may also be involved in the regulation of *CCA1* and *LHY* expression.

Collectively, we propose a model to demonstrate how the core circadian clock components are regulated by H3K4 demethylation and histone deacetylation (Figure 6). The histone modification complex containing *LDL1/2* and *HDA6* can interact

with both morning accumulated *CCA1/LHY* (Hung et al., 2018) and evening accumulated *TOC1*. The transcription repressors *CCA1* and *LHY* can recruit the *LDL1/2-HDA6* complex to their target loci including *TOC1* in the morning (Hung et al., 2018). Furthermore, *TOC1* can also recruit the histone modification complex to its targets such as *CCA1* and *LHY* in the evening.

## DATA AVAILABILITY

All datasets generated for this study are included in the manuscript and/or the **Supplementary Files**.

## AUTHOR CONTRIBUTIONS

F-YH, KW, and YC designed the research. F-YH, F-FC, and J-HC performed the research. F-YH, CL, CC, and KW analyzed the data. F-YH and KW wrote the article.

## FUNDING

This work was supported by the Ministry of Science and Technology of the Republic of China (105-2311-B-002-012-MY3 and 106-2313-B-002-003- to KW) and National Taiwan University (106R891501 and 107L893101 to KW). This work was also supported by funding from the Agriculture and Agri-Food Canada A-base and the National Science and Engineering Research Council of Canada (RGPIN/04625-2017 to YC).

## ACKNOWLEDGMENTS

We thank Prof. S.-H. Wu. and J.-F. Wu (Academia Sinica) for sharing the *toc1-101* mutant and *CCA1pro::LUC* plasmid construct. We are grateful to the Technology Commons, College of Life Science, National Taiwan University for the convenient use of the Bio-Rad real-time PCR system, the confocal spectral microscope imaging, and Delta-vision systems.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2019.00233/full#supplementary-material>

## REFERENCES

- Alabadi, D., Oyama, T., Yanovsky, M. J., Harmon, F. G., Mas, P., and Kay, S. A. (2001). Reciprocal regulation between *TOC1* and *LHY/CCA1* within the *Arabidopsis* circadian clock. *Science* 293, 880–883. doi: 10.1126/science.1061320
- Bustin, S. A., Benes, V., Garson, J. A., Hellems, J., Huggett, J., Kubista, M., et al. (2009). The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin. Chem.* 55, 611–622. doi: 10.1373/clinchem.2008.112797
- Chen, L. T., Luo, M., Wang, Y. Y., and Wu, K. (2010). Involvement of *Arabidopsis* histone deacetylase *HDA6* in ABA and salt stress response. *J. Exp. Bot.* 61, 3345–3353. doi: 10.1093/jxb/erq154
- Earley, K., Lawrence, R. J., Pontes, O., Reuther, R., Enciso, A. J., Silva, M., et al. (2006). Erasure of histone acetylation by *Arabidopsis* *HDA6* mediates large-scale gene silencing in nucleolar dominance. *Genes Dev.* 20, 1283–1293. doi: 10.1101/gad.1417706
- Edgar, R. S., Green, E. W., Zhao, Y., Van Ooijen, G., Olmedo, M., Qin, X., et al. (2012). Peroxiredoxins are conserved markers of circadian rhythms. *Nature* 485, 459–464. doi: 10.1038/nature11088



- Gendron, J. M., Pruneda-Paz, J. L., Doherty, C. J., Gross, A. M., Kang, S. E., and Kay, S. A. (2012). *Arabidopsis* circadian clock protein, TOC1, is a DNA-binding transcription factor. *Proc. Natl. Acad. Sci. U.S.A.* 109, 3167–3172. doi: 10.1073/pnas.1200355109
- Hazen, S. P., Schultz, T. F., Pruneda-Paz, J. L., Borevitz, J. O., Ecker, J. R., and Kay, S. A. (2005). LUX ARRHYTHMO encodes a Myb domain protein essential for circadian rhythms. *Proc. Natl. Acad. Sci. U.S.A.* 102, 10387–10392. doi: 10.1073/pnas.0503029102
- Hemmes, H., Henriques, R., Jang, I. C., Kim, S., and Chua, N. H. (2012). Circadian clock regulates dynamic chromatin modifications associated with *Arabidopsis* CCA1/LHY and TOC1 transcriptional rhythms. *Plant Cell Physiol.* 53, 2016–2029. doi: 10.1093/pcp/pcs148
- Huang, P. H., Chen, C. H., Chou, C. C., Sargeant, A. M., Kulp, S. K., Teng, C. M., et al. (2011). Histone deacetylase inhibitors stimulate histone H3 lysine 4 methylation in part via transcriptional repression of histone H3 lysine 4 demethylases. *Mol. Pharmacol.* 79, 197–206. doi: 10.1124/mol.110.067702
- Huang, W., Perez-Garcia, P., Pokhilko, A., Millar, A. J., Antoshechkin, I., Riechmann, J. L., et al. (2012). Mapping the core of the *Arabidopsis* circadian clock defines the network structure of the oscillator. *Science* 336, 75–79. doi: 10.1126/science.1219075
- Hung, F. Y., Chen, F. F., Li, C., Chen, C., Lai, Y. C., Chen, J. H., et al. (2018). The *Arabidopsis* LDL1/2-HDA6 histone modification complex is functionally associated with CCA1/LHY in regulation of circadian clock genes. *Nucleic Acids Res.* 46, 10669–10681. doi: 10.1093/nar/gky749
- Jiang, D., Yang, W., He, Y., and Amasino, R. M. (2007). *Arabidopsis* relatives of the human lysine-specific Demethylase1 repress the expression of *FWA* and *FLOWERING LOCUS C* and thus promote the floral transition. *Plant Cell* 19, 2975–2987. doi: 10.1105/tpc.107.052373
- Joshi, P., Greco, T. M., Guise, A. J., Luo, Y., Yu, F., Nesvizhskii, A. I., et al. (2013). The functional interactome landscape of the human histone deacetylase family. *Mol. Syst. Biol.* 9:672. doi: 10.1038/msb.2013.26
- Kamioka, M., Takao, S., Suzuki, T., Taki, K., Higashiyama, T., Kinoshita, T., et al. (2016). Direct repression of evening genes by CIRCADIAN CLOCK-ASSOCIATED1 in the *Arabidopsis* circadian clock. *Plant Cell* 28, 696–711. doi: 10.1105/tpc.15.00737
- Khochbin, S., Verdel, A., Lemercier, C., and Seigneurin-Berny, D. (2001). Functional significance of histone deacetylase diversity. *Curr. Opin. Genet. Dev.* 11, 162–166. doi: 10.1016/S0959-437X(00)00174-X
- Kim, W. Y., Geng, R., and Somers, D. E. (2003). Circadian phase-specific degradation of the F-box protein ZTL is mediated by the proteasome. *Proc. Natl. Acad. Sci. U.S.A.* 100, 4933–4938. doi: 10.1073/pnas.0736949100
- Lee, M. G., Wynder, C., Cooch, N., and Shiekhhattar, R. (2005). An essential role for CoREST in nucleosomal histone 3 lysine 4 demethylation. *Nature* 437, 432–435. doi: 10.1038/nature04021
- Li, C. L., Chen, C., Gao, L., Yang, S. G., Nguyen, V., Shi, X. J., et al. (2015). The *Arabidopsis* SWI2/SNF2 chromatin remodeler BRAHMA regulates polycomb function during vegetative development and directly activates the flowering repressor gene SVP. *PLoS Genet.* 11:e1004944. doi: 10.1371/journal.pgen.1004944
- Li, C. L., Gu, L. F., Gao, L., Chen, C., Wei, C. Q., Qiu, Q., et al. (2016). Concerted genomic targeting of H3K27 demethylase REF6 and chromatin-remodeling ATPase BRM in *Arabidopsis*. *Nat. Genet.* 48, 687–693. doi: 10.1038/ng.3555
- Liu, X., Yang, S., Zhao, M., Luo, M., Yu, C. W., Chen, C. Y., et al. (2014). Transcriptional repression by histone deacetylases in plants. *Mol. Plant* 7, 764–772. doi: 10.1093/mp/ssu033
- Liu, X. C., Yu, C. W., Duan, J., Luo, M., Wang, K. C., Tian, G., et al. (2012). HDA6 directly interacts with DNA methyltransferase MET1 and maintains transposable element silencing in *Arabidopsis*. *Plant Physiol.* 158, 119–129. doi: 10.1104/pp.111.184275
- Lu, Q., Tang, X., Tian, G., Wang, F., Liu, K., Nguyen, V., et al. (2010). *Arabidopsis* homolog of the yeast *TREX-2* mRNA export complex: components and anchoring nucleoporin. *Plant J.* 61, 259–270. doi: 10.1111/j.1365-313X.2009.04048.x
- Luo, M., Yu, C. W., Chen, F. F., Zhao, L., Tian, G., Liu, X., et al. (2012). Histone deacetylase HDA6 is functionally associated with AS1 in repression of KNOX genes in *Arabidopsis*. *PLoS Genet.* 8:e1003114. doi: 10.1371/journal.pgen.1003114
- Malapeira, J., Khaitova, L. C., and Mas, P. (2012). Ordered changes in histone modifications at the core of the *Arabidopsis* circadian clock. *Proc. Natl. Acad. Sci. U.S.A.* 109, 21540–21545. doi: 10.1073/pnas.1217022110
- Mas, P., Alabadi, D., Yanovsky, M. J., Oyama, T., and Kay, S. A. (2003a). Dual role of TOC1 in the control of circadian and photomorphogenic responses in *Arabidopsis*. *Plant Cell* 15, 223–236.
- Mas, P., Kim, W. Y., Somers, D. E., and Kay, S. A. (2003b). Targeted degradation of TOC1 by ZTL modulates circadian function in *Arabidopsis thaliana*. *Nature* 426, 567–570.
- McClung, C. R., and Gutiérrez, R. A. (2010). Network news: prime time for systems biology of the plant circadian clock. *Curr. Opin. Genet. Dev.* 20, 588–598. doi: 10.1016/j.gde.2010.08.010
- Murfett, J., Wang, X. J., Hagen, G., and Guilfoyle, T. J. (2001). Identification of *Arabidopsis* histone deacetylase HDA6 mutants that affect transgene expression. *Plant Cell* 13, 1047–1061. doi: 10.1105/tpc.13.5.1047
- Nagel, D. H., Doherty, C. J., Pruneda-Paz, J. L., Schmitz, R. J., Ecker, J. R., and Kay, S. A. (2015). Genome-wide identification of CCA1 targets uncovers an expanded clock network in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 112, E4802–E4810. doi: 10.1073/pnas.1513609112
- Nakamichi, N., Kiba, T., Henriques, R., Mizuno, T., Chua, N. H., and Sakakibara, H. (2010). PSEUDO-RESPONSE REGULATORS 9, 7, and 5 are transcriptional repressors in the *Arabidopsis* circadian clock. *Plant Cell* 22, 594–605. doi: 10.1105/tpc.109.072892
- Nusinow, D. A., Helfer, A., Hamilton, E. E., King, J. J., Imaizumi, T., Schultz, T. F., et al. (2011). The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl growth. *Nature* 475, 398–402. doi: 10.1038/nature10182
- Para, A., Farre, E. M., Imaizumi, T., Pruneda-Paz, J. L., Harmon, F. G., and Kay, S. A. (2007). PRR3 is a vascular regulator of TOC1 stability in the *Arabidopsis* circadian clock. *Plant Cell* 19, 3462–3473. doi: 10.1105/tpc.107.054775
- Perales, M., and Más, P. (2007). A functional link between rhythmic changes in chromatin structure and the *Arabidopsis* biological clock. *Plant Cell* 19, 2111–2123. doi: 10.1105/tpc.107.050807
- Pokhilko, A., Fernandez, A. P., Edwards, K. D., Southern, M. M., Halliday, K. J., and Millar, A. J. (2012). The clock gene circuit in *Arabidopsis* includes a repressilator with additional feedback loops. *Mol. Syst. Biol.* 8:574. doi: 10.1038/msb.2012.6
- Probst, A. V., Fagard, M., Proux, F., Mourrain, P., Boutet, S., Earley, K., et al. (2004). *Arabidopsis* histone deacetylase HDA6 is required for maintenance of transcriptional gene silencing and determines nuclear organization of rDNA repeats. *Plant Cell* 16, 1021–1034. doi: 10.1105/tpc.018754
- Pruneda-Paz, J. L., Breton, G., Para, A., and Kay, S. A. (2009). A functional genomics approach reveals CHE as a component of the *Arabidopsis* circadian clock. *Science* 323, 1481–1485. doi: 10.1126/science.1167206
- Salomé, P. A., Weigel, D., and McClung, C. R. (2010). The role of the *Arabidopsis* morning loop components CCA1, LHY, PRR7, and PRR9 in temperature compensation. *Plant Cell* 22, 3650–3661. doi: 10.1105/tpc.110.079087
- Schaffer, R., Ramsay, N., Samach, A., Corden, S., Putterill, J., Carre, I. A., et al. (1998). The late elongated hypocotyl mutation of *Arabidopsis* disrupts circadian rhythms and the photoperiodic control of flowering. *Cell* 93, 1219–1229. doi: 10.1016/S0092-8674(00)81465-8
- Wang, L., Kim, J., and Somers, D. E. (2013). Transcriptional corepressor TOPLESS complexes with pseudoresponse regulator proteins and histone deacetylases to regulate circadian transcription. *Proc. Natl. Acad. Sci. U.S.A.* 110, 761–766. doi: 10.1073/pnas.1215010110
- Wang, Y., Wu, J. F., Nakamichi, N., Sakakibara, H., Nam, H. G., and Wu, S. H. (2011). LIGHT-REGULATED WD1 and PSEUDO-RESPONSE REGULATOR9 form a positive feedback regulatory loop in the *Arabidopsis* circadian clock. *Plant Cell* 23, 486–498. doi: 10.1105/tpc.110.081661

- Wang, Y., Zhang, H., Chen, Y., Sun, Y., Yang, F., Yu, W., et al. (2009). LSD1 is a subunit of the NuRD complex and targets the metastasis programs in breast cancer. *Cell* 138, 660–672. doi: 10.1016/j.cell.2009.05.050
- Wang, Z. Y., and Tobin, E. M. (1998). Constitutive expression of the *CIRCADIAN CLOCK ASSOCIATED 1 (CCA1)* gene disrupts circadian rhythms and suppresses its own expression. *Cell* 93, 1207–1217. doi: 10.1016/S0092-8674(00)81464-6
- Wu, K., Zhang, L., Zhou, C., Yu, C. W., and Chaikam, V. (2008). HDA6 is required for jasmonate response, senescence and flowering in *Arabidopsis*. *J. Exp. Bot.* 59, 225–234. doi: 10.1093/jxb/erm300
- Yu, C. W., Liu, X., Luo, M., Chen, C., Lin, X., Tian, G., et al. (2011). HISTONE DEACETYLASE6 interacts with FLOWERING LOCUS D and regulates flowering in *Arabidopsis*. *Plant Physiol.* 156, 173–184. doi: 10.1104/pp.111.174417
- Yu, C. W., Tai, R., Wang, S. C., Yang, P., Luo, M., Yang, S., et al. (2017). HISTONE DEACETYLASE6 acts in concert with histone methyltransferases SUVH4, SUVH5, and SUVH6 to regulate transposon silencing. *Plant Cell* 29, 1970–1983. doi: 10.1105/tpc.16.00570

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Hung, Chen, Li, Chen, Chen, Cui and Wu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.