



Published in final edited form as:

J Autism Dev Disord. 2016 May ; 46(5): 1513–1527. doi:10.1007/s10803-013-1957-x.

The mechanisms underlying the ASD advantage in visual search

Zsuzsa Kaldy, Ivy Giserman, Alice S. Carter, and Erik Blaser

University of Massachusetts Boston, Department of Psychology, 100 Morrissey Blvd., Boston, MA, USA, 02125

Abstract

A number of studies have demonstrated that individuals with Autism Spectrum Disorders (ASD) are faster or more successful than typically developing control participants at various visual-attentional tasks (for reviews, see Dakin & Frith, 2005; Simmons, *et al.*, 2009). This “ASD advantage” was first identified in the domain of visual search by Plaisted and colleagues (Plaisted, O’Riordan, & Baron-Cohen, 1998). Here we survey the findings of visual search studies from the past 15 years that contrasted the performance of individuals with and without ASD. Although there are some minor caveats, the overall consensus is that - across development and a broad range of symptom severity - individuals with ASD reliably outperform controls on visual search.

The etiology of the ASD advantage has not been formally specified, but has been commonly attributed to ‘enhanced perceptual discrimination’, a superior ability to visually discriminate between targets and distractors in such tasks (e.g. O’Riordan, 2000). As well, there is considerable evidence for impairments of the attentional network in ASD (for a review, see Keehn, Muller, & Townsend, 2013). We discuss some recent results from our laboratory that support an attentional, rather than perceptual explanation for the ASD advantage in visual search. We speculate that this new conceptualization may offer a better understanding of some of the behavioral symptoms associated with ASD, such as over-focusing and restricted interests.

Keywords

ASD advantage; visual search; over-focusing; phasic attention

Alongside the well-known social and communicative deficits of Autism Spectrum Disorder (ASD) stands an ‘ASD advantage’ characterized by superior performance on many perceptual and attentional tasks, including the embedded figures task (Shah & Frith, 1983) and the block design task (Shah & Frith, 1993; for reviews, see Dakin & Frith, 2005; Simmons, *et al.*, 2009). While most of the research and the public’s awareness have been focused on the impairments in ASD, some scientists (e.g. Mottron, 2011), non-profit companies and many individuals living with autism are actively fighting this one-sided view. This perspective has been featured in the popular press, most recently, for example, in *The New York Times* (Cook: The Autism advantage, Nov. 29, 2012).

Our goal in this paper is twofold. First, we provide a targeted ‘mini-review’ of the empirical findings from a paradigm where the ASD advantage has been studied most extensively: visual search. Second, we present the two most influential theories that have been proposed to explain the ASD advantage in these search tasks. One of them asserts that the cause is primarily perceptual: that bottom-up, low-level processes are functioning atypically in ASD. The other explanation suggests that it is the attention system that is atypical in ASD; a tendency to ‘over-focus’, while perhaps disadvantageous in some contexts, is beneficial to visual search. While not mutually exclusive, these explanations can be hard to tease apart, as both predict reductions in reaction times and increases in accuracy for finding hidden targets in visual search tasks. We will then introduce a recent result from our laboratory, using pupillometry as a measure of attentional engagement that supports an attention-based explanation. Finally, we show how differences in attentional processing can be related to clinical observations. Throughout this review, wherever possible, we will highlight developmental trends.

Visual search

There are several psychophysical paradigms that have been developed to study search abilities. The most ubiquitous is Anne Treisman’s (Treisman & Gelade, 1980). In this paradigm, two conditions have been contrasted: ‘single feature’, ‘pop-out’ (Julesz, 1981) search, and ‘feature conjunction’ search. In a single feature search task an array of items is shown in which a ‘target’ item (the item that is to be searched for) has a unique feature that distinguishes it from a homogeneous set of distractors (e.g. a red disk target in a field of blue disk distractors). The classic signature of single feature search is that the amount of time it takes individuals to find the target is not significantly affected by the set size of the distractors (a red target in a field of 20 blue distractors is found as quickly as in a field of five); the target ‘pops out’ automatically and effortlessly. However, to find an item that is unique among the distractors by virtue of having a conjunction of two different feature dimensions (e.g. a red disk among a distractor set containing both blue disks and red triangles) typically requires an effortful search of the items in the display in a more or less serial fashion; the target no longer pops out. Therefore, search times in feature conjunction search tasks typically vary linearly with the number of distractors. While research suggests that the attentional mechanisms involved in single versus feature conjunction search are not categorically different (see e.g. Duncan & Humphreys, 1989; Wolfe, 1998), and that target-distractor similarity and the perceptual characteristics of the target itself together determine the efficiency of search (Wolfe & Horowitz, 2004), the classic paradigm using shape and color as features is still often used as a way to systematically contrast easy and difficult attentional demands. In the current theory of visual search the distinction is simply between efficient and inefficient search (Wolfe & Horowitz, 2004). For our purposes, the distinctions between easy vs. difficult and efficient vs. inefficient search map onto the single feature/feature conjunction distinction.

Visual search in ASD: the findings

Kate Plaisted and her colleagues were the first to demonstrate the ASD advantage in visual search in 1998 (Plaisted, O’Riordan & Baron-Cohen, 1998). They found that 7–10-year-old

children with autism demonstrated faster reaction times in featureconjunction tasks than verbal-ability matched typically developing (TD) children. The group extended these findings to nonverbal IQ-matched and age-matched controls (O’Riordan, 2000; O’Riordan, Plaisted, Driver, & Baron-Cohen, 2001) and adults (O’Riordan, 2004). Since this seminal report and its follow-ups, a number of studies have contrasted the visual search skills of participants with and without ASD. We have summarized the participant characteristics, basic elements of the tasks employed, and main results of 22 experimental studies from the past fifteen years (see Table 1).

Overwhelmingly, these studies demonstrate that compared to control participants, individuals with ASD are both faster and more accurate (though accuracy is typically just measured to ensure that shorter reaction times do not just reflect a speed-accuracy tradeoff) at identifying a target hidden amongst a number of distractors (‘set size’). This finding has been consistent across ages (from 2.5-year-old children to adults) and the spectrum of symptom severity. In children, the ASD advantage is consistently present even in age-matched control groups. Even the presence of subclinical autism-like traits in normally developed adults (as measured by the Autism Quotient) correlates with better visual search performance. In terms of differences across task design, the ASD advantage is present at different difficulty levels (though more reliably so in ‘conjunction’ search and more difficult, inefficient feature searches) and for various perceptual features (color, shape, and orientation).

There are only a few exceptions to these findings. In Ashwin, Wheelwright, and Baron-Cohen’s (2006) task, the target and distractors in the search task consisted of threatening, happy, and neutral faces. In a series of four experiments, adults with ASD were slower than controls to find a discrepant face in an array of faces. However, these findings are in line with previous research demonstrating deficits related to facial and emotional recognition in individuals with ASD (see Harms, Martin & Wallace, 2010, for a review).

In Baldassi *et al.* (2009), the search performance of the ASD group was not significantly different from TD controls. The paradigm they used though had some salient differences from the search task employed in the other studies reviewed here. Instead of RT or accuracy of target detection, they measured orientation discrimination thresholds in briefly presented (200 ms) displays presented centrally or peripherally. The target was either presented alone (set size 1) or with flankers (set size 6 or 9). Relative to their detection thresholds, both groups were equally slowed by the addition of flankers; no apparent ASD advantage for visual search. Intriguingly, in a different manifestation of the ASD advantage, when the set of items was presented in the periphery, children with ASD showed no crowding effect (a significant impairment in orientation discrimination of the target when it is surrounded by flankers, an effect that was present in TD children). These findings reveal the importance of certain task demands, such as employing brief (such as 200 ms) onset-offset stimuli.

Caron, Mottron, Berthiaume and Dawson (2006) tested typical adults and adults with high-functioning autism. Within each group, they compared a subgroup that was selected based on their exceptional score on the Block Design Task (BDT). They found that those who did exceptionally well on the BDT were faster at conjunction search, independent of diagnosis;

at face value, this would indicate a lack of an ASD advantage. However, BDT scores highly correlated with visual search performance in this study, so matching individuals by BDT score is a proxy for matching them for search; no differences would be expected. The ASD advantage is found in general because the base rate of individuals that score high on BDT and do well on search is (apparently, our survey supports) higher in ASD than TD populations.

Finally, Iarocci *et al.* (2005) found no significant difference between the performance of children with and without ASD in a local/global visual search task. Here, since the ASD group (mean age 8 years) was well above average in non-verbal IQ, the matching TD control group was almost a 1.5 years older. Similarly, Riby *et al.* (2012) found no group differences in a task where older children with and without ASD had to look for a specific object in an array of random objects. Since in this task the salience of the distractors was much more heterogeneous, it is possible that children with ASD got more distracted by particular objects in the displays.

In sum, even though some studies reviewed above have pointed out certain limitations, the ASD advantage in visual search stands on firm empirical ground. That said, it is clear that further investigation is required to pinpoint the task and stimulus parameters that have greatest influence over the expression of the ASD advantage. We will now turn to the discussion of the two prominent theories on the mechanisms underlying the ASD advantage in visual search.

Two theories of the ASD advantage in visual search

I Perceptual enhancement

The first theory of the ASD advantage in visual search was that individuals with ASD have enhanced perceptual discrimination (Plaisted *et al.*, 1998; O’Riordan, 2000; O’Riordan & Plaisted, 2001). Similarly, the Enhanced Perceptual Functioning model proposed by Mottron and his colleagues (Mottron & Burack, 2001; Mottron *et al.*, 2006, for a more recent extension of the model, see Mottron *et al.*, 2013) is based on the idea that both low-level (discrimination) and mid-level (pattern detection) perceptual processes are enhanced in ASD (this model has not been specifically applied to visual search performance (apart from Experiment 4 of Caron *et al.*, 2006), but it is relevant to this discussion). Models based on perceptual enhancement have considerable intuitive appeal and face validity, after all, any manipulation – physical or perceptual – that increases target discriminability (decreasing target-distractor similarity) can increase performance; target–distractor discriminability is a principal rate-determining factor in visual search (Duncan & Humphreys, 1989; Wolfe, 1994).

While there is ample evidence to support that individuals with ASD show higher sensitivity in certain perceptual domains (for a summary, see Mottron, Dawson, Soulieres, Hubert & Burack, 2006), there is no reliable evidence for higher sensitivity to spatial or color contrast in children or adults with ASD (Koh, Milne, & Dobkins, 2010; for a summary, see Simmons *et al.*, 2009), nor evidence for higher visual acuity in general (Bölte *et al.*, 2012, Falkmer *et al.*, 2011). Regarding color perception in ASD, a recent study found *reduced* sensitivity for

color in children with high-functioning autism compared to controls matched on age and nonverbal ability, and no significant differences were found in luminance sensitivity (Franklin *et al.*, 2010).

Examining the studies in Table 1, it is clear that most invoke some form of perceptual enhancement to explain the ASD advantage. Most though do not claim to have direct evidence for perceptual enhancement per se, but instead choose it after a ‘process of elimination’ of competing explanations, or by analogy to similar studies that have also invoked it (as we ourselves did in Kaldy *et al.*, 2011). Of the papers reviewed here, only two measured the discrimination thresholds for their visual search stimuli (in both cases, the parameters of Gabor patches), with one finding no correlation between thresholds and search performance (Brock *et al.*, 2011), and another finding significantly higher thresholds in their Autism group (Baldassi, *et al.*, 2009). In particular, in the Brock, Xu and Brooks (2011) study, the authors tested whether visual search performance for a target defined by the conjunction of spatial frequency and orientation correlates with lower visual discrimination thresholds for the same features. Their sample was undergraduate students who completed the Autism Spectrum Quotient (AQ) questionnaire (Baron-Cohen *et al.*, 2001) before the psychophysical tests. Brock *et al.* (2011) found that adults with higher AQ were faster in visual search, but that visual search performance did not correlate with discrimination thresholds. To date, this has been the most direct test of the enhanced discrimination explanation behind visual search performance, which, if it can be replicated with individuals with and without ASD, can pose a problem for the enhanced perceptual discrimination model as an explanation for the ASD advantage in visual search.

II. Atypical attention

Another theory attributes the superior skills in ASD to an atypically functioning attentional system. Under this view, a tendency to over-focus, and a resistance to disengagement is responsible for superior visual search¹. In a recent review, Keehn, Muller & Townsend (2012) summarized the extant research on attentional functioning in ASD following Posner’s model that distinguishes the alerting, orienting and executive control networks of attention (Posner & Petersen, 1990; Petersen & Posner, 2012). The alerting network has two dissociable functions: *tonic* alertness provides a baseline sensitivity level to incoming information (equivalent to vigilance) and *phasic* alertness is the more transient state that the organism is in when it is actively engaged in a task. The orienting network, on the other hand, selects information from the sensory input by engaging, disengaging and re-engaging attention (whether covertly or overtly, endogenously or exogenously). Finally, the executive control network consists of set shifting (or task switching), working memory and inhibition.

Keehn *et al.* (2013) finds evidence for atypical function in the all three systems in ASD, but puts problems with attentional disengagement at the origin of the developmental cascade that leads to dysfunctional arousal regulation by the alerting system, which in turn engenders

¹It is important to note that an opposite trend, namely increased distractibility has also been reported in ASD. Burack (1994) tested a small group of extremely low-functioning ($N = 12$, mean IQ = 49.5) adults with ASD and suggested that there is a general selective attention deficit in autism. Many authors since this early report have demonstrated (e.g. Remington, Swettenham, Campbell, & Coleman, 2009) evidence for the contrary. A general selective attention deficit is also in contrast with the findings of the large body of research on ASD advantage in visual search reviewed in this paper.

over-focused attention, which explains enhanced visual search abilities. With respect to the other systems, impairments of the executive control network appear later in development so are viewed as secondary to other symptoms of the disorder, and while there are ASD-related attentional disengagement problems in the orienting system as measured by the classic ‘gap-overlap’ task (Saslow, 1967), it does not manifest itself in visual search paradigms. Landry and Bryson (2004) found that children with ASD were slower in the ‘overlap’ condition (in this condition a simple shape is presented centrally briefly, then another stimulus appears in one of two lateral positions. The dependent measure is the saccadic latency to the second stimulus. This is contrasted with the ‘gap’ condition, where the first stimulus disappears before the appearance of the second one) than mental age-matched controls, while there were no differences in the gap condition. Impaired disengagement in the same task was found in older children with ASD (van der Geest, Kemner, Camfferman, Verbaten, & van Engeland, 2001) and in adults with autism (Kawakubo *et al.*, 2007) and importantly, in infants at risk for developing autism (Zwaigenbaum, *et al.*, 2005, Elsabbagh *et al.*, 2009). For example, Zwaigenbaum *et al.* (2005) found that slower disengagement of gaze at 12 months significantly predicted ASD outcome at 2 years of age, and notably, all the infants’ whose disengagement score declined between 6 and 12 months were on the spectrum by 2 years of age. In spite of these findings, children with ASD often show shorter fixation times than controls in visual search (Joseph *et al.*, 2009), and they have been shown to disengage from items at a faster pace or at the same pace (Kaldy *et al.*, 2011) as TD children. Whatever role the orienting network may play in ASD, it does not seem to impede attentional disengagement – as manifested in gaze behavior – in visual search.

An account where atypical function of the alerting system leads to (advantageously) overly focused attention finds independent support from work on the Locus Coeruleus (LC; the brainstem area that regulates noradrenergic activity in cortex). Tonic activity of the LC modulates a diffuse, exploratory attentional state that facilitates task switching, while phasic activity modulates a focused attentional state that facilitates performance on fixed, well-defined tasks (Sara, 2009). This link can be seen during direct manipulation of the LC. In monkeys, local microinfusion of clonidine to increase LC phasic activity increases performance on a visual task, while a suppressive agent (pilocarpine) has the reduces it (Aston-Jones & Cohen, 2005). In humans, administration of modafinil to increase LC phasic activity increases task-related activity in cognitive control areas (shown by fMRI) and improves performance on a visual task (Minzenberg, Watrous, Yoon, Ursu, & Carter, 2008). Recent work has implicated the LC in ASD etiology, in humans (Mehler & Purpura, 2009) and rat models (Darling *et al.*, 2011). In fact, it has been speculated that the LC may be in a persistent hyperphasic state in ASD (Aston-Jones, Iba, Clayton, Rajkowski, & Cohen, 2007). Indeed, administration of venlafaxine to regulate LC activity effectively treats some of the attention-related symptoms of ASD (Hollander, Kaplan, Cartwright, & Reichman, 2000).

It is possible, then, that a dysregulated, hyperphasic LC predisposes individuals with ASD to over-focused attention, thereby increasing performance on tasks that benefit from focused attention and reduced distractibility (like visual search), while potentially decreasing performance on tasks that require shifts of attentional engagement. Fortunately, there is a way to gain insight into LC activity (and therefore attentional state) during a task: pupil

dilation is a biomarker of LC activity (Aston-Jones & Cohen, 2005). Gradual changes in pupil size reflect the tonic activity of the LC while dilations time-linked to task events reflect its phasic activity.

Pupillometry focusing on the phasic, task-related response has long been used as a sensitive, real-time physiological measure of cognitive effort (Kahneman & Beatty, 1966; Beatty, 1982). For instance, when attempting to read an incongruent word in the Stroop task, the pupil dilates (Laeng, Orbo, Holmlund, & Miozzo, 2011). Porter, Troscianko and Gilchrist (2007) found a clear link between search difficulty and mental effort as shown by larger pupil dilations in inefficient (feature-conjunction) vs. efficient (single feature) search. (Recently, Nassar *et al.* (2012) demonstrated a causal link between mental effort (as measured by pupil dilations) and performance in a challenging cognitive task.) This relationship to cognitive effort even holds in infants. For instance, 8-month-olds will show greater pupil dilation when presented with impossible events (Jackson & Sirois, 2009). Pupil responses in ASD and TD groups have been investigated by Anderson, Colombo and their colleagues (Anderson, Colombo, & Shaddy, 2006; Anderson & Colombo, 2009; Anderson, Colombo & Unruh, 2012). They found that tonic pupil size is significantly elevated in children with ASD. However, pupillometry has not been used to investigate the phasic, task-related responses in ASD or to elucidate the mechanisms behind the ASD advantage.

In our recent study (Kaldy, Kraper, Carter, & Blaser, 2011), we found that the ASD advantage in visual search is present as early as 2.5 years of age. We developed a version of the classic visual search paradigm that contrasted single-feature search and feature-conjunction search with varying set sizes (using shape and color as features), that did not require following verbal instructions, making it ideal for toddlers with weak receptive language skills. We tested 17 toddlers with ASD (who tended to be on the severe end of the symptom spectrum – in fact, 15 of them met the criteria for autism) and 17 age-matched typically developing children. Using a Tobii T120 eyetracker, we measured success rate: the percent of trials were toddlers able to find the target within the 4 s presentation period. Our main finding was that (especially in the more attention-demanding feature-conjunction task) toddlers with ASD outperformed controls (by up to a factor of two).

We analyzed pupil dilation from that study (Blaser, Eglington, & Kaldy, 2012; Blaser, Eglington, Carter, & Kaldy, submitted) and found that the toddlers with ASD had exaggerated task-related pupil dilations; the LC was indeed more frequently phasic during search than in controls – evidence of a focused attentional state. Toddlers with ASD are predisposed, in a sense, to be ‘on task’, for visual search. Our analysis showed that typically developing toddlers can focus too, but they tend to do so less frequently during a test session. In short, children with ASD do not search better than TD controls, they are simply more likely to search with effort in any given trial.

This is a parsimonious explanation that accounts for the ASD advantage observed in our study. It could conceivably account for similar effects in other studies (or at least account for some of the variance). After all, if an ASD group exerts cognitive effort more consistently in a visual search task, this will result in better performance – a result that would otherwise be tempting to attribute to enhanced perception. It is important to note that this greater

attentional engagement need not be evident in other measures. In our study we examined gaze behavior alongside task performance, yet the ASD and the TD groups did not differ in the number of fixations they made during a trial, the percent of trials rejected for never having fixated an item, the total time spent dwelling on items, or even the amount of time it took for them to get to the target on successful trials. Without pupillometry, we would have lacked the insight to reject perceptual enhancement as the default explanation (which we had invoked in Kaldy, *et al.*, 2011). This attentional explanation has received some further support from a recent ERP study (Milne, Dunn, Freeth, & Rosas-Martinez, 2013). In a sample of neurotypical adults, Milne and colleagues found that a late ERP component (P3b) in an attentional task significantly predicted visual search efficiency (while earlier components, that reflect perceptual processing, did not).

We hypothesize that this predisposition to intense attentional focus in ASD comes at the cost of resistance to task disengagement. This link is examined in our final section.

Over-focusing and resistance to task disengagement: clinical observations

Both retrospective parental reports on children with ASD (Baranek, 1999) and prospective experimental studies of infants at genetic risk for ASD (Zwaigenbaum *et al.*, 2009, Elsabbagh *et al.*, 2009) demonstrate that attentional dysfunction is one of the most reliable early signs of ASD among affected children. Early atypical attentional symptoms in the social domain include limited initiation and maintenance of eye contact (e.g., Gilberg *et al.*, 1990), reduced social orienting, reduced and atypical joint attention, including deficits in both (a) following others' gaze or pointing and (b) initiating shared attention with others (e.g., showing) as well as unusual and repetitive object-oriented play that often involves atypical visual inspection of or peering at objects (Ozonoff *et al.*, 2008). In many of these atypical attentional behaviors the social and the attentional substrates are deeply intertwined and it is not surprising that social reward circuitry is also implicated in understanding these early atypicalities (Dawson *et al.*, 2001).

Consistent with the heterogeneity of presentations, Liss, Saulnier, Fein & Kinsbourne (2006) reported that 43% of parents of children with ASD ($N=144$) who were surveyed about their child's sensory, motor and attention regulation reported overfocused attention. Consistent with our hypothesis of heightened phasic alerting activity, these authors linked these parental observations of over-focused attention with hyperarousal. Importantly, attention regulation is critical not only for social engagement and communication, but also for regulation of emotion, as gaze shifting is one of the earliest emerging emotion regulation strategies (Manglesdorf, Shapiro, & Marzolf, 1995).

Early differences in attention-modulation may contribute to dynamic, developmental cascades, in which brain and behavioral functioning may be constrained through limited age-typical social interactions (Dawson 2008; Elsabbagh & Johnson 2007, 2010). Specifically, if infants are more likely to be in heightened states of phasic alerting, which may be experienced by interactive partners as over-focusing their attention on (non-social) objects to the exclusion of social information in their environments, intrinsic susceptibilities

that confer an Autism advantage in visual search and an Autism disadvantage in facial processing and social communication may be intensified over time.

To the extent that very young infants with or at risk for ASD choose to pursue repetitive object play (Ozonoff *et al.*, 2008) and fail to acquire joint attention, imitation, and other social communication skills that support typical language acquisition (Mundy, Sigman & Kasari, 1990; Rogers, Hepburn, Stackhouse, & Wehner, 2003), atypical development may be exacerbated. Ozonoff and colleagues (Ozonoff *et al.*, 2008) documented the presence of elevated repetitive and restricted object play as early as 12 months of age and three other studies characterized atypical use of objects in the toddler/preschool period among children with ASD (Bruckner & Yoder, 2007; Mottron *et al.*, 2007; Wetherby *et al.*, 2004). Across these studies, the kinds of behaviors observed included reduced exploration as well as rotating, spinning, twirling, rolling, tapping, banging, rubbing, lining up and unusual visual inspection of or peering at objects. Of great interest for understanding the impact of the ASD advantage in clinical presentation, Ozonoff and colleagues (Ozonoff *et al.*, 2008) reported that the most common repetitive behavior was atypical visual inspection, which was present in seven of the nine 12-month-old infants later diagnosed with ASD. Moreover, there was stability in repetitive behaviors between 12 and 36 months of age and atypical repetitive behaviors at 12 months were associated with lower social-communication scores on the ADOS and lower developmental outcomes on the Mullen Scales of Early Learning.

Consistent with relations between restricted and repetitive object play and broader development, restricted object use, which was defined as the number of toys that children engaged with in a differentiated manner when a set array of toys was presented, was associated with poorer joint attention, social engagement, and imitation skills (Bruckner & Yoder, 2007). Moreover, Wetherby and colleagues (2004) reported that the rate of repetitive behaviors in the second year of life was significantly correlated with ASD outcome at 36 months of age. Sasson and colleagues (Sasson, Turner-Brown, Holtzclaw, Lam, & Bodfish, 2008) used a passive viewing task of social and nonsocial stimuli, and within the second set they included a subset of stimuli chosen to be of high interest to children with ASD (e.g., trains, planes). In both school-age and preschool-age children (Sasson *et al.*, 2011), the stimuli selected for high interest in children in ASD were, in fact, more visually engaging for this group. Importantly, in the older children, longer exploration of non-social stimuli was associated with parent report of increased repetitive behaviors.

Thus, these studies not only highlight the early emergence of restricted and repetitive behaviors in children affected with ASD, but also provide preliminary evidence in support of the potentially important role that visual attention is playing in developmental outcomes.

Dynamic, developmental theories hold promise for early intervention efforts as it may be possible to use deeper understanding of emerging attentional proclivities in children at risk or evidencing early signs to redirect them toward more typical learning experiences that can promote optimal social and communicative development. Thus, just as visual search performance has been associated with concurrent symptom severity (e.g. Joseph *et al.*, 2009), it might inform decisions regarding intervention response and/or aspects of developmental course.

Acknowledgments

This research was supported by National Institutes of Health Grant 2R15EY017985 awarded to EB and ZK and by the University of Massachusetts Presidential Science and Technology Award awarded to ZK, EB and ASC.

REFERENCES

- Almeida RA, Dickinson J, Maybery MT, Badcock JC, Badcock DR. Visual search performance in the autism spectrum II: The radial frequency search task with additional segmentation cues. *Neuropsychologia*. 2010; 48:4117–4124. [PubMed: 20946906]
- Almeida RA, Dickinson J, Maybery MT, Badcock JC, Badcock DR. Visual search targeting either local or global perceptual processes differs as a function of autistic-like traits in the typically developing population. *Journal of Autism and Developmental Disorders*. 2013; 43:1272–1286. [PubMed: 23054202]
- Anderson CJ, Colombo J, Shaddy D. Visual scanning and pupillary responses in young children with autism spectrum disorder. *Journal of Clinical and Experimental Neuropsychology*. 2006; 28:1238–1256. [PubMed: 16840248]
- Anderson CJ, Colombo J. Larger tonic pupil size in young children with autism spectrum disorder. *Developmental Psychobiology*. 2009; 51:207–211. [PubMed: 18988196]
- Anderson CJ, Colombo J, Unruh KE. Pupil and salivary indicators of autonomic dysfunction in autism spectrum disorder. *Developmental Psychobiology*. 2012
- Ashwin E, Wheelwright S, Baron-Cohen S. Finding a face in the crowd: Testing the anger superiority effect in Asperger Syndrome. *Brain and Cognition*. 2006; 61:78–95. [PubMed: 16455174]
- Aston-Jones G, Cohen JD. An Integrative Theory of Locus Coeruleus-Norepinephrine Function: Adaptive Gain and Optimal Performance. *Annual Review of Neuroscience*. 2005; 28:403–450.
- Aston-Jones, G.; Iba, M.; Clayton, E.; Rajkowski, J.; Cohen, J. The locus coeruleus and regulation of behavioral flexibility and attention: Clinical implications. In: Ordway, GA.; Schwartz, MA.; Frazer, A., editors. *Brain norepinephrine: Neurobiology and therapeutics*. New York, NY US: Cambridge University Press; 2007. p. 196-235.
- Baldassi S, Pei F, Megna N, Recupero G, Viespoli M, Iglizzo R, Tancredi R, Muratori F, Cioni G. Search superiority in autism within, but not outside the crowding regime. *Vision Research*. 2009; 49:2151–2156. [PubMed: 19524608]
- Baron-Cohen S, Wheelwright S, Skinner R, Martin J, Clubley E. The Autism-Spectrum Quotient (AQ): Evidence from Asperger syndrome/highfunctioning autism, males and females, scientists and mathematicians. *Journal of Autism and Developmental Disorders*. 2001; 31:5–17. [PubMed: 11439754]
- Bayliss AP, Kritikos A. Brief report: Perceptual load and the autism spectrum in typically developed individuals. *Journal of Autism and Developmental Disorders*. 2011; 41:1573–1578. [PubMed: 21188489]
- Beatty J. Task-evoked pupillary responses, processing load, and the structure of processing resources. *Psychological Bulletin*. 1982; 91:276–292. [PubMed: 7071262]
- Blaser, E.; Eglinton, L.; Kaldy, Z. Toddlers with ASD are better at visual search without trying harder: A pupillometric study; Poster presented at the Annual Meeting of the Vision Sciences Society; May 11–16, 2012; Naples, FL. 2012.
- Blaser E, Eglinton L, Carter AS, Kaldy Z. Pupillometry reveals that the Locus Coeruleus compels focused attention in Autism Spectrum Disorder. submitted.
- Bölte S, Schlitt S, Gapp V, Hainz D, Schirman S, Poustka F, Walter H. A close eye on the eagle-eyed visual acuity hypothesis of autism. *Journal of Autism and Developmental Disorders*. 2012; 42:726–733. [PubMed: 21660498]
- Brock J, Xu JY, Brooks KR. Individual differences in visual search: Relationship to autistic traits, discrimination thresholds, and speed of processing. *Perception*. 2011; 40:739–742. [PubMed: 21936301]
- Bruckner C, Yoder P. Restricted object use in young children with autism: Definition and construct validity. *Autism*. 2007; 11:161–171. [PubMed: 17353216]

- Caron MJ, Mottron LL, Berthiaume CC, Dawson MM. Cognitive mechanisms, specificity and neural underpinnings of visuospatial peaks in autism. *Brain*. 2006; 129:1789–1802. [PubMed: 16597652]
- Cook, J. The autism advantage. *The New York Times Magazine*. 2012 Nov 29. Retrieved from <http://www.nytimes.com>
- Dakin S, Frith U. Vagaries of visual perception in autism. *Neuron*. 2005; 48:497–507. [PubMed: 16269366]
- Darling RD, Alzghoul L, Zhang J, Khatri N, Paul IA, Simpson KL, Lin RS. Perinatal citalopram exposure selectively increases locus coeruleus circuit function in male rats. *Journal of Neuroscience*. 2011; 31:16709–16715. [PubMed: 22090498]
- Dawson G. Early behavioral intervention, brain plasticity, and the prevention of autism spectrum disorder. *Development and Psychopathology*. 2008; 20:775–803. [PubMed: 18606031]
- Dawson G, Osterling J, Rinaldi J, Carver L, McPartland J. Brief report: Recognition memory and stimulus-reward associations: Indirect support for the role of ventromedial prefrontal dysfunction in autism. *Journal of Autism and Developmental Disorders*. 2001; 31:337–341. [PubMed: 11518486]
- Duncan J, Humphreys GW. Visual search and stimulus similarity. *Psychological Review*. 1989; 96:433–458. [PubMed: 2756067]
- Elsabbagh M, Johnson MH. Infancy and autism: progress, prospects, and challenges. *Progress in Brain Research*. 2007; 164:355–383. [PubMed: 17920442]
- Elsabbagh M, Johnson MH. Getting answers from babies about autism. *Trends In Cognitive Sciences*. 2010; 14:81–87. [PubMed: 20074996]
- Elsabbagh M, Volein A, Holmboe K, Tucker L, Csibra G, Baron-Cohen S, Johnson MH. Visual orienting in the early broader autism phenotype: Disengagement and facilitation. *Journal of Child Psychology and Psychiatry*. 2009; 50:637–642. [PubMed: 19298466]
- Falkmer M, Stuart GW, Danielsson H, Bram S, Lönebrink M, Falkmer T. Visual acuity in adults with Asperger’s syndrome: No evidence for “eagleeyed” vision. *Biological Psychiatry*. 2011; 70:812–816. [PubMed: 21885036]
- Franklin A, Sowden P, Notman L, Gonzalez-Dixon M, West D, Alexander I, White A. Reduced chromatic discrimination in children with autism spectrum disorders. *Developmental Science*. 2010; 13:188–200. [PubMed: 20121875]
- Gillberg C, Ehlers S, Schaumann H, Jakobsson G. Autism under age 3 years: A clinical study of 28 cases referred for autistic symptoms in infancy. *Journal of Child Psychology and Psychiatry*. 1990; 31:921–934. [PubMed: 2246342]
- Harms MB, Martin A, Wallace GL. Facial emotion recognition in autism spectrum disorders: A review of behavioral and neuroimaging studies. *Neuropsychology Review*. 2010; 20:290–322. [PubMed: 20809200]
- Hollander E, Kaplan A, Cartwright C, Reichman D. Venlafaxine in children, adolescents, and young adults with autism spectrum disorders: An open retrospective clinical report. *Journal of Child Neurology*. 2000; 15:132–135. [PubMed: 10695900]
- Iarocci G, Burack JA, Shore DI, Mottron L, Enns JT. Global-local visual processing in high functioning children with autism: Structural vs. implicit task biases. *Journal of Autism and Developmental Disorders*. 2006; 36:117–129. [PubMed: 16397823]
- Iarocci G, Burack JA, Shore DI, Mottron L, Enns JT. Global-local visual processing in high functioning children with autism: Structural vs. implicit task biases. *Journal of Autism and Developmental Disorders*. 2006; 36:117–129. [PubMed: 16397823]
- Jackson I, Sirois S. Infant cognition: Going full factorial with pupil dilation. *Developmental Science*. 2009; 12:670–679. [PubMed: 19635092]
- Jarrold C, Gilchrist ID, Bender A. Embedded figures detection in autism and typical development: Preliminary evidence of a double dissociation in relationships with visual search. *Developmental Science*. 2005; 8:344–351. [PubMed: 15985068]
- Joseph RM, Keehn B, Connolly C, Wolfe JM, Horowitz TS. Why is visual search superior in autism spectrum disorder? *Developmental Science*. 2009; 12:1083–1096. [PubMed: 19840062]
- Julesz B. Textons, the elements of texture perception, and their interactions. *Nature*. 1981; 290:91–97. [PubMed: 7207603]

- Kahneman D, Beatty J. Pupil diameter and load on memory. *Science*. 1966; 154:1583–1585. [PubMed: 5924930]
- Kaldy Z, Kraper C, Carter AS, Blaser E. Toddlers with autism spectrum disorder are more successful at visual search than typically developing toddlers. *Developmental Science*. 2011; 14:980–988. [PubMed: 21884314]
- Kawakubo Y, Kasai K, Okazaki S, Hosokawa-Kakurai M, Watanabe K, Kuwabara H, Maekawa H. Electrophysiological abnormalities of spatial attention in adults with autism during the gap overlap task. *Clinical Neurophysiology*. 2007; 118:1464–1471. [PubMed: 17532260]
- Keehn B, Joseph RM. Impaired prioritization of novel onset stimuli in autism spectrum disorder. *Journal of Child Psychology and Psychiatry*. 2008; 49:1296–1303. [PubMed: 19120708]
- Keehn B, Müller RA, Townsend J. Atypical attentional networks and the emergence of autism. *Neuroscience & Biobehavioral Reviews*. 2013; 37:164–183. [PubMed: 23206665]
- Koh HC, Milne E, Dobkins K. Spatial contrast sensitivity in adolescents with autism spectrum disorders. *Journal of Autism and Developmental Disorders*. 2010; 40:978–87. [PubMed: 20213250]
- Laeng B, Ørbo M, Holmlund T, Miozzo M. Pupillary Stroop effects. *Cognitive Processing*. 2011; 12:13–21. [PubMed: 20865297]
- Landry R, Bryson SE. Impaired disengagement of attention in young children with autism. *Journal of Child Psychology and Psychiatry*. 2004; 45:1115–1122. [PubMed: 15257668]
- Liss M, Saulnier C, Fein D, Kinsbourne M. Sensory and attention abnormalities in autistic spectrum disorders. *Autism*. 2006; 10:155–172. [PubMed: 16613865]
- Mangelsdorf SC, Shapiro JR, Marzolf D. Developmental and temperamental differences in emotional regulation in infancy. *Child Development*. 1995; 66:1817–1828. [PubMed: 8556901]
- Mehler MF, Purpura DP. Autism, fever, epigenetics and the locus coeruleus. *Brain Research Reviews*. 2009; 59:388–392. [PubMed: 19059284]
- Milne E, Dunn SA, Freeth M, Rosas-Martinez L. Visual search performance is predicted by the degree to which selective attention to features modulates the ERP between 350 and 600ms. *Neuropsychologia*. 2013; 51:1109–1118. [PubMed: 23499721]
- Minzenberg MJ, Watrous AJ, Yoon JH, Ursu S, Carter CS. Modafinil shifts human locus coeruleus to low-tonic, high-phasic activity during functional MRI. *Science*. 2008; 322:1700–1702. [PubMed: 19074351]
- Mottron L. Changing perceptions: The power of autism. *Nature*. 2011; 479:33–35. [PubMed: 22051659]
- Mottron, L.; Burack, JA. Enhanced perceptual functioning in the development of autism. In: Burack, JA.; Charman, T.; Yirmiya, N.; Zelazo, PR., editors. *The development of autism: Perspectives from theory and research*. Mahwah, NJ US: Lawrence Erlbaum Associates Publishers; 2001. p. 131-148.
- Mottron L, Bouvet L, Bonnel A, Samson F, Burack JA, Dawson M, Heaton P. Veridical mapping in the development of exceptional autistic abilities. *Neuroscience & Biobehavioral Reviews*. 2013; 37:209–228. [PubMed: 23219745]
- Mottron L, Dawson M, Soulières I, Hubert B, Burack J. Enhanced Perceptual Functioning in Autism: An Update, and Eight Principles of Autistic Perception. *Journal of Autism And Developmental Disorders*. 2006; 36:27–43. [PubMed: 16453071]
- Mottron L, Mineau S, Bernier CS, Berthiaume C, Dawson M, Lemay, Faubert J. Lateral glances toward moving stimuli among young children with autism: Early regulation of locally oriented perception? *Development and Psychopathology*. 2007; 19:23–36. [PubMed: 17241482]
- Mundy P, Sigman M, Kasari C. A longitudinal study of joint attention and language development in autistic children. *Journal of Autism and Developmental Disorders*. 1990; 20:115–128. [PubMed: 2324051]
- Nassar MR, Rumsey KM, Wilson RC, Parikh K, Heasley B, Gold JJ. Rational regulation of learning dynamics by pupil-linked arousal systems. *Nature Neuroscience*. 2012; 15:1040–1046. [PubMed: 22660479]
- O’Riordan M. Superior modulation of activation levels of stimulus representations does not underlie superior discrimination in autism. *Cognition*. 2000; 77:81–96. [PubMed: 10986363]

- O'Riordan MA. Superior visual search in adults with autism. *Autism*. 2004; 8:229–248. [PubMed: 15358868]
- O'Riordan MA, Plaisted KC, Driver J, Baron-Cohen S. Superior visual search in autism. *Journal of Experimental Psychology: Human Perception and Performance*. 2001; 27:719–730. [PubMed: 11424657]
- O'Riordan M, Plaisted K. Enhanced discrimination in autism. *The Quarterly Journal of Experimental Psychology A: Human Experimental Psychology*. 2001; 54A:961–979. [PubMed: 11765744]
- Ozonoff S, Macari S, Young GS, Goldring S, Thompson M, Rogers SJ. Atypical object exploration at 12 months of age is associated with autism in a prospective sample. *Autism*. 2008; 12:457–472. [PubMed: 18805942]
- Petersen SE, Posner MI. The attention system of the human brain: 20 years after. *Annual Review of Neuroscience*. 2012; 35:73–89.
- Plaisted K, O'Riordan M, Baron-Cohen S. Enhanced visual search for a conjunctive target in autism: a research note. *Journal of Child Psychology and Psychiatry*. 1998; 39:777–783. [PubMed: 9690940]
- Porter G, Troscianko T, Gilchrist ID. Effort during visual search and counting: insights from pupillometry. *The Quarterly Journal of Experimental Psychology*. 2007; 60:211–229. [PubMed: 17455055]
- Posner MI, Petersen SE. The attention system of the human brain. *Annual Review of Neuroscience*. 1990; 13:25–42.
- Remington AM, Swettenham JG, Lavie N. Lightening the load: Perceptual load impairs visual detection in typical adults but not in autism. *Journal of Abnormal Psychology*. 2012; 121:544–551. [PubMed: 22428792]
- Remington A, Swettenham J, Campbell R, Coleman M. Selective attention and perceptual load in autism spectrum disorder. *Psychological Science*. 2009; 20:1388–1393. [PubMed: 19843262]
- Riby DM, Brown PH, Jones N, Hanley M. Brief report: Faces cause less distraction in autism. *Journal of Autism and Developmental Disorders*. 2012; 42:634–639. [PubMed: 21553149]
- Rogers SJ, Hepburn SL, Stackhouse T, Wehner E. Imitation performance in toddlers with autism and those with other developmental disorders. *Journal of Child Psychology and Psychiatry*. 2003; 44:763–781. [PubMed: 12831120]
- Sara SJ. The locus coeruleus and noradrenergic modulation of cognition. *Nature Reviews Neuroscience*. 2009; 10:211–223. [PubMed: 19190638]
- Saslow MG. Effects of components of displacement step stimuli upon latency for saccadic eye movement. *Journal of The Optical Society Of America*. 1967; 57:1024–1029. [PubMed: 6035296]
- Sasson NJ, Turner-Brown LM, Holtzclaw TN, Lam KSL, Bodfish JW. Children with autism demonstrate circumscribed attention during passive viewing of complex social and nonsocial picture arrays. *Autism Research*. 2008; 1:31–42. [PubMed: 19360648]
- Sasson N, Elison J, Turner-Brown L, Dichter G, Bodfish J. Brief report: circumscribed attention in young children with autism. *Journal of Autism and Developmental Disorders*. 2011; 41:242–247. [PubMed: 20499147]
- Shah A, Frith U. An islet of ability in Autistic children. *Journal of Child Psychology and Psychiatry*. 1983; 4:613–620. [PubMed: 6630333]
- Shah A, Frith U. Why do autistic individuals show superior performance on the block design task? *Journal of Child Psychology and Psychiatry*. 1993; 34:1351–1364. [PubMed: 8294523]
- Simmons DR, Robertson AE, McKay LS, Toal E, McAleer P, Pollick FE. Vision in autism spectrum disorders. *Vision Research*. 2009; 49:2705–2739. [PubMed: 19682485]
- Treisman AM, Gelade G. A feature-integration theory of attention. *Cognitive Psychology*. 1980; 12:97–136. [PubMed: 7351125]
- Van der Geest JN, Kemner C, Camfferman G, Verbaten MN, Van Engeland H. Eye movements, visual attention, and autism: A saccadic reaction time study using the gap and overlap paradigm. *Biological Psychiatry*. 2001; 50:614–619. [PubMed: 11690597]
- Wetherby AM, Woods J, Allen L, Cleary J, Dickinson H, Lord C. Early indicators of autism spectrum disorders in the second year of life. *Journal of Autism and Developmental Disorders*. 2004; 34:473–493. [PubMed: 15628603]

- Wolfe JM. Guided Search 2.0: A revised model of visual search. *Psychonomic Bulletin and Review*. 1994; 1:202–238. [PubMed: 24203471]
- Wolfe JM. What can 1,000,000 trials tell us about visual search? *Psychological Science*. 1998; 9:33–39.
- Wolfe JM, Horowitz TS. What attributes guide the deployment of visual attention and how do they do it? *Nature Reviews Neuroscience*. 2004; 5:1–7.
- Zwaigenbaum L, Bryson S, Lord C, Rogers S, Carter A, Carver L, Yirmiya N. Clinical assessment and management of toddlers with suspected autism spectrum disorder: Insights from studies of high-risk infants. *Pediatrics*. 2009; 123:1383–1391. [PubMed: 19403506]
- Zwaigenbaum L, Bryson S, Rogers T, Roberts W, Brian J, Szatmari P. Behavioral manifestations of autism in the first year of life. *International Journal of Developmental Neuroscience*. 2005; 23:143–152. [PubMed: 15749241]

Table 1

A summary of 22 studies measuring visual search performance in Autism Spectrum Conditions. The studies are listed in the order of the participants' age.

Reference	Clinical group	Comparison group	Task	Stimuli	Outcome Measure	Results, group comparison ('>' means better)	Possible Explanation
Kaldy et al., 2011	ASD (age=2.5, n=17, 3 F)	Typically Developing (age=2.5, n=17, 10 F)	Feature & conjunction search	Objects, varying color and shape	Accuracy	ASD > TD*	"It is likely that 'enhanced discrimination' (O'Riordan & Plaisted, 2001; Joseph et al., 2009) exaggerates the differences between the target and distractors...."
Iarocci et al., 2005	HFA (age=7.9, n=12, 2 F) [Expt 1]; HFA (age=7.8, n=20, F; not spec.) [Expt 2]	TD (age=9.3, n=12, F; not spec.), matched on non-verbal IQ & TD (age=8.1, n=12, F; not spec.), matched on verbal IQ [Expt 1]; TD (age=8.4, n=20, F; not spec.), matched on non-verbal IQ & TD (age=7.8, n=20, F; not spec.), matched on verbal IQ [Expt 2]	Local & global conjunction search [Expt 1]; Conjunction search, probability of local/global target manipulated [Expt 2]	Search elements comprised of 4 black/white dots [Expt 1]; Navon-style geometric shapes [Expt 2]	RT	HFA = TD [Expt 1]; HFA > TD [Expt 2]	"...children with autism are equally adept as their typically developing cohorts in accessing targets at either the local or the global level of visual structure in a search task"; "The children with autism were better able than their typically developing peers to overcome any task independent biases to either local or global processing levels and to tune themselves to the implicit demands of the task."
O'Riordan, 2001	Autism (age=8.4, n=12, F; not spec.) [Expt 1]; Autism (age=8.8, n=12, F; not spec.) [Expt 2]	TD (age=8.2, n=12, F; not spec.), matched on RCPM [Expt 1]; TD (age=8.3, n=12, F; not spec.), matched on RCPM [Expt 2]	Feature & conjunction search [Expt 1]; Difficult feature search [Expt 2]	Letters, varying character and color	RT	Autism > TD [Expt 1]; Autism > TD [Expt 2]	"Possible explanations... include an enhanced ability of individuals with autism to discriminate between items, a superior ability to excite target items and/or inhibit distractor items in autism, or superior inhibition-of-return mechanisms."
O'Riordan, 2000	Autism (age=8.6, n=11, F; not spec.) [Expt 1]; Autism (age=8.5, n=12, F; not spec.) [Expt 2]	TD (age=8.6, n=12, F; not spec.), matched on RCPM [Expt 1]; TD (age=8.5, n=12, F; not spec.), matched on RCPM [Expt 2]	Conjunctive search with negative priming [Expt 1]; Conjunctive search with positive priming [Expt 2]	Letters, varying character and color	RT	Autism > TD [Expt 1]; Autism > TD [Expt 2]	"By a process of elimination these results suggest that enhanced discrimination in autism might instead result from differences in bottom-up, low-level perceptual processing of incoming stimuli."
Plaisted et al., 1998	ASD (age=8.8, n=8, F; not spec.)	TD (age=7.8, n=8, F; not spec.), matched on verbal IQ	Feature & conjunction search	Letters, varying character and color	RT	ASD > TD*	"A simpler way of explaining the current result is to suggest that individuals with autism show superior visuospatial skills (Shah, 1988) and that

Reference	Clinical group	Comparison group	Task	Stimuli	Outcome Measure	Results, group comparison ('>' means better)	Possible Explanation
O'Riordan & Plautsch, 2001	Autism (age=9.2, n=15, 3 F) [Expt 1]; Autism (age=9.0, n=13, 2 F) [Expt 2]	TD (age=8.6, n=15, 3 F), matched on RCPM [Expt 1]; TD (age=8.5, n=13, 2 F), matched on RCPM [Expt 2]	Conjunction search [Expt 1]; Feature search with target-distractor similarity manipulation [Expt 2]	Rectangles, of varying color, orientation, and shape [Expt 1]; Letters, varying character and color [Expt 2]	RT	Autism > TD	"This work suggests that individuals with autism have an enhanced ability to discriminate between items and that this difference in processing underlies their superior performance on visual search tasks."
Baldassi et al., 2009	HFA (age=11.2, n=12, 0 F)	TD (age=12.4, n=15, 0 F), matched on IQ	Feature search with brief presentations [Expt 1]; Orientation discrimination with crowding [Expt 2]; Discrimination thresholds also measured	Gabor patches, of varying orientation	Accuracy (relative discrimination thresholds)	HFA = TD [Expt 1]; HFA > TD (i.e. lower relative discrimination thresholds in HFA group) [Expt 2]	"If we relate the present psychophysical results with the electrophysiological finding that observers with ASD lack the neural signature of one of the two key integration tasks, it is then legitimate to raise the suggestion that the ability to resist to crowding is due to the lack of 'compulsory' integration in observers with ASD."
Jarrold et al., 2005	Autism (age=12.4, n=18, F; not spec.)	TD (age=6.5, n=18, F not specified), matched on RCPM	Difficult feature & conjunction search	'Clown' figures, of varying color and letter-character shape	RT	Autism > TD	"The fact that detection in this group relates to feature search, which we argue relies more on target detection than grouping, suggests that it may be enhanced discrimination of target features that underlies superior embedded figures performance in autism."
Riby et al., 2012	Autism (age=12.9; n=28; 4 F)	TD (age=10.3; n=28; 6 F), matched on RCPM	Object search	Real-world object images (butterfly target; various distractors including a face of varying emotion)	RT	Autism = TD	"...individuals with autism show significantly less face distraction than individuals who are developing typically...;...more experimental work is required to elucidate whether the pattern seen here for individuals with autism is due to general visual perception and attention atypicalities, rather than being 'face-specific'."
Joseph et al., 2009	ASD (age=14.7, n=21, 4 F)	TD (age=14.2, n=21; 4 F), matched on non-verbal IQ	Spatial configuration search (static & dynamic)	Letters, varying character	RT and gaze behavior	ASD > TD	"Thus, it is likely that the ASD advantage reflected faster processing of individual

Reference	Clinical group	Comparison group	Task	Stimuli	Outcome Measure	Results, group comparison (> means better)	Possible Explanation
Keehn & Joseph, 2008	ASD (age=14.1, n=22, 3 F)	TD (age=14.4, n=22, 4 F), matched on non-verbal IQ	(with preview) conjunction search	Letters, varying character and color	RT and gaze behavior	ASD > TD (just in 'Target Old' condition)	items after they had been selected by attention."; "...our findings suggest that non-search factors in general and enhanced visual perception in particular account for the visual search advantage exhibited by individuals with ASD." "...children with ASD showed no difference in RT to search targets occurring as new elements as compared to targets occurring as old elements, indicating that their attention was not preferentially directed to novel onsets"; "...findings provide fairly compelling evidence of a specific anomaly of attention modulation that could explain a range of associated behavioral features of ASD, including tendencies toward over-selectivity and perseveration in attentional focus...."
Caron et al., 2006	HFA with high BDT score (age=23.3, n=8, F: not spec.); HFA with average BDT score (age=18.9, n=8, F: not spec.)	Typical adults with high BDT score (age=16.9, n=8, F: not spec.); Typical adults with average BDT score (age=18.6, n=10, F: not spec.)	Feature & conjunction search	Two-part geometric shapes	RT	High scores in Block Design Test correlate with faster RT (in both ASD and typical group)	"...in a significant proportion of autistic individuals a superior visual input issuing from early stages of visual processing increases the level of performance of subsequent feed-forward flow of visual information as visual selective attention [visual search]..."
O'Riordan, 2004	HFA (age=22, n=10, F: not spec.)	Typical adults (age=22, n=10, F: not spec.), matched on IQ	Feature & conjunction search [Expt 1]; Difficult feature search [Expt 2]; Conjunction search, target-distractor similarity manipulated [Expt 3]	Letters, varying color [Expt 1]; Letter parts [Expt 2]; Letters, varying character and color [Expt 3]	RT	HFA > Typical adults* [Expt 1]; HFA > Typical adults [Expt 2]; HFA > Typical adults [Expt 3] (just for larger set sizes)	"The results showed that adults with autism were not slowed to the same extent as controls by an increase in target-distractor similarity, thus supporting the notion of superior discrimination in adults with autism."; "The present study suggests that superior unique item detection resulting from enhanced discrimination is

Reference	Clinical group	Comparison group	Task	Stimuli	Outcome Measure	Results, group comparison ('>' means better)	Possible Explanation
Remington et al., 2009	ASD (age=23.5, n=14, 7 F)	Typical adults (age=26.6, n=23, 10 F), matched on non-verbal IQ	Response-competition letter search in presence of a special 'neutral' or 'incompatible' distractor)	Letters, varying character	RT	ASD = Typical adults (search); ASD > Typical adults for concurrent detection task	present in adults with autism as well as children." "The results presented here seem to suggest that the control and ASD groups had different perceptual capacities.... Findings regarding increased perceptual capacity may also explain the superior visual search abilities that have consistently been found among individuals with ASD."
Remington et al., 2012	ASD (age=24.1, n=14, 3 F)	Typical adults (age=25.75, n=14, 8 F), matched on non-verbal IQ	Dual task: easy/hard letter search coupled with concurrent detection task	Letters, varying character	RT	ASD = Typical adults (search task); ASD > Typical adults (detection task)	"The present findings demonstrate that ASD is associated with enhanced perceptual capacity in a visual detection task."; "...the high level of detection that the participants with ASD maintained even under higher levels of load clearly suggests a perceptual advantage."
Collignon et al., 2011	Autism (Adults, age=24.5, n=16, 1 F)	Typical adults (age=21, n=19, 0 F), matched on IQ	Difficult feature search	Line segments, of varying orientation	RT	Autism > Typical adults	"Our results also demonstrated... more efficient unimodal visual search performance.... This well replicated ability has been demonstrated to have a perceptual rather than attentional origin (Joseph et al., 2009), and has been associated with multiple other perceptual superiorities (Caron et al., 2006)."
Ashwin et al., 2006	HFA & Asperger Syndrome (Adults, age=25.9, n=1 HFA/17 AS, 0 F)	Typical adults (age=27.3, n=18, 0 F), matched on IQ	Emotional face search	Schematic faces, of varying emotional expressions	RT	Typical adults > Autism	"... while [the ASD group] showed the anger superiority effect, they may be processing the faces using a different strategy or style."; "These findings suggest that... people more likely to process information using a 'cognitive style' characterised by enhanced local feature

Reference	Clinical group	Comparison group	Task	Stimuli	Outcome Measure	Results, group comparison ('>' means better)	Possible Explanation
Bayliss & Kritikos, 2011	High AQ (Adults, age=20.1, n=86, 63 F)	Low AQ (Adults, n=93, 69 F)	Response-competition letter search (letter search in presence of a special 'neutral' or 'incompatible' distractor)	Letters, of varying character	RT	High AQ = Low AQ (search); High AQ group shows greater interference effect	detection over holistic processing. "This finding suggests that High AQ participants have higher perceptual capacity than Low AQ participants."; "....competition at the level of response or attentional selection is the clearest explanation for the group differences (as opposed to, for example, enhanced peripheral perception in the high AQ group."
Brook et al., 2011	High AQ (Adults, age=22, n=38, 24 F)	Low AQ (drawn from same group as High AQ)	Conjunction search, target-distractor similarity manipulated; Discrimination thresholds also measured	Gabor patches, of varying orientation and spatial frequency	RT	High AQ > Low AQ	"...we found no evidence linking performance on visual-search tasks to the ability to make fine-grained discriminations along the stimulus dimensions the differentiated targets from distractors."; "AQ scores were not directly correlated with... discrimination thresholds.... The link between visual search and sub-clinical autism traits appears to be mediated by some other cognitive variable...."
Almeida et al., 2010	High AQ (Adults, age not spec., n=25, 15 F)	Low AQ (Adults, age not spec., n=20, 10 F), matched on IQ	Difficult feature search	Radial frequency closed contours, of varying shape	RT	High AQ > Low AQ	"...those higher on the autism spectrum are more efficient searchers, displaying superior detection of a single feature within a larger scene."
Almeida et al., 2013	High AQ (Adults, Age not spec., n=19, 15 F)	Low AQ (Adults, age not spec., n=25, 10 F), matched on IQ	Difficult feature search	Radial frequency contours, of varying shape	RT	High AQ > Low AQ	"High AQ search superiority may be explained by enhanced focal processing of curvature in the early stages of the form vision pathway and superior global detection of shape primitives.... more probable is the presence of a superior search process that allows elements to be processed more efficiently, which enables a consistent search advantage at both levels of processing."

Reference	Clinical group	Comparison group	Task	Stimuli	Outcome Measure	Results, group comparison ('>' means better)	Possible Explanation
Milne et al., 2013	High AQ (Adults, Age range: 18–37 years, n=20, number of F not spec.)	Low AQ (Adults, Age range: 18–37 years, n=16, number of F not spec.)	Conjunction search	Letters, varying character and color	RT	High AQ > Low AQ	"visual search efficiency in ASD may arise from atypicality of a higher level process rather than from differences in low-level perceptual function."

(Abbreviations: AQ: Autism Spectrum Quotient (Baron-Cohen *et al.* 2001), BDT: Block Design Task; F: Female; HFA: High Functioning Autism, RT: Reaction Time; TD: Typically Developing.)