

The Mukai pairing. I. A categorical approach

Andrei Căldăraru and Simon Willerton

ABSTRACT. We study the Hochschild homology of smooth spaces, emphasizing the importance of a pairing which generalizes Mukai’s pairing on the cohomology of K3 surfaces. We show that integral transforms between derived categories of spaces functorially induce linear maps on homology. Adjoint functors induce adjoint linear maps with respect to the Mukai pairing. We define a Chern character with values in Hochschild homology, and we discuss analogues of the Hirzebruch–Riemann–Roch theorem and the Cardy Condition from physics. This is done in the context of a 2-category which has spaces as its objects and integral kernels as its 1-morphisms.

CONTENTS

Introduction	62
1. The 2-category of kernels	67
1.1. A reminder on 2-categories	67
1.2. The 2-category $\mathcal{V}ar$	70
2. Serre functors	70
2.1. The Serre functor on $\mathbf{D}(X)$	71
2.2. Serre kernels and the Serre functor for $\mathbf{D}(X \times Y)$	71
3. Adjoint kernels	72
3.1. Adjunctions in 2-categories	72
3.2. Left and right adjoints of kernels	75
3.3. Partial traces	76
3.4. Adjunction as a 2-functor	77
4. Induced maps on homology	78

Received July 30, 2007.

2000 *Mathematics Subject Classification*. 18E30 (primary), 14F05, 81T45.

Key words and phrases. Hochschild homology, Mukai pairing, two-category of varieties, Cardy condition, integral kernel, Fourier–Mukai transform.

AC’s initial work on this project was supported by an NSF postdoctoral fellowship, and by travel grants and hospitality from the University of Pennsylvania, the University of Salamanca, Spain, and the Newton Institute in Cambridge, England. AC’s current work is supported by the National Science Foundation under Grant No. DMS-0556042. SW has been supported by a WUN travel bursary and a Royal Society Conference grant.

4.1.	Hochschild cohomology	78
4.2.	Hochschild homology	78
4.3.	Push-forward and pull-back	79
5.	The Mukai pairing and adjoint kernels	80
6.	The Chern character	82
6.1.	Definition of the Chern character	82
6.2.	The Chern character as a map on K-theory	82
6.3.	The Chern character and inner products	84
6.4.	Example	85
7.	Open-closed TQFTs and the Cardy Condition	85
7.1.	Open-closed 2d TQFTs	86
7.2.	Open-closed 2d TQFTs with D-branes	87
7.3.	The open-closed 2d TQFT from a Calabi–Yau manifold	88
7.4.	The Baggy Cardy Condition	89
Appendix A.	Duality and partial trace	91
A.1.	Polite duality	91
A.2.	Reflexively polite kernels	95
References		97

Introduction

The purpose of the present paper is to introduce the Mukai pairing on the Hochschild homology of smooth, proper spaces. This pairing is the natural analogue, in the context of Hochschild theory, of the Poincaré pairing on the singular cohomology of smooth manifolds.

Our approach is categorical. We start with a geometric category, whose objects will be called *spaces*. For a space X we define its Hochschild homology which is a graded vector space $\mathrm{HH}_\bullet(X)$ equipped with the nondegenerate *Mukai pairing*. We show that this structure satisfies a number of properties, the most important of which are *functoriality* and *adjointness*.

The advantage of the categorical approach is that the techniques we develop apply in a wide variety of geometric situations, as long as an analogue of Serre duality is satisfied. Examples of categories for which our results apply include compact complex manifolds, proper smooth algebraic varieties, proper Deligne–Mumford stacks for which Serre duality holds, representations of a fixed finite group, and compact “twisted spaces” in the sense of [3]. We expect the same construction to work for categories of Landau–Ginzburg models [16], but at the moment we do not know if this context satisfies all the required properties.

The Hochschild structure. In order to define the Hochschild structure of a space we need notation for certain special kernels which will play a

fundamental role in what follows. For a space X , denote by Id_X and Σ_X^{-1} the objects of $\mathbf{D}(X \times X)$ given by

$$\text{Id}_X := \Delta_* \mathcal{O}_X \quad \text{and} \quad \Sigma_X^{-1} := \Delta_* \omega_X^{-1}[-\dim X],$$

where $\Delta : X \rightarrow X \times X$ is the diagonal map, and ω_X^{-1} is the anticanonical line bundle of X . When regarded as kernels, these objects induce the identity functor and the inverse of the Serre functor on $\mathbf{D}(X)$, respectively. We shall see in the sequel that Id_X can be regarded as the identity 1-morphism of X in a certain 2-category $\mathcal{V}\text{ar}$.

The *Hochschild structure* of the space X then consists of the following data:

- the graded ring $\text{HH}^\bullet(X)$, the Hochschild cohomology ring of X , whose i -th graded piece is defined as

$$\text{HH}^i(X) := \text{Hom}_{\mathbf{D}(X \times X)}^i(\text{Id}_X, \text{Id}_X);$$

- the graded left $\text{HH}^\bullet(X)$ -module $\text{HH}_\bullet(X)$, the Hochschild homology module of X , defined as

$$\text{HH}_i(X) := \text{Hom}_{\mathbf{D}(X \times X)}^{-i}(\Sigma_X^{-1}, \text{Id}_X);$$

- a nondegenerate graded pairing $\langle -, - \rangle_M$ on $\text{HH}_\bullet(X)$, the *generalized Mukai pairing*.

The above definitions of Hochschild homology and cohomology agree with the usual ones for quasiprojective schemes (see [5]). The pairing is named after Mukai, who was the first to introduce a pairing satisfying the main properties below, on the total cohomology of complex K3 surfaces [15].

Properties of the Mukai pairing. The actual definition of the Mukai pairing is quite complicated and is given in Section 5. We can, however, extricate the fundamental properties of Hochschild homology and of the Mukai pairing.

Functoriality: Integral kernels induce, in a functorial way, linear maps on Hochschild homology. Explicitly, to any integral kernel $\Phi \in \mathbf{D}(X \times Y)$ we associate, in Section 4.3, a linear map of graded vector spaces

$$\Phi_* : \text{HH}_\bullet(X) \rightarrow \text{HH}_\bullet(Y),$$

and this association is functorial with respect to composition of integral kernels (Theorem 6).

Adjointness: For any adjoint pair of integral kernels $\Psi \dashv \Phi$, the induced maps on homology are themselves adjoint with respect to the Mukai pairing:

$$\langle \Psi_* v, w \rangle_M = \langle v, \Phi_* w \rangle_M$$

for $v \in \text{HH}_\bullet(Y)$, $w \in \text{HH}_\bullet(X)$ (Theorem 8).

The following are then consequences of the above basic properties:

Chern character: In all geometric situations there is a naturally defined object $\mathbf{1} \in \mathrm{HH}_0(\mathrm{pt})$. An element \mathcal{E} in $\mathbf{D}(X)$ can be thought of as the kernel of an integral transform $\mathrm{pt} \rightarrow X$, and using functoriality of homology we define a Chern character map

$$\mathrm{ch}: K_0(X) \rightarrow \mathrm{HH}_0(X), \quad \mathrm{ch}(\mathcal{E}) = \mathcal{E}_*(\mathbf{1}).$$

For a smooth proper variety the Hochschild–Kostant–Rosenberg isomorphism identifies $\mathrm{HH}_0(X)$ and $\bigoplus_{\mathbb{P}} H^{\mathbb{P}, \mathbb{P}}(X)$; our definition of the Chern character matches the usual one under this identification [5].

Semi-Hirzebruch–Riemann–Roch Theorem: For $\mathcal{E}, \mathcal{F} \in \mathbf{D}(X)$ we have

$$\langle \mathrm{ch}(\mathcal{E}), \mathrm{ch}(\mathcal{F}) \rangle_{\mathrm{M}} = \chi(\mathcal{E}, \mathcal{F}) = \sum_i (-1)^i \dim \mathrm{Ext}_{\mathcal{X}}^i(\mathcal{E}, \mathcal{F}).$$

Cardy Condition: The Hochschild structure appears naturally in the context of open-closed topological quantum field theories (TQFTs). The Riemann–Roch theorem above is a particular case of a standard constraint in these theories, the Cardy Condition. We briefly discuss open-closed TQFTs, and we argue that the natural statement of the Cardy Condition in the B-model open-closed TQFT is always satisfied, even for spaces which are not Calabi–Yau (Theorem 16).

The 2-categorical perspective. In order to describe the functoriality of Hochschild homology it is useful to take a macroscopic point of view using a 2-category called $\mathcal{V}\mathrm{ar}$. One way to think of this 2-category is as something half-way between the usual category consisting of spaces and maps, and $\mathcal{C}\mathrm{at}$, the 2-category of (derived) categories, functors and natural transformations. The 2-category $\mathcal{V}\mathrm{ar}$ has spaces as its objects, has objects of the derived category $\mathbf{D}(X \times Y)$ — considered as integral kernels — as its 1-morphisms from X to Y , and has morphisms in the derived category as its 2-morphisms.

One consequence of thinking of spaces in this 2-category is that whereas in the usual category of spaces and maps two spaces are equivalent if they are isomorphic, in $\mathcal{V}\mathrm{ar}$ two spaces are equivalent precisely when they are Fourier–Mukai partners. This is the correct notion of equivalence in many circumstances, thus making $\mathcal{V}\mathrm{ar}$ an appropriate context in which to work.

This point of view is analogous to the situation in Morita theory in which the appropriate place to work is not the category of algebras and algebra morphisms, but rather the 2-category of algebras, bimodules and bimodule morphisms. In this 2-category two algebras are equivalent precisely when they are Morita equivalent, which again is the pertinent notion of equivalence in many situations.

Many facts about integral transforms can be stated very elegantly as facts about the 2-category $\mathcal{V}\mathrm{ar}$. For example, the fact that every integral transform between derived categories has both a left and right adjoint is an immediate consequence of the more precise fact — proved exactly the same way — that every integral kernel has both a left and right adjoint in

$\mathcal{V}\text{ar}$. Here the definition of an adjoint pair of 1-morphisms in a 2-category is obtained from one of the standard definitions of an adjoint pair of functors by everywhere replacing the word ‘functor’ by the word ‘1-morphism’ and the words ‘natural transformation’ by the word ‘2-morphism’.

The Hochschild cohomology of a space X has a very natural description in terms of the 2-category $\mathcal{V}\text{ar}$: it is the “second homotopy group of $\mathcal{V}\text{ar}$ based at X ”, which means that it is $2\text{-Hom}_{\mathcal{V}\text{ar}}(\text{Id}_X, \text{Id}_X)$, the set of 2-morphisms from the identity 1-morphism at X to itself. Unpacking this definition for $\mathcal{V}\text{ar}$ one obtains precisely $\text{Ext}_{X \times X}^\bullet(\mathcal{O}_\Delta, \mathcal{O}_\Delta)$, one of the standard definitions of Hochschild cohomology. By analogy with homotopy groups, given a kernel $\Phi: X \rightarrow Y$, i.e., a “path” in $\mathcal{V}\text{ar}$, one might expect an induced map $\text{HH}^\bullet(X) \rightarrow \text{HH}^\bullet(Y)$ obtained by “conjugating with Φ ”. However, this does not work, as the analogue of the “inverse path to Φ ” needed is a simultaneous left and right adjoint of Φ , and such a thing does not exist in general as the left and right adjoints of Φ differ by the *Serre kernels* of X and Y .

The Hochschild homology $\text{HH}_\bullet(X)$ of a space X can be given a similar natural definition in terms of $\mathcal{V}\text{ar}$ — it is $2\text{-Hom}_{\mathcal{V}\text{ar}}(\Sigma_X^{-1}, \text{Id}_X)$ the set of 2-morphisms from the inverse Serre kernel of X to the identity 1-morphism at X . In this case, the idea of “conjugating by a kernel $\Phi: X \rightarrow Y$ ” does work as the Serre kernel in the definition exactly compensates the discrepancy between the left and right adjoints of Φ .

The functoriality of Hochschild homology can be expressed by saying that HH_\bullet is a functor into the category of vector spaces from the Grothendieck category of the 2-category $\mathcal{V}\text{ar}$ (i.e., the analogue of the Grothendieck group of a 1-category) whose objects are spaces and whose morphisms are *isomorphism classes* of kernels. One aspect of this which we do not examine here is related to the fact that this Grothendieck category is actually a monoidal category with certain kinds of duals for objects and morphisms, and that Hochschild homology is a monoidal functor. The Mukai pairing is then a manifestation of the fact that spaces are self-dual in this Grothendieck category. Details will have to appear elsewhere.

There is an alternative categorical approach to defining Hochschild homology and cohomology. This approach uses the notion of enhanced triangulated categories of Bondal and Kapranov [1], which are triangulated categories arising as homotopy categories of differential-graded (dg) categories. In [18], Toën argued that the Hochschild cohomology $\text{HH}^\bullet(X)$ of a space X can be regarded as the cohomology of the dg-algebra of dg-natural transformations of the identity functor on the dg-enhancement of $\mathbf{D}(X)$. It seems reasonable to expect that a similar construction can be used to define the Hochschild homology $\text{HH}_\bullet(X)$ as dg-natural transformations from the inverse of the Serre functor to the identity. However, since the theory of Serre functors for dg-categories is not yet fully developed, we chose to use the language of the 2-category $\mathcal{V}\text{ar}$, where all our results can be made precise.

String diagram notation. As 2-categories are fundamental to the functoriality, and they are fundamentally 2-dimensional creatures, we adopt a 2-dimensional notation. The most apt notation in this situation appears to be that of *string diagrams*, which generalizes the standard notation used for monoidal categories in quantum topology. String diagrams are Poincaré dual to the usual arrow diagrams for 2-categories. The reader unfamiliar with these ideas should be aware that the pictures scattered through this paper form rigorous notation and are not just mnemonics.

Note. This paper supersedes the unpublished paper [4], in which it was stated that hopefully the correct categorical context could be found for the results therein. This paper is supposed to provide the appropriate context.

Synopsis. The paper is structured as follows. The first section is devoted to the study of integral transforms and of the 2-category $\mathcal{V}ar$. In the next section we review the Serre functor and Serre trace on the derived category, and we use these in Section 3 to study adjoint kernels in $\mathcal{V}ar$. In Section 4 we introduce the maps between Hochschild homology groups associated to a kernel, and we examine their functorial properties. The Mukai pairing is defined in Section 5, where we also prove its compatibility with adjoint functors. In Section 6 we define the Chern character and we prove the Semi-Hirzebruch–Riemann–Roch theorem. We conclude with Section 7 where we review open-closed TQFTs, and we discuss the Cardy Condition. An appendix contains some of the more technical proofs.

Notation. Throughout this paper k will denote an algebraically closed field of characteristic zero and $\mathbf{D}(X)$ will denote the bounded derived category of coherent sheaves on X . Categories will be denoted by bold letters, such as \mathbf{C} , and the names of 2-categories will have a script initial letter, such as $\mathcal{V}ar$.

The base category of spaces. We fix for the remainder of the paper a geometric category, whose objects we shall call *spaces*. It is beyond the purpose of this paper to list the axioms that this category needs to satisfy, but the following categories can be used:

- smooth projective schemes over k ;
- smooth proper Deligne-Mumford stacks over k ;
- smooth projective schemes over k , with an action of a fixed finite group G , along with G -equivariant morphisms;
- twisted spaces in the sense of [3], i.e., smooth projective schemes over k , enriched with a sheaf of Azumaya algebras.

For any space X as above, the category of coherent sheaves on X makes sense, and the standard functors (push-forward, pull-back, sheaf-hom, etc.) are defined and satisfy the usual compatibility relations.

Acknowledgments. We have greatly benefited from conversations with Jonathan Block, Tom Bridgeland, Andrew Kresch, Aaron Lauda, Eyal Markman, Mircea Mustața, Tony Pantev, Justin Roberts, Justin Sawon, and Sarah Witherspoon. Many of the ideas in this work were inspired by an effort to decipher the little known but excellent work [10] of Nikita Markarian. Greg Moore and Tom Bridgeland suggested the connection between the Semi-Hirzebruch–Riemann–Roch theorem and the Cardy Condition in physics.

1. The 2-category of kernels

In this section we introduce the 2-category $\mathcal{V}ar$, which provides the natural context for the study of the structure of integral transforms between derived categories of spaces. The objects of $\mathcal{V}ar$ are spaces, 1-morphisms are kernels of integral transforms, and 2-morphisms are maps between these kernels. Before introducing $\mathcal{V}ar$ we remind the reader of the notion of a 2-category and we explain the string diagram notation of which we will have much use.

1.1. A reminder on 2-categories. We will review the notion of a 2-category at the same time as introducing the notation we will be using. Recall that a 2-category consists of three levels of structure: objects; 1-morphisms between objects; and 2-morphisms between 1-morphisms. It is worth mentioning a few examples to bear in mind during the following exposition.

1. The first example is the 2-category $\mathcal{C}at$ of categories, functors and natural transformations.
2. The second example is rather a family of examples. There is a correspondence between 2-categories with one object \star and monoidal categories. For any monoidal category the objects and morphisms give respectively the 1-morphisms and 2-morphisms of the corresponding 2-category.
3. The third example is the 2-category $\mathcal{A}lg$ with algebras over some fixed commutative ring as its objects, with the set of A - B -bimodules as its 1-morphisms from A to B , where composition is given by tensoring over the intermediate algebra, and with bimodule morphisms as its 2-morphisms.

There are various ways of notating 2-categories: the most common way is to use *arrow diagrams*, however the most convenient way for the ideas in this paper is via *string diagrams* which are Poincaré dual to the arrow diagrams. In this subsection we will draw arrow diagrams on the left and string diagrams on the right to aid the reader in the use of string diagrams.

Recall the idea of a 2-category. For any pair of objects X and Y there is a collection of morphisms $1\text{-Hom}(X, Y)$; if $\Phi \in 1\text{-Hom}(X, Y)$ is a 1-morphism

then it is drawn as below.

$$\dot{Y} \xleftarrow{\Phi} \dot{X} \qquad \frac{\Phi}{Y \bullet X}$$

These 1-dimensional pictures will only appear as the source and target of 2-morphism, i.e., the top and bottom of the 2-dimensional pictures we will be using. In general 1-morphisms will be denoted by their identity 2-morphisms, see below.

If $\Phi, \Phi' \in \mathbf{1-Hom}(X, Y)$ are parallel 1-morphisms — meaning simply that they have the same source and target — then there is a set of 2-morphisms $\mathbf{2-Hom}(\Phi, \Phi')$ from Φ to Φ' . If $\alpha \in \mathbf{2-Hom}(\Phi, \Phi')$ is a 2-morphism then it is drawn as below.

The diagram shows two 1-morphisms, Φ and Φ' , between objects \dot{Y} and \dot{X} . Φ is represented by a curved arrow pointing from \dot{X} to \dot{Y} , and Φ' is represented by a curved arrow pointing from \dot{Y} to \dot{X} . A 2-morphism α is shown as a vertical double arrow pointing upwards from Φ to Φ' .

The string notation shows a vertical line representing the 1-morphism Φ from Y to X . A box labeled α is placed on this line, with a vertical line representing the 1-morphism Φ' passing through the box.

At this point make the very important observation that diagrams are read from right to left and from bottom to top.

There is a *vertical composition* of 2-morphisms so that if $\alpha: \Phi \Rightarrow \Phi'$ and $\alpha': \Phi' \Rightarrow \Phi''$ are 2-morphisms then the vertical composite $\alpha' \circ_v \alpha: \Phi \Rightarrow \Phi''$ is defined and is denoted as below.

The diagram shows two 1-morphisms, Φ and Φ'' , between objects \dot{Y} and \dot{X} . Φ is represented by a curved arrow pointing from \dot{X} to \dot{Y} , and Φ'' is represented by a curved arrow pointing from \dot{Y} to \dot{X} . A 2-morphism α is shown as a vertical double arrow pointing upwards from Φ to Φ' , and a 2-morphism α' is shown as a vertical double arrow pointing upwards from Φ' to Φ'' . The vertical composite $\alpha' \circ_v \alpha$ is shown as a vertical double arrow pointing upwards from Φ to Φ'' .

The string notation shows a vertical line representing the 1-morphism Φ from Y to X . A box labeled α is placed on this line, with a vertical line representing the 1-morphism Φ' passing through the box. Another box labeled α' is placed on the Φ' line, with a vertical line representing the 1-morphism Φ'' passing through the box.

The string notation shows a vertical line representing the 1-morphism Φ from Y to X . A box labeled $\alpha' \circ_v \alpha$ is placed on this line, with a vertical line representing the 1-morphism Φ'' passing through the box.

This vertical composition is strictly associative so that $(\alpha'' \circ_v \alpha') \circ_v \alpha = \alpha'' \circ_v (\alpha' \circ_v \alpha)$ whenever the three 2-morphisms are composable. Moreover, there is an identity 2-morphism $\text{Id}_\Phi: \Phi \Rightarrow \Phi$ for every 1-morphism Φ so that $\alpha \circ_v \text{Id}_\Phi = \alpha = \text{Id}_{\Phi'} \circ_v \alpha$ for every 2-morphism $\alpha: \Phi \Rightarrow \Phi'$. This means that for every pair of objects X and Y , the 1-morphisms between them are the objects of a category $\mathbf{Hom}(X, Y)$, with the 2-morphisms forming the morphisms. In the string notation the identity 2-morphisms are usually just drawn as straight lines.

The diagram shows a 1-morphism Φ between objects \dot{Y} and \dot{X} . Φ is represented by a curved arrow pointing from \dot{X} to \dot{Y} . An identity 2-morphism Id_Φ is shown as a vertical double arrow pointing upwards from Φ to Φ .


The string notation shows a vertical line representing the 1-morphism Φ from Y to X .

There is also a composition for 1-morphisms, so if $\Phi: X \rightarrow Y$ and $\Psi: Y \rightarrow Z$ are 1-morphisms then the composite $\Psi \circ \Phi: X \rightarrow Z$ is defined and is denoted as below.

$$\dot{Z} \xleftarrow{\Psi} \dot{Y} \xleftarrow{\Phi} \dot{X} \equiv \dot{Z} \xleftarrow{\Psi \circ \Phi} \dot{X} \qquad \frac{\Psi}{Z \bullet Y} \frac{\Phi}{Y \bullet X} \equiv \frac{\Psi \circ \Phi}{Z \bullet X}$$


Again, these pictures will only appear at the top and bottom of 2-morphisms. This composition of 1-morphisms is not required to be strictly associative, but it is required to be associative up to a *coherent* 2-isomorphism. This means that for every composable triple Θ, Ψ and Φ of 1-morphisms there is a specified 2-isomorphism $(\Theta \circ \Psi) \circ \Phi \xrightarrow{\cong} \Theta \circ (\Psi \circ \Phi)$ and these 2-isomorphisms have to satisfy the so-called pentagon coherency condition which ensures that although $\Theta \circ \Psi \circ \Phi$ is ambiguous, it can be taken to mean either $(\Theta \circ \Psi) \circ \Phi$ or $\Theta \circ (\Psi \circ \Phi)$ without confusion. The up-shot of this is that parentheses are unnecessary in the notation.

Each object X also comes with an identity 1-morphism Id_X , but again, in general, one does not have equality of $\text{Id}_Y \circ \Phi, \Phi$ and $\Phi \circ \text{Id}_X$, but rather the identity 1-morphisms come with coherent 2-isomorphisms $\text{Id}_Y \circ \Phi \xrightarrow{\cong} \Phi$, and $\Phi \circ \text{Id}_X \xrightarrow{\cong} \Phi$. Again this means that in practice the identities can be neglected in the notation: so although we could denote the identity 1-morphism with, say, a dotted line, we choose not to. This is illustrated below.



A *strict* 2-category is one in which the coherency 2-isomorphisms for associativity and identities are themselves all identities. So the 2-category $\mathcal{C}\text{at}$ of categories, functors and natural transformations is a strict 2-category.


The last piece of structure that a 2-category has is the *horizontal composition* of 2-morphisms. If $\Phi, \Phi': X \rightarrow Y$ and $\Psi, \Psi': Y \rightarrow Z$ are 1-morphisms, and $\alpha: \Phi \Rightarrow \Phi'$ and $\beta: \Psi \Rightarrow \Psi'$ are 2-morphisms, then $\beta \circ_h \alpha: \Psi \circ \Phi \Rightarrow \Psi' \circ \Phi'$ is defined, and is notated as below.



The horizontal and vertical composition are required to obey the *interchange law*.

$$(\beta' \circ_v \beta) \circ_h (\alpha' \circ_v \alpha) = (\beta' \circ_h \alpha') \circ_v (\beta \circ_h \alpha).$$

This means that the following diagrams are unambiguous.



It also means that 2-morphisms can be ‘slid past’ each other in the following sense.

From now on, string diagrams will be drawn without the grey borders, and labels will be omitted if they are clear from the context.

1.2. The 2-category $\mathcal{V}\text{ar}$. The 2-category $\mathcal{V}\text{ar}$, of spaces and integral kernels, is defined as follows. The objects are spaces, as defined in the introduction, and the hom-category $\mathbf{Hom}_{\mathcal{V}\text{ar}}(X, Y)$ from a space X to a space Y is the derived category $\mathbf{D}(X \times Y)$, which is to be thought of as the category of integral kernels from X to Y . Explicitly, this means that the 1-morphisms in $\mathcal{V}\text{ar}$ from X to Y are objects of $\mathbf{D}(X \times Y)$ and the 2-morphisms from Φ to Φ' are morphisms in $\mathbf{Hom}_{\mathbf{D}(X \times Y)}(\Phi, \Phi')$, with vertical composition of 2-morphisms just being usual composition in the derived category.

Composition of 1-morphisms in $\mathcal{V}\text{ar}$ is defined using the convolution of integral kernels: if $\Phi \in \mathbf{D}(X \times Y)$ and $\Psi \in \mathbf{D}(Y \times Z)$ are 1-morphisms then define the convolution $\Psi \circ \Phi \in \mathbf{D}(X \times Z)$ by

$$\Phi \circ \Psi := \pi_{XZ,*}(\pi_{YZ}^* \Psi \otimes \pi_{XY}^* \Phi),$$

where π_{XZ} , π_{XY} and π_{YZ} are the projections from $X \times Y \times Z$ to the appropriate factors. The horizontal composition of 2-morphisms is similarly defined. Finally, the identity 1-morphism $\text{Id}_X: X \rightarrow X$ is given by $\mathcal{O}_\Delta \in \mathbf{D}(X \times X)$, the structure sheaf of the diagonal in $X \times X$.

The above 2-category is really what is at the heart of the study of integral transforms, and it is entirely analogous to $\mathcal{A}\text{lg}$, the 2-category of algebras described above. For example, the Hochschild cohomology groups of a space X arise as the second homotopy groups of the 2-category $\mathcal{V}\text{ar}$, at X :

$$\begin{aligned} \mathbf{HH}^\bullet(X) &:= \text{Ext}_{X \times X}^\bullet(\mathcal{O}_\Delta, \mathcal{O}_\Delta) \cong \mathbf{Hom}_{\mathbf{D}(X \times X)}^\bullet(\mathcal{O}_\Delta, \mathcal{O}_\Delta) \\ &=: 2\text{-Hom}_{\mathcal{V}\text{ar}}(\text{Id}_X, \text{Id}_X). \end{aligned}$$

There is a 2-functor from $\mathcal{V}\text{ar}$ to $\mathcal{C}\text{at}$ which encodes integral transforms: this 2-functor sends each space X to its derived category $\mathbf{D}(X)$, sends each kernel $\Phi: X \rightarrow Y$ to the corresponding integral transform $\bar{\Phi}: \mathbf{D}(X) \rightarrow \mathbf{D}(Y)$, and sends each map of kernels to the appropriate natural transformation. Many of the statements about integral transforms have better formulations in the language of the 2-category $\mathcal{V}\text{ar}$.

2. Serre functors

In this section we review the notion of the Serre functor on $\mathbf{D}(X)$ and then show how to realise the Serre functor on the derived category $\mathbf{D}(X \times Y)$ using 2-categorical language.

2.1. The Serre functor on $\mathbf{D}(X)$. If X is a space then we consider the functor

$$S: \mathbf{D}(X) \rightarrow \mathbf{D}(X); \quad \mathcal{E} \mapsto \omega_X[\dim X] \otimes \mathcal{E},$$

where ω_X is the canonical line bundle of X . Serre duality then gives natural, bifunctorial isomorphisms

$$\eta_{\mathcal{E}, \mathcal{F}}: \mathrm{Hom}_{\mathbf{D}(X)}(\mathcal{E}, \mathcal{F}) \xrightarrow{\sim} \mathrm{Hom}_{\mathbf{D}(X)}(\mathcal{F}, S\mathcal{E})^\vee$$

for any objects $\mathcal{E}, \mathcal{F} \in \mathbf{D}(X)$, where $-\vee$ denotes the dual vector space.

A functor such as S , together with isomorphisms as above, was called a *Serre functor* by Bondal and Kapranov [2] (see also [17]). From this data, for any object $\mathcal{E} \in \mathbf{D}(X)$, define the *Serre trace* as follows:

$$\mathrm{Tr}: \mathrm{Hom}(\mathcal{E}, S\mathcal{E}) \rightarrow k; \quad \mathrm{Tr}(\alpha) := \eta_{\mathcal{E}, \mathcal{E}}(\mathrm{Id}_{\mathcal{E}})(\alpha).$$

Note that from this trace we can recover $\eta_{\mathcal{E}, \mathcal{F}}$ because

$$\eta_{\mathcal{E}, \mathcal{F}}(\alpha)(\beta) = \mathrm{Tr}(\beta \circ \alpha).$$

We also have the commutativity identity

$$\mathrm{Tr}(\beta \circ \alpha) = \mathrm{Tr}(S\alpha \circ \beta).$$

Yet another way to encode this data is as a perfect pairing, the *Serre pairing*:

$$\langle -, - \rangle_S: \mathrm{Hom}(\mathcal{E}, \mathcal{F}) \otimes \mathrm{Hom}(\mathcal{F}, S\mathcal{E}) \rightarrow k; \quad \langle \alpha, \beta \rangle_S := \mathrm{Tr}(\beta \circ \alpha).$$

2.2. Serre kernels and the Serre functor for $\mathbf{D}(X \times Y)$. We are interested in kernels and the 2-category $\mathcal{V}\mathrm{ar}$, so are interested in Serre functors for product spaces $X \times Y$, and these have a lovely description in the 2-categorical language. We can now define one of the key objects in this paper.

Definition. For a space X , the *Serre kernel* $\Sigma_X \in 1\text{-Hom}_{\mathcal{V}\mathrm{ar}}(X, X)$ is defined to be $\Delta_*\omega_X[\dim X] \in \mathbf{D}(X \times X)$, the kernel inducing the Serre functor on X . Similarly the *anti-Serre kernel* $\Sigma_X^{-1} \in 1\text{-Hom}_{\mathcal{V}\mathrm{ar}}(X, X)$ is defined to be $\Delta_*\omega_X^{-1}[-\dim X] \in \mathbf{D}(X \times X)$.

Notation. In string diagrams the Serre kernel will be denoted by a dashed-dotted line, while the anti-Serre kernel will be denoted by a dashed-dotted line with a horizontal bar through it. For example, for kernels $\Psi, \Phi: X \rightarrow Y$, a kernel morphism $\alpha: \Phi \circ \Sigma_X \Rightarrow \Sigma_Y \circ \Psi \circ \Sigma_X^{-1}$ will be denoted

$$\begin{array}{c} \vdots \\ \vdots \\ \vdots \\ \Psi \\ \vdots \\ \vdots \\ \vdots \\ \alpha \\ \vdots \\ \vdots \\ \vdots \\ \Phi \\ \vdots \\ \vdots \\ \vdots \end{array}$$

The Serre kernel can now be used to give a natural description, in the 2-category language, of the Serre functor on the product $X \times Y$.

Proposition 1. *For spaces X and Y the Serre functor $S_{X \times Y}: \mathbf{D}(X \times Y) \rightarrow \mathbf{D}(X \times Y)$ can be taken to be $\Sigma_Y \circ - \circ \Sigma_X$.*

Proof. The Serre functor on $\mathbf{D}(X \times Y)$ is given by

$$S_{X \times Y}(\Phi) = \Phi \otimes \pi_X^* \omega_X \otimes \pi_Y^* \omega_Y [\dim X + \dim Y].$$

However, observe that if $\Phi \in \mathbf{D}(X \times Y)$ and $\mathcal{E} \in \mathbf{D}(X)$ then

$$\Phi \circ \Delta_* \mathcal{E} \cong \Phi \otimes \pi_X^* \mathcal{E},$$

where $\pi_X: X \times Y \rightarrow X$ is the projection. This is just a standard application of the base-change and projection formulas. Similarly if $\mathcal{F} \in \mathbf{D}(Y)$ then $\Delta_* \mathcal{F} \circ \Phi \cong \pi_Y^* \mathcal{F} \otimes \Phi$. From this the Serre functor can be written as

$$S_{X \times Y}(\Phi) = \Sigma_Y \circ \Phi \circ \Sigma_X. \quad \square$$

This means that the Serre trace map on $X \times Y$ is a map

$$\mathrm{Tr}: 2\text{-Hom}_{\mathcal{V}\mathrm{ar}}(\Phi, \Sigma_Y \circ \Phi \circ \Sigma_X) \rightarrow k$$

which can be pictured as

$$\mathrm{Tr} \left(\begin{array}{c} \vdots \quad | \quad \vdots \\ \boxed{} \\ \vdots \quad | \quad \vdots \end{array} \right) \in k,$$

where the Serre kernel is denoted by the dashed-dotted line.

We will see below that we have ‘partial trace’ operations which the Serre trace factors through.

3. Adjoint kernels

The reader is undoubtedly familiar with the notion of adjoint functors. It is easy and natural to generalize this from the context of the 2-category $\mathcal{C}\mathrm{at}$ of categories, functors and natural transformations to the context of an arbitrary 2-category. In this section it is shown that every kernel, considered as a 1-morphism in the 2-category $\mathcal{V}\mathrm{ar}$, has both a left and right adjoint: this is a consequence of Serre duality, and is closely related to the familiar fact that every integral transform functor has both a left and right adjoint functor.

Using these notions of left and right adjoints we define *left partial trace maps*, and similarly *right partial trace maps*. These can be viewed as partial versions of the Serre trace map. This construction is very much the heart of the paper.

3.1. Adjunctions in 2-categories. The notion of an adjunction in a 2-category simultaneously generalizes the notion of an adjunction between functors and the notion of a duality between objects of a monoidal category. As it is the former that arises in the context of integral transforms, we will use that as the motivation, but will come back to the latter below.

The most familiar definition of adjoint functors is as follows. For categories \mathbf{C} and \mathbf{D} , an adjunction $\Psi \dashv \Phi$ between functors $\Psi: \mathbf{D} \rightarrow \mathbf{C}$ and $\Phi: \mathbf{C} \rightarrow \mathbf{D}$ is the specification of a natural isomorphism

$$t_{\mathbf{a},\mathbf{b}}: \text{Hom}_{\mathbf{C}}(\Psi(\mathbf{a}), \mathbf{b}) \xrightarrow{\sim} \text{Hom}_{\mathbf{D}}(\mathbf{a}, \Phi(\mathbf{b}))$$

for every $\mathbf{a} \in \mathbf{D}$ and $\mathbf{b} \in \mathbf{C}$.

It is well known (see [8, page 91]) that this definition is equivalent to an alternative definition of adjunction which consists of the specification of unit and counit natural morphisms, namely

$$\eta: \text{Id}_{\mathbf{D}} \Rightarrow \Phi \circ \Psi \quad \text{and} \quad \epsilon: \Psi \circ \Phi \Rightarrow \text{Id}_{\mathbf{C}},$$

such that the composite natural transformations

$$\Psi \xrightarrow{\text{Id}_{\Psi} \circ \eta} \Psi \circ \Phi \circ \Psi \xrightarrow{\epsilon \circ \text{Id}_{\Psi}} \Psi \quad \text{and} \quad \Phi \xrightarrow{\eta \circ \text{Id}_{\Phi}} \Phi \circ \Psi \circ \Phi \xrightarrow{\text{Id}_{\Phi} \circ \epsilon} \Phi$$

are respectively the identity natural transformation on Ψ and the identity natural transformation on Φ .

It is straightforward to translate between the two different definitions of adjunction. Given η and ϵ as above, define

$$t_{\mathbf{a},\mathbf{b}}: \text{Hom}_{\mathbf{C}}(\Psi(\mathbf{a}), \mathbf{b}) \xrightarrow{\sim} \text{Hom}_{\mathbf{D}}(\mathbf{a}, \Phi(\mathbf{b}))$$

by $t_{\mathbf{a},\mathbf{b}}(f) := \Phi(f) \circ \eta_{\mathbf{a}}$. The inverse of $t_{\mathbf{a},\mathbf{b}}$ is defined similarly using the counit ϵ . Conversely, to get the unit and counit from the natural isomorphism of hom-sets, define $\eta_{\mathbf{a}} := t_{\mathbf{a},\Psi(\mathbf{a})}(\text{Id}_{\Psi(\mathbf{a})})$ and define ϵ similarly.

The definition involving the unit and counit is stated purely in terms of functors and natural transformations — without mentioning objects — thus it generalizes immediately to arbitrary 2-categories.

Definition. If \mathcal{C} is a 2-category, X and Y are objects of \mathcal{C} , and

$$\Phi: X \rightarrow Y \quad \text{and} \quad \Psi: Y \rightarrow X$$

are 1-morphisms, then an *adjunction* between Φ and Ψ consists of two 2-morphisms

$$\eta: \text{Id}_Y \Rightarrow \Phi \circ \Psi \quad \text{and} \quad \epsilon: \Psi \circ \Phi \Rightarrow \text{Id}_X,$$

such that

$$(\epsilon \circ_h \text{Id}_{\Psi}) \circ_v (\text{Id}_{\Psi} \circ_h \eta) = \text{Id}_{\Psi} \quad \text{and} \quad (\text{Id}_{\Phi} \circ_h \epsilon) \circ_v (\eta \circ_h \text{Id}_{\Phi}) = \text{Id}_{\Phi}.$$

Given such an adjunction we write $\Psi \dashv \Phi$.

It is worth noting that this also generalizes the notion of duality in a monoidal category, that is to say two objects are dual in a monoidal category if and only if the corresponding 1-morphisms are adjoint in the corresponding 2-category-with-one-object. Indeed, taking this point of view, May and Sigurdsson [12] refer to what is here called adjunction as duality.

It is at this point that the utility of the string diagram notation begins to be seen. Given an adjunction $\Psi \dashv \Phi$ the counit $\epsilon: \Psi \circ \Phi \Rightarrow \text{Id}$ and the unit $\eta: \text{Id} \Rightarrow \Phi \circ \Psi$ can be denoted as follows:

$$\begin{array}{c} \text{Id}_X \\ \vdots \\ \boxed{\epsilon} \\ \Psi \quad \Phi \end{array} \quad \text{and} \quad \begin{array}{c} \Phi \quad \Psi \\ \boxed{\eta} \\ \vdots \\ \text{Id}_Y \end{array} .$$

However, adopting the convention of denoting the identity one-morphism by omission, it is useful just to draw the unit and counit as a cup and a cap respectively:

$$\begin{array}{c} \Psi \quad \Phi \\ \cup \end{array} := \begin{array}{c} \text{Id}_X \\ \vdots \\ \boxed{\epsilon} \\ \Psi \quad \Phi \end{array} \quad \text{and} \quad \begin{array}{c} \Phi \quad \Psi \\ \cup \end{array} := \begin{array}{c} \Phi \quad \Psi \\ \boxed{\eta} \\ \vdots \\ \text{Id}_Y \end{array} .$$

The relations become the satisfying

$$\begin{array}{c} \Psi \quad \Phi \\ \cup \end{array} = \begin{array}{c} \Psi \\ | \\ \Psi \end{array} \quad \text{and} \quad \begin{array}{c} \Phi \quad \Psi \\ \cup \end{array} = \begin{array}{c} \Phi \\ | \\ \Phi \end{array} .$$

Adjunctions in 2-categories, as defined above, do correspond to isomorphisms of certain hom-sets but in a different way to the classical notion of adjunction. Namely, if $\Theta: Z \rightarrow Y$ and $\Xi: Z \rightarrow X$ are two other 1-morphisms, then an adjunction $\Psi \dashv \Phi$ as above gives an isomorphism

$$\begin{array}{c} \Xi \\ | \\ \boxed{\alpha} \\ \Psi \quad \Theta \end{array} \mapsto \begin{array}{c} \Phi \quad \Xi \\ \cup \\ \boxed{\alpha} \\ \Psi \quad \Theta \end{array} .$$

The inverse isomorphism uses the counit in the obvious way.


In a similar fashion, for $\hat{\Theta}: Y \rightarrow Z$ and $\hat{\Xi}: X \rightarrow Z$ two 1-morphisms, one obtains an isomorphism

$$2\text{-Hom}(\hat{\Theta} \circ \Phi, \hat{\Xi}) \xrightarrow{\sim} 2\text{-Hom}(\hat{\Theta}, \hat{\Xi} \circ \Psi),$$

for which the reader is encouraged to draw the relevant pictures. It is worth noting that with respect to the previous isomorphism, Ψ and Φ have swapped sides in all senses.

Adjunctions are unique up to a canonical isomorphism by the usual argument. This means that if Ψ and Ψ' are, say, both left adjoint to Φ , then there is a canonical isomorphism $\Psi \xrightarrow{\sim} \Psi'$. This is pictured below and it is

easy to check that this is an isomorphism.



Adjunctions are natural in the sense that they are preserved by 2-functors, so, for instance, given a pair of adjoint kernels in $\mathcal{V}ar$, the corresponding integral transforms are adjoint functors.

3.2. Left and right adjoints of kernels. In an arbitrary 2-category a given 1-morphism might or might not have a left or a right adjoint, but in the 2-category $\mathcal{V}ar$ every 1-morphism, that is every kernel, has both a left and a right adjoint. We will see below that for a kernel $\Phi: X \rightarrow Y$ there are adjunctions

$$\Phi^\vee \circ \Sigma_Y \dashv \Phi \dashv \Sigma_X \circ \Phi^\vee,$$

where $\Phi^\vee: Y \rightarrow X$ means the object $\underline{\text{Hom}}_{\mathbf{D}(X \times Y)}(\Phi, \mathcal{O}_{X \times Y})$ considered as an object in $\mathbf{D}(Y \times X)$. This should be compared with the fact that if M is an A - B -bimodule then M^\vee is naturally a B - A -bimodule. We shall see that the two adjunctions above are related in some very useful ways.

Proposition 2. *If X is a space and $\Delta: X \rightarrow X \times X$ is the diagonal embedding then $\Delta_*: \mathbf{D}(X) \rightarrow \mathbf{D}(X \times X)$, the push-forward on derived categories, is a monoidal functor where $\mathbf{D}(X)$ has the usual monoidal tensor product \otimes and $\mathbf{D}(X \times X)$ has the composition \circ as the monoidal structure.*


Proof. The proof is just an application of the projection formula. □

This has the following immediate consequence.


Lemma 3. *If \mathcal{E} and \mathcal{F} are dual as objects in $\mathbf{D}(X)$ then $\Delta_*\mathcal{E}$ and $\Delta_*\mathcal{F}$ are both left and right adjoint to each other as 1-morphisms in $\mathcal{V}ar$.*

In order not to hold-up the flow of the narrative, the proofs of the remaining results from this section have been relegated to Appendix A.

We begin with some background on the Serre kernel Σ_X . Recall from Section 2 that the anti-Serre kernel Σ_X^{-1} is defined to be $\Delta_*\omega_X^{-1}[-\dim X]$, and that the Serre kernel Σ_X is denoted by a dashed-dotted line, while the anti-Serre kernel Σ_X^{-1} is denoted by a dashed-dotted line with a horizontal bar. As ω and ω^{-1} are inverse with respect to \otimes the above proposition means that Σ_X and Σ_X^{-1} are inverse with respect to \circ , thus we have maps



satisfying the following relations



and all obvious variations thereof.

In the appendix it is shown that for a kernel Φ there are natural morphisms $\epsilon_\Phi: \Phi \circ \Sigma_X \circ \Phi^\vee \rightarrow \mathcal{O}_\Delta$ (pronounced “mepsilon”) and $\gamma_\Phi: \Sigma_X^{-1} \rightarrow \Phi^\vee \circ \Phi$, denoted in the following fashion, where the solid, upward oriented lines are labelled with Φ and the solid, implicitly downward oriented lines are labelled by Φ^\vee :

$$\epsilon_\Phi: \begin{array}{c} \text{---} \\ \uparrow \\ \text{---} \end{array} \text{ and } \gamma_\Phi: \begin{array}{c} \text{---} \\ \downarrow \\ \text{---} \end{array}.$$

The main property of ϵ_Φ and γ_Φ is that if we define $\epsilon_\Phi, \bar{\epsilon}_\Phi, \eta_\Phi$ and $\bar{\eta}_\Phi$ via

$$\begin{array}{cc} \eta_\Phi: \begin{array}{c} \text{---} \\ \uparrow \\ \text{---} \end{array} := \begin{array}{c} \text{---} \\ \uparrow \\ \text{---} \end{array}, & \bar{\eta}_\Phi: \begin{array}{c} \text{---} \\ \downarrow \\ \text{---} \end{array} := \begin{array}{c} \text{---} \\ \downarrow \\ \text{---} \end{array}, \\ \epsilon_\Phi := \epsilon_{\Phi^\vee} = \begin{array}{c} \text{---} \\ \uparrow \\ \text{---} \end{array}, & \bar{\epsilon}_\Phi := \epsilon_\Phi = \begin{array}{c} \text{---} \\ \uparrow \\ \text{---} \end{array}, \end{array}$$

then these are the units and counits of adjunctions

$$\Phi^\vee \circ \Sigma_Y \dashv \Phi \dashv \Sigma_X \circ \Phi^\vee.$$

3.3. Partial traces. We can now define the important notion of partial traces.

Definition. For a kernel $\Phi: X \rightarrow Y$, and kernels $\Psi, \Theta: Z \rightarrow X$ define the *left partial trace*

$$2\text{-Hom}(\Phi \circ \Theta, \Sigma_Y \circ \Phi \circ \Psi) \rightarrow 2\text{-Hom}(\Theta, \Sigma_X \circ \Psi)$$

as

$$\begin{array}{c} \Psi \\ \uparrow \\ \alpha \\ \uparrow \\ \Theta \end{array} \mapsto \begin{array}{c} \Psi \\ \uparrow \\ \alpha \\ \uparrow \\ \Theta \end{array}.$$

Similarly we define a *right partial trace*

$$2\text{-Hom}(\Theta' \circ \Phi, \Psi' \circ \Phi \circ \Sigma_X) \rightarrow 2\text{-Hom}(\Theta', \Psi' \circ \Sigma_Y)$$

as

$$\begin{array}{c} \Psi' \\ \uparrow \\ \alpha' \\ \uparrow \\ \Theta' \end{array} \mapsto \begin{array}{c} \Psi' \\ \uparrow \\ \alpha' \\ \uparrow \\ \Theta' \end{array}.$$

The following key result, proved in the appendix, says that taking partial trace does not affect the Serre trace.

Theorem 4. *For a kernel $\Phi: X \rightarrow Y$, a kernel $\Psi: Z \rightarrow X$ and a kernel morphism $\alpha \in \text{Hom}(\Phi \circ \mathcal{F}, \Sigma_Y \circ \Phi \circ \Psi \circ \Sigma_Z)$ then the left partial trace of α has the same Serre trace as α , i.e., pictorially*

$$\text{Tr} \left(\begin{array}{c} \Psi \\ \uparrow \\ \alpha \\ \downarrow \\ \Psi \end{array} \right) = \text{Tr} \left(\begin{array}{c} \Psi \\ \uparrow \\ \alpha \\ \uparrow \\ \Psi \end{array} \right).$$

The analogous result holds for the right partial trace.

3.4. Adjunction as a 2-functor. As shown in Section 3.2, in the 2-category $\mathcal{V}\text{ar}$ every 1-morphism, that is every kernel, $\Phi: X \rightarrow Y$ has a right adjoint $\Sigma_X \circ \Phi^\vee$. This can be extended to a ‘right adjunction 2-functor’ $\tau_R: \mathcal{V}\text{ar}^{\text{coop}} \rightarrow \mathcal{V}\text{ar}$, where $\mathcal{V}\text{ar}^{\text{coop}}$ means the contra-opposite 2-category of $\mathcal{V}\text{ar}$, which is the 2-category with the same collections of objects, morphisms and 2-morphisms, but in which the direction of the morphisms and the 2-morphisms are reversed.

Before defining τ_R , however, it is perhaps useful to think of the more familiar situation of a one-object 2-category with right adjoints, i.e., a monoidal category with (right) duals. So if \mathcal{C} is a monoidal category in which each object \mathfrak{a} has a dual \mathfrak{a}^\vee with evaluation map $\epsilon_{\mathfrak{a}}: \mathfrak{a}^\vee \otimes \mathfrak{a} \rightarrow \mathbf{1}$ and coevaluation map $\eta_{\mathfrak{a}}: \mathbf{1} \rightarrow \mathfrak{a} \otimes \mathfrak{a}^\vee$, then for any morphism $f: \mathfrak{a} \rightarrow \mathfrak{b}$ define $f^\vee: \mathfrak{b}^\vee \rightarrow \mathfrak{a}^\vee$ to be the composite:

$$\mathfrak{b}^\vee \xrightarrow{\text{Id} \otimes \eta} \mathfrak{b}^\vee \otimes \mathfrak{a} \otimes \mathfrak{a}^\vee \xrightarrow{\text{Id} \otimes f \otimes \text{Id}} \mathfrak{b}^\vee \otimes \mathfrak{b} \otimes \mathfrak{a}^\vee \xrightarrow{\epsilon \otimes \text{Id}} \mathfrak{a}^\vee.$$

This gives rise to a functor $(-)^\vee: \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}$.

Now return to the case of interest and define $\tau_R: \mathcal{V}\text{ar}^{\text{coop}} \rightarrow \mathcal{V}\text{ar}$ as follows. On spaces define $\tau_R(X) := X$. On a kernel $\Phi: X \rightarrow Y$ define $\tau_R(\Phi) := \Sigma_X \circ \Phi^\vee$. Finally, on morphisms of kernels define it as illustrated:

$$\tau_R \left(\begin{array}{c} \uparrow \\ \Phi' \\ \alpha \\ \uparrow \\ \Phi \end{array} \right) := \begin{array}{c} \Phi' \\ \uparrow \\ \alpha \\ \uparrow \\ \Phi \end{array}$$

It is a nice exercise for the reader to check that this is a 2-functor.

Clearly a left adjoint 2-functor $\tau_L: \mathcal{V}\text{ar}^{\text{coop}} \rightarrow \mathcal{V}\text{ar}$ can be similarly created by defining it on a kernel $\Phi: X \rightarrow Y$ by $\tau_L(\Phi) := \Phi^\vee \circ \Sigma_Y$ and by defining it on morphisms of kernels by

$$\tau_L \left(\begin{array}{c} \uparrow \\ \Phi' \\ \alpha \\ \uparrow \\ \Phi \end{array} \right) := \begin{array}{c} \Phi' \\ \uparrow \\ \alpha \\ \uparrow \\ \Phi \end{array}$$

4. Induced maps on homology

In this section we define $\mathrm{HH}_\bullet(X)$, the Hochschild homology of a space X , and show that given a kernel $\Phi: X \rightarrow Y$ we get pull-back and push-forward maps, $\Phi^*: \mathrm{HH}_\bullet(Y) \rightarrow \mathrm{HH}_\bullet(X)$ and $\Phi_*: \mathrm{HH}_\bullet(X) \rightarrow \mathrm{HH}_\bullet(Y)$, such that if Φ is right adjoint to Ψ then $\Psi^* = \Phi_*$.

4.1. Hochschild cohomology. First recall that for a space X , one way to define its Hochschild *cohomology* is as

$$\mathrm{HH}^\bullet(X) := \mathrm{Ext}_{X \times X}^\bullet(\mathcal{O}_\Delta, \mathcal{O}_\Delta).$$

However, the ext-group is just the hom-set $\mathrm{Hom}_{\mathbf{D}(X \times X)}^\bullet(\mathcal{O}_\Delta, \mathcal{O}_\Delta)$ which by the definition of $\mathcal{V}\mathrm{ar}$ is just $2\text{-}\mathrm{Hom}_{\mathcal{V}\mathrm{ar}}^\bullet(\mathrm{Id}_X, \mathrm{Id}_X)$. In terms of diagrams, we can thus denote an element $\varphi \in \mathrm{HH}^\bullet(X)$ as

$$\boxed{\varphi}.$$

Note that the grading is not indicated in the picture, but this should not give rise to confusion.

4.2. Hochschild homology. Now we define $\mathrm{HH}_\bullet(X)$ the Hochschild *homology* of a space X as follows:

$$\mathrm{HH}_\bullet(X) := 2\text{-}\mathrm{Hom}_{\mathcal{V}\mathrm{ar}}^\bullet(\Sigma_X^{-1}, \mathrm{Id}_X)$$

or, in more concrete terms,

$$\mathrm{HH}_\bullet(X) = \mathrm{Ext}_{X \times X}^\bullet(\Sigma_X^{-1}, \mathcal{O}_\Delta).$$

Thus an element $w \in \mathrm{HH}_\bullet(X)$ will be denoted

$$\boxed{w},$$

where again the shifts are understood.

It is worth taking a moment to compare this with other definitions of Hochschild homology, such as that of Weibel [19]. He defines the Hochschild homology of a space X as $\mathrm{H}^\bullet(X, \Delta^* \mathcal{O}_\Delta)$, where as usual by Δ^* we mean the left-derived functor. This cohomology group is naturally identified with the hom-set $\mathrm{Hom}_X^\bullet(\mathcal{O}_X, \Delta^* \mathcal{O}_\Delta)$ which is isomorphic to $\mathrm{Hom}_{X \times X}^\bullet(\Delta_! \mathcal{O}_X, \mathcal{O}_\Delta)$ where $\Delta_!$ is the left-adjoint of Δ^* . Direct calculation shows that $\Delta_! \mathcal{O}_X \cong \Sigma_X^{-1}$ and so our definition is recovered. Another feasible definition of Hochschild homology is $\mathrm{H}^\bullet(X \times X, \mathcal{O}_\Delta \otimes \mathcal{O}_\Delta)$, and this again is equivalent to our definition as there is the isomorphism $\Sigma_X^{-1} \cong \mathcal{O}_\Delta^\vee$.

4.3. Push-forward and pull-back. For spaces X and Y and a kernel $\Phi: X \rightarrow Y$ define the push-forward on Hochschild homology

$$\Phi_*: \mathrm{HH}_\bullet(X) \rightarrow \mathrm{HH}_\bullet(Y)$$

as follows:

$$\Phi_* \left(\begin{array}{c} \boxed{w} \\ \vdots \end{array} \right) := \begin{array}{c} \text{---} \circlearrowleft \text{---} \\ \text{---} \boxed{w} \text{---} \\ \text{---} \circlearrowright \text{---} \\ \vdots \end{array} \cdot \Phi.$$

For the reader still unhappy with diagrams, for $v \in \mathrm{Hom}^\bullet(\Sigma_X^{-1}, \mathrm{Id}_X)$, define $\Phi_*(v) \in \mathrm{Hom}^\bullet(\Sigma_Y^{-1}, \mathrm{Id}_Y)$ as the following composite, which is read from the above diagram by reading upwards from the bottom:

$$\Sigma_Y^{-1} \xrightarrow{\gamma} \Phi \circ \Phi^\vee \xrightarrow{\mathrm{Id} \circ \eta^{\mathrm{old}}} \Phi \circ \Sigma_X^{-1} \circ \Sigma_X \circ \Phi^\vee \xrightarrow{\mathrm{Id} \circ \mathrm{vold} \circ \mathrm{old}} \Phi \circ \Sigma_X \circ \Phi^\vee \xrightarrow{\circ} \mathrm{Id}_Y.$$

Similarly define the pull-back $\Phi^*: \mathrm{HH}_\bullet(Y) \rightarrow \mathrm{HH}_\bullet(X)$ as follows:

$$\Phi^* \left(\begin{array}{c} \boxed{v} \\ \vdots \end{array} \right) := \begin{array}{c} \text{---} \circlearrowleft \text{---} \\ \text{---} \boxed{v} \text{---} \\ \text{---} \circlearrowright \text{---} \\ \vdots \end{array} \cdot \Phi.$$

These operations depend only on the isomorphism class of the kernel as shown by the following.

Proposition 5. *If kernels Φ and $\hat{\Phi}$ are isomorphic then they give rise to equal push-forwards and equal pull-backs: $\Phi_* = \hat{\Phi}_*$ and $\Phi^* = \hat{\Phi}^*$.*

Proof. This follows immediately from the fact that the 2-morphisms γ_Φ , $\gamma_{\hat{\Phi}}$, \circ_Φ and $\circ_{\hat{\Phi}}$ of Section 3.2 are natural and thus commute with the given kernel isomorphism $\Phi \cong \hat{\Phi}$. \square

The push-forward and pull-back operations are functorial in the following sense.

Theorem 6 (Functoriality). *If $\Phi: X \rightarrow Y$ and $\Psi: Y \rightarrow Z$ are kernels then the push-forwards and pull-backs compose appropriately, namely:*

$$(\Psi \circ \Phi)_* = \Psi_* \circ \Phi_*: \mathrm{HH}_\bullet(X) \rightarrow \mathrm{HH}_\bullet(Z)$$

and

$$(\Psi \circ \Phi)^* = \Phi^* \circ \Psi^*: \mathrm{HH}_\bullet(Z) \rightarrow \mathrm{HH}_\bullet(X).$$

Proof. This follows from the fact that the right adjunction τ_R is a 2-functor. The adjoint of $\Psi \circ \Phi$ is canonically $\tau_R(\Phi) \circ \tau_R(\Psi)$, i.e., is $\Sigma_X \circ \Phi^\vee \circ \Sigma_Y \circ \Psi^\vee$. This means that the unit of the adjunction $\mathrm{Id} \Rightarrow \Psi \circ \Phi \circ \Sigma_X \circ \Phi^\vee \circ \Sigma_Y \circ \Psi^\vee$

Note that this differs slightly from the given definition, but we have used the uniqueness of adjoints. The above isomorphisms allow the definition of the Mukai pairing as follows.

Definition. The *Mukai pairing* on the Hochschild homology of a space X is the map

$$\langle -, - \rangle_M: \mathrm{HH}_\bullet(X) \otimes \mathrm{HH}_\bullet(X) \rightarrow k,$$

defined by

$$\langle v, v' \rangle_M := \mathrm{Tr}(\tau_R(v) \circ \tau_L(v')).$$

Diagrammatically, this is

$$\left\langle \begin{array}{c} \boxed{v} \\ \vdots \\ \vdots \end{array}, \begin{array}{c} \boxed{v'} \\ \vdots \\ \vdots \end{array} \right\rangle_M := \mathrm{Tr} \left(\begin{array}{c} \boxed{v} \quad \boxed{v'} \\ \vdots \quad \vdots \\ \vdots \quad \vdots \end{array} \right).$$

Observe that as τ_R and τ_L are both isomorphisms and as the Serre pairing is nondegenerate, it follows that the Mukai pairing is nondegenerate.

We can now easily show that adjoint kernels give rise to adjoint maps between the corresponding Hochschild homology groups.

Theorem 8 (Adjointness). *If $\Phi: X \rightarrow Y$ and $\Psi: Y \rightarrow X$ are adjoint kernels, $\Psi \dashv \Phi$, then the corresponding push forwards are adjoint with respect to the Mukai pairing in the sense that for all $w \in \mathrm{HH}_\bullet(X)$ and $v \in \mathrm{HH}_\bullet(Y)$ we have*

$$\langle \Psi_*(v), w \rangle_M = \langle v, \Phi_*(w) \rangle_M.$$

Proof. Note first that $\Psi_* = \Phi^*$, by Theorem 7. Thus

$$\begin{aligned} \langle \Psi_*(v), w \rangle_M &= \langle \Phi^*(v), w \rangle_M = \mathrm{Tr} \left(\begin{array}{c} \boxed{v} \quad \boxed{w} \\ \vdots \quad \vdots \\ \vdots \quad \vdots \end{array} \right) \\ &= \mathrm{Tr} \left(\begin{array}{c} \boxed{v} \quad \boxed{w} \\ \vdots \quad \vdots \\ \vdots \quad \vdots \end{array} \right) = \mathrm{Tr} \left(\begin{array}{c} \boxed{v} \quad \boxed{w} \\ \vdots \quad \vdots \\ \vdots \quad \vdots \end{array} \right) \\ &= \langle v, \Phi_*(w) \rangle_M. \quad \square \end{aligned}$$

Corollary 9. *If the integral kernel $\Phi: X \rightarrow Y$ induces an equivalence on derived categories, then $\Phi_*: \mathrm{HH}_\bullet(X) \rightarrow \mathrm{HH}_\bullet(Y)$ is an isometry.*

Proof. If Φ induces an equivalence, then it has a left adjoint $\Psi: Y \rightarrow X$ which induces the inverse, so $\Psi \circ \Phi \cong \mathrm{Id}_X$, and we know that $(\mathrm{Id}_X)_*$ is the identity map. Thus

$$\langle \Phi_*v, \Phi_*w \rangle_M = \langle \Psi_*\Phi_*v, w \rangle_M = \langle (\Psi \circ \Phi)_*v, w \rangle_M = \langle v, w \rangle_M. \quad \square$$

6. The Chern character

In this section we define the Chern character map $\text{ch}: K_0(X) \rightarrow \text{HH}_0(X)$. We discuss the relationship between our construction and the one of Markarian [10, Definition 2]. Then we show that the Chern character maps the Euler pairing to the Mukai pairing: we call this the Semi-Hirzebruch–Riemann–Roch Theorem.

6.1. Definition of the Chern character. Suppose X is a space, and \mathcal{E} is an object in $\mathbf{D}(X)$. Consider \mathcal{E} as an object of $\mathbf{D}(\text{pt} \times X)$, i.e., as a kernel $\text{pt} \rightarrow X$, so there is an induced linear map

$$\mathcal{E}_*: \text{HH}_\bullet(\text{pt}) \rightarrow \text{HH}_\bullet(X).$$

Now, because the Serre functor on a point is trivial, $\text{HH}_0(\text{pt})$ is canonically identifiable with $\text{Hom}_{\text{pt} \times \text{pt}}(\mathcal{O}_{\text{pt}}, \mathcal{O}_{\text{pt}})$ so there is a distinguished class $\mathbf{1} \in \text{HH}_0(\text{pt})$ corresponding to the identity map. Define the Chern character of \mathcal{E} as

$$\text{ch}(\mathcal{E}) := \mathcal{E}_*(\mathbf{1}) \in \text{HH}_0(X).$$

Graphically this has the following description:

$$\text{ch}(\mathcal{E}) := \begin{array}{c} \circlearrowleft \\ \mathcal{E} \uparrow \\ \vdots \end{array}$$

Naturality of push-forward leads to the next theorem.

Theorem 10. *If X and Y are spaces and $\Phi: X \rightarrow Y$ is a kernel then the diagram below commutes.*

$$\begin{array}{ccc} \mathbf{D}(X) & \xrightarrow{\Phi \circ -} & \mathbf{D}(Y) \\ \downarrow \text{ch} & & \downarrow \text{ch} \\ \text{HH}_0(X) & \xrightarrow{\Phi_*} & \text{HH}_0(Y). \end{array}$$

Proof. Let \mathcal{E} be an object of $\mathbf{D}(X)$. We will regard it either as an object in $\mathbf{D}(X)$, or as a kernel $\text{pt} \rightarrow X$, and similarly we will regard $\Phi \circ \mathcal{E}$ either as an object in $\mathbf{D}(Y)$ or as a kernel $\text{pt} \rightarrow Y$. By Theorem 6 we have

$$\Phi_* \text{ch}(\mathcal{E}) = \Phi_*(\mathcal{E}_*(\mathbf{1})) = (\Phi \circ \mathcal{E})_* \mathbf{1} = \text{ch}(\Phi \circ \mathcal{E}). \quad \square$$

6.2. The Chern character as a map on K-theory. To show that the Chern character descends to a map on K-theory we give a characterization of the Chern character similar to that of Markarian [10].

For any object $\mathcal{E} \in \mathbf{D}(X)$, which is to be considered an object of $\mathbf{D}(\text{pt} \times X)$, there are the following two maps:

$$\iota_{\mathcal{E}}: \text{HH}_\bullet(X) \rightarrow \text{Hom}_{\mathbf{D}(X \times X)}^\bullet(\mathcal{E}, \Sigma_X \circ \mathcal{E}); \quad \begin{array}{c} \boxed{v} \\ \vdots \end{array} \mapsto \begin{array}{c} \boxed{v} \\ \vdots \end{array} \begin{array}{c} \circlearrowleft \\ \mathcal{E} \end{array}$$

and

$$\iota_{\mathcal{E}}^{\bullet} : \mathrm{Hom}_{\mathbf{D}(X \times X)}^{\bullet}(\mathcal{E}, \mathcal{E}) \rightarrow \mathrm{HH}_{\bullet}(X); \quad \begin{array}{c} \uparrow \\ \boxed{\varphi} \\ \downarrow \\ \mathcal{E} \end{array} \mapsto \begin{array}{c} \uparrow \\ \boxed{\varphi} \text{ pt} \\ \downarrow \\ \mathcal{E} \\ \downarrow \\ X \end{array}.$$

Recall that the Mukai pairing is a nondegenerate pairing on $\mathrm{HH}_{\bullet}(X)$ and that the Serre pairing is a perfect pairing between $\mathrm{Hom}_{\mathbf{D}(X \times X)}^{\bullet}(\mathcal{E}, \Sigma_X \circ \mathcal{E})$ and $\mathrm{Hom}_{\mathbf{D}(X \times X)}^{\bullet}(\mathcal{E}, \mathcal{E})$. With respect to these pairings the two maps $\iota_{\mathcal{E}}^{\bullet}$ and $\iota_{\mathcal{E}}$ are adjoint in the following sense.

Proposition 11. *For $\varphi \in \mathrm{Hom}^{\bullet}(\mathcal{E}, \mathcal{E})$ and $v \in \mathrm{HH}_{\bullet}(X)$ the following equality holds:*

$$\langle v, \iota_{\mathcal{E}}^{\bullet} \varphi \rangle_M = \langle \iota_{\mathcal{E}} v, \varphi \rangle_S.$$

Proof. Here in the third equality we use the invariance of the Serre trace under the partial trace map.

$$\begin{aligned} \langle v, \iota_{\mathcal{E}}^{\bullet} \varphi \rangle_M &= \left\langle \begin{array}{c} \boxed{v} \\ \downarrow \\ \mathcal{E} \end{array}, \begin{array}{c} \boxed{\varphi} \text{ pt} \\ \downarrow \\ \mathcal{E} \\ \downarrow \\ X \end{array} \right\rangle_M := \mathrm{Tr} \left(\begin{array}{c} X \\ \downarrow \\ \boxed{v} \\ \downarrow \\ \mathcal{E} \end{array} \begin{array}{c} \boxed{\varphi} \text{ pt} \\ \downarrow \\ \mathcal{E} \\ \downarrow \\ X \end{array} \right) \\ &= \mathrm{Tr} \left(\begin{array}{c} X \\ \downarrow \\ \boxed{v} \\ \downarrow \\ \mathcal{E} \end{array} \begin{array}{c} \uparrow \\ \boxed{\varphi} \\ \downarrow \\ \mathcal{E} \\ \downarrow \\ \text{pt} \end{array} \right) = \mathrm{Tr} \left(\begin{array}{c} \boxed{v} \\ \downarrow \\ X \end{array} \begin{array}{c} \uparrow \\ \boxed{\varphi} \\ \downarrow \\ \mathcal{E} \\ \downarrow \\ \text{pt} \end{array} \right) \\ &=: \left\langle \begin{array}{c} \boxed{v} \\ \downarrow \\ \mathcal{E} \end{array}, \begin{array}{c} \uparrow \\ \boxed{\varphi} \\ \downarrow \\ \mathcal{E} \end{array} \right\rangle_S =: \langle \iota_{\mathcal{E}} v, \varphi \rangle_S. \quad \square \end{aligned}$$

Note that, using this, the Chern character could have been defined as

$$\mathrm{ch}(\mathcal{E}) := \iota_{\mathcal{E}}^{\bullet}(\mathrm{Id}_{\mathcal{E}}).$$

Then from the above proposition the following is immediate.

Lemma 12. *For $v \in \mathrm{HH}_0(X)$ and $\mathcal{E} \in \mathbf{D}(X)$ there is the equality*

$$\langle v, \mathrm{ch}(\mathcal{E}) \rangle_M = \mathrm{Tr}(\iota_{\mathcal{E}}(v)),$$

and this defines $\mathrm{ch}(\mathcal{E})$ uniquely.

The fact that the Chern character descends to a function on the K-group can now be demonstrated.

Proposition 13. *For $\mathcal{E} \in \mathbf{D}(X)$ the Chern character $\mathrm{ch}(\mathcal{E})$ depends only on the class of \mathcal{E} in $K_0(X)$. Thus the Chern character can be considered as a map*

$$\mathrm{ch}: K_0(X) \rightarrow \mathrm{HH}_0(X).$$

Proof. It suffices to show that if $\mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H} \rightarrow \mathcal{F}[1]$ is an exact triangle in $\mathbf{D}(X)$, then

$$\mathrm{ch}(\mathcal{F}) - \mathrm{ch}(\mathcal{G}) + \mathrm{ch}(\mathcal{H}) = 0$$

in $\mathrm{HH}_0(X)$.

For $\mathcal{G}, \mathcal{H} \in \mathbf{D}(X)$, $\alpha: \mathcal{G} \rightarrow \mathcal{H}$ and $v \in \mathrm{HH}_\bullet(X)$ the diagram on the left commutes as it expresses the equality on the right:

$$\begin{array}{ccc} \mathcal{G} & \xrightarrow{\alpha} & \mathcal{H} \\ \downarrow \iota_{\mathcal{G}}(v) & & \downarrow \iota_{\mathcal{H}}(v) \\ \Sigma_X \circ \mathcal{G} & \xrightarrow{\mathrm{Id}_{\Sigma_X} \circ \alpha} & \Sigma_X \circ \mathcal{H} \end{array} \quad ; \quad \begin{array}{c} \boxed{v} \\ \downarrow \\ \boxed{\alpha} \\ \downarrow \\ \mathcal{G} \end{array} \xrightarrow{\mathcal{H}} \begin{array}{c} \boxed{\alpha} \\ \downarrow \\ \mathcal{H} \\ \downarrow \\ \boxed{v} \end{array}.$$

In other words, from an element $v \in \mathrm{HH}_\bullet(X)$ we get $\tau_{\mathbf{R}}(v) \in \mathrm{Hom}(\mathrm{Id}_X, \Sigma_X)$, which in turn gives rise to a natural transformation between the functors $\mathrm{Id}_{\mathbf{D}(X)}: \mathbf{D}(X) \rightarrow \mathbf{D}(X)$ and $\Sigma_X \circ -: \mathbf{D}(X) \rightarrow \mathbf{D}(X)$. This leads to a map of triangles

$$\begin{array}{ccccccc} \mathcal{F} & \longrightarrow & \mathcal{G} & \longrightarrow & \mathcal{H} & \longrightarrow & \mathcal{F}[1] \\ \downarrow \iota_{\mathcal{F}}(v) & & \downarrow \iota_{\mathcal{G}}(v) & & \downarrow \iota_{\mathcal{H}}(v) & & \downarrow \iota_{\mathcal{F}}(v)[1] \\ \Sigma_X \mathcal{F} & \longrightarrow & \Sigma_X \mathcal{G} & \longrightarrow & \Sigma_X \mathcal{H} & \longrightarrow & \Sigma_X \mathcal{F}[1]. \end{array}$$

Observe that if we represent the morphism v by an actual map of complexes of injectives, and the objects \mathcal{F} , \mathcal{G} and \mathcal{H} by complexes of locally free sheaves, then the resulting maps in the above diagram commute on the nose (no further injective or locally free resolutions are needed), so we can apply [11, Theorem 1.9] to get

$$\mathrm{Tr}_X(\iota_{\mathcal{F}}(v)) - \mathrm{Tr}_X(\iota_{\mathcal{G}}(v)) + \mathrm{Tr}_X(\iota_{\mathcal{H}}(v)) = 0.$$

Therefore, by the lemma above, for any $v \in \mathrm{HH}_\bullet(X)$,

$$\langle v, \mathrm{ch}(\mathcal{F}) - \mathrm{ch}(\mathcal{G}) + \mathrm{ch}(\mathcal{H}) \rangle_{\mathbf{M}} = 0.$$

Since the Mukai pairing on $\mathrm{HH}_0(X)$ is nondegenerate, we conclude that

$$\mathrm{ch}(\mathcal{F}) - \mathrm{ch}(\mathcal{G}) + \mathrm{ch}(\mathcal{H}) = 0. \quad \square$$

6.3. The Chern character and inner products. One reading of the Hirzebruch–Riemann–Roch Theorem is that it says that the usual Chern character map $\mathrm{ch}: K_0 \rightarrow \mathbf{H}^\bullet(X)$ is a map of inner product spaces when $K_0(X)$ is equipped with the Euler pairing (see below) and $\mathbf{H}^\bullet(X)$ is equipped with the pairing $\langle x_1, x_2 \rangle := (x_1 \cup x_2 \cup \mathrm{td}_X) \cap [X]$. It is shown in [5] that the Hochschild homology Chern character composed with the Hochschild–Kostant–Rosenberg map I_{HKR} gives the usual Chern character:

$$K_0 \xrightarrow{\mathrm{ch}} \mathrm{HH}_0(X) \xrightarrow{I_{\mathrm{HKR}}} \bigoplus_{\mathbf{p}} \mathrm{H}^{\mathbf{p}, \mathbf{p}}(X).$$

Here we show that the Hochschild homology Chern character is an inner product map when $\mathrm{HH}_\bullet(X)$ is equipped with the Mukai pairing.

First recall that the Euler pairing on $K_0(X)$ is defined by

$$\chi(\mathcal{E}, \mathcal{F}) := \sum_i (-1)^i \dim \text{Ext}_X^i(\mathcal{E}, \mathcal{F}).$$

Theorem 14 (Semi-Hirzebruch–Riemann–Roch). *The Chern character*

$$\text{ch}: K_0 \rightarrow \text{HH}_0(X)$$

is a map of inner product spaces: in other words, for $\mathcal{E}, \mathcal{F} \in \mathbf{D}(X)$ we have

$$\langle \text{ch}(\mathcal{E}), \text{ch}(\mathcal{F}) \rangle_M = \chi(\mathcal{E}, \mathcal{F}).$$

Proof. The first thing to do is to get an interpretation of the Euler pairing. Considering the totality of Ext-groups $\text{Ext}^\bullet(\mathcal{E}, \mathcal{F})$ as a graded vector space, the Euler characteristic is just the graded dimension of $\text{Ext}^\bullet(\mathcal{E}, \mathcal{F})$, which is to say it is the trace of the identity map on $\text{Ext}^\bullet(\mathcal{E}, \mathcal{F})$. Moreover, if $\mathbf{R}\pi_*: \mathbf{D}(X) \rightarrow \mathbf{D}(\text{pt})$ is the derived functor coming from the map $X \rightarrow \text{pt}$, then

$$\text{Ext}^\bullet(\mathcal{E}, \mathcal{F}) \cong \mathbf{H}^\bullet(X, \mathcal{E}^\vee \otimes \mathcal{F}) \cong \mathbf{R}\pi_*(\mathcal{E}^\vee \otimes \mathcal{F})$$

and the latter is just the composition $\mathcal{E}^\vee \circ \mathcal{F}$, where \mathcal{E}^\vee and \mathcal{F} are considered as kernels respectively $X \rightarrow \text{pt}$ and $\text{pt} \rightarrow X$. Thus, using the invariance of the Serre trace under the partial trace map,


$$\begin{aligned} \chi(\mathcal{E}, \mathcal{F}) &= \text{Tr}(\text{Id}_{\mathcal{E}^\vee \circ \mathcal{F}}) = \text{Tr} \left(\begin{array}{ccc} \text{pt} & X & \text{pt} \\ \mathcal{E} \downarrow & & \uparrow \mathcal{F} \end{array} \right) = \text{Tr} \left(\begin{array}{c} \text{pt} \\ \mathcal{E} \downarrow \quad \uparrow \mathcal{F} \\ \text{pt} \end{array} \right) \\ &= \left\langle \begin{array}{c} \text{pt} \\ \mathcal{E} \downarrow \quad \uparrow \mathcal{F} \\ \text{pt} \end{array}, \begin{array}{c} \text{pt} \\ \mathcal{E} \downarrow \quad \uparrow \mathcal{F} \\ \text{pt} \end{array} \right\rangle_M = \langle \text{ch}(\mathcal{E}), \text{ch}(\mathcal{F}) \rangle_M. \quad \square \end{aligned}$$

6.4. Example. To have a noncommutative example at hand, consider the case when G is a finite group acting trivially on a point. The orbifold BG is defined to be the global quotient $[\cdot/G]$ and then the category of coherent sheaves on the orbifold BG is precisely the category of finite dimensional representations of G . One can naturally identify $\text{HH}_0(BG)$ with the space of conjugation invariant functions on G , and the Chern character of a representation ρ is precisely the representation-theoretic character of ρ . See [20] for details.


7. Open-closed TQFTs and the Cardy Condition

We conclude with a discussion of open-closed topological field theories in the B-model and we prove that a condition holds for Hochschild homology which is equivalent to the Cardy Condition in the Calabi–Yau case. Appropriate references for open-closed 2d topological field theories include Moore–Segal [14], Costello [7] and Lauda–Pfeiffer [9].

7.1. Open-closed 2d TQFTs. Consider the open and closed 2-cobordism category 2Cob_{oc} whose objects are oriented, compact one-manifolds — in other words, disjoint unions of circles and intervals — and whose morphisms are (diffeomorphism classes of) cobordisms-with-corners between the source and target one-manifolds. A morphism can be drawn as a vertical cobordism, from the source at the bottom to the target at the top. As well as parts of the boundary being at the top and the bottom, there will be parts of the boundary in between, corresponding to the fact that this is a cobordism with corners. An example is shown below.




Disjoint union makes 2Cob_{oc} into a symmetric monoidal category and an open-closed two-dimensional topological quantum field theory (2d TQFT) is defined to be a symmetric monoidal functor from 2Cob_{oc} to some appropriate symmetric monoidal target category, which we will take to be the category of vector spaces or the category of graded vector spaces. The category 2Cob_{oc} has a simple description in terms of generators and relations which means that there is a reasonably straight forward classification of open-closed 2d TQFTs up to equivalence. This is what we will now describe. The following morphisms generate 2Cob_{oc} as a symmetric monoidal category.




We will come back to the relations below.

To specify an open-closed 2d TQFT up to equivalence on objects it suffices to specify the image \mathbb{C} of the circle and the image \mathbb{O} of the interval. The former is called the space of closed string states and the latter is called the space of open-string states. Using the four *planar* generating morphisms pictured above, together with the relations between them, it transpires that \mathbb{O} , the space of open-string states is precisely a symmetric, but not-necessarily commutative Frobenius algebra. This means that it is a unital algebra with a nondegenerate, symmetric, invariant inner product. It is useful to note here that the inner product is symmetric because the two surfaces pictured below are diffeomorphic, however these surfaces are *not* ambient isotopic — so one cannot be deformed to the other in three-space while the bottom boundary is fixed.



On the other hand, the first four generating morphisms, along with their relations, mean that \mathcal{C} , the space of closed string states, is a *commutative* Frobenius algebra. The last two morphisms mean that there are maps $i_*: \mathcal{C} \rightarrow \mathcal{O}$ and $i^*: \mathcal{O} \rightarrow \mathcal{C}$, and by the relations these are adjoint with respect to the pairings on these spaces. Moreover, i_* is an algebra map, such that its image lies in the centre of \mathcal{O} . The final relation that these must satisfy is the *Cardy Condition*. In terms of the generators pictured above this is the following relation:




Note again that these surfaces are diffeomorphic but not isotopic in three-space. In terms of maps, writing $\mu: \mathcal{O} \otimes \mathcal{O} \rightarrow \mathcal{O}$ and $\delta: \mathcal{O} \rightarrow \mathcal{O} \otimes \mathcal{O}$ for the product and coproduct of the open string state space and writing $\tau: \mathcal{O} \otimes \mathcal{O} \rightarrow \mathcal{O} \otimes \mathcal{O}$ for the symmetry in the target category, the Cardy Condition is the equality of maps from \mathcal{O} to \mathcal{O} :

$$\mu \circ \tau \circ \delta = i_* \circ i^*.$$

We will have reason to use an equivalent condition below.

To summarize, having an open-closed 2d TQFT is equivalent to having the data of a commutative Frobenius algebra \mathcal{C} , a symmetric Frobenius algebra \mathcal{O} , and an algebra map $i_*: \mathcal{C} \rightarrow \mathcal{O}$ with central image, such that the Cardy Condition is satisfied.

7.2. Open-closed 2d TQFTs with D-branes. A more interesting model of string theory is obtained when we specify a set of ‘boundary conditions’ or ‘D-branes’ for the open strings. For a mathematician this just means a set of labels for the boundary points of objects. So fix a set Λ of labels, and consider the category $2\text{Cob}_{\text{oc}}^\Lambda$ of open-closed cobordisms such that the objects are compact, oriented one-manifolds with the boundary points labelled with elements of the set Λ , and morphisms having their internal boundaries labelled compatibly with their boundaries. Here is an example of a morphism from the union of the circle and the interval labelled (B, A) , to the interval labelled (B, A) .



Now a Λ -labelled open-closed TQFT is a symmetric monoidal functor to some appropriate target category which we will again take to be the category of vector spaces or the category of graded vector spaces. Moreover, the category $2\text{Cob}_{\text{oc}}^\Lambda$ is similarly generated by morphisms as listed above, but now they must all be labelled, and the relations are just labelled versions of the previous relations. Thus we can similarly classify Λ -labelled open-closed TQFTs. Once again the image of the circle is a commutative Frobenius

algebra, \mathcal{C} . However, rather than getting a single vector space \mathcal{O} associated to an interval, we get a vector space \mathcal{O}_{BA} associated to each ordered pair (B, A) of elements of Λ ; so we do not get a single Frobenius algebra, but rather something which could be called a ‘Frobenius algebra with many objects’ or a ‘Frobenius algebroid’, but, for the reason explained below, such a thing is commonly known as a Calabi–Yau category. It is a category in the following sense. We take the category whose objects are parametrized by Λ and, for $A, B \in \Lambda$, the morphism set $\text{Hom}(A, B)$ is taken to be \mathcal{O}_{BA} (this is consistent with us reading diagrams from right to left). The composition $\mu_{CBA}: \mathcal{O}_{CB} \otimes \mathcal{O}_{BA} \rightarrow \mathcal{O}_{CA}$ is given by the image of the appropriately labelled version of the morphism pictured.



The Frobenius or Calabi–Yau part of the structure is a — possibly graded — perfect pairing $\mathcal{O}_{AB} \otimes \mathcal{O}_{BA} \rightarrow \mathbf{k}$: the grading degree of this map is called the dimension of the Calabi–Yau category.

So to specify a labelled open-closed 2d TQFT it suffices to specify a commutative Frobenius algebra \mathcal{C} , a Calabi–Yau category \mathcal{O} and an algebra map $i_A: \mathcal{C} \rightarrow \mathcal{O}_{AA}$ with central image, for each object A , such that the labelled version of the Cardy Condition holds.

7.3. The open-closed 2d TQFT from a Calabi–Yau manifold. Associated to a Calabi–Yau manifold X there are two standard 2d TQFTs coming from string theory, imaginatively named the A-model and the B-model: it is the B-model we will be interested in here. In the B-model the boundary conditions are supposed to be “generated” by complex submanifolds of X so the boundary conditions are taken to be complexes of coherent sheaves on X ; the open string category is then supposed to be the derived category of coherent sheaves on X . This is indeed a Calabi–Yau category, which is why such categories are so named: for each \mathcal{E} and \mathcal{F} , the requisite pairing $\text{Hom}_{\mathbf{D}(X)}^{\bullet}(\mathcal{E}, \mathcal{F}) \otimes \text{Hom}_{\mathbf{D}(X)}^{\bullet}(\mathcal{F}, \mathcal{E}) \rightarrow \mathbf{k}[-\dim X]$ comes from the Serre pairing as a Calabi–Yau manifold is precisely a manifold with a trivial canonical bundle.

According to the physics, the closed string state space \mathcal{C} should be

$$\text{Hom}_{\mathbf{D}(X \times X)}^{\bullet}(\mathcal{O}_{\Delta}, \mathcal{O}_{\Delta}),$$

in other words, the Hochschild cohomology algebra $\text{HH}^{\bullet}(X)$. As X is Calabi–Yau, a trivialization of the canonical bundle induces an isomorphism between Hochschild cohomology and Hochschild homology, up to a shift. This means that the closed string space \mathcal{C} has both the cohomological product and the Mukai pairing, and these make \mathcal{C} into a Frobenius algebra.

We need to specify the algebra maps $i_{\mathcal{E}}: \mathcal{C} \rightarrow \mathcal{O}_{\mathcal{E}\mathcal{E}}$. These are maps

$$i_{\mathcal{E}}: \text{Hom}^{\bullet}(\mathcal{O}_{\Delta}, \mathcal{O}_{\Delta}) \rightarrow \text{Hom}^{\bullet}(\mathcal{E}, \mathcal{E})$$

which can be given by interpreting \mathcal{E} as a kernel $\text{pt} \rightarrow X$ and taking $i_{\mathcal{E}}$ to be convolution with the identity on \mathcal{E} . This is given diagrammatically on an element $\varphi \in \text{Hom}^{\bullet}(\mathcal{O}_{\Delta}, \mathcal{O}_{\Delta})$ as follows.

$$\boxed{\varphi} \mapsto \begin{array}{c} X \\ \boxed{\varphi} \\ \mathcal{E} \end{array} \Bigg| \begin{array}{c} \text{pt} \\ \cdot \end{array} .$$

At this point it should be noted that \mathcal{C} is to be thought of as the centre of the category \mathcal{O} . The notion of centre is generalized from algebras to categories by taking the centre of a category to be the natural transformations of the identity functor; however, in a 2-category an appropriate notion of the centre of an object is the set of 2-endomorphisms of the identity morphism on that object. This means that \mathcal{C} is the centre, in this sense, of the category \mathcal{O} in the 2-category $\mathcal{V}\text{ar}$.

The map going the other way, $i^{\mathcal{E}}: \text{Hom}^{\bullet}(\mathcal{E}, \mathcal{E}) \rightarrow \text{Hom}^{\bullet}(\mathcal{O}_{\Delta}, \mathcal{O}_{\Delta})$ is given by taking the trace, namely for $e \in \text{Hom}_{\mathbf{D}(X)}(\mathcal{E}, \mathcal{E})$ the map is given by

$$\begin{array}{c} X \\ \uparrow \text{pt} \\ \boxed{e} \\ \downarrow \mathcal{E} \end{array} \mapsto X \begin{array}{c} \boxed{e} \\ \uparrow \text{pt} \\ \mathcal{E} \end{array} .$$

This definition relies on the fact that X is Calabi–Yau, so that the Serre kernel is, up to a shift, just the identity 1-morphism Id_X .

An argument similar to Proposition 11 shows that $i_{\mathcal{E}}$ and $i^{\mathcal{E}}$ are adjoint. In order to argue that we indeed have an open-closed TQFT it remains to show that the Cardy Condition holds. In fact, we will prove a more general statement, the Baggy Cardy Condition.

7.4. The Baggy Cardy Condition. In the case of a manifold X that is not necessarily Calabi–Yau we don’t have the same coincidence of structure as above: we no longer have a Frobenius algebra $\text{HH}^{\bullet}(X)$; rather we have an algebra $\text{HH}^{\bullet}(X)$ and an inner product space $\text{HH}_{\bullet}(X)$. This means that we can not formulate the Cardy Condition as it stands. We now state a condition which makes sense for an arbitrary, non-Calabi–Yau manifold and which is equivalent to the Cardy Condition in the Calabi–Yau case.


Theorem 15. *Suppose that \mathcal{O} is a Calabi–Yau category and \mathcal{C} is an inner product space, such that for each $A \in \mathcal{O}$ there are adjoint maps $i^A: \mathcal{O}_{AA} \rightarrow \mathcal{C}$ and $i_A: \mathcal{C} \rightarrow \mathcal{O}_{AA}$. Then the Cardy Condition*

$$\mu_{\text{BAB}} \circ \tau \circ \delta_{\text{ABA}} = i_{\text{B}} \circ i^{\text{A}}$$

is equivalent to the following equality holding for all $\mathbf{a} \in \mathcal{O}_{AA}$ and $\mathbf{b} \in \mathcal{O}_{BB}$, where the map ${}_a\mathbf{m}_b: \mathcal{O}_{AB} \rightarrow \mathcal{O}_{AB}$ is the map obtained by precomposing with \mathbf{a} and postcomposing with \mathbf{b} :


$$\left\langle i^{\text{B}} _ , i^{\text{A}} _ \right\rangle_{\mathcal{C}} = \text{Tr} _ \mathbf{m} _ .$$

Proof. The first thing to do is examine the left-hand side of the Cardy Condition. As \mathcal{O} is a Calabi–Yau category there is the following equality of morphisms $\mathcal{O}_{AA} \rightarrow \mathcal{O}_{BB}$.




Note that this does not require any reference to \mathcal{C} , but it does fundamentally require the symmetry of the inner product. This is reflected in the fact that the surfaces underlying the above pictures are diffeomorphic but not ambient isotopic.

This means that the Cardy Condition is equivalent to the following equality.



By the nondegeneracy of the inner product on \mathcal{O}_{BB} this is equivalent to the equality of two maps $\mathcal{O}_{BB} \otimes \mathcal{O}_{AA} \rightarrow k$ which are drawn as follows.



The right-hand side is instantly identifiable as $\langle i^B, i^A \rangle_{\mathcal{C}}$. The left-hand side is identifiable as the trace of the triple composition map $\mathcal{O}_{BB} \otimes \mathcal{O}_{BA} \otimes \mathcal{O}_{AA} \rightarrow \mathcal{O}_{BA}$ which gives the required result. \square

We can now show that the alternative condition given in the above theorem holds for the derived category and Hochschild homology of *any* space: in particular, the Cardy Condition holds for Calabi–Yau spaces.

Theorem 16 (The Baggy Cardy Condition). *Let X be a space, let \mathcal{E} and \mathcal{F} be objects in $\mathbf{D}(X)$ and consider morphisms*

$$e \in \mathrm{Hom}_{\mathbf{D}(X)}(\mathcal{E}, \mathcal{E}) \quad \text{and} \quad f \in \mathrm{Hom}_{\mathbf{D}(X)}(\mathcal{F}, \mathcal{F}).$$

Define the operator

$${}_f m_e: \mathrm{Hom}_{\mathbf{D}(X)}^{\bullet}(\mathcal{E}, \mathcal{F}) \rightarrow \mathrm{Hom}_{\mathbf{D}(X)}^{\bullet}(\mathcal{E}, \mathcal{F})$$

to be postcomposition by f and precomposition by e . Then we have

$$\mathrm{Tr} {}_f m_e = \left\langle \iota^{\mathcal{E}}(e), \iota^{\mathcal{F}}(f) \right\rangle_M,$$

where $\iota^{\mathcal{E}}, \iota^{\mathcal{F}}$ are the maps defined in Section 6.2, and Tr denotes the (super) trace.

Proof. The proof is very similar to the proof of the Semi-Hirzebruch–Riemann–Roch Theorem (Theorem 14). The first thing to observe is that $\mathrm{Hom}(\mathcal{E}, \mathcal{F}) \cong \mathcal{E}^\vee \circ \mathcal{F}$ and that ${}_f m_e$ is just $e^\vee \circ f$. However, we have $\tau_R(e) = \Sigma_{\mathrm{pt}} \circ e^\vee$ and Σ_{pt} is trivial so $e^\vee = \tau_R(e)$. Putting this together with the invariance of the Serre trace under the partial trace we get the following sequence, and hence the required result.

$$\begin{aligned}
\mathrm{Tr} \, {}_f m_e &= \mathrm{Tr} \left(\begin{array}{c} \begin{array}{ccc} & X & \\ \downarrow & & \uparrow \\ \mathrm{pt} & \boxed{e^\vee} & \boxed{f} & \mathrm{pt} \\ \downarrow & & \downarrow \\ \mathcal{E} & & \mathcal{F} \end{array} \\ \downarrow \\ \begin{array}{ccc} \text{circle } \mathcal{E} & & \text{circle } \mathcal{F} \\ \downarrow & & \downarrow \\ \mathcal{E} & & \mathcal{F} \end{array} \end{array} \right) = \mathrm{Tr} \left(\begin{array}{c} \text{circle } \mathcal{E} \text{ with } \boxed{e^\vee} \text{ and } \text{circle } \mathcal{F} \text{ with } \boxed{f} \\ \downarrow \\ \mathcal{E} \text{ and } \mathcal{F} \end{array} \right) \\
&= \mathrm{Tr} \left(\begin{array}{c} \text{circle } \mathcal{E} \text{ with } \boxed{e} \text{ and } \text{circle } \mathcal{F} \text{ with } \boxed{f} \\ \downarrow \\ \mathcal{E} \text{ and } \mathcal{F} \end{array} \right) = \mathrm{Tr} \left(\begin{array}{c} \text{circle } \mathcal{E} \text{ with } \boxed{e} \text{ and } \text{circle } \mathcal{F} \text{ with } \boxed{f} \\ \downarrow \\ \mathcal{E} \text{ and } \mathcal{F} \end{array} \right) \\
&= \left\langle \begin{array}{c} \text{circle } \mathcal{E} \text{ with } \boxed{e} \\ \downarrow \\ \mathcal{E} \end{array}, \begin{array}{c} \text{circle } \mathcal{F} \text{ with } \boxed{f} \\ \downarrow \\ \mathcal{F} \end{array} \right\rangle_M = \left\langle \iota^{\mathcal{E}}(e), \iota^{\mathcal{F}}(f) \right\rangle_M. \quad \square
\end{aligned}$$

Observe that the Semi-Hirzebruch–Riemann–Roch Theorem is a direct consequence of the Baggly Cardy Condition, with $e = \mathrm{Id}_{\mathcal{E}}$, $f = \mathrm{Id}_{\mathcal{F}}$.

Appendix A. Duality and partial trace

In this appendix we show that given a kernel $\Phi: X \rightarrow Y$ and its dual kernel $\Phi^\vee: Y \rightarrow X$ there are canonical 2-morphisms

$$\Sigma_X^{-1} \rightarrow \Phi^\vee \circ \Phi \quad \text{and} \quad \Phi \circ \Sigma_X \circ \Phi^\vee \rightarrow \mathrm{Id}_Y$$

giving rise to a variety of natural adjunctions satisfying a number of compatibility relations.

The notion of duality in $\mathcal{V}\mathrm{ar}$ is seen to be a middle-ground between the operations τ_L and τ_R : it is an involution, unlike τ_L and τ_R , but it does not respect composition, which τ_L and τ_R do.

A.1. Polite duality. Recall from Section 2.2 that for every space X there is the Serre kernel $\Sigma_X: X \rightarrow X$ such that for spaces X and Y the functor

$$\Sigma_Y \circ - \circ \Sigma_X: \mathbf{Hom}_{\mathcal{V}\mathrm{ar}}(X, Y) \rightarrow \mathbf{Hom}_{\mathcal{V}\mathrm{ar}}(X, Y)$$

is a Serre functor for the category $\mathbf{Hom}_{\mathcal{V}\mathrm{ar}}(X, Y)$.

Definition. If $\Phi: X \rightarrow Y$ and $\Phi^\dagger: Y \rightarrow X$ are kernels then a *polite duality* between them, denoted $\Phi \overset{\vee}{\longleftrightarrow} \Phi^\dagger$, consists of adjunctions as follows (numbered as shown)

$$\begin{aligned}
\Phi^\dagger \circ \Sigma_Y \dashv_1 \Phi \dashv_2 \Sigma_X \circ \Phi^\dagger, \\
\Phi \circ \Sigma_X \dashv_3 \Phi^\dagger \dashv_4 \Sigma_Y \circ \Phi,
\end{aligned}$$

such that the following compatibility relations hold.

The symbol \circ is pronounced “mepsilon” and the names \circ and γ come from the shape of the morphisms.

Just as above, using the relation (1+4) from the units η_1 and η_4 , and the counits ϵ_1 and ϵ_4 we obtain the common morphisms denoted as follows:

$$\gamma_{\Phi^\vee} : \begin{array}{c} \text{---} \\ \curvearrowright \\ \text{---} \\ \vdots \end{array} \quad \circ_{\Phi^\vee} : \begin{array}{c} \text{---} \\ \curvearrowleft \\ \text{---} \\ \vdots \end{array}.$$

So the four units and four counits are obtained from these two \circ -shaped and two γ -shaped morphisms.

Relations (1+2) and (3+4) in the definition of polite duality are essentially equivalent to the invariance of the Serre trace under a partial trace in the following sense. Given a polite duality $\Phi \overset{\vee}{\longleftrightarrow} \Phi^\dagger$ and a morphism $\alpha \in \text{Hom}(\Phi \circ \Theta, \Sigma \circ \Phi \circ \Psi)$ we can define the left partial trace in $\text{Hom}(\Theta, \Sigma \circ \Psi)$ with respect to Φ^\dagger as drawn below:

$$\begin{array}{c} \Psi \\ \uparrow \\ \text{---} \\ \alpha \\ \text{---} \\ \Theta \end{array}.$$

Similarly we can define a right partial trace when the Φ is on the right rather than on the left. So the following diagram is the right partial trace with respect to Φ and Φ^\dagger of a morphism $\alpha' \in \text{Hom}(\Theta' \circ \Phi, \Psi' \circ \Phi \circ \Sigma)$:

$$\begin{array}{c} \Psi' \\ \uparrow \\ \text{---} \\ \alpha' \\ \text{---} \\ \Theta' \end{array}.$$

Theorem 17. *If Φ and Φ^\dagger form a politely dual pair of kernels, then the Serre trace is invariant under partial trace with respect to Φ :*

$$\text{Tr} \left(\begin{array}{c} \Psi \\ \uparrow \\ \text{---} \\ \alpha \\ \text{---} \\ \Psi \end{array} \right) = \text{Tr} \left(\begin{array}{c} \Psi \\ \uparrow \\ \text{---} \\ \alpha \\ \text{---} \\ \Psi \end{array} \right); \quad \text{Tr} \left(\begin{array}{c} \Psi' \\ \uparrow \\ \text{---} \\ \alpha' \\ \text{---} \\ \Psi' \end{array} \right) = \text{Tr} \left(\begin{array}{c} \Psi' \\ \uparrow \\ \text{---} \\ \alpha' \\ \text{---} \\ \Psi' \end{array} \right).$$

Proof. We will just prove the left partial trace case. Relation (1+2) says that for $\beta \in \text{Hom}(\Psi, \Phi \circ \Theta)$ and $\gamma \in (\Theta, \Sigma \circ \Phi^\dagger \circ \Sigma \circ \Psi \circ \Sigma)$ we have the equality

$$\text{Tr} \left(\begin{array}{c} \Psi \\ \uparrow \\ \text{---} \\ \gamma \\ \text{---} \\ \Theta \\ \uparrow \\ \text{---} \\ \beta \\ \text{---} \\ \Psi \end{array} \right) = \text{Tr} \left(\begin{array}{c} \Psi \\ \uparrow \\ \text{---} \\ \gamma \\ \text{---} \\ \Theta \\ \uparrow \\ \text{---} \\ \beta \\ \text{---} \\ \Psi \end{array} \right).$$

Proof. For kernels $\Theta \in \mathbf{D}(Z \times X)$ and $\Psi \in \mathbf{D}(Z \times Y)$ consider the sequence of isomorphisms

$$\begin{aligned} & \mathrm{Hom}_{\mathbf{D}(Z \times Y)}(\Psi, \Phi \circ \Theta) \\ & \cong \mathrm{Hom}_{\mathbf{D}(Z \times Y)}(\Psi, \pi_{ZY,*}(\pi_{ZX}^* \Theta \otimes \pi_{XY}^* \Phi)) \\ & \cong \mathrm{Hom}_{\mathbf{D}(Z \times X)}(\pi_{ZX,!}(\pi_{ZY}^* \Psi \otimes \pi_{YX}^* \Phi^\vee), \Theta) \\ & = \mathrm{Hom}_{\mathbf{D}(Z \times X)}(\pi_{ZX,*}(\pi_{ZY}^* \Psi \otimes \pi_{YX}^*(\Phi^\vee \otimes \omega_Y[\dim Y])), \Theta) \\ & = \mathrm{Hom}_{\mathbf{D}(Z \times Y)}((\Phi^\vee \circ \Sigma_Y) \circ \Psi, \Theta). \end{aligned}$$

Taking $Z = Y$, $\Theta = \Phi^\vee \circ \Sigma_Y$ and $\Psi = \mathrm{Id}_Y$ yields a morphism *of kernels*

$$\mathrm{Id}_Y \rightarrow \Phi \circ \Phi^\vee \circ \Sigma_Y;$$

in a similar fashion we obtain a morphism

$$\Phi^\vee \circ \Sigma_Y \circ \Phi \rightarrow \mathrm{Id}_X.$$

These two morphisms satisfy the identities needed to make $\Phi^\vee \circ \Sigma_Y$ the left adjoint of Φ . Proposition 19 gives the result. \square

A.2. Reflexively polite kernels. We still need to address one more compatibility between the dualities constructed above. Given a kernel Φ , the previous proposition yields a polite duality

$$\Phi \overset{\vee}{\longleftarrow} \Phi^\vee.$$

Given any polite duality $\Phi \overset{\vee}{\longleftarrow} \Phi^\dagger$, we get, symmetrically, a polite duality $\Phi^\dagger \overset{\vee}{\longleftarrow} \Phi$ by switching the adjunctions 1 and 3 and the adjunctions 2 and 4. Thus there is a natural polite duality

$$\Phi^\vee \overset{\vee}{\longleftarrow} \Phi.$$

On the other hand, applying Proposition 20 to the kernel Φ^\vee we get a polite duality $\Phi^\vee \overset{\vee}{\longleftarrow} \Phi^{\vee\vee}$ and then, using the canonical identification $\Phi^{\vee\vee} \cong \Phi$, we get *another* polite duality

$$\Phi^\vee \overset{\vee}{\longleftarrow} \Phi.$$

The fundamental question is whether these two polite dualities are the same.

Definition. Let $\Phi: X \rightarrow Y$ be a kernel. We shall say that Φ is *reflexively polite* if the two dualities above are equal.

Immediately we get the following result.

Proposition 21. *If a kernel Φ is reflexively polite then so is Φ^\vee .*

Proposition 22. *If $\Phi: X \rightarrow Y$ and $\Psi: Y \rightarrow Z$ are reflexively polite kernels then so is $\Psi \circ \Phi$.*

Proof. This follows at once from the fact that the adjunctions defined in Proposition 18 are canonical. \square

Proposition 23. *Suppose that $\Phi: X \rightarrow Y$ and $\Psi: X' \rightarrow Y'$ are reflexively polite kernels then so is $\Phi \boxtimes \Psi: X \times X' \rightarrow Y \times Y'$, the kernel defined by*

$$\Phi \boxtimes \Psi := \pi_{XY}^* \Phi \otimes \pi_{X'Y'}^* \Psi.$$

Proof. This is obvious once one realises that all operations decompose with respect to the box product operation, and the canonical bundle of a product of spaces is the box product of the canonical bundles of the factors. \square

Proposition 24. *Let $\Delta: X \rightarrow X \times X$ denote the diagonal map, let \mathcal{E} be an object in $\mathbf{D}(X)$, then the kernel $\Delta_* \mathcal{E}: X \rightarrow X$ is reflexively polite.*

Proof. Serre duality gives a natural identification of kernels $X \rightarrow X$

$$(\Delta_* \mathcal{E})^\vee \cong \Delta_*(\mathcal{E}^\vee \otimes \omega_X^{-1}[-\dim X]).$$

The result now follows from Proposition 2 and the fact that the functor given by tensoring with an object of the derived category is left and right adjoint to the one obtained by tensoring with the dual object. \square

Proposition 25. *Let X be a space, then the kernel \mathcal{O}_X considered as a kernel $X \rightarrow \mathbf{pt}$ is reflexively polite, and so also is $\mathcal{O}_X^\vee: \mathbf{pt} \rightarrow X$.*

Proof. First, observe that checking reflexive duality amounts to checking the equality of two quadruples of adjunctions. In each quadruple, any one of the adjunctions determines the remaining three, and thus in order to check the equality it suffices to check that one of the compatibilities (1+2), (1+3), etc. holds with one adjunction chosen from one duality, and the other one from the second duality.

Furthermore, adjunctions are completely determined by their respective units and counits, and these are determined by considered functors *between derived categories* of the form

$$\Phi \circ -: \mathbf{D}(Z \times X) \rightarrow \mathbf{Hom}_{\mathcal{Y}_{\text{ar}}}(Z, X) \rightarrow \mathbf{Hom}_{\mathcal{Y}_{\text{ar}}}(Z, \mathbf{pt}) := \mathbf{D}(Z),$$

and analogues for $\Phi^\vee \circ -$, for various choices of the space Z . Thus, if we argue that the desired equality holds for the adjunctions between these induced functors (for arbitrary choice of Z), we will have argued that Φ is reflexively polite.

For ease of notation, write Φ for the kernel $\mathcal{O}_X: X \rightarrow \mathbf{pt}$. For a given space Z , let $\pi_Z: X \times Z \rightarrow Z$ denote the projection. The functors $\Phi \circ -$ and $\Phi^\vee \circ -$ are then naturally identified with $\pi_{Z,*}$ and π_Z^* , respectively. Adjunctions 1 and 2 from the polite duality $\Phi \overset{\vee}{\leftarrow} \Phi^\vee$ correspond to the classical adjunctions

$$\pi_Z^* \dashv \pi_{Z,*} \dashv \pi_Z^!$$

while the same ones from the polite duality $\Phi^\vee \overset{\vee}{\leftarrow} \Phi^{\vee\vee}$ correspond to

$$\pi_{Z,!} \dashv \pi_Z^* \dashv \pi_{Z,*}.$$

Indeed, the standard way (see [6, Theorem 4.6]) to define the adjunctions $\pi_{Z,*} \dashv \pi_Z^!$ and $\pi_{Z,!} \dashv \pi_Z^*$ is to require them to satisfy the analogue of condition (1+2) from the definition of a polite duality.

The condition of polite duality can now be stated as the statement that the composite isomorphism

$$\begin{aligned} \mathrm{Hom}_{X \times Z}(\Theta, \pi_Z^* \Psi) &\cong \mathrm{Hom}_Z(\pi_{Z,!} \Theta, \Psi) = \mathrm{Hom}_Z(\pi_{Z,*} \mathbf{S}(\Theta), \Psi) \\ &\cong \mathrm{Hom}_{X \times Z}(\mathbf{S}(\Theta), \pi_Z^! \Psi) = \mathrm{Hom}_{X \times Z}(\mathbf{S}(\Theta), \mathbf{S}(\pi_Z^* \Psi)) \end{aligned}$$

is the one induced by the functor $\mathbf{S}(-) = - \otimes \pi_X^* \omega_X[\dim X]$, where π_X denotes the projection from $X \times Z$ to X . This fact corresponds to the fact that the diagram marked (!) below commutes

$$\begin{array}{ccccccc} \mathrm{Hom}(\Theta, \pi^* \Psi) & \xrightarrow{\sim} & \mathrm{Hom}(\mathbf{S}^{-1} \pi_* \mathbf{S} \Theta, \Psi) & \xrightarrow{\sim} & \mathrm{Hom}(\pi_* \mathbf{S} \Theta, \mathbf{S} \Psi) & \xrightarrow{\sim} & \mathrm{Hom}(\mathbf{S} \Theta, \mathbf{S} \pi_* \Psi) \\ \sim \downarrow & \text{Serre} & \sim \downarrow & (!) & \sim \downarrow & \text{Serre} & \sim \downarrow \\ \mathrm{Hom}(\pi^* \Psi, \mathbf{S} \Theta)^\vee & \xrightarrow{\sim} & \mathrm{Hom}(\Psi, \pi_* \mathbf{S} \Theta)^\vee & = & \mathrm{Hom}(\Psi, \pi_* \mathbf{S} \Theta)^\vee & \xrightarrow{\sim} & \mathrm{Hom}(\pi^* \Psi, \Theta)^\vee. \end{array}$$

□

Proposition 26. *Let $\pi_X: X \times Y \rightarrow X$ be the projection, and (abusing notation) denote by $\pi_{X,*}: X \times Y \rightarrow X$ and $\pi_X^*: X \rightarrow X \times Y$ the kernels represented in $\mathbf{D}(X \times X \times Y)$ by the structure sheaf of the graph of π_X . Then $\pi_{X,*}$ and π_X^* are reflexively polite.*

Proof. Both kernels are of the form $\mathcal{O}_{\Delta_X} \boxtimes \mathcal{O}_Y$, and the result then follows from Propositions 25, 24, and 23. □

Theorem 27. *Every kernel is reflexively polite.*

Proof. With notation as in Proposition 26, any kernel $\Phi: X \rightarrow Y$ decomposes as

$$\Phi = \pi_{Y,*} \circ (- \otimes \Phi) \circ \pi_X^*.$$

Since the claim of the theorem holds for each individual kernel in the decomposition (Propositions 26 and 24), the result follows from Proposition 18. □

References

- [1] BONDAL, A. I., KAPRANOV, M. M. Enhanced triangulated categories. *Mat. Sb.* **181** (1990), no. 5, 669–683; translation in *Math. USSR-Sb.* **70** (1991), no. 1, 93–107. MR1055981 (91g:18010), Zbl 0729.18008.
- [2] BONDAL, A. I., KAPRANOV, M. M. Representable functors, Serre functors, and reconstructions. *Izv. Akad. Nauk SSSR Ser. Mat.* **53** (1989), no. 6, 1183–1205, 1337; translation in *Math. USSR-Izv.* **35** (1990), no. 3, 519–541. MR1039961 (91b:14013), Zbl 0703.14011.
- [3] CĂLDĂRARU, ANDREI. Derived categories of twisted sheaves on Calabi–Yau manifolds. Ph.D. thesis, Cornell University, 2000. <http://www.math.wisc.edu/~andreic/publications/ThesisSingleSpaced.pdf>.
- [4] CĂLDĂRARU, ANDREI. The Mukai pairing, I: the Hochschild structure. Preprint arXiv:math.AG/0308079.

- [5] CĂLDĂRARU, ANDREI. The Mukai pairing. II. The Hochschild–Kostant–Rosenberg isomorphism. *Adv. Math.* **194** (2005), no. 1, 34–66. MR2141853 (2006a:14029), Zbl 1098.14011. arXiv:math.AG/0308080.
- [6] CĂLDĂRARU, ANDREI. Derived categories of sheaves: a skimming. *Snowbird lectures in algebraic geometry*, 43–75. *Contemp. Math.*, 388. *Amer. Math. Soc., Providence, RI*, 2005. MR2182889 (2006h:14022).
- [7] COSTELLO, KEVIN. Topological conformal field theories and Calabi–Yau categories. *Adv. Math.* **210** (2007), no. 1, 165–214. MR2298823 (2008f:14071), Zbl 1171.14038. arXiv:math.AG/0412149.
- [8] GELFAND, SERGEI I.; MANIN, YURI I. Methods of homological algebra. Second edition. Springer Monographs in Mathematics, *Springer-Verlag, Berlin*, 2003. xx+372 pp. ISBN: 3-540-43583-2. MR1950475 (2003m:18001), Zbl 1006.18001.
- [9] LAUDA, AARON D.; PFEIFFER, HENDRYK. Open-closed strings: Two-dimensional extended TQFTs and Frobenius algebras. *Topology Appl.* **155** (2008), no. 7, 623–666. MR2395583 (2009g:57049), Zbl 1158.57038. arXiv:math/0510664.
- [10] MARKARIAN, NIKITA. The Atiyah class, Hochschild cohomology and the Riemann–Roch theorem. arXiv:math.AG/0610553 (earlier version available as Max-Planck preprint MPIM2001-52).
- [11] MAY, J. PETER. The additivity of traces in triangulated categories. *Adv. Math.* **163** (2001), no. 1, 34–73. MR1867203 (2002k:18019), Zbl 1007.18012.
- [12] MAY, J. P.; SIGURDSSON, J. Parametrized homotopy theory. *Mathematical Surveys and Monographs*, 132. *American Mathematical Society, Providence, RI*, 2006. x+441 pp. ISBN: 978-0-8218-3922-5; 0-8218-3922-5. MR2271789 (2007k:55012), Zbl 1119.55001.
- [13] MOORE, GREGORY W. Lectures on branes, K-theory and RR charges — two variations on the theme of 2d TFT. *Clay Mathematical Institute Lectures*. Available at <http://www.physics.rutgers.edu/~gmoore/clay.html>.
- [14] MOORE, GREGORY W.; SEGAL, GRAEME. D-branes and K-theory in 2D topological field theory. arXiv:hep-th/0609042.
- [15] MUKAI, SHIGERU. Moduli of vector bundles on K3 surfaces, and symplectic manifolds. *Sugaku* **39** (1987) 216–235; translated as *Sugaku Expositions* **1** (1988) 139–174. MR0922020 (89h:32057), Zbl 0651.14003, Zbl 0685.14021.
- [16] ORLOV, D. O. Triangulated categories of singularities and D-branes in Landau–Ginzburg models. *Tr. Mat. Inst. Steklova* **246** (2004), *Algebr. Geom. Metody, Svyazi i Prilozh.*, 240–262; translation in *Proc. Steklov Inst. Math.* **2004**, no. 3 (246), 227–248. MR2101296 (2006i:81173), Zbl 1101.81093.
- [17] REITEN, I.; VAN DEN BERGH, M. Noetherian hereditary abelian categories satisfying Serre duality. *J. Amer. Math. Soc.* **15** (2002), no. 2, 295–366. MR1887637 (2003a:18011), Zbl 0991.18009.
- [18] TOËN, BERTRAND. The homotopy theory of dg-categories and derived Morita theory. *Invent. Math.* **167** (2007), no. 3, 615–667. MR2276263 (2008a:18006), Zbl 1118.18010. arXiv:math.AG/0408337.
- [19] WEIBEL, CHARLES. Cyclic homology for schemes. *Proc. Amer. Math. Soc.* **124** (1996) 1655–1662. MR1277141 (96h:19003), Zbl 0855.19002.
- [20] WILLERTON, SIMON. In preparation.

MATHEMATICS DEPARTMENT, UNIVERSITY OF WISCONSIN–MADISON, 480 LINCOLN DRIVE,
MADISON, WI 53706–1388, USA
andreic@math.wisc.edu

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF SHEFFIELD, HICKS BUILDING,
HOUNSFIELD ROAD, SHEFFIELD, S3 7RH, UK
S.Willerton@sheffield.ac.uk

This paper is available via <http://nyjm.albany.edu/j/2010/16-6.html>.