

# 1 **The northern sector of the Last British Ice Sheet: maximum extent** 2 **and demise**

3  
4 Tom Bradwell<sup>1</sup>, Martyn Stoker<sup>1</sup>, Nick Golledge<sup>1</sup>, Christian Wilson<sup>1</sup>, Jon Merritt<sup>1</sup>,  
5 Dave Long<sup>1</sup>, Jez Everest<sup>1</sup>, Ole B. Hestvik<sup>2</sup>, Alan Stevenson<sup>1</sup>, Alun Hubbard<sup>3</sup>, Andrew  
6 Finlayson<sup>1</sup>, Hannah Mathers<sup>1,4</sup>

7  
8 <sup>1</sup> *British Geological Survey, Murchison House, West Mains Road, Edinburgh, EH9 3LA, UK*

9 <sup>2</sup> *Olex AS, Pirsenteret, N-7462 Trondheim, Norway*

10 <sup>3</sup> *Institute of Geography and Earth Sciences, University of Wales, Aberystwyth, Ceredigion,*  
11 *SY23 3DB, UK*

12 <sup>4</sup> *Department of Geographical and Earth Sciences, East Quadrangle, University Avenue,*  
13 *University of Glasgow, Glasgow, G12 8QQ, UK*

## 14 15 **Abstract**

16 Strongly divided opinion has led to competing, apparently contradictory, views on the  
17 timing, extent, flow configuration and decay mechanism of the last British Ice Sheet.  
18 We review the existing literature and reconcile some of these differences using  
19 remarkable new seabed imagery. This bathymetric data provides unprecedented  
20 empirical evidence of confluence and subsequent separation of the last British and  
21 Fennoscandian Ice Sheets. Critically, it also allows a viable pattern of ice-sheet  
22 disintegration to be proposed for the first time. Covering the continental shelf around  
23 the northern United Kingdom, extensive echosounder data reveals striking  
24 geomorphic evidence – in the form of tunnel valleys and moraines – relating to the  
25 former British and Fennoscandian Ice Sheets. The pattern of tunnel valleys in the  
26 northern North Sea Basin and the presence of large moraines on the West Shetland  
27 Shelf, coupled with stratigraphic evidence from the Witch Ground Basin, all suggest  
28 that at its maximum extent a grounded ice sheet flowed from SE to NW across the  
29 northern North Sea Basin, terminating at the continental shelf edge. The zone of  
30 confluence between the British and much larger Fennoscandian Ice Sheets was  
31 probably across the northern Orkney Islands, with fast-flowing ice in the Fair Isle  
32 Channel focusing sediment delivery to the Rona and Foula Wedges. This period of  
33 maximum confluent glaciation (c. 30-25 ka BP) was followed by a remarkable period  
34 of large-scale ice-sheet re-organisation. We present evidence suggesting that as sea-  
35 level rose, a large marine embayment opened in the northern North Sea Basin, as far  
36 south as the Witch Ground Basin, forcing the two ice sheets to decouple rapidly along  
37 a north-south axis east of Shetland. As a result, both ice-sheets rapidly adjusted to  
38 new quasi-stable margin positions forming a second distinct set of moraines (c. 24-18  
39 ka BP). The lobate overprinted morphology of these moraines on the mid-shelf west  
40 of Orkney and Shetland indicates that the re-organisation of the British Ice Sheet was  
41 extremely dynamic – probably dominated by a series of internally forced readvances.  
42 Critically, much of the ice in the low-lying North Sea Basin may have disintegrated  
43 catastrophically as decoupling progressed in response to rising sea levels. Final-stage  
44 deglaciation was marked by near-shore ice streaming and increasing topographic  
45 control on ice-flow direction. Punctuated retreat of the British Ice Sheet continued  
46 until c. 16 ka BP when, following the North Atlantic iceberg-discharge event  
47 (Heinrich-1), ice was situated at the present-day coastline in NW Scotland.

48 **Key words:** UK continental shelf, North Sea, ice stream, sea-level rise, deglaciation

49

1 **Introduction**

2 At present, views vary widely regarding the extent, thickness and geometry of the last  
3 British Ice Sheet (BIS), and its interaction with the neighbouring Fennoscandian Ice  
4 Sheet (FIS). Nowhere is this problem better highlighted than in the North Sea Basin.  
5 Whilst some authors have claimed that, at its maximum, the last BIS coalesced with  
6 the Fennoscandian Ice Sheet (FIS) in the North Sea Basin (e.g. Sejrup et al., 1994;  
7 2005; Carr et al., 2006); others have contended that it terminated only a short distance  
8 offshore (e.g. Sutherland, 1984; Bowen et al., 2002). These differences in ice-sheet  
9 extent are contradictory and appear impossible to reconcile (cf. Bowen et al., 2002;  
10 Hall et al., 2003). Resolving this impasse has important implications, not only for  
11 underpinning past changes in the geometry and dynamics of the last BIS, but also for  
12 environmental change on a global scale. Questions regarding Northern Hemisphere  
13 ice-sheet volume, concomitant sea-level changes and the potential impact of  
14 meltwater and calving flux on the North Atlantic thermohaline circulation are central  
15 to our understanding of the coupled ocean-atmosphere system.

16

17 The glacial history of the continental shelf east of the UK, predominantly comprising  
18 the North Sea Basin, remains relatively poorly understood (Figure 1). Previous  
19 reconstructions for the northern North Sea area (north of 57°N) depict both ice-free  
20 (e.g. Sutherland, 1984; Boulton et al., 1985; Bowen et al., 2002) and ice-covered  
21 scenarios (e.g. Sissons, 1967; Boulton et al., 1977; Sejrup et al., 1994). Only recently  
22 has opinion begun to converge on the idea of a glaciated North Sea Basin in Late  
23 Weichselian times (Marine Isotope Stage 2) (Sejrup et al., 2005; Carr et al., 2006).  
24 This view has been strongly reinforced by the recent identification of mega-scale  
25 glacial lineations (MSGs) in the central part of the northern North Sea (Graham et  
26 al., 2007). These sub-surface lineations, imaged on 3D seismic profiles in the Witch  
27 Ground Basin, represent the signature of fast-flow within a grounded ice sheet. Their  
28 presence and orientation are compatible with a major ice stream draining the northern  
29 North Sea fed by the coalescent British and Fennoscandian ice sheets. Stratigraphic  
30 evidence from British Geological Survey (BGS) boreholes in the Witch Ground Basin  
31 supports the interpretation of a palaeo-ice stream flowing over deformable sediment,  
32 and crucially constrains the timing of ice flow to between <42 ka <sup>14</sup>C BP and ~22 ka  
33 <sup>14</sup>C BP (i.e. <47-26 cal ka BP). On this basis, Graham et al. (2007) place the most  
34 likely period of ice streaming in the Late Weichselian (MIS 2). These recent

1 important findings, and those of Carr et al. (2006), echo the views first proposed by  
2 Sejrup et al. (1994) and developed in subsequent papers (Sejrup et al., 2003; Sejrup et  
3 al., 2005). Although the exact timing of the glaciation needs refining, unequivocal  
4 evidence that a grounded Weichselian ice sheet occupied the North Sea Basin, to at  
5 least 58.5°N, has now been demonstrated (Graham, 2007; Graham et al., 2007).

6  
7 Previous work on the palaeoglaciology of the western margins of the FIS has  
8 highlighted the role of the Norwegian Channel in concentrating flow along the west  
9 coast of Norway. The Norwegian Channel palaeo-ice stream, first identified in the  
10 1990s, drained much of the western sector of the FIS and operated during the last  
11 glacial cycle (MIS 2) (Sejrup et al., 1994, 1998; Rise et al., 2004; Ottesen et al.,  
12 2005). However, geophysical data from the seafloor ~50 km west of the Norwegian  
13 Channel has also revealed the signature of streaming grounded ice flowing NW on the  
14 Egersundbanken (Stalsberg et al., 2003). These glacial lineations indicate that at some  
15 periods during the Late Weichselian the FIS was not confined by the Norwegian  
16 Channel and may have flowed into the North Sea Basin. Graham et al. (2007) inferred  
17 ice flow from the SW sector of the FIS across the North Sea Basin to explain the  
18 presence of MSGs in the northern North Sea. This evidence is supported by sensitive  
19 computer modelling experiments of the last FIS which simulate the ice sheet  
20 periodically overflowing the Norwegian Channel during periods of maximum  
21 glaciation (Boulton and Hagdorn, 2006).

22  
23 In this paper we present a new model for the deglaciation of the continental shelf  
24 around the northern United Kingdom based on a synthesis of new offshore imagery.  
25 This model allows us to explain and subsequently reconcile much of the previously  
26 published evidence relating to the former extent of the last BIS and FIS. We first  
27 review the existing literature pertaining to the extent and demise of the northern sector  
28 of the last BIS. We then present virtually complete coverage of echosounder data  
29 from the continental shelf around the northern United Kingdom, held within the Olex  
30 database ([www.olex.no](http://www.olex.no)). This phenomenal dataset allows an unprecedented view of  
31 the seafloor geomorphology, enabling some of the controversies outlined in our  
32 review to be resolved. Focusing principally on geomorphic evidence from the  
33 Hebrides Shelf, West Shetland Shelf and the North Sea Basin, we describe and  
34 interpret the morphology and distribution of many previously unreported glacial

1 features. Based on these findings we propose a new, glaciologically plausible,  
2 reconstruction for the northern half of the last BIS, showing the extent and flow  
3 configuration of the ice sheet at Last Glacial Maximum and at two prominent stages  
4 during deglaciation. Finally, we discuss the causes of ice-sheet break up; the likely  
5 feedbacks within the ocean-cryosphere system; and the wider implications of our  
6 findings.

7

8 [FIGURE 1 HERE]

9

10 **Terminology**

11 This paper refers to the last ice-sheet glaciation to have affected the British Isles.  
12 Many onshore workers refer to this as the Late Devensian glaciation (UK) (e.g.  
13 Shotton, 1977; Sissons, 1980; Merritt et al., 1995; Ballantyne et al., 1998; Carr et al.,  
14 2006) or Late Weichselian glaciation (Norway) (e.g. Sejrup et al., 1994, 2003; 2005;  
15 Stalsberg et al., 2003; Ottesen et al., 2005); whilst most marine geologists use Marine  
16 Isotopic Stages to classify glacial events (i.e. MIS 2) (e.g. Stoker et al., 1993; Gatliff  
17 et al., 1994). Unfortunately, confusion still surrounds the use of the term ‘Last Glacial  
18 Maximum (LGM)’ when referring to the British Ice Sheet. The term LGM has been  
19 used by various workers to refer to two temporally distinct periods: 18-24 ka BP<sup>1</sup>  
20 (Boulton et al., 1991; Mix et al., 2001; Bowen et al., 2002) and 25-35 ka BP (Sejrup  
21 et al., 2005; Carr et al., 2006). This has led to debate over the exact timing of the Last  
22 Glacial Maximum in Britain, Ireland and western Scandinavia.

23

24 In this paper, to avoid confusion, we define the term ‘LGM’ with reference to the last  
25 global ice-sheet maximum, determined from sea-level records to have been between  
26 30-22 ka BP – culminating c. 26 ka BP (Peltier and Fairbanks, 2006). With reference  
27 to the British Ice Sheet, the LGM encompasses the period of extensive shelf-wide  
28 glaciation identified by previous workers (e.g. Stoker et al., 1993; Sejrup et al., 1994,  
29 2000, 2005; Knutz et al., 2001; Kroon et al., 2002; Peck et al., 2007). Although  
30 loosely defined chronologically, this period of maximum glaciation probably occurred  
31 between ~30-25 ka BP (i.e. during the earliest part of the Late Devensian (MIS 2)  
32 (Figure 2).

33 [FIGURE 2 HERE]

34 Footnote 1: unless otherwise stated, dates are expressed in calendar (sidereal) years (i.e. 18 ka BP = 18,000 years before present).

1 **Record of glaciation**

2 In this section we review the key morphological and chronological evidence for ice-  
3 sheet glaciation, both onshore and offshore northern Britain – providing a context in  
4 which to place our new observations. Dates are quoted in both radiocarbon years ( $^{14}\text{C}$   
5 BP) and calibrated to calendar years before present (cal BP) where appropriate.  
6 Calibrations were performed using CALIB 5.0 (Stuiver et al., 2005) and Fairbanks et  
7 al. (2005).

8

9 *Onset of glaciation*

10 The record of ice rafting offshore northern Britain dates from the Late Pliocene (~2.5  
11 Ma), although expansive glaciation of the continental shelf probably did not occur  
12 until the early Mid-Pleistocene, about 0.44 Ma (MIS 12) (Shackleton et al., 1984;  
13 Cameron et al., 1987; Stoker et al., 1994). In the central and northern North Sea  
14 Basin, and on the Hebrides and West Shetland shelves, seismic reflection profiles  
15 reveal stacked glacial sequences that imply recurrent glaciation of the continental  
16 shelf between MIS 12 and MIS 2 (Skinner and Gregory, 1983; Stoker et al., 1985,  
17 1993, 1994; Cameron et al., 1987; Sejrup et al., 1987, 1991, 1994, 2000, 2005;  
18 Johnson et al., 1993; Gatliff et al., 1994; Holmes, 1997). This has resulted in a glacial  
19 succession that is locally several hundreds of metres thick.

20

21 Controversy surrounds the timing of onset of the ‘Late Devensian glaciation’ (MIS 2)  
22 in Britain. Traditionally the presence of ice-free conditions in Scotland during the  
23 Middle Devensian has been based on a handful of key sites. Organic lake sediments  
24 beneath a till at Tolsta Head in northern Lewis yielded a radiocarbon date of  $27,333 \pm$   
25  $240$   $^{14}\text{C}$  BP (Von Weymarn and Edwards, 1973); whilst organic deposits beneath till  
26 at Sourlie, near Glasgow, suggest that an ice-free, periglacial environment prevailed  
27 here at around the same time ( $29\text{-}33$  ka  $^{14}\text{C}$  BP =  $34\text{-}38$  ka cal BP) (Jardine et al.,  
28 1988). These two sites are often used to indicate that much of Scotland was ice free in  
29 MIS 3 (e.g. Gordon and Sutherland, 1993).

30

31 Based on numerous cosmogenic dates from Ireland, Bowen et al. (2002) speculate  
32 that the BIS had reached its maximum size by 37 ka BP, and in doing so challenged  
33 the validity of the radiocarbon dates from Sourlie and elsewhere. However, six AMS  
34 dates of between 34,480 and 28,050  $^{14}\text{C}$  BP on *Carex* fruit and Coleopteran fragments

1 have been recently reported from Balglass, close to the southern end of the Loch  
2 Lomond basin (Brown et al., 2007). The presence of these glacitectonised organic  
3 deposits constrain the onset of the last regional glaciation in Scotland to c. 31.5 ka <sup>14</sup>C  
4 BP (36.5 ka cal BP), because they are located close to an important conduit of the last  
5 ice sheet and they occur between two distinct tills, the lower one being weathered. A  
6 detailed re-investigation of the fossil flora and fauna at Sourlie by Bos et al. (2004)  
7 produced four new conventional radiocarbon dates of between 33.3 and 29.3 ka <sup>14</sup>C  
8 BP, reinforcing the original findings of Jardine et al. (1988). Furthermore, a re-  
9 investigation of the organic horizon within the Tolsta Head deposits (Whittington and  
10 Hall, 2002) has yielded seven AMS dates spanning the period 31,700 ka to 26,150 <sup>14</sup>C  
11 BP (31-37 ka cal BP), similar in age to those from Balglass and Sourlie. On the basis  
12 of these results it is evident that much of Scotland was ice free towards the end of  
13 MIS 3, with build up of the last BIS occurring from 35-32 ka cal BP onwards.

14

15 It is worth noting that the presence of reindeer bones found in limestone caves near  
16 Inchnadamph and radiocarbon dated to between ~22-32 ka <sup>14</sup>C BP (c. 25-37 ka cal  
17 BP) suggest that the NW Highlands were substantially ice-free at this time (Lawson,  
18 1984; Murray et al., 1993). However, these dates are likely to be anomalously young,  
19 as demonstrated by bone remains on carbonate geology elsewhere (Jacobi et al.,  
20 1998; Hedges and Millard, 1995). Furthermore, improved analytical techniques  
21 (ultrafiltration) have recently resulted in revised <sup>14</sup>C ages for Pleistocene faunal  
22 remains from the UK (Higham et al., 2006), casting further doubt on the radiocarbon  
23 ages of the Inchnadamph bones.

24

### 25 *Offshore evidence*

26 Key indicators of glacial activity on the UK continental shelf include several  
27 generations of tunnel valleys in the North Sea (Cameron et al., 1987; Wingfield, 1989,  
28 1990; Ehlers and Wingfield, 1991; Huuse and Lykke-Andersen, 2000; Praeg, 2003;  
29 Lonergan et al., 2006), and moraines preserved to the NE and NW of Britain (e.g.  
30 Rokoengen et al., 1982; Stoker et al., 1985, 1993; Selby, 1989; Hall and Bent, 1990;  
31 Stoker and Holmes, 1991; Austin and Kroon, 1996). The continuation of these glacial  
32 sequences across the shelf edge to the north and west of Britain indicates that ice-  
33 marginal and proglacial processes have contributed to the growth of the slope aprons  
34 bordering the Hebrides and West Shetland shelves. This is expressed by the

1 development of large trough-mouth fans, including the Barra-Donegal and Sula Sgeir  
2 Fans and the Rona and Foula Wedges (Stoker, 1995, 2002; Bulat and Long, 2001;  
3 Davison and Stoker, 2002; Holmes et al., 2003; Long et al., 2004) (Figures 1, 3).

4  
5 [FIGURE 3 HERE]

6  
7 The recognition of subglacial till in BGS borehole 77/2 in the Witch Ground Basin  
8 (Figure 1), dated at between 43 and 22 ka <sup>14</sup>C BP led Sejrup et al. (1994) to propose  
9 coalescence of the BIS and FIS during the interval 35-26 cal ka BP. A confluent BIS-  
10 FIS is also consistent with micromorphological studies that have recognised a much  
11 more extensive cover of subglacial till within the central and northern North Sea  
12 glacial succession (Carr et al., 2006). Graham et al. (2007) have recently reported  
13 unambiguous evidence for grounded ice crossing the northern North Sea Basin. From  
14 3D seismic data, they identify highly elongate (up to 150:1) MSGs – formed on top  
15 of the subglacial till unit described by Sejrup et al. (1994) in BGS borehole 77/2. The  
16 MSGs are orientated NW–SE, and help define a palaeo-ice stream, at least 30 to 50  
17 km wide and >90 km in length (the extent of the 3D image). Graham et al. (2007)  
18 refer to this as the Witch Ground palaeo-ice stream, and infer that it was sourced from  
19 the FIS overspilling the Norwegian Channel during the LGM (MIS 2).

20  
21 A number of discrete overdeepened troughs have been identified crossing the  
22 Hebrides and West Shetland shelves, linking the Scottish hinterland to the adjacent  
23 continental slope (Stoker, 1990, 1995; Stoker et al., 1993). The location of these  
24 major cross-shelf troughs is marked by the landward indentation of the 100 m  
25 bathymetric contour. Significantly, each of these pathways links the Scottish  
26 hinterland to a specific trough-mouth fan on the adjacent continental slope (Figure 1).  
27 The best studied of these pathways links The Minch to the Sula Sgeir Fan. In total,  
28 this trough is about 200 km long, up to 50 km wide, and contains locally thick  
29 accumulations (50–150 m) of subglacial and proglacial sediments, including basal till,  
30 multiple ice-contact sequences, stratified proglacial outwash and glacial marine  
31 sediments (Stoker et al., 1994; Stoker and Bradwell, 2005). The occurrence of highly  
32 elongate (up to 70:1) MSGs, identified at several stratigraphic levels within the  
33 trough infill, has been cited as evidence of a palaeo-ice stream. This fast-flow corridor  
34 – The Minch palaeo-ice stream – was responsible for draining about 15 000 km<sup>2</sup> of

1 the NW sector of the BIS and probably operated during several Mid- to Late  
2 Pleistocene glaciations, including the LGM (Stoker and Bradwell, 2005; Bradwell et  
3 al., 2007).

4  
5 The Barra–Donegal Fan represents a major focus of glacial sediment (Knutz et al.,  
6 2001; Wilson et al., 2002), most probably fed by ice streams that periodically crossed  
7 the continental shelf, draining much of western Scotland and northwest Ireland. Ice-  
8 rafted debris recovered from the Barra Fan, sourced from British volcanic rocks,  
9 suggest extensive glaciation c. 45 ka BP prior to full glaciation c. 27 ka BP (Knutz et  
10 al., 2001; Peck et al., 2007). Further north, the Rona and Foula Wedges also represent  
11 trough-mouth fans probably fed by focused flow zones acting between northern  
12 mainland Scotland and Shetland during the LGM (Stoker et al., 1993; Davison,  
13 2005).

14  
15 *Onshore evidence*

16 Onshore landform evidence confirms that during the last ice-sheet glaciation several  
17 large fast-flow zones dominated the northern sector of the BIS. An ice stream sourced  
18 in the western Scottish Highlands stretched to the Moray Firth, invading the coastal  
19 lowlands of Moray, Banffshire and Buchan (Merritt et al., 2003). This same ice  
20 stream also flowed north-westwards across Caithness and Orkney (Hall and Bent,  
21 1990), laying down shelly tills, and rafts of Mesozoic strata and Pleistocene marine  
22 sediment dredged up from the seafloor (Sutherland, 1984; Gordon and Sutherland,  
23 1993). Shells, dated by AMS, from within the ice-stream till in northern Caithness  
24 indicate that it was laid down after 46 ka <sup>14</sup>C BP (Auton, 2003). The pronounced  
25 deflection of ice flow in the Moray Firth NW across Caithness was most likely caused  
26 by the presence of Scandinavian ice in the central North Sea (Peach and Horne, 1879;  
27 Sissons, 1967; Hall and Whittington, 1989; Sutherland, 1984). In NW Scotland a  
28 powerful ice stream sourced in the NW Highlands, Skye and Lewis flowed north  
29 along The Minch and then northwest onto the continental shelf (Stoker and Bradwell,  
30 2005; Bradwell et al., 2007). An ice stream probably also occupied the Vale of  
31 Strathmore, Firth of Forth and adjoining lowlands, flowing broadly northeast towards  
32 the North Sea Basin (Golledge and Stoker, 2006). Other ice streams have been  
33 identified within the northern sector of the last British Ice Sheet: flowing northeast



1 from the Cairngorms (Hall and Glasser, 2003) and along the Tweed Valley towards  
2 the North Sea (Clapperton, 1970; Everest et al., 2005).

3  
4 Glacial evidence on Orkney and Shetland indicates that an ice sheet sourced in  
5 Scandinavia occupied the northern North Sea Basin. The presence of rare Norwegian  
6 erratics on Sanday (Orkney), Fair Isle and on southernmost Shetland has long been  
7 taken as evidence that a far-travelled ice sheet crossed these islands (Peach and  
8 Horne, 1879, 1880; Finlay, 1926; Birnie et al., 1993). Ice-flow indicators on Orkney,  
9 most notably striae and ice-directional bedforms, clearly suggest the passage of a  
10 powerful ice sheet in a west-northwesterly direction, even on the northernmost islands  
11 (Peach and Horne, 1880, 1983; Wilson et al., 1935; Rae, 1976). The presence, on Fair  
12 Isle, of large-scale bedrock grooves and streamlined forms trending in the same  
13 orientation (WNW) is key, as this island has not been affected by glaciers since the  
14 LGM (Mykura, 1976). This evidence is incompatible with the deflection of British ice  
15 from the Moray Firth on glaciological grounds, but is entirely consistent with glacial  
16 overriding from the southeast by a coalescent BIS-FIS. Although most workers have  
17 accepted that this ice-sheet configuration existed (e.g. Wilson et al., 1935; Boulton et  
18 al., 1977, 2002; Flinn, 1978; Sutherland, 1984; Sejrup et al., 1994; Carr et al., 2006),  
19 some have argued that it predated the LGM (Sutherland, 1984; Bowen, 1989; Bowen  
20 et al., 2002; Hall et al., 2003). Much of this debate rests on the evidence for a local  
21 Shetland ice cap, thought by many to equate to the LGM. However, there is no reason  
22 why the presence of a locally nourished ice cap on Shetland is incompatible with the  
23 islands being overwhelmed by an ice sheet during the early part of the same  
24 glaciation, as has previously been suggested (e.g. Flinn, 1978; Stoker et al., 1993;  
25 Ross, 1996; Carr et al., 2006; Golledge et al., 2008).

### 26 27 *Extent of glaciation*

28 The maximum extent of the last ice sheet to cover northern Britain is currently poorly  
29 defined (Clark et al., 2004; Evans et al., 2005). The most comprehensive attempts at  
30 ice-sheet reconstruction on the continental shelf, based on offshore stratigraphy and  
31 geomorphology, are those of Stoker et al. (1993, 1994) and Hall et al. (2003). These  
32 reconstructions incorporate geomorphological indicators – end moraines – preserved  
33 on the Hebrides and West Shetland shelves (Figure 1) (Selby, 1989; Stoker and  
34 Holmes, 1991), and suggest an expansive LGM, reaching the continental-shelf edge in

1 most places except, perhaps, on the northern Hebrides Shelf. Here the presence of  
2 undisturbed glacimarine deposits on the outer shelf, with Amino-acid ratios typical of  
3 MIS 4 (Stoker et al., 1993), suggests that the grounded ice-sheet limit was on the mid-  
4 shelf during LGM (Stoker and Bradwell, 2005). However, the reliability and precision  
5 of this dating technique has been seriously questioned (McCarroll, 2002). Expansive  
6 glaciation across the southern Hebrides Shelf is supported by increased sedimentation  
7 on the Barra Fan in MIS3 (Kroon et al., 2000; Knutz et al., 2001, 2002; Wilson et al.,  
8 2002), followed by a major increase c. 26 ka BP (Peck et al., 2007) associated with  
9 widespread shelf-edge glaciation at LGM. Other published reconstructions and  
10 models, which propose that the last ice sheet terminated only a short distance offshore  
11 at its maximum, are therefore probably too conservative (e.g. Sutherland, 1984;  
12 Bowen et al., 1986, 2002; Lambeck, 1993, 1995; Ballantyne et al., 1998; Stone et al.,  
13 1998).

14

15 Prior to 1980, textbook reconstructions showed the FIS crossing Shetland during the  
16 last glaciation and deflecting the BIS northwest across Orkney (e.g. Sissons, 1976;  
17 Boulton et al., 1977). However, oil-related exploration of the central North Sea Basin  
18 in the 1970s and 1980s revealed an apparent absence of Late Devensian/Late  
19 Weichselian tills (e.g. Cameron et al., 1987). This led to Sutherland's (1984)  
20 reconstruction of an independent BIS of restricted size terminating at the Wee Bankie  
21 and Bosies Bank moraines off eastern Scotland and on Lewis, NW of the Scottish  
22 mainland (Figure 1). Radiocarbon dates between 21.7 ka and 17.7 ka <sup>14</sup>C BP,  
23 obtained from lignitised wood sampled from glacimarine deposits adjacent to the  
24 eastern margin of the Wee Bankie moraine (Holmes, 1977), have traditionally formed  
25 the basis for interpreting these moraines as the maximum eastern limit of the last BIS,  
26 at around 18-22 ka BP (Sutherland, 1984; Boulton et al., 1985, 1991, 2002; Cameron  
27 et al., 1987; Hall and Bent, 1990; Lambeck, 1991; Bowen et al., 2002; Clark et al.,  
28 2004). However, such a limit fails to reconcile the occurrence of subglacial tunnel  
29 valleys largely formed during the Late Devensian (MIS 2), located east of these  
30 moraines (Wingfield, 1989; Ehlers and Wingfield, 1991; Lonergan et al., 2006).  
31 Many of the tunnel valleys remain exposed at the present-day sea bed (e.g. Devil's  
32 Hole Deeps and Fladen Deeps (Figure 1) and, hence, must have formed before ice  
33 retreated to the Wee Bankie and Bosies Bank moraines.

34

1 Unglaciaded enclaves such as those proposed by Sutherland (1984) in Caithness and  
2 Buchan have subsequently appeared on many published reconstructions of the  
3 northern sector of the last BIS (e.g. Bowen et al., 1986; Nesje and Sejrup, 1988;  
4 Lambeck, 1993, 1995; Bowen et al., 2002), some long after Sutherland's evidence  
5 had been seriously questioned by Peacock (1985), and then refuted by Stoker et al.  
6 (1993), Merritt et al. (2003) and Hall et al. (2003).

7  
8 Controversy also surrounds the vertical extent and thickness of the last BIS. Although  
9 high-level trimline evidence from mountains has been used to reconstruct the ice-  
10 sheet surface in NW Scotland (Ballantyne et al., 1998), the validity of trimlines as ice-  
11 sheet surface indicators has been seriously questioned (Fabel et al., 2002; Boulton and  
12 Hagdorn, 2006; Shennan et al., 2006; Kleman and Glasser, 2007). Cosmogenic-  
13 isotope analyses by Stone et al. (1998) were unable to determine whether mountain  
14 summits in NW Scotland were buried beneath non-erosive cold-based ice or if they  
15 had experienced long subaerial exposure histories. It is therefore uncertain whether  
16 ice-free areas existed in Scotland during the LGM, when the ice sheet terminated on  
17 the continental shelf.

18  
19 Evidence of glacial activity directly attributed to the FIS is preserved in the  
20 northeastern part of the North Sea, where the Tampen Ridge (Figure 1) is interpreted  
21 as a lateral moraine associated with the western flank of the Norwegian Channel  
22 palaeo-ice stream. This moraine has been attributed to a minor readvance of the ice  
23 margin on the flank of the Norwegian Channel (Sejrup et al., 1994, 2000).  
24 Radiocarbon dates from shells indicate a maximum age of  $18.86 \pm 2.6$  ka  $^{14}\text{C}$  BP for  
25 the moraine (Rokoengen et al., 1982).

26  
27 Palaeo-ice streams that link the source regions of the BIS to the continental shelf and  
28 slope are recognised both east and west of Scotland (Clapperton, 1970; Everest et al.,  
29 2005; Stoker and Bradwell, 2005; Golledge and Stoker, 2006). Palaeo-ice stream  
30 bedforms have strong morphological expression on the extant landsurface and sea  
31 bed, suggesting that these ice streams probably operated during deglaciation of the  
32 last BIS. Significant dates and other key elements relating to deglaciation of the  
33 northern sector of the last BIS are summarised below.

1 A sequence of glacial marine sediments overlies the MSGL surface in the Witch Ground  
2 Basin, dated at between ~22 and 13 ka <sup>14</sup>C BP (Sejrup et al., 1994; Graham et al.,  
3 2007). These sediments record repeated iceberg scouring, which has been taken to  
4 indicate the presence of a marine embayment – an ice-free enclave – within the  
5 central and northern North Sea from about 22 ka <sup>14</sup>C BP (~26 ka cal BP) onwards. It  
6 has been suggested that the Wee Bankie and Bosies Bank moraines were formed in  
7 this deglacial phase (Carr et al., 2006). Sporadic readjustments of the receding ice  
8 margin offshore NE Scotland may have resulted in readvances, between about 18 and  
9 16 ka <sup>14</sup>C BP (Merritt et al., 2003), correlated with the Tampen Readvance in the NE  
10 North Sea (~20 ka cal BP; Sejrup et al., 1994, 2000) and possibly the Dimlington  
11 Stadial advance in eastern England (Rose, 1985; Wintle and Catt, 1985). However,  
12 the preservation of shoreline fragments, such as the Main Perth Shoreline, and the  
13 occurrence of shelly glacial marine sediment overlying till in NE Scotland indicate that  
14 the adjacent coastal area may have been largely ice free by 15–16 ka <sup>14</sup>C BP  
15 (Cullingford and Smith, 1980; Armstrong et al., 1985; Hall and Jarvis, 1989),  
16 although this is still debated (cf. McCabe et al. 2007; Peacock et al., 2007). Offshore  
17 SW Norway, the Norwegian Channel was deglaciated by about 15.1 ka <sup>14</sup>C BP (~18  
18 ka cal BP)(Lehmann et al., 1991).

19

20 North of Shetland, shell fragments from a diamicton (probably till), equivalent in  
21 stratigraphic status to the large sea-floor moraines west of Shetland (Figure 1), date  
22 from 17.8 ka <sup>14</sup>C BP (~21 ka cal BP)(Ross, 1996); whereas the earliest organic  
23 deposits (overlying till) on Shetland date from 13 ka <sup>14</sup>C BP or a little earlier (Hoppe,  
24 1974; Birnie et al., 1993). On the southern Hebrides Shelf, the formation of the St  
25 Kilda moraine banks is dated to between 22.48 and 15.65 ka <sup>14</sup>C BP (Selby, 1989;  
26 Austin and Kroon, 1996). In The Minch region, recently performed cosmogenic <sup>10</sup>Be  
27 analyses of glacially deposited boulders on the east coast of Harris yield exposure  
28 ages of 15.4 to 17.6 ka cal BP (Stone and Ballantyne, 2006). Cosmogenic <sup>10</sup>Be  
29 analyses of boulders on an ice-sheet moraine in Gairloch, Wester Ross (Figure 1),  
30 yield a comparable age (15-17 ka cal BP: Everest et al., 2006). These data, combined  
31 with environmental information from marine fauna in borehole 78/4 east of Lewis  
32 (Graham et al., 1990) (Figure 2) and from several short cores south and west of St  
33 Kilda (Austin and Kroon, 1996) indicate open marine conditions from >12,785 ka <sup>14</sup>C  
34 BP (c.15 ka cal BP) in The Minch and from >13.5 ka <sup>14</sup>C BP around St Kilda. These

1 dates provide minimum ages for deglaciation of the continental shelf off northern  
2 Britain. By 15 ka cal BP arctic open-water conditions existed on the continental shelf  
3 when the ice-sheet margin was situated at, or close to, the present-day coastline of  
4 NW Scotland (Everest et al., 2006; Stoker et al., 2006; Stone and Ballantyne, 2006).

5  
6 Following LGM it is thought that the eastern margin of the last ice sheet retreated  
7 slowly in a cold climate, owing to precipitation starvation (Sutherland, 1984). Dates  
8 from eastern Scotland suggest that the ice-sheet margin had retreated to the present-  
9 day coastline in Buchan by about 18 ka cal BP (Peacock and Merritt, 2000; Merritt et  
10 al., 2003). During deglaciation a number of prominent push moraines were formed  
11 including one at Ardersier, near Inverness, probably dating from c. 15 ka cal BP  
12 (Merritt et al., 1995). This event has been tentatively correlated with the Killard Point  
13 Readvance in Ireland, which in turn has been linked with the North Atlantic iceberg-  
14 discharge event Heinrich-1 (McCabe et al., 1998, 2005). Deglaciation of the west  
15 coast of Scotland was also punctuated by a series of ice-marginal stillstands or  
16 oscillations. The Wester Ross Readvance, dated to c.15-17 ka cal BP, may have also  
17 been in response to wider events in the North Atlantic (i.e. Heinrich-1) (Everest et al.,  
18 2006).

## 19 20 21 22 23 **Data Compilation and Methods**

24 This section outlines the datasets and techniques used to elucidate the geomorphology  
25 of the UK continental shelf (Figures 4, 5).

26  
27 [FIGURE 4 HERE]

### 28 29 *Offshore data*

30 The marine dataset (Figure 4) is part of the Olex bathymetric database compiled,  
31 processed and managed by the Norwegian company Olex AS ([www.olex.no](http://www.olex.no)). The  
32 sea-bed image is based upon echosounder data acquired mainly by commercial fishing  
33 vessels, but also including data from research vessels. The datasets are contributed  
34 voluntarily, the data is then individually merged with the central dataset, after which  
35 the contributor has access to all of the shared bathymetry. The data are located by

1 global positioning systems (GPS) and the positional error is generally less than 10 m.  
2 The database represents the earth's surface as a series of 5 x 5 m cells. Vertical  
3 resolution is 1 m in water depths >100 m and 0.1 m at depths <100 m. Horizontal  
4 datum is WGS84; vertical reference is equinoctial spring low water from predicted  
5 tides. The speed of sound in water is harmonised to 1500 ms<sup>-1</sup>. This means that  
6 relative depths are highly accurate with an error range of only 1-2%. The density of  
7 soundings depends on instrumental output rates; ideally systems record one sounding  
8 for every echosounder value. Depth position is adjusted for installation offsets and  
9 timing between the echosounder and the GPS. The strength of the Olex system lies in  
10 the integration of data contributed by a number of users over several years. Any depth  
11 errors are minimised during database compilation and processing by comparing an  
12 individual contributor with the large number of other soundings covering the same  
13 area. The resultant bathymetric surface can be viewed as 2D contours, 2D shaded  
14 relief, 3D views or as 2D profiles. A 2D shaded relief map, with illumination from the  
15 north, is shown in Figure 4.

16

#### 17 *Onshore data*

18 The onshore topography (Figure 4) is from NEXTMap Britain (Intermap  
19 Technologies, 2003). Heights are acquired from an aircraft using interferometric  
20 synthetic aperture radar (IFSAR) with a vertical resolution of +/- 1 m (95%) and a  
21 grid cell size of 5 m. Artefacts in areas of steep slopes are possible where the data are  
22 degraded either by the rate of change in elevation or by 'shadows' on the far side of a  
23 positive object or on the nearside of a steep depression. The dataset is provided in UK  
24 Ordnance Survey GB36 projection (as derived from OSTN97) and referenced to  
25 Ordnance Datum. The surface model is not vertically exaggerated but is illuminated  
26 from the NW at an angle of 45° to highlight the relief.

27

#### 28 *Methods*

29 The processed echosounder data were georectified and merged in a geospatial  
30 database, from which a sea-bed surface model was generated (Figure 4). The use of  
31 three-dimensional vector data allowed surface models to be illuminated and viewed  
32 from any angle (using Olex or ESRI software) thus enabling clear, accurate  
33 identification of sea-bed morphology. All positive and negative, linear, bathymetric  
34 features within the offshore area (Figure 1) were digitized on screen in ArcGIS 9.0

1 (ESRI). Onshore landforms were not digitized. The data capture method used is  
2 similar to that outlined by Stokes and Clark (2003) and Golledge and Stoker (2006).  
3 To overcome problems of azimuth bias, identified by Smith and Clark (2005), digital  
4 surface models were illuminated first from the northeast and then from the northwest.  
5 Digitizing scales varied between 1:50,000 and 1:200,000. Landform dimensions, such  
6 as length, height and width were measured digitally within the GIS. The final dataset  
7 was output in map format at A0 size, whereby some generalization was made for  
8 cartographic clarity (Figure 5).

9

10 [FIGURE 5 HERE]

11

12 **Results**

13

14 Here we describe the morphology and spatial distribution of sea-bed landforms  
15 imaged in the bathymetric Olex dataset (Figure 4). Relevant geographical information  
16 is shown in Figure 1.

17

18 The continental shelf around the UK is presently less than 160 m below sea level,  
19 with the exception of isolated deeps that incise to depths locally in excess of 200 m  
20 (Figure 1). The Witch Ground Basin and the Norwegian Channel are the most notable  
21 large bathymetric depressions on the continental shelf between Britain and Norway  
22 reaching depths of 155 and 415 m respectively. The shelf break to the north and west  
23 of Britain generally occurs around 200 m below present-day sea level and slopes at  
24 between 1 and 6 degrees (Figure 1).

25

26 The Olex data reveal channels and ridges across the majority of the sea bed around  
27 northern Britain, in particular: around St Kilda; west of Orkney and Shetland; in the  
28 Moray Firth; offshore Strathmore; and flanking the western margin of the Norwegian  
29 Channel. By contrast, the Witch Ground Basin in the northern North Sea is largely  
30 devoid of landforms. Systematic analysis of the Olex dataset has resulted in over 700  
31 individual landform elements being digitised in a GIS:– 174 negative linear  
32 topographic features (channels) and 537 positive linear topographic features (ridges)  
33 (Figure 5).

34

1 *Channels*

2 The digitised negative features fall broadly into two groups: A) a north to northwest-  
3 trending population widely distributed across the northern North Sea Basin; B) and a  
4 second set that lie close to the present coastline of eastern Scotland (Figure 5).

5

6 Group A comprises 67 channels ranging from 3 to 50 km in length. The channels have  
7 a strongly consistent orientation, trending north to northwest, and are relatively evenly  
8 spaced across a zone about 300 km west to east and 450 km from north to south. The  
9 majority of the channels occur around the northern margin of the Witch Ground  
10 Basin; whilst others are located on the southern edge of the basin; and west of the  
11 Norwegian Channel (Figure 3). This group of major channels includes the Fladen  
12 Deeps (Figure 1) which are cut up to 280 m below sea level, with individual channels  
13 up to 4.5 km in width (Andrews et al., 1990; Johnson et al., 1993) (Figure 6).

14

15 [FIGURE 6 HERE]

16

17 Group B consists of two main sets of channels in geographically separate areas. The  
18 first set of 26 channels flanking the east coast of mainland Scotland, trend NNE  
19 roughly parallel to the present coastline. These features range in length from 2 to 30  
20 km, are relatively evenly spaced, and incise an area 40 x 65 km. The channels are cut  
21 up to 120 m below sea level and range from 1.5 to 3 km wide (Thomson and Eden,  
22 1977; Golledge and Stoker, 2006). Many have branching, sinuous courses. The  
23 second set of 60 channels, occur in the outer Moray Firth and are tightly grouped in an  
24 area covering 60 x 65 km. They trend broadly west to east and range in length from  
25 1.5 to 58 km. Many have branching, sinuous courses. Approximately three quarters of  
26 the channels are greater than 10 km in length. The longest of these, the Southern  
27 Trench (Figure 1), is 40 km long has a maximum width of 9 km and is locally up to  
28 200 m deep (Andrews et al., 1990). Numerous other isolated channels, with similar  
29 dimensions to those above, occur on the mid-shelf west of Shetland and in the Fair  
30 Isle Channel.

31

32 In both of these groups, thalwegs do not exhibit consistent downstream deepening but  
33 undulate along the length of the channel (Figure 6). Most channels in the northern  
34 North Sea Basin begin or terminate abruptly; many possess branching tributaries. It



1 should also be noted that whereas these channels retain bathymetric expression, due to  
2 only partial sediment infill, they form part of a more extensive system of north to  
3 northwest-trending channels in the central and northern North Sea Basin that have  
4 been subsequently buried by sediment (Stoker et al., 1985; Gatliff et al., 1994; Praeg,  
5 2003; Fitch et al., 2005; Lonergan et al., 2006). Whether open or infilled, the bases of  
6 all these channels generally lie between 50 and 100 m below the surrounding sea bed.  
7 Consequently, the area of the North Sea Basin dissected by large-scale channels is  
8 much greater than shown by the Olex data alone.

9

### 10 *Ridges*

11 Over 500 crestlines were digitised from the Olex dataset. These are broadly divided  
12 into three groups based on geographical setting (Figure 5).

13

14 Group 1: This group comprises the westernmost landforms, occurring on the  
15 outermost shelf. They are large, curvilinear, gently arcuate ridges, concave to the east,  
16 trending approximately northeast-southwest. Most are broad features, 2 to 10 km  
17 wide, with moderately well-defined seabed expression. The two longest of these  
18 ridges run unbroken for over 60 km; between St Kilda and the Flannan Isles, and  
19 close to the shelf break west of Shetland (Figure 5). Previously, some of these broad  
20 ridges have been mapped from seismic profiles as shelf-edge ice-sheet moraines  
21 (Stoker et al., 1993; Stoker and Holmes, 1991) (Figures 1, 3).

22

23 Group 2: This group comprises the mid-shelf ridges to the north and west of Scotland.  
24 Most of these are large, strongly arcuate ridges, with well-defined expression. These  
25 ridges can be further subdivided based on location: (a) those west of Orkney and  
26 Shetland; (b) those north and east of Shetland; (c) those around St Kilda. A further  
27 subset of more-linear Group 2 ridges occurs adjacent to the eastern margin of the  
28 Norwegian Channel.

29

30 [FIGURE 7 HERE]

31

32 Those ridges on the mid-shelf, west of Orkney and Shetland are typically 10 to 100  
33 km in length, and often occur as nested, lobate forms. The striking pattern of ridges,  
34 NW of Orkney is strongly concentric from west to east (Figure 7). Here, up to 10

1 large nested ridges occur within a horizontal distance of 60 km. The outer ridges are  
2 broad features, up to 6 km wide and 50 m high, some display pronounced crenulate or  
3 zig-zag morphologies. Many of the ridges show overprinting patterns. The innermost  
4 ridge is a delicate sharply defined feature, <1 km wide, with simple arcuate plan form.  
5 This innermost ridge forms a remarkable unbroken loop ~70 km in length. Further  
6 south, due west of Orkney, numerous large curvilinear ridges trend perpendicular to  
7 the coastline of mainland Scotland. Here too, overprinting can be seen between these  
8 ridges and those to the north. Further north, due west of Shetland (Figure 8),  
9 numerous ridges form a separate set of sub-concentric nested arcs and loops.  
10 Overprinting of one ridge on another can be clearly seen in places, particularly on the  
11 sea bed NW of Foula. The innermost mid-shelf ridges west of Shetland are  
12 approximately parallel to the present-day coastline.

13

14 The greatest concentration of large ridges on the UK continental shelf occurs north  
15 and east of Shetland (Figures 4, 5, 8). The northernmost ridges have irregular outlines  
16 with highly crenulated, distinctly lobate form. The largest of these extends for 30 km,  
17 is between 1 and 3 km wide, and locally exceeds 30 m in height. The nested pattern of  
18 these lobate ridges follows a broadly north-south axis. Overprinting of ridges on one  
19 another can be clearly seen.

20

21 Two morphologically similar mid-shelf ridges occur on the seabed southwest of St  
22 Kilda on the Hebrides Shelf (Figure 5). These large features are 30 to 40 km in length  
23 and 2 to 5 km wide. Like those west of Orkney and Shetland they have pronounced  
24 lobate and crenulated plan form, and are concave towards the east.

25

26 A further set of curvilinear, subparallel Group 2 ridges flank the western margin of the  
27 Norwegian Channel trending approximately southeast-northwest. This high-density  
28 cluster of ridges lies on the relatively flat sea-bed high west of the Norwegian  
29 Channel and east of Shetland (Figure 5). The ridges range in length from 2 to 55 km,  
30 and occur within a broad zone c. 600 km long and c. 100 km wide. The majority have  
31 subtle sea-bed expression; many are less than 10 m high. Individual ridges are linear  
32 or curvilinear in plan. To the north, a well-defined and distinct linear ridge occurs  
33 along the Norwegian Channel margin – it has been mapped previously as the Tampen  
34 Ridge (Figure 1) (Sejrup et al., 1994) and is formed of a stratigraphic unit 50 m thick

1 and up to 7 km wide. This ridge can be traced for ~200 km along the western edge of  
2 the Norwegian Channel (Figure 5).

3

4 Group 3: The final group of ridges are those on the inner shelf, closer to the present  
5 day coastline of Scotland, generally east of Groups 1 and 2. These smaller-scale  
6 features range in length from 500 m to 25 km and are typically <500 m wide and <20  
7 m high. They are widely geographically distributed and many occur as concentrations  
8 of closely spaced ridges (Figure 5).

9

10 Immediately east of Shetland, numerous closely spaced linear and curvilinear ridges  
11 occur, trending NW-SE. These features are generally concave towards the south, and  
12 are best preserved within a topographic sea-bed trough, although many extend beyond  
13 its lateral margins (Figure 8). Some of the landforms within the trough have clearly  
14 visible zigzag morphologies whilst others are more linear. Crestline orientations are  
15 strongly oblique to the trough margins and cut across topographic undulations. The  
16 ridges range in length from 1 to 20 km, and exhibit intra-ridge spacings of 700 to  
17 2000 m. Most ridges are well-defined features with vertical sea bed expression of  
18 between 10 and 20 m.

19

20 [FIGURE 8 HERE]

21

22 Southeast of Shetland few clear landforms are resolved on the Olex sea-bed image.  
23 However, a cluster of subparallel linear ridges occur around 1°E, 59°N. These  
24 features range from 1.5 to 20 km in length, have heights generally <10 m, widths  
25 <100 m, and exhibit consistent spacing distances of 2.5 to 3 km. Within this  
26 population there are also shorter, WNW-ESE-trending linear ridges 1.5 to 6 km in  
27 length (Figure 3).

28

29 Northeast of Orkney is a cluster of short ridges, 800 m to 6 km in length, aligned  
30 WSW-ENE. As with those east of Shetland, these 1 to 2 km-spaced ridges occur  
31 within a topographic trough, but are aligned oblique to its margins (Figure 3).

32

33 In the generally flat ground northeast of the Moray Firth and northwest of the Witch  
34 Ground Basin, a spatially distributed population of nearly 100 ridges is mapped. Most

1 are aligned generally northwest-southeast, are curvilinear in plan and less than 10 m  
2 in height. These ridges range in length from 1 to 35 km (Figure 5), with the exception  
3 of the large, arcuate, outermost ridge in this group which can be traced for 65 km.

4  
5 Other well-defined small-scale Group 3 ridges occur in coastal parts of The Minch,  
6 off NW Scotland; adjacent to western Orkney; and in the Fair Isle Channel (Figure 3).  
7 These features are considerably smaller than the ridges seen on the mid- and outer  
8 shelf, being less than 10 m high and <1000 m wide. They typically range from 5 to 20  
9 km in length and generally occur within ~50 km of the present-day coastline. The  
10 features in The Minch and adjacent to Orkney are broadly parallel to the coastline,  
11 whereas some of the ridges off the east coast of Shetland are perpendicular to the  
12 coast. An assemblage of small, sharply defined, subparallel ridges occur in the Fair  
13 Isle Channel. They are distinctive linear features, less than 10 m high, extending over  
14 30-40 km and trending NNE-SSW (Figure 8).

15  
16 Finally it should be noted that large parts of the seabed with good echosounder  
17 coverage have a poor landform record. Most notably, parts of the shelf north of  
18 Orkney and southwest of Shetland; the inner shelf immediately west of Lewis; the  
19 shelf to the southwest of Barra; and the central core of the Witch Ground Basin. With  
20 the exception of the Witch Ground Basin, which is underlain by thick Quaternary  
21 sediment (Long et al., 1986), the absence of landforms is probably related to the  
22 presence of crystalline bedrock at seabed in these areas (Johnson et al., 1993; Stoker  
23 et al., 1993).

## 24 25 26 **Landform interpretation**

27 Sea-bed mapping based on the Olex dataset reveals over 700 geomorphological  
28 features relating to glaciation of the continental shelf around the northern UK (Figures  
29 4, 5). We interpret these features by comparison with glacial landsystems described  
30 elsewhere (e.g. Evans, 2003), coupled with inferences from existing  
31 seismostratigraphic data within the study area (e.g. Stoker and Holmes, 1991;  
32 Davison, 2005; Sejrup et al., 2005). On the basis of their morphology, distribution and  
33 setting we interpret the vast majority of positive linear features as end moraines  
34 marking the margins of former ice masses. Whilst some of the smaller near-shore

1 features (Group 3) are distinctly straight, the majority of mapped moraines are  
2 arcuate, lobate or curvilinear in form (Groups 1 and 2) – typical of features formed at  
3 grounded terrestrial ice margins (e.g. Benn and Evans, 1998; Colgan et al., 2003).  
4 Most studies agree that end moraine morphology closely reflects the geometry and  
5 position of former ice margins. Consequently, their distribution, orientation and size  
6 are commonly used in glaciological inversion models to reconstruct the decay  
7 dynamics of former ice sheets and glaciers (e.g. Kleman et al., 1997; Clark et al.,  
8 2004).

9 The population of straight, sub-parallel, sharp-crested ridges identified in the vicinity  
10 of Orkney and Shetland, by contrast, more closely resemble moraines formed at the  
11 subaqueous grounding line of a marine-terminating ice-sheet margin, and  
12 consequently are interpreted as De Geer-type moraines (e.g. Sollid, 1989; Larsen et  
13 al., 1991; Blake, 2000; Linden and Moller, 2005).

14 We interpret the negative linear features on the sea-bed as glacial meltwater channels,  
15 largely on the basis of their morphology (Figure 3). Many of these landforms have  
16 been previously described as tunnel valleys by numerous authors (e.g. Wingfield,  
17 1990; Praeg, 2003; Lonergan et al., 2006). Although the exact mechanism of tunnel  
18 valley formation is still debated, there is general agreement that these major erosional  
19 features are formed by subglacial meltwater flowing more-or-less parallel to former  
20 ice flow (Ó Cofaigh, 1996). There is also evidence to suggest that the features in the  
21 central North Sea Basin were repeatedly occupied (Lonergan et al., 2006).

22 On the basis of the geomorphology and distribution of the features, described above,  
23 we have grouped the mapped features into three main assemblages. Assemblage 1  
24 includes the large, broadly curvilinear moraines on the shelf edge (Group 1), and the  
25 tunnel valleys (Group A) in the northern North Sea Basin that trend broadly  
26 perpendicular to these moraines. Assemblage 2 comprises the strongly arcuate, lobate,  
27 and often convolute, Group 2 moraines found on the mid-shelf. Assemblage 3  
28 includes the Group 3 moraines and Group B channels on the inner shelf, principally  
29 east and north of the Scotland. We now interpret each of these 3 assemblages in turn,  
30 with reference to the last British and Fennoscandian ice sheets.

31

1 *Assemblage 1:* The major moraines on the northwestern UK continental shelf margin  
2 represent substantial volumes of Pleistocene sediment relating to former, coherent,  
3 shelf-wide ice-sheet glaciation. These moraines occur in direct association with major  
4 shelf-edge fans such as the Sula Sgeir Fan, the Rona and Foula Wedges and the Barra-  
5 Donegal Fan – themselves representing large volumes of sediment deposited over a  
6 prolonged period (Stoker et al., 1993) (Figures 1, 3). We believe that this coupling of  
7 shelf-edge moraines and trough-mouth fans is best interpreted as reflecting the  
8 position of a formerly extensive continental ice sheet with overall flow towards the  
9 Atlantic Ocean in the west and northwest of the study area. This geometry is  
10 consistent with orientations of tunnel valleys and buried mega-scale glacial lineations  
11 (MSGs), identified in the North Sea Basin by Graham et al. (2007). When used as  
12 the basis for a glaciological inversion model, these three components – moraines,  
13 tunnel valleys, and MSGs – strongly suggest, at maximum stage, an ice-sheet  
14 surface across the northern North Sea Basin declining in altitude from southeast to  
15 northwest. By implication, the former BIS and FIS must have been confluent at this  
16 time.

17 *Assemblage 2:* The mid-shelf moraines on the West Shetland Shelf and those NE of  
18 Shetland must have formed after the extensive shelf-edge glaciation described above,  
19 most probably during a significant retreat stage. By this time the ice sheet had receded  
20 and thinned sufficiently to allow topography to influence ice-flow somewhat. It is  
21 notable that there are very few moraines of this stage on the low-lying shelf northwest  
22 of the Witch Ground Basin. The mid-shelf moraines to the west of Orkney and  
23 Shetland are lobate and highly convolute in morphology; many are strongly  
24 reminiscent of push moraines and thrust-block complexes formed at the margins of  
25 modern surging glaciers (Figure 7) (e.g. Clayton et al., 1985; Evans et al., 1999;  
26 Evans and Rea, 2003). Although their exact genesis cannot be determined at this point  
27 in time, the large number and density of these features shows that dynamically  
28 oscillating lobes were common within the last BIS. Importantly, the lobate margins  
29 defined by these moraines show that the ice-sheet terminus was grounded, highly  
30 irregular and quite different to the form of the terminus represented by the older,  
31 curvilinear, shelf-edge moraines (Figures 2 & 6). The lobate, bifurcating, overprinting  
32 nature of these younger moraines indicates that ice-marginal oscillations were a  
33 feature of mid-shelf deglaciation. This assertion is confirmed by seismic profiles

1 showing considerable readvances during overall recession within the moraine  
2 sequence NW of Orkney (Figure 6) (Stoker and Holmes, 1991). The convolute, often  
3 zigzag, form of these moraines is intriguing and may reflect marginal flow variations  
4 or the outline of highly crevassed palaeo-ice margins. Significantly, the vast majority  
5 of the moraines in Assemblage 2 are concave towards the UK landmass, even east of  
6 Shetland, suggesting that, by this stage, the influence of Scandinavian ice on the  
7 overall geometry of the BIS was considerably diminished. At this time an ice sheet  
8 appears to have occupied the ground north and west of the Witch Ground Basin and a  
9 separate ice sheet still occupied the Norwegian Channel and its western margin.  
10 However, the lowest part of the northern North Sea Basin, between the retreating BIS  
11 and FIS, was probably an ice-free marine embayment.

12 *Assemblage 3:* The third assemblage of landforms include generally smaller-scale  
13 near-shore moraines and meltwater channels that vary in alignment in different areas  
14 – compare the moraines and channels adjacent to the Moray Firth and Strathmore  
15 coasts, for example (Figures 1 & 3). The largest features – the Bosies Bank and Wee  
16 Bankie moraine complexes – are perhaps the earliest moraines of this stage, marking  
17 the transition into a more stable phase of deglaciation from the dynamically  
18 oscillating behaviour immediately preceding it. During this period the Witch Ground  
19 Basin was ice free and is known to have begun accumulating glaciomarine muds (c.  
20 22-19 ka <sup>14</sup>C BP) (Gatliff et al., 1994; Sejrup et al., 1994; 2004; Graham et al., 2007).  
21 On the basis of highly elongate landforms in NW Scotland and Strathmore, on land  
22 and on the seabed, and the deep channels at the mouth of the Moray Firth, it is likely  
23 that ice streams were still active within the BIS at this time (Merritt et al., 1995;  
24 Stoker and Bradwell, 2005; Golledge and Stoker, 2006). These fast-flowing outlets  
25 may have been prone to periods of flow instability – as highlighted by Merritt et al.  
26 (1995) in the Moray Firth region. Ice streaming was probably highly effective at  
27 discharging mass from the centres of ice sheet accumulation and therefore strongly  
28 conditioned recession of the BIS during this time.

29 To summarise, we interpret the sea-bed landforms on the UK continental shelf to  
30 show a complex pattern of ice-sheet deglaciation. Extensive shelf-edge glaciation  
31 occurred first, at LGM, characterised by confluence of the British and Fennoscandian  
32 Ice Sheets. This coalescent ice sheet flowed broadly northwestwards across the

1 northern North Sea transporting vast volumes of sediment to the continental slope,  
2 forming prominent moraines and shelf-edge fans. Subsequent oscillatory retreat of  
3 both ice sheets led to separation and the re-establishment of ice-free marine conditions  
4 in the North Sea Basin. At this time, the BIS margin was probably highly irregular –  
5 dominated by numerous, large, dynamic lobes. Final-stage deglaciation of the  
6 independent BIS was marked by near-shore ice streaming and increasing topographic  
7 control on ice-flow directions.

8

## 9 **Discussion**

10 Our results show that the glacial geomorphology on the sea bed around the northern  
11 United Kingdom preserves the footprint of a retreating ice sheet that significantly re-  
12 organised during its decay. Whilst each of the three stages identified in this study have  
13 been broadly suggested by previous workers (e.g. Boulton et al., 1977; Sejrup et al.,  
14 1994, 2005, Carr et al., 2006; Graham, 2007), no investigations to date have explicitly  
15 attempted to unravel the complexities inherent in such an ice-sheet retreat pattern.  
16 Specifically, no-one has yet identified the zone of confluence at ice-sheet maximum,  
17 or the zone of pull-apart where the British and Fennoscandian ice sheets separated, or  
18 the mechanism by which this happened. We believe that the Olex dataset provides the  
19 best empirical evidence for ice-sheet confluence and subsequent separation and,  
20 critically, enables a viable hypothesis of British Ice Sheet disintegration to be  
21 proposed for the first time.

22 The bathymetric data show a broad, weakly sinuous depression extending north from  
23 the Witch Ground Basin to the continental shelf margin east of Shetland (Figure 6).  
24 The relative absence of moraines in this area, together with the bifurcating pattern of  
25 retreat inferred from moraines to the east and west of this broad depression, suggest  
26 that this zone may have been the focus for ice-sheet separation. The present  
27 bathymetry indicates that the seafloor in this area would have been only 10-30 m  
28 below sea level during the last global sea-level minimum (at LGM). The conjoined  
29 BIS-FIS was probably sufficiently thick during early deglaciation to have been  
30 grounded in this area, despite some possible tidewater calving. However, as  
31 deglaciation proceeded, we propose that the interplay between sea-level and the ice-  
32 sheet margin along this corridor would have been crucial.



1 [FIGURE 9 HERE]

2 Calving is known to be responsible for the majority of mass loss in contemporary ice  
3 sheets (Reeh, 1968; Paterson, 1994), and so was probably equally important in palaeo-  
4 ice sheets such as the BIS-FIS. Floating ice is considerably more vulnerable to rapid  
5 collapse than grounded ice, since the highly viscous nature of ice means that stresses  
6 are transmitted throughout an ice shelf almost instantly. ‘Debuttressing’ occurs when  
7 ice shelves calve resulting in an acceleration of grounded ice flow and an increase in  
8 ice supply to the calving front, consequently leading to further mass loss (e.g. Mercer,  
9 1978; Paterson, 1994; Alley et al., 2005).

10 In light of the above, it is evident that if calving were to initiate along a sector of the  
11 margin of the former BIS-FIS, significant mass might be drawn-down from the  
12 interior of the ice sheet in the North Sea Basin. Overall surface lowering would be  
13 greatest nearer to the calving front, with lesser effects in grounded ice away from the  
14 margins. Initial calving may have been triggered by glacio-isostatic depression that  
15 produced rising relative sea levels in the period after the maximal ice volume occurred  
16 (cf. P.U. Clark et al., 2004). We suggest that, an initially small calving sector of the  
17 confluent BIS-FIS margin propagated southward, governed by the greater water  
18 depths east of Shetland and in the Witch Ground Basin, leading to the development of  
19 a large calving bay in the northern North Sea Basin (Figure 9). Rapid recession forced  
20 both grounded ice masses to re-organise, as surface lowering close to the calving bay  
21 led to a change in overall surface slope of the BIS. Changing flow directions as a  
22 result of this ice sheet re-organisation probably led to the formation of well-developed  
23 moraines north and east of Shetland. That this period of re-organisation is preserved  
24 in the geomorphic record as large, lobate, readvance moraines may suggest that the  
25 readjusting ice sheet was in a state of considerable disequilibrium. It was also during  
26 this episode that the De Geer moraines around Orkney and Shetland were formed –  
27 indicative of a tidewater margin. Final separation of the ice sheets occurred when sea  
28 levels had risen sufficiently to inundate the Witch Ground Basin, causing  
29 glaciomarine deposition. The subsequent marginal retreat of the British Ice Sheet may  
30 have been relatively rapid – dominated in eastern Scotland by ice stream activity in  
31 the Moray Firth, the Firth of Forth and Strathmore.

1 We have proposed a new model for the geometry of the confluent BIS-FIS at its  
2 maximum. We have also described the subsequent ice-sheet separation,  
3 reconfiguration and demise based on entirely new empirical evidence (Figures 4, 5).  
4 The validity of this hypothesis can be evaluated by comparison with other global  
5 accounts of ice sheet demise. The Laurentide Ice Sheet has been investigated  
6 extensively, and the repeated delivery of ice from its interior to its calving bays – via  
7 ‘Binge-Purge’ cycles – is now widely accepted (Heinrich, 1988; MacAyeal, 1993;  
8 Andrews and Maclean, 2003). Indeed, calving bays along the eastern Laurentide Ice  
9 Sheet margin are thought to have been instrumental in its eventual disintegration  
10 (Hughes et al., 1977; Thomas, 1977; Hughes, 2002; Shaw et al., 2006). Similar  
11 mechanisms have been proposed for the Fennoscandian Ice Sheet (Hoppe, 1948) and  
12 for the Barents Sea Ice Sheet (Siegert et al., 2002). Modern analogues also exist: the  
13 Larsen B Ice Shelf in West Antarctica collapsed rapidly in 2002 leading to the  
14 development of a large calving bay and changing the flow dynamics of the  
15 surrounding glaciers (Rack and Rott, 2004; Scambos et al., 2004). The fast-flowing  
16 Pine Island and Thwaites Glaciers on the Amundsen Coast of West Antarctica are  
17 currently undergoing rapid thinning and increased draw-down as a result of recent  
18 sea-level rise (Payne et al., 2004; Thomas et al., 2004). It remains uncertain how the  
19 flow dynamics of the West Antarctic Ice Sheet will be affected by these rapid  
20 changes. Modern glaciers in Greenland are experiencing accelerated calving rates  
21 associated with enhanced melting, rapid flow and thinning (Zwally et al., 2002). In  
22 Alaska, the Columbia Glacier is also known to have experienced a recent abrupt  
23 increase in calving rate as a result of surface lowering (van der Veen, 1996). Hence,  
24 we suggest, it would be unusual if disintegration of the marine-terminating sectors of  
25 the BIS had **not** involved substantial calving and subsequent draw-down. Studies in  
26 Ireland lend further plausibility to this suggestion. McCabe and co-workers postulate  
27 that Irish Sea deglaciation was triggered by an isostatically induced rise in relative sea  
28 level, perhaps occurring in as little as 500 years (McCabe and Clark, 1998; P.U. Clark  
29 et al., 2004; McCabe et al., 2005); whilst in the Irish Midlands, Delaney (2002)  
30 describes scenarios in which flooding of a central area of low ground was  
31 instrumental in the re-organisation of the Irish Ice Sheet. We propose that the calving-  
32 bay mechanism provides a plausible explanation for the geomorphic evidence present  
33 in the study area described here. Furthermore, that calving and unzipping was  
34 principally focused along a north-south axis east of Shetland extending as far south as

1 the Witch Ground Basin (Figure 9). This new insight provides a radically revised  
2 framework for BIS research, and a testable hypothesis for future modelling  
3 experiments.

#### 6 **Synthesis : combining the evidence**

7 Seismostratigraphic evidence from the North Sea shows that a grounded ice sheet  
8 flowed approximately northwest across the Witch Ground Basin (Graham et al.,  
9 2007). During this extensive period of ice-sheet glaciation, ice from Scandinavia  
10 coalesced with the BIS in the North Sea Basin (Sejrup et al., 2005). The presence of  
11 buried MSGLs in the Witch Ground Basin, dated stratigraphically to 22-28 ka <sup>14</sup>C BP,  
12 have been used to infer the presence of fast ice-sheet flow in this area. However, no  
13 surface expression of fast flow exists in this area today (Figure 4).

15 We present a reconstruction of the conjoined BIS-FIS during the LGM, around 25-30  
16 ka BP (Figure 10). During periods of maximum glaciation, when the FIS overspilled  
17 the Norwegian Channel, we suggest that Fennoscandian ice flowed west into the  
18 North Sea Basin and coalesced with ice from the eastern margins of the BIS. This  
19 coalescent, convergent flow resulted in a fast-flow corridor across deformable  
20 sediment in the lowest part of the basin (CZ, Figure 10). The presence of NW-  
21 orientated convergent MSGLs in the Witch Ground Basin (Graham et al., 2007); a  
22 widespread deformable bed; and the presence of large shelf-edge fans (Rona and  
23 Foula Wedges); are all consistent with a palaeo-ice stream flowing NW across the  
24 northern North Sea Basin. The reconstructed coalescent FIS-BIS is reinforced by its  
25 glaciological plausibility (Figure 8); with convergent flow focused in the low-lying  
26 corridor between Orkney and Shetland. The presence of strongly streamlined bedrock  
27 forms on Orkney, Shetland and Fair Isle indicate the former passage of a powerful ice  
28 sheet in a northwesterly direction (Peach and Horne, 1879, 1880; Hoppe, 1974;  
29 Mykura, 1976; Golledge et al., 2008). The presence of rare Norwegian erratics is also  
30 consistent with a far-travelled ice stream crossing these islands during the last  
31 glaciation (Finlay, 1926; Mykura, 1976; Birnie et al., 1993).

32  
33 [FIGURE 10 HERE]

1 Although this scenario has been previously challenged by numerous authors, we think  
2 opposing views can be reconciled when new geomorphological dataset from the  
3 whole continental shelf is considered. Examination of the wider picture during this  
4 glacial maximum event shows a concordant ice-flow pattern across much of northern  
5 and eastern Scotland. Glacial bedforms and striae in Strathmore and the inner Moray  
6 Firth suggest ice streams flowing in a northeasterly direction before being deflected  
7 northwest (Figure 10). The deflection of the Moray Firth ice stream northwest across  
8 Caithness has often been cited as evidence that the FIS occupied the northern North  
9 Sea Basin at this time (Peach and Horne, 1879; Hall and Bent, 1990; Carr et al.,  
10 2006). Far-travelled erratics, glacial striae and elongate bedforms in Orkney  
11 indicating WNW flow are all best explained by our revised BIS reconstruction – with  
12 ice flowing NE from the northeast Scottish mainland, before converging and being  
13 deflected to the NW by the more powerful FIS. At this time the zone of confluence  
14 between the BIS and the much larger FIS probably lay across the northern Orkney  
15 Islands (Figure 10).

16

17 The timing of confluent shelf-edge BIS-FIS glaciation is not well constrained.  
18 However, it is likely to have occurred after Heinrich Event 3 and before Heinrich  
19 Event 2 (i.e. between 32-24 ka BP). Seismostratigraphic evidence from the West  
20 Shetland Shelf shows that the moraines within the Otter Bank Formation formed  
21 during the Late Weichselian (Stoker et al., 1993). Sediment sequences on the Rona  
22 and Foula Wedges also demonstrate ice-proximal deposition at the shelf edge during  
23 the Late Weichselian (MIS 2) (Stoker et al., 1993, 1994). This evidence, combined  
24 with other moraines mapped in this study, indicate that the last BIS-FIS reached the  
25 continental-shelf edge west of Orkney and Shetland at LGM. Previously published  
26 radiocarbon dates from marine fauna place the moraines south of St Kilda, associated  
27 with the Barra-Donegal Fan, within the same period (i.e. MIS 2) (Selby, 1989;  
28 Peacock et al., 1992). The Barra Fan largely comprises sediment sourced in SW  
29 Scotland and NW Ireland and shows a major increase in ice-rafted debris c. 27 ka BP  
30 (Kroon et al., 2000; Knutz et al., 2001; Peck et al., 2007). NW of Lewis, however,  
31 moraines at the shelf edge adjacent to the Sula Sgeir Fan, are overlain by deposits of  
32 probable MIS 4 age (Stoker, 1995). However, given the wider evidence for shelf-edge  
33 MIS 2 glaciation to the north and south, it is likely that this Amino-acid age-  
34 assessment is in need of revision. Other ice-sheet flow lines relating to the BIS at

1 maximum stage are inferred based on bathymetry, seafloor geomorphology and  
2 unpublished BGS mapping. Some of those relating to the FIS have been taken from  
3 previously published work (Sejrup et al., 1994; Ottesen et al., 2005). Crucially, we  
4 suggest that our new ice-sheet reconstruction is the most likely scenario for the BIS at  
5 LGM (Figure 10), and that this scenario is entirely consistent with most of the  
6 previously published evidence.

7  
8 Following extensive shelf-edge glaciation, at LGM, the BIS underwent a period of  
9 large-scale re-organization (Figure 11). We propose that this ice-mass re-organization  
10 was largely the result of rapidly rising sea levels (Figure 9), probably owing to glacio-  
11 isostatic loading, causing major changes in ice-sheet configuration and flow  
12 dynamics. It is possible that this re-organisation was closely associated with the North  
13 Atlantic iceberg-discharge event – Heinrich Event 2 (c. 24 ka BP; cf. Bond et al.,  
14 1993; Peck et al., 2007). However, current dating constraints do not permit more  
15 certainty on this linkage. During this stage, both ice sheets underwent rapid recession,  
16 we suggest largely as a result of calving into a marine embayment. As this embayment  
17 opened, along a north-south axis east of Shetland, both ice-sheet margins receded c.  
18 100 km back towards higher ground, probably over a relatively short period of time  
19 (Figure 11). Sea water had inundated the Witch Ground Basin by ~20 ka <sup>14</sup>C BP (~24  
20 ka cal BP) (Sejrup et al., 1994; Graham et al., 2007). As the BIS re-organized, ice  
21 divides migrated rapidly. Large ice-sheet-lobe moraines on the mid-shelf west of  
22 Orkney suggest that this northwestern margin was dynamically unstable during  
23 overall ice-sheet re-organization. The innermost moraine loop off Orkney has sharp  
24 fresh-looking morphology and is clearly the youngest ice-sheet moraine in this suite.  
25 However, the absence of similar lobate moraines on the seafloor east of Orkney  
26 suggest that these islands did not host an independent active ice cap during this time,  
27 and that the dynamic behaviour was probably a short-lived consequence of ice-sheet  
28 readjustment. By contrast, large lobate moraines north, east and west of Shetland  
29 suggest that these islands maintained an independent, dynamic ice cap during this  
30 deglacial stage (Figure 11) – a model in line with many previous findings (e.g. Hoppe,  
31 1974; Mykura, 1976; Long et al., 2004; Carr et al., 2006; Golledge et al., 2008). The  
32 large size and linear morphology of the Bosies Bank and Wee Bankie moraine  
33 complexes suggest a considerable period of ice-sheet stability. We suggest that these  
34 moraines, previously attributed to the ‘Last Glacial Maximum’ (e.g. Sutherland, 1984;

1 Bowen et al., 2002), represent a readvance of the BIS in the aftermath of major ice-  
2 sheet re-organization (Figure 11). This latter episode probably occurred after Heinrich  
3 Event 2 and before Heinrich Event 1, between c.18-24 ka BP.

4  
5 [FIGURE 11 HERE]

6  
7 In summary, we have reconstructed a new model for the glaciation and subsequent  
8 deglaciation of the northern UK and the surrounding continental shelf (Figures 10,  
9 11). At LGM the reconstructed BIS was coalescent with the FIS in the northern North  
10 Sea and was drained by several ice streams – the most dominant of these, fed from  
11 western Norway and eastern Scotland, flowed NW across the Witch Ground Basin to  
12 the continental-shelf edge west of Shetland. We propose that following relative sea-  
13 level rise, the BIS-FIS broke apart and re-organized into two independent ice sheets  
14 which may have remained unstable for some time as they adjusted to new boundary  
15 conditions (Figure 11). Widespread, dynamic, ice-margin oscillations probably  
16 occurred in response to this re-organisation. Final-stage deglaciation was marked by  
17 ice-sheet thinning and increasing topographic control on ice-flow dynamics in the  
18 BIS. This stage was also typified by near-shore ice streaming and numerous ice-  
19 marginal oscillations in east and west Scotland.

20  
21 We believe that this 3-stage model explains the long-standing apparent contradiction  
22 between those who have argued for ice-free conditions in the North Sea Basin at  
23 LGM and those who have argued against. Crucially, our new interpretation, based on  
24 shelf-wide sea-bed imagery (Olex), suggests that the period of ice-sheet separation  
25 was one of dynamic instability – ultimately leading to large-scale ice-sheet collapse in  
26 the North Sea Basin.

## 27 28 29 **Conclusions**

30  
31 Access to the entire Olex echosounder dataset has revealed the geomorphology of the  
32 sea floor around the northern UK in unprecedented detail. From this, and the wider  
33 evidence, we have drawn the following conclusions regarding the British Ice Sheet at  
34 Last Glacial Maximum:

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

1. A large number of previously unidentified geomorphic features relating to the last ice sheet occur on the continental shelf around the northern UK.
2. The BIS was coalescent with the FIS during a relatively recent phase of extensive ice-sheet glaciation. NW-oriented MSGLs of Late Devensian/Late Weichselian age and NW-oriented tunnel valleys in the northern North Sea, coupled with large shelf-edge moraines west of Shetland, north and west of Lewis and south of St Kilda, dated to MIS 2, all indicate an extensive glaciation between c. 30-25 ka BP. This was also a time when large shelf-edge fans received a major increase in glacially derived sedimentation. This interpretation is consistent with previously published geological evidence from the western sector of the FIS. We propose that at LGM the zone of ice-sheet confluence probably stretched from 1°deg E longitude, NW across the northern Orkney Islands, to the shelf edge west of Shetland. Fast-flowing confluent ice in the Fair Isle Channel during this time would have focused sediment delivery to the Rona and Foula Wedges.
3. Ice-sheet retreat occurred primarily by calving as a result of rising sea levels. We suggest that, during abrupt relative sea-level rise around the time of Heinrich-2 (c. 24 ka BP), a large marine embayment opened in the northern North Sea, as far south as the Witch Ground Basin. This marine embayment changed the entire configuration of the two ice sheets forcing them to decouple rapidly along a north-south axis east of Shetland. Similarities are striking between this scenario and the ongoing, rapid, break-up of marine-terminating parts of the West Antarctic Ice Sheet.
4. The northern sector of the last BIS underwent a remarkable period of ice-sheet re-organisation in response to the opening of a large marine embayment centred on the Witch Ground Basin. Dynamic ice-front oscillations – deduced from overprinted lobate moraine patterns and seismic stratigraphy – occurred during this period as the BIS rapidly adjusted to new quasi-stable margin positions.
5. A period of relative stability followed, when the BIS margin was situated in the vicinity of the Bosies Bank–Wee Bankie moraine complex off eastern Scotland. This ice-sheet configuration probably equates to the traditional

1 'LGM' c. 18-24 ka BP – with an ice-free North Sea Basin and a substantial ice  
2 cap on Shetland.

3 6. Ice-sheet thinning and punctuated retreat continued, increasing topographic  
4 control and invigorating several near-shore ice streams. Around 16 ka BP,  
5 following Heinrich Event 1, the BIS margin was stably situated at the present-  
6 day coastline in NW Scotland. Subsequent ice-sheet retreat was probably  
7 slower and took place primarily by melting, rather than calving.

8 7. We suggest that this new 3-stage reconstruction of the northern half of the last  
9 BIS offers a glaciologically plausible model which reconciles much of the  
10 previously published, apparently contradictory, evidence from the wider area.

11

12

### 13 **Acknowledgements**

14 We thank Clive Auton and Chris Thomas for their comments on an earlier version of  
15 this manuscript. The comments of Andrew Miall and three anonymous reviewers  
16 helped to clarify aspects of this work. Bob Gatliff and Martin Smith are thanked for  
17 their continuing supporting of this project. Published with the permission of the  
18 executive Director, BGS (NERC).

19

20

### 21 **References**

22

- 23 Alley, R. B., Clark, P.U., Huybrechts, P., Joughin, I., 2005. Ice-sheet and sea-level  
24 changes. *Science* 310, 456–460.
- 25 Andrews, I.J., Long, D., Richards, P.C., Thomson, A.R., Brown, S., Chesher, J.A.,  
26 McCormac, M., 1990. United Kingdom offshore regional report: The Geology  
27 of the Moray Firth. HMSO for the British Geological Survey. London, 93pp.
- 28 Andrews, J.T., Maclean, B., 2003. Hudson Strait ice streams: a review of stratigraphy,  
29 chronology and links with North Atlantic Heinrich events. *Boreas* 32, 4-17.
- 30 Armstrong, M., Paterson, I.B., Browne, M.A.E., 1985. Geology of the Perth and  
31 Dundee District. 1:50,000 geological sheets 48W, 48E, 49. *Memoirs of the*  
32 *British Geological Survey (Scotland)*, London, HMSO, 108pp.
- 33 Austin, W.E.N., Kroon, D., 1996. Late glacial sedimentology, foraminifera and stable  
34 isotope stratigraphy of the Hebridean Continental Shelf, northwest Scotland. .  
35 In: J.T. Andrews, Austin, W.E.N., Bergsten, H. and Jennings, A.E. (Editor),  
36 Late Quaternary Palaeoceanography of the North Atlantic Margins. *Geological*  
37 *Society, London*, pp187-213.
- 38 Auton, C.A., 2003. The Quaternary and Devonian geology of Sheet 115 (Reay).  
39 *British Geological Survey Commissioned Report, CR/03/030*, pp68.



- 1 Ballantyne, C.K., McCarroll, D., Nesje, A., Dahl, S.O., Stone, J.O., Fifield, L.K.,  
2 1998. High-resolution reconstruction of the last ice sheet in NW Scotland  
3 *Terra Nova* 10, 63–67.
- 4 Benn, D.I., Evans, D.J.A., 1998. *Glaciers and Glaciation*. London: Arnold, 734pp.
- 5 Birnie, J., Gordon, J.E., Bennett, K., Hall, A.M., 1993. *The Quaternary of Shetland:*  
6 *Field Guide*. Cambridge: Quaternary Research Association.
- 7 Blake, K. P., 2000. Common origin for De Geer moraines of variable composition in  
8 Raudvassdalen, northern Norway. *Journal of Quaternary Science* 15, 633–644.
- 9 Bond, G., Broecker, W., Johnsen, S., McManus, J., Labeyrie, L., Jourzel, J., Bonani,  
10 G., 1993. Correlations between climate records from North Atlantic sediments  
11 and Greenland ice. *Nature*, 365, 143-147.
- 12 Bos J., A.A, Dickson, J.H., Coope, G.R., Jardine, W.G., 2004. Flora, fauna and  
13 climate of Scotland during the Weichselian Middle Pleniglacial—  
14 Palynological, macrofossil and coleopteran investigations. *Palaeogeography,*  
15 *Palaeoclimatology, Palaeoecology* 204, 65–100.
- 16 Boulton, G.S., Hagdorn, M., 2006. Glaciology of the British Isles during the last  
17 glacial cycle: form, flow, streams and lobes. *Quaternary Science Reviews* 25,  
18 3359-3390.
- 19 Boulton, G.S., Jones, A.S., Clayton, K.M., Kenning, M.J., 1977. A British ice sheet  
20 model and patterns of glacial erosion and deposition. In: Shotton, F.W. (Ed.),  
21 *British Quaternary Studies: Recent Advances*. Oxford University Press,  
22 Oxford, pp231–246.
- 23 Boulton, G. S., Smith, G. D., Jones, A. S., Newsome, J., 1985. Glacial geology and  
24 glaciology of the last mid-latitude ice sheets. *Journal of the Geological*  
25 *Society*, London 142, 447–474.
- 26 Boulton, G. S., Peacock, J. D., Sutherland, D. G., 1991. Quaternary. In: Craig, G. Y.  
27 (Ed.), *Geology of Scotland*, 3<sup>rd</sup> edition. The Geological Society, London,  
28 pp503-543.
- 29 Boulton, G.S., Peacock, J.D., Sutherland, D.G., 2002. Quaternary. In: Trewin, N.H.  
30 (Ed.) *The Geology of Scotland*, 4<sup>th</sup> edition. The Geological Society, London,  
31 pp409-431.
- 32 Bowen, D.Q. 1989. The last interglacial-glacial cycle in the British Isles. *Quaternary*  
33 *International*, 3-4, 41-7.
- 34 Bowen, D.Q. 1999. A revised correlation of Quaternary deposits in the British Isles.  
35 *Geological Survey Special Report*, 23.
- 36 Bowen, D.Q., McCabe, A.M., Rose, J., Sutherland, D., 1986. Correlation of  
37 Quaternary Glaciations in England, Ireland, Scotland and Wales. In: Sibrava,  
38 B., Bowen, D.Q., Richmond, G.M. (Eds.), *Quaternary Glaciations in the*  
39 *Northern Hemisphere*. International Geological Correlation Programme  
40 Project No. 24, *Quaternary Science Reviews* 5, 199–340.
- 41 Bowen, D.Q., Phillips, F.M., McCabe, A.M., Knutz, P.C., Sykes, G.A., 2002. New  
42 data for the Last Glacial Maximum in Great Britain and Ireland. *Quaternary*  
43 *Science Reviews* 21, 89-101.
- 44 Bradwell, T., Stoker, M.S., Larter, R., 2007. Geomorphological signature and flow  
45 dynamics of The Minch palaeo-ice stream, NW Scotland. *Journal of*  
46 *Quaternary Science*, 22, 609-622.
- 47 Brown., E.J., Rose, J., Coope, R. Lowe, J.J., 2007. An MIS 3 age organic deposit  
48 from Balglass Burn, central Scotland: palaeoenvironmental significance and  
49 implications for the timing of the onset of the LGM ice sheet in the vicinity of  
50 the British Isles. *Journal of Quaternary Science* 22, 295-308.

- 1 Bulat, J., Long, D., 2001. Images of the seabed in the Faroe-Shetland Channel from  
2 commercial 3D seismic data. *Marine Geophysical Researches* 22, 345-367.
- 3 Cameron, T.D.J, Stoker, M.S., Long, D., 1987. The history of Quaternary  
4 sedimentation in the UK sector of the North Sea basin. *Journal of the*  
5 *Geological Society, London* 144, 43-58.
- 6 Carr, S.J., Holmes, R., van der Meer, J. J. M., Rose, J., 2006. The Last Glacial  
7 Maximum in the North Sea Basin: micromorphological evidence of extensive  
8 glaciation. *Journal of Quaternary Science* 21, 131-153
- 9 Clapperton, C.M., 1970. The pattern of deglaciation in part of North Northumberland.  
10 *Transactions of the Institute of British Geographers* 45, 67-78.
- 11 Clark, C.D., Evans, D.J.A., Khatwa, A., Bradwell, T., Jordan, C.J., Marsh, S.H.,  
12 Mitchell, W.A., Bateman, M.D., 2004. BRITICE: Map and GIS database of  
13 landforms and features related to the last British Ice Sheet. *Boreas* 33, 359-  
14 375.
- 15 Clark, P. U., McCabe, A. M., Mix, A.C., Weaver, A. J., 2004. Rapid rise of sea level  
16 19,000 years ago and its global implications. *Science* 304, 1141–1144.
- 17 Clayton, L., Teller, J.T., Attig, J.W., 1985. Surging of the southwestern part of the  
18 Laurentide Ice Sheet. *Boreas* 14, 235-241.
- 19 Colgan, P.M., Mickelson, D.M., Cutler, P.M., 2003. Ice-marginal terrestrial  
20 landsystems: southern Laurentide Ice Sheet margin. In: Evans, D.J.A. (Ed.),  
21 *Glacial Landsystems*. London: Arnold, pp111-142.
- 22 Cullingford, R.A., Smith, D.E., 1980. Late Devensian raised shorelines in Angus and  
23 Kincardineshire, Scotland. *Boreas* 9, 21-38.
- 24 Davison, S., 2005. Reconstructing the Last Pleistocene (Late Devensian) Glaciation  
25 on the Continental Margin of Northwest Britain. Unpublished PhD thesis.  
26 University of Edinburgh.
- 27 Davison, S., Stoker, M.S., 2002. Late Pleistocene glacially-influenced deep-marine  
28 sedimentation off NW Britain: implications for the rock record. *Special*  
29 *Publication Geological Society of London* 203, 129-148.
- 30 Delaney, C., 2002. Sedimentology of a glaciofluvial landsystem, Lough Ree area,  
31 Central Ireland: implications for ice margin characteristics during Devensian  
32 deglaciation. *Sedimentary Geology* 149, 111–126.
- 33 Ehlers, J., Wingfield, R., 1991. The extension of the Late Weichselian/Late Devensian  
34 ice sheets in the North Sea Basin. *Journal of Quaternary Science* 6, 313-326.
- 35 Evans, D.J.A., 2003. *Glacial Landsystems*. Hodder Arnold, London, 532 pp.
- 36 Evans, D.J.A., Rea, B.R., 2003. Surging glacier landsystem. In: Evans, D.J.A. *Glacial*  
37 *Landsystems*. Hodder Arnold, London, pp259-289.
- 38 Evans, D.J.A., Lemmen, D.S., Rea, B.R., 1999. Glacial landsystems of the southwest  
39 Laurentide ice sheet: Modern Icelandic analogues. *Journal of Quaternary*  
40 *Science* 14, 673-679.
- 41 Evans, D.J.A. Clark, C.D., Mitchell, W.A., 2005. The last British Ice Sheet: A review  
42 of the evidence utilised in the compilation of the Glacial Map of Britain. *Earth*  
43 *Science Reviews* 70, 252-312.
- 44 Everest, J. D., Bradwell, T., Gollidge, N.R., 2005. Subglacial bedforms of the Tweed  
45 palaeo-ice stream. *Scottish Geographical Journal* 121, 163-173.
- 46 Everest, J.D., Bradwell, T., Fogwill, C.J., Kubik, P.W., 2006. Cosmogenic <sup>10</sup>Be age  
47 constraints from the Wester Ross Readvance Moraine: insights into British Ice  
48 Sheet behaviour. *Geografiska Annaler* 88A, 9-17.

- 1 Fabel, D., Stroeven, A.P., Harbor, J., Kleman, J., Elmore, D., Fink, D., 2002.  
2 Landscape preservation under ice sheets. *Earth and Planetary Science Letters*  
3 201, 397-406.
- 4 Fairbanks, R.G., Richard J Mortlock, R.A., Chiu, T-C., Cao, L., Kaplan, A.,  
5 Guilderson, T.P., Fairbanks, T.W., Bloom, A.L., 2005. Marine radiocarbon  
6 calibration curve spanning 0 to 50,000 years B.P. based on paired  
7  $^{230}\text{Th}/^{234}\text{U}/^{238}\text{U}$  and  $^{14}\text{C}$  dates on pristine corals. *Quaternary Science Reviews*  
8 24, 1781-1796.
- 9 Finlay, T.M., 1926. A tongsbergite boulder from the boulder clay of Shetland.  
10 *Transactions of the Edinburgh Geological Society* 12, 180.
- 11 Fitch, S., Thomson, K., Gaffney, V., 2005. Late Pleistocene and Holocene  
12 depositional systems and the palaeogeography of the Dogger Bank, North Sea.  
13 *Quaternary Research* 64,185-196.
- 14 Flinn, D., 1978. The most recent glaciation of the Orkney-Shetland Channel and  
15 adjacent regions. *Scottish Journal of Geology* 14, 109-123.
- 16 Gatliff, R.W., Richards, P.C., Smith, K., Graham, C.C., McCormac, M., Smith,  
17 N.J.P., Jeffery, D., Long, D., Cameron, T.D.J., Evans, D., Stevenson, A.G.,  
18 Bulat, J., Ritchie, J.D., 1994. British Geological Survey: United Kingdom  
19 offshore regional report: the geology of the central North Sea, London:  
20 HMSO.
- 21 Golledge, N.R., Stoker, M., 2006. A palaeo-ice stream of the British Ice Sheet in  
22 eastern Scotland *Boreas* 35, 231-243.
- 23 Golledge, N. R., Finlayson, A., Bradwell, T., Everest, J. D. 2008. The last glaciation  
24 of Shetland, North Atlantic. *Geografiska Annaler* 90A, tbc.
- 25 Gordon, J. E., Sutherland, D. G., 1993. *Quaternary of Scotland*. Chapman and Hall,  
26 London, 695pp.
- 27 Graham, A.G.C., 2007. Reconstructing Pleistocene glacial environments in the central  
28 North Sea using 3D seismic and borehole data. Unpublished PhD thesis.  
29 Imperial College London.
- 30 Graham, A. G. C., Lonergan, L., Stoker, M. S., 2007. Evidence for Late Pleistocene  
31 ice stream activity in the Witch Ground Basin, central North Sea, from 3D  
32 seismic reflection data. *Quaternary Science Reviews* 26, 627-643.
- 33 Graham, D.K., Harland, R., Gregory, D.M., Long, D., Morton, A.C., 1990. The  
34 biostratigraphy and chronostratigraphy of BGS Borehole 78/4, North Minch.  
35 *Scottish Journal of Geology*, 26, 65–75.
- 36 Hall, A. M., Bent, A. J. A., 1990. The limits of the last British ice sheet in northern  
37 Scotland and the adjacent shelf. *Quaternary Newsletter* 61, 2-12.
- 38 Hall, A., Glasser, N.F., 2003. Reconstructing the basal thermal regime of an ice  
39 stream in a landscape of selective linear erosion: Glen Avon, Cairngorm  
40 Mountains, Scotland. *Boreas* 32, 191-208.
- 41 Hall, A M., Jarvis, J., 1989. A preliminary report on the Late Devensian glaciomarine  
42 deposits around St Fergus, Grampian Region. *Quaternary Newsletter* 59, 5-7.
- 43 Hall, A M., Whittington, G., 1989. Late Devensian glaciation of southern Caithness.  
44 *Scottish Journal of Geology* 25, 307-324.
- 45 Hall, A.M., Peacock, J.D., Connell, E.R., 2003. New data for the Last Glacial  
46 Maximum in Great Britain and Ireland: a Scottish perspective on the paper by  
47 Bowen et al. (2002). *Quaternary Science Reviews* 22, 1551–1554.
- 48 Hedges, R.E.M., Millard, A.R., 1995. Bones and groundwater: towards the modelling  
49 of diagenetic processes. *Journal of Archaeological Science* 22, 155-64.

- 1 Heinrich, H., 1988. Origin and consequences of cyclic ice rafting in the northeast  
2 Atlantic Ocean during the past 130,000 years. *Quaternary Research* 29, 412-  
3 152.
- 4 Higham, T.F.G., Jacobi, R.M., Ramsey, C.B., 2006. AMS Radiocarbon dating of  
5 ancient bone using ultrafiltration. *Radiocarbon* 48, 179-195.
- 6 Holmes, R., 1977. Quaternary deposits of the central North Sea, 5. Quaternary  
7 geology of the UK sector of the North Sea between 56° and 58° North. Report  
8 of the Institute of Geological Sciences 77/14.
- 9 Holmes, R., 1997. Quaternary stratigraphy: the offshore record. In: J.E. Gordon  
10 (Editor), *Reflections on the Ice Age in Scotland*. Scottish Natural Heritage,  
11 Glasgow, pp72–94.
- 12 Holmes, R., Bulat, J., Hamilton, I., Long, D., 2003. Morphology of an ice-sheet limit  
13 and constructional glacially fed slope front, Faeroe-Shetland margin. In: J.W.  
14 Mienert, P. (Ed.), *European Margin Sediment Dynamics: Side-Scan Sonar and*  
15 *Seismic Images*. Springer-Verlag, Berlin, pp149–151.
- 16 Hoppe, G., 1948. Isrecessionen fran Norbottens Kustland i belysning av de glaciala  
17 formelementen. *Geographica Skriften fran Uppsala Universitets Geografiska*  
18 *Institutione*. 20, 112pp.
- 19 Hoppe, G., 1974. The glacial history of the Shetland Islands. Institute of British  
20 geographers Special Publication No. 7.
- 21 Hughes, T., 2002. Calving bays. *Quaternary Science Reviews* 21, 267–282.
- 22 Huuse, M., Lykke-Andersen, H., 2000. Overdeepened Quaternary valleys in the  
23 eastern Danish North Sea: morphology and origin. *Quaternary Science*  
24 *Reviews* 19, 1233-1253.
- 25 Jacobi, R.M., Rowe, P.J., Gilmour, M.A., Grün, R., Atkinson, T.C., 1998.  
26 Radiometric dating of the Middle Palaeolithic tool industry and associated  
27 fauna of Pin Hole Cave, Creswell Crags, England. *Journal of Quaternary*  
28 *Science* 13, 29-42.
- 29 Jardine, W.G., Dickson, J.H., Haughton, P.D.W., Harkness, D.D., Bowen D.Q.,  
30 Sykes, G.A., 1988. A late Middle Devensian interstadial site at Sourlie, near  
31 Irvine, Strathclyde. *Scottish Journal of Geology* 24, 288–295.
- 32 Johnson, H., Richards, P.C., Long, D., Graham, C.C., 1993. United Kingdom offshore  
33 regional report: the geology of the northern North Sea., HMSO, London.
- 34 Johnstone, G. S., Mykura, W., 1989. *The Northern Highlands of Scotland*, 4<sup>th</sup>  
35 Edition. HMSO for the British Geological Survey, Edinburgh.
- 36 Kleman, J., Glasser, N., 2007. The subglacial thermal organisation (STO) of ice  
37 sheets. *Quaternary Science Reviews*, 26: 587-597.
- 38 Kleman, J., Hättestrand, C., Borgstrom, I., Stroeven, A., 1997. Fennoscandian  
39 palaeoglaciology reconstructed using a glacial geological inversion model.  
40 *Journal of Glaciology* 43, 283–299.
- 41 Knutz, P.C., Austin, W.E.N., Jones, E.J.W., 2001. Millennial-scale depositional  
42 cycles related to British Ice Sheet variability and North Atlantic  
43 paleocirculation since 45 kyr B.P., Barra Fan, U.K. margin. *Paleoceanography*  
44 16, 53-64.
- 45 Knutz, P.K., Jones, E.J., Austin, W.E.N., van Weering, T.C.E., 2002. Glacimarine  
46 slope sedimentation, contourite drifts and bottom current pathways on the  
47 Barra Fan, UK North Atlantic margin. *Marine Geology* 188,129–146.
- 48 Kroon, D., Shimmield, G., Austin, W.E.N., Derrick, S., Knutz, P., Shimmield, T.,  
49 2000. Century- to millennial-scale sedimentological-geochemical records of

- 1 glacial–Holocene sediment variations from the Barra Fan (NE Atlantic).  
2 *Journal of the Geological Society, London* 157, 643–653.
- 3 Lambeck, K., 1991. Glacial rebound and sea level change in the British Isles. *Terra*  
4 *Nova* 3, 379–389.
- 5 Lambeck, K., 1993. Glacial rebound of the British Isles: 2. A high-resolution, high  
6 precision model. *Geophysical Journal International* 15, 960–990.
- 7 Lambeck, K., 1995. Late Devensian and Holocene shorelines of the British Isles and  
8 North Sea from models of glacio-hydro-isostatic rebound. *Journal of the*  
9 *Geological Society, London* 152, 437–448.
- 10 Larsen, E., Longva, O., Follestad, B.A., 1991. Formation of De Geer moraines and  
11 implications for deglaciation dynamics. *Journal of Quaternary Science* 6, 263–  
12 277.
- 13 Lawson, T. J., 1984. Reindeer in the Scottish Quaternary. *Quaternary Newsletter* 42,  
14 1–7.
- 15 Lehmann, S.J., Jones, G.A., Keigwin, L.D., Andersen, E.S., Butenkoi, G., Østmo,  
16 S.R., 1991. Initiation of Fennoscandian ice-sheet retreat during the last  
17 deglaciation. *Nature* 349, 513–516.
- 18 Lindén, M., Möller, P. 2005. Marginal formation of De Geer moraines and their  
19 implications to the dynamics of grounding-line recession. *Journal of*  
20 *Quaternary Science* 20, 113–133.
- 21 Lonergan, L., Maidment, S.C.R., Collier, J.S., 2006. Pleistocene subglacial tunnel  
22 valleys in the central North Sea basin: 3-D morphology and evolution. *Journal*  
23 *of Quaternary Science* 21, 891–903.
- 24 Long, D., Bent, A., Harland, R., Gregory, D.M., Graham, D.K., Morton, A.C., 1986.  
25 Late Quaternary palaeontology, sedimentology and geochemistry of a  
26 vibrocore from the Witch Ground Basin, central North Sea. *Marine Geology*  
27 73, 109–123.
- 28 Long, D., Bulat, J., Stoker, M.S., 2004. Sea bed morphology of the Faro-Shetland  
29 Channel derived from 3D seismic datasets. In: Davies, R.J., Cartwright, J.A.,  
30 Stewart, S.A., Lappin, M. and Underhill, J.R. (Eds.). *3D Seismic Technology:*  
31 *application to the exploration of sedimentary basins.* Geological Society of  
32 London Memoir 29, 53–61.
- 33 MacAyeal, D. R., 1993. Binge/purge oscillations of the Laurentide Ice Sheet as a  
34 cause of the North Atlantic's Heinrich events. *Paleoceanography* 8, 775–784.
- 35 McCabe, A.M., Clark, P.U., 1998. Ice-sheet variability around the North Atlantic  
36 Ocean during the last glaciation. *Nature* 392, 373–377.
- 37 McCabe, A. M., Knight, J., McCarron, S. 1998. Evidence for Heinrich event 1 in the  
38 British Isles. *Journal of Quaternary Science* 13, 549–568.
- 39 McCabe, A.M., Clark, P.U., Clark, J. 2005. AMS <sup>14</sup>C dating of deglacial events in the  
40 Irish Sea Basin and other sectors of the British–Irish ice sheet. *Quaternary*  
41 *Science Reviews* 24, 1673–1690.
- 42 McCabe, A.M., Clark, P.U., Smith, D.E., Dunlop, P. 2007. A revised model for the  
43 last deglaciation of eastern Scotland. *Journal of the Geological Society,*  
44 *London* 164, 313–316.
- 45 McCarroll, D. 2002. Amino-acid geochronology and the British Pleistocene: secure  
46 stratigraphical framework or a case of circular reasoning? *Journal of*  
47 *Quaternary Science* 17, 647–651.
- 48 Mercer, J.H., 1978. West Antarctic ice sheet and CO<sup>2</sup> greenhouse effect: a threat of  
49 disaster. *Nature* 271, 321–325.

- 1 Merritt, J.W., Auton, C.A., Firth, C.R., 1995. Ice-proximal glaciomarine  
2 sedimentation and sea-level change in the Inverness area, Scotland - A review  
3 of the deglaciation of a major ice stream of the British Late Devensian Ice-  
4 Sheet. *Quaternary Science Reviews* 14, 289-329.
- 5 Merritt, J.W., Auton, C.A., Connell, E.R., Hall, A.M., Peacock, J.D., 2003. Cainozoic  
6 geology and landscape evolution of north-east Scotland. *British Geological*  
7 *Survey, Edinburgh*, 178pp.
- 8 Mix, A.C., Bard, E., Schneider, R., 2001. Environmental Processes of the Ice Age:  
9 land, ocean, glaciers (EPILOG). *Quaternary Science Reviews* 20, 627–657.
- 10 Mykura, W. 1976. Orkney and Shetland. *British Regional Geology*. Institute  
11 of Geological Sciences/HMSO, Edinburgh, 149pp.
- 12 Murray, N.E., Bonsall, C., Sutherland, D.G., Lawson, T.J., Kitchener, A.C., 1993.  
13 Further radiocarbon determinations on reindeer remains of middle and late  
14 Devensian age from the Creag nan Uamh caves, Assynt, N.W. Scotland.  
15 *Quaternary Newsletter* 70, 1–10.
- 16 Mykura, W. 1976. Orkney and Shetland. *British Regional Geology*. Edinburgh,  
17 HMSO
- 18 Nesje A., Sejrup H. P., 1988. Late Weichselian/Devensian ice sheets in the North Sea  
19 and adjacent land areas. *Boreas* 17, 371-384.
- 20 Nielsen, T., Rasmussen, T.L., Ceramicola, S., Kuijpers, A., 2007. Quaternary  
21 sedimentation, margin architecture and ocean circulation variability around the  
22 Faroe Islands, North Atlantic. *Quaternary Science Reviews*, in press.  
23 doi:10.1016/j.quascirev.2006.12.005.
- 24 Nygard, A., Sejrup, H.P. Haflidason, H., Bryn, P., 2005. The glacial North Sea Fan,  
25 southern Norwegian Margin: Architecture and evolution from the upper  
26 continental slope to the deep-sea basin. *Marine and Petroleum Geology* 22, 71-  
27 84.
- 28 Ó Cofaigh, C., 1996. Tunnel valley genesis. *Progress in Physical Geography* 20, 1-19.
- 29 Ottesen, D., Dowdeswell, J.A., Rise, L., 2005. Submarine landforms and the  
30 reconstruction of fast-flowing ice streams within a large Quaternary ice sheet:  
31 The 2500-km-long Norwegian-Svalbard margin (57°-80°N). *Geological*  
32 *Society of America Bulletin* 117, 1033-1050.
- 33 Paterson, W.S.B., 1994. *The Physics of Glaciers*. 3<sup>rd</sup> edition. Oxford: Pergamon,  
34 480pp.
- 35 Payne, A. J., Vieli, A., Shepherd, A., Wingham, D. J., Rignot, E., 2004. Recent  
36 dramatic thinning of largest West-Antarctic ice stream triggered by oceans.  
37 *Geophysical Research Letters* 31, L23401, 1–4.
- 38 Peach, B.N., Horne, J., 1879. The glaciation of the Shetland Islands. *Quarterly Journal*  
39 *of the Geological Society of London* 35, 778-811.
- 40 Peach, B.N., Horne, J., 1880. The glaciation of the Orkney Islands. *Quarterly Journal*  
41 *of the Geological Society of London* 36, 648-63.
- 42 Peach, B.N., Horne, J., 1893. On the occurrence of shelly boulder clay in North  
43 Ronaldsay, Orkney. *Transactions of the Edinburgh Geological Society* 6, 309-  
44 13.
- 45 Peacock, J.D., 1985. Comments on the Quaternary deposits and landforms of Scotland  
46 and the neighbouring shelves: a review. *Quaternary Science Reviews* 4, 1-2.
- 47 Peacock, J.D., Merritt, J.W., 2000. Glacial deposits at the Boyne Bay Limestone  
48 Quarry, Portsoy, and their place in the late Pleistocene history of northeast  
49 Scotland. *Journal of Quaternary Science* 15, 543-555.

- 1 Peacock, J.D., Austin, W. E. N., Selby, I., Graham, D. K., Harland, R., Wilkinson, I.  
2 P., 1992. Late Devensian and Flandrian palaeoenvironmental changes on the  
3 Scottish continental shelf west of the Outer Hebrides. *Journal of Quaternary*  
4 *Science* 7, 145-161.
- 5 Peacock, J.D., Armstrong, M., Browne, M.A.E., Golledge, N.R. and Stoker, M.S.  
6 2007. The deglaciation of eastern Scotland: a response to McCabe et al.  
7 (2007). *Journal of the Geological Society, London* 164, 1261
- 8 Peck, V. L., Hall, I. R., Zahn, R., Grousset, F., Hemming, S. R. 2007. The relationship  
9 of Heinrich events and their European precursors over the past 60 ka BP: a  
10 multi-proxy ice-rafted debris provenance study in the North East Atlantic.  
11 *Quaternary Science Reviews* 26, 862–875.
- 12 Peltier, W.R., Fairbanks, R.G., 2006. Global glacial ice volume and Last Glacial  
13 Maximum duration from an extended Barbados sea level record. *Quaternary*  
14 *Science Reviews* 25, 3322-3337.
- 15 Praeg, D., 2003. Seismic imaging of mid-Pleistocene tunnel-valleys in the North Sea  
16 Basin-high resolution from low frequencies. *Journal of Applied Geophysics*,  
17 53, 273-298.
- 18 Rack, W., Rott, H., 2004. Pattern of retreat and disintegration of Larsen B ice shelf,  
19 Antarctic Peninsula, *Annals of Glaciology* 39, 505-510.
- 20 Rae, D.A., 1976. Aspects of glaciation in Orkney. Unpublished PhD thesis. University  
21 of Liverpool.
- 22 Reeh, N., 1968. On the calving of ice from floating glaciers and ice shelves. *Journal*  
23 *of Glaciology* 7, 215–232.
- 24 Rise, L., Olesen, O., Rokoengen, K., Ottesen, D., Riis, F., 2004. Mid-Pleistocene ice  
25 drainage pattern in the Norwegian Channel imaged by 3D seismic. *Quaternary*  
26 *Science Reviews* 23, 2323-2335.
- 27 Robinson, M., Ballantyne, C.K. 1979. Evidence for a glacial advance predating the  
28 Loch Lomond Advance in Wester Ross. *Scottish Journal of Geology* 15, 271-  
29 277.
- 30 Rokoengen, K., Løfaldli, M., Rise, L., Løken, T., Carlsen, R., 1982. Description and  
31 dating of a submerged beach in the northern North Sea Marine. *Geology* 50, 1-  
32 2.
- 33 Rose, J., 1985. The Dimlington Stadial/Dimlington Chronozone: a proposal for  
34 naming the main glacial episode of the Late Devensian in Britain. *Boreas* 14,  
35 225-230.
- 36 Ross, H., 1996. The last glaciation of Shetland. Unpublished PhD thesis. University of  
37 St Andrews.
- 38 Scambos, T.A., Bohlander, J.A., Shuman, C. A., Skvarca, P., 2004. Glacier  
39 acceleration and thinning after ice shelf collapse in the Larsen B embayment,  
40 Antarctica. *Geophysical Research Letters*, 31, L18402 doi:10.1029/2004  
41 GL020670, 2004.
- 42 Sejrup, H.P., Aarseth, I., Ellingsen, K. L., Reither, E., Jansen, E., Løvlie, R., Bent,  
43 A., Brigham-Grette, J., Larsen, E., Stoker, M., 1987. Quaternary stratigraphy  
44 of the Fladen area, central North Sea: A multidisciplinary study. *Journal of*  
45 *Quaternary Science* 2, 35-58.
- 46 Sejrup, H.P., Aarseth, I., Haflidason, H., 1991. The Quaternary succession in the  
47 northern North Sea. *Marine Geology* 101, 103-111.
- 48 Sejrup, H.P., Haflidason, H., Aarseth, I., King, E., Forsberg, C.F., Long, D.,  
49 Rokoengen, K., 1994. Late Weichselian glaciation history of the northern  
50 North Sea. *Boreas* 23, 1–13.

- 1 Sejrurp, H.P., Landvik, J. Y., Larsen, E., Janocko, J., Eiriksson, J., King, E., 1998.  
2 The Jaeren area, a border zone of the Norwegian channel ice stream.  
3 Quaternary Science Reviews 17, 801-812.
- 4 Sejrurp, H.P., Larsen, E., Landvik, J., King, E.L., Haflidason, H., Nesje, A. 2000.  
5 Quaternary glaciations in southern Fennoscandia: evidence from south-  
6 western Norway and the northern North Sea region. Quaternary Science  
7 Reviews, 19, 7, 667-685.
- 8 Sejrurp, H.P., Larsen, E., Haflidason, H., Berstad, I.M., Hjelstuen, B.O., Jonsdottir,  
9 H.E., King, E.L., Landvik, J., Longva, O., Nygård, A., Ottesen, D., Raunholm,  
10 S., Rise, L., Stalsberg, K., 2003. Configuration, history and impact of the  
11 Norwegian Channel Ice Stream. Boreas 32, 18-36.
- 12 Sejrurp, H. P., Hjelstuen, B. O., Dahlgren, K. I. T., Haflidason, H., Kuijpers, A.,  
13 Nygard, A., Praeg, D., Stoker, M. S., Vorren, T. O., 2005. Pleistocene glacial  
14 history of the NW European continental margin. Marine and Petroleum  
15 Geology 22, 1111–1129.
- 16 Selby, I., 1989. Quaternary Geology of the Hebridean Continental Margin.  
17 Unpublished PhD thesis. University of Nottingham.
- 18 Shackelton, N.J., Backman, J., Zimmerman, H., Kent, D.V., Hall, M.A., Roberts,  
19 D.G., Schnitker, D.G., Baldauf, J.G., Despraires, A., Homrighausen, R.,  
20 Huddlestun, P., Kenne, J.B., Kaltenback, A.J., Krumslek, K.A.O., Morton,  
21 A.C., Murray, J.W., Westberg-Smith, J., 1984. Oxygen isotope calibration of  
22 the onset of ice-rafting and history of glaciation in the North Atlantic region.  
23 Nature 307, 620-623.
- 24 Shaw, J., Piper, D.J.W., Fader, G.B.J., King, E.L., Todd, B.J., Bell, T., Batterson,  
25 M.J., Liverman, D.G.E., 2006. A conceptual model of the deglaciation of  
26 Atlantic Canada. Quaternary Science Reviews 25, 2059–2081.
- 27 Shennan, I., Bradley, S., Milne, G., Brooks, A., Bassett, S., Hamilton, S., 2006.  
28 Relative sea-level changes, glacial isostatic modelling and ice-sheet  
29 reconstructions from the British Isles since the Last Glacial Maximum. Journal  
30 of Quaternary Science 21, 585-599.
- 31 Shotton, F.W., 1977. The English Midlands. INQUA Excursion Guide A2, Xth  
32 INQUA Congress, Birmingham, 51 pp.
- 33 Siegert, M. J., Dowdeswell, J. A., Svendsen, J. I., Elverhøi, A., 2002. The Eurasian  
34 Arctic during the last Ice Age. American Scientist 90, 32–39.
- 35 Sissons, J.B., 1967. The Evolution of Scotland's Scenery. Olivier and Boyd,  
36 Edinburgh, 259 pp.
- 37 Sissons, J.B., 1980. The glaciation of the Outer Hebrides (Letter). Scottish Journal of  
38 Geology 16, 81-4.
- 39 Skinner, A.C., Gregory, D.M., 1983. Quaternary stratigraphy in the northern North  
40 Sea. Boreas 12, 145-152.
- 41 Smith, M.J., Clark, C.D., 2005. Methods for the visualization of digital elevation  
42 models for landform mapping. Earth Surface Processes and Landforms 30,  
43 885-900.
- 44 Sollid, J. L., 1989. Comments on the genesis of De Geer moraines. Norsk Geografisk  
45 Tidsskrift 43, 45–47.
- 46 Stalsberg, K., Larsen, E., Ottesen, D., Sejrurp, H.P., 2003a. Middle to Late  
47 Weichselian Norwegian Channel Ice Stream deposits and morphology on  
48 Jæren, south-western Norway and the eastern North Sea area Boreas, 32, 149-  
49 166.



- 1 Stoker, M.S., 1990. Glacially-influenced sedimentation on the Hebridean slope, north-  
2 western United Kingdom continental margin. In: Dowdeswell, J.A. and  
3 Scourse, J.D. (Eds.), *Glacimarine Environments: Processes and Sediments*.  
4 Geological Society, London, Special Publications 53, 349–362.
- 5 Stoker, M.S., 1995. The influence of glacially-influenced sedimentation on slope-apron  
6 development on the continental margin off Northwest Britain. In: R.A.  
7 Scrutton, Stoker, M.S., Shimmiel, G.B., Tudhope, A.W. (Eds.), *The*  
8 *Tectonics, Sedimentation and Paleoceanography of the North Atlantic Region*.  
9 Geological Society, London, Special Publications, London, pp159–177.
- 10 Stoker, M.S., 2002. Late Neogene development of the UK Atlantic margin. . In: A.G.  
11 Doré, Cartwright, J.A., Stoker, M.S, Turner, J.P., White, N. (Eds.),  
12 *Exhumation of the North Atlantic margin: Timing, mechanisms and*  
13 *implications for petroleum exploration*. Geological Society, London, Special  
14 Publications London, pp313–329.
- 15 Stoker, M.S., In press. Cenozoic sedimentary rocks. In: United Kingdom offshore  
16 regional report: the geology of the Rockall Basin. Hitchen, K. (ed.) British  
17 Geological Survey, HMSO, London.
- 18 Stoker, M.S., Bradwell, T., 2005. The Minch palaeo-ice stream, NW sector of the  
19 British–Irish Ice Sheet. *Journal of the Geological Society*, London 162, 425-  
20 428.
- 21 Stoker, M.S., Holmes, R., 1991. Submarine end-moraines as indicators of Pleistocene  
22 ice-limits off northwest Britain. *Journal of the Geological Society*, London  
23 148, 431-434.
- 24 Stoker, M.S., Varming, T., In press. Cenozoic sedimentary rocks. In: United Kingdom  
25 offshore regional report: the geology of the Faroe-Shetland Basin. Ritchie J.  
26 (Ed.). British Geological Survey, HMSO, London.
- 27 Stoker, M.S., Long, D., Fyfe, J.A., 1985. A revised Quaternary stratigraphy for the  
28 central North Sea. Report of the British Geological Survey, 17/2. Edinburgh.
- 29 Stoker, M.S., Hitchen, K., Graham, C.C., 1993. United Kingdom offshore report: the  
30 geology of the Hebrides and West Shetland shelves, and adjacent deep-water  
31 areas. London: HMSO for the British Geological Survey.
- 32 Stoker, M.S., Leslie, A.B., Scott, W.D., Briden, J.C., Hine, N.M., Harland, R.,  
33 Wilkinson, I.P., Evans, D., Ardur, D.A. 1994. A record of late Cenozoic  
34 stratigraphy, sedimentation and climate change from the Hebrides Slope, NE  
35 Atlantic Ocean. *Journal of the Geological Society*, London 151, 235-249.
- 36 Stoker, M.S., Bradwell, T., Wilson, C., Harper, C., Smith, D., Brett, C., 2006. Pristine  
37 fjord landsystem revealed on the sea bed in the Summer Isles region, NW  
38 Scotland. *Scottish Journal of Geology* 42, 89–99.
- 39 Stokes, C. R., Clark, C. D., 2003. The Dubawnt Lake palaeo-ice stream: evidence for  
40 dynamic ice-sheet behaviour on the Canadian Shield and insights regarding  
41 the controls on ice-stream location and vigour. *Boreas* 32, 263-279.
- 42 Stone, J.O., Ballantyne, C.K., Fifield, L.K., 1998. Exposure dating and validation of  
43 periglacial weathering limits, northwest Scotland. *Geology* 26, 87-590.
- 44 Stone, J.O., Ballantyne, C.K., 2006. Dimensions and deglacial chronology of the  
45 Outer Hebrides Ice Cap, northwest Scotland: implications of cosmic-ray  
46 exposure dating. *Journal of Quaternary Science* 21, 75-84.
- 47 Stuiver, M., Reimer, P.J., Reimer, R.W., 2005. CALIB 5.0.1 (Computer program and  
48 documentation). Available from: <http://radiocarbon.pa.qub.ac.uk/calib/>.
- 49 Sutherland, D.G., 1984. The Quaternary deposits and landforms of Scotland and the  
50 neighbouring shelves: A review. *Quaternary Science Reviews* 3, 157-254.

1 Thomas, R. H., 1977. Calving bay dynamics and ice-sheet retreat up the St. Lawrence  
2 valley system. *Geographie Physique et Quaternaire* 31, 347–356.

3 Thomas, R., Rignot, E., Casassa, G., Kanagaratnam, P., Acuña, C., Akins, T.,  
4 Brecher, H., Frederick, E., Gogineni, P., Krabill, W., Manizade, S.,  
5 Ramamoorthy, H., Rivera, A., Russell, R., Sonntag, J., Swift, R., Yungel, J.,  
6 Zwally J. 2004. Accelerated Sea-Level Rise from West Antarctica. *Science*  
7 306, 255-258.

8 Thomson, M.E., Eden, R.A., 1977. Quaternary deposits of the central North Sea, 3:  
9 The Quaternary sequence in the west-central North Sea, British Geological  
10 Survey.

11 van der Veen, C.J. 1996. Tidewater calving. *Journal of Glaciology* 42, 375–385.

12 von Weymarn, J., Edwards, K.J., 1973. Interstadial site on the island of Lewis,  
13 Scotland. *Nature* 246, 473-474.

14 Walker, M.J.C., Bjorck, S., Lowe, J.J., Cwynar, L.C., Johnsen, S., Knudsen, K.-L.,  
15 Wohlfarth, B., INTIMATE group, 1999. Isotopic ‘events’ in the GRIP ice  
16 core: a stratotype for the Late Pleistocene. *Quaternary Science Reviews* 18,  
17 1143-1150.

18 Whittington, G., Hall, A.M., 2002. The Tolsta Interstadial, Scotland: a correlation  
19 with D-O cycles GI-8 to GI-5? *Quaternary Science Reviews* 21, 901-915.

20 Wilson, L.J., Austin, W.E.N., Jansen, E., 2002. The last British Ice Sheet: growth,  
21 maximum extent and deglaciation. *Polar Research* 21, 243-250.

22 Wilson, G.V., Edwards, W., Knox, J., Jones, R.C.B., Stevens, J.V., 1935. The  
23 Geology of the Orkneys. *Memoirs of the Geological Survey of Scotland*,  
24 HMSO, Edinburgh, 205 pp.

25 Wingfield, R., 1989. Glacial incisions indicating Middle and Upper Pleistocene ice  
26 limits off Britain. *Terra Nova* 1, 31-52.

27 Wingfield, R., 1990. The origin of major incisions within the Pleistocene deposits of  
28 the North Sea. *Marine Geology* 91, 1-2, 31-52.

29 Wintle, A.G., Catt, J.A., 1985. Thermoluminescence dating of Dimlington Stadial  
30 deposits in eastern England. *Boreas* 14, 3, 231-234.

31 Zwally, H. J., Abdalati, W., Herring, T., Larson, K., Saba, J., Steffen, K., 2002.  
32 Surface melt-induced acceleration of Greenland Ice Sheet flow. *Science* 297,  
33 218–222.

34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

**Figure captions**

**Figure 1.** Map showing the general bathymetry and main glacio-geological sea-bed landforms around the northern UK. Data sources are given in the legend. Extent of the study area (Figure 4) and key placenames are also shown.

**Figure 2.** Summary of Late Quaternary chronostratigraphic stages (UK) and their correlation with Marine Isotope Stages (modified from Bowen, 1999 and Walker et al., 1999). Dates are in calendar years BP. The timing of the Last Glacial Maximum (LGM) in NW Europe is also shown. Note: diagram is not to scale and correlations are not fully resolved.

**Figure 3.** BGS airgun profile and interpreted line drawing across the West Shetland margin showing the geometry and stratigraphic architecture of the Rona Wedge (see Figure 1 for location). Inset image shows the sea bed morphology of the West Shetland slope (A-A' on the profile) derived from first returns from 3D exploration surveys (cf. Long et al., 2004). The Glacial Unconformity (GU) is interpreted to represent the instigation of shelf-wide glaciation during the early Mid-Pleistocene (Stoker, 1995, 2002). Prominent ridges preserved on the outer shelf represent submarine moraine banks, up to 50 m high and 6 km wide. The West Shetland Slope surface morphology displays abundant glacial debris flows sourced from an ice-sheet margin at the continental shelf edge (Davison and Stoker, 2002). Abbreviations: SBM, sea bed multiple; WD, water depth.

**Figure 4.** Merged onshore-offshore (topographic-bathymetric) surface model for the northern UK and surrounding continental shelf. Offshore data – Olex AS; Onshore data – derived from NEXTMap Britain digital surface model (Intermap Technologies, 2003). See text for details of data collection and image production.

**Figure 5.** Sea-bed landforms on the northern UK continental shelf mapped from the Olex dataset (Figure 4). Solid lines – positive linear features (ridges, primarily moraines); dashed lines – negative linear features (channels, primarily tunnel valleys). There is some generalisation, in places, for the sake of cartographic clarity. Colours denote feature groups outlined in Key (1, 2, 3, A, B).

**Figure 6.** (A) Olex bathymetry of the northern North Sea showing the morphology and distribution of large tunnel valleys (Fladen Deep). For location see Figure 5. Most authors relate these features to subglacial hydrological phenomena (e.g. Ó Cofaigh, 1996; Lonergan et al., 2006). (B) Rose diagram showing the preferred orientation of the tunnel valleys in the northern North Sea. (C) Oblique view, looking SE from point X, along a well-developed sea-floor tunnel valley. The feature is >100 m deep, 1 km wide and over 30 km long. Note the undulating long profile and abrupt initiation/termination. 5x vertical exaggeration. Faint orthogonal linear features are data artefacts.

1 **Figure 7.** (A) Olex bathymetry and (B) geomorphological map of the sea bed  
2 NW of Orkney. For location see Figure 5. Black polygons – large ridges; black lines –  
3 narrow ridges. The large concentric ridges with irregular, in places overprinted,  
4 morphology are interpreted as recessional moraines relating to a grounded, highly  
5 dynamic, oscillatory lobe of the last BIS. Note the sharp-crested coherent morphology  
6 of the innermost arcuate loop – strongly reminiscent of a terrestrial thrust-block or  
7 push moraine. (C) BGS seismic profile and interpreted line drawing of submarine  
8 moraine banks along line X-X' (modified after Stoker and Holmes, 1991). The  
9 moraine banks are up to 50 m high and 6 km wide; the crests of the banks are  
10 currently at water depths of 100-150 m. The moraines are largely acoustically  
11 structureless. The observation that the most landward moraine imaged in the seismic  
12 profile overlies layered strata which post-date the middle moraine reflects a  
13 considerable readvance of the BIS during the general phase of overall recession.


14  
15 **Figure 8.** (A) Olex bathymetry and (B) geomorphological map of the sea bed NE  
16 of Shetland. For location see Figure 5. Black polygons – large ridges; black lines –  
17 narrow ridges; trough margin shown as triangle-ornamented line. The high density of  
18 arcuate, convoluted, occasionally overprinting ridges NE of Unst are interpreted as  
19 recessional moraines of a grounded, highly dynamic, oscillatory lobe of the last BIS.  
20 The broadly concentric configuration of moraines around Shetland supports the notion  
21 of an independent ice cap on the islands during the latter stages of the last glaciation.  
22 (C) Close-up image of subparallel more-linear ridges within the trough east of  
23 Shetland. Based on their geomorphological expression, these are interpreted as De  
24 Geer moraines – strongly suggestive of a tidewater glacier margin. (D) Olex  
25 bathymetry and geomorphological map of closely spaced low-elevation ridges in the  
26 Fair Isle channel, also interpreted as De Geer moraines.

27  
28 **Figure 9.** Olex bathymetry of the northern North Sea showing coloured  
29 bathymetric slices at various elevations: (i) 160-190 m below present day; (ii) 120-  
30 150 m below present day; and (iii) 90-120 m below present day sea level. Darker  
31 tones are deeper water. The pattern of sea-level rise associated with ice-sheet  
32 deglaciation would have followed this sequence, resulting in a large north-south  
33 marine embayment in the northern North Sea Basin. Note: No corrections for glacio-  
34 isostatic loading or forebulge development have been incorporated into these sea-level  
35 scenarios.

36  
37 **Figure 10.** Reconstruction of the confluent British (BIS) and Fennoscandian ice  
38 sheets (FIS) at LGM (30-25 ka BP), with hypothesised flow lines. Solid lines inferred  
39 from this study; dashed flow lines (FIS) taken from Sejrup et al. (1994) and Ottesen et  
40 al. (2005). Dark shading shows approximate zone of confluence (CZ); small arrows  
41 show MSGL orientations recorded (from west to east) by Bradwell et al. (2007),  
42 Stalsberg et al. (2003) and Graham et al. (2007). Hatching denotes trough-mouth fans.  
43 FB denotes possible ice sheet frozen-bed patches in Buchan and on Lewis. Sea-bed  
44 contours refer to present-day bathymetry (see Figure 1). Faeroe Ice cap not shown,  
45 extent uncertain (see Nielsen et al., 2007).

46  
47 **Figure 11.** Reconstruction of British and Fennoscandian ice sheets during LGM  
48 deglaciation. Three sequential stages are shown: 1 (black line) – calving bay  
49 initiation; lobes dominate western margin. 2 (grey line) – calving bay well developed;  
50 dynamic ice-sheet separation and re-organisation ongoing. 3 (shaded fill) – separate

1 BIS and FIS; ice free central North Sea Basin; independent Shetland ice cap.  
2 Hypothesised flow lines shown for final stage only; those for FIS taken from Sejrup et  
3 al. (1994) and Ottesen et al. (2005). Sea-bed contours refer to present-day bathymetry  
4 (see Figure 1).  
5  
6



Key to abbreviations:

**Trough Mouth Fans**

- BD Barra-Donagel Fan
- SS Sula Sgeir Fan
- R Rona Wedge
- F Foula Wedge
- S Suduroy Fan
- Sa Sandoy Fan
- Sk Skeivi Fan
- NS North Sea Fan

**Moraines**

- BB Bosies Bank
- WB Wee Bankie
- OB Otter Bank
- TR Tampen Ridge
- WRR Wester Ross Moraine

**Bathymetric Deeps**

- FD Fladen Deeps
- DHD Devil's Hole Deeps
- ST Southern Trench

**Islands**

- StK St Kilda
- FL Flannan Islands
- H Harris
- NR North Rona
- FI Fair Isle
- Fa Foula

**Data sources**

- Moraines  
*Robinson and Ballantyne (1979)*  
*Cameron et al. (1987)*  
*Hall and Bent (1990)*  
*Selby (1989)*  
*Stoker (1990)*  
*Stoker and Holmes (1991)*  
*Davison (2005)*

- Trough-Mouth Fans  
*Sejrup et al. (2005)*  
*Nygaard et al. (2005)*  
*Stoker and Varming (in press)*  
*Stoker (in press)*



Box showing extent and orientation of mega-scale glacial lineations, mapped by Graham et al. (2007)

Period	Marine Isotope Stage	Stage		Age (ka)
QUATERNARY	1	HOLOCENE		11.5
	2	DEVENSIAN = Weichselian	LATE	
			<i>LGM</i>	31
	3		MID	58
	4		EARLY	
	5	IPSWICHIAN = Eemian		116

