

The On-Line Encyclopedia of Integer Sequences



Neil J. A. Sloane

Introduction

The *OEIS*[®] (or *On-Line Encyclopedia of Integer Sequences*[®])¹ is a freely accessible database of number sequences, now in its 54th year, and online since 1995. It contains over 300,000 entries, and for each one gives a definition, properties, references, computer programs, tables, etc., as appropriate. It is widely referenced: a webpage² lists over 6,000 works that cite it, and often say things like “this theorem would not exist without the help of the *OEIS*.” It has been called one of the most useful mathematical sites in the Web.

The main use is to serve as a dictionary or fingerprint file for identifying number sequences (and when you find the sequence you are looking for, you will understand why the *OEIS* is so popular). If your sequence is not recognized, you see a message saying that if the sequence is of general interest, you should submit it for inclusion in the database. The resulting queue of new submissions is a continual source of lovely problems.

I described the *OEIS* in a short article in the September 2003 issue of these *Notices*. The most significant changes since then took place in 2009, when a nonprofit foundation³ was set up to own and maintain the *OEIS*, and in 2010 when the *OEIS* was moved off my home page at AT&T Labs to a commercial host. The format has also changed: since 2010 the *OEIS* has been a refereed “wiki.” Four people played a crucial role in the transition: Harvey P. Dale and Nancy C. Eberhardt helped set up the Foundation, Russell S. Cox wrote the software, and David L. Applegate helped move the *OEIS*. The *OEIS* would probably not exist today but for their help.

All submissions, of new sequences and updates, are now refereed by volunteer editors. One of the rewards of being an editor is that you see a constant flow of new problems, often submitted by nonmathematicians, which frequently contain juicy-looking questions that are begging to be investigated.

This article will describe a selection of recent sequences, mostly connected with unsolved problems.

Sequences in the *OEIS* are identified by a 6-digit number prefixed by A. [A000001](#) is the number of groups of order n , [A000002](#) is Kolakoski’s sequence, and so on. When we were approaching a quarter of a million entries, the editors voted to decide which sequence would become [A250000](#). The winner was the Peaceable Queens sequence, described in the next section, and the runner-up was the “circles in the plane” sequence [A250001](#) discussed after that. The n th term of the sequence under discussion is usually denoted by $a(n)$.

Neil J. A. Sloane is president of The *OEIS* Foundation, Inc. and a visiting scientist at Rutgers University. His email address is njasloane@gmail.com.

¹oeis.org.

²oeis.org/wiki/Works_Citing_OEIS.

³The *OEIS* Foundation, Inc., oeisf.org.

For permission to reprint this article, please contact: reprint-permission@ams.org.

DOI: <http://dx.doi.org/10.1090/noti1734>



Figure 1. One of three solutions to the Peaceable Queens problem on a 5×5 board, illustrating $a(5) = 4$.




Figure 2. A solution to the Peaceable Queens problem on an 8×8 board, illustrating $a(8) = 9$. (There are actually 10 white queens here but only 9 count since the numbers of white and black queens must be equal. Any one of the white queens could be omitted.)

Peaceable Queens

In [A250000](#), $a(n)$ is the maximal number m such that it is possible to place m white queens and m black queens on an $n \times n$ chess board so that no queen attacks a queen of the opposite color. These are peaceable queens. This is a fairly new problem with some striking pictures,




Figure 3. A solution to the Peaceable Queens problem on an 11×11 board, illustrating $a(11) = 17$.




Figure 4. A conjectured solution to the Peaceable Queens problem on a 20×20 board, found by Bob Selcoe, showing that $a(20) \geq 58$.

an interesting conjecture, and a satisfactorily nonviolent theme. It was posed by Robert A. Bosch in 1999, as a variation on the classical problem of finding the number of ways to place n queens on an $n \times n$ board so that they do not attack each other ([A000170](#)). It was added to the OEIS in 2014 by Donald E. Knuth, and a number of people have contributed to the entry since then. Only thirteen terms are known:

n :	1	2	3	4	5	6	7	8	9	10	11	12	13
$a(n)$:	0	0	1	2	4	5	7	9	12	14	17	21	24

Figures 1-4 show examples of solutions for $n = 5, 8, 11$ and (conjecturally) 20.

For larger values of n , the best solutions presently known were found by Benoît Jubin and concentrate the queens into four pentagonal regions, as shown in Figure 5

(and generalize the arrangement shown in Figure 4). This construction gives a lower bound of $\lfloor 7n^2/48 \rfloor$, a formula which in fact matches all the best arrangements known so far except $n = 5$ and 9. It would be nice to know if this construction really does solve the problem!




Figure 5. A general construction for the Peaceable Queens problem found by Benoît Jubin, showing that for large n , $a(n) \geq \lfloor 7n^2/48 \rfloor$, a formula which might be exact for all $n > 9$.

Circles in the Plane

The runner-up in the competition for [A250000](#) is now [A250001](#): here $a(n)$ is the number of ways to draw n circles in the affine plane. Two circles must be disjoint or meet in two distinct points (tangential contacts are not permitted), and three circles may not meet at a point.⁴ The sequence was proposed by Jonathan Wild, a professor of music at McGill University, who found the values $a(1) = 1$, $a(2) = 3$, $a(3) = 14$, $a(4) = 173$, and, jointly with Christopher Jones, $a(5) = 16951$ (see Figures 6-8).

Wild and Jones have found that there are complications which first appear when five circles are being considered: here there are arrangements which theoretically could exist if one considered only the intersections between circles, but which cannot actually be drawn using circles. For example, start with four circles arranged in a chain, each one overlapping its two neighbors, and label the overlaps a, b, c, d (see Figure 9). Suppose we try to add a fifth circle that meets all four circles but avoids their overlaps, encloses overlaps b and d , but does not enclose overlaps a or c . This can be drawn if the fifth circle

⁴The circles may have different radii. Two arrangements are considered the same if one can be continuously changed to the other while keeping all circles circular (although the radii may be continuously changed), without changing the multiplicity of intersection points, and without a circle passing through an intersection point. Turning the whole configuration over is allowed.




Figure 6. The fourteen ways to draw three circles in the affine plane.




Figure 7. Eight of the 173 ways to draw four circles. For the full set of 173 drawings, see [A250001](#).




Figure 8. Seven further ways (out of 173) to draw four circles.

is flattened to an ellipse, but it can be shown that the arrangement cannot be realized with five circles. There are twenty-six such unrealizable arrangements of five circles, which can be ruled out by ad hoc arguments.

The delicate configurations like those in Figure 8 are very appealing. It would be interesting to see all 17142 arrangements of five or fewer circles displayed along the Great Wall of China.




Figure 9. A hypothetical arrangement of five circles that can only be realized if one or more of the circles is distorted.

Lexicographically Earliest Cube-Free Binary Sequence

There is an obvious way to sort integer sequences $a(1), a(2), a(3), a(4), \dots$ into lexicographic order. A number of recent entries in the *OEIS* are defined to be the lexicographically earliest sequence of nonnegative or positive integers satisfying certain conditions.

For example, one of the first results in the subject now called “Combinatorics on Words” was Axel Thue’s 1912 theorem that the “Thue-Morse sequence”

$$T = 0, 1, 1, 0, 1, 0, 0, 1, 1, 0, 0, 1, \\ 0, 1, 1, 0, 1, 0, 0, 1, 0, 1, 1, 0, 0, 1, \dots$$

[\(A010060\)](#) contains no substring of the form XXX , that is, T is *cube-free*. T can be defined as a fixed point of the mapping $0 \rightarrow 01, 1 \rightarrow 10$; alternatively, by taking $a(n)$ to be the parity of the number of 1s in the binary expansion of n . One hundred and five years later, David W. Wilson asked for the lexicographically earliest cube-free sequence of 0s and 1s. Using a back-tracking algorithm, he found what appear to be the first 10000 terms, which begin

$$(1) \quad 0, 0, 1, 0, 0, 1, 0, 1, 0, 0, 1, 0, 0, 1, \\ 1, 0, 0, 1, 0, 0, 1, 0, 1, 0, 0, 1, \dots$$

This is now [A282317](#).

There is no difficulty in showing that the sequence exists.⁵ To see this, make the set S of all infinite binary sequences $a = (a(1), a(2), \dots)$ into a metric space by defining $d(a, b)$ to be 0 if $a = b$, or 2^{-i} if a and b first differ at position i . This identifies S with the Cantor set in $[0, 1)$. The subset $F \subset S$ of infinite cube-free sequences is nonempty and has an infimum c say. It is easy to show that the complement $S \setminus F$, sequences that contain a cube, is an open set in this topology, so F is closed and $c \in F$.

So far only the first 999 terms of [A282317](#) have been verified to be correct (by showing that there is at least one infinite cube-free sequence with that beginning). The rest of the 10000 terms are only conjectural. It would be nice to know more. In particular, does this sequence have an alternative construction? There is no apparent formula or recurrence, which seems surprising.

The EKG and Yellowstone Sequences

To continue the “lexicographically earliest” theme, many recent entries in the *OEIS* are defined to be the lexicographically earliest sequence $a(1), a(2), \dots$ of distinct positive integers satisfying certain divisibility conditions.

The first task here is usually to show that there are no missing numbers, i.e., that the sequence is a permutation of the positive integers. Sequences of this type were studied in a 1983 paper by Erdős, Freud, and Hegyvári, which included the examples [A036552](#) ($a(2n) = \text{smallest missing number}$, $a(2n+1) = 2a(2n)$) and [A064736](#) ($a(2n+2) = \text{smallest missing number}$, $a(2n+1) = a(2n) \cdot a(2n+2)$). For these two it is clear that there are no missing numbers. This is less obvious, but still true, for Jonathan Ayres’s *EKG sequence*, [A064413](#), defined to be the lexicographically earliest sequence of distinct positive integers such that

$$\gcd(a(n-1), a(n)) > 1 \text{ for all } n \geq 3.$$

This begins

$$1, 2, 4, 6, 3, 9, 12, 8, 10, 5, 15, 18, \\ 14, 7, 21, 24, 16, 20, 22, 11, 33, 27, \dots$$

The proof that it is a permutation is omitted—it is similar to the proof for the Yellowstone sequence given below.

Next, one can investigate the rate of growth. In the case of [A064413](#), the points appear to lie roughly on three curved lines (Figure 10), although the following conjecture of Lagarias, Rains, and Sloane (2002) is still open.

Plot of [A064413](#)




Figure 10. The first 10000 terms of the EKG sequence, so named because locally this graph resembles an EKG. Every number appears exactly once.

Conjecture 1. In the EKG sequence [A064413](#), if $a(n)$ is neither a prime nor three times a prime then

$$a(n) \sim n \left(1 + \frac{1}{3 \log n} \right);$$

if $a(n)$ is a prime then

$$a(n) \sim \frac{1}{2} n \left(1 + \frac{1}{3 \log n} \right);$$

and if $a(n)$ is 3 times a prime then

$$a(n) \sim \frac{3}{2} n \left(1 + \frac{1}{3 \log n} \right).$$

Furthermore, if the sequence is a permutation, one can also try to study its cycle structure. However, this often leads to very difficult questions, similar to those encountered in studying the Collatz conjecture, and we can’t do much more than collect experimental data. Typically there is a set of finite cycles, and one or more apparently infinite cycles, but we can’t prove that the apparently infinite cycles really are infinite, nor that they are distinct. See the entries for [A064413](#) and [A098550](#) for examples.

The definition of the *Yellowstone sequence* (Reinhard Zumkeller, 2004, [A098550](#), [1]) is similar to that of the EKG sequence, but now the requirement is that, for $n > 3$,

$$\gcd(a(n-2), a(n)) > 1 \text{ and } \gcd(a(n-1), a(n)) = 1.$$

This begins

$$1, 2, 3, 4, 9, 8, 15, 14, 5, 6, 25, 12, 35, \\ 16, 7, 10, 21, 20, 27, 22, 39, 11, \dots$$

Figure 11 shows terms $a(101) = 47$ through $a(200) = 279$, with successive points joined by lines.

Theorem 2. The Yellowstone sequence [A098550](#) is a permutation of the positive integers.

⁵Thanks to Jean-Paul Allouche for this argument.

The proof is typical of the arguments used to prove that several similar sequences are permutations, including the EKG sequence above.

Proof. There are several steps.

(i) The sequence is infinite. (For $pa(n-2)$ is always a candidate for $a(n)$, where p is a prime larger than any divisor of $a(i)$, $i < n$.)

(ii) There are infinitely many different primes that divide the terms of the sequence. (If not, there is a prime p such that all terms are products of primes less than p . Using (i), find a term $a(n) > p^2$, and let q be a common prime factor of $a(n-2)$ and $a(n)$. But now $pq < p^2 < a(n)$ is a smaller candidate for $a(n)$, a contradiction.)

(iii) For any prime p , some term is divisible by p . (For if not, no prime $q > p$ can divide any $a(n)$: if $a(n) = kq$ is the first multiple of q to appear, kp would be a smaller candidate for $a(n)$. This contradicts (ii).)

(iv) For any prime p , p divides infinitely many terms. (If not, let p^i be larger than any multiple of p in the sequence, and choose a prime $q > p^i$. Again we obtain a contradiction.)

(v) Every prime p is a term in the sequence. (Suppose not, and using (i), choose n_0 such that $a(n) > p$ for all $n > n_0$. Using (iv), find $a(n) = kp$, $k > 1$, for some $n > n_0$. But then $a(n+2) = p$, a contradiction.)

(vi) All numbers appear. For if not, let k be the smallest missing number, and choose n_0 so that all of $1, \dots, k-1$ have occurred in $a(1), \dots, a(n_0)$. Let p be a prime dividing k . Since, by (iv), p divides infinitely many terms, there is a number $n_1 > n_0$ such that $\gcd(a(n_1), k) > 1$. This forces

(2) $\gcd(a(n), k) > 1$ for all $n \geq n_1$.

(If not, there would be some $j \geq n_1$ where $\gcd(a(j), k) > 1$ and $\gcd(a(j+1), k) = 1$, which would lead to $a(j+2) = k$.) But (2) is impossible, because we know from (v) that infinitely many of the $a(n)$ are primes. \square




Figure 11. Plot of terms $a(101)$ through $a(200)$ of the Yellowstone sequence. The sequence has a downward spike to $a(n)$ when $a(n)$ is a prime, and larger upward spikes (the “geysers,” which suggests the name for this sequence) two steps later.

The growth of this sequence is more complicated than that of the EKG sequence. Figure 12 shows the first 300,000 terms, without lines connecting the points. The points appear to fall on or close to a number of distinct curves. There is a conjecture in [1, p. 5] that would explain these curves.

Three Further Lexicographically Earliest Sequences

Here are three further examples of this type, all of which are surely permutations of the positive integers. For the first there is a proof, for the second there is “almost” a proof, but the third may be beyond reach.

The first (Leroy Quet, 2007, [A127202](#)) is the lexicographically earliest sequence of distinct positive integers such that

$$\gcd(a(n-1), a(n)) \neq \gcd(a(n-2), a(n-1)) \quad \text{for } n \geq 3.$$

It begins

1, 2, 4, 3, 6, 5, 10, 7, 14, 8, 9, 12, 11, 22,
13, 26, 15, 18, 16, 17, 34, 19, ...

For the second (Rémy Sigrist, 2017, [A280864](#)), the definition is

if a prime p divides $a(n)$, then it divides
exactly one of $a(n-1)$ and $a(n+1)$, for $n \geq 2$,

and the initial terms are

1, 2, 4, 3, 6, 8, 5, 10, 12, 9, 7, 14, 16, 11,
22, 18, 15, 20, 24, 21, 28, 26, ...

The proof that the first is a permutation is similar to that for the Yellowstone sequence, although a bit more involved (see [A127202](#)). The second struck me as one of




Figure 12. Scatterplot of the first 300,000 terms of the Yellowstone sequence. The primes lie on the lowest line (labeled “p”), the even numbers on the second line (“E”), the majority of the odd composite numbers on the third line (“C”), and the $3p$, $5p$, $7p$, $11p, \dots$ points on the higher lines. The lines are not actually straight, except for the red line $f(x) = x$, which is included for reference.

those “drop everything and work on this” problems that are common hazards when editing new submissions to the *OEIS*. However, after several months, I could prove that every prime and every even number appears, and that if p is an odd prime then there are infinitely many odd multiples of p (see [A280864](#) for details), but I could not prove that every odd number appears. The missing step feels like it is only a couple of cups of coffee away, and I’m hoping that some reader of this article will complete the proof.

The third example (Henry Bottomley, 2000, [A055265](#)) is the lexicographically earliest sequence of distinct positive integers such that $a(n-1) + a(n)$ is a prime for $n \geq 2$:

1, 2, 3, 4, 7, 6, 5, 8, 9, 10, 13, 16, 15,
14, 17, 12, 11, 18, 19, 22, 21, 20, ...

The terms appear to lie on or near the line $a(n) = n$, but the proof that every number appears may be difficult because it involves the gaps between the primes.

Two-Dimensional Lexicographically Earliest Arrays

The *OEIS* is primarily a database of sequences $(a_n, n \geq n_0)$. However, triangles of numbers are included by reading them by rows. Pascal’s triangle becomes 1, 1, 1, 1, 2, 1, 1, 3, 3, 1, 1, 4, 6, 4, 1, ... which (without the extra spaces) is [A007318](#). Doubly-indexed arrays $(T_{m,n}, m \geq m_0, n \geq n_0)$ are converted to sequences by reading them by antidiagonals (in either the upwards or downwards directions, or both). So an array $(T_{m,n}, m \geq 0, n \geq 0)$ might become $T_{0,0}, T_{1,0}, T_{0,1}, T_{2,0}, T_{1,1}, T_{0,2}, \dots$. For example, the table of Nim-sums $m \oplus n$:

(3)

0	1	2	3	4	5	6	7	...
1	0	3	2	5	4	7	6	...
2	3	0	1	6	7	4	5	...
3	2	1	0	7	6	5	4	...
4	5	6	7	0	1	2	3	...
5	4	7	6	1	0	3	2	...
6	7	4	5	2	3	0	1	...
7	6	5	4	3	2	1	0	...
.

produces the sequence [A003987](#):

0, 1, 1, 2, 0, 2, 3, 3, 3, 3, 4, 2, 0, 2, 4,
5, 5, 1, 1, 5, 5, 6, 4, 6, 0, 6, 4, 6, ...

Doubly-indexed doubly-infinite arrays $(T_{m,n}, m \in \mathbb{Z}, n \in \mathbb{Z})$ can become sequences by reading them in a spiral around the origin, in say a counter-clockwise direction: $T_{0,0}, T_{1,0}, T_{1,1}, T_{0,1}, T_{-1,1}, T_{-1,0}, T_{-1,-1}, T_{0,-1}, \dots$ (cf. Figure 13).

There are many “lexicographically earliest” versions of these arrays. For example, the Nim-sum array (3) has an equivalent definition: scan along upwards antidiagonals, filling in each cell with the smallest nonnegative number that is neither in the row to the left of that cell nor in the column above it.

A variation on the Nim-sum array was proposed by Alec Jones in 2016, as a kind of “infinite Sudoku array.” This array $(T_{m,n}, m \geq 0, n \geq 0)$ is to be filled in by upwards antidiagonals, always choosing the smallest positive integer such that no row, column, diagonal, or antidiagonal contains a repeated term. The top left corner of the array is:

(4)

1	3	2	6	4	5	10	11	...
2	4	5	1	8	3	6	12	...
3	1	6	2	9	7	5	4	...
4	2	3	5	1	8	9	7	...
5	7	1	4	2	6	3	15	...
6	8	9	7	5	10	4	16	...
7	5	4	3	6	14	8	9	...
8	6	7	9	11	4	13	3	...
.

The resulting sequence [\(A269526\)](#) is

1, 2, 3, 3, 4, 2, 4, 1, 5, 6, 5, 2, 6, 1, 4,
6, 7, 3, 2, 8, 5, 7, 8, 1, 5, 9, 3, 10, ...

This array has many interesting properties. If we subtract 1 from each entry, the entries are the Nim-values for a game played with two piles of counters, of sizes m and n , and reminiscent of Wythoff’s game (see [A004481](#), [A274528](#)).

But the main question about the array (4) is, are the individual rows, columns, and diagonals of this array permutations of \mathbb{N} ? (The antidiagonals are obviously not,

9	←	16	←	2	←	4	←	7	←	14	←	11	←	12	←	1	←	5	←	8
↓																				↑
17		8	←	15	←	14	←	13	←	12	←	9	←	10	←	6	←	7		3
↓	↓																	↑	↑	
1		2		4	←	11	←	10	←	3	←	8	←	7	←	9		13		15
↓	↓	↓														↑	↑	↑	↑	
8		9		7		3	←	5	←	6	←	1	←	2		4		12		11
↓	↓	↓	↓												↑	↑	↑	↑	↑	
11		12		8		1		2	←	4	←	3		6		5		10		14
↓	↓	↓	↓	↓								↑	↑	↑	↑	↑	↑	↑	↑	
15		7		6		5		3		1	→	2		4		8		11		12
↓	↓	↓	↓	↓	↓									↑	↑	↑	↑	↑	↑	
14		10		3		2		4	→	5	→	6	→	1		7		9		13
↓	↓	↓	↓	↓	↓										↑	↑	↑	↑	↑	
7		11		9		6	→	1	→	2	→	4	→	5	→	3		8		10
↓	↓	↓																↑	↑	
4		13		5	→	7	→	8	→	9	→	10	→	11	→	12	→	6		1
↓	↓	↓																	↑	
12		14	→	10	→	9	→	6	→	13	→	5	→	3	→	15	→	16	→	7
↓																				
10	→	15	→	1	→	12	→	16	→	8	→	14	→	13	→	11	→	18	→	17

Figure 13. [A274640](#): choose the smallest positive number so that no row, column, or diagonal contains a repeat. Are the rows, columns, diagonals permutations of \mathbb{N} ?

since they are finite sequences.) It is easy to see that each column is a permutation. In column $c \geq 0$, a number k will eventually be the smallest missing number and will appear in some cell in that column, unless there is a copy of k to the North-West, West, or South-West of that cell. But there are at most c copies of k in all the earlier columns, so eventually k will appear.

The rows are also permutations, although the proof is less obvious. Consider row $r \geq 0$, and suppose k never appears. There are at most r copies of k in the earlier rows, and these can affect only a bounded portion of row r . Consider a cell (r, n) , $n \geq 0$ large. If k is not to appear in that cell, there must be a copy of k in the antidiagonal to the South-West. So in the triangle bounded by row r , column 0, and the antidiagonal through (r, n) , there must be at least $n + 1 - r$ copies of k . Imagine these k s replaced by chess queens. By construction they are mutually nonattacking. But it is known ([6, Problem 252], or [A274616](#)) that on a triangular half-chessboard of side n , there can be at most $2n/3 + 1$ mutually nonattacking queens, which for large n leads to a contradiction.

As to the diagonals, although they appear to be permutations, this is an open question. The argument using nonattacking queens breaks down because the diagonal of the half-chessboard contains only half as many squares as the sides. Even the main diagonal, [A274318](#),

1, 4, 6, 5, 2, 10, 8, 3, 7, 9, 16, 26, 29,
22, 20, 23, 28, 38, 12, 32, 46, 13, 14, 11, 15, ... ,

is not presently known to be a permutation of \mathbb{N} .

The spiral version of this array is even more frustrating. This array $((T(m, n), m \in \mathbb{Z}, n \in \mathbb{Z}),$ [A274640](#), proposed by Zak Seidov and Kerry Mitchell in June 2016), is constructed in a counterclockwise spiral, filling in each cell with the smallest positive number such that no row, column, or diagonal contains a repeated term (Figures 13, 14). (“Diagonal” now means any line of cells of slope ± 1 .)

Although it seems very plausible that every row, column, and diagonal is a permutation of \mathbb{N} , now there are no proofs at all. The eight spokes through the center are sequences [A274924](#)–[A274931](#). For example, the row through the central cell is

..., 14, 25, 13, 17, 10, 15, 7, 6, 5, 3,
1, 2, 4, 8, 11, 12, 16, 9, 19, 24, 22, ... ,

which is [A274928](#) reversed followed by [A274924](#). Is it a permutation of \mathbb{N} ? We do not know.

Fun With Digits

Functions of the digits of numbers have always fascinated people,⁶ and one such function was in the news in 2017. The idea underlying this story and several related sequences is to start with some simple function $f(n)$ of

⁶Although in *A Mathematician's Apology*, G. H. Hardy, referring to the fact that 1089 and 2178 are the smallest numbers which when written backwards are nontrivial multiples of themselves (cf. [A008919](#)), remarked that this fact was “likely to amuse amateurs,” but was not of interest to mathematicians.




Figure 14. Colored representation of central 200×200 portion of the spiral in Figure 13: the colors represent the values, ranging from black (smallest) to white (largest).

the digits of n in some base, iterate it, and watch what happens.

For the first example we write n as a product of prime powers, $n = p_1^{e_1} p_2^{e_2} \dots$ with the p_i in increasing order, and define $f(n)$ to be the decimal concatenation $p_1 e_1 p_2 e_2 \dots$, where we omit any exponents e_i that are equal to 1. So $f(7) = f(7^1) = 7$, $f(8) = f(2^3) = 23$.

The initial values of $f(n)$ ([A080670](#)) are

n : 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 ...
 $f(n)$: 1 2 3 22 5 23 7 23 32 25 11 223 13 27 35 24 17 232 ...

If we start with a positive number n and repeatedly apply f , in many small cases we rapidly reach a prime (or 1).⁷ For example, $9 = 3^2 \rightarrow 32 = 2^5 \rightarrow 25 = 5^2 \rightarrow 52 = 2^2 13 \rightarrow 2213$, a prime. Define $F(n)$ to be the prime that is eventually reached, or -1 if the iteration never reaches a prime. The value -1 will occur if the iterates are unbounded or if they enter a cycle of composite numbers. The initial values of $F(n)$ ([A195264](#)) are

1, 2, 3, 211, 5, 23, 7, 23, 2213, 2213, 11,
223, 13, 311, 1129, 233, 17, 17137, 19,

$F(20)$ is currently unknown (after 110 steps the trajectory of twenty has stalled at a 192-digit number which has not yet been factored). At a DIMACS conference in October 2014 to celebrate the fiftieth anniversary of the start of what is now the OEIS, John H. Conway offered \$US1000

⁷In what follows we will tacitly assume $n \geq 2$, to avoid having to repeatedly say “(or 1).”

for a proof or disproof of his conjecture that the iteration of f will always reach a prime.

However, in June 2017 James Davis found a number $D_0 = 13532385396179$ whose prime factorization is $13 \cdot 53^2 \cdot 3853 \cdot 96179$, and so clearly $f(D_0) = D_0$ and $F(D_0) = -1$.

The method used by James Davis to find D_0 is quite simple. Suppose $n = m \cdot p$ is fixed by f , where p is a prime greater than all the prime factors of m . Then $f(n) = f(m)10^y + p$, where y is the number of digits in p . From $f(n) = n$ we have $p = \frac{f(m)10^y}{m-1}$. Assuming $p \neq 2, 5$, this implies that p divides $f(m)$, and setting $x = f(m)/p$, we find that $m = x10^y + 1$ with $p = \frac{f(m)}{x}$ prime. A computer easily finds the solution $x = 1407, y = 5, m = 140700001, p = 96179$, and so $n = D_0$.

No other composite fixed points are known, and David J. Seal has recently shown that there is no composite fixed point less than D_0 . It is easy, however, to find numbers whose trajectory under f ends at D_0 , by repeatedly finding a prime prefix of the previous number, as shown by the example⁸ $D_1 = 13^{532385396179}$ with $f(D_1) = D_0$. So presumably there are infinitely many n with $F(n) = -1$.

Consideration of the analogous questions in other bases might have suggested that counterexamples to Conway's question could exist. We will use subscripts to indicate the base (so $4_{10} = 100_2$). The base-2 analog of f , f_2 (say), is defined by taking $f_2(p_1^{e_1} p_2^{e_2} \dots)$ to be the concatenation $p_1 e_1 p_2 e_2 \dots$, as before (again omitting any e_i that are 1), except that now we write the p_i and e_i in base 2 and interpret the concatenation as a base-2 number. For example, $f_2(8) = f_2(2^3) = 1011_2 = 11_{10}$.

The initial values of $f_2(n)$ (A230625) are

n : 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 ...
 $f_2(n)$: 1 2 3 10 5 11 7 11 14 21 11 43 13 23 29 20 17 46 ...

and the base-2 analog of F , F_2 (A230627) is the prime (or 1) that is reached when f_2 is repeatedly applied to n , or -1 if no prime (or 1) is reached:

1, 2, 3, 31, 5, 11, 7, 11, 23, 31, 11, 43, 13, 23, 29, 251, 17, 23, ...

Now there is a fairly small composite fixed point, namely 255987, found by David J. Seal. Sean A. Irvine and Chai Wah Wu have also studied this sequence, and the present status is that $F_2(n)$ is known for all n less than 12388. All numbers in this range reach 1, a prime, the composite number 255987, or one of the two cycles $1007 \leftrightarrow 1269$ or $1503 \leftrightarrow 3751$. The numbers for which $F_2(n) = -1$ are 217, 255, 446, 558, ... (A288847). Initially it appeared that 234 might be on this list, but Irvine found that after 104 steps the trajectory reaches the 51-digit prime

350743229748317519260857777660944018966290406786641.

Home Primes

A rather older problem arises if we change the definition of $f(n)$ slightly, making $f(8) = 222$ rather than 23. So if $n = p_1 \cdot p_2 \cdot p_3 \cdot \dots$, where $p_1 \leq p_2 \leq p_3 \leq \dots$, then $f(n)$ is the decimal concatenation $p_1 p_2 p_3 \dots$ (A037276). In 1990, Jeffrey Heleen studied the analog of $F(n)$ for this

function: that is, $F(n)$ is the prime reached if we start with n and repeatedly apply f , or -1 if no prime is ever reached (A037274).

The trajectory of 8 now takes 14 steps to reach a prime (the individual prime factors here have been separated by spaces):

8 \rightarrow 2 2 2 \rightarrow 2 3 37 \rightarrow 3 19 41 \rightarrow 3 3 3 7 13 13 \rightarrow 3 11123771
 \rightarrow 7 149 317 941 \rightarrow 229 31219729 \rightarrow 11 2084656339
 \rightarrow 3 347 911 118189 \rightarrow 11 613 496501723 \rightarrow
 \rightarrow 97 130517 917327 \rightarrow 53 1832651281459
 \rightarrow 3 3 3 11 139 653 3863 5107
 \rightarrow 3331113965338635107,

the last number being a prime.

Since $f(n) > n$ if n is composite, now there cannot be any composite fixed points nor any cycles of length greater than 1. The only way for $F(n)$ to be -1 is for the trajectory of n to be unbounded. This appears to be a harder problem than the one in the previous section, since so far no trajectory has been proved to be unbounded. The first open case is $n = 49$, which after 119 iterations has reached a 251-digit composite number (see A056938). The completion of the factorization for step 117 took 765 days by the general number field sieve, and at the time (December 2014) was one of the hardest factorizations ever completed.

Power Trains

A third choice for $f(n)$ was proposed by John H. Conway in 2007: he called it the *power train* map. If the decimal expansion of n is $d_1 d_2 d_3 \dots d_k$ (with $0 \leq d_i \leq 9, 0 < d_1$), then $f(n) = d_1^{d_2} \cdot d_2^{d_3} \dots$, ending with $\dots \cdot d_k$ if k is odd, or with $\dots \cdot d_{k-1}^{d_k}$ if k is even (A133500). We take 0^0 to be 1. For example, $f(39) = 3^9 = 19683$, $f(623) = 6^2 \cdot 3 = 108$. Conway observed that $2592 = 2^{5 \cdot 9^2}$ is a nontrivial fixed point, and asked me if there were any others. I found one more: $n = 2^{46} \cdot 3^6 \cdot 5^{10} \cdot 7^2 = 24547284284866560000000000$, for which $f(n) = 2^4 \cdot 5^4 \cdot 7^2 \cdot 8^4 \cdot 2^8 \cdot 4^8 \cdot 6^6 \cdot 5^6 \cdot 0^0 \cdot 0^0 \cdot 0^0 \cdot 0^0 = n$. The eleven known fixed points (including the trivial values 1, ..., 9) form A135385, and it is known that there are no further terms below 10^{100} . Maybe this is a hint that for all of the functions $f(n)$ that have just been mentioned, there may be only a handful of genuinely exceptional values?

A Memorable Prime

If you happen to need an explicit 20-digit prime in a hurry, it is useful to remember that although $1,121 = 11^2, 12321 = 111^2, 1234321 = 1111^2, \dots$, and $12345678987654321 = 11111111^2$ are not primes, the next term in A173426 is a prime,

12345678910987654321.

As David Broadhurst remarked on the Number Theory Mailing List in August 2015, this is a memorable prime! He also pointed out that on probabilistic grounds, there should be infinitely many values of n such that the decimal

⁸Found by Hans Havermann.

concatenation of the numbers 1 up through n followed by $n - 1$ down through 1 is a prime. Shortly afterwards, Shyam Sunder Gupta found what is presumably the next prime in the sequence, corresponding to $n = 2446$, the 17350-digit probable prime 1234567..244524462445..7654321. Serge Batalov has shown that there are no further terms with $n < 60000$. What is the next term? The values 10, 2446 are not enough to create an *OEIS* entry.

A Missing Prime

The previous question naturally led me to wonder what the first prime is in the simpler sequence (A007908):

1, 12, 123, 1234, ..., 12345678910, 1234567891011, ... ,

formed by the decimal concatenation of the numbers 1 through n . In *Unsolved Problems in Number Theory*, Richard K. Guy reports that this question was already asked by Charles Nicol and John Selfridge. However, although the same probabilistic argument suggests that there should be an infinite number of primes of this type, not a single one is known. I asked several friends to help with the search, and as a result this task was taken up by the folks who run the GIMP (or Great Internet Mersenne Prime) search, and there is now a webpage⁹ that shows the current status of the search for the first prime. As of August 2017 the search seems to have stalled, the present status being that all the potential values of n through 344869 failed (obviously many values of n can be ruled out by congruence conditions). In this range the candidates have about two million digits. One estimate suggests that there is a probability of about 0.5 that a prime will be found with $n < 10^6$, so it would be good to resume this search.

Post's Tag System

In his recent book *Elements of Mathematics: From Euclid to Gödel*¹⁰ John Stillwell mentions that Emil L. Post's tag system from the 1930s is still not understood. Post asked the following question. Take a finite string, or word, S of 0s and 1s, and if it begins with 0, append 00 to the end of S and delete the first three symbols, or if it begins with 1, append 1101 to the end of S and delete the first three symbols. When this process is iterated, eventually one of three things will happen: either S will reach the empty word (S dies), S will enter a loop (S cycles), or S will keep growing for ever (S blows up). For example, $S = 1000$ reaches the empty word ϵ at the 7th step:

$$1000 \rightarrow 01101 \rightarrow 0100 \rightarrow 000 \rightarrow 00 \rightarrow 0 \rightarrow \epsilon,$$

whereas 100100 enters a cycle of length six (indicated by parentheses) after 15 steps:

⁹mersenneforum.org/showthread.php?t=20527.

¹⁰A superb successor to Felix Klein's 1908, *Elementary Mathematics from an Advanced Standpoint*.

(5)

$$\begin{aligned} 100100 &\rightarrow 1001101 \rightarrow 11011101 \rightarrow 111011101 \\ &\rightarrow 0111011101 \rightarrow 101110100 \rightarrow 1101001101 \\ &\rightarrow 10011011101 \rightarrow 110111011101 \\ &\rightarrow 1110111011101 \rightarrow 01110111011101 \\ &\rightarrow 1011101110100 \rightarrow 11011101001101 \\ &\rightarrow 111010011011101 \rightarrow 0100110111011101 \\ &\rightarrow (011011101110100 \rightarrow 01110111010000 \\ &\rightarrow 1011101000000 \rightarrow 11010000001101 \\ &\rightarrow 100000011011101 \rightarrow 0000110111011101). \end{aligned}$$

Post was hoping to find an algorithm which, given S , would determine which of these outcomes would occur. He did not succeed.

Post called this process a 'tag system.' It can be generalized by considering initial words over an alphabet of size M (rather than 2), allowing any fixed set \mathcal{A} of M tag words to be appended (rather than 00 and 1101), and deleting some fixed number P of initial symbols at each step (not necessarily 3). In 1961, Marvin Minsky showed that such a generalized tag system could simulate a Turing machine. By choosing an appropriate alphabet, an appropriate set \mathcal{A} of tag words to be appended, and an appropriate value of P (in fact $P = 2$ will do), any computable function can be simulated. So, because of the undecidability of the Halting Problem, for general tag systems it is impossible to predict which initial words will blow up.

But what about Post's original tag system? Could this simulate a Turing machine (by encoding the problem in the initial word S)? At first this seems very unlikely, but the Cook-Wolfram theorem that the one-dimensional cellular automaton defined by Rule 110 can simulate a Turing machine (by encoding the problem in the starting state) suggests that it might be possible. If it is possible, there must be some initial words that blow up (again because of the Halting Problem).

In early 2017, when I read Stillwell's book, the *OEIS* contained three sequences related to the original tag system, based on the work of Peter Asveld and submitted by Jeffrey Shallit: A284116, giving the maximal number of words in the 'trajectory' of any initial word S of length n (18 terms were known), and two sequences connected with the especially interesting starting word σ_n of length $3n$ consisting of n copies of 100. A284119(n) is defined to be the number of words in the trajectory of σ_n before it enters a cycle or dies, or -1 if the trajectory blows up, and A284121(n) is the length of the cycle, or 1 if the trajectory dies, or -1 if the trajectory blows up. For example, from (5) we see that A284119(2) = 15 and A284121(2) = 6. Shallit had extended Asveld's work and had found 43 terms of the two last-mentioned sequences.

I then added many further sequences based on tag systems discussed by Asveld, Liesbeth De Mol, Shigeru Watanabe, and others, and appealed to contributors to the *OEIS* to extend them.

The most interesting response came from Lars Blomberg, who investigated the trajectory of σ_n for

$n \leq 110$. On September 9, 2017, he reported that every σ_n for $n \leq 110$ had either died or cycled after at most 13 million terms, except for σ_{110} , which after 38.10^{11} steps had reached a word of length 10^7 and was still growing. This was exciting news! Could σ_{110} be the first word to be discovered that blew up?¹¹ Sadly, on October 4, 2017, Blomberg reported that after 43913328040672 steps σ_{110} had terminated in the empty word.

Figure 15 displays the remarkable graph (technically, a pin plot) of the number of steps for σ_n to either die or cycle for $n \leq 200$. Figure 16 shows the lengths of the successive words in the trajectory of σ_{110} .




Figure 15. Pin plot illustrating Lars Blomberg’s remarkable discovery that the Post tag system started at the word $(100)^{110}$ takes an exceptionally long time (43913328040672 steps) to converge.




Figure 16. Lengths of successive words in trajectory of $(100)^{110}$ under the Post tag system. The numbers on the horizontal axis are spaced at multiples of 10^{12} .

In the past six months Blomberg has continued this investigation and has determined the fate of σ_n for all $n \leq 6075$. The new record-holder for the number of steps before the trajectory dies is now held by σ_{4974} , which takes 57042251906801 steps, while σ_{110} is in second place.

¹¹Of course the fact that the same number 110 was involved could not possibly be anything more than a coincidence.

Of course it is still possible that some initial word S , not necessarily of the form σ_n , will blow up, but this seems increasingly unlikely. So Post’s tag system probably does not simulate a Turing machine.

The question as to which σ_n die and which cycle remains a mystery. Up to $n = 6075$, Blomberg’s results show that about one-sixth of the values of n die and five-sixths cycle. The precise values can be found in [A291792](#). It would be nice to understand this sequence better.

Coordination Sequences

This final section is concerned with *coordination sequences*, which arise in crystallography and in studying tiling problems, have beautiful illustrations, and lead to many unsolved mathematical questions.




Figure 17. A portion of the Cairo tiling.

The “Cairo” tiling, so called because it is said to be used on many streets in that city, is shown in Figure 17. Let G denote the corresponding infinite graph (with vertices for points where three or more tiles meet, and edges between two vertices where two tiles meet). The figure is also a picture of the graph.

The distance between vertices $P, Q \in G$ is defined to be the number of edges in the shortest path joining them. The *coordination sequence* of G with respect to a vertex $P \in G$ is then the sequence $a(n)$ ($n \geq 0$) giving the number of vertices Q at distance n from P . Coordination sequences have been studied by crystallographers for many years [5].

The graph of the Cairo tiling has two kinds of vertices, trivalent (where three edges meet) and tetravalent. As can be seen from Figure 17, the coordination sequence with respect to a tetravalent vertex begins 1, 4, 8, 12, 16, 20, 24, ..., which appears to be the same as the coordination sequence [A008574](#) for a vertex in the familiar square grid. This observation seemed to be new. Chaim Goodman-Strauss and I thought that such a simple fact should have a simple proof, and we developed an elementary “coloring book” procedure [4] which not only proved this result but also established a number of conjectured formulas for coordination sequences of other tilings mentioned in entries in the *OEIS*. The “coloring book” drawing of the Cairo graph centered at a tetravalent vertex is shown in Figure 18. This coloring makes it easy to prove that the coordination sequence is given by $a(n) = 4n$ for $n \geq 1$ (see [4] for details).




Figure 18. The “coloring book” method applied to a tetravalent vertex (the red dot) in the Cairo tiling, used to prove that the coordination sequence is the same as that for the square grid.

For a trivalent vertex in the Cairo tiling, the coordination sequence is

1, 3, 8, 12, 15, 20, 25, 28, 31, 36,
41, 44, 47, 52, 57, 60, 63, 68, ...

(this is now [A296368](#)), and we [4] show that for $n \geq 3$, $a(n) = 4n$ if n is odd, $4n - 1$ if $n \equiv 0 \pmod{4}$, and $4n + 1$ if $n \equiv 2 \pmod{4}$.

One can similarly define coordination sequences for other two- and higher-dimensional structures, and the *OEIS* presently contains over 7000 such sequences (mostly without formulas). Many more could be added. There are many excellent websites with lists of tilings and crystals. Brian Galebach’s website¹² is especially important, as it includes pictures of all “ k -uniform” tilings with $k \leq 6$, with over 1000 tilings. Darrah Chavey’s article [2] and the Michael Hartley and Printable Paper websites¹³ have many further pictures, and the RCSR and ToposPro databases¹⁴ have thousands more.

Only last week (on May 4, 2018), Rémy Sigrist investigated the Ammann-Beenker (or “octagonal”) tiling shown in Figure 19, an aperiodic tiling with eight-fold rotational symmetry about the central point.




Figure 19. Ammann-Beenker (or “octagonal”) tiling.

Sigrist determined the initial terms of the coordination sequence with respect to the central vertex ([A303981](#)):

(6) 1, 8, 16, 32, 32, 40, 48, 72, 64, 96, 80,
104, 112, 112, 128, 152, ...

Figure 20 shows the vertices at distances 0, 1, 2, ..., 6 from the center.




Figure 20. Illustrating the coordination sequence for the Ammann-Beenker tiling, showing the vertices at distances 0 through 6 from the central vertex.

¹²probabilitysportsports.com/tilings.html.

¹³www.dr-mikes-math-games-for-kids.com/archimedean-graph-paper.html, <https://www.printablepaper.net/category/graph>.

¹⁴rcsr.net, topospro.com.

No formula or growth estimate is presently known for this sequence. However, earlier this year Anton Shutov and Andrey Maleev determined the asymptotic behavior of the coordination sequence (A302176) with respect to a vertex with five-fold rotational symmetry in a certain Penrose tiling. So we end with a question: Can the Shutov-Maleev approach be used to find the asymptotic growth of (6)? Of course an explicit formula would be even nicer.

Acknowledgments

Now that the *OEIS* is a wiki, many volunteer editors help maintain it.¹⁵ Besides the *OEIS* contributors already mentioned in this article, I would like to thank Jörg Arndt, Michael De Vlieger, Charles Greathouse IV, Alois P. Heinz, Michel Marcus, Richard J. Mathar, Don Reble, Jon E. Schoenfeld, and Allan C. Wechsler for their help. (Many other names could be added to this list.)

References

- [1] D. L. APPLGATE, H. HAVERMANN, B. SELCOE, V. SHEVELEV, N. J. A. SLOANE, and R. ZUMKELLER, The Yellowstone Permutation, *J. Integer Seqs.*, **18** (2015), #15.6.7. [MR3360900](#)
- [2] D. CHAVEY, Tilings by regular polygons II: A catalog of tilings, *Computers & Mathematics with Applications*, **17**:1–3 (1989), 147–165. [MR994197](#)
- [3] P. ERDŐS, R. FREUD, and N. HEGYVÁRI, Arithmetical properties of permutations of integers, *Acta Math. Hungar.*, **41** (1983), 169–176. [MR704537](#)
- [4] C. GOODMAN-STRAUSS and N. J. A. SLOANE, A coloring book approach to finding coordination sequences, *Acta. Cryst. A*, to appear: arXiv:1803.08530, 2018.
- [5] M. O'KEEFE and B. G. HYDE, Plane nets in crystal chemistry, *Phil. Trans. Royal Soc. London, Series A, Mathematical and Physical Sciences*, **295**:1417 (1980), 553–618. [MR569155](#)
- [6] P. VANDERLIND, R. K. GUY, and L. C. LARSON, *The Inquisitive Problem Solver*, MAA Press, 2002. [MR1917371](#)

Image Credits

Figures 1–5, 10, 11, 13, 15 courtesy of Michael De Vlieger (who also designed the “Peace to the Max” T-shirt in Figure 3).

Figure 6 courtesy of Jessica Gonzalez.

Figures 7–9 courtesy of Jonathan Wild.

Figure 12 courtesy of Hans Havermann.

Figure 14 courtesy of Kerry Mitchell.

Figure 16 courtesy of Lars Blomberg.

Figures 17 and 18 courtesy of Chaim Goodman-Strauss.

Figure 19 from The Tilings Encyclopedia, <https://tilings.math.uni-bielefeld.de>. Used under the Creative Commons Attribution-

NonCommercial-ShareAlike 2.0 Generic License.

Figure 20 courtesy of Rémy Sigris.

Author photo courtesy of Susanna Stevens Cuyler.



Neil J. A. Sloane

ABOUT THE AUTHOR

Neil J. A. Sloane started collecting sequences in 1964. Every advance in computing since then has made the task easier; an apt subtitle to this article could be “From punched cards to wiki in fifty-four years.” The Unix operating system was the greatest breakthrough of all, especially grep, the programmable editor ex, and pipes.

¹⁵And more are needed—if interested, please contact the author.