

The Ordovician history of the Iapetus Ocean in Britain: new palaeomagnetic constraints

T. H. TORSVIK and A. TRENCH

Journal of the Geological Society 1991; v. 148; p. 423-425

doi: 10.1144/gsjgs.148.3.0423

Email alerting service

click [here](#) to receive free e-mail alerts when new articles cite this article

Permission request

click [here](#) to seek permission to re-use all or part of this article

Subscribe

click [here](#) to subscribe to Journal of the Geological Society or the Lyell Collection

Notes

Downloaded by on January 23, 2012

SHORT PAPER

The Ordovician history of the Iapetus Ocean in Britain: new palaeomagnetic constraints

T. H. TORSVIK¹ & A. TRENCH²

Department of Earth Sciences, University of Oxford, Oxford, OX1 3PR, UK

¹ *Present address: Geological Survey of Norway, P.B. 3006 Lade N-7002 Trondheim, Norway*

² *Present address: Dept of Geology, University of Western Australia, Nedlands, Perth, WA 6009, Australia*

New late Tremadoc–early Arenig palaeomagnetic results from SW Wales imply that S Britain (part of Eastern Avalonia) occupied a southerly latitude of *c.* 60°S in early Ordovician times. When combined with Scottish Ordovician palaeomagnetic data, which indicate a 15°S latitude, the results indicate that the British sector of the Iapetus Ocean reached a latitudinal width of *c.* 5000 km in Tremadoc–Arenig times, which was reduced to *c.* 3300 km by Llanvirn–Llandeilo (mid-Ordovician) times.

The new data resolve two previous controversies in Palaeozoic palaeogeography. First, the high southerly palaeolatitude links Avalonia to Gondwana, marginal to W Africa, thus reconciling conflicting reconstructions based upon either palaeomagnetic or faunal/facies evidence alone. Second, reliable Llanvirn palaeomagnetic data imply that Avalonia had rifted northwards by Arenig time, whereas Armorica remained proximal to northern Africa throughout the Ordovician. The combined data therefore establish that Avalonia and Armorica formed separate micro-continents when rifting from Gondwana.

Palaeomagnetic, biogeographic and palaeoclimatic evidence are in agreement in establishing the existence of the Iapetus Ocean (Wilson 1966; Harland & Gayer 1972) in Ordovician times, but estimates of its size have varied substantially (Dewey 1969, 1971; McKerrow & Cocks 1976; Smith *et al.* 1981; Briden *et al.* 1984; McKerrow 1988; Torsvik *et al.* 1990a; Scotese & McKerrow 1990). In particular, biogeographic considerations have consistently been interpreted to represent wider oceanic separation than that indicated by palaeomagnetic data. This discrepancy is explicable in that early palaeomagnetic studies may not have employed sufficiently detailed demagnetization experiments to remove Late Palaeozoic secondary magnetizations of shallow inclination (see Briden & Mullan 1984, for discussion).

Recently, however, a palaeomagnetic result from the Shelve inlier (McCabe & Channell 1990), as well as a re-compilation of palaeomagnetic data from S Britain (Trench & Torsvik 1991), indicates that the British sector of the Iapetus Ocean covered approximately 30° of latitude in mid-Ordovician (Llanvirn–Llandeilo) times (N Britain 15°S, S Britain 45°S). Prior to the Llanvirn however, no reliable palaeomagnetic data exist for Southern Britain.

In early Ordovician times (Tremadoc–Arenig), faunal evidence indicates Iapetus to have reached its maximum extent (McKerrow & Cocks 1976, 1986), and early Ordovician sedimentary facies suggest a provenance link with the Armorican Massif (Fortey & Owens 1987; Noblet *et*

al. 1990), from which palaeomagnetic data indicate high southerly latitudes (Perroud & Van der Voo 1985; Perroud *et al.* 1986). Therefore, although several palaeogeographic reconstructions depict S Britain as a peri-Gondwanan block in early Ordovician time (Pickering *et al.* 1988; Scotese & McKerrow 1990), no palaeomagnetic data have been available to quantify its palaeo-position. Similarly, palaeomagnetic data have been insufficient to discriminate whether S Britain (part of Eastern Avalonia; Soper *et al.* 1987; McKerrow 1988) formed a constituent of the Armorican plate (Armorican and Iberian Massifs: Van der Voo 1979; Perroud *et al.* 1984) during the Ordovician period.

To address these uncertainties, 15 sites were sampled within a sequence of late Tremadoc–early Arenig andesitic lavas and volcanoclastic sediments at Treffgarne, Dyfed, SW Wales (Traynor 1988). This sequence dips on average 30° to the north. Low-grade metamorphism (Oliver 1988) and tectonism is most likely Acadian (Woodcock *et al.* 1988), although Hercynian deformation cannot be excluded (Hancock *et al.* 1981).

Palaeomagnetic results. Stepwise-thermal demagnetization experiments (161 samples) identify two magnetization components. The lower unblocking-temperature component (200–500°C) has sub-horizontal to shallow negative inclination due south. We interpret this component, termed *S*, as the Permo-Carboniferous (Hercynian) overprint observed elsewhere in Wales (McClelland-Brown 1983; Briden & Mullan 1984; McCabe & Channell 1990). The higher unblocking-temperature component (>500°C, Fig. 1a), termed *T*, is primary, and has the following remanence properties.

(1) Reversely-magnetized polarity (tilt-corrected declination = 298°, inclination = 75°, $\alpha_{95} = 5.5^\circ$, observed in 10 sites; Fig. 1b: Pole N56°, E306° & $dp/dm = 9/10$).

(2) Clasts from an intra-formational conglomerate (Thomas & Cox 1924) reveal high-unblocking magnetizations which are directionally-consistent within, but differ between, individual boulders (Fig. 1c). These magnetizations have equivalent unblocking temperatures to component *T*. Conversely, component *S* displays consistent directions within the boulders.

(3) Laboratory unblocking temperatures cover both magnetite and hematite ranges, but generally display discrete unblocking between 520–550°C (Fig. 1a), which indicates the dominance of low-Ti titanomagnetite formed during high-temperature deuteric oxidation in volcanic rocks.

Discussion. The new palaeomagnetic data imply that Southern Britain occupied a latitude of *c.* 60°S in late Tremadoc–early Arenig times (Fig. 2). Compared with Ordovician data from the Scottish terranes (*c.* 15°S, Torsvik *et al.* 1990a), this implies that the intervening Iapetus Ocean reached a latitudinal width of near 5000 km (Fig. 2).

A comparison with Ordovician palaeomagnetic data from Gondwana (Van der Voo 1988; Bachtadse & Briden 1990), which allow the reconstruction of the west Gondwanan margin, suggests Eastern Avalonia to have been positioned close to west Africa, possibly in the vicinity of Mauritania (Fig. 2) while Armorica was attached to NW Africa (Torsvik *et al.* 1990a). This scenario attributes late




Fig. 1. (a) Representative orthogonal projection: Andesite lava. Primary (*T*) magnetization component is indicated. Open (Closed) squares refer to projection in the vertical (horizontal) plane. In-Situ co-ordinates. Component *S* not resolved in this example.

(b) Site-mean component (*T*) directions after structural tilt correction. Cone of 95% confidence about the total mean direction is dotted. Equal-angle projection.

(c) Clast-mean component (*T*) directions from an intra-formational volcanic conglomerate. Cones of 95% confidence are indicated for each clast. Number of samples from each clast is shown. In (b) and (c), solid (open) squares represent lower (upper) hemisphere directions.

Precambrian arc rocks in both Avalonia and Armorica (e.g. Scotese & McKerrow 1990) to a peripheral Gondwanan arc. At this time, Baltica was confined to southerly latitudes of 30–60° (Torsvik *et al.* 1990b), whereas Laurentia and Siberia (Mongolian margin facing north; Scotese & McKerrow 1990; Khramov *et al.* 1981) retained more equatorial latitudes (Fig. 2).

From this early Ordovician scenario, Eastern Avalonia then had rifted away from Gondwana into mid-southerly latitudes by mid-Ordovician (late Llanvirn–early Llandeilo) times (Fig. 3). This movement history is clearly recorded by palaeomagnetic data from volcanic rocks of the early Llanvirn Shelf (51°S; McCabe & Channell 1990) and the late Llanvirn Builth inliers (35°S; Briden & Mullan 1984; Trench *et al.* 1991) respectively. Conversely, the African Gondwanan margin and Armorica remained in high




Fig. 2. Continental reconstruction for early Ordovician (late Tremadoc–early Arenig) times based on palaeomagnetic data (optional longitude). Southern Britain and other Avalonian segments positioned in *latitude* according to the Treffgarne pole. A preferred palaeo-orientation has been used to maintain the facing of the Iapetus margin in Britain. In Figs 2 & 3, Western Avalonia (WA) is positioned according to data from Eastern Avalonia (EA) in a Bullard *et al.* fit (1965), since no structurally reliable early and mid-Ordovician data exist for Western Avalonia. Gondwana positioned according to Van der Voo (1988) (pole: 34°N, 007°E, African co-ordinates), Laurentia and Northern Britain combined (Torsvik *et al.* 1990b, pole: 13°S, 29°E, European co-ordinates in a Bullard *et al.* fit (1965)), Baltica (Torsvik *et al.* 1990b, pole: 31°N, 086°E), Armorica (A) (Torsvik *et al.* 1990a, table 7, 490 Ma pole: 30°N, 334°E), Siberia according to a mean pole of 30°N, 330°E (Torsvik *et al.* 1990a). Equal-area polar projection.




Fig. 3. Continental reconstruction for mid-Ordovician (late Llanvirn–early Llandeilo) times. Latitudinal positioning of Gondwana and Siberia as Fig. 2. Avalonia from Trench & Torsvik (1991, table 2, 470 Ma pole: 12°N, 23°E), Baltica (Torsvik *et al.* 1990b, combined path X and Y, 470 Ma pole: 21°N, 32°E), Armorica (Torsvik *et al.* 1990a, table 7, 470 Ma pole: 33°N, 345°E), Laurentia and Northern Britain (Torsvik *et al.* 1990b, pole: 22°S, 19°E, European co-ordinates). Equal-area polar projection.

southerly latitudes throughout the Ordovician period as implied by palaeomagnetic data (Perroud & Van der Voo 1985; Perroud *et al.* 1986; Van der Voo 1988; Bachtadse & Briden 1990) and the presence of Ordovician tillites (Scotese & Barrett 1990).

Conclusions.

(1) Avalonia, Armorica (Armorican–Iberian Massifs) and probably other European Massifs formed marginal parts of Gondwana in the vicinity of NW Africa in early Ordovician times, separated by the Iapetus Ocean and the Tornquist Sea from Laurentia and Baltica respectively,

(2) Latitudinal separation across the British sector of Iapetus was approximately 5000 km in late Tremadoc–early Arenig times, reduced to *c.* 3300 km by Llanvirn–Llandeilo times,

(3) Avalonia rifted away from Gondwana late in the early Ordovician, Arenig, and later collided with Baltica and Laurentia, ultimately forming Euramerica by late Silurian–early Devonian times. Conversely, Armorica remained attached to Gondwana until at least late Ordovician–early Silurian times.

A.T. held a Natural Environment Research Council Fellowship at Oxford. T.H.T. acknowledges support from the Norwegian Research Council for Science and Humanities. We also thank W. S. McKerrow and J.-J. Traynor. Norwegian International Lithosphere Contribution (126).

References

- BACHTADSE, V. & BRIDEN, J. C. 1990. Palaeomagnetic constraints on the position of Gondwana during Ordovician to Devonian times. In McKERROW, W. S. & SCOTESE, C. R. (eds) *Palaeozoic Palaeogeography and Biogeography*, Geological Society London Memoir, **12**, 43–48.
- BRIDEN, J. C. & MULLAN, A. J. 1984. Superimposed Recent, Permian–Carboniferous and Ordovician palaeomagnetic remanence in the Builth volcanic Series, Wales. *Earth and Planetary Science Letters*, **69**, 413–421.
- BRIDEN, J. C., TURNELL, H. B. & WATTS, D. R. 1984. British palaeomagnetism, Iapetus Ocean, and the Great Glen Fault. *Geology*, **12**, 428–431.
- BULLARD, E. C., EVERETT, J. E. & SMITH, A. G. 1965. The fit of the continents around the Atlantic. *Royal Society of London, Philosophical Transactions Series*, **A258**, 41–51.
- COCKS, L. R. M. & FORTEY, R. A. 1982. Faunal evidence for oceanic separations in the Palaeozoic of Britain. *Journal of the Geological Society, London*, **139**, 465–478.
- DEWEY, J. F. 1969. Evolution of the Appalachian–Caledonian orogen. *Nature*, **222**, 124–8.
- 1971. A model for the Lower Palaeozoic evolution of the southern margin of the early Caledonides in Scotland and Ireland. *Scottish Journal of Geology*, **7**, 219–40.
- FORTEY, R. A. & OWENS, R. M. 1987. The Arenig in South Wales. *Bulletin of the British Museum of Natural History (Geology)*, **41**, 3, 69–307.
- HANCOCK, P. L., DUNNE, W. M. & TRINGHAM, M. E. 1981. Variscan structures in southwest Wales. *Geologie en Mijnbouw*, **60**, 81–88.
- HARLAND, W. B., & GAYER, R. A. 1972. The Arctic Caledonides and earlier Oceans. *Geological Magazine*, **109**, 283–314.
- KHRAMOV, A. N., PETROVA, G. N. & PECHERSKY, D. M. 1981. Palaeomagnetism of the Soviet Union. In: McELHINNY, M. & VALENCIO, D. A. (eds) *Palaeoreconstructions of the Continents*. Geodynamic Series, **2**, American Geophysical Union, 177–194.
- MCCABE, C. & CHANNELL, J. E. T. 1990. Palaeomagnetic results from volcanic rocks of the Shelve inlier, Wales: evidence for a wide Late Ordovician Iapetus Ocean in Britain. *Earth and Planetary Science Letters*, **96**, 458–468.
- MCCLELLAND-BROWN, E. A. 1983. Palaeomagnetic studies of fold development in the Pembroke Old Red Sandstone. *Tectonophysics*, **98**, 131–149.
- MCKERROW, W. S. 1988. The development of the Iapetus Ocean from the Arenig to the Wenlock. In: HARRIS, A. L. & FETTES, D. J. (eds) *The Caledonian–Appalachian Orogen*. Geological Society, London, Special Publication, **38**, 405–415.
- & COCKS, L. R. M. 1976. Progressive faunal migration across the Iapetus Ocean. *Nature*, **267**, 237–239.
- & — 1986. Oceans, island arc and olistostromes: the use of fossils in distinguishing sutures, terranes and environments around the Iapetus Ocean. *Journal of the Geological Society, London*, **143**, 185–191.
- NOBLET, CH. & LEFORT, J. P. 1990. Sedimentological evidence for a limited separation between Armonica and Gondwana during the Early Ordovician. *Geology*, **18**, 303–306.
- OLIVER, G. J. H. 1988. Arenig to Wenlock regional metamorphism in the Paratectonic Caledonides of the British Isles: a review. In: HARRIS, A. L. & FETTES, D. J. (eds) *The Caledonian–Appalachian Orogen*. Geological Society, London, Special Publication, **38**, 347–363.
- PERROUD, H. & VAN DER VOO, R., 1985. Palaeomagnetism of the late Ordovician Thouars Massif, Vendée Province, France. *Journal of Geophysical Research*, **90**, 4611–4625.
- , BONHOMMET, N. & THEBAULT, J. P. 1986. Palaeomagnetism of the Ordovician Moulin de Chateaupanne formation, Vendée, western France. *Geophysical Journal of the Royal Astronomical Society*, **85**, 573–582.
- , VAN DER VOO, R. & BONHOMMET, N. 1984. Palaeozoic evolution of the Armorican plate on the basis of palaeomagnetic data. *Geology*, **12**, 579–582.
- PICKERING, K. T., BASSETT, M. G. & SIVETER, D. J. 1988. Late Ordovician–early Silurian destruction of the Iapetus Ocean: Newfoundland, British Isles and Scandinavia—a discussion. *Transactions of the Royal Society of Edinburgh Earth Sciences*, **79**, 361–382.
- SCOTESE, C. R. & BARRETT, S. F. 1990. Gondwana's movement over the South Pole during the Palaeozoic: evidence from lithological indicators of climate. In: McKERROW, W. S. & SCOTESE, C. R. (eds) *Palaeozoic Palaeogeography and Biogeography*. Geological Society, London, Memoir, **12**, 75–85.
- & McKERROW, W. S. 1990. Revised World maps and introduction. In: McKERROW, W. S. & SCOTESE, C. R. (eds) *Palaeozoic Palaeogeography and Biogeography*. Geological Society, London, Memoir, **12**, 1–21.
- SMITH, A. G., HURLEY, A. M. & BRIDEN, J. C. 1981. *Phanerozoic Palaeo-continental World Maps*. Cambridge University Press.
- SOPER, N. J., WEBB, B. C. & WOODCOCK, N. H. 1987. Late Caledonian (Acadian) transpression in north-west England: Timing, geometry and geotectonic significance. *Proceedings of the Yorkshire Geological Society*, **46**, 175–192.
- THOMAS, H. H. & COX, A. H. 1924. The volcanic series of Trefgarn, Roch and Ambleston. *Quarterly Journal of the Geological Society of London*, **80**, 520–548.
- TORSVIK, T. H., SMETHURST, M. A., BRIDEN, J. C. & STURT, B. A., 1990a. A review of Palaeozoic palaeomagnetic data from Europe and their palaeogeographical implications. In: McKERROW, W. S. & SCOTESE, C. R. (eds) *Palaeozoic Palaeogeography and Biogeography*, Geological Society London Memoir, **12**, 25–41.
- , OLESEN, O., RYAN, P. D. & TRENCH, A. 1990b. On the palaeogeography of Baltica during the Palaeozoic: New palaeomagnetic data from the Scandinavian Caledonides. *Geophysical Journal International*, **103**, 261–279.
- TRAYNOR, J.-J. 1988. The Arenig in South Wales: Sedimentary and volcanic processes during the initiation of a marginal basin. *Geological Journal*, **23**, 275–292.
- TRENCH, A. & TORSVIK, T. H. 1991. A revised Palaeozoic apparent polar wander path for Southern Britain (Eastern Avalonia). *Geophysical Journal International*, **104**, 227–233.
- , —, SMETHURST, M. A., WOODCOCK, N. H. & METCALFE, R. 1991. A palaeomagnetic study of the Builth Wells/Llandrindod Wells Ordovician Inlier, Wales. *Geophysical Journal International* (in press).
- VAN DER VOO, R. 1979. Palaeozoic assembly of Pangea: a new plate tectonic model for the Taconic, Caledonian and Hercynian orogenies. *EOS, Transactions of AGU*, **60**, 241.
- 1988. Palaeozoic Palaeogeography of North America, Gondwana, and intervening displaced terranes: Comparison of palaeomagnetism with palaeoclimatology and biogeographical patterns. *Geological Society of America Bulletin*, **100**, 311–324.
- WILSON, J. T. 1966. Did the Atlantic close and then reopen?. *Nature*, **211**, 676–679.
- WOODCOCK, N. H., AWAN, M. A., JOHNSON, T. E., MACKIE, A. H. & SMITH, R. D. A. 1988. Acadian tectonics of Wales during Avalonia/Laurentia convergence. *Tectonics*, **7**, 483–495.