

 Open access • Journal Article • DOI:10.1112/S002557930001069X

The p -periodicity of the groups $GL(n, \mathcal{O}_S(K))$ and $SL(n, \mathcal{O}_S(K))$

— [Source link](#) 

B. Bürgisser, B. Eckmann

Published on: 01 Jun 1984 - Mathematika (London Mathematical Society)

Related papers:

- [On the Stability and Gelfand Property of Symmetric Pairs](#)
- [SL2 over group rings of cyclic groups](#)
- [The homotopy types of SU\(n\)-gauge groups over \$S_6\$](#)
- [Homological vanishing for the Steinberg representation](#)
- [Criterion for the Equality of Norm Groups of Idele Groups of Algebraic Number Fields](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/the-p-periodicity-of-the-groups-gl-n-o-s-k-and-sl-n-o-s-k-3glo5kuy8h>

The p-periodicity of the groups $GL(n, \mathcal{O}_s(K))$ and $SL(n, \mathcal{O}_s(K))$

Journal Article**Author(s):**

Bürgisser, B.; Eckmann, B.

Publication date:

1984

Permanent link:

<https://doi.org/10.3929/ethz-b-000422878>

Rights / license:

[In Copyright - Non-Commercial Use Permitted](#)

Originally published in:

Mathematika 31(1), <https://doi.org/10.1112/S002557930001069X>

THE p -PERIODICITY OF THE GROUPS $\mathrm{GL}(n, \mathcal{O}_S(K))$ AND $\mathrm{SL}(n, \mathcal{O}_S(K))$

B. BÜRGISSER AND B. ECKMANN

§1. *Introduction.* 1.1. In this paper we investigate the p -periodicity of the S -arithmetic groups $G = \mathrm{GL}(n, \mathcal{O}_S(K))$ and $G_1 = \mathrm{SL}(n, \mathcal{O}_S(K))$ where $\mathcal{O}_S(K)$ is the ring of S -integers of a number field K (cf. [12, 13]; S is a finite set of places in K including the infinite places). These groups are known to be virtually of finite (cohomological) dimension, and thus the concept of p -periodicity is defined; it refers to a rational prime p and to the p -primary component $\hat{H}^i(G, A, p)$ of the Farrell–Tate cohomology $\hat{H}^i(G, A)$ with respect to an arbitrary G -module A . We recall that \hat{H}^i coincides with the usual cohomology H^i for all i above the virtual dimension of G , and that in the case of a finite group (i.e., a group of virtual dimension zero) the \hat{H}^i , $i \in \mathbb{Z}$, are the usual Tate cohomology groups. The group G is called p -periodic if $\hat{H}^i(G, A, p)$ is periodic in i , for all A , and the smallest corresponding period is then simply called the p -period of G . If G has no p -torsion, the p -primary component of all its \hat{H}^i is 0, and thus G is trivially p -periodic.

We shall determine the rational primes p for which the above S -arithmetic groups are p -periodic, and compute the value of the p -period.

Partial results in that direction have been obtained earlier [3]. The present procedure is simpler and yields complete answers.

1.2. Our method is based on the following fact. Let G be any group of virtually finite dimension, and N a torsion-free normal subgroup of finite index in G . If G/N is p -periodic with p -period m_p , then G itself is p -periodic with p -period dividing m_p (see Section 5). In the case of the S -arithmetic groups G and G_1 above we take for N or N_1 , respectively, the principal congruence subgroup of G or G_1 , with respect to a certain prime ideal P of $\mathcal{O}_S(K)$. This prime ideal can be chosen in such a way that N and N_1 are torsion-free and that the absolute norm $\mathfrak{N}(P) = |\mathcal{O}_S(K)/P| = q$ is a rational prime suitable for our purpose. Then

$$G_1/N_1 \cong \mathrm{SL}(n, \mathbb{F}_q) \subset G/N \subset \mathrm{GL}(n, \mathbb{F}_q).$$

Thus the task is reduced essentially to investigating the p -periodicity of the finite groups $\mathrm{GL}(n, \mathbb{F}_q)$ and $\mathrm{SL}(n, \mathbb{F}_q)$. It turns out (Section 4) that both these groups are p -periodic if $\frac{1}{2}n < h_p(q) \leq n$, where $h_p(q)$ is the order of the residue class of q in $(\mathbb{Z}/p\mathbb{Z})^*$; and that then the p -period is $2h_p(q)$.

The “suitable choice” of P is such that, in addition to rendering N and N_1 torsion-free, its norm $\mathfrak{N}(P) = q$ fulfills $h_q(p) = \phi_\kappa(p)$, the degree over K of the p -th cyclotomic extension $K(\zeta_p)$ of K . It then follows that G and G_1 are p -periodic for $\frac{1}{2}n < \phi_\kappa(p) \leq n$ with p -period dividing $2\phi_\kappa(p)$.

1.3. The existence of such a prime ideal is guaranteed by a *number-theoretic lemma* which we formulate and prove in Section 2, in a slightly more general version than actually needed (Lemma 2.2).

Let p be an odd rational prime, and r a positive integer. There exist infinitely many prime ideals P in $O_S(K)$ such that $\mathfrak{N}(P)$ is a rational prime q whose residue class has order $\phi_K(p^r)$ in $(\mathbb{Z}/p^r\mathbb{Z})^*$.

This lemma is useful also for other applications, in particular, in computations concerning the projective class group of certain arithmetic groups (see [7]), and in connection with topological problems as mentioned in [4].

1.4. In order to obtain, for the appropriate rational primes p , the precise value of the p -period of the groups G and G_1 we exhibit certain finite subgroups; they are obtained as semi-direct products of the group of p -th roots of unity with the Galois group of $K(\zeta_p)$ over K . Since quite generally any subgroup of a p -periodic group is also p -periodic, with p -period dividing that of the group, we thus get lower bounds for the p -periods of G and G_1 . It turns out that they agree with the upper bounds $2\phi_K(p)$ except for the special case $SL(\phi_K(p), O_S(K))$. The final results (Theorems 5.2 and 5.4 with Remarks) are as follows.

The groups $GL(n, O_S(K))$, $n > 0$, and $SL(n, O_S(K))$, $n > 2$, are p -periodic for all rational primes p with $\frac{1}{2}n < \phi_K(p) \leq n$; the p -period is $2\phi_K(p)$ except for $SL(\phi_K(p), O_S(K))$ where it is either $\phi_K(p)$ or $2\phi_K(p)$ depending on the number field K . For $\phi_K(p) \leq \frac{1}{2}n$ they are not p -periodic, and for $\phi_K(p) > n$ they have no p -torsion. The group $SL(2, O_S(K))$ is periodic (i.e., p -periodic for all p) with period 2 or 4.

§2. *The number-theoretic lemma.* 2.1. We consider an algebraic number field K and its ring of integers $O(K)$. Let $\mathfrak{N}(I)$ denote the absolute norm $|O(K)/I|$ of the ideal I in $O(K)$.

LEMMA 2.1. *Let p be an odd prime number and r a positive integer. There exist infinitely many prime ideals P of $O(K)$ such that $\mathfrak{N}(P) = q$ is a prime number whose residue class has order $\phi_K(p^r)$ in $(\mathbb{Z}/p^r\mathbb{Z})^*$.*

Proof. The Galois group $\text{Gal}(K(\zeta_{p^r})/K)$ is cyclic of order $\phi_K(p^r)$; let σ be a generator, i.e. $\sigma(\zeta_{p^r}) = \zeta_{p^r}^s$ where the order of the residue class of s in $(\mathbb{Z}/p^r\mathbb{Z})^*$ is $\phi_K(p^r)$.

We shall use results and notations of [11], Chapters IV and V. We consider the following “modulus” m . Let m_∞ be the product of all real places of K , and $m_0 = p^r O(K)$, and $m = m_0 m_\infty$. Let $K_{m,1}$ be defined by

$$K_{m,1} = \{x/y; x, y \in O(K) \text{ with } xO(K) \text{ and } yO(K) \text{ relatively prime to } m_0 \text{ and } x/y \equiv 1 \pmod{m}\};$$

and I_K^m the subgroup of the ideal group of K generated by all prime ideals not dividing m_0 . The Artin map

$$\phi : I_K^m \rightarrow \text{Gal}(K(\zeta_{p^r})/K)$$

is surjective, and its kernel contains the image $i(K_{m,1})$ of the embedding of $K_{m,1}$ in the ideal group by the reciprocity law for $(K(\zeta_{p^r}), K, m)$. Take $J \in I_K^m$ such that $\phi(J) = \sigma$. Then $\phi^{-1}(\sigma) = J \ker \phi \supset J i(K_{m,1})$. By the generalized Dirichlet theorem

there are in $\phi^{-1}(\sigma)$ infinitely many prime ideals, even if we require them to be of relative degree 1 (over \mathbb{Z}).

Let P be such a prime ideal of $O(K)$. The Frobenius automorphism

$$\left(\frac{K(\zeta_{p^r})/K}{P}\right)$$

is equal to $\sigma \in \text{Gal}(K(\zeta_{p^r})/K)$. Since the relative degree of P is 1, we have $O(K)/P \cong \mathbb{Z}/q\mathbb{Z}$ where q is the rational prime over which P lies ($P \cap \mathbb{Z} = q\mathbb{Z}$). The Frobenius automorphism

$$\left(\frac{\mathbb{Q}(\zeta_{p^r})/\mathbb{Q}}{q}\right)$$

is the restriction of σ to $\mathbb{Q}(\zeta_{p^r})$; i.e.,

$$\zeta_{p^r}^q = \sigma(\zeta_{p^r}) = \zeta_{p^r}^s,$$

whence $q \equiv s \pmod{p^r}$. Thus q has order $\phi_K(p^r)$ in $(\mathbb{Z}/p^r\mathbb{Z})^*$.

2.2. We now consider the ring of S -integers $O_S(K)$ in K . Let Σ be the set of all places of K and S a subset of Σ containing Σ^∞ , the set of infinite places. Then

$$O_S(K) = \bigcap_{Q \in \Sigma - S} O_Q$$

where O_Q is the valuation ring of Q . Hence $O_S(K)$ is a Dedekind ring with quotient field K .

If S above is a finite set then (cf. [12] or [13]) $GL(n, O_S(K))$ is virtually of finite dimension.

LEMMA 2.2. *Let S be a finite set of places including Σ^∞ . Then the assertion of Lemma 2.1 also holds for $O_S(K)$.*

Indeed, all the prime ideals P occurring in Lemma 2.1, except for finitely many of them, generate prime ideals $P' = PO_S(K)$ of $O_S(K)$, and $\mathfrak{R}(P') = |O_S(K)/P'| = |O(K)/P| = \mathfrak{R}(P)$.

§3. Finite subgroups. 3.1. Notation. R is an integrally closed domain of characteristic zero, K its field of quotients, ζ_m a primitive m -th root of unity in an algebraic closure of K , $\phi_K(m) = [K(\zeta_m):K]$, $Z_m = \langle \zeta_m \rangle$ the group of all m -th roots of unity, $C_k = \langle t \rangle$ any multiplicative cyclic group of order k with generator t (m, k are arbitrary natural numbers).

Let p be a rational prime, and let $C_{\phi_K(p)}$ operate on Z_p through the isomorphism $C_{\phi_K(p)} \cong \text{Gal}(K(\zeta_p)/K)$ which maps t to a generator σ of the Galois group.

PROPOSITION 3.1. *The semi-direct product $Z_p \rtimes C_{\phi_K(p)}$ is p -periodic with p -period $2\phi_K(p)$.*

Proof. Obviously Z_p is a p -Sylow subgroup of $G = Z_p \rtimes C_{\phi_K(p)}$. Since it is cyclic, G is p -periodic (cf. [8, Chap. XII]). The p -period is given (cf. [14]) by $2|N_G(Z_p)/C_G(Z_p)|$ where N_G denotes the normalizer, C_G the centralizer in G . Now $N_G(Z_p) = G$ and $C_G(Z_p) = Z_p$, and hence the p -period is $2\phi_K(p)$.

3.2. The group in Proposition 3.1 can be embedded in $GL(\phi_K(p), R)$, as follows. Since the irreducible polynomial in $K[x]$ of ζ_p is of degree $\phi_K(p)$ and has coefficients in R , the R -module $R[\zeta_p]$ is free with basis $1, \zeta_p, \dots, \zeta_p^{\phi_K(p)-1}$. We can thus identify $GL(\phi_K(p), R)$ with the group of R -module automorphisms $\text{Aut}_R R[\zeta_p]$. Multiplication μ_{ζ_p} with ζ_p is an element of that group, and so is any element σ^s of $\text{Gal}(K(\zeta_p)/K)$ if restricted to $R[\zeta_p]$.

We consider the subgroup $S = \{\mu_{\zeta_p}^r \sigma^s; 0 \leq r < p, 0 \leq s < \phi_K(p)\}$ of $\text{Aut}_R R[\zeta_p]$. The map $Z_p \rtimes C_{\phi_K(p)} \rightarrow S$ given by $\zeta_p \mapsto \mu_{\zeta_p}, t \mapsto \sigma$ is easily seen to be an isomorphism. Thus $Z_p \rtimes C_{\phi_K(p)}$ is realized as a subgroup of $GL(\phi_K(p), R)$, and therefore also of $GL(n, R)$ for all $n \geq \phi_K(p)$.

THEOREM 3.2. *For a rational prime p with $\phi_K(p) \leq n$ the group $GL(n, R)$ contains a finite subgroup which is p -periodic with p -period $2\phi_K(p)$.*

3.3. We now turn to the special linear groups over R . Since $SL(n, R)$ contains $GL(n-1, R)$ as a subgroup ($n > 1$) there is, for all p with $\phi_K(p) < n$, a finite subgroup in $SL(n, R)$ which is p -periodic with p -period $2\phi_K(p)$. Some special arguments are needed in the case where $\phi_K(p) = n (> 1)$.

We can identify $SL(\phi_K(p), R)$ with the subgroup $\text{Aut}_R R[\zeta_p]_1$ of $\text{Aut}_R R[\zeta_p]$ consisting of all automorphisms with determinant 1. The determinant of μ_{ζ_p} is a p -th root of 1 in K and hence $\neq 1$ since $\phi_K(p) > 1$. As for the generator σ of $\text{Gal}(K(\zeta_p)/K)$, it has determinant $(-1)^{\phi_K(p)-1}$, indeed σ can be viewed as a cyclic permutation of a suitable basis of $K(\zeta_p)$ over K . Thus for odd $\phi_K(p) > 1$ the group S above actually lies in $\text{Aut}_R R[\zeta_p]_1$. If $\phi_K(p)$ is even, $S_1 = S \cap \text{Aut}_R R[\zeta_p]_1$ has index 2 in S ; this group S_1 is p -periodic with p -period $\phi_K(p)$.

If $\phi_K(p)$ is even there are, however, also cases where one can have in $\text{Aut}_R R[\zeta_p]_1$ a finite p -periodic subgroup S_2 with p -period $2\phi_K(p)$. This is so if there exists in $R[\zeta_p]$ a unit u with relative norm $\mathfrak{N}_{K(\zeta_p)/K}(u) = -1$. Indeed let again μ_u be multiplication in $R[\zeta_p]$ by u . This automorphism has determinant -1 ; thus $\mu_u \sigma$ has determinant 1 and generates in $\text{Aut}_R R[\zeta_p]_1$ a cyclic subgroup of order $2\phi_K(p)$ (since $(\mu_u \sigma)^{\phi_K(p)} = -\text{identity}$). We put

$$S_2 = \{\mu_{\zeta_p}^r (\mu_u \sigma)^s, 0 \leq r < p, 0 \leq s < 2\phi_K(p)\}.$$

This subgroup of $\text{Aut}_R R[\zeta_p]_1$ is isomorphic to $Z_p \rtimes C_{2\phi_K(p)}$ where the generator t of $C_{2\phi_K(p)}$ acts on Z_p through $t \mapsto \sigma$. The computation analogous to that in the proof of Proposition 3.1 shows that S_2 is p -periodic with p -period $2\phi_K(p)$.

In summary we have

THEOREM 3.3. (a) *For all p with $\phi_K(p) < n$, and for $\phi_K(p) = n$ if $\phi_K(p)$ is odd > 1 , the group $SL(n, R)$ contains a finite subgroup which is p -periodic with p -period $2\phi_K(p)$.*

(b) *If $\phi_K(p)$ is even, then $SL(\phi_K(p), R)$ contains a finite subgroup which is p -periodic with p -period $\phi_K(p)$. If there is in $R[\zeta_p]$ a unit u with $\mathfrak{N}_{K(\zeta_p)/K}(u) = -1$, there exists even a finite subgroup with p -period $2\phi_K(p)$.*

§4. *The p -periodicity of $GL(n, \mathbb{F}_q)$ and $SL(n, \mathbb{F}_q)$.* 4.1. As usual \mathbb{F}_{q^n} denotes the field of q^n elements; we recall that

$$|GL(n, \mathbb{F}_q)| = q^{n(n-1)/2} \prod_{i=1}^n (q^i - 1) = (q-1)|SL(n, \mathbb{F}_q)|.$$

Let p and q be different rational primes. We denote by $h_p(q)$ the order of the residue class of q in $(\mathbb{Z}/p\mathbb{Z})^*$. If $h = h_p(q)$ then p divides $q^h - 1$ but none of the other factors in $|GL(h, \mathbb{F}_q)|$. Let p^a be the highest power of p dividing $q^h - 1$, i.e., dividing $|GL(h, \mathbb{F}_q)|$, and let S_p be a p -Sylow subgroup of $GL(h, \mathbb{F}_q)$.

PROPOSITION 4.1. *The group S_p is cyclic; the centralizer of S_p in $GL(h, \mathbb{F}_q)$ has index h in the normalizer.*

Proof. We write G for $GL(h, \mathbb{F}_q)$ and identify G with the group of \mathbb{F}_q -vector space automorphisms of \mathbb{F}_{q^h} . For $x \in \mathbb{F}_{q^h}^*$ let μ_x be multiplication with x in \mathbb{F}_{q^h} ; it is an element of $G = \text{Aut}_{\mathbb{F}_q}(\mathbb{F}_{q^h})$. Let g be a generator of the cyclic group $\mathbb{F}_{q^h}^*$ and $f = g^{(q^h - 1)/p^a}$. Then $\mu_f \in G$ is of order p^a and generates a p -Sylow subgroup S_p of G .

To prove the second part we show that $N_G(S_p)/C_G(S_p)$ is isomorphic to $\text{Gal}(\mathbb{F}_{q^h}/\mathbb{F}_q)$ and hence of order h . Indeed $\text{Gal}(\mathbb{F}_{q^h}/\mathbb{F}_q)$ is contained in G and one easily checks (cf. [6], Lemma 3.2 or [10], Chap. II, §7) that

$$N_G(S_p) = \{ \mu_x \gamma; \quad x \in \mathbb{F}_{q^h}^*, \quad \gamma \in \text{Gal}(\mathbb{F}_{q^h}/\mathbb{F}_q) \},$$

and

$$C_G(S_p) = \{ \mu_x; \quad x \in \mathbb{F}_{q^h}^* \}.$$

Thus $C_G(S_p)$ is the kernel of the obvious map $N_G(S_p) \rightarrow \text{Gal}(\mathbb{F}_{q^h}/\mathbb{F}_q)$ and the assertion follows.

4.2. From Proposition 4.1 it follows that $GL(h, \mathbb{F}_q)$, $h = h_p(q)$, is p -periodic with p -period $2h$. We shall show that the same holds for $GL(n, \mathbb{F}_q)$ if $\frac{1}{2}n < h \leq n$.

Let $B \in GL(h, \mathbb{F}_q)$ be a matrix of order p^a , generating S_p . Then

$$B' = \begin{pmatrix} B & 0 \\ 0 & E \end{pmatrix},$$

where E is the $(n-h) \times (n-h)$ unit matrix, has order p^a in $GL(n, \mathbb{F}_q)$. The assumption $n < 2h$ guarantees that p^a is the highest power of p dividing $|GL(n, \mathbb{F}_q)|$. Thus B' generates a cyclic p -Sylow subgroup S'_p of $GL(n, \mathbb{F}_q)$. The normalizer of S'_p is given by the matrices

$$\left\{ \begin{pmatrix} N & 0 \\ 0 & D \end{pmatrix}; \quad N \in N_{GL(h, \mathbb{F}_q)}(S_p), \quad D \in GL(n-h, \mathbb{F}_q) \right\},$$

and similarly for the centralizer of S'_p . It immediately follows that the index of the centralizer of S'_p in the normalizer is again h ; thus the p -period of $GL(n, \mathbb{F}_q)$ is $2h$.

4.3. The remaining cases $n < h$ and $n \geq 2h$ are easy.

If $n < h = h_p(q)$ then p does not divide $|GL(n, \mathbb{F}_q)|$; i.e., $GL(n, \mathbb{F}_q)$ has no p -torsion.

If $n \geq 2h$ we take an embedding

$$\text{GL}(h, \mathbb{F}_q) \times \text{GL}(h, \mathbb{F}_q) \subset \text{GL}(2h, \mathbb{F}_q) \subset \text{GL}(n, \mathbb{F}_q).$$

Since p divides $|\text{GL}(h, \mathbb{F}_q)|$ there is a cyclic subgroup C_p in $\text{GL}(h, \mathbb{F}_q)$. Thus $\text{GL}(n, \mathbb{F}_q)$ contains a subgroup $C_p \times C_p$ and can therefore not be p -periodic.

4.4. We now turn to the group $\text{SL}(n, \mathbb{F}_q)$, first for $n \geq 3$, and show that all the p -periodicity statements for $\text{GL}(n, \mathbb{F}_q)$ above also hold for $\text{SL}(n, \mathbb{F}_q)$, $n \geq 3$.

We may, of course, assume q odd. So $\text{SL}(n, \mathbb{F}_q)$, being a subgroup of $G = \text{GL}(n, \mathbb{F}_q)$, is p -periodic for $\frac{1}{2}n < h \leq n$, $h = h_p(q)$, with p -period dividing $2h$. The crucial case is again $\text{SL}(h, \mathbb{F}_q)$; by assumption $h > \frac{1}{2}n > 1$.

We write G_1 for $\text{SL}(h, \mathbb{F}_q)$ and identify G_1 with $\text{Aut}_1(\mathbb{F}_{q^h})_1$ where the index 1 refers to determinant 1. With notations as in 4.1 the automorphism μ_f has determinant 1 since p does not divide $q-1 = |\mathbb{F}_q^*|$. Thus the cyclic group S_p generated by μ_f lies in G_1 . Its normalizer is $N_G(S_p) \cap G_1$ and its centralizer is $C_G(S_p) \cap G_1$.

For the generator g of $\mathbb{F}_{q^h}^*$ the determinant $\det \mu_g$ is $g^{(q^h-1)/(q-1)} \in \mathbb{F}_q^*$; and for the generator $\sigma \in \text{Gal}(\mathbb{F}_{q^h}/\mathbb{F}_q)$, $\det \sigma = (-1)^{h-1} \in \mathbb{F}_q^*$ since σ may be viewed as a cyclic permutation of order h . Thus the elements $\mu_x \gamma$, $x \in \mathbb{F}_{q^h}^*$, $\gamma \in \text{Gal}(\mathbb{F}_{q^h}/\mathbb{F}_q)$, of $N_G(S_p)$ have determinant 1 in the following cases.

If h is odd: $x = g^{r(q-1)}$, $0 \leq r < (q^h-1)/(q-1)$; $\gamma = \sigma^s$, $0 \leq s < h$.

If h is even: $x = g^{r(q-1)}$, $0 \leq r < (q^h-1)/(q-1)$; $\gamma = \sigma^{2s}$, $0 \leq s < \frac{1}{2}h$,

and $x = g^{r(q-1) + \frac{1}{2}(q-1)}$, $0 \leq r < (q^h-1)/(q-1)$; $\gamma = \sigma^{2s+1}$, $0 \leq s < \frac{1}{2}h$.

The elements μ_x , $x \in \mathbb{F}_{q^h}^*$, of $C_G(S_p)$ have determinant 1, if, and only if, $x = g^{r(q-1)}$, $0 \leq r < (q^h-1)/(q-1)$. A simple count shows that the index of the centralizer in the normalizer is h ; hence the p -period of $\text{SL}(n, \mathbb{F}_q)$, $n \geq 3$, is $2h$.

4.5. We summarize as follows.

THEOREM 4.2. *Let p and q be different prime numbers, and $h = h_p(q)$ the order of q in $(\mathbb{Z}/p\mathbb{Z})^*$. If $\frac{1}{2}n < h \leq n$, then the groups $\text{GL}(n, \mathbb{F}_q)$, $n \geq 1$, and $\text{SL}(n, \mathbb{F}_q)$, $n \geq 3$, are p -periodic with p -period $2h$.*

Remark 4.3. (a) For $\frac{1}{2}n \geq h = h_p(q)$ the groups in Theorem 4.2 are not p -periodic.

(b) For $n < h$ they have no p -torsion.

Indeed, (a) is proved in 4.3 for $\text{GL}(n, \mathbb{F}_q)$. If $h \geq 2$ ($n \geq 4$), then p does not divide $q-1 = |\mathbb{F}_q^*|$, and the subgroup $C_p \times C_p$ mentioned in 4.3 actually lies in $\text{SL}(n, \mathbb{F}_q)$. If $h = 1$ a special argument is needed for $\text{SL}(n, \mathbb{F}_q)$, $n \geq 3$. In that case p divides $q-1$; let $x \in \mathbb{F}_{q-1}^*$ be of order p . The matrices

$$\begin{pmatrix} x^r & 0 & 0 \\ 0 & x^s & 0 \\ 0 & 0 & x^{-r-s} \end{pmatrix}$$

with $0 \leq r, s < p$ constitute a subgroup of $SL(3, \mathbb{F}_q)$ isomorphic to $C_p \times C_p$. Thus $SL(n, \mathbb{F}_q)$, $n \geq 3$, is not p -periodic in that case. The result (b) is proved in §4.3.

Remark 4.4. $SL(2, \mathbb{F}_q)$ is well known to be p -periodic for all p . The q -period is $q-1$ for odd q , and 2 for $q = 2$. For p dividing $q^2 - 1$ the p -period is 4.

§5. *Finite quotients. Main results.* 5.1. We now turn to the groups $G = GL(n, O_S(K))$ and $G_1 = SL(n, O_S(K))$ described in Section 1. K is a number field, S a finite set of places including the infinite places, $O_S(K)$ the ring of S -integers of K .

We choose, by virtue of Lemma 2.2, a prime ideal P of $O_S(K)$ such that $\mathfrak{N}(P)$ is a prime number $q > 2^{[K:\mathbb{Q}]}$, and that $h_p(q) = \phi_K(p)$; p is a given prime number and $h_p(q)$ is the order of q in $(\mathbb{Z}/p\mathbb{Z})^*$. Then $O_S(K)/P \cong \mathbb{F}_q$, and reducing all matrix entries modulo P yields canonical maps $\psi : G \rightarrow GL(n, \mathbb{F}_q)$ and $\psi_1 : G_1 \rightarrow SL(n, \mathbb{F}_q)$. Their kernels are the respective congruence subgroups modulo P , $N \subset G$ and $N_1 \subset G_1$. Due to the choice of P they are torsion-free (cf. [2], for example). The map ψ_1 is known to be surjective ([1], p.267), i.e., we have

$$G_1/N \cong SL(n, \mathbb{F}_q) \subset \text{Im } \psi \subset GL(n, \mathbb{F}_q).$$

As shown in Section 4 both $SL(n, \mathbb{F}_q)$ and $GL(n, \mathbb{F}_q)$ are p -periodic with p -period $2h_p(q) = 2\phi_K(p)$ for all prime numbers p with $\frac{1}{2}n < \phi_K(p) \leq n$; thus the same holds for G/N and G_1/N_1 .

PROPOSITION 5.1. *There exists a prime ideal P in $O_S(K)$ such that the congruence subgroups modulo P , $N \subset G$ and $N_1 \subset G_1$, are torsion-free and such that the finite quotients G/N and G_1/N_1 are p -periodic with p -period $2\phi_K(p)$ for all p with $\frac{1}{2}n < \phi_K(p) \leq n$.*

5.2. We now invoke a general result concerning the Farrell–Tate cohomology of a group G of virtually finite dimension. Let N be a torsion-free normal subgroup of finite index in G such that G/N is p -periodic with p -period m_p ; then G itself is p -periodic with p -period dividing m_p . In the case, where G admits a projective resolution which is finitely generated in all dimensions, this result is proved in [2] using the construction of a complete resolution for G from a complete resolution for G/N , cf. [2] or [9]. Actually the result holds without any finiteness condition (see [5]); in the present context this generality is not needed since the above finiteness condition holds for $GL(n, O_S(K))$ and $SL(n, O_S(K))$ according to Borel–Serre (see [13], e.g.).

It thus follows that our groups G and G_1 are p -periodic for the appropriate prime numbers p , and that the p -period divides $2\phi_K(p)$.

5.3. To obtain the precise value of the p -period we use the finite subgroups constructed in Section 3. By Theorems 3.2 and 3.3 the groups $G = GL(n, O_S(K))$, $n \geq \phi_K(p)$, and $G_1 = SL(n, O_S(K))$, $n > \phi_K(p)$ contain a finite subgroup which has p -period $2\phi_K(p)$. Thus, for $\frac{1}{2}n < \phi_K(p) \leq n$ (or $< n$ respectively) the p -period of $GL(n, O_S(K))$ and $SL(n, O_S(K))$ respectively is equal to $2\phi_K(p)$. The case $SL(\phi_K(p), O_S(K))$ is discussed in 5.4 below.

THEOREM 5.2. *The groups $\mathrm{GL}(n, O_S(K))$, $\frac{1}{2}n < \phi_K(p) \leq n$, and $\mathrm{SL}(n, O_S(K))$, $\frac{1}{2}n < \phi_K(p) < n$, are p -periodic with p -period $2\phi_K(p)$.*

Remark 5.3. The groups $\mathrm{GL}(n, O_S(K))$ and $\mathrm{SL}(n, O_S(K))$ have p -torsion, if, and only if, $\phi_K(p) \leq n$, see [3]. Using this fact one can, if $n \geq 2\phi_K(p)$, easily find a subgroup of these groups (for $\mathrm{SL}(n, O_S(K))$ assuming $n \geq 3$) isomorphic to $C_p \times C_p$. Therefore they are not p -periodic if $\frac{1}{2}n \geq \phi_K(p)$.

5.4. In the special case $\mathrm{SL}(\phi_K(p), O_S(K))$ all the above arguments remain valid except that Theorem 3.3 yields, in general, the two possibilities $\phi_K(p)$ or $2\phi_K(p)$ for the p -period. If $\phi_K(p)$ is odd and greater than one, the p -period is $2\phi_K(p)$, by Theorem 3.3(a). If $\phi_K(p)$ is even, the precise value depends on the norm map $\mathfrak{N}_{K(\zeta_p)/K}$. By Theorem 3.3(b) the period is again $2\phi_K(p)$, if there exists in $O_S(K)[\zeta_p]$ a unit u with $\mathfrak{N}_{K(\zeta_p)/K}(u) = -1$.

THEOREM 5.4. *The group $\mathrm{SL}(\phi_K(p), O_S(K))$, $\phi_K(p) > 1$, is p -periodic with p -period $\phi_K(p)$ or $2\phi_K(p)$. If $\phi_K(p)$ is odd or, more generally, if there is in $O_S(K)[\zeta_p]$ a unit with norm -1 over K , then the p -period is $2\phi_K(p)$.*

Remark 5.5. If there is no element in $K(\zeta_p)$ with norm -1 over K , then the p -period of $\mathrm{SL}(\phi_K(p), O_S(K))$ is $\phi_K(p)$. This follows from the computations in [6], Section 8. The condition is fulfilled, in particular, if K has an embedding in \mathbb{R} . Thus $\mathrm{SL}(p-1, \mathbb{Z})$, for example, is p -periodic with p -period $p-1$ (this case appears in [3] and is obtained by an entirely different method).

References

1. H. Bass. *Algebraic K-theory* (Benjamin, New York–Amsterdam, 1968).
2. B. Bürgisser. Gruppen virtuell endlicher Dimension und Periodizität der Cohomologie. *Diss. ETH*, Nr. 6425 (Zürich, 1979).
3. B. Bürgisser. On the p -periodicity of arithmetic subgroups of general linear groups. *Comment. Math. Helv.*, 55 (1980), 499–509.
4. B. Bürgisser. Finite p -periodic quotients of general linear groups. *Math. Ann.*, 256 (1981), 121–132.
5. B. Bürgisser. Yoneda product in Farrell Tate cohomology and periodicity. Preprint *FIM, ETH Zürich* (1981).
6. B. Bürgisser. The p -torsion of the Farrell Tate cohomology of $\mathrm{GL}(\phi_K(p), O(K))$ and $\mathrm{SL}(\phi_K(p), O(K))$. Preprint *FIM, ETH Zürich* (1981).
7. B. Bürgisser. On the projective class group of arithmetic groups. *Math. Zeitschrift*, 184 (1983), 339–357.
8. H. Cartan and S. Eilenberg. *Homological algebra* (Princeton University Press, 1956).
9. F. T. Farrell. An extension of Tate cohomology to a class of infinite groups. *J. Pure Appl. Algebra*, 10 (1977), 153–161.
10. B. Huppert. *Endliche Gruppen I. Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellung*, 134 (Springer, Berlin–Heidelberg–New York, 1967).
11. G. J. Janusz. *Algebraic number fields. Pure and Appl. Math.*, 55 (Academic Press, New York, 1973).
12. J.-P. Serre. *Cohomologie des groupes discrets. Prospects in Math.*, Ann. Math. Study, 70 (1971), 77–169.
13. J.-P. Serre. *Arithmetic groups. Proc. of the Sept. 1977 Durham conference on homological and combinatorial techniques in group theory*. Edited by C. T. C. Wall. (Cambridge University Press, 1979).
14. R. G. Swan. The p -period of a finite group. *Illinois J. Math.*, 4 (1960), 341–346.
15. R. G. Swan and E. G. Evans. *K-theory of finite groups and orders. Lecture Notes in Math.*, 149 (Springer, Berlin–Heidelberg–New York, 1970).

Dr. B. Bürgisser,
Eidgenössische Technische Hochschule,
Mathematik,
ETH-Zentrum,
CH-8092 Zürich,
Switzerland.

20G10: *GROUP THEORY AND GENERALIZATIONS; Linear algebraic groups; Cohomology theory.*

Prof. B. Eckmann,
Eidgenössische Technische Hochschule,
Mathematik,
ETH-Zentrum,
CH-8092 Zürich,
Switzerland.

Received on the 4th of August, 1983.