

University of Wollongong

Research Online

Faculty of Science, Medicine and Health -
Papers: part A

Faculty of Science, Medicine and Health

1-1-2014

The P2X7 receptor channel: recent developments and the use of P2X7 antagonists in models of disease

Rachael Bartlett

University of Wollongong, rb401@uowmail.edu.au

Leanne Stokes

RMIT University

Ronald Sluyter

University of Wollongong, rsluyter@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/smhpapers>



Part of the [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Bartlett, Rachael; Stokes, Leanne; and Sluyter, Ronald, "The P2X7 receptor channel: recent developments and the use of P2X7 antagonists in models of disease" (2014). *Faculty of Science, Medicine and Health - Papers: part A*. 1818.

<https://ro.uow.edu.au/smhpapers/1818>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

The P2X7 receptor channel: recent developments and the use of P2X7 antagonists in models of disease

Abstract

The P2X7 receptor is a trimeric ATP-gated cation channel found predominantly, but not exclusively, on immune cells. P2X7 activation results in a number of downstream events, including the release of proinflammatory mediators and cell death and proliferation. As such, P2X7 plays important roles in various inflammatory, immune, neurologic and musculoskeletal disorders. This review focuses on the use of P2X7 antagonists in rodent models of neurologic disease and injury, inflammation, and musculoskeletal and other disorders. The cloning and characterization of human, rat, mouse, guinea pig, dog, and Rhesus macaque P2X7, as well as recent observations regarding the gating and permeability of P2X7, are discussed. Furthermore, this review discusses polymorphic and splice variants of P2X7, as well as the generation and use of P2X7 knockout mice. Recent evidence for emerging signaling pathways downstream of P2X7 activation and the growing list of negative and positive modulators of P2X7 activation and expression are also described. In addition, the use of P2X7 antagonists in numerous rodent models of disease is extensively summarized. Finally, the use of P2X7 antagonists in clinical trials in humans and future directions exploring P2X7 as a therapeutic target are described.

Keywords

CMMB

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Bartlett, R., Stokes, L. & Sluyter, R. (2014). The P2X7 receptor channel: recent developments and the use of P2X7 antagonists in models of disease. *Pharmacological Reviews*, 66 (3), 638-675.

ASSOCIATE EDITOR: DAVID R. SIBLEY

The P2X7 Receptor Channel: Recent Developments and the Use of P2X7 Antagonists in Models of Disease

Rachael Bartlett, Leanne Stokes, and Ronald Sluyter

School of Biological Sciences, University of Wollongong, New South Wales, Australia and Illawarra Health and Medical Research Institute, Wollongong, New South Wales, Australia (R.B., R.S.); and Health Innovations Research Institute, School of Medical Sciences, RMIT University, Bundoora, Victoria, Australia (L.S.)

Abstract	639
I. The P2X7 Receptor	639
A. Structure and Function of the P2X7 Receptor	639
B. Distribution of the P2X7 Receptor	640
C. The P2X7 Receptor in Health and Disease	640
II. The Rodent P2X7 Receptor	641
III. The Human P2X7 Receptor	641
IV. P2X7 Receptors of Other Mammalian Species	642
A. The Guinea Pig P2X7 Receptor	642
B. The Canine P2X7 Receptor	642
C. The Rhesus Macaque P2X7 Receptor	642
V. Gating of the P2X7 Receptor Channel	643
VI. Permeability of the P2X7 Receptor	643
VII. P2X7 Receptor-Dependent Signaling Pathways	644
A. Cytokine Release	644
B. Reactive Oxygen Species Formation	644
C. Protease Activation and Release	644
D. Prostaglandin Release	645
E. Glutamate Efflux	645
F. Transcription Activation	645
G. Cell Proliferation	645
H. Phagocytosis	645
I. Summary	646
VIII. Variants of the P2X7 Receptor	646
A. P2X7 Single Nucleotide Polymorphisms	646
B. P2X7 Splice Isoforms	648
C. P2X7 Variants in P2X7 Knockout Mice	649
IX. Modulators of P2X7 Receptor Activation	649
X. Use of Modulators of P2X7 Receptor In Vivo	653
A. P2X7 Antagonists in Rodent Models of Neurologic Disease	653
B. P2X7 Antagonists in Rodent Models of Neurologic Injury	658
C. P2X7 Antagonists in Rodent Models of Compound-Induced Inflammation	660
D. P2X7 Antagonists in Rodent Models of Musculoskeletal Disorders	661
E. P2X7 Antagonists in Rodent Models of Disorders Associated with Other Tissues	661
XI. P2X7 Antagonists in Human Clinical Trials	661

This work was supported by an Australian Postgraduate Award from the University of Wollongong (to R.B.); the University of Wollongong Global Challenges Program (to R.B.); the National Health and Medical Research Council (to L.S.), an RMIT University Vice Chancellor's Research Fellowship (to L.S.); the American Kennel Club Canine Health Foundation (to R.S. and L.S.); the University Research Committee of the University of Wollongong (to R.S.); and the Centre for Medical and Molecular Bioscience (to R.S.).

The authors declare no conflict of interest.

Address correspondence to: Ronald Sluyter, Illawarra Health and Medical Research Institute, Wollongong, NSW 2522, Australia.

E-mail address: rsluyter@uow.edu.au

dx.doi.org/10.1124/pr.113.008003.

XII. Modulators of P2X7 Receptor Expression	667
XIII. Conclusions and Future Directions	667
Acknowledgments	668
References	668

Abstract—The P2X7 receptor is a trimeric ATP-gated cation channel found predominantly, but not exclusively, on immune cells. P2X7 activation results in a number of downstream events, including the release of proinflammatory mediators and cell death and proliferation. As such, P2X7 plays important roles in various inflammatory, immune, neurologic and musculoskeletal disorders. This review focuses on the use of P2X7 antagonists in rodent models of neurologic disease and injury, inflammation, and musculoskeletal and other disorders. The cloning and characterization of human, rat, mouse, guinea pig, dog, and Rhesus macaque P2X7, as well as recent observations

regarding the gating and permeability of P2X7, are discussed. Furthermore, this review discusses polymorphic and splice variants of P2X7, as well as the generation and use of P2X7 knockout mice. Recent evidence for emerging signaling pathways downstream of P2X7 activation and the growing list of negative and positive modulators of P2X7 activation and expression are also described. In addition, the use of P2X7 antagonists in numerous rodent models of disease is extensively summarized. Finally, the use of P2X7 antagonists in clinical trials in humans and future directions exploring P2X7 as a therapeutic target are described.

I. The P2X7 Receptor

A. Structure and Function of the P2X7 Receptor

ATP and other nucleotides, as well as nucleosides, are important extracellular signaling molecules that operate through a complex purinergic signaling network (Burnstock, 1972, 2006). This network is composed of a number of membrane receptors and ectoenzymes, which includes the P2X7 receptor (Yegutkin, 2008). The P2X7 receptor is encoded by the *P2RX7* gene and belongs to the P2X family of trimeric ligand-gated cation channels, of which there are seven distinct members (P2X1–7) (Coddou et al., 2011). Of the P2X family, the P2X7 monomeric subunit is the largest, with a length of 595 amino acids for the human, rat, mouse, dog, and Rhesus macaque receptors (Surprenant et al., 1996; Rassendren et al., 1997; Chessell et al., 1998; Roman et al., 2009; Bradley et al., 2011b). Each subunit is characterized by relatively short and long intracellular amino and carboxyl (C) termini, respectively, as well as two hydrophobic membrane-spanning segments (transmembrane domains) separated by a long glycosylated extracellular ATP-binding domain. The trimeric structures of the human and Rhesus macaque P2X7 receptors are supported by atomic computer modeling (Roger et al., 2010b; Bradley et al., 2011b; Jiang et al., 2013), based on the crystal structure

of the related zebrafish P2X4 receptor (Gonzales et al., 2009; Kawate et al., 2009). To the best of our knowledge, similar modeling experiments have not been carried out for rodent or canine P2X7 receptors. However, the structures of rodent and canine P2X7 receptors are likely to be similar to that of primate P2X7, as these receptors share 77–85% sequence identity to human P2X7 (Table 1). Furthermore, the study of native rat P2X7 complexes supports the trimeric arrangement of rodent P2X7 (Nicke, 2008). More recently, the crystal structure of the zebrafish P2X4 receptor in complex with ATP has been reported (Hattori and Gouaux, 2012). This agonist-bound structure provides new insights into the mechanism of P2X receptor activation and may be useful for the development of new pharmacological agents.

Activation of P2X7 by extracellular ATP allows for the passage of small cations, including Ca^{2+} , Na^{+} , and K^{+} , across the plasma membrane (Surprenant et al., 1996; Rassendren et al., 1997; Chessell et al., 1998). However, prolonged ATP stimulation leads to the formation of a larger reversible pore, which allows for the uptake of organic ions (Steinberg and Silverstein, 1987; Surprenant et al., 1996; Rassendren et al., 1997; Chessell et al., 1998) (see section VI). Activation of P2X7 by ATP, or the alternate, rodent-specific ligand nicotinamide adenine dinucleotide (NAD), results in a number of cell-specific

ABBREVIATIONS: A438079, 3-[[5-(2,3-dichlorophenyl)-1H-tetrazol-1-yl]methyl]pyridine; A740003, *N*-(1-((cyanoimino)(5-quinolinylamino)methyl) amino)-2,2-dimethylpropyl)-2-(3,4-dimethoxyphenyl)acetamide; A839977, 1-(2,3-dichlorophenyl)-*N*-[[2-(2-pyridinyloxy)phenyl]methyl]-1H-tetrazol-5-amine; AACBA, *N*-(adamantan-1-ylmethyl)-5-[(3*R*-amino-pyrrolidin-1-yl)methyl]-2-chloro-benzamide; ART, ADP-ribosyltransferase; ATP γ S, adenosine-5'-*O*-(3-thiotriphosphate); AZ10606120, *N*-[2-[[2-[(2-hydroxyethyl)amino]ethyl]amino]-5-quinoliny]-2-tricyclo[3.3.1.1^{3,3}.1^{3,7}]dec-1-ylacetamide; AZ11645373, 3-[1-[[[3'-nitro[1,1'-biphenyl]-4-yl]oxy]methyl]-3-(4-pyridinyl)propyl]-2,4-thiazolidinedione; BBG, Brilliant Blue G; BzATP, 2'-(3')-*O*-4-benzoylbenzoyl)-ATP; CE-224,535, 2-(4-chloro-3-[3-(1-hydroxycycloheptyl)propanoyl]phenyl)-4-[(2*R*)-2-hydroxy-3-methoxy-propyl]-1,2,4-triazine-3,5-dione; CFA, complete Freund's adjuvant; GSK1370319A, *N*-[[2,4-dichlorophenyl)methyl]-1-methyl-5-oxo-1-prolinamide; GSK, GlaxoSmithKline; HEK, human embryonic kidney; IL, interleukin; IFN- γ , interferon- γ ; JNJ-47965567, *N*-[(4-(4-phenyl-piperazin-1-yl)tetrahydro-2*H*-pyran-4-yl)methyl]-2-(phenyl-thio) nicotinamide; KN-62, 1-[*N*,*O*-bis(5-isoquinolinesulfonyl)-*N*-methyl-1-tyrosyl]-4-phenylpiperazine; KO, knockout; LPS, lipopolysaccharide; MED1011, 2-[1-(6-aminopurin-9-yl)-2-osso-etosil]prop-2-enale; 2MeSATP, 2-methylthio-ATP; NAD, nicotinamide adenine dinucleotide; NCBI, National Center for Biotechnology Information; oATP, periodate-oxidized ATP; PGE₂, prostaglandin E₂; PPADS, pyridoxal phosphate-6-azophenyl-2,4-disulfonic acid; ROS, reactive oxygen species; SNP, single nucleotide polymorphism; Sp1, specificity protein 1 transcription factor; TM2, second transmembrane domain; TNF- α , tumor necrosis factor- α ; WT, wild-type.

TABLE 1
Agonist profiles of mammalian P2X7

Species	Sequence Identity	EC ₅₀			References
		ATP	BzATP	ATP γ S	
	%		μ M		
Human	100	96	5	>100	Bianchi et al., 1999
Rhesus macaque	96	800	58	N.D.	Bradley et al., 2011b
Dog	85	1148	21	N.D.	Roman et al., 2009
Rat	80	85	4	>1000	Surprenant et al., 1996
Mouse (BALB/c)	80	200	60	>1000	Donnelly-Roberts et al., 2009
Mouse (C57BL/6)	80	162	36	>1000	Donnelly-Roberts et al., 2009
Guinea pig	77	603	>200	N.D.	Fonfria et al., 2008

N.D., not determined.

downstream signaling events, many of which were established or are supported by rodent models of disease (see sections VII and X). These downstream signaling events are dependent on a variety of factors, including cell type, extracellular conditions, and the concentration of extracellular ATP (Burnstock, 2007). The presence of ectonucleotidases, which degrade ATP and other nucleotides, regulate the concentration and duration of availability of P2 receptor agonists in the extracellular space (Yegutkin, 2008). Thus, P2X7 receptors are possibly only activated after injury, infection, or in tumor micro-environments when the concentration of ATP increases locally or when ectonucleotidases are downregulated (Lenertz et al., 2011). Alternatively, it is possible that currently unknown allosteric modulators might act on P2X7 in vivo to decrease its K_m for ATP, thus allowing the activation of P2X7 at lower nucleotide concentrations. There is also emerging evidence that P2X7 activation stimulates ATP release, further complicating the study of the ATP/P2X7 signaling axis. P2X7-mediated ATP release has been observed from human embryonic kidney (HEK)-293 cells transfected with rat or human P2X7 (Pellegatti et al., 2005), from astrocytes during Ca^{2+} signal transmission (Suadicani et al., 2006), from osteoclasts undergoing fusion (Pellegatti et al., 2011) and in culture (Brandao-Burch et al., 2012), and from melanoma cells after γ -irradiation (Ohshima et al., 2010).

B. Distribution of the P2X7 Receptor

P2X7 is widely distributed throughout the mammalian body (see Burnstock and Knight, 2004). This receptor was originally thought to be restricted to cells of the hematopoietic lineages; this includes macrophages, dendritic cells, monocytes, lymphocytes, and erythrocytes, as well as osteoclasts, mast cells, and eosinophils. However, it is now evident that P2X7 is present on cells from other lineages, including osteoblasts, fibroblasts, endothelial cells, and epithelial cells. Furthermore, P2X7 is present on cells in the central and peripheral nervous systems, including microglia, astrocytes, oligodendrocytes, and Schwann cells (Sperlágh et al., 2006). In addition, there are reports of the presence of P2X7 on some populations of neurons, including those from the spinal cord, cerebellum, hypothalamus, and substantia nigra (see Lenertz

et al., 2011; Wiley et al., 2011). Although P2X7 has been identified on a number of cell types, less is known about the relative distribution of P2X7 between different cell types within whole tissues. Recently, transgenic P2X7 reporter mice expressing enhanced green fluorescent protein downstream of the *P2RX7* promoter have been generated (Engel et al., 2012; Garcia-Huerta et al., 2012; Jimenez-Pacheco et al., 2013), which is an important step toward addressing the distribution of P2X7 in vivo. For example, these mice have been used to localize P2X7 expression in the central nervous system after prolonged seizures, with P2X7 identified to be upregulated on neurons, but not upregulated on microglia or astrocytes, in the hippocampus and neocortex (Engel et al., 2012; Jimenez-Pacheco et al., 2013). Although the P2X7 expression on microglia and astrocytes was weak in these studies, P2X7 expression in these cell types is well established from immunohistochemical studies (see Verkhratsky et al., 2012). In addition, transgenic P2X7 green fluorescent protein reporter mice have been used to localize P2X7 on cells from the cerebral cortex and pons of newborn mice (Engel et al., 2012; Garcia-Huerta et al., 2012). P2X7 expression was also demonstrated in peritoneal macrophages and the spleen, validating the reliability of these mice (Garcia-Huerta et al., 2012).

C. The P2X7 Receptor in Health and Disease

P2X7 has been associated with a number of diseases (see Miller et al., 2011; Sluyter and Stokes, 2011; Volonté et al., 2012). In particular, a number of human P2X7 single nucleotide polymorphisms (SNPs) or haplotypes that decrease or increase receptor function have been associated with various infectious, musculoskeletal, psychiatric, inflammatory, and cardiovascular diseases (see section VIII). In addition to diseases associated with SNPs, studies of human tissue or mouse models suggest that P2X7 may play important roles in a number of inflammatory, immune, neurologic, or musculoskeletal disorders. Such disorders include multiple sclerosis (Yiangou et al., 2006), amyotrophic lateral sclerosis (Yiangou et al., 2006), Alzheimer's disease (Ryu and McLarnon, 2008; Diaz-Hernandez et al., 2012), Huntington's disease (Díaz-Hernández et al., 2009), cancer (Ghiringhelli et al., 2009; Adinolfi

et al., 2012), ischemia (Arbeloa et al., 2012; Chu et al., 2012), neuropathic and inflammatory pain (Chessell et al., 2005), rheumatoid arthritis (Portales-Cervantes et al., 2010; Bhattacharya et al., 2011), glomerulonephritis (Taylor et al., 2009b), pulmonary fibrosis (Riteau et al., 2010), and graft-versus-host disease (Wilhelm et al., 2010).

Given the potential importance of P2X7 in health and disease, considerable effort has gone into characterizing the presence and function of mammalian P2X7 receptors, and into the generation of selective P2X7 antagonists and the investigation of the potential therapeutic efficacy of such compounds in rodent models of disease. This paper aims to review recent developments regarding mammalian P2X7 receptors, with an emphasis on the use of P2X7 antagonists in rodent models of disease.

II. The Rodent P2X7 Receptor

Rodent P2X7 was first cloned from rat brain (Surprenant et al., 1996), followed by the cloning of P2X7 from NTW8 murine microglia soon after (Chessell et al., 1998). Compared with other P2X receptors, P2X7 requires at least one log higher ATP concentrations for activation (Table 1). Within this concentration range, ATP induces channel and pore formation in cells transfected with recombinant rat or mouse P2X7 (Surprenant et al., 1996; Chessell et al., 1998). With recombinant rat P2X7, the agonist order of potency was 2'(3')-O-4-benzoylbenzoyl-ATP (BzATP) \gg ATP $>$ 2-methylthio-ATP (2MeSATP) $>$ adenosine-5'-O-(3-thiotriphosphate) (ATP γ S) \gg ADP (Surprenant et al., 1996). However, a more recent study failed to confirm 2MeSATP and ATP γ S as agonists of rat P2X7 (Donnelly-Roberts et al., 2009). Moreover, this same study failed to establish these same two nucleotides as agonists of murine P2X7 (Donnelly-Roberts et al., 2009). Notably, there are reported differences in P2X7 agonist sensitivities between these two species, with rat P2X7 being 10-fold more sensitive to BzATP and ATP than mouse P2X7 (Young et al., 2007).

In rodents, P2X7 can also be activated via a unique ATP-independent pathway involving the transfer of an ADP-ribose group from NAD to P2X7 (Adriouch et al., 2001; Seman et al., 2003). ADP-ribosylation, a post-translational modification of proteins, is an important regulatory mechanism. In prokaryotes, ADP-ribosylation of target proteins is responsible for the adverse effects of a number of bacterial toxins in host cells (Ludden, 1994; Aktories, 2011). In eukaryotes, ADP-ribosylation has been implicated in transcriptional regulation, cell division, intracellular energy metabolism, neuronal signaling, and inflammation (Abd Elmageed et al., 2012; Hassler and Ladurner, 2012). Similar to ATP-induced P2X7 activation, NAD-induced activation of P2X7 results in Ca²⁺ flux, pore formation, phosphatidyserine exposure, shedding of CD62L, cell shrinkage, DNA fragmentation, and apoptosis (Adriouch et al., 2001; Seman et al., 2003). These NAD-induced P2X7 responses are observed

in T cells, but they are not observed in macrophages (Adriouch et al., 2001; Hong et al., 2009). However, although the majority of studies investigating NAD-induced activation of P2X7 have been performed with murine T cells, there is also evidence that this pathway occurs in murine astrocytes (Wang et al., 2012) and rat retinal microvessels (Liao and Puro, 2006). Although NAD induces similar downstream events to that seen after ATP-induced activation of P2X7, much lower NAD concentrations are required for receptor activation compared with ATP (EC₅₀ for phosphatidyserine exposure 2 and 100 μ M, respectively) (Seman et al., 2003). It is noteworthy that NAD and ATP, released through the lysis of erythrocytes, are able to activate P2X7, suggesting that sufficient concentrations of these ligands can be released to activate P2X7 within rodents (Scheuplein et al., 2009).

The transfer of ADP-ribose from NAD to P2X7 is catalyzed by the cell-surface ectoenzyme ADP-ribosyltransferase (ART) 2 (Adriouch et al., 2001), the crystal structure of which has been determined for rat (Mueller-Dieckmann et al., 2002). Of note, this pathway of P2X7 activation does not occur in humans because of the absence of ART2 orthologs (Haag et al., 1994). In rodents, there are two isoforms of ART2, termed ART2.1 and ART2.2. These isoforms are thought to play a role in the reported differences in sensitivities to NAD-mediated P2X7 activation between mouse strains and cell types. First, NAD induces Ca²⁺ flux, but not apoptosis, in T cells derived from C57BL/6 mice, whereas both these NAD-mediated P2X7 responses are observed in T cells derived from BALB/c mice (Adriouch et al., 2001; Hong et al., 2009). C57BL/6 mice have deficient expression of ART2.1 due to a premature stop codon in the *Art2a* gene (Kanaitzuka et al., 1997), which may explain these observations (Adriouch et al., 2001; Hong et al., 2009). Alternatively, these differences may also be explained by a known loss-of-function SNP present in C57BL/6, but not BALB/c, P2X7 (see section VIII). Second, ART2.2 has been reported on T cells, but not macrophages, from a number of mouse strains (Okamoto et al., 1998; Koch-Nolte et al., 1999; Hong et al., 2007). Moreover, bone marrow-derived macrophages from BALB/c mice do not constitutively express any ecto-ART subtypes; however, ART2.1 is upregulated on these cells in response to proinflammatory mediators (Hong et al., 2007). Despite this, these macrophages remain unresponsive to NAD when primed (Hong et al., 2009). Rather, NAD appears to potentiate ATP-induced P2X7 activation in these cells (Hong et al., 2009). Together, these results indicate that ART2.2 is important for NAD-mediated P2X7 activation and that different P2X7 signaling mechanisms operate in lymphoid and myeloid leukocytes, at least in mice.

III. The Human P2X7 Receptor

Human P2X7 was first cloned from a human monocytic cDNA library (Rassendren et al., 1997) using

information from the cloned rat P2X7 sequence. Several pharmacological differences exist between human and rodent P2X7, including varying EC₅₀ values for agonists ATP and BzATP (10- to 25-fold higher for humans) and sensitivity to other compounds. The agonist profile for human P2X7 is BzATP >> ATP > 2MeSATP > ATP γ S >> ADP (Gargett et al., 1997; Donnelly-Roberts et al., 2009). Differences are also apparent in the electrophysiological profile between human and rat P2X7, with faster deactivation observed with human P2X7 expressed in HEK-293 cells activated by either ATP or BzATP (Rassendren et al., 1997). The C terminus was responsible for this effect because a chimeric receptor with human ectodomain and rat C terminus displayed an increased deactivation time, similar to rat P2X7 (Rassendren et al., 1997). More recently, Roger and colleagues (2008, 2010a) identified differences in the intracellular regulation of human and rat P2X7 by calmodulin. Rat P2X7 displays a Ca²⁺-calmodulin dependent facilitation of the ATP-induced inward current via a C-terminal calmodulin binding site, whereas human P2X7 only displays a smaller Ca²⁺-independent facilitation (Roger et al., 2010a).

IV. P2X7 Receptors of Other Mammalian Species

The P2X7 receptor has also been cloned from the guinea pig, dog, and Rhesus macaque. Details of recombinant and native P2X7 from these mammalian species are limited, but nevertheless highlight the potential of these species for the future study of P2X7-related diseases and in the mandatory testing and evaluation of preclinical drugs targeting P2X7.

A. The Guinea Pig P2X7 Receptor

Guinea pig P2X7 was first cloned from a guinea pig brain cDNA library (Fonfria et al., 2008). Unlike the other cloned mammalian P2X7 receptors, guinea pig P2X7 is 594 amino acids in length because of the absence of a glutamic acid at position 77 (Fonfria et al., 2008). The EC₅₀ value for ATP for guinea pig was similar to that of human P2X7, but in contrast to human (and rat) P2X7, BzATP was a weak partial agonist of guinea pig P2X7 (Fonfria et al., 2008). Notably, BzATP-induced responses were observed only in sucrose medium (Fonfria et al., 2008). The capacity of other nucleotides and NAD to activate this rodent P2X7 has not been assessed. Moreover, the electrophysiological characteristics of recombinant guinea pig P2X7 are yet to be reported, although P2X7-mediated currents have been detected in guinea pig myenteric neurons (Valdez-Morales et al., 2011). Further studies of native P2X7 within this species are required before guinea pigs can serve as a model of P2X7-related disorders and preclinical drug testing.

B. The Canine P2X7 Receptor

Canine P2X7 was first cloned from a dog heart cDNA library from a nondisclosed breed (Roman et al., 2009).

EC₅₀ values for both BzATP and ATP of canine P2X7 were similar to that of human P2X7; however, BzATP was only a partial agonist of canine P2X7 (Roman et al., 2009). In contrast, activation of P2X7 in canine erythrocytes identified that ATP, relative to BzATP, was a partial agonist of native canine P2X7 (Stevenson et al., 2009). This finding is similar to that of native human P2X7, in which ATP was also reported to be a partial agonist relative to BzATP (Gargett et al., 1997). The reasons for these differences between recombinant and native canine P2X7 remain unknown but may relate to the functional assays employed or the cell types studied. The electrophysiological profile varies between human and canine recombinant P2X7, with a slower onset and decline in inward currents for canine P2X7 than human P2X7 expressed in HEK-293 cells activated by either ATP or BzATP (Roman et al., 2009). The capacity of other nucleotides to activate recombinant canine P2X7 has not been assessed. However, study of native P2X7 in canine erythrocytes demonstrates that 2MeSATP and ATP γ S, but not ADP or UTP, may also be (partial) agonists of canine P2X7 (Sluyter et al., 2007). NAD also failed to induce ethidium⁺ uptake into canine lymphocytes and monocytes (Stevenson et al., 2009), suggesting that NAD is not an agonist of canine P2X7.

Some attempts have been made to determine the tissue distribution of P2X7 in dogs. Functional P2X7 has been reported in erythrocytes, T cells, B cells, and monocytes (Sluyter et al., 2007; Stevenson et al., 2009; Jalilian et al., 2012a), as well as in kidney epithelial cells (Jalilian et al., 2012b). Of note, P2X7 activation induces the release of interleukin (IL)-1 β from canine monocytes (Jalilian et al., 2012a) and whole blood (Roman et al., 2009; Spildrejorde et al., 2014). Collectively, these studies support the potential use of dogs in future studies of P2X7. In this regard, laboratories of GlaxoSmithKline (GSK) and Pfizer have already used dogs to assess the pharmacokinetics and bioavailability of P2X7 antagonists, including 2-oxo-*N*-(phenylmethyl)-4-imidazolidinecarboxamide analogs (Abberley et al., 2010) and 2-(4-chloro-3-[3-(1-hydroxycycloheptyl)propanoyl]phenyl)-4-[(2*R*)-2-hydroxy-3-methoxy-propyl]-1,2,4-triazine-3,5-dione (CE-224,535) (Duplantier et al., 2011), respectively. The pharmacokinetics and bioavailability of these compounds in dogs were similar to that observed for monkeys (Abberley et al., 2010; Duplantier et al., 2011) and thus potentially similar to that of humans.

C. The Rhesus Macaque P2X7 Receptor

Rhesus macaque P2X7 was the first recombinant P2X7 to be synthesized from a published sequence rather than being generated via traditional cloning techniques (Bradley et al., 2011b). Both BzATP and ATP induce robust currents in cells transfected with Rhesus macaque P2X7, with EC₅₀ values similar to

that of human P2X7 (Bradley et al., 2011b). The capacity of other nucleotides or NAD to activate Rhesus macaque P2X7 has not been assessed. In regard to the latter, it is likely that NAD will also not activate Rhesus macaque P2X7, because the ART2 ortholog appears to be absent in primates (Haag et al., 1994). P2X7 protein is present in the retinal neurons of the Rhesus macaque (Ishii et al., 2003), but functional evidence and the tissue distribution of native P2X7 within these animals are lacking. Collectively, this limits the potential utility of these nonhuman primates as a model of P2X7-related diseases and pre-clinical testing of drugs targeting this receptor at this time.

V. Gating of the P2X7 Receptor Channel

The binding of agonist molecules to ligand-gated ion channels transduces a signal to the channel gate, the region of the protein involved in opening the channel pore and allowing ionic flux to occur. Similar to other P2X receptors, P2X7 receptors are trimeric in structure (Nicke, 2008; Jiang et al., 2013) and contain three intersubunit nucleotide binding sites with residues from two individual monomers both contributing to the binding site. Six helical transmembrane domains span the membrane, and it is thought that the second transmembrane domains (TM2) of each P2X7 monomer form a physical gate to the flow of ions (reviewed in Jiang et al., 2013). Structural data from P2X4 (Hattori and Gouaux, 2012) suggest a rotation of the transmembrane domains after agonist binding opening the channel pore. The external portion of TM2 has been implicated as a channel gating domain; residues Ile332 to Val343 in P2X2 correspond to residues Val335 to Leu346 in P2X7 (Li et al., 2008). Once the TM2 domains have moved, the channel pore is fully open and conducts current, the directionality of this being dictated by the membrane potential. After agonist binding and gating rearrangements, monophasic or biphasic currents can be measured by standard whole-cell electrophysiology (Yan et al., 2008, 2011; Khadra et al., 2013). Mono- and biphasic currents through P2X7 channels are distinguished by the agonist concentration and whether the channel is “naive” to agonist or not (Khadra et al., 2013). However, P2X7 gating is more complex than other ion channels because of the secondary permeability pathway (see section VI). Single channel recordings of P2X7 channels have also been used to investigate how this ion channel operates. Markwardt and colleagues measured a single channel conductance of 9–13 pS for P2X7 in *Xenopus laevis* oocytes (Riedel et al., 2007a) and have developed extensive kinetic models for the transitions between open and closed states. These kinetic models were recently extended to a 16-state Markov model, taking into account both low-high affinity agonist binding and sensitization/desensitization states (Khadra et al., 2013).

VI. Permeability of the P2X7 Receptor

P2X7 was originally identified as the P2Z receptor, capable of permeabilizing cells after ATP addition. This defining feature of P2X7 activation was thought to be unique to this receptor for many years. However it is now apparent that activation of other purinergic receptors (P2X2, P2X4, P2X5, and P2X2/5), and indeed other ligand-gated ion channels (*N*-methyl-D-aspartate receptor and the transient receptor potential cation channel subfamily A member 1 and subfamily V member 1) can also permeabilize cells (Virginio et al., 1999; Chaumont and Khakh, 2008; Chung et al., 2008; Thompson et al., 2008; Banke et al., 2010; Compan et al., 2012b). Despite these observations, it should be noted that P2X7 is the only P2X receptor where this process is consistently observed. The secondary pore pathway permits the entry of normally impermeant organic cations such as *N*-methyl-D-glucamine⁺, ethidium⁺, and YO-PRO-1²⁺. Organic anions such as lucifer yellow and carboxyfluorescein can also enter cells, although whether this involves the same permeation pathway as for cations is unclear. In addition to ion channels, the marine toxin maitotoxin (Schilling et al., 1999) and a sustained elevation of intracellular Ca²⁺ using the ionophore ionomycin (Faria et al., 2009) can also induce the opening of a pore pathway permitting impermeant dye entry.

The search to understand the mechanism by which P2X7 permits large cation/anion uptake is still ongoing, as is the debate regarding whether this is an intrinsic property of P2X7 or whether it requires at least one other protein. Several studies have shown that the unique and characteristic long C-terminal tail of P2X7 is crucial for pore formation. Truncation of P2X7 abolishes dye uptake without affecting ion channel currents (Surprenant et al., 1996; Rassendren et al., 1997; Smart et al., 2003). Single point mutations due to polymorphisms can also dramatically affect the dye uptake response (Gu et al., 2001; Adriouch et al., 2002; Wiley et al., 2003). In addition, there is reported to be some dependence on second messenger pathways, including a role for Ca²⁺ (Donnelly-Roberts et al., 2004; Faria et al., 2005). These studies and others (reviewed in Rokic and Stojilkovic, 2013) have attempted to dissect out ion channel and secondary pore pathways. A breakthrough in mechanistic understanding came with the identification of pannexin-1 as a secondary protein involved in dye uptake (Pelegriin and Surprenant, 2006). However, there is now growing evidence suggesting that pannexin-1 cannot be the major pore pathway used by P2X7. For example, no defect in ATP-mediated YO-PRO-1²⁺ uptake was observed in macrophages from pannexin-1 knockout (KO) mice (Qu et al., 2011), siRNA knockdown of pannexin-1 in mouse macrophages failed to affect P2X7 responses (Alberto et al., 2013), and pharmacological blockade of pannexin-1 did not affect ATP-induced dye uptake in transfected HEK-293 cells or human monocytes (Bhaskaracharya et al., 2014).

Electrophysiological recordings of P2X7 currents suggest that biphasic currents reflect pore dilation (Yan et al., 2008; Khadra et al., 2013), and therefore a large proportion of the inward current is via the secondary permeability pathway. However, biphasic currents have not been recorded under single channel recording conditions (Riedel et al., 2007b). Recently, Browne et al. (2013) provided evidence for direct permeation of large molecules through the P2X7 channel pore. By introducing cysteine mutations into TM2, which lines the conductance pathway of the P2X7 channel pore, the direct access of methanethiosulfonate reagents to these residues after ATP treatment was demonstrated. Inhibition of P2X7 ion channel currents using methanethiosulfonate reagents also blocked fluorescent dye uptake (Browne et al., 2013).

Given the importance of the secondary permeability pathway in signaling events downstream of P2X7 activation (see section VII), it is necessary that a clear understanding of this secondary pathway is established. Moreover, we must also consider the physiologic relevance of this additional pore pathway allowing influx/efflux of large molecules. Is this a mechanism for cells to take up ATP or other molecules required for an increased metabolic demand to respond to tissue injury or inflammation? Is this a mechanism for ATP release to drive further purinergic signaling? Or does this pathway provide a way of secreting chemical transmitter molecules such as glutamate? Cervetto and colleagues (2013a) recently demonstrated that glutamate efflux from P2X7-transfected HEK-293 cells was dependent on the long C terminus of rat P2X7 and the Cys-rich domain of the proximal C terminus, implicating the secondary pore pathway in efflux events.

VII. P2X7 Receptor-Dependent Signaling Pathways

A number of cell type-specific signaling pathways have been associated with P2X7 activation. These include the activation of the caspase-1-containing inflammasome NLRP3 and the subsequent processing and release of proinflammatory interleukin-1 (IL-1) cytokines, as well as the formation of reactive oxygen and nitrogen species and the formation of phagolysosomes and subsequent killing of intracellular pathogens (see Hewinson and Mackenzie, 2007; Miller et al., 2011; Di Virgilio, 2013). Here we will review recent developments in established signaling pathways and novel signaling pathways associated with P2X7.

A. Cytokine Release

In addition to the secretion of leaderless cytokines IL-1 β (IL-1F2) and IL-18 (IL-1F4), P2X7 activation has also been shown to induce release of IL-1 α (IL-1F1) (Pelegri et al., 2008), IL-1 receptor antagonist (IL-1F3) (Wilson et al., 2004, 2007; Glas et al., 2009) and IL-36 α

(IL-1F6) (Martin et al., 2009). This places P2X7 as a major physiologic regulator of secretion of the IL-1 family of cytokines. A recent comprehensive investigation into the mechanism underlying NLRP3 inflammasome activation and IL-1 β secretion by diverse stimuli, including P2X7, suggests K⁺ efflux is a common signal activating the NLRP3 complex (Muñoz-Planillo et al., 2013). Extracellular osmolarity and cell volume regulation also plays a role in the activation of NLRP3 inflammasome (Compan et al., 2012a), highlighting that it is the cellular microenvironment that is crucial for switching on cytokine secretion.

B. Reactive Oxygen Species Formation

The generation of reactive oxygen species (ROS) after ATP activation of P2X7 is now well established in macrophages (Lenertz et al., 2009; Moore and MacKenzie, 2009), microglia (Apolloni et al., 2013; Bartlett et al., 2013), submandibular gland cells (Seil et al., 2008), and erythroid cells (Wang and Sluyter, 2013). Activation of NADPH oxidase 2 by P2X7 was a common feature found in all cell types studied, although there is evidence that ATP, possibly via P2X7 activation, can also enhance mitochondrial reactive oxygen species (Nakahira et al., 2011). One downstream consequence of increased ROS generation in myeloid cells is the activation of the NLRP3 inflammasome and IL-1 β secretion (reviewed in Gross et al., 2011; Tschopp, 2011). However, in a more recent study, ROS were not found to play a role in NLRP3 activation in bone marrow-derived macrophages (Muñoz-Planillo et al., 2013). Rather, a reduction of the intracellular K⁺ concentration was required. Despite this, there is evidence that K⁺ channel activity can be positively modulated by ROS (Ichinari et al., 1996; Avshalumov and Rice, 2003; Sesti et al., 2010), and it may be that ROS are able to induce NLRP3 activation in some cell types via this signaling process.

C. Protease Activation and Release

Activation of P2X7 receptors can also lead to a number of membrane-related changes including membrane blebbing, microparticle and exosome release, multinucleated cell formation, reversible phosphatidylserine exposure, and the activation of membrane metalloproteases resulting in the shedding of cell-surface molecules (see Qu and Dubyak, 2009; Wiley et al., 2011). Recent work has identified additional cell surface molecules shed from the cell surface or released after activation of P2X7. This includes the chemokine CXCL16 (Pupovac et al., 2013a), the adhesion molecule CD44 (Lin et al., 2012), soluble amyloid precursor protein (Delarasse et al., 2011), and the IL-6 receptor (Garbers et al., 2011), in addition to CD62L (L-selectin), CD23 (the low-affinity IgE receptor) (Gu et al., 1998), and CD27 (Moon et al., 2006). Two metalloproteinases, ADAM10 and ADAM17, are the main effector molecules associated with these P2X7-mediated shedding events (Garbers et al., 2011;

Pupovac et al., 2013a). Many physiologically relevant substrates are cleaved by these proteases, and it will be interesting to determine if P2X7 can induce cleavage of all ADAM10/17 substrates or only a subset.

P2X7 activation can also induce the release of various enzymes. Andrei and colleagues (2004) first identified that ATP could induce lysosome exocytosis from human monocytes and measured cathepsin D secretion as a readout. It is now clear that lysosome secretion of cathepsins is a P2X7-mediated pathway likely regulated by increased intracellular Ca^{2+} concentrations. Cathepsins B, D, L, and S are also secreted in this manner from monocytes and macrophages (Lopez-Castejon et al., 2010). P2X7 activation can also stimulate the release of metalloprotease-9 (Gu and Wiley, 2006); however, the specific release mechanism is yet to be ascertained. Therefore, P2X7 can regulate several families of proteases, namely caspases (Ferrari et al., 1999), as well as cathepsins and matrix metalloproteinases.

D. Prostaglandin Release

Prostaglandin E_2 (PGE_2), thromboxane B_2 , and leukotriene B_4 are prostanoids induced by P2X7 signaling in murine macrophages (Barberà-Cremades et al., 2012). Phospholipases mediate the hydrolysis of arachidonic acid from various phospholipids, and P2X7 is known to activate cytosolic phospholipase A_2 (Andrei et al., 2004) and Ca^{2+} -insensitive phospholipase A_2 isoforms (El Ouailiti et al., 2012), in addition to phospholipase D (el-Moatassim and Dubyak, 1992). Arachidonic acid is then a substrate for cyclooxygenase and prostaglandin E synthases in the biosynthesis of PGE_2 . PGE_2 plays an important role in inflammation, fever, and pain, and therefore is likely to be an important downstream signaling pathway from P2X7 in vivo. Given the activation of P2X7 by ATP in areas of tissue damage, this may place P2X7 as a potential alternative anti-inflammatory target to cyclooxygenase (Barberà-Cremades et al., 2012).

E. Glutamate Efflux

Early reports exist showing that P2X7 induces non-vesicular release of excitatory amino acids such as glutamate and aspartate from murine astrocytes (Duan et al., 2003; Fellin et al., 2006). This finding was recently extended with the demonstration that P2X7 activation could induce glutamate efflux from rat cortical nerve terminals (Marcoli et al., 2008) and from HEK-293 cells expressing recombinant rat P2X7 (Cervetto et al., 2013a). Such contributions to extrasynaptic and/or synaptic glutamate concentrations may alter neurotransmission and could contribute to changes in neuroplasticity and behavior.

F. Transcription Activation

P2X7 can influence gene expression through activation of a variety of transcription factors. Several studies have identified nuclear factor- κB p65 as a transcription

factor regulated by P2X7 in microglia (Ferrari et al., 1997) and osteoclasts (Korcok et al., 2004). In monocytes, P2X7 activation can also induce cAMP response element-binding protein activation (Gavala et al., 2008) and FosB/activating protein-1 (Gavala et al., 2010), and in T cells, P2X7 activation can induce the translocation of nuclear factor of activated T cells to the nucleus (Yip et al., 2009). Therefore P2X7 may regulate gene expression in response to extracellular ATP, perhaps to reprogram cellular metabolism in times of stress or to switch on genes required for the inflammatory response.

G. Cell Proliferation

P2X7 is historically known for its ability of ATP to induce cell death through apoptosis (Zanovello et al., 1990; Chow et al., 1997; Ferrari et al., 1999). A role in cell proliferation would seem to be at odds with this, but evidence suggests that in certain cell types, activation of P2X7 is more likely involved with proliferation than cell death. This may be due to expression of particular splice variants (see section VIII), and this aspect certainly warrants further investigation in T cells and microglia. In T cells, release of autocrine ATP during activation through the T-cell receptor activates P2X7 and enhances secretion of the T-cell growth factor IL-2, through stimulation of the nuclear factor of activated T-cells transcription factor (Yip et al., 2009). Baricordi and colleagues (1996) first suggested a role for P2X7 in mitogen-induced T-cell proliferation. This same group subsequently went on to show that transfection of cell lines with full-length P2X7 or a truncated splice variant of this receptor increased cell proliferation, especially in the absence of serum (Baricordi et al., 1999; Adinolfi et al., 2005, 2010). In microglia, blocking P2X7 pharmacologically or overexpressing a pore-mutant (G345Y) to abolish secondary permeability reduced the proliferative response (Bianco et al., 2006; Monif et al., 2009). It will be interesting to determine if the trophic effect in microglia is also due to transcriptional regulation of a growth factor.

H. Phagocytosis

A novel cellular signaling process associated with P2X7 is the regulation of phagocytosis (Wiley and Gu, 2012), a process by which foreign particles or organisms are taken up into cells. Transfection of HEK-293 cells, which are not normally phagocytic cells, with DNA encoding P2X7 introduces the ability to take up 1- μm latex beads or fluorescently labeled bacteria (Gu et al., 2010). Gu and colleagues (2009, 2010, 2011, 2012) propose that P2X7 is acting as a scavenger receptor. Typically, scavenger receptors, such as CD36, recognize modified lipoproteins, such as low-density lipoproteins (Endemann et al., 1993), and although it is noted that P2X7 has a similar two transmembrane domain structure to CD36 (Gu et al., 2011, 2012), whether P2X7 can recognize such lipoproteins is not yet known. The ability of P2X7-expressing cells to take up particles does not appear

to require classic activation of the ion channel by extracellular ATP. In fact, stimulation of the channel with ATP switches off this phagocytic pathway (Gu et al., 2010). This suggests that particle uptake does not require classic signaling through the ion channel or the secondary pore pathway, but may use a novel signaling pathway due to association of P2X7 with the cytoskeleton (Gu et al., 2009, 2010). This is an exciting development in P2X7 cell biology, and further investigations may reveal other ligands for this receptor.

I. Summary

In summary, P2X7 regulates ionic flux, protease activation, membrane events, and various secretion events. This is a diverse series of signaling pathways commonly involved in inflammation. Because these signaling pathways are cell type specific, it is highly likely that each cell type expressing P2X7 has a different complement of downstream effectors. This information is critical as we try to understand the role of P2X7 in disease and the potential for this receptor as a therapeutic target.

VIII. Variants of the P2X7 Receptor

In the last 10 years, particular attention has been drawn to the importance and contributions of P2X7 variants to the diversity of P2X7 receptor-mediated responses in health and disease, bringing the purinergic field to a point where these variants can no longer be ignored. In particular, these studies have highlighted the need for the consideration of variants in current and future in vivo rodent studies and human clinical trials. The two main causes of P2X7 variants, P2X7 SNPs and splice isoforms, as well as their implications for rodent studies, will be discussed below. The human P2X7 receptor will be discussed first because the largest body of work has been done in this species.

A. P2X7 Single Nucleotide Polymorphisms

The human *P2RX7* gene is highly polymorphic, with over 1500 SNPs reported in the National Center for Biotechnology Information (NCBI) SNP database (www.ncbi.nlm.nih.gov/sites/entrez, accessed December 27, 2013). However, the majority of these SNPs are intronic, with approximately 150 nonsynonymous (or missense) SNPs reported in both this database and the Ensembl Genome Browser (asia.ensembl.org/index.html, accessed December 27, 2013). Of these nonsynonymous SNPs, a small number have been investigated to date that lead to altered human P2X7 function. A loss-of-function effect has been reported for SNPs V76A, R117W, G150R, E186K, L191P, R276H, R307Q, T357S, E496A, and I568N (Gu et al., 2001, 2004; Wiley et al., 2003; Shemon et al., 2006; Roger et al., 2010b; Stokes et al., 2010). Because of the absence of frequency data it is not clear whether R117W, E186K, or L191P represent SNPs or mutations. Investigations into the underlying mechanisms

behind the loss-of-function effect of individual SNPs have revealed differences depending on the site of the mutations. Residue 307, originally thought to contribute to ATP binding (Gu et al., 2004), may instead be involved in hydrogen bonding and salt bridge structures in the P2X7 ectodomain (Jiang et al., 2013). Residue 568 lies within a trafficking motif, and mutant 568N-P2X7 receptors do not efficiently traffic to the cell membrane (Wiley et al., 2003). It is currently unclear how V76A, R117W, G150R, L191P, R276H, and E496A mutations cause a loss-of-function effect. SNPs in regulatory and intronic regions have also been identified, including the upstream promoter region (−762; rs2393799) (Li et al., 2002), an intronic splicing site at the exon1/intron1 boundary (rs35933842) (Skarratt et al., 2005), and the 3'-untranslated region affecting a microRNA binding site (rs1653625) (Rahman et al., 2010). However, only rs35933842 has been associated with a loss-of-function effect.

A gain-of-function effect has been reported for three SNPs: H155Y, H270R, and A348T (Cabrini et al., 2005; Stokes et al., 2010). The underlying mechanisms behind these gain-of-function effects have not been established. There is some evidence to suggest that the amino acid variation at 155 increases cell surface expression of P2X7 (Bradley et al., 2011a); however, this was not observed in the initial report of this SNP (Cabrini et al., 2005). One of the most interesting SNPs in human *P2RX7* is Q460R in the C terminus. In isolation this mutation does little to the functional response, but our investigations revealed coinheritance with gain-of-function SNPs in a particular haplotype (Stokes et al., 2010), perhaps explaining the association of this SNP with various disorders, including bipolar disorder (Barden et al., 2006), depression (Lucae et al., 2006), and Sjogren's syndrome (Lester et al., 2013).

In the last 5 years many studies have investigated the association of SNPs in *P2RX7* with a number of different diseases. Some studies have used individual key SNPs such as E496A or Q460R, whereas others have used a more informative 12 SNP approach. Loss-of-function haplotypes have been linked to susceptibility to extrapulmonary tuberculosis infection (Fernando et al., 2007), enhanced bone mineral density loss in postmenopausal women (Gartland et al., 2012; Jorgensen et al., 2012), and an increased risk of age-related macular degeneration due to reduced phagocytosis of cellular debris (Gu et al., 2013). Inheritance of E496A has been linked to a reduced risk of ischemic stroke and ischemic heart disease (Gidlof et al., 2012). The gain-of-function A348T SNP is protective for *Toxoplasmosis gondii* infection (Jamieson et al., 2010) and is found at an increased frequency in Arabic patients with rheumatoid arthritis (Al-Shukaili et al., 2011) and patients with major depressive disorder (Lucae et al., 2006). Q460R, which is in linkage disequilibrium with A348T, was found to be increased in patients with

Sjogren's syndrome and was linked to increased seropositive disease (Lester et al., 2013).

In summary, gain-of-function haplotypes of human P2X7 may protect against infection and premature bone loss, but may increase the risk of developing certain inflammatory and psychiatric disorders. On the converse, carrying loss-of-function haplotypes of P2X7 may protect against inflammatory disorders, such as heart disease, but may increase susceptibility to some infections.

A number of SNPs have similarly been reported in the murine *P2RX7* gene region, with a survey of the NCBI SNP database revealing >1700 SNPs (accessed December 27, 2013). The majority of these SNPs are intronic. With the Ensembl Genome Browser (accessed December 27, 2013), 34 SNPs were revealed to be present in the coding region of the full-length murine *P2RX7* gene, 10 of which are nonsynonymous (Fig. 1).

Of the nonsynonymous SNPs reported for murine P2X7 in the NCBI SNP database and the Ensembl Genome Browser, only the P451L SNP has been characterized. This naturally occurring mutation leads to the replacement of the proline residue at position 451 by a leucine residue in the cytoplasmic tail of murine P2X7 (Adriouch et al., 2002). The distribution of the allelic versions of this SNP in different mouse strains is outlined in Fig. 2. The P451L SNP was originally identified based on studies of P2X7 function in T cells from different strains of mice, whereby T cells from BALB/c mice (451P) had significantly higher sensitivity to ATP than those from




Fig. 2. Allelic versions of the loss-of-function P2X7 P451L SNP in different mouse strains (based on data from Adriouch et al., 2002; Sorge et al., 2012; Syberg et al., 2012). The 451P allele is also observed in rats and humans. Strains in italics are wild mouse strains. NZW, New Zealand White; NOD, nonobese diabetic.

C57BL/6 mice (451L) (Adriouch et al., 2002). By transfecting HEK-293 cells with either of the two variants (451P or 451L), it was demonstrated that P2X7 channel and pore function were markedly reduced in cells expressing the mutated 451L allele. This was later confirmed in another study, in which HEK-293 cells expressing 451L had reduced cation dye uptake and Ca^{2+} flux compared with cells expressing 451P (Young et al., 2006). However, 451L was not found to be associated with reduced surface protein expression or maximum agonist-induced currents in this latter study. In contrast, human astrocytoma 1321N1 cells expressing 451L are reported to have fully functional P2X7, with a similar pharmacological profile to those expressing 451P (Donnelly-Roberts et al., 2009).

Differences in murine P2X7 function encoded by the 451P and 451L alleles are further supported by two additional studies with BALB/c and C57BL/6 mice, respectively. First, thymocytes from C57BL/6 mice display significantly impaired P2X7-induced cell death compared with those from BALB/c thymocytes (Le Stunff et al., 2004). In contrast, P2X7-induced phospholipase D stimulation was similar in thymocytes from both strains (Le Stunff et al., 2004). This suggests that P2X7-induced cell death and phospholipase D stimulation are controlled through distinct domains of P2X7. Second, astrocytes from C57BL/6 mice have attenuated ATP-induced pannexin-1 currents, ATP release, and intracellular Ca^{2+} waves compared with astrocytes from BALB/c mice, all of which were thought to be mediated by P2X7 (Suadicani et al., 2009). Of note, using a Src inhibitor and electrophysiological, biochemical, and fluorescence imaging techniques, this same study also provided evidence that P451L is located within the SH3 domain, a Src tyrosine kinase-binding site. This further supports the potential influence of this SNP on domain structure and function of the murine P2X7 receptor.




Fig. 1. Nonsynonymous SNPs reported in the coding region of the murine *P2RX7* gene. SNP, SNP ID, and alleles are as follows: S6N, rs227977285, and G/A; V32I, rs229167308, and G/A; E70Q, rs238213067, and G/C; G113D, rs37541158, and G/A; T117M, rs238065774, and C/T; R270Q, rs220799687, and G/A; P396R, rs37237164, and C/G; P451L, rs48804829, and C/T; R486H, rs37867325, and G/A; H514R, rs36331015, and A/G. Yellow and red circles refer to functionally characterized and noncharacterized SNPs, respectively. SNPs were identified using Ensembl Genome Browser (asia.ensembl.org/index.html, accessed December 27, 2013).

The P451L SNP has been correlated with disease in two studies. First, genome-wide linkage analysis studies have found an association between the P451L SNP and neuropathic pain in mice (Sorge et al., 2012). In this study, mice from a number of different strains expressing the 451L allele showed less mechanical allodynia, a symptom of neuropathic pain, after spared nerve injury compared with those with the 451P allele. Second, an association between the P451L SNP and bone phenotype has been reported (Syberg et al., 2012). Strains with the 451L allele (C57BL/6 and DBA/2J) had weaker femurs, lower amounts of the bone resorption marker C-telopeptide collagen, and lower bone mineral densities compared with those strains with the 451P allele (BALB/cJ and 129X1/SvJ). However, the cellular bases for these differences were not established, whereas other measured bone parameters were not associated with the P451L SNP.

In addition to the naturally occurring P451L SNP, nonsynonymous mutations have also been identified in murine cell lines. For example, L11F, T221A, and T283M mutations have been found in P2X7 cloned from the NTW8 microglial cell line (Chessell et al., 1998; Young et al., 2006). These mutations have been associated with reduced function in this cloned receptor compared with the function of recombinant P2X7 cloned from a C57BL/6 mouse (Adriouch et al., 2002). By introducing point mutations, the T283M SNP, but not L11F or T221A SNPs, was found to be responsible for this low P2X7 function, with expression of the 283M variant leading to both reduced channel and pore activity (Young et al., 2006). In comparison, the 283T allele is present in P2X7 sequences from other mouse strains, as well as the rat and human sequences. Thus, the T283M mutation may be specific to the NTW8 cell line or a result of cloning.

Further survey of the NCBI SNP database and Ensembl Genome Browser (both accessed December 27, 2013) reveals few nonsynonymous SNPs in the coding region of the other cloned mammalian P2X7 receptors. Nonsynonymous SNPs in the rat or guinea pig *P2RX7* gene have not been reported. Two nonsynonymous SNPs, L103F and P452S, have been reported in the canine *P2RX7* gene. In the other cloned mammalian P2X7 receptors, a phenylalanine residue resides at amino acid position 103. Thus, it is likely that 103F may also represent the more common variant in dogs and that the SNP at this position may be more accurately defined as F103L rather than the reported L103F. Nevertheless the cloned canine P2X7, which contains a leucine residue at amino acid position 103, forms a functional receptor (Roman et al., 2009), although its relative effect compared with canine P2X7 containing residue 103F remains unknown. The P452S SNP in the canine *P2RX7* gene is located at the equivalent position as the partial loss of function SNP, P451L, of the murine *P2RX7* gene. The functional effect of this canine SNP remains unknown.

Finally, one nonsynonymous SNP, I75M, has been reported in the Rhesus macaque *P2RX7* gene. An isoleucine residue is found at this amino acid position in each of the other cloned mammalian P2X7 receptors, and the functional effect of a methionine at this position remains to be determined.

B. *P2X7* Splice Isoforms

Alternative splicing is a process that enables one gene to produce multiple protein isoforms, with a number of alternative P2X7 splice variants identified for the human and rodent receptor to date. In humans, seven naturally occurring splice variants were initially reported and termed P2X7B to P2X7H (Cheewatrakoolpong et al., 2005), with P2X7A referring to the original full-length 595 amino acid protein (Rassendren et al., 1997). These include truncated variants that have an alternate short C terminus (P2X7B, C, E, and G), variants that lack exons responsible for coding parts of the extracellular domain (P2X7C–F) and variants that have additional exons, thought to result in P2X7 protein lacking the first transmembrane domain (P2X7G and H) (see Sluyter and Stokes, 2011). An additional two naturally occurring splice variants have been identified in humans, termed P2X7I and P2X7J (Feng et al., 2006). P2X7I codes for a null allele and is a result of SNP rs35933842 (Skarratt et al., 2005), whereas P2X7J is another truncated variant (Feng et al., 2006). These ten variants of human P2X7 differ in distribution, functional characteristics, and predicted physiologic roles. Of these, P2X7B is the best studied and is able to form functional channels but not the large pores that are often associated with inflammation and cell death (Adinolfi et al., 2010). P2X7J is similarly unable to form pores and also displays reduced ligand-binding ability and channel function and oligomerizes with P2X7A to form nonfunctional P2X7 heterotrimers (Feng et al., 2006).

Naturally occurring P2X7 splice variants have also been identified in rodents. The first reported, originally termed P2X7(K) (Nicke et al., 2009) or P2X7K by others (Masin et al., 2012), results from a novel exon 1 in the rodent *P2RX7* gene. This leads to a receptor with an alternate amino-terminal and first transmembrane domain (Nicke et al., 2009). In contrast to the human P2X7 variants mentioned above, P2X7K is a fully functional variant, with an 8-fold higher sensitivity to P2X7 agonists, slower deactivation, and increased propensity to form P2X7 pores compared with P2X7A. Furthermore, it was recently reported that this variant is sensitive to activation by extracellular NAD, whereas P2X7A is not (Schwarz et al., 2012). This is the first report suggesting that NAD is unable to activate P2X7A, which complicates the interpretation of earlier NAD studies. In this study, T cells were found to preferentially express P2X7K, whereas macrophages preferentially express P2X7A. This may account for differences in P2X7 signaling between these cell types in rodents (Hong

et al., 2009; Xu et al., 2012) and suggests that P2X7 variants may mediate the differential sensitivity of macrophages and lymphocytes to NAD. However, the relative contribution of different P2X7 splice variants on NAD-mediated P2X7 activation in other cell types, such as astrocytes, has not been investigated. To the best of our knowledge, a similar variant to P2X7K has not been described in humans.

C-terminal truncated P2X7 variants have also been identified in mice (Masin et al., 2012). These variants, termed P2X7 13B and P2X7 13C, are encoded by genes with alternative exon 13s, resulting in truncated or alternate short C termini, respectively. P2X7 13B was detected in all tissues tested, with strongest expression in the brain, salivary glands, and spleen, whereas P2X7 13C was present in the brain, lungs, and salivary glands. When expressed in HEK-293 cells, both variants displayed low cell surface expression and channel function and no pore formation. This low cell surface expression of P2X7 13B was due to inefficient trafficking to the cell surface and, similar to the human variant P2X7J (Feng et al., 2006), appeared to inhibit the function of full-length P2X7 through hetero-oligomerization (Masin et al., 2012). However, equivalent variants of human P2X7B-J are yet to be reported in rodents. Furthermore, the influence of these variants on P2X7 in vivo or even native P2X7 on primary cells has not been widely examined, and this may be a confounder in preclinical studies in rodents, particularly if the sensitivity of these variants to P2X7 antagonists varies.

C. P2X7 Variants in P2X7 Knockout Mice

There are currently two strains of P2X7 KO mice available: the GSK line generated by Chessell et al. (2005) (see also Sim et al., 2004) and the Pfizer line generated by Solle et al. (2001). The GSK KO strain was generated by inserting a lacZ transgene and neomycin cassette into exon 1, whereas the Pfizer KO was generated by inserting a neomycin cassette into exon 13, replacing part of the C terminus of the receptor. These models have been extensively used for demonstrating P2X7 involvement in inflammatory and neuropathic pain (GSK KO), cytokine production, inflammation, bone formation, and a number of disorders (Pfizer KO) (see Volonté et al., 2012). However recent evidence, coinciding with the discovery of P2X7 splice variants, indicates that these mice may not be true knockouts.

First, molecular and biochemical analyses indicated that the widely expressed P2X7K splice variant discussed above escaped deletion in the GSK KO, with highest expression in the spleen (Nicke et al., 2009). This is likely because of the fact that exon 1 of the *P2RX7* gene was targeted in the generation of this KO strain, whereas the P2X7K variant encodes a novel exon 1. Although P2X7 activity is absent in macrophages and dendritic cells from this KO strain, T cells displayed higher levels of P2X7 activity than that of control mice

(Taylor et al., 2009a). Second, molecular and biochemical analyses indicated that P2X7 13B and 13C C-terminal truncated variants, as well as a C-terminal truncated hybrid variant containing a short sequence of the targeting vector used to disrupt exon 13 of the *P2RX7* gene, escaped deletion in the Pfizer KO mouse (Masin et al., 2012). However, unlike T cells from the P2X7K-expressing GSK KO, those from the Pfizer KO have reduced receptor function (Taylor et al., 2009a).

The tissue-specific expression of functional P2X7 in these KO strains may explain the controversial staining of P2X7 in brain tissue from KO animals, which is identical to that of wild-type (WT) littermates (Sim et al., 2004; Sánchez-Nogueiro et al., 2005). In these studies, P2X7 antibodies may be detecting splice variants, which have escaped deletion, rather than P2X7-like proteins with similar immunogenicity or other proteins through nonspecific staining. In addition, these variants may explain some of the phenotypic and functional differences observed between these two KO strains (Fig. 3). Although P2X7 variants have escaped deletion in both the GSK and Pfizer KOs, overall evidence indicates that the P2X7A variant has been knocked out in both strains. Despite this, caution needs to be undertaken when interpreting results of in vivo studies that use these KO mice.

IX. Modulators of P2X7 Receptor Activation

There are a number of P2X7 antagonists commercially available, all of which vary in terms of chemical structure, mode of antagonism, specificity, and species selectivity. The first generation of compounds available that was capable of antagonizing P2X7 were generally nonspecific, with many also capable of inhibiting other P2X and in some cases P2Y receptors or other molecules (see Gever et al., 2006). These early antagonists include Reactive Blue 2, which was first used in the 1970s (Kerr and Krantis, 1979); suramin and Brilliant Blue G (BBG), which were first used in the 1980s (Dunn and Blakeley, 1988; Soltoff et al., 1989); and pyridoxal phosphate-6-azophenyl-2-4-disulfonic acid (PPADS), periodate-oxidized ATP (oATP), and 1-[*N,O*-bis(5-isoquinolinesulfonyl)-*N*-methyl-*l*-tyrosyl]-4-phenylpiperazine (KN-62), which were first used in the 1990s (Lambrecht et al., 1992; Murgia et al., 1993; Gargett and Wiley, 1997). Following on from their identification as P2X7 antagonists, structural modifications of these compounds were carried out in an attempt to improve their properties and specificity (see Romagnoli et al., 2008; Jiang, 2012). For example, MRS 2540 is based upon the structure of KN-62, and exhibits enhanced potency at both human and rodent P2X7 (Lee et al., 2008a; Hu et al., 2010). Of the earlier compounds, BBG remains one of the most useful. This compound is generally considered a specific P2X7 antagonist, blocking rat P2X7 in the nanomolar compared with micromolar range or not at all




Fig 3. Phenotypic and functional differences between GlaxoSmithKline (GSK) and Pfizer knockout (KO) mice compared with wild-type (WT) counterparts. Differences in P2X7 isoforms expressed, bone phenotype (mineral density, volume, and thickness), P2X7-mediated responses in T cells (pore formation, nucleotide-induced CD62L shedding, phosphatidyserine exposure, and cell death), and experimental autoimmune encephalomyelitis (EAE) between P2X7 KO mice generated by GSK and Pfizer compared with WT. ^a(Solle et al., 2001; Sim et al., 2004; Nicke et al., 2009; Masin et al., 2012); ^b(Gartland et al., 2003; Ke et al., 2003); ^c(Labasi et al., 2002; Taylor et al., 2008, 2009a); ^d(Chen and Brosnan, 2006; Sharp et al., 2008).

for other rat P2X receptors (P2X1, P2X2, P2X3, P2X2/3, P2X4, and P2X1/5) (Jiang et al., 2000). More recent data indicate that BBG also blocks pannexin 1 (Qiu and Dahl, 2009), complicating its use as a P2X7 antagonist. oATP is also widely used to antagonize P2X7, but has also been shown to inhibit other P2X receptors (including P2X2 and P2X3) (Evans et al., 1995) and attenuate pro-inflammatory signaling through P2 receptor-independent mechanisms (Beigi et al., 2003). However, the ability of oATP to bind irreversibly to P2X7 via covalent modification of the receptor (Murgia et al., 1993) makes this antagonist an attractive blocker of P2X7 in vivo. Likewise, PPADS, which can also bind irreversibly to P2X receptors (Nicke et al., 1998), also remains an attractive blocker of P2X7 in vivo.

In the last two decades, a second generation of more specific antagonists has been developed (see Friedle et al., 2010; Bhattacharya et al., 2011; Jiang, 2012). This has been aided by high-throughput screening of large chemical libraries and has largely occurred alongside reports indicating important roles for P2X7 in pain and inflammation (Romagnoli et al., 2008). These newer drug-like antagonists are particularly important for the purinergic field, because the first generation of antagonists was generally designed for in vitro, rather than therapeutic, use. Furthermore, many of the original antagonists are subject to degradation when used in vivo or are predicted to have poor pharmacokinetics (Friedle et al., 2010; Jiang, 2012), problems avoided by many of the second generation P2X7 antagonists. Second generation P2X7 antagonists that have been characterized include the tetrazole/triazole-based compounds 3-[[5-

(2,3-dichlorophenyl)-1*H*-tetrazol-1-yl]methyl]pyridine (A438079) (Nelson et al., 2006) and 1-(2,3-dichlorophenyl)-*N*-[[2-(2-pyridinyloxy)phenyl]methyl]-1*H*-tetrazol-5-amine (A839977) (Honore et al., 2009), the adamantane amides *N*-[2-[[2-[(2-hydroxyethyl)amino]ethyl]amino]-5-quinoliny]-2-tricyclo[3.3.1.1^{3,7}]dec-1-ylacetamide (AZ10606120) (Michel et al., 2007), and *N*-(adamantan-1-ylmethyl)-5-[(3*R*-amino-pyrrolidin-1-yl)methyl]-2-chloro-benzamide (AACBA; also known as GSK314181) (Broom et al., 2008), the cyanoguanidine derivative *N*-(1-[[[cyanoimino](5-quinolinylamino) methyl] amino]-2,2-dimethylpropyl)-2-(3,4-dimethoxyphenyl)acetamide (A740003) (Honore et al., 2006), the cyclic imide 3-[1-[[[3'-nitro[1,1'-biphenyl]-4-yl]oxy]methyl]-3-(4-pyridinyl)propyl]-2,4-thiazolidinedione (AZ11645373) (Stokes et al., 2006), and the nicotinamide derivative *N*-[[4-(4-phenyl-piperazin-1-yl)tetrahydro-2*H*-pyran-4-yl]-methyl]-2-(phenyl-thio) nicotinamide (JNJ-47965567) (Letavic et al., 2013). The concentration responses of some of these specific antagonists (including A438079 and AZ11645373) have been determined at native murine P2X7 receptors (Bartlett et al., 2013; Pupovac et al., 2013b), which is useful for their in vivo application. However, despite the availability of these newer P2X7 antagonists, many in vivo studies still use first generation antagonists (see section X). Moreover, there are a number of additional potential antagonists identified that have yet to be fully characterized (Donnelly-Roberts and Jarvis, 2007).

One factor confounding pharmacological studies of P2X7 is the differential activity of P2X7 antagonists at receptors from different species (Table 2). For example, KN-62 at nanomolar concentrations potently inhibits

all cloned mammalian P2X7 receptors, except rat, when expressed in HEK-293 cells (Fonfria et al., 2008; Donnelly-Roberts et al., 2009; Roman et al., 2009; Bradley et al., 2011b). In comparison, the commonly used in vivo P2X7 antagonist BBG is at least a 3-fold more potent antagonist of rat, murine, canine, or guinea pig P2X7 than human P2X7 (Fonfria et al., 2008; Donnelly-Roberts et al., 2009; Roman et al., 2009). Although these and similar studies reveal important insights into the specificity and selectivity of P2X7 antagonists, many are limited to examining recombinant P2X7 expressed in mammalian cells. It is unknown if the functional characteristics of P2X7 in these studies deviate from that of native P2X7 and, if so, to what degree. In addition, in vitro pharmacological results, and in particular IC₅₀ values, are highly dependent on a number of factors. These include the assay (e.g., whether studying channel opening or pore formation), assay buffer (e.g., presence of divalent cations and bovine serum albumin can be inhibitory), host cell or cell type, agonist concentration, assay temperature, and antagonist mechanism of action (e.g., competitive or noncompetitive, reversible or irreversible, orthosteric or allosteric) (Hibell et al., 2001; Donnelly-Roberts and Jarvis, 2007). Of the antagonists listed in Table 2, PPADS, BBG, Reactive Blue, KN-62, and AZ10606120 are noncompetitive inhibitors of P2X7 (Jiang et al., 2000; Hibell et al., 2001; Honore et al., 2006; Felix et al., 2012). In contrast, A438079, A740003, and AZ11645373 are competitive inhibitors of P2X7 (Alcaraz et al., 2003; Nelson et al., 2006; Yan et al., 2011), whereas the mechanism of action of JNJ-47965567 remains to be determined. Collectively, all of the above have implications for direct pharmacological comparisons between in vitro studies but, more importantly, for the use of these antagonists in vivo. In this regard, the in vivo

cellular environment may result in drastically different outcomes of P2X7 antagonists to those outcomes observed in vitro. In addition, an efficacious antagonist in an in vivo rodent model may not translate to a human therapeutic because of the differential activity of P2X7 antagonists at receptors from different species. Furthermore, differences between individuals may highly impact upon the efficacy of these compounds. Thus a number of factors need to be considered when using P2X7 antagonists in vivo.

In addition to the above experimental differences, variations in amino acid homology in the ectodomain or presumed antagonist binding site between species may be one explanation for the differential activity of P2X7 agonists and antagonists at receptors from different species (Young et al., 2007; Michel et al., 2008). For example, one ectodomain residue (N284D) accounts for the 10-fold difference in ATP sensitivity between rat and mouse P2X7 (Young et al., 2007), whereas two other ectodomain residues (F95L and R126G) are important for the different antagonist effects of KN-62 and PPADS, respectively, at the human and rat receptor (Michel et al., 2008). Furthermore, P2X7 variants, including SNPs and splice isoforms, are likely to play a role in the differential activity of these compounds both between and within species. For example, *N*-[(2,4-dichlorophenyl)methyl]-1-methyl-5-oxo-1-prolinamide (GSK1370319A) has an almost 7-fold higher IC₅₀ value for ATP-induced IL-1 β in human blood cultures from individuals with the A348T gain-of-function SNP than those from individuals with the E496A loss-of-function SNP (McHugh et al., 2012). Thus, given that human P2X7 is highly polymorphic (see section VIII), this has ramifications for the future use of P2X7 antagonists in treating human disorders. In particular, a need for prospective genotyping and tailor-made drug regimens

TABLE 2
Antagonist profiles of mammalian P2X7

Species	IC ₅₀									References
	PPADS	BBG	Reactive Blue	KN-62	A438079	A740003	AZ10606120	AZ11645373	JNJ-47965567	
	μM									
Human	1.2	1.9	5.4	0.2	0.9	0.09	0.003	0.05	0.005	Donnelly-Roberts et al., 2009; Michel et al., 2009; Roman et al., 2009; Bhattacharya et al., 2013
Rhesus macaque	N.D.	N.D.	N.D.	0.1	0.3	N.D.	0.004	0.02	0.003	Bradley et al., 2011b; Bhattacharya et al., 2013
Dog	N.D.	0.05	N.D.	0.01	3.9	N.D.	0.06	0.04	0.005	Michel et al., 2009; Roman et al., 2009; Bhattacharya et al., 2013
Rat	1.2	0.6	2.8	>100	0.1	0.1	0.03	>20	0.063	Donnelly-Roberts et al., 2009; Michel et al., 2009; Bhattacharya et al., 2013
Mouse (BALB/c)	4.9	0.2	21	0.2	0.6	0.7	0.6 ^a	1.6 ^a	0.03 ^a	Donnelly-Roberts et al., 2009; Michel et al., 2009; Bhattacharya et al., 2013
Mouse (C57BL/6)	40	0.5	18	0.6	0.6	1.7	0.6 ^a	1.6 ^a	0.03 ^a	Donnelly-Roberts et al., 2009; Michel et al., 2009; Bhattacharya et al., 2013
Guinea pig	0.2	0.02	N.D.	0.1	N.D.	N.D.	N.D.	1.2	N.D.	Fonfria et al., 2008; Michel et al., 2009

N.D., not determined.

^aMouse strain not disclosed.

is highlighted. Likewise, P2X7 variants also need to be considered when investigating the pharmacological blockade of P2X7 in rodents, with many studies to date overlooking the existence of such variants.

In addition to the nonspecific and specific P2X7 antagonists listed above, a number of other compounds have been shown to block P2X7 activation, many with IC₅₀ values in the micromolar range (Table 3). Naturally derived compounds such as the traditional Chinese herb emodin (Liu et al., 2010; Jelassi et al., 2013; Zhu et al., 2014), the plant-derived alkaloids berberine and sanguinine (Shemon et al., 2004), niphatoxin C, sytlissadine A and sytlissadine B from the marine sponge (Buchanan et al., 2007a,b), and the estrogen hormone 17 β -estradiol (Cario-Toumaniantz et al., 1998) can each block P2X7. The effects of other naturally derived compounds on P2X7 such as chelerythrine and colchicine are discussed below. Moreover, a number of enzyme inhibitors can block P2X7. Inhibitors of the Ca²⁺/calmodulin-dependent protein kinase (KN-62) (Gargett and Wiley, 1997), protein kinase C [chelerythrine and Ro 31-8220 (2-[1-(3-(amidinothio) propyl)-1*H*-indol-3-yl]-3-(1-methylindol-3-yl) maleimide methanesulfonate)] (Shemon et al., 2004, 2007), mitogen-activated protein kinases [SB 203580 4-(4-fluorophenyl)-2-(4-methylsulfinylphenyl)-5-(4-pyridyl)-1*H*-imidazole] and SB 202190 [4-(4-fluorophenyl)-2-(4-hydroxyphenyl)-5-(4-pyridyl)-1*H*-imidazole] (Michel et al., 2006), a vascular endothelial growth factor receptor tyrosine kinase (Compound P [*N*-[4-(1-methylethyl)phenyl]-4-[2-(3-pyridinyl)ethyl]-1-phthala-zinamine]) (Shemon et al., 2008), and phospholipase D (CAY10593 [*N*-[2-[4-(5-chloro-2,3-dihydro-2-oxo-1*H*-benzimidazol-1-yl)-1-piperidinyl]-1-methylethyl]-2-naphthalenecarboxamide], CAY10594

[*N*-[2-(4-oxo-1-phenyl-1, 3, 8-triazaspiro[4, 5]dec-8-yl)ethyl]-2-naphthalenecarboxamide], and halopemide) (Pupovac et al., 2013b) can each directly block P2X7. In this regard, studies investigating signaling pathways downstream of P2X7 activation through the use of enzyme inhibitors need to ensure that such compounds do not directly impair P2X7. A number of therapeutic drugs can also impair P2X7 activation. The antipsychotic drugs, prochlorperazine and trifluoperazine and to a lesser extent triflupromazine can each impair human, but not rodent, P2X7 channel and pore activation (Hempel et al., 2013). Colchicine, a drug used to treat gout and familial Mediterranean fever, can inhibit P2X7 pore but not channel activation (Marques-da-Silva et al., 2011). More recently, probenecid, which is used occasionally in the treatment of gout, has been found to block both the human P2X7 channel and pore (Bhaskaracharya et al., 2014). Thus, again caution needs to be made when assigning inhibitory actions of these drugs in both in vitro and in vivo studies. Furthermore, the possibility remains that the therapeutic action or unwanted side effects of such drugs may be due, in part, to P2X7 blockade.

In addition to chemical compounds, P2X7 can be blocked by various cations (Jiang, 2009). In particular, the divalent cations Ca²⁺ and Mg²⁺, at concentrations found in vivo, can inhibit P2X7 activation in vitro (Surprenant et al., 1996; Chessell et al., 1998). The ability of these divalent cations to impair P2X7 is attributed to the ability of these cations to chelate the free acid form of ATP (or ATP⁴⁻), the presumed form of ATP responsible for P2X7 activation (Cockcroft and Gomperts, 1979; Li et al., 2013). However, evidence

TABLE 3
Natural and other pharmacological compounds that impair P2X7 activation

Compound	Species	IC ₅₀ μ M	References
Natural compounds			
Emodin	Human, rat	3, 1	Liu et al., 2010; Jelassi et al., 2013; Zhu et al., 2014
Berberine	Human	N.D.	Shemon et al., 2004
Sanguinine	Human	N.D.	Shemon et al., 2004
Niphatoxin C	Human	>10	Buchanan et al., 2007b
Sytlissadine A	Human	1	Buchanan et al., 2007a
Sytlissadine B	Human	2	Buchanan et al., 2007a
17 β -Estradiol	Human	3	Cario-Toumaniantz et al., 1998
Enzyme inhibitors			
KN-62	Human, mouse	0.1, 1	Gargett and Wiley, 1997; Donnelly-Roberts et al., 2009
Chelerythrine	Human	6	Shemon et al., 2004
Ro 31-8220	Human	N.D.	Shemon et al., 2007
Compound P	Human, dog	5, N.D.	Shemon et al., 2008
SB-203580	Human	0.4–4	Michel et al., 2006
SB-202190	Human	0.6–16	Michel et al., 2006
CAY10593	Human	2	Pupovac et al., 2013b
CAY10594	Human	>10	Pupovac et al., 2013b
Halopemide	Human	>10	Pupovac et al., 2013b
Therapeutic drugs			
Colchicine	Human, mouse, rat	N.D.	Marques-da-Silva et al., 2011
Prochlorperazine	Human	1	Hempel et al., 2013
Trifluoperazine	Human	2	Hempel et al., 2013
Triflupromazine	Human	>50	Hempel et al., 2013
Probenecid	Human	200	Bhaskaracharya et al., 2014

N.D., not determined.

suggests that these and other divalent cations can also impair P2X7 by allosteric modulation (Acuna-Castillo et al., 2007; Liu et al., 2008; Moore and Mackenzie, 2008; Yan et al., 2011).

Finally, it should be noted that a number of compounds potentiate ATP-induced P2X7 activation. The human antimicrobial peptide LL-37 can potentiate both human and mouse P2X7 activation (Elssner et al., 2004). In contrast, the murine antimicrobial peptide CRAMP can enhance human but impair murine P2X7 activation (Seil et al., 2010). The antibiotic polymyxin B can potentiate human P2X7 activation (Ferrari et al., 2004), whereas the antiparasitic ivermectin can potentiate human but not murine P2X7 (Norenberg et al., 2012). A variety of lipids, including lysophosphidylcholine and sphingoglycosylphosphorylcholine, can also enhance human and murine P2X7 (Michel and Fonfria, 2007). The anesthetics ketamine, propofol, and thiopental have all been shown to increase P2X7-mediated currents (Nakanishi et al., 2007). Finally, a drug screen of 1040 approved low molecular weight drug compounds recently identified that the antihistamine clemastine can potentiate human P2X7 activation (Norenberg et al., 2011). Collectively, the expanding list of compounds that can either block or potentiate P2X7 activation highlights the promiscuous nature of this receptor and its ability to bind a large number of structurally diverse compounds.

X. Use of Modulators of P2X7 Receptor In Vivo

Pharmacological blockade of P2X7 has proved, by and large, to be efficacious in a range of rodent models of disease and injury. These studies have been comprehensively summarized in Tables 4–8. In these studies, a number of P2X7 antagonists with varying specificities have been used, at a range of concentrations, with varying injection sites and regimens. However, the efficacy of these compounds as presented in this review is potentially skewed, because studies in which P2X7 antagonists have had no effect may not have been published. Furthermore, some studies may have initially investigated the efficacy of multiple antagonists but only reported the outcomes of those that appeared beneficial. Although the majority of rodent studies report positive effects, there have been conflicting outcomes between groups for some models of human disease (e.g., status epilepticus, brain tumors, ischemia, and spinal cord injury, Tables 4 and 5). In many of the early studies listed in Tables 4–8, the antagonists used were relatively nonspecific, being either general P2X or P2 antagonists (e.g., PPADS, suramin, and oATP). Thus, the outcomes of these studies cannot definitively be associated with P2X7 but with P2 receptors in general. Furthermore, it is likely that some studies that used nonspecific P2 antagonists have been overlooked because these reports did not specifically discuss P2X7 (or the former term P2Z).

Of the first generation antagonists, oATP was the main compound used for inhibiting P2X7 in vivo, followed by BBG and PPADS. In the last 6 years, a number of more specific antagonists have been used in in vivo studies, supporting the work done in earlier studies. Of these, A438079 was most commonly used, with generally positive outcomes reported. However, despite the availability of these newer antagonists, many recent studies are still using first generation antagonists, in particular BBG and oATP. This is likely because of their relatively low cost, effectiveness in earlier studies, and, in the case of BBG, its ability to cross the blood-brain barrier (Peng et al., 2009). Further studies with additional antagonists will be important for supporting a role for P2X7 in disease. This will be particularly relevant before considering antagonizing P2X7 in human clinical trials. The following sections summarize and report the parameters and outcomes of pharmacological blockade of P2X7 in rodent models of disease and injury (Tables 4–8). The outcomes of in vivo studies using P2X7 KO mice in similar models of disease (which support antagonist studies) are also discussed.

A. P2X7 Antagonists in Rodent Models of Neurologic Disease

To the best of our knowledge, the first reported use of a P2X7 antagonist in a rodent model of neurologic disease was in 2007 (Matute et al., 2007). Since then, there have been a number of in vivo studies investigating the effects of pharmacological blockade of P2X7 in rodent models of neurologic disease (Table 4). These have largely emerged alongside evidence suggesting a role for P2X7 in central nervous system disorders (see Takenouchi et al., 2010). Overall, P2X7 antagonists reduced disease scores and symptoms in rodent models of multiple sclerosis (experimental autoimmune encephalomyelitis), status epilepticus, brain tumors (glioma), autism spectrum disorders, mania, depression, and a range of protein folding diseases (including Alzheimer's, Huntington's, Parkinson's and prion disease, and amyotrophic lateral sclerosis).

Studies using P2X7 KO mice have supported the outcomes of P2X7 antagonist studies in models of neurologic disease. First, in an Alzheimer's disease model, injection of amyloid β resulted in an accumulation of IL-1 β in WT, but not P2X7 KO mice (Sanz et al., 2009). This correlates with the positive outcomes reported in two different Alzheimer's disease models that used P2X7 antagonists (Ryu and McLarnon, 2008; Diaz-Hernandez et al., 2012). Second, in models of depression, P2X7 KO mice displayed antidepressant-like profiles in comparison with WT controls (Basso et al., 2009; Boucher et al., 2011; Csölle et al., 2013). Two different P2X7 antagonists (BBG and AZ10606120) also induced an antidepressant phenotype in WT mice (Csölle et al., 2013). However, the antagonist JNJ-47965567 had no effect on models of depression, although this antagonist was

TABLE 4
Antagonism of P2X7 in rodent models of neurologic disease

Disease/Model	Species and Strain	Sex	Start Age or Size	Injection Site	Antagonist & Dose	Injection Regimen	Outcomes	References
Experimental autoimmune encephalomyelitis	C57BL/6 mice	N.S.	N.S.	i.p. (oATP) or p.o. (BBG)	2.5 mg/kg oATP or 10 mg/kg BBG	oATP every 12 hours or BBG daily from 21 to 40 days post- MOG/IFA injection	Restored axonal conductance, reduced demyelination, and improved neurologic score	Matute et al., 2007
Experimental autoimmune encephalomyelitis	C57BL/6 mice	F	8–9 weeks	i.v.	1 mg of oATP	Once 10 days post-MOG/ CFA injection	Reduced demyelination and disease score	Lang et al., 2010
Alzheimer's disease	Sprague-Dawley rats	M	280– 300 g	i.p.	50 mg/kg BBG	With amyloid- β followed by daily injections for 7 days	Reduced P2X7 immunoreactivity, microgliosis, astrogliosis, and albumin leakage and improved neuronal viability	Ryu and McLarnon, 2008
Alzheimer's disease	J20 transgenic mice	N.S.	3–4 months	i.p.	45.5 mg/kg BBG	Every 48 hours for 4 months	Reduced hippocampal amyloid plaques and glycogen synthase kinase 3 activity	Diaz- Hernandez et al., 2012
Huntington's disease	R6/1 transgenic mice	N.S.	8 months	i.p.	45.5 mg/kg BBG or 34.2 mg/kg A438079	BBG every 48 hours and A438079 every 24 hours for 4 weeks	BBG and A438079 prevented weight loss. BBG, but not A438079, improved motor coordination and/ or prevented neuronal apoptosis	Díaz- Hernández et al., 2009
Parkinson's disease	Sprague-Dawley rats	M	8–12 weeks	i.p.	30 mg/kg A438079	1 hour before and 1 hour after 6-OHDA injection	Partially prevented striatal dopaminergic depletion but not associated with a reduction of dopaminergic cell loss	Marcellino et al., 2010
Prion disease	C57BL/6 mice	N.S.	N.S.	i.p.	100 mg/kg BBG	3 \times /week for 3 weeks, then 2 \times /week for 4 weeks at 100 dpi (Fukuoka- 1 strain)	Prevented cerebral protease-resistant prion protein formation but did not prevent disease progression	Iwamaru et al., 2012
Amyotrophic lateral sclerosis	B6SJL-TgN SOD1/G93A1Gur mice	M and F	90 days	i.p.	45.5 mg/kg BBG	Every 48 hours	Improved motor function (rotarod and grip-strength) in both sexes but had no effect on survival, delayed body weight loss in males	Cervetto et al., 2013b
Pilocarpine-induced seizures	C57BL/6J mice	M	8–13 weeks	i.c.v.	5 mM oATP, 10 μ M A438079 or 10 μ M A740003	Infused over 1 week before PILO injection (rate N.S.)	Increased seizure susceptibility and induced epileptiform discharge	Kim and Kang, 2011
Pilocarpine-induced status epilepticus	Sprague-Dawley rats	M	9–11 weeks	i.c.v.	5 mM oATP	0.5 μ l/h for 1 week beginning 3 days before PILO injection	Reduced neutrophilic and monocytic infiltrates in frontoparietal cortex but not in piriform cortex	Kim et al., 2010
Pilocarpine-induced status epilepticus	Sprague-Dawley rats	M	9–11 weeks	i.c.v.	5 mM oATP, 10 μ M A438079 or 10 μ M A740003	0.5 μ l/h for 2 weeks beginning 3 days before PILO injection	Reduced TNF- α in dentate granule cells and increased neuronal death	Kim et al., 2011

(continued)

TABLE 4—Continued

Disease/Model	Species and Strain	Sex	Start Age or Size	Injection Site	Antagonist & Dose	Injection Regimen	Outcomes	References
Kainic acid-induced status epilepticus	C57BL/6 mice	M	20–25 g	i.c.v.	1.75 nmol of A438079, 1 pmol of BBG, or 1.75 and 3.5 nmol of PPADS	5 min before or 15 min or 1 hour after kainic acid injection	A438079 (pre- or post-treatment) and BBG (pretreatment), but not PPADS, reduced seizure time, seizure-induced neuronal death, and reactive microglia	Engel et al., 2012
Kainic acid-induced status epilepticus	C57BL/6 mice	N.S.	Adult	i.c.v.	1.75 nmol of A438079 or 1 pmol of BBG	2 μ l 10 min before and 60 min after kainic acid injection	A438079 reduced electrographic and clinical seizure severity, A438079 and BBG reduced seizure-induced neuronal death	Jimenez-Pacheco et al., 2013
Glioma	Sprague-Dawley rats	M	250–300 g	i.p.	50 mg/kg BBG	Immediately after implantation of C6 cells, then 2 \times /day	Inhibited tumor growth	Ryu et al., 2011
Glioma	Sprague-Dawley rats	F	200–300 g	i.c.	10 μ g of oATP	Coinjected with C6 cells	Reduced activated microglia, macrophages, and MIP-1 α or MCP-1-expressing cells at tumor site and reduced tumor size 7 dpi	Fang et al., 2011
Autism spectrum disorders	C57BL/6J mice	M and F	6 weeks	i.p.	10 or 20 mg/kg suramin	Weekly	Improved social behavior and sensorimotor coordination deficits, prevented cerebellar Purkinje cell loss, and restored normal body temperature, synaptosomal structure, and various metabolic/signaling parameters	Naviaux et al., 2013
Amphetamine-induced hyperactivity (mania)	Sprague-Dawley rats	M	Adult	s.c.	30 mg/kg JNJ-47965567	1 hour before d-amphetamine sulfate injection	Reduced hyperactivity but not basal locomotor activity	Bhattacharya et al., 2013
Depression	Sprague-Dawley rats	M	N.S.	s.c.	3, 10, or 30 mg/kg JNJ-47965567	24 hours and 30 min before forced swim test	No effect on immobility, climbing, or swimming behavior	Bhattacharya et al., 2013
Depression	C57BL/6 mice	M	2–3 months	i.p.	2 mg/kg AZ10606120 or 50 mg/kg BBG	1 \times /day for 4 days or once before testing/treatment	Resulted in an antidepressant phenotype	Csölle et al., 2013

6-OHDA, 6-hydroxydopamine; BBG, Brilliant Blue G; CAMKII, calcium-calmodulin kinase II; CFA, complete Freund's adjuvant; dpi, days postinfection/injection; EGFR, epidermal growth factor receptor; ERK1/2, extracellular response kinase 1 and 2; HIF-1 α , hypoxia induce factor-1; i.c., intracerebral; IFA, incomplete Freund's adjuvant; MCP-1, monocyte chemoattractant protein-1; MIP-1 α , macrophage inflammatory protein-1 α ; MOG, myelin oligodendrocyte glycoprotein_{35–55}; N.S., not stated; oATP, periodate-oxidized ATP; PILO, pilocarpine; PPADS, pyridoxal phosphate-6-azophenyl-2,4-disulfonic acid; TNF, tumor necrosis factor; VEGF, vascular endothelial growth factor.

efficacious and slightly effective, respectively, in models of mania and neuropathic pain (Bhattacharya et al., 2013). To the best of our knowledge, this is the only documented use of JNJ-47965567, and it is possible that the injection regimen used was not sufficient to see an efficacious response in this disease model.

The role for P2X7 in neurologic disease in rodents, however, is complicated by other varying results with

P2X7 antagonists or P2X7 KO mice. For example, in a single study of experimental autoimmune encephalomyelitis (Matute et al., 2007), two different P2X7 antagonists were efficacious. In support of this finding, the incidence of experimental autoimmune encephalomyelitis was reduced 4-fold in GSK P2X7 KO mice compared with WT mice (Sharp et al., 2008). However, Pfizer P2X7 KO mice developed more severe experimental

TABLE 5
Antagonism of P2X7 in rodent models of neurologic injury

Disease/Model	Species and Strain	Sex	Start Age or Size	Injection Site	Antagonist and Dose	Injection Regimen	Outcomes	References
Focal cerebral ischemia and excitotoxic brain injury	C57/SV129 mice	M	25–30 g	i.c.v.	3 mM oATP, 30 μ M or 10 mM PPADS, or 2 mM KN-62	1 μ l before and 30 min after MCAO and 0.5 μ l before methanoglutamate infusion	Antagonists did not affect ischemic or excitotoxic cell death	Le Feuvre et al., 2003
Focal cerebral ischemia	Wistar rats	M	270–290 g	i.p.	10 and 100 mg/kg Reactive Blue 2	5 min after MCAO	Improved neurologic score and reduced infarct volume	Melani et al., 2006
Focal cerebral ischemia	Wistar rats	M	260–300 g	i.c.v.	100 mM oATP	2 μ l 60 min after onset of MCAO	Exacerbated behavioral dysfunction and ischemic neural injury	Yanagisawa et al., 2008
Focal cerebral ischemia	Wistar rats	M	270–300 g	i.p.	30 mg/kg BBG	Every 12 hours for 3 days beginning 30 min after MCAO	Reduced histologic brain damage and neurologic score	Arbeloa et al., 2012
Focal cerebral ischemia	C57BL/6 and Balb/c mice	M	20–28 g	i.v.	35 mg/kg BBG	Daily for 3 days after MCAO	Increased endogenous ciliary neurotrophic factor but no effect on histologic brain damage	Kang et al., 2013
Global cerebral ischemia/reperfusion	Sprague-Dawley rats	M	260–320 g	i.c.v.	10 μ g of BBG, 1 μ g of oATP, or 3 μ g of A438079	2 μ l at 0.5 μ l/min 10 min before four-vessel occlusion	Reduced neuron death, neuronal DNA cleavage, glial activation, hippocampal cytokine overexpression, and behavioral deficits and improved survival rate	Chu et al., 2012
Anoxia-induced brain injury	Sprague-Dawley rats	N.S.	2–3 weeks	i.p.	10 mM oATP	100 μ l 1 hour before anoxia-induced brain ischemia and reperfusion injury	Inhibited whole-brain aquaporin 4 downregulation	Lee et al., 2008b
Subarachnoid hemorrhage	Sprague-Dawley rats	M	280–320 g	i.p.	30 mg/kg BBG	0.5, 24, and 48 hours after subarachnoid hemorrhage	Improved neurobehavioral functions, reduced brain edema, subcortical neuronal apoptosis, p38 MAPK activation and cleaved caspase-3, and increased Bcl-2	Chen et al., 2013a
Subarachnoid hemorrhage	Sprague-Dawley rats	M	280–320 g	i.p.	5, 30, or 100 mg/kg BBG	30 min after subarachnoid hemorrhage	30, but not 5 or 100, mg/kg improved neurologic score and brain edema, and decreased caspase-1, IL-1 β , IL-18 and neutrophil infiltration	Chen et al., 2013b
Brain trauma	Sprague-Dawley rats	M	Adult	Brain (cortex)	2 mm ³ foam soaked in 1 mM suramin	After corticectomy into surgery site	Reduced MCP-1 expression	Panenka et al., 2001
Mechanical injury-induced astrogliosis	Wistar rats	M	280–320 g	Brain (nucleus accumbens)	0.03 nmol PPADS	1 μ l at 12 μ l/h 15 min before or with agonists	Reduced injury- and agonist-induced astrocyte activation	Frankie et al., 2003
Mechanical injury-induced microglial activation	Wistar rats	M	280–320 g	Brain (nucleus accumbens)	30 pmol of PPADS or 1 pmol of BBG	1 μ l at a rate of 12 μ l/h before P2X agonists	Reduced injury-induced P2X receptor expression and inhibited agonist-induced microglial activation	Frankie et al., 2007
Spinal cord injury	Sprague-Dawley rats	F	220–250 g	i.s.	2 nM oATP and PPADS	2 μ l immediately before or 1 h after weight-drop	Improved functional recovery and reduced cell death	Wang et al., 2004

(continued)

TABLE 5—Continued

Disease/Model	Species and Strain	Sex	Start Age or Size	Injection Site	Antagonist and Dose	Injection Regimen	Outcomes	References
Spinal cord injury	Sprague-Dawley rats	N.S.	Adult	i.v.	10 or 50 mg/kg BBG	10–15 min after weight-drop and once daily for the next 2 days	Improved hindlimb motor function recovery and reduced spinal cord tissue loss	Peng et al., 2009
Spinal cord injury	Fischer rats	F	220–250 g	i.s.	10 mM PPADS or 0.2 mM BBG	30 min after weight-drop	Did not improve locomotion or secondary histopathological changes after 7 weeks	Marcello et al., 2012
Spinal cord injury	Sprague-Dawley rats	F	220–250 g	i.v.	10 or 50 mg/kg BBG	15 min after weight-drop and then at 24 and 72 hours	Did not improve motor recovery or histopathological outcome after 6 weeks	Marcello et al., 2012
Nerve fusion	Sprague-Dawley rats	F	N.S.	Nerve bathed directly	100 μ M Brilliant Blue	After sciatic nerve transection and before nerve rejoining	Decreased compound action potential amplitudes and improved behavioral recovery	Rodriguez-Feo et al., 2013
Neuropathic pain	Rats	N.S.	N.S.	i.p.	10–300 μ mol/kg A438079	7–14 days after spinal nerve ligation	Dose-dependently reversed mechanical allodynia	Nelson et al., 2006
Neuropathic pain	Sprague-Dawley rats	M	200–300 g	i.p. or p.o.	4–105 (i.p.) or 142 mg/kg (p.o.) A740003	2 weeks after nerve ligation/constriction or vincristine infusion	Dose-dependently attenuated mechanical allodynia in all models	Honore et al., 2006
Neuropathic pain	Sprague-Dawley rats	M	250–400 g	i.p. or i.v.	10–300 μ mol/kg A438079	2 weeks after nerve ligation/constriction or vincristine infusion	Dose-dependently attenuated mechanical allodynia in all models (i.p.) and reduced activity of spinal neurons after spinal nerve ligation (i.v.)	McGarraughy et al., 2007
Neuropathic pain	C57BL/6 mice	N.S.	8 weeks	i.p.l.	4 mg/kg oATP	Daily, starting day of sciatic nerve constriction	Reduced hyperalgesia	Fulgenzi et al., 2008
Neuropathic pain	Sprague-Dawley rats	M	200–250 g	s.c.	30 mg/kg AACBA	22–24 days after spinal nerve ligation	Did not alter mechanical hypersensitivity	Broom et al., 2008
Neuropathic pain	Rats	N.S.	N.S.	i.p. or p.o.	10–100 (i.p.) or 100 μ mol/kg (p.o.) Compound 53	After spinal nerve ligation, once (i.p.) or twice daily for 4 days (p.o.) before testing	Reduced tactile allodynia	Perez-Medrano et al., 2009
Neuropathic pain	Wistar rats	M	250–350 g	i.p.	6.25–50 mg/kg PPADS or 1–100 mg/kg BBG	7 days after partial ligation of sciatic nerve	Slightly attenuated mechanical allodynia at lower doses	Andó et al., 2010
Neuropathic pain	Sprague-Dawley rats	N.S.	180–220 g	i.t.	10 μ M BBG	10 μ l daily for 14 days starting right after chronic constriction of sciatic nerve	Partially attenuated mechanical allodynia and thermal hypersensitivity	He et al., 2012
Neuropathic pain	Sprague-Dawley rats	M	180–350 g	i.t.	300 μ M oATP or 20 μ M BBG	5 μ l/min for 4 min 0.5 hours before tetanic stimulation of sciatic nerve	Prevented spinal long-term potentiation and reduced mechanical allodynia; BBG also inhibited the upregulation of microglial P2X7 and expression of Iba-1, phosphorylated p38, IL-1 β and GluR1	Chu et al., 2010
Neuropathic pain	Sprague-Dawley rats	N.S.	N.S.	s.c.	3, 30, or 100 mg/kg JNJ-47965567	Once after spinal nerve ligation	reduced mechanical allodynia after 30–60 min	Bhattacharya et al., 2013
Orofacial pain	Sprague-Dawley rats	M	200–250 g	i.t.	35 μ g of A438079	Immediately after CCI of the infraorbital nerve	Inhibited tactile allodynia/hyperalgesia and prevented TNF- α upregulation in trigeminal sensory nuclear complex	Ito et al., 2013

(continued)

TABLE 5—Continued

Disease/Model	Species and Strain	Sex	Start Age or Size	Injection Site	Antagonist and Dose	Injection Regimen	Outcomes	References
Acute thermal nociception	Sprague-Dawley rats	M	200–300 g	i.p.	142 mg/kg A740003	30 min before hot-plate test	Did not affect paw withdrawal latencies in response to heat	Honore et al., 2006
Acute thermal nociception	Wistar rats	M	170–300 g	i.p.	6.25–50 mg/kg PPADS or 3–100 mg/kg BBG	30 min before hot-plate test	25 and 50 mg/kg PPADS—elevated heat threshold; 10 mg/kg BBG reduced heat threshold	Andó et al., 2010
Morphine tolerance	Sprague-Dawley rats	M	180–350 g	i.t.	25 mg/kg BBG	2×/day for 5 days, 30 min before each morphine dose	Reduced tolerance to morphine analgesia	Chen et al., 2012
BzATP-induced IL-1 β release	Sprague-Dawley rats	M	280–320 g	s.c.	30 or 100 mg/kg JNJ-47965567	30 min before Bz-ATP	100, but not 30, mg/kg attenuated IL-1 β release from hippocampus	Bhattacharya et al., 2013

BzATP; 2'-(3'-O-(4-benzoylbenzoyl)ATP; CCI, chronic constriction injury; GFAP, glial fibrillary acidic protein; Iba-1, ionized calcium-binding adapter molecule 1; IL, interleukin; i.p.i., intraplantar; i.s., intraspinal; MAPK, mitogen-activated protein kinase; MCAO, middle cerebral artery occlusion; MCP-1, monocyte chemoattractant protein-1; N.S., not stated; PKC, protein kinase C; TNF, tumor necrosis factor.

autoimmune encephalomyelitis than WT mice (Chen and Brosnan, 2006). In another example, Pfizer P2X7 KO mice were more susceptible to seizures than WT littermates in a model of status epilepticus (Kim and Kang, 2011). This finding was supported by the use of three P2X7 antagonists (oATP, A438079, and A740003), each of which enhanced disease symptoms in status epilepticus (Kim and Kang, 2011; Kim et al., 2011). In contrast, opposite outcomes were reported using two antagonists (BBG and A438079) in this disorder (Engel et al., 2012; Jimenez-Pacheco et al., 2013).

B. P2X7 Antagonists in Rodent Models of Neurologic Injury

In addition to neurologic diseases, there have been a number of studies investigating the effects of pharmacological blockade of P2X7 in rodent models of neurologic injuries (Table 5) induced by both focal and global ischemia/reperfusion, excitotoxicity, subarachnoid hemorrhage, and mechanical trauma (e.g., corticectomy, stab wound, and weight drop models). In addition, there have been a number of pain models investigated, including nerve ligation- and chemical-induced models of neuropathic pain and acute pain induced by heat. Lastly, P2X7 blockade in a model of morphine tolerance has been investigated. P2X7 antagonists were largely efficacious in these models, reducing injury scores, symptoms, and pain. In these studies, pain is often characterized by an increased sensitivity to painful stimuli (hyperalgesia) or a pain reaction in response to normally innocuous stimuli (allodynia). Notable differences were observed for models of ischemic and excitotoxic brain injury and mechanical-induced spinal cord injury. In these studies, P2X7 antagonists were reported to either exacerbate or reduce symptoms, despite similar protocols. Furthermore, these conflicting outcomes were observed despite the use of multiple antagonists in some studies (including Le Feuvre et al., 2003; Wang et al., 2004; Chu et al., 2012; Marcillo et al., 2012). Thus, notwithstanding given differences between laboratories, the cause of these opposing observations remains unknown.

The efficacy of P2X7 antagonists in models of ischemic brain injury and neuropathic pain are supported by similar studies using P2X7 KO mice. In one study, neurologic deficits, neuronal death, and the overproduction of Fas ligand (CD95L), Fas-associated protein with death domain and caspase-8 were reduced in P2X7 KO mice compared with WT littermates after microsphere embolism (Lu et al., 2012). Likewise, P2X7 antagonists were efficacious in models of ischemia in three independent studies (Melani et al., 2006; Arbeloa et al., 2012; Chu et al., 2012). These findings contrast with another study in which P2X7 KO mice had similar amounts of cell death induced by temporary cerebral ischemia compared with WT mice (Le Feuvre et al., 2003). In this same study, P2X7 antagonists (oATP, PPADS, and KN-62) similarly had no effect on ischemic cell death. However an IL-1 receptor antagonist reduced

TABLE 6
Antagonism of P2X7 in rodent models of compound-induced inflammation

Disease/Model	Species and Strain	Sex	Start Age or Size	Injection Site	Antagonist and Dose	Injection Regimen	Outcomes	References
Formalin-induced pain	Sprague-Dawley rats	M	100–200 g	s.c., hindpaw	50–500 nmol of suramin or 5–500 nmol of PPADS	Coadministered with formalin or formalin/ATP	Dose-dependently inhibited amplification of the formalin response (flinching) by ATP but not to formalin alone	Sawynok and Reid, 1997
Formalin-induced pain	Sprague-Dawley rats	M	250–400 g	i.p.	30–300 μ mol/kg A438079	After injecting formalin into hind paw	Dose-dependently attenuated nocifensive behaviors (paw flinching, licking, guarding)	McGaraughty et al., 2007
Formalin-induced pain	BALB/cJ mice	N.S.	6–8 weeks	s.c.	300 μ mol/kg A438079	30 min before formalin injection	Attenuated nocifensive behaviors (licking, biting)	Hansen et al., 2011
Acetic acid-induced visceral pain	C57BL/6 mice	N.S.	8 weeks	i.p.	4 mg/kg oATP	30 min before inducing visceral pain	Reduced writhing responses	Fulgenzi et al., 2008
LPS-induced inflammation	Wistar rats	M	280–320 g	i.p.	5, 25, or 100 mg/kg suramin; 5 or 25 mg/kg PPADS; or 40 or 100 mg/kg BBG	5 min before i.p. LPS injection	Attenuated fever (all), increased plasma IL-1 β and IL-6 (PPADS, BBG), and increased plasma TNF- α (BBG)	Gourine et al., 2005
LPS-induced inflammation	Wistar rats	M	200–300 g	s.c.	1–30 mg/kg AACBA	1 hour before i.p. LPS injection	Dose-dependently reduced plasma IL-6 release	Broom et al., 2008
LPS-induced inflammation	Wistar rats	M	200–300 g	i.p.	25 mg/kg PPADS, 100 mg/kg BBG, or 0.9 mg/kg oATP	30 min (PPADS and BBG) or 2 hours (oATP) before i.p. LPS injection	Reduced hippocampal IL-1 β release; BBG also reduced basal release	Csölle and Sperlágh, 2010
LPS-induced inflammation	C57Bl/6J mice	M	2–3 months	i.p.	100 mg/kg BBG, 0.9 mg/kg oATP	30 min (BBG) or 2 hours (oATP) before i.p. LPS injection	Reduced hippocampal IL-1 β release; also reduced basal release	Csölle and Sperlágh, 2010
LPS-induced inflammation	C57BL/6 mice	N.S.	N.S.	i.p.	100 μ mol/kg A438079	1 hour before i.p. LPS injection	Impaired body temperature increase	Barberà-Cremades et al., 2012
LPS-induced CNS inflammation	Sprague-Dawley rats	M	240–260 g	i.c.v.	1 μ M oATP	30 min before striatal LPS injection	Reduced activated microglia, phosphorylation of p38 MAPK, iNOS production, nitration of proteins, lipid peroxidation, oxidative DNA damage, and caspase-3 activation and increased neuronal survival	Choi et al., 2007
LPS-induced CNS inflammation	Wistar rats	M	N.S.	i.t.	50 μ g of oATP or 50 μ g of A438079	With or without LPS 24 h after 1st i.t. LPS injection	Attenuated mechanical hyperalgesia and prevented phosphorylation of p38 MAPK in microglia	Clark et al., 2010
Carrageenan-induced inflammation	Wistar rats	M	~250 g	i.p.l., p.o., or i.v.	50–200 μ M oATP	3 hours after i.p.l. carrageenan injection	Reduced thermal hyperalgesia (all routes), expression of chemokines and P2X7, and macrophage infiltrates (i.p.l, i.v.)	Fulgenzi et al., 2005

(continued)

TABLE 6—Continued

Disease/Model	Species and Strain	Sex	Start Age or Size	Injection Site	Antagonist and Dose	Injection Regimen	Outcomes	References
Carrageenan-induced inflammation	Sprague-Dawley rats	M	200–300 g	i.p.	14–142 mg/kg A740003	90 min after i.p.l. carrageenan injection	Dose-dependently reduced thermal hyperalgesia and paw edema	Honore et al., 2006
Carrageenan-induced inflammation	Sprague-Dawley rats	M	200–250 g	s.c.	3–100 mg/kg AACBA	30 min before (prophylactic) or 1–1.5 hours after (therapeutic) i.p.l. carrageenan injection	Prophylactically and therapeutically reduced mechanical hypersensitivity, and therapeutically reduced paw swelling	Broom et al., 2008
Carrageenan-induced inflammation	Albino Wistar rats	M	200–250 g	TMJ	100, 300, and 900 μ g of A438079	Coinjected with carrageenan	Did not alter TMJ hyperalgesia	Teixeira et al., 2010a
Carrageenan-induced inflammation	Wistar rats	M	200–250 g	s.c., hindpaw	0.5, 2, or 6 μ g/paw oATP or 100 or 300 μ g/paw A438079	Coinjected with carrageenan	Reduced hyperalgesia and the concentration of TNF- α , IL-6, and CINC, but not IL-1 β , in hindpaw	Teixeira et al., 2010b
MO-induced inflammation	Sprague-Dawley rats	M	275–420 g	i.t.	1 μ M BBG or 100 μ M oATP	Continuous superfusion over MDH for 2 hours	Attenuated MDH central sensitization	Itoh et al., 2011
CO-induced irritant contact dermatitis	Swiss mice	M	6–8 weeks	i.p.	80 μ mol/kg A438079	2 hours after CO application	Reduced neutrophil infiltration, myeloperoxidase levels, IL-1 β , Gr1 ⁺ cell recruitment, dendritic cells, and macrophages and decreased ear edema	Da Silva et al., 2013
Venom-induced pain and inflammation	Sprague-Dawley rats	M	250–300 g	i.t.	0.1, 1, or 10 μ g of A438079	10 min before venom injection	Dose-dependently inhibited spontaneous flinching and mechanical allodynia but not edema	Zhou et al., 2012

CINC, cytokine-induced neutrophil chemoattractant 1; CNS, central nervous system; CO, croton oil; IL, interleukin; iNOS, inducible nitric-oxide synthase; i.p.l., intraplantar; LPS, lipopolysaccharide; MAPK, mitogen-activated protein kinase; MDH, medullary dorsal horn; MO, mustard oil; N.S., not stated; TMJ, temporomandibular joint; TNF, tumor necrosis factor.

cell death, suggesting IL-1 α and IL-1 β can be secreted via a P2X7-independent mechanism during cerebral ischemia in mice. In models of neuropathic pain, P2X7 KO mice lacked hypersensitivity to thermal and mechanical stimuli compared with WT mice (Chessell et al., 2005; Fulgenzi et al., 2008; Hansen et al., 2011). These studies support those that used P2X7 antagonists, in which a range of antagonists attenuated neuropathic pain (Table 5).

C. P2X7 Antagonists in Rodent Models of Compound-Induced Inflammation

Most laboratories studying the effects of P2X7 antagonists on inflammation *in vivo* have done so by relatively simple rodent models involving the injection of compounds such as formalin, acetic acid, lipopolysaccharide (LPS), carrageenan, mustard oil, croton oil, or venom (Table 6). In these models, inflammation has largely been assessed at either a cellular level or by using measures of pain, fever, or edema, three of the classic hallmarks of inflammation. In the majority of studies, administration of P2X7 antagonists reduced inflammatory processes. Of

particular note was the separate use of multiple antagonists in some studies. These studies reported an antagonist-dependent reduction in LPS- (Gourine et al., 2005; Clark et al., 2010; Csölle and Sperlágh, 2010), carrageenan- (Teixeira et al., 2010b), and mustard oil-induced inflammation (Itoh et al., 2011) and further support the benefits of targeting P2X7 in inflammation.

In addition to the use of multiple antagonists, the use of P2X7 KO mice, which demonstrate attenuated compound-induced inflammatory responses compared with WT counterparts, also supports the benefits of targeting P2X7 in inflammation. The majority of P2X7 KO studies investigating the role of P2X7 on inflammation *in vivo* have done so by injecting LPS. In these studies, LPS induced mechanical hypersensitivity, fever, hyporeactivity in response to phenylephrine, increased p38 and nuclear factor- κ B phosphorylation, increased cytokines [IL-1 β , IL-10, and tumor necrosis factor- α (TNF- α)], and decreased mean arterial pressure and pressor responses to noradrenaline in WT mice, all of which were attenuated or absent in P2X7 KO mice (Mingam et al., 2008; Clark et al., 2010; Barberà-Cremades et al., 2012; Chiao et al.,

2013; Kim et al., 2013). These findings contrast with the results of one study, which found that LPS induced similar levels of IL-1 β and TNF- α in the serum of WT and P2X7 KO mice (He et al., 2013). In this study, these cytokines were attenuated in NLRP3 KO mice, suggesting that these cytokines can be secreted via a P2X7-independent mechanism during LPS-induced inflammation in mice or that compensatory mechanisms operate in P2X7 KO mice. The role of P2X7 in visceral pain has also been investigated in P2X7 KO mice. In one study, acetic acid-induced visceral pain, as measured by writhing behavior, was largely absent in P2X7 KO mice compared with WT littermates (Fulgenzi et al., 2008). This same study also observed greatly reduced acetic acid-induced visceral pain in WT mice after pretreatment with the P2X7 antagonist oATP.

D. P2X7 Antagonists in Rodent Models of Musculoskeletal Disorders

The efficacy of P2X7 antagonists has also been examined in rodent models of musculoskeletal disorders such as arthritis, bone cancer, and Duchenne muscular dystrophy (Table 7). The majority of these studies examined inflammatory arthritis, for which there are two main types of rodent models employed. In the first model, intraplantar injection of Complete Freund's adjuvant (CFA) is used to induce inflammatory pain and swelling of the paws. This model is also used to investigate compound-induced inflammation, but is included here because of its relevance to musculoskeletal disorders. In the second model, collagen-induced or anti-collagen antibody-induced arthritis results in an autoimmune response against collagen and subsequent joint damage. These are generally used as models of rheumatoid arthritis, resulting in pathology similar to that of the human disease (Williams, 2012). A range of P2X7 antagonists has resulted in improved experimental outcomes in models of inflammatory arthritis (Table 7). In addition, GSK P2X7 KO mice do not develop CFA-induced inflammatory hypersensitivity (Chessell et al., 2005), whereas Pfizer P2X7 KO mice have reduced incidence and severity of anticollagen antibody-induced arthritis compared with WT littermates (Labasi et al., 2002; Fulgenzi et al., 2008). As a result of these and other studies, the first approved drugs that target P2X7 have undergone clinical trials in rheumatoid arthritis patients (see section XI).

E. P2X7 Antagonists in Rodent Models of Disorders Associated with Other Tissues

Although the majority of in vivo P2X7 antagonist studies have focused on neurologic and musculoskeletal inflammation and inflammatory disorders, there are a number of studies that have looked at the effects of pharmacological blockade of P2X7 in disorders associated with other tissues (Table 8). P2X7 antagonists reduced symptoms in rodent models of disorders affecting the bladder (hemorrhagic cystitis), colon (carcinoma), eyes

(retinitis pigmentosa, photoreceptor degeneration, retinal ganglion cell death, retinal pigmented epithelium degeneration, subretinal hemorrhage, and optic nerve injury), heart (transplantation, cardiac transplant rejection, and myocardial ischemia), kidney (glomerulonephritis, lupus nephritis, hypertension, and renal injury), liver (obesity and hepatotoxicity), lung (chemical and allergic inflammation or chronic obstructive pulmonary disease), pancreas (autoimmune diabetes, islet allograft rejection, and pancreatitis), and skin (dermatitis, graft-versus-host disease, and melanoma). Altogether these studies used a range of antagonists, demonstrating the widespread effects of P2X7 in a number of tissues and expanding the number of disorders that may benefit from P2X7 blockade. Furthermore, a number of these studies also had supporting P2X7 KO mouse data alongside P2X7 antagonist data, further implicating P2X7 in a number of disease processes. These include models of hemorrhagic cystitis (Martins et al., 2012), glomerulonephritis (Taylor et al., 2009b), hypertension and renal injury (Ji et al., 2012), lung inflammation (Lucattelli et al., 2011), allergic airway inflammation (Müller et al., 2011), pancreatitis (Hoque et al., 2011), and graft-versus-host disease (Wilhelm et al., 2010).

XI. P2X7 Antagonists in Human Clinical Trials

Because of the potential benefits of P2X7 blockade in rodent studies, the use of drugs targeting P2X7 in humans has commenced for a number of disorders, including arthritis, pain, and multiple sclerosis (Baas, 2012). Of these trials, only two have been published as full-length reports, with both reporting the clinical efficacy and safety of the P2X7 antagonists AZD9056 or CE-224,535 in rheumatoid arthritis patients (Keystone et al., 2012; Stock et al., 2012). However, both of the trials failed to find any therapeutic benefit with these compounds (Keystone et al., 2012; Stock et al., 2012). The first of these trials were two phase II studies in patients with rheumatoid arthritis treated with the AstraZeneca compound, AZD9056 (Keystone et al., 2012). Despite a significant reduction in swollen and tender joint count in the phase IIa study, there was no significant benefit on disease in the phase IIb study. With the exception of one patient, who required hospitalization for nausea and vomiting after treatment with AZD9056, there were no significant side effects. The second clinical trial reported was a phase IIa study in patients with rheumatoid arthritis treated with the Pfizer compound, CE-224,535 (Stock et al., 2012). There was no significant benefit on disease with this compound. Furthermore, treatment with CE-224,535 resulted in a small but acceptable increase in the rate of nausea and diarrhea. In both clinical trials there were no serious infections or deaths related to adverse events (Keystone et al., 2012; Stock et al., 2012).

TABLE 7
Antagonism of P2X7 in rodent models of musculoskeletal disorders

Disease/Model	Species and Strain	Sex	Start Age or Size	Injection Site	Antagonist and Dose	Injection Regimen	Outcomes	References
CFA-induced inflammation	Wistar rats	M	~250 g	i.p.l.	56–336 μ M oATP	24 hours after i.p.l. CFA injection	Dose-dependently increased paw pressure threshold (reduced hyperalgesia) in inflamed paws	Dell'Antonio et al., 2002b
CFA-induced inflammation	Wistar rats	M	~250 g	i.p.l. or i.v.	168 μ M oATP	24 hours after i.p.l. CFA or 3 hours after i.v. CFA/fucoidin injection	Increased paw pressure threshold (reduced hyperalgesia) and reduced edema and ATP levels in inflamed paws	Dell'Antonio et al., 2002a
CFA-induced inflammation	Sprague-Dawley rats	M	200–300 g	i.p.	14–142 mg/kg A740003	47.5 hours after i.p.l. CFA injection	Dose-dependently reduced thermal hyperalgesia but not paw edema	Honore et al., 2006
CFA-induced inflammation	Sprague-Dawley rats	M	200–300 g	i.p.	30–300 μ mol/kg A839977	48 hours after i.p.l. CFA injection	Dose-dependently reduced thermal hyperalgesia	Honore et al., 2009
CFA-induced inflammation	BALB/c and IL-1 α β KO mice	M	20–25 g	s.c.	10–100 μ mol/kg A839977	48 hours after i.p.l. CFA injection	Dose-dependently reduced thermal hyperalgesia in WT but not IL-1 α β KO mice	Honore et al., 2009
CFA-induced inflammation	Wistar rats	M	200–250 g	i.p.	3–100 mg/kg BBG	48 hours after i.p.l. CFA injection	Dose-dependently reduced hyperalgesia but had no effect on edema	Andó et al., 2010
CFA/MO-induced chronic arthritis	B6129SF2/J WT and TNFR1/R2 KO mice	N.S.	6–8 weeks (20–25 g)	i.t.	50 μ g of A438079	20 weeks after knee joint cavity CFA injection	Reduced thermal hyperalgesia and mechanical allodynia for >3 hours in TNFR1/R2 KO mice	Westlund et al., 2012
Anticollagen antibody-induced arthritis	C57BL/6 mice	M	8 weeks	i.p.l., p.o., or i.p.	4 mg/kg oATP	10 days after inducing arthritis	Increased paw withdrawal latency (i.e., reduced hyperalgesia) (all routes) up to 24 hours post-treatment	Fulgenzi et al., 2008
Collagen-induced arthritis	Lewis rats	F	125–175 g	s.c.	30 mg/kg AACBA	Before collagen injection and once a day after for 21 days (prophylactic) or once a day from 13–21 days after injection (therapeutic)	Prophylactically, but not therapeutically, reduced clinical and histologic scores	Broom et al., 2008
Collagen-induced arthritis	DBA/1J mice	M	8 weeks	i.p.	3 mM oATP	100 μ l daily for 10 days starting at onset of visible clinical signs	Reduced clinical score, cartilage erosion, infiltrating T cells, and serum type II collagen IgG autoantibodies	Ardissone et al., 2011
Bone cancer pain	BALB/cJ and C3H/HeN mice	N.S.	6–8 weeks	s.c.	300 μ mol/kg A438079	14 days post bone cancer operation	Did not improve limb use score or weight-bearing ratio after acute treatment in models with (BALB/cJ) and without (C3H/HeN) astrocyte activation	Hansen et al., 2011
Duchenne muscular dystrophy	<i>mdx</i> transgenic mice	M	3 weeks	i.p.	125 mg/kg Coomassie Brilliant Blue	Every 3 days from 3–14 weeks of age	Reduced number of degeneration-regeneration cycles in skeletal muscles	Young et al., 2012
Duchenne muscular dystrophy	<i>mdx</i> transgenic mice	M	9 weeks	i.p.	45.5 mg/kg BBG	Every 48 hours for 3 weeks	Blocked CD62L shedding from blood T cells and restored their capability to migrate to the heart	Cascahalho et al., 2012

IL, interleukin; i.p.l., intraplantar; i.t., intrathecal; MO, mustard oil; N.S., not stated; p.o., per os; TNFR, tumor necrosis factor receptor.

TABLE 8
Antagonism of P2X7 in rodent models of disorders associated with other tissues

Disease/Model	Species and Strain	Sex	Start Age or Size	Injection Site	Antagonist and Dose	Injection Regimen	Outcomes	References
Bladder Cyclophosphamide-induced hemorrhagic cystitis	Swiss and C57/BL6 mice	M	25–30 g	i.p.	50, 100, and 200 $\mu\text{mol/kg}$ A438079	30 min before and 4 hours after cyclophosphamide injection	Reduced nociceptive behavior scores, c-Fos staining in the spinal cord and brain, edema and hemorrhage indices, tissue IL-1 β , and TNF- α levels and macrophage migration	Martins et al., 2012
Colon Colon carcinoma	BALB/c mice	F	5–7 weeks, 15–18 g	Tumor	600 μM oATP	200 μl 5, and 13 days after P2X7-overexpressing CT26 cell injection	Reduced tumor growth, angiogenesis, and VEGF staining	Adinolfi et al., 2012
Eyes Retinitis pigmentosa	rd1 transgenic mice	N.S.	14–21 days	i.vi.	20 mM PPADS (~2 mM in retina)	At day 14	Increased photoreceptor survival by 30%	Puthussery and Fletcher, 2009
ATP-induced photoreceptor degeneration	Sprague-Dawley rats	N.S.	8–12 weeks	i.vi.	20 mM PPADS (~1 mM in retina)	10 min before ATP injection	Protected photoreceptors against apoptosis	Puthussery and Fletcher, 2009
BzATP-induced retinal ganglion cell death	Long Evans rats	N.S.	14–26 days	i.vi.	10 mM BBG or 10 μM MRS 2540	1 μl of coinjected with BzATP over 20 s	Inhibited ganglion cell death	Hu et al., 2010
Pressure-induced retinal ganglion cell death	Wistar rats	M	10–12 weeks	i.vi.	1–100 μM oATP or 0.3–300 nM BBG	5 μl after elevating intraocular pressure	Dose-dependently reduced cell loss and P2X7, TNF- α , IL-1 β , and IL-6 mRNA	Sugiyama et al., 2013
Retinal pigmented epithelium degeneration	C57BL/6J mice	N.S.	N.S.	i.vi.	1 μl of A740003	24 hours before and 72 hours after inducing RPE degeneration	Protected against A β RNA-induced RPE degeneration	Kerur et al., 2013
Subretinal hemorrhage (age-related macular degeneration)	C57BL6/Jcl mice	M	8 weeks	Subretinal space	50 μM BBG	With autologous blood	Reduced TUNEL-detectable photoreceptor cell apoptosis	Notomi et al., 2013
Optic nerve crush injury	Wistar rats	M	280–320 g	Vitreous body	10–300 μM oATP or 0.003–30 μM BBG	Immediately after injury	oATP (30 μM) and BBG (0.003–3 μM) protected retinal ganglion cells from death and suppressed P2X7 up-regulation	Kakurai et al., 2013
Optic nerve injury	Piebald Virol Glaxo rats	F	160–200 g	s.c.	1 mM oATP	0.5 $\mu\text{l/h}$ for 2 weeks after partial transection of optic nerve	In combination with lomerizine and INQ improved injury parameters and visual function	Savigni et al., 2013

(continued)

TABLE 8—Continued

Disease/Model	Species and Strain	Sex	Start Age or Size	Injection Site	Antagonist and Dose	Injection Regimen	Outcomes	References
Heart Heart transplantation	C57BL/6 and BALB/c mice	N.S.	N.S.	i.p.	250 µg of oATP	1 ×/day for 14 days	Induced long-term transplant survival (80%) and anergy toward graft antigens, and reduced Th1/Th17 cells	Vergani et al., 2013b
Chronic heart cardiac transplant rejection	C57BL/6 mice	N.S.	N.S.	i.p.	250 µg of oATP	1 ×/day for 14 days	Prevented coronary vasculopathy, reduced cellular infiltrates and Th1/Th17 cells	Vergani et al., 2013b
Myocardial ischemia	Sprague-Dawley rats	N.S.	180–230 g	i.p.	30 mg/kg BBG	1 ×/day for 20 days 24 hours after occluding left coronary artery	Reduced elevation of systolic blood pressure, heart rate, cardiac enzymes, serum TNF-α and IL-6, and P2X7	Tu et al., 2013
Myocardial ischemia	Sprague-Dawley rats	N.S.	180–230 g	i.g.	1 mg/kg oATP	1 ×/day for 20 days	Reduced elevation of systolic blood pressure, heart rate, TNF-α, IL-6, serum cardiac enzymes, P2X7, and improved Q wave	Liu et al., 2013
Myocardial ischemia	Sprague-Dawley rats	M	180–220 g	i.p.	30 mg/kg BBG	1 ×/day for 7 days after occluding left coronary artery	Reduced elevation of P2X7 and p-ERK1/2 in superior cervical ganglia neurons	Kong et al., 2013
Kidney Antibody-mediated glomerulonephritis	Wistar Kyoto rats	M	180–220 g	N.S.	100 and 300 µmol/kg A438079	On day nephrotoxic serum injected then 2 ×/day for 7 days	Prevented glomerulonephritis, with reduced fibrinoid necrosis, proteinuria and glomeruli macrophage infiltration (300 but not 100 µmol/kg)	Taylor et al., 2009b
Lupus nephritis	MRL/lpr mice	F	12 weeks	i.p.	45.5 mg/kg BBG	Every 48 hours for 8 weeks	Reduced renal damage, NLRP3, ASC, caspase 1-p20, IL-1β, immune complex deposition, anti-dsDNA antibodies, Th17:Treg cell ratio and mortality	Zhao et al., 2013
Lupus nephritis	NZM2328 mice	F	3 months	i.p.	45.5 mg/kg BBG	Every 48 hours for 3–5 weeks after adenovirus-expressing interferon-α injection	Delayed proteinuria and reduced urinary albumin/creatinine ratio, serum BUN, anti-dsDNA antibodies, serum, and renal IL-1β, and C3 and IgG deposits	Zhao et al., 2013

(continued)

TABLE 8—Continued

Disease/Model	Species and Strain	Sex	Start Age or Size	Injection Site	Antagonist and Dose	Injection Regimen	Outcomes	References
Salt-sensitive hypertension and renal injury	Dahl salt-sensitive rats	N.S.	4–10 weeks (high-salt diet), antagonist start N.S.	i.p.	50 mg/kg BBG or 100 mg/kg A438079	Every day for 4 weeks (BBG) or 7 days (A438079)	Improved creatinine clearance and attenuated salt-sensitive hypertension, interstitial fibrosis (BBG), macrophage and T-cell infiltration, and urinary protein excretion (BBG, A438079)	Ji et al., 2012
Renal Injury	Fischer and Lewis rats	M	N.S.	i.v.	50 μ g/min/100 g BBG	Acute infusion	Reduced mean arterial blood pressure, renal vascular resistance, and pressure-diuresis threshold in Fischer but not Lewis rats	Menzies et al., 2013
Liver								
Diet-induced obesity	DIO C57BL6/J mice	M	10 weeks (high-fat diet), 16 weeks for antagonist	i.p.	300 μ mol/kg A438079	1 hour before CCl ₄ injection	Inhibited formation of protein radicals and tyrosine nitration, and decreased TNF- α and MCP-2 release from Kupffer cells	Chatterjee et al., 2012
Acetaminophen-induced hepatotoxicity	C57BL/6 mice	M	5–8 weeks	i.p.	300 mM/kg (2 mg) A438079	1 hour before or 2 hour after i.p. APAP injection	Pre- and post-treatment prevented acute liver injury and inflammation	Hoque et al., 2012
Acetaminophen-induced hepatotoxicity	C57BL/6	M	8 weeks	i.p.	80 mg/kg A438079	1 hour before i.p. APAP	Attenuated liver injury (plasma ALT activity and necrosis), GSH depletion, protein adducts, JNK activation, and P450 activities	Xie et al., 2013
Lung								
Chemical irritant-induced cough	Hartley guinea pigs	M	300–350 g	Aerosol	50 μ M PPADS	Exposed for 2 min, 5 min before ATP and irritants	No effect on ATP-induced enhancement of coughing	Kamei et al., 2005
Cigarette smoke-induced lung inflammation	C57/Bl6 mice	M	6–8 weeks	i.t.	10 μ M KN-62	Before cigarette smoke exposure, 5 times over 3 consecutive days	Reduced macrophage and neutrophil infiltrates and cytokines in bronchoalveolar lavage fluid	Lucattelli et al., 2011
Cigarette smoke- and LPS-induced airway inflammation	C57BL/6 mice	M	N.S.	p.o.	30–1000 mg/kg A438079	1 hour before first exposure and 1 hour after second exposure each day for 3 days	Dose-dependently inhibited airway neutrophilia induced by cigarette smoke but not LPS	Eltom et al., 2011
LPS-induced airway inflammation	Human CD39 overexpressing C57BL/6J mice	N.S.	8–12 weeks	i.t.	100 μ M A438079	50 μ l with LPS	Reduced neutrophil and lymphocyte, but not macrophage, recruitment; reduced keratinocyte chemoattractant levels 2 hours after LPS	Th��atre et al., 2012

(continued)

TABLE 8—Continued

Disease/Model	Species and Strain	Sex	Start Age or Size	Injection Site	Antagonist and Dose	Injection Regimen	Outcomes	References
OVA/alum-induced allergic airway inflammation	BALB/c and C57/BL6 mice	F (chronic) N.S. (acute)	6–8 weeks	i.p. (acute) or i.t. (chronic)	1 μ M KN-62	Before OVA exposure on days 19–21 after sensitization (acute) or 3 days per week for 2 weeks (chronic)	Reduced airway eosinophilia, goblet cell hyperplasia, bronchial hyperresponsiveness to methacholine, and airway remodeling	Müller et al., 2011
Pancreas Autoimmune diabetes	RIP-GP transgenic mice	N.S.	N.S.	i.v.	1 mg of oATP	4 days after LCMV-WE infection	Suppressed diabetes in 75% of treated mice by affecting T-cell proliferation	Lang et al., 2010
Islet allograft rejection	C57BL/6 and BALB/c mice	N.S.	N.S.	i.p.	250 μ g of oATP	1 \times /day for 14 days	Prolonged allograft survival and reduced severity of autoimmune response (IFN- γ , IL-6, cellular infiltrates, and Th1 cells)	Vergani et al., 2013a
Acute pancreatitis	C57BL/6 mice	M	5–8 weeks	s.c.	300 μ mol/kg A438079	1 hour before 1st cerulean injection	Reduced pancreatic edema and leukocyte infiltrate	Hoque et al., 2011
Skin Allergic contact dermatitis	C57BL/6 mice	N.S.	N.S.	Pinna of ear	3.6 μ g of KN-62	4 hours before TNCB sensitization	Prevented sensitization and contact hypersensitivity	Weber et al., 2010
Graft-versus-host disease	BALB/c mice	N.S.	6–12 weeks	i.p.	10 μ mol of PPADS or 1 μ mol of KN-62	Days 0–10	Reduced mortality, disease severity (both), donor T-cell expansion, and IFN- γ production (PPADS)	Wilhelm et al., 2010
Melanoma	C57BL/6 mice	F	5–7 weeks, 15–18 g	Tumor	300 nM AZ10606120	100 μ l 5, 7, 9, and 13 days after B16 cell injection	Reduced tumor growth, VEGF staining, and vessel formation	Adinolfi et al., 2012
Melanoma	C57BL/6 mice	M	6 weeks	i.p.	0.5 mg of oATP	5, 15, and 25 days after B16 cell injection	Reduced tumor growth	Hattori et al., 2012

ALT, alanine aminotransferase; APAP, acetaminophen; BUN, blood urea nitrogen; DIO, diet-induced obesity; ERK, extracellular signal-regulated kinase; GSH, glutathione; i.g., intragastric; IL, interleukin; INQ, 2-[7-(H-imidazol-1-yl)-6-nitro-2,3-dioxo-1,2,3,4-tetrahydro quinoxalin-1-yl]acetic acid; i.p., intraperitoneal; i.v., intravenous; JNK, c-jun-N-terminal kinase; LCMV-WE, lymphocytic choriomeningitis virus, strain WE; MCP-2, monocyte chemoattractant protein 2; OVA, ovalbumin; RPE, retinal pigmented epithelium; STAT3, signal transducer and activator of transcription 3; Th1, T helper type 1; TNCB, 2,4,6-trinitrochloro-1-benzene; TNF, tumor necrosis factor; TUNEL, terminal deoxynucleotidyl transferase dUTP nick end labeling.

As a result of these studies, both groups concluded that P2X7 does not appear to be a useful target in rheumatoid arthritis (Keystone et al., 2012; Stock et al., 2012). However, the possibility remains that the efficacy of these P2X7 antagonists is dependent on *P2RX7* genotype, as indicated in other *in vitro* studies using GSK1370319A (McHugh et al., 2012). Thus, it may be essential to stratify clinical data by *P2RX7* genotype to detect therapeutic benefits of P2X7 blockade. The possibility also remains that P2X7 blockade is of no therapeutic benefit in patients who are refractory to methotrexate or sulfasalazine, because both studies used a patient cohort already nonresponsive to these therapies. Thus, it may be of value to explore if the therapeutic benefit of P2X7 blockade is influenced by methotrexate or sulfasalazine resistance. In the meantime, the trial of AZD9056 or CE-224,535 in other human disorders and the trial of other compounds, such as those from GSK (Abberley et al., 2010) and Johnson & Johnson (Bhattacharya et al., 2013), await.

XII. Modulators of P2X7 Receptor Expression

P2X7 is widely distributed throughout the mammalian body (see section I); however, the relative expression and distribution of this receptor between cell types and tissues have not been fully elucidated. As also noted above (section I), future studies of P2X7 reporter mice expressing enhanced green fluorescent protein (Engel et al., 2012; Garcia-Huerta et al., 2012; Jimenez-Pacheco et al., 2013) will be useful for visualizing P2X7 expression patterns in both healthy and diseased animals. However, studies comparing the relative expression of P2X7 on cell types between mice and humans are lacking, and the possibility remains that this expression may not be comparable. In humans, P2X7 is present in decreasing amounts on monocytes, B cells and T cells (Gu et al., 2000), whereas in dogs P2X7 is present in greatest amounts on T cells and is present at lower amounts on monocytes and B cells (Stevenson et al., 2009). Nevertheless, modulation of P2X7 expression may provide an alternate therapeutic strategy in addition to pharmacological blockade.

A range of inflammatory mediators can modulate P2X7 expression and its subsequent function on various cell types. Proinflammatory mediators such as interferon- γ (IFN- γ), TNF- α , and LPS can upregulate P2X7 on macrophages (Humphreys and Dubyak, 1998), umbilical vein endothelial cells (Wilson et al., 2007), and epithelial cells (Welter-Stahl et al., 2009). Conversely, anti-inflammatory mediators including transforming growth factor- β 1, IL-4, and IL-10 can downregulate P2X7 on macrophages (Lemaire and Leduc, 2003; Gadeock et al., 2010). As a result of these and other studies, the possibility remains that such mediators may be used to regulate P2X7 expression and its subsequent function in a range of disorders. Other compounds may also be of value in

modulating P2X7 expression. Recently, the dialdehydic compound 2-[1-(6-aminopurin-9-yl)-2-oxo-*etossil*prop-2-enale (MED1011) was developed and shown to downregulate the expression and function of P2X7 in macrophages by causing the internalization of P2X7 in these cells (Muzzachi et al., 2013). Moreover, the medicinal extract from Aloe vera has been shown to downregulate P2X7 expression in macrophages to attenuate IL-1 β secretion (Budai et al., 2013). Alternatively, P2X7 expression could potentially be down modulated by use of RNA interference. Small interfering RNA has been used to specifically downregulate P2X7 in mice and rats to attenuate diseases such as brain injury (Chen et al., 2013a) and nephritis (Zhao et al., 2013). Finally, P2X7 expression and its subsequent function could be downregulated by targeting the specificity protein 1 transcription factor (Sp1), which plays an important role in the transcriptional regulation of P2X7 (Garcia-Huerta et al., 2012). Of note, targeting Sp1 by small hairpin RNA or the antibiotic mithramycin, which blocks Sp1 activity, downregulates P2X7 *in vitro* (Garcia-Huerta et al., 2012).

XIII. Conclusions and Future Directions

Rodent models are valuable tools for investigating the role of P2X7 in health and disease. However, there are a number of factors that need to be considered when interpreting the results of such studies in the context of human disease. First, unlike human P2X7, rodent P2X7 can be activated by the alternate ligand NAD to induce similar downstream signaling events to that of ATP activation. Second, P2X7 variants, such as the murine P451L SNP and splice isoforms P2X7K, P2X7 13B, and P2X7 13C, contribute to the diversity of receptor-mediated responses, thus adding further complexity to P2X7 studies in rodents. This is particularly highlighted by the currently available P2X7 KO mice, in which some of these variants have escaped deletion. Identification of P2X7 in the guinea pig, dog, and Rhesus macaque, in addition to the rat and mouse, highlights the potential of these species for the future study of P2X7-related diseases and in the mandatory testing and evaluation of preclinical drugs targeting P2X7.

In general, a variety of P2X7 antagonists have been reported to be efficacious in rodent models of neurologic disease and injury, compound-induced inflammation, musculoskeletal disorders, and disorders associated with other tissues. In these studies, there was no clear consensus on the use of P2X7 antagonists in terms of doses or injection sites, with a variety of regimens successfully employed. Although nonspecific P2X7 antagonists were used in a number of these studies, the outcomes of some have been confirmed in more recent years through the use of more specific P2X7 antagonists. This suggests that P2X7 may play a role in a wide variety of diseases, implicating this receptor as a potential

therapeutic target in a broad range of settings. However, differential activity of P2X7 antagonists is observed at receptors between individuals and species, presumably as a result of P2X7 variants and variations in amino acid homology in antagonist binding sites. Thus, this may be a limitation of translating preclinical studies of P2X7 antagonists in rodents to clinical trials in humans.

Although rodent models and P2X7 antagonists have been valuable tools for investigating P2X7 in vivo, the following considerations may prove useful in future studies. First, the use of more selective P2X7 antagonists will support the current in vivo literature proposing a role for P2X7 in disease and injury. However, this may be limited for many laboratories because of the lack of availability or high cost of such antagonists. Second, although a number of commonly used antagonists are branded as P2X7 specific, there is still an absence of comprehensive information on the specificity of these compounds against other P2 receptor members, especially for mouse P2X7. In addition, data are lacking on the effects of many of these antagonists on recombinant murine variants. It will be important to obtain more complete pharmacological profiles for some of the widely used P2X7 antagonists, as previously done at a number of human and rat P2X subtypes (Bianchi et al., 1999; Donnelly-Roberts et al., 2009). Third, the generation of a complete P2X7 KO mouse, to negate the effect of escaped P2X7 variants, will be of value for future investigations. Fourth, the use of P2X7 antagonists in conjunction with various KO mice may be useful for determining the mechanisms of P2X7 action in vivo. For example, the use of the P2X7 antagonist A839977 in IL-1 α KO mice, which lack genes for both IL-1 α and IL-1 β , has suggested that the positive effects of P2X7 inhibition in inflammatory pain models are mediated by blocking IL-1 β release (Honore et al., 2009). Fifth, with the emergence of biologics (therapeutic antibodies) (Hatcher et al., 2011), anti-P2X7 antibodies represent promising tools for investigating the roles of P2X7 in disease and injury. In this regard, an anti-P2X7 monoclonal antibody was recently generated and demonstrated to inhibit mast cell activation and prevent the resulting intestinal inflammation in mice (Kurashima et al., 2012). Moreover, an anti-P2X7 monoclonal antibody has long been known to impair human P2X7 (Buell et al., 1998). Of note, this same antibody is capable of binding to rat and guinea pig P2X7 in transfected HEK-293 and human osteosarcoma U-2 OS cells (Smart et al., 2003; Fonfria et al., 2008), but whether this antibody is capable of blocking the activation of this receptor in these species remains unknown. Furthermore, single-domain antibodies, or nanobodies, have been shown to be promising tools for inhibiting specific membrane proteins (Pardon et al., 2014). In one pioneer study, an ART2.2-specific single-domain antibody was generated and shown to completely block ART2.2 enzymatic activity and NAD-induced cytotoxicity in vivo (Koch-Nolte et al., 2007). Thus, these antibodies could be

used indirectly and reversibly to block P2X7 activation in mice. Likewise, this same technology is being applied to develop single-domain antibodies to P2X7 to directly target or inhibit this receptor (Laeremans et al., 2010). Finally, despite the development and patenting of a number of P2X7 antagonists for the treatment of human disease (Gunosewoyo and Kassiou, 2010), we are still awaiting results on the efficacy of many of these antagonists in humans, as well as the outcomes of current clinical trials, which will further influence future attempts to target P2X7.

Acknowledgments

The authors thank J. J. Yerbury, M. L. Sanderson-Smith, and A. Pupovac (University of Wollongong) for critical review.

Authorship Contributions

Wrote or contributed to the writing of the manuscript: Bartlett, Stokes, Sluyter.

References

- Abberley L, Bebius A, Beswick PJ, Billinton A, Collis KL, Dean DK, Fonfria E, Gleave RJ, Medhurst SJ, and Michel AD, et al. (2010) Identification of 2-oxo-N-(phenylmethyl)-4-imidazolidinecarboxamide antagonists of the P2X7(7) receptor. *Bioorg Med Chem Lett* **20**:6370–6374.
- Abd Elmageed ZY, Naura AS, Errami Y, and Zerfaoui M (2012) The poly(ADP-ribose) polymerases (PARPs): new roles in intracellular transport. *Cell Signal* **24**:1–8.
- Acuña-Castillo C, Coddou C, Bull P, Brito J, and Huidobro-Toro JP (2007) Differential role of extracellular histidines in copper, zinc, magnesium and proton modulation of the P2X7 purinergic receptor. *J Neurochem* **101**:17–26.
- Adinolfi E, Callegari MG, Ferrari D, Bolognesi C, Minelli M, Wiekowski MR, Pinton P, Rizzuto R, and Di Virgilio F (2005) Basal activation of the P2X7 ATP receptor elevates mitochondrial calcium and potential, increases cellular ATP levels, and promotes serum-independent growth. *Mol Biol Cell* **16**:3260–3272.
- Adinolfi E, Cirillo M, Woltersdorf R, Falzoni S, Chiozzi P, Pellegatti P, Callegari MG, Sandoña D, Markwardt F, and Schmalzing G, et al. (2010) Trophic activity of a naturally occurring truncated isoform of the P2X7 receptor. *FASEB J* **24**:3393–3404.
- Adinolfi E, Raffaghello L, Giuliani AL, Cavazzini L, Capece M, Chiozzi P, Bianchi G, Kroemer G, Pistoia V, and Di Virgilio F (2012) Expression of P2X7 receptor increases in vivo tumor growth. *Cancer Res* **72**:2957–2969.
- Adriouch S, Dox C, Welge V, Seman M, Koch-Nolte F, and Haag F (2002) Cutting edge: a natural P451L mutation in the cytoplasmic domain impairs the function of the mouse P2X7 receptor. *J Immunol* **169**:4108–4112.
- Adriouch S, Ohlrogge W, Haag F, Koch-Nolte F, and Seman M (2001) Rapid induction of naive T cell apoptosis by ecto-nicotinamide adenine dinucleotide: requirement for mono(ADP-ribosyl)transferase 2 and a downstream effector. *J Immunol* **167**:196–203.
- Aktories K (2011) Bacterial protein toxins that modify host regulatory GTPases. *Nat Rev Microbiol* **9**:487–498.
- Al-Shukaili A, Al-Kaabi J, Hassan B, Al-Araimi T, Al-Tobi M, Al-Kindi M, Al-Maniri A, Al-Gheilani A, and Al-Ansari A (2011) P2X7 receptor gene polymorphism analysis in rheumatoid arthritis. *Int J Immunogenet* **38**:389–396.
- Alberto AVP, Faria RX, Couto CGC, Ferreira LGB, Souza CAM, Teixeira PCN, Fróes MM, and Alves LA (2013) Is pannexin the pore associated with the P2X7 receptor? *Naunyn Schmiedeberg's Arch Pharmacol* **386**:775–787.
- Alcaraz L, Baxter A, Bent J, Bowers K, Braddock M, Cladingboel D, Donald D, Fagura M, Furber M, and Laurent C, et al. (2003) Novel P2X7 receptor antagonists. *Bioorg Med Chem Lett* **13**:4043–4046.
- Andó RD, Méhécz B, Gyires K, Illes P, and Sperlágh B (2010) A comparative analysis of the activity of ligands acting at P2X and P2Y receptor subtypes in models of neuropathic, acute and inflammatory pain. *Br J Pharmacol* **159**:1106–1117.
- Andrei C, Margiocco P, Poggi A, Lotti LV, Torrisi MR, and Rubartelli A (2004) Phospholipases C and A₂ control lysosome-mediated IL-1 β secretion: Implications for inflammatory processes. *Proc Natl Acad Sci USA* **101**:9745–9750.
- Apolloni S, Parisi C, Pesaresi MG, Rossi S, Carri MT, Cozzolino M, Volonté C, and D'Ambrosi N (2013) The NADPH oxidase pathway is dysregulated by the P2X7 receptor in the SOD1-G93A microglia model of amyotrophic lateral sclerosis. *J Immunol* **190**:5187–5195.
- Arbeola J, Pérez-Samartín A, Gottlieb M, and Matute C (2012) P2X7 receptor blockade prevents ATP excitotoxicity in neurons and reduces brain damage after ischemia. *Neurobiol Dis* **45**:954–961.
- Ardissone V, Radaelli E, Zaratin P, Ardizzone M, Ladel C, Gattorno M, Martini A, Grassi F, and Traggiai E (2011) Pharmacologic P2X purinergic receptor antagonism in the treatment of collagen-induced arthritis. *Arthritis Rheum* **63**:3323–3332.
- Avshalumov MV and Rice ME (2003) Activation of ATP-sensitive K⁺ (K(ATP)) channels by H₂O₂ underlies glutamate-dependent inhibition of striatal dopamine release. *Proc Natl Acad Sci USA* **100**:11729–11734.
- Baas T (2012) Paradoxical P2X7. *SciBX* **5**:5–7.

- Banke TG, Chaplan SR, and Wickenden AD (2010) Dynamic changes in the TRPA1 selectivity filter lead to progressive but reversible pore dilation. *Am J Physiol Cell Physiol* **298**:C1457–C1468.
- Barberá-Cremades M, Baroja-Mazo A, Gomez AI, Machado F, Di Virgilio F, and Pelegrín P (2012) P2X7 receptor-stimulation causes fever via PGE₂ and IL-1 β release. *FASEB J* **26**:2951–2962.
- Barden N, Harvey M, Gagné B, Shink E, Tremblay M, Raymond C, Labbé M, Villeneuve A, Rochette D, and Bordeleau L, et al. (2006) Analysis of single nucleotide polymorphisms in genes in the chromosome 12Q24.31 region points to P2RX7 as a susceptibility gene to bipolar affective disorder. *Am J Med Genet B Neuro-psychiatr Genet* **141B**:374–382.
- Baricordi OR, Ferrari D, Melchiorri L, Chiozzi P, Hanau S, Chiari E, Rubini M, and Di Virgilio F (1996) An ATP-activated channel is involved in mitogenic stimulation of human T lymphocytes. *Blood* **87**:682–690.
- Baricordi OR, Melchiorri L, Adinolfi E, Falzoni S, Chiozzi P, Buell G, and Di Virgilio F (1999) Increased proliferation rate of lymphoid cells transfected with the P2X(7) ATP receptor. *J Biol Chem* **274**:33206–33208.
- Bartlett R, Yerbury JJ, and Sluyter R (2013) P2X7 receptor activation induces reactive oxygen species formation and cell death in murine EOC13 microglia. *Mediators Inflamm* **2013**:271813.
- Basso AM, Bratcher NA, Harris RR, Jarvis MF, Decker MW, and Rueter LE (2009) Behavioral profile of P2X7 receptor knockout mice in animal models of depression and anxiety: relevance for neuropsychiatric disorders. *Behav Brain Res* **198**:83–90.
- Beigi RD, Kertesz SB, Aquilina G, and DUBYAK GR (2003) Oxidized ATP (oATP) attenuates proinflammatory signaling via P2 receptor-independent mechanisms. *Br J Pharmacol* **140**:507–519.
- Bhaskaracharya A, Dao-Ung P, Jalilian I, Spildrejerde M, Skarratt KK, Fuller SJ, Sluyter R, and Stokes L (2014) Probenecid blocks human P2X7 receptor-induced dye uptake via a pannexin-1 independent mechanism. *PLoS ONE* **9**:e93058.
- Bhattacharya A, Neff RA, and Wickenden AD (2011) The physiology, pharmacology and future of P2X7 as an analgesic drug target: hype or promise? *Curr Pharm Biotechnol* **12**:1698–1706.
- Bhattacharya A, Wang Q, Ao H, Shoblock JR, Lord B, Aluisio L, Fraser I, Nepomuceno D, Neff RA, and Welty N, et al. (2013) Pharmacological characterization of a novel centrally permeable P2X7 receptor antagonist: JNJ-47965567. *Br J Pharmacol* **170**:624–640.
- Bianchi BR, Lynch KJ, Touma E, Niforatos W, Burgard EC, Alexander KM, Park HS, Yu H, Metzger R, and Kowaluk E, et al. (1999) Pharmacological characterization of recombinant human and rat P2X receptor subtypes. *Eur J Pharmacol* **376**:127–138.
- Bianco F, Ceruti S, Colombo A, Fumagalli M, Ferrari D, Pizzirani C, Matteoli M, Di Virgilio F, Abbracchio MP, and Verderio C (2006) A role for P2X7 in microglial proliferation. *J Neurochem* **99**:745–758.
- Boucher AA, Arnold JC, Hunt GE, Spiro A, Spencer J, Brown C, McGregor IS, Bennett MR, and Kassiu M (2011) Resilience and reduced c-Fos expression in P2X7 receptor knockout mice exposed to repeated forced swim test. *Neuroscience* **189**:170–177.
- Bradley HJ, Baldwin JM, Goli GR, Johnson B, Zou J, Sivaprasadarao A, Baldwin SA, and Jiang L-H (2011a) Residues 155 and 348 contribute to the determination of P2X7 receptor function via distinct mechanisms revealed by single-nucleotide polymorphisms. *J Biol Chem* **286**:8176–8187.
- Bradley HJ, Browne LE, Yang W, and Jiang LH (2011b) Pharmacological properties of the rhesus macaque monkey P2X7 receptor. *Br J Pharmacol* **164** (2b):743–754.
- Brandao-Burch A, Key ML, Patel JJ, Arnett TR, and Orriss IR (2012) The P2X7 receptor is an important regulator of extracellular ATP levels. *Front Endocrinol (Lausanne)* **3**:41.
- Broom DC, Matson DJ, Bradshaw E, Buck ME, Meade R, Coombs S, Matchett M, Ford KK, Yu W, and Yuan J, et al. (2008) Characterization of N-(adamantan-1-ylmethyl)-5-(3R-amino-pyrrolidin-1-yl)methyl-2-chloro-benzamide, a P2X7 antagonist in animal models of pain and inflammation. *J Pharmacol Exp Ther* **327**:620–633.
- Browne LE, Compan V, Bragg L, and North RA (2013) P2X7 receptor channels allow direct permeation of nanometer-sized dyes. *J Neurosci* **33**:3557–3566.
- Buchanan MS, Carroll AR, Addepalli R, Avery VM, Hooper JN, and Quinn RJ (2007a) Natural products, stylissadines A and B, specific antagonists of the P2X7 receptor, an important inflammatory target. *J Org Chem* **72**:2309–2317.
- Buchanan MS, Carroll AR, Addepalli R, Avery VM, Hooper JN, and Quinn RJ (2007b) Niphatoxin C, a cytotoxic tripyridine alkaloid from *Callispongia* sp. *J Nat Prod* **70**:2040–2041.
- Budai MM, Varga A, Milesz S, Tózsér J, and Benkő S (2013) Aloe vera downregulates LPS-induced inflammatory cytokine production and expression of NLRP3 inflammasome in human macrophages. *Mol Immunol* **56**:471–479.
- Buell G, Chessell IP, Michel AD, Collo G, Salazzo M, Herren S, Gretener D, Grahames C, Kaur R, and Kosco-Vilbois MH, et al. (1998) Blockade of human P2X7 receptor function with a monoclonal antibody. *Blood* **92**:3521–3528.
- Burnstock G (1972) Purinergic nerves. *Pharmacol Rev* **24**:509–581.
- Burnstock G (2006) Pathophysiology and therapeutic potential of purinergic signaling. *Pharmacol Rev* **58**:58–86.
- Burnstock G (2007) Purine and pyrimidine receptors. *Cell Mol Life Sci* **64**:1471–1483.
- Burnstock G and Knight GE (2004) Cellular distribution and functions of P2 receptor subtypes in different systems. *Int Rev Cytol* **240**:31–304.
- Cabrini G, Falzoni S, Forchap SL, Pellegatti P, Balboni A, Agostini P, Cuneo A, Castoldi G, Baricordi OR, and Di Virgilio F (2005) A His-155 to Tyr polymorphism confers gain-of-function to the human P2X7 receptor of human leukemic lymphocytes. *J Immunol* **175**:82–89.
- Cario-Toumaniantz C, Loirand G, Ferrier L, and Pacaud P (1998) Non-genomic inhibition of human P2X7 purinoceptor by 17 β -oestradiol. *J Physiol* **508**:659–666.
- Cascabulho CM, Bani Corrêa C, Cotta-de-Almeida V, and Henriques-Pons A (2012) Defective T-lymphocyte migration to muscles in dystrophin-deficient mice. *Am J Pathol* **181**:593–604.
- Cervetto C, Alloisio S, Frattaroli D, Mazzotta MC, Milanese M, Gavazzo P, Passalacqua M, Nobile M, Maura G, and Marcoli M (2013a) The P2X7 receptor as a route for non-exocytotic glutamate release: dependence on the carboxyl tail. *J Neurochem* **124**:821–831.
- Cervetto C, Frattaroli D, Maura G, and Marcoli M (2013b) Motor neuron dysfunction in a mouse model of ALS: gender-dependent effect of P2X7 antagonism. *Toxicology* **311**:69–77.
- Chatterjee S, Rana R, Corbett J, Kadiiska MB, Goldstein J, and Mason RP (2012) P2X7 receptor-NADPH oxidase axis mediates protein radical formation and Kupffer cell activation in carbon tetrachloride-mediated steatohepatitis in obese mice. *Free Radic Biol Med* **52**:1666–1679.
- Chaumont S and Khakh BS (2008) Patch-clamp coordinated spectroscopy shows P2X2 receptor permeability dynamics require cytosolic domain rearrangements but not Panx-1 channels. *Proc Natl Acad Sci USA* **105**:12063–12068.
- Cheewatrakoolpong B, Gilchrist H, Anthes JC, and Greenfeder S (2005) Identification and characterization of splice variants of the human P2X7 ATP channel. *Biochem Biophys Res Commun* **332**:17–27.
- Chen L and Brosnan CF (2006) Exacerbation of experimental autoimmune encephalomyelitis in P2X7R^{-/-} mice: evidence for loss of apoptotic activity in lymphocytes. *J Immunol* **176**:3115–3126.
- Chen ML, Cao H, Chu YX, Cheng LZ, Liang LL, Zhang YQ, and Zhao ZQ (2012) Role of P2X7 receptor-mediated IL-18/IL-18R signaling in morphine tolerance: multiple glial-neuronal dialogues in the rat spinal cord. *J Pain* **13**:945–958.
- Chen S, Ma Q, Krafft PR, Chen Y, Tang J, Zhang J, and Zhang JH (2013a) P2X7 receptor antagonism inhibits p38 mitogen-activated protein kinase activation and ameliorates neuronal apoptosis after subarachnoid hemorrhage in rats. *Crit Care Med* **41**:e466–e474.
- Chen S, Ma Q, Krafft PR, Hu Q, Rolland W 2nd, Sherchan P, Zhang J, Tang J, and Zhang JH (2013b) P2X7R/cryopyrin inflammasome axis inhibition reduces neuroinflammation after SAH. *Neurobiol Dis* **58**:296–307.
- Chessell IP, Hatcher JP, Bountra C, Michel AD, Hughes JP, Green P, Egerton J, Murfin M, Richardson J, and Peck WL, et al. (2005) Disruption of the P2X7 purinoceptor gene abolishes chronic inflammatory and neuropathic pain. *Pain* **114**:386–396.
- Chessell IP, Simon J, Hibell AD, Michel AD, Barnard EA, and Humphrey PPA (1998) Cloning and functional characterisation of the mouse P2X7 receptor. *FEBS Lett* **439**:26–30.
- Chiao CW, da Silva-Santos JE, Giachini FR, Tostes RC, Su MJ, and Webb RC (2013) P2X7 receptor activation contributes to an initial upstream mechanism of lipopolysaccharide-induced vascular dysfunction. *Clin Sci (Lond)* **125**:131–141.
- Choi HB, Ryu JK, Kim SU, and McLarnon JG (2007) Modulation of the purinergic P2X7 receptor attenuates lipopolysaccharide-mediated microglial activation and neuronal damage in inflamed brain. *J Neurosci* **27**:4957–4968.
- Chow SC, Kass GEN, and Orrenius S (1997) Purines and their roles in apoptosis. *Neuropharmacology* **36**:1149–1156.
- Chu K, Yin B, Wang J, Peng G, Liang H, Xu Z, Du Y, Fang M, Xia Q, and Luo B (2012) Inhibition of P2X7 receptor ameliorates transient global cerebral ischemia/reperfusion injury via modulating inflammatory responses in the rat hippocampus. *J Neuroinflammation* **9**:69.
- Chu YX, Zhang Y, Zhang YQ, and Zhao ZQ (2010) Involvement of microglial P2X7 receptors and downstream signaling pathways in long-term potentiation of spinal nociceptive responses. *Brain Behav Immun* **24**:1176–1189.
- Chung M-K, Güler AD, and Caterina MJ (2008) TRPV1 shows dynamic ionic selectivity during agonist stimulation. *Nat Neurosci* **11**:555–564.
- Clark AK, Staniland AA, Marchand F, Kaaan TKY, McMahon SB, and Malcangio M (2010) P2X7-dependent release of interleukin-1 β and nociception in the spinal cord following lipopolysaccharide. *J Neurosci* **30**:573–582.
- Cockcroft S and Gomperts BD (1979) Activation and inhibition of calcium-dependent histamine secretion by ATP ions applied to rat mast cells. *J Physiol* **296**:229–243.
- Coddou C, Yan Z, Obsil T, Huidobro-Toro JP, and Stojilkovic SS (2011) Activation and regulation of purinergic P2X receptor channels. *Pharmacol Rev* **63**:641–683.
- Compan V, Baroja-Mazo A, López-Castejón G, Gomez AI, Martínez CM, Angosto D, Montero MT, Herranz AS, Bazán E, and Reimers D, et al. (2012a) Cell volume regulation modulates NLRP3 inflammasome activation. *Immunity* **37**:487–500.
- Compan V, Ulmann L, Stelmashenko O, Chemin J, Chaumont S, and Rassendren F (2012b) P2X2 and P2X5 subunits define a new heteromeric receptor with P2X7-like properties. *J Neurosci* **32**:4284–4296.
- Csölle C, Baranyi M, Zsilla G, Kittel A, Gölöncsér F, Illes P, Papp E, Vizi ES, and Sperlágh B (2013) Neurochemical changes in the mouse hippocampus underlying the antidepressant effect of genetic deletion of P2X7 receptors. *PLoS ONE* **8**:e66547.
- Csölle C and Sperlágh B (2010) Peripheral origin of IL-1 β production in the rodent hippocampus under in vivo systemic bacterial lipopolysaccharide (LPS) challenge and its regulation by P2X(7) receptors. *J Neuroimmunol* **219**:38–46.
- da Silva GL, Sperotto NDM, Borges TJ, Bonorino C, Takyia CM, Coutinho-Silva R, Campos MM, Zanin RF, and Morrone FB (2013) P2X7 receptor is required for neutrophil accumulation in a mouse model of irritant contact dermatitis. *Exp Dermatol* **22**:184–188.
- Delarasse C, Auger R, Gonnord P, Fontaine B, and Kanellopoulos JM (2011) The purinergic receptor P2X7 triggers α -secretase-dependent processing of the amyloid precursor protein. *J Biol Chem* **286**:2596–2606.
- Dell'Antonio G, Quattrini A, Cin ED, Fulgenzi A, and Ferrero ME (2002a) Relief of inflammatory pain in rats by local use of the selective P2X7 ATP receptor inhibitor, oxidized ATP. *Arthritis Rheum* **46**:3378–3385.
- Dell'Antonio G, Quattrini A, Dal Cin E, Fulgenzi A, and Ferrero ME (2002b) Antinociceptive effect of a new P(2Z)/P2X7 antagonist, oxidized ATP, in arthritic rats. *Neurosci Lett* **327**:87–90.
- Di Virgilio F (2013) The therapeutic potential of modifying inflammasomes and NOD-like receptors. *Pharmacol Rev* **65**:872–905.

- Diaz-Hernandez JI, Gomez-Villafuertes R, León-Otegui M, Hontecillas-Prieto L, Del Puerto A, Trejo JL, Lucas JJ, Garrido JJ, Gualix J, and Miras-Portugal MT, et al. (2012) In vivo P2X7 inhibition reduces amyloid plaques in Alzheimer's disease through GSK3 β and secretases. *Neurobiol Aging* **33**:1816–1828.
- Diaz-Hernández M, Diez-Zaera M, Sánchez-Nogueiro J, Gómez-Villafuertes R, Canals JM, Alberch J, Miras-Portugal MT, and Lucas JJ (2009) Altered P2X7-receptor level and function in mouse models of Huntington's disease and therapeutic efficacy of antagonist administration. *FASEB J* **23**:1893–1906.
- Donnelly-Roberts DL and Jarvis MF (2007) Discovery of P2X7 receptor-selective antagonists offers new insights into P2X7 receptor function and indicates a role in chronic pain states. *Br J Pharmacol* **151**:571–579.
- Donnelly-Roberts DL, Namovic MT, Faltynek CR, and Jarvis MF (2004) Mitogen-activated protein kinase and caspase signaling pathways are required for P2X7 receptor (P2X7R)-induced pore formation in human THP-1 cells. *J Pharmacol Exp Ther* **308**:1053–1061.
- Donnelly-Roberts DL, Namovic MT, Han P, and Jarvis MF (2009) Mammalian P2X7 receptor pharmacology: comparison of recombinant mouse, rat and human P2X7 receptors. *Br J Pharmacol* **157**:1203–1214.
- Duan S, Anderson CM, Keung EC, Chen Y, Chen Y, and Swanson RA (2003) P2X7 receptor-mediated release of excitatory amino acids from astrocytes. *J Neurosci* **23**:1320–1328.
- Dunn PM and Blakeley AGH (1988) Suramin: a reversible P2-purinoceptor antagonist in the mouse vas deferens. *Br J Pharmacol* **93**:243–245.
- Duplantier AJ, Dombroski MA, Subramanyam C, Beaulieu AM, Chang SP, Gabel CA, Jordan C, Kalgutkar AS, Kraus KG, and Labasi JM, et al. (2011) Optimization of the physicochemical and pharmacokinetic attributes in a 6-azauracil series of P2X7 receptor antagonists leading to the discovery of the clinical candidate CE-224,535. *Bioorg Med Chem Lett* **21**:3708–3711.
- el-Moatassim C and Dubyak GR (1992) A novel pathway for the activation of phospholipase D by P_{2z} purinergic receptors in BAC1.2F5 macrophages. *J Biol Chem* **267**:23664–23673.
- El Ouaali M, Seil M, and Dehaye JP (2012) Activation of calcium-insensitive phospholipase A₂ (iPLA₂) by P2X₇ receptors in murine peritoneal macrophages. *Prostaglandins Other Lipid Mediat* **99**:116–123.
- Ellsner A, Duncan M, Gavrilin M, and Wewers MD (2004) A novel P2X7 receptor activator, the human cathelicidin-derived peptide LL37, induces IL-1 β processing and release. *J Immunol* **172**:4987–4994.
- Eltom S, Stevenson CS, Rastrick J, Dale N, Raemdonck K, Wong S, Catley MC, Belvisi MG, and Birrell MA (2011) P2X7 receptor and caspase 1 activation are central to airway inflammation observed after exposure to tobacco smoke. *PLoS ONE* **6**:e24097.
- Endemann G, Stanton LW, Madden KS, Bryant CM, White RT, and Protter AA (1993) CD36 is a receptor for oxidized low density lipoprotein. *J Biol Chem* **268**:11811–11816.
- Engel T, Gomez-Villafuertes R, Tanaka K, Mesuret G, Sanz-Rodriguez A, Garcia-Huerta P, Miras-Portugal MT, Henshall DC, and Diaz-Hernandez M (2012) Seizure suppression and neuroprotection by targeting the purinergic P2X7 receptor during status epilepticus in mice. *FASEB J* **26**:1616–1628.
- Evans RJ, Lewis C, Buell G, Valera S, North RA, and Surprenant A (1995) Pharmacological characterization of heterologously expressed ATP-gated cation channels (P2x purinoceptors). *Mol Pharmacol* **48**:178–183.
- Fang KM, Wang YL, Huang MC, Sun SH, Cheng H, and Tzeng SF (2011) Expression of macrophage inflammatory protein-1 α and monocyte chemoattractant protein-1 in glioma-infiltrating microglia: involvement of ATP and P2X₇ receptor. *J Neurosci Res* **89**:199–211.
- Faria RX, Defarias FP, and Alves LA (2005) Are second messengers crucial for opening the pore associated with P2X7 receptor? *Am J Physiol Cell Physiol* **288**:C260–C271.
- Faria RX, Reis RAM, Casabulho CM, Alberto AVP, de Farias FP, Henriques-Pons A, and Alves LA (2009) Pharmacological properties of a pore induced by raising intracellular Ca²⁺. *Am J Physiol Cell Physiol* **297**:C28–C42.
- Felix RA, Martin S, Pinion S, and Crawford DJ (2012) Development of a comprehensive set of P2 receptor pharmacological research compounds. *Purinergic Signal* **8** (Suppl 1):101–112.
- Fellin T, Pozzan T, and Carmignoto G (2006) Purinergic receptors mediate two distinct glutamate release pathways in hippocampal astrocytes. *J Biol Chem* **281**:4274–4284.
- Feng YH, Li X, Wang L, Zhou L, and Gorodeski GI (2006) A truncated P2X7 receptor variant (P2X7-j) endogenously expressed in cervical cancer cells antagonizes the full-length P2X7 receptor through hetero-oligomerization. *J Biol Chem* **281**:17228–17237.
- Fernando SL, Saunders BM, Sluyter R, Skarratt KK, Goldberg H, Marks GB, Wiley JS, and Britton WJ (2007) A polymorphism in the P2X7 gene increases susceptibility to extrapulmonary tuberculosis. *Am J Respir Crit Care Med* **175**:360–366.
- Ferrari D, Los M, Bauer MKA, Vandenaebelle P, Wesselborg S, and Schulze-Osthoff K (1999) P2Z purinoreceptor ligation induces activation of caspases with distinct roles in apoptotic and necrotic alterations of cell death. *FEBS Lett* **447**:71–75.
- Ferrari D, Pizzirani C, Adinolfi E, Forchap S, Sitta B, Turchet L, Falzoni S, Minelli M, Baricordi R, and Di Virgilio F (2004) The antibiotic polymyxin B modulates P2X7 receptor function. *J Immunol* **173**:4652–4660.
- Ferrari D, Wesselborg S, Bauer MKA, and Schulze-Osthoff K (1997) Extracellular ATP activates transcription factor NF-kappaB through the P2Z purinoreceptor by selectively targeting NF-kappaB p65. *J Cell Biol* **139**:1635–1643.
- Fonfria E, Clay WC, Levy DS, Goodwin JA, Roman S, Smith GD, Condreay JP, and Michel AD (2008) Cloning and pharmacological characterization of the guinea pig P2X7 receptor orthologue. *Br J Pharmacol* **153**:544–556.
- Franke H, Krügel U, Grosche J, and Illes P (2003) Immunoreactivity for glial fibrillary acidic protein and P2 receptor expression on astrocytes in vivo. *Drug Dev Res* **59**:175–189.
- Franke H, Schepper C, Illes P, and Krügel U (2007) Involvement of P2X and P2Y receptors in microglial activation in vivo. *Purinergic Signal* **3**:435–445.
- Friedle SA, Curet MA, and Watters JJ (2010) Recent patents on novel P2X(7) receptor antagonists and their potential for reducing central nervous system inflammation. *Recent Patents CNS Drug Discov* **5**:35–45.
- Fulgenzi A, Dell'Antonio G, Foglieni C, Dal Cin E, Ticozzi P, Franzone JS, and Ferrero ME (2005) Inhibition of chemokine expression in rat inflamed paws by systemic use of the antihyperalgesic oxidized ATP. *BMC Immunol* **6**:18.
- Fulgenzi A, Ticozzi P, Gabel CA, Dell'Antonio G, Quattrini A, Franzone JS, and Ferrero ME (2008) Periodate oxidized ATP (oATP) reduces hyperalgesia in mice: involvement of P2X7 receptors and implications for therapy. *Int J Immunopathol Pharmacol* **21**:61–71.
- Gadeock S, Tran JN, Georgiou JG, Jalilian I, Taylor RM, Wiley JS, and Sluyter R (2010) TGF- β 1 prevents up-regulation of the P2X7 receptor by IFN- γ and LPS in leukemic THP-1 monocytes. *Biochim Biophys Acta* **1798**:2058–2066.
- Garbers C, Jänner N, Chalaris A, Moss ML, Floss DM, Meyer D, Koch-Nolte F, Rose-John S, and Scheller J (2011) Species specificity of ADAM10 and ADAM17 proteins in interleukin-6 (IL-6) trans-signaling and novel role of ADAM10 in inducible IL-6 receptor shedding. *J Biol Chem* **286**:14804–14811.
- García-Huerta P, Diaz-Hernandez M, Delicado EG, Pimentel-Santillana M, Miras-Portugal MT, and Gómez-Villafuertes R (2012) The specificity protein factor Sp1 mediates transcriptional regulation of P2X7 receptors in the nervous system. *J Biol Chem* **287**:44628–44644.
- Gargett CE, Cornish JE, and Wiley JS (1997) ATP, a partial agonist for the P2Z receptor of human lymphocytes. *Br J Pharmacol* **122**:911–917.
- Gargett CE and Wiley JS (1997) The isoquinoline derivative KN-62 a potent antagonist of the P2Z-receptor of human lymphocytes. *Br J Pharmacol* **120**:1483–1490.
- Garland A, Buckley KA, Hipskind RA, Perry MJ, Tobias JH, Buell G, Chessell I, Bowler WB, and Gallagher JA (2003) Multinucleated osteoclast formation in vivo and in vitro by P2X7 receptor-deficient mice. *Crit Rev Eukaryot Gene Expr* **13**:243–253.
- Garland A, Skarratt KK, Hocking LJ, Parsons C, Stokes L, Jørgensen NR, Fraser WD, Reid DM, Gallagher JA, and Wiley JS (2012) Polymorphisms in the P2X7 receptor gene are associated with low lumbar spine bone mineral density and accelerated bone loss in post-menopausal women. *Eur J Hum Genet* **20**:559–564.
- Gavala ML, Hill LM, Lenertz LY, Karta MR, and Bertics PJ (2010) Activation of the transcription factor FosB/activating protein-1 (AP-1) is a prominent downstream signal of the extracellular nucleotide receptor P2RX7 in monocytic and osteoblastic cells. *J Biol Chem* **285**:34288–34298.
- Gavala ML, Pfeiffer ZA, and Bertics PJ (2008) The nucleotide receptor P2RX7 mediates ATP-induced CREB activation in human and murine monocytic cells. *J Leukoc Biol* **84**:1159–1171.
- Gever JR, Cockayne DA, Dillon MP, Burnstock G, and Ford AP (2006) Pharmacology of P2X channels. *Pflugers Arch* **452**:513–537.
- Ghiringhelli F, Apetoh L, Tesniere A, Aymeric L, Ma Y, Ortiz C, Vermaelen K, Panaretakis T, Mignot G, and Ullrich E, et al. (2009) Activation of the NLRP3 inflammasome in dendritic cells induces IL-1 β -dependent adaptive immunity against tumors. *Nat Med* **15**:1170–1178.
- Gidlöf O, Smith JG, Melander O, Lövkvist H, Hedblad B, Engström G, Nilsson P, Carlsson J, Berglund G, and Olsson S, et al. (2012) A common missense variant in the ATP receptor P2X7 is associated with reduced risk of cardiovascular events. *PLoS ONE* **7**:e37491.
- Glas R, Sauter NS, Schulthess FT, Shu L, Oberholzer J, and Maedler K (2009) Purinergic P2X7 receptors regulate secretion of interleukin-1 receptor antagonist and beta cell function and survival. *Diabetologia* **52**:1579–1588.
- Gonzales EB, Kawate T, and Gouaux E (2009) Pore architecture and ion sites in acid-sensing ion channels and P2X receptors. *Nature* **460**:599–604.
- Gourine AV, Poputnikov DM, Zhernosek N, Melenchuk EV, Gerstberger R, Spyer KM, and Gourine VN (2005) P2 receptor blockade attenuates fever and cytokine responses induced by lipopolysaccharide in rats. *Br J Pharmacol* **146**:139–145.
- Gross O, Thomas CJ, Guarda G, and Tschopp J (2011) The inflammasome: an integrated view. *Immunol Rev* **243**:136–151.
- Gu B, Bendall LJ, and Wiley JS (1998) Adenosine triphosphate-induced shedding of CD23 and L-selectin (CD62L) from lymphocytes is mediated by the same receptor but different metalloproteases. *Blood* **92**:946–951.
- Gu BJ, Baird PN, Vessey KA, Skarratt KK, Fletcher EL, Fuller SJ, Richardson AJ, Guymer RH, and Wiley JS (2013) A rare functional haplotype of the P2RX4 and P2RX7 genes leads to loss of innate phagocytosis and confers increased risk of age-related macular degeneration. *FASEB J* **27**:1479–1487.
- Gu BJ, Duce JA, Valova VA, Wong B, Bush AI, Petrou S, and Wiley JS (2012) P2X7 receptor-mediated scavenger activity of mononuclear phagocytes toward non-opsonized particles and apoptotic cells is inhibited by serum glycoproteins but remains active in cerebrospinal fluid. *J Biol Chem* **287**:17318–17330.
- Gu BJ, Rathsam C, Stokes L, McGeachie AB, and Wiley JS (2009) Extracellular ATP dissociates nonmuscle myosin from P2X(7) complex: this dissociation regulates P2X(7) pore formation. *Am J Physiol Cell Physiol* **297**:C430–C439.
- Gu BJ, Saunders BM, Jurisk C, and Wiley JS (2010) The P2X7-nonmuscle myosin membrane complex regulates phagocytosis of nonopsonized particles and bacteria by a pathway attenuated by extracellular ATP. *Blood* **115**:1621–1631.
- Gu BJ, Saunders BM, Petrou S, and Wiley JS (2011) P2X(7) is a scavenger receptor for apoptotic cells in the absence of its ligand, extracellular ATP. *J Immunol* **187**:2365–2375.
- Gu BJ, Sluyter R, Skarratt KK, Shemon AN, Dao-Ung L-P, Fuller SJ, Barden JA, Clarke AL, Petrou S, and Wiley JS (2004) An Arg307 to Gln polymorphism within the ATP-binding site causes loss of function of the human P2X7 receptor. *J Biol Chem* **279**:31287–31295.
- Gu BJ and Wiley JS (2006) Rapid ATP-induced release of matrix metalloproteinase 9 is mediated by the P2X7 receptor. *Blood* **107**:4946–4953.

- Gu BJ, Zhang W, Worthington RA, Sluyter R, Dao-Ung P, Petrou S, Barden JA, and Wiley JS (2001) A Glu-496 to Ala polymorphism leads to loss of function of the human P2X7 receptor. *J Biol Chem* **276**:11135–11142.
- Gu BJ, Zhang WY, Bendall LJ, Chessell IP, Buell GN, and Wiley JS (2000) Expression of P2X₇ purinoceptors on human lymphocytes and monocytes: evidence for nonfunctional P2X₇ receptors. *Am J Physiol Cell Physiol* **279**:C1189–C1197.
- Gunosewoyo H and Kassiou M (2010) P2X purinergic receptor ligands: recently patented compounds. *Expert Opin Ther Pat* **20**:625–646.
- Haag F, Koch-Nolte F, Kühn M, Lorenzen S, and Thiele HG (1994) Premature stop codons inactivate the RT6 genes of the human and chimpanzee species. *J Mol Biol* **243**:537–546.
- Hansen RR, Nielsen CK, Nasser A, Thomsen SIM, Eghorn LF, Pham Y, Schulenburg C, Syberg S, Ding M, and Stojilkovic SS, et al. (2011) P2X7 receptor-deficient mice are susceptible to bone cancer pain. *Pain* **152**:1766–1776.
- Hassler M and Ladurner AG (2012) Towards a structural understanding of PARP1 activation and related signalling ADP-ribosyl-transferases. *Curr Opin Struct Biol* **22**:721–729.
- Hatcher JP, Chessell IP, and Hughes JP (2011) Biologics: the next-generation therapeutics for analgesia? *Expert Rev Neurother* **11**:1653–1658.
- Hattori F, Ohshima Y, Seki S, Tsukimoto M, Sato M, Takenouchi T, Suzuki A, Takai E, Kitani H, and Harada H, et al. (2012) Feasibility study of B16 melanoma therapy using oxidized ATP to target purinergic receptor P2X7. *Eur J Pharmacol* **695**:20–26.
- Hattori M and Gouaux E (2012) Molecular mechanism of ATP binding and ion channel activation in P2X receptors. *Nature* **485**:207–212.
- He WJ, Cui J, Du L, Zhao YD, Burnstock G, Zhou HD, and Ruan HZ (2012) Spinal P2X₇ receptor mediates microglia activation-induced neuropathic pain in the sciatic nerve injury rat model. *Behav Brain Res* **226**:163–170.
- He Y, Franchi L, and Núñez G (2013) TLR agonists stimulate Nlrp3-dependent IL-1 β production independently of the purinergic P2X₇ receptor in dendritic cells and in vivo. *J Immunol* **190**:334–339.
- Hempel C, Nörenberg W, Sobottka H, Urban N, Nicke A, Fischer W, and Schaefer M (2013) The phenothiazine-class antipsychotic drugs prochlorperazine and trifluoperazine are potent allosteric modulators of the human P2X₇ receptor. *Neuropharmacology* **75**:365–379.
- Hewinson J and Mackenzie AB (2007) P2X₇ receptor-mediated reactive oxygen and nitrogen species formation: from receptor to generators. *Biochem Soc Trans* **35**:1168–1170.
- Hibell AD, Thompson KM, Xing M, Humphrey PPA, and Michel AD (2001) Complexities of measuring antagonist potency at P2X₇ receptor orthologs. *J Pharmacol Exp Ther* **296**:947–957.
- Hong S, Brass A, Seman M, Haag F, Koch-Nolte F, and DUBYAK GR (2007) Lipopolysaccharide, IFN- γ , and IFN- β induce expression of the thiol-sensitive ART2.1 Ecto-ADP-ribosyltransferase in murine macrophages. *J Immunol* **179**:6215–6227.
- Hong S, Schwarz N, Brass A, Seman M, Haag F, Koch-Nolte F, Schilling WP, and DUBYAK GR (2009) Differential regulation of P2X₇ receptor activation by extracellular nicotinamide adenine dinucleotide and ecto-ADP-ribosyltransferases in murine macrophages and T cells. *J Immunol* **183**:578–592.
- Honore P, Donnelly-Roberts D, Namovic M, Zhong C, Wade C, Chandran P, Zhu C, Carroll W, Perez-Medrano A, and Iwakura Y, et al. (2009) The antihyperalgesic activity of a selective P2X₇ receptor antagonist, A-839977, is lost in IL-1 α knockout mice. *Behav Brain Res* **204**:77–81.
- Honore P, Donnelly-Roberts D, Namovic MT, Hsieh G, Zhu CZ, Mikusa JP, Hernandez G, Zhong C, Gauvin DM, and Chandran P, et al. (2006) A-740003 [N-(1-[(cyanoinimino)(5-quinolinylamino) methyl]amino-2,2-dimethylpropyl)-2-(3,4-dimethoxyphenyl) acetamide], a novel and selective P2X₇ receptor antagonist, dose-dependently reduces neuropathic pain in the rat. *J Pharmacol Exp Ther* **319**:1376–1385.
- Hoque R, Sohail M, Malik A, Sarwar S, Luo Y, Shah A, Barrat F, Flavell R, Gorelick F, and Husain S, et al. (2011) TLR9 and the NLRP3 inflammasome link acinar cell death with inflammation in acute pancreatitis. *Gastroenterology* **141**:358–369.
- Hoque R, Sohail MA, Salhanick S, Malik AF, Ghani A, Robson SC, and Mehal WZ (2012) P2X₇ receptor-mediated purinergic signaling promotes liver injury in acetaminophen hepatotoxicity in mice. *Am J Physiol Gastrointest Liver Physiol* **302**:G1171–G1179.
- Hu H, Lu W, Zhang M, Zhang X, Argall AJ, Patel S, Lee GE, Kim YC, Jacobson KA, and Laties AM, et al. (2010) Stimulation of the P2X₇ receptor kills rat retinal ganglion cells in vivo. *Exp Eye Res* **91**:425–432.
- Humphreys BD and DUBYAK GR (1998) Modulation of P2X₇ nucleotide receptor expression by pro- and anti-inflammatory stimuli in THP-1 monocytes. *J Leukoc Biol* **64**:265–273.
- Ichinari K, Kakei M, Matsuoka T, Nakashima H, and Tanaka H (1996) Direct activation of the ATP-sensitive potassium channel by oxygen free radicals in guinea-pig ventricular cells: its potentiation by MgADP. *J Mol Cell Cardiol* **28**:1867–1877.
- Ishii K, Kaneda M, Li H, Rockland KS, and Hashikawa T (2003) Neuron-specific distribution of P2X₇ purinergic receptors in the monkey retina. *J Comp Neurol* **459**:267–277.
- Ito G, Suekawa Y, Watanabe M, Takahashi K, Inubushi T, Murasaki K, Hirose N, Hiyama S, Uchida T, and Tanne K (2013) P2X₇ receptor in the trigeminal sensory nuclear complex contributes to tactile allodynia/hyperalgesia following trigeminal nerve injury. *Eur J Pain* **17**:185–199.
- Itoh K, Chiang CY, Li Z, Lee JC, Dostrovsky JO, and Sessle BJ (2011) Central sensitization of nociceptive neurons in rat medullary dorsal horn involves purinergic P2X₇ receptors. *Neuroscience* **192**:721–731.
- Iwamura Y, Takenouchi T, Murayama Y, Okada H, Imamura M, Shimizu Y, Hashimoto M, Mohri S, Yokoyama T, and Kitani H (2012) Anti-prion activity of brilliant blue G. *PLoS ONE* **7**:e37896.
- Jalilian I, Peranec M, Curtis BL, Seavers A, Spildrejorde M, Sluyter R, and Sluyter R (2012a) Activation of the damage-associated molecular pattern receptor P2X₇ induces interleukin-1 β release from canine monocytes. *Vet Immunol Immunopathol* **149**:86–91.
- Jalilian I, Spildrejorde M, Seavers A, Curtis BL, McArthur JD, and Sluyter R (2012b) Functional expression of the damage-associated molecular pattern receptor P2X₇ on canine kidney epithelial cells. *Vet Immunol Immunopathol* **150**:228–233.
- Jamieson SE, Peixoto-Rangel AL, Hargrave AC, Roubaix LAD, Mui EJ, Boulter NR, Miller EN, Fuller SJ, Wiley JS, and Castellucci L, et al. (2010) Evidence for associations between the purinergic receptor P2X₇ (P2RX7) and toxoplasmosis. *Genes Immun* **11**:374–383.
- Jelassi B, Anchelín M, Chamouton J, Cayuela ML, Clarysse L, Li J, Goré J, Jiang LH, and Roger S (2013) Anthraquinone emodin inhibits human cancer cell invasiveness by antagonizing P2X₇ receptors. *Carcinogenesis* **34**:1487–1496.
- Ji X, Naito Y, Hirokawa G, Weng H, Hiura Y, Takahashi R, and Iwai N (2012) P2X₇ receptor antagonism attenuates the hypertension and renal injury in Dahl salt-sensitive rats. *Hypertens Res* **35**:173–179.
- Jiang LH (2012) P2X receptor-mediated ATP purinergic signaling in health and disease. *Cell Health Cytoskeleton* **4**:83–101.
- Jiang LH (2009) Inhibition of P2X₇ receptors by divalent cations: old action and new insight. *Eur Biophys J* **38**:339–346.
- Jiang LH, Baldwin JM, Roger S, and Baldwin SA (2013) Insights into the molecular mechanisms underlying mammalian P2X₇ receptor functions and contributions in diseases, revealed by structural modeling and single nucleotide polymorphisms. *Front Pharmacol* **4**:55.
- Jiang LH, Mackenzie AB, North RA, and Surprenant A (2000) Brilliant blue G selectively blocks ATP-gated rat P2X₇ receptors. *Mol Pharmacol* **58**:82–88.
- Jimenez-Pacheco A, Mesuret G, Sanz-Rodriguez A, Tanaka K, Mooney C, Conroy R, Miras-Portugal MT, Diaz-Hernandez M, Henshall DC, and Engel T (2013) Increased neocortical expression of the P2X₇ receptor after status epilepticus and anticonvulsant effect of P2X₇ receptor antagonist A-438079. *Epilepsia* **54**:1551–1561.
- Jørgensen NR, Husted LB, Skarratt KK, Stokes L, Tofteng CL, Kvist T, Jensen J-EB, Eiken P, Brixen K, and Fuller S, et al. (2012) Single-nucleotide polymorphisms in the P2X₇ receptor gene are associated with post-menopausal bone loss and vertebral fractures. *Eur J Hum Genet* **20**:675–681.
- Kakurai K, Sugiyama T, Kurimoto T, Oku H, and Ikeda T (2013) Involvement of P2X₇ receptors in retinal ganglion cell death after optic nerve crush injury in rats. *Neurosci Lett* **534**:237–241.
- Kamei J, Takahashi Y, Yoshikawa Y, and Saitoh A (2005) Involvement of P2X₇ receptor subtypes in ATP-induced enhancement of the cough reflex sensitivity. *Eur J Pharmacol* **528**:158–161.
- Kanaituka T, Bertoll R, Stevens LA, Moss J, Sardinha D, Rajan TV, Zipris D, Mordes JP, Greiner DL, and Rossini AA (1997) Expression in BALB/c and C57BL/6 mice of R6-1 and R6-2 ADP-ribosyltransferases that differ in enzymatic activity: C57BL/6 R6-1 is a natural transferase knockout. *J Immunol* **159**:2741–2749.
- Kang SS, Keasey MP, and Hagg T (2013) P2X₇ receptor inhibition increases CNTF in the subventricular zone, but not neurogenesis or neuroprotection after stroke in adult mice. *Transl Stroke Res* **4**:533–545.
- Kawate T, Michel JC, Birdsong WT, and Gouaux E (2009) Crystal structure of the ATP-gated P2X₇ ion channel in the closed state. *Nature* **460**:592–598.
- Ke HZ, Qi H, Weidema AF, Zhang Q, Panupinthu N, Crawford DT, Grasser WA, Paralkar VM, Li M, and Audoly LP, et al. (2003) Deletion of the P2X₇ nucleotide receptor reveals its regulatory roles in bone formation and resorption. *Mol Endocrinol* **17**:1356–1367.
- Kerr D and Krantis A (1979) A new class of ATP antagonist. *Proc Aust Physiol Pharmacol Soc* **10**:156P.
- Kerur N, Hirano Y, Tarallo V, Fowler BJ, Bastos-Carvalho A, Yasuma T, Yasuma R, Kim Y, Hinton DR, and Kirschning CJ, et al. (2013) TLR-independent and P2X₇-dependent signaling mediate Alu RNA-induced NLRP3 inflammasome activation in geographic atrophy. *Invest Ophthalmol Vis Sci* **54**:7395–7401.
- Keystone EC, Wang MM, Layton M, Hollis S, and McInnes IB; D1520C00001 Study Team (2012) Clinical evaluation of the efficacy of the P2X₇ purinergic receptor antagonist AZD9056 on the signs and symptoms of rheumatoid arthritis in patients with active disease despite treatment with methotrexate or sulphasalazine. *Ann Rheum Dis* **71**:1630–1635.
- Khadra A, Tomić M, Yan Z, Zemkova H, Sherman A, and Stojilkovic SS (2013) Dual gating mechanism and function of P2X₇ receptor channels. *Biophys J* **104**:2612–2621.
- Kim JE and Kang TC (2011) The P2X₇ receptor-pannexin-1 complex decreases muscarinic acetylcholine receptor-mediated seizure susceptibility in mice. *J Clin Invest* **121**:2037–2047.
- Kim JE, Ryu HJ, and Kang TC (2011) P2X₇ receptor activation ameliorates CA3 neuronal damage via a tumor necrosis factor- α -mediated pathway in the rat hippocampus following status epilepticus. *J Neuroinflammation* **8**:62.
- Kim WI, Ryu HJ, Kim JE, Seo CH, Lee BC, Choi IG, and Kang TC (2013) Differential nuclear factor-kappa B phosphorylation induced by lipopolysaccharide in the hippocampus of P2X₇ receptor knockout mouse. *Neurosci Res* **35**:369–381.
- Kim JE, Ryu HJ, Yeo SI, and Kang TC (2010) P2X₇ receptor regulates leukocyte infiltrations in rat frontoparietal cortex following status epilepticus. *J Neuroinflammation* **7**:65.
- Koch-Nolte F, Duffy T, Nissen M, Kahl S, Killeen N, Ablamunits V, Haag F, and Leiter EH (1999) A new monoclonal antibody detects a developmentally regulated mouse ecto-ADP-ribosyltransferase on T cells: subset distribution, inbred strain variation, and modulation upon T cell activation. *J Immunol* **163**:6014–6022.
- Koch-Nolte F, Reyelt J, Schössow B, Schwarz N, Scheuplein F, Rothenburg S, Haag F, Alzogaray V, Cauerhff A, and Goldbaum FA (2007) Single domain antibodies from llama effectively and specifically block T cell ecto-ADP-ribosyltransferase ART2.2 in vivo. *FASEB J* **21**:3490–3498.
- Kong F, Liu S, Xu C, Liu J, Li G, Li G, Gao Y, Lin H, Tu G, and Peng H, et al. (2013) Electrophysiological studies of upregulated P2X₇ receptors in rat superior cervical ganglia after myocardial ischemic injury. *Neurochem Int* **63**:230–237.
- Korocok J, Raimundo LN, Ke HZ, Sims SM, and Dixon SJ (2004) Extracellular nucleotides act through P2X₇ receptors to activate NF-kappaB in osteoclasts. *J Bone Miner Res* **19**:642–651.

- Kurashima Y, Amiya T, Nochi T, Fujisawa K, Haraguchi T, Iba H, Tsutsui H, Sato S, Nakajima S, and Iijima H, et al. (2012) Extracellular ATP mediates mast cell-dependent intestinal inflammation through P2X7 purinoceptors. *Nat Commun* **3**: 1034.
- Labasi JM, Petrushova N, Donovan C, McCurdy S, Lira P, Payette MM, Brissette W, Wicks JR, Audoly L, and Gabel CA (2002) Absence of the P2X7 receptor alters leukocyte function and attenuates an inflammatory response. *J Immunol* **168**: 6436–6445.
- Laeremans T, Stortelers C, Nolte F, Gonzalez M, Assuncao J, and Van Rompaey P (2010) inventors, Ablynx, assignee. Genetic immunization for producing immunoglobulins against cell-associated antigens such as P2X7, CXCR7 or CXCR4, Ablynx N.V. Patent WO2010070145. 2010 Dec 2.
- Lambrecht G, Friebe T, Grimm U, Windscheif U, Bungardt E, Hildebrandt C, Baumert HG, Spatz-Kümbel G, and Mutschler E (1992) PPADS, a novel functionally selective antagonist of P2 purinoceptor-mediated responses. *Eur J Pharmacol* **217**:217–219.
- Lang PA, Merkle D, Funkner P, Shaabani N, Meryk A, Krings C, Barthuber C, Recher M, Brück W, and Häussinger D, et al. (2010) Oxidized ATP inhibits T-cell-mediated autoimmunity. *Eur J Immunol* **40**:2401–2408.
- Le Feuvre RA, Brough D, Touzani O, and Rothwell NJ (2003) Role of P2X7 receptors in ischemic and excitotoxic brain injury in vivo. *J Cereb Blood Flow Metab* **23**: 381–384.
- Le Stunff H, Auger R, Kanellopoulos J, and Raymond MN (2004) The Pro-451 to Leu polymorphism within the C-terminal tail of P2X7 receptor impairs cell death but not phospholipase D activation in murine thymocytes. *J Biol Chem* **279**: 16918–16926.
- Lee GE, Joshi BV, Chen W, Jeong LS, Moon HR, Jacobson KA, and Kim YC (2008a) Synthesis and structure-activity relationship studies of tyrosine-based antagonists at the human P2X7 receptor. *Bioorg Med Chem Lett* **18**:571–575.
- Lee M, Lee SJ, Choi HJ, Jung YW, Frøkiær J, Nielsen S, and Kwon TH (2008b) Regulation of AQP4 protein expression in rat brain astrocytes: role of P2X7 receptor activation. *Brain Res* **1195**:1–11.
- Lemaire I and Leduc N (2003) Purinergic P2X7 receptor function in lung alveolar macrophages: Pharmacologic characterization and bidirectional regulation by Th1 and Th2 cytokines. *Drug Dev Res* **59**:118–127.
- Lenertz LY, Gavala ML, Hill LM, and Bertics PJ (2009) Cell signaling via the P2X7(7) nucleotide receptor: linkage to ROS production, gene transcription, and receptor trafficking. *Purinergic Signal* **5**:175–187.
- Lenertz LY, Gavala ML, Zhu Y, and Bertics PJ (2011) Transcriptional control mechanisms associated with the nucleotide receptor P2X7, a critical regulator of immunologic, osteogenic, and neurologic functions. *Immunol Res* **50**:22–38.
- Lester S, Stokes L, Skarratt KK, Gu BJ, Sivils KL, Lessard CJ, Wiley JS, and Rischmueller M (2013) Epistasis with HLA DR3 implicates the P2X7 receptor in the pathogenesis of primary Sjögren's syndrome. *Arthritis Res Ther* **15**:R71.
- Letavic MA, Lord B, Bischoff F, Hawryluk NA, Pieters S, Rech JC, Sales Z, Velter AI, Ao H, and Bonaventure P, et al. (2013) Synthesis and pharmacological characterization of two novel, brain penetrating P2X7 antagonists. *ACS Med Chem Lett* **4**: 419–422.
- Li CM, Campbell SJ, Kumararatne DS, Hill AV, and Lammas DA (2002) Response heterogeneity of human macrophages to ATP is associated with P2X7 receptor expression but not to polymorphisms in the P2RX7 promoter. *FEBS Lett* **531**: 127–131.
- Li M, Chang TH, Silberberg SD, and Swartz KJ (2008) Gating the pore of P2X receptor channels. *Nat Neurosci* **11**:883–887.
- Li M, Silberberg SD, and Swartz KJ (2013) Subtype-specific control of P2X receptor channel signaling by ATP and Mg²⁺. *Proc Natl Acad Sci USA* **110**:E3455–E3463.
- Liao SD and Puro DG (2006) NAD⁺-induced vasotoxicity in the pericyte-containing microvasculature of the rat retina: effect of diabetes. *Invest Ophthalmol Vis Sci* **47**: 5032–5038.
- Lin C, Ren S, Zhang L, Jin H, Sun J, and Zuo Y (2012) Extracellular ATP induces CD44 shedding from macrophage-like P388D1 cells via the P2X7 receptor. *Hematol Oncol* **30**:70–75.
- Liu J, Li G, Peng H, Tu G, Kong F, Liu S, Gao Y, Xu H, Qiu S, and Fan B, et al. (2013) Sensory-sympathetic coupling in superior cervical ganglia after myocardial ischemic injury facilitates sympathoexcitatory action via P2X7 receptor. *Purinergic Signal* **9**:463–479.
- Liu L, Zou J, Liu X, Jiang LH, and Li J (2010) Inhibition of ATP-induced macrophage death by emodin via antagonizing P2X7 receptor. *Eur J Pharmacol* **640**:15–19.
- Liu X, Surprenant A, Mao HJ, Roger S, Xia R, Bradley H, and Jiang LH (2008) Identification of key residues coordinating functional inhibition of P2X7 receptors by zinc and copper. *Mol Pharmacol* **73**:252–259.
- Lopez-Castejon G, Theaker J, Pelegrin P, Clifton AD, Braddock M, and Surprenant A (2010) P2X7(7) receptor-mediated release of cathepsins from macrophages is a cytokine-independent mechanism potentially involved in joint diseases. *J Immunol* **185**:2611–2619.
- Lu YM, Tao RR, Huang JY, Li LT, Liao MH, Li XM, Fukunaga K, Hong ZH, and Han F (2012) P2X7 signaling promotes microsphere embolism-triggered microglia activation by maintaining elevation of Fas ligand. *J Neuroinflammation* **9**:172.
- Lucae S, Salyakina D, Barden N, Harvey M, Gagné B, Labbé M, Binder EB, Uhr M, Paez-Pereda M, and Sillaber I, et al. (2006) P2RX7, a gene coding for a purinergic ligand-gated ion channel, is associated with major depressive disorder. *Hum Mol Genet* **15**:2438–2445.
- Lucattelli M, Cicko S, Müller T, Lommatzsch M, De Cunto G, Cardini S, Sundas W, Grimm M, Zeiser R, and Dürk T, et al. (2011) P2X7 receptor signaling in the pathogenesis of smoke-induced lung inflammation and emphysema. *Am J Respir Cell Mol Biol* **44**:423–429.
- Ludden PW (1994) Reversible ADP-ribosylation as a mechanism of enzyme regulation in procarboxylates. *Mol Cell Biochem* **138**:123–129.
- Marcellino D, Suárez-Boomgaard D, Sánchez-Reina MD, Aguirre JA, Yoshitake T, Yoshitake S, Hagman B, Kehr J, Agnati LF, and Fuxe K, et al. (2010) On the role of P2X(7) receptors in dopamine nerve cell degeneration in a rat model of Parkinson's disease: studies with the P2X(7) receptor antagonist A-438079. *J Neural Transm* **117**:681–687.
- Marcillo A, Frydel B, Bramlett HM, and Dietrich WD (2012) A reassessment of P2X7 receptor inhibition as a neuroprotective strategy in rat models of contusion injury. *Exp Neurol* **233**:687–692.
- Marcoli M, Cervetto C, Paluzzi P, Guarnieri S, Alloisio S, Thellung S, Nobile M, and Maura G (2008) P2X7 pre-synaptic receptors in adult rat cerebrocortical nerve terminals: a role in ATP-induced glutamate release. *J Neurochem* **105**:2330–2342.
- Marques-da-Silva C, Chaves MM, Castro NG, Coutinho-Silva R, and Guimaraes MZ (2011) Colchicine inhibits cationic dye uptake induced by ATP in P2X2 and P2X7 receptor-expressing cells: implications for its therapeutic action. *Br J Pharmacol* **163**:912–926.
- Martin U, Scholler J, Gurgel J, Renshaw B, Sims JE, and Gabel CA (2009) Externalization of the leaderless cytokine IL-1F6 occurs in response to lipopolysaccharide/ATP activation of transduced bone marrow macrophages. *J Immunol* **183**:4021–4030.
- Martins JP, Silva RBM, Coutinho-Silva R, Takiya CM, Battastini AMO, Morrone FB, and Campos MM (2012) The role of P2X7 purinergic receptors in inflammatory and nociceptive changes accompanying cyclophosphamide-induced haemorrhagic cystitis in mice. *Br J Pharmacol* **165**:183–196.
- Masin M, Young C, Lim K, Barnes SJ, Xu XJ, Marshall V, Brutkowski W, Mooney ER, Gorecki DC, and Murrell-Lagnado R (2012) Expression, assembly and function of novel C-terminal truncated variants of the mouse P2X7 receptor: re-evaluation of P2X7 knockouts. *Br J Pharmacol* **165**:978–993.
- Matute C, Torre I, Pérez-Cerdá F, Pérez-Samartín A, Alberdi E, Etxebarria E, Arranz AM, Ravid R, Rodríguez-Antigüedad A, and Sánchez-Gómez M, et al. (2007) P2X(7) receptor blockade prevents ATP excitotoxicity in oligodendrocytes and ameliorates experimental autoimmune encephalomyelitis. *J Neurosci* **27**:9525–9533.
- McGaraughty S, Chu KL, Namovic MT, Donnelly-Roberts DL, Harris RR, Zhang XF, Shieh CC, Wismer CT, Zhu CZ, and Gauvin DM, et al. (2007) P2X7-related modulation of pathological nociception in rats. *Neuroscience* **146**:1817–1828.
- McHugh SM, Roman S, Davis B, Koch A, Pickett AM, Richardson JC, Miller SR, Wetten S, Cox CJ, and Karpe F, et al. (2012) Effects of genetic variation in the P2RX7 gene on pharmacodynamics of a P2X(7) receptor antagonist: a prospective genotyping approach. *Br J Clin Pharmacol* **74**:376–380.
- Melani A, Amadio S, Gianfriddo M, Vannucchi MG, Volonté C, Bernardi G, Pedata F, and Sancesario G (2006) P2X7 receptor modulation on microglial cells and reduction of brain infarct caused by middle cerebral artery occlusion in rat. *J Cereb Blood Flow Metab* **26**:974–982.
- Menzies RI, Unwin RJ, Dash RK, Beard DA, Cowley AW Jr, Carlson BE, Mullins JJ, and Bailey MA (2013) Effect of P2X4 and P2X7 receptor antagonism on the pressure diuresis relationship in rats. *Front Physiol* **4**:305.
- Michel AD, Chambers LJ, Clay WC, Condreay JP, Walter DS, and Chessell IP (2007) Direct labelling of the human P2X7 receptor and identification of positive and negative cooperativity of binding. *Br J Pharmacol* **151**:103–114.
- Michel AD, Clay WC, Ng SW, Roman S, Thompson K, Condreay JP, Hall M, Holbrook J, Livermore D, and Senger S (2008) Identification of regions of the P2X(7) receptor that contribute to human and rat species differences in antagonist effects. *Br J Pharmacol* **155**:738–751.
- Michel AD and Fonfria E (2007) Agonist potency at P2X7 receptors is modulated by structurally diverse lipids. *Br J Pharmacol* **152**:523–537.
- Michel AD, Ng SW, Roman S, Clay WC, Dean DK, and Walter DS (2009) Mechanism of action of species-selective P2X(7) receptor antagonists. *Br J Pharmacol* **156**: 1312–1325.
- Michel AD, Thompson KM, Simon J, Boyfield I, Fonfria E, and Humphrey PP (2006) Species and response dependent differences in the effects of MAPK inhibitors on P2X(7) receptor function. *Br J Pharmacol* **149**:948–957.
- Miller CM, Boulter NR, Fuller SJ, Zakrzewski AM, Lees MP, Saunders BM, Wiley JS, and Smith NC (2011) The role of the P2X₇ receptor in infectious diseases. *PLoS Pathog* **7**:e1002212.
- Mingam R, De Smedt V, Amédée T, Bluthé RM, Kelley KW, Dantzer R, and Layé S (2008) In vitro and in vivo evidence for a role of the P2X7 receptor in the release of IL-1 β in the murine brain. *Brain Behav Immun* **22**:234–244.
- Monif M, Reid CA, Powell KL, Smart ML, and Williams DA (2009) The P2X7 receptor drives microglial activation and proliferation: a trophic role for P2X7R pore. *J Neurosci* **29**:3781–3791.
- Moon H, Na H-Y, Chong KH, and Kim TJ (2006) P2X7 receptor-dependent ATP-induced shedding of CD27 in mouse lymphocytes. *Immunol Lett* **102**:98–105.
- Moore SF and Mackenzie AB (2008) Species and agonist dependent zinc modulation of endogenous and recombinant ATP-gated P2X7 receptors. *Biochem Pharmacol* **76**:1740–1747.
- Moore SF and MacKenzie AB (2009) NADPH oxidase NOX2 mediates rapid cellular oxidation following ATP stimulation of endotoxin-primed macrophages. *J Immunol* **183**:3302–3308.
- Mueller-Diekmann C, Scheuermann T, Wursthorn K, Schröder J, Haag F, Schulz GE, and Koch-Nolte F (2002) Expression, purification, crystallization and preliminary X-ray analysis of rat ecto-ADP-ribosyltransferase 2 (ART2.2). *Acta Crystallogr D Biol Crystallogr* **58**:1211–1213.
- Müller T, Vieira RP, Grimm M, Dürk T, Cicko S, Zeiser R, Jakob T, Martin SF, Blumenthal B, and Sorichter S, et al. (2011) A potential role for P2X7R in allergic airway inflammation in mice and humans. *Am J Respir Cell Mol Biol* **44**: 456–464.
- Muñoz-Planillo R, Kuffa P, Martínez-Colón G, Smith BL, Rajendiran TM, and Núñez G (2013) K⁺ efflux is the common trigger of NLRP3 inflammasome activation by bacterial toxins and particulate matter. *Immunity* **38**:1142–1153.
- Murgia M, Hanau S, Pizzo P, Rippla M, and Di Virgilio F (1993) Oxidized ATP. An irreversible inhibitor of the macrophage purinergic P2Z receptor. *J Biol Chem* **268**: 8199–8203.
- Muzzachi S, Blasi A, Ciani E, Favia M, Cardone RA, Marzulli D, Reshkin SJ, Merizzi G, Casavola V, and Soletti A, et al. (2013) MEDI1101: a new dialdehyde compound

- regulating P2 \times 7 receptor cell surface expression in U937 cells. *Biol Cell* **105**: 399–413.
- Nakahira K, Haspel JA, Rathinam VA, Lee SJ, Dolinay T, Lam HC, Englert JA, Rabinovitch M, Cernadas M, and Kim HP, et al. (2011) Autophagy proteins regulate innate immune responses by inhibiting the release of mitochondrial DNA mediated by the NALP3 inflammasome. *Nat Immunol* **12**:222–230.
- Nakanishi M, Mori T, Nishikawa K, Sawada M, Kuno M, and Asada A (2007) The effects of general anesthetics on P2X7 and P2Y receptors in a rat microglial cell line. *Anesth Analg* **104**:1136–1144.
- Naviaux RK, Zolkipli Z, Wang L, Nakayama T, Naviaux JC, Le TP, Schuchbauer MA, Rogac M, Tang Q, and Dugan LL, et al. (2013) Antipurinergic therapy corrects the autism-like features in the poly(IC) mouse model. *PLoS ONE* **8**:e57380.
- Nelson DW, Gregg RJ, Kort ME, Perez-Medrano A, Voight EA, Wang Y, Grayson G, Namovic MT, Donnelly-Roberts DL, and Niforatos W, et al. (2006) Structure-activity relationship studies on a series of novel, substituted 1-benzyl-5-phenyltetrazole P2X7 antagonists. *J Med Chem* **49**:3659–3666.
- Nicke A (2008) Homotrimeric complexes are the dominant assembly state of native P2X7 subunits. *Biochem Biophys Res Commun* **377**:803–808.
- Nicke A, Bäumert HG, Rettinger J, Eichele A, Lambrecht G, Mutschler E, and Schmalzing G (1998) P2X1 and P2X3 receptors form stable trimers: a novel structural motif of ligand-gated ion channels. *EMBO J* **17**:3016–3028.
- Nicke A, Kuan YH, Masin M, Rettinger J, Marquez-Klaka B, Bender O, Górecki DC, Murrell-Lagnado RD, and Soto F (2009) A functional P2X7 splice variant with an alternative transmembrane domain 1 escapes gene inactivation in P2X7 knock-out mice. *J Biol Chem* **284**:25813–25822.
- Nörenberg W, Hempel C, Urban N, Sobottka H, Illes P, and Schaefer M (2011) Clemastine potentiates the human P2X7 receptor by sensitizing it to lower ATP concentrations. *J Biol Chem* **286**:11067–11081.
- Nörenberg W, Sobottka H, Hempel C, Plötz T, Fischer W, Schmalzing G, and Schaefer M (2012) Positive allosteric modulation by ivermectin of human but not murine P2X7 receptors. *Br J Pharmacol* **167**:48–66.
- Notomi S, Hisatomi T, Murakami Y, Terasaki H, Sonoda S, Asato R, Takeda A, Ikeda Y, Enaida H, and Sakamoto T, et al. (2013) Dynamic increase in extracellular ATP accelerates photoreceptor cell apoptosis via ligation of P2RX7 in subretinal hemorrhage. *PLoS ONE* **8**:e53338.
- Ohshima Y, Tsukimoto M, Takenouchi T, Harada H, Suzuki A, Sato M, Kitani H, and Kojima S (2010) γ -Irradiation induces P2X(7) receptor-dependent ATP release from B16 melanoma cells. *Biochim Biophys Acta* **1800**:40–46.
- Okamoto S, Azhpa O, Yu Y, Russo E, and Dennert G (1998) Expression of ADP-ribosyltransferase on normal T lymphocytes and effects of nicotinamide adenine dinucleotide on their function. *J Immunol* **160**:4190–4198.
- Panena W, Jijon H, Herx LM, Armstrong JN, Feighan D, Wei T, Yong VW, Ransohoff RM, and MacVicar BA (2001) P2X7-like receptor activation in astrocytes increases chemokine monocyte chemoattractant protein-1 expression via mitogen-activated protein kinase. *J Neurosci* **21**:7135–7142.
- Pardon E, Laeremans T, Triest S, Rasmussen SG, Wohlkönig A, Ruf A, Muylderma S, Hol WG, Kobilka BK, and Steyaert J (2014) A general protocol for the generation of Nanobodies for structural biology. *Nat Protoc* **9**:674–693.
- Pelegrin P, Barroso-Gutierrez C, and Surprenant A (2008) P2X7 receptor differentially couples to distinct release pathways for IL-1 β in mouse macrophage. *J Immunol* **180**: 7147–7157.
- Pelegrin P and Surprenant A (2006) Pannexin-1 mediates large pore formation and interleukin-1 β release by the ATP-gated P2X7 receptor. *EMBO J* **25**: 5071–5082.
- Pellegatti P, Falzoni S, Donvito G, Lemaire I, and Di Virgilio F (2011) P2X7 receptor drives osteoclast fusion by increasing the extracellular adenosine concentration. *FASEB J* **25**:1264–1274.
- Pellegatti P, Falzoni S, Pinton P, Rizzuto R, and Di Virgilio F (2005) A novel recombinant plasma membrane-targeted luciferase reveals a new pathway for ATP secretion. *Mol Biol Cell* **16**:3659–3665.
- Peng W, Cotrina ML, Han X, Yu H, Bekar L, Blum L, Takano T, Tian GF, Goldman SA, and Nedergaard M (2009) Systemic administration of an antagonist of the ATP-sensitive receptor P2X7 improves recovery after spinal cord injury. *Proc Natl Acad Sci USA* **106**:12489–12493.
- Perez-Medrano A, Donnelly-Roberts DL, Honore P, Hsieh GC, Namovic MT, Peddi S, Shuai Q, Wang Y, Faltynek CR, and Jarvis MF, et al. (2009) Discovery and biological evaluation of novel cyanoguanidine P2X(7) antagonists with analgesic activity in a rat model of neuropathic pain. *J Med Chem* **52**:3366–3376.
- Portales-Cervantes L, Niño-Moreno P, Doniz-Padilla L, Baranda-Candido L, Garcia-Hernández M, Salgado-Bustamante M, González-Amaro R, and Portales-Pérez D (2010) Expression and function of the P2X(7) purinergic receptor in patients with systemic lupus erythematosus and rheumatoid arthritis. *Hum Immunol* **71**: 818–825.
- Pupovac A, Foster CM, and Slyuter R (2013a) Human P2X7 receptor activation induces the rapid shedding of CXCL16. *Biochem Biophys Res Commun* **432**: 626–631.
- Pupovac A, Stokes L, and Slyuter R (2013b) CAY10593 inhibits the human P2X7 receptor independently of phospholipase D1 stimulation. *Purinergic Signal* **9**: 609–619.
- Puthussery T and Fletcher E (2009) Extracellular ATP induces retinal photoreceptor apoptosis through activation of purinoceptors in rodents. *J Comp Neurol* **513**: 430–440.
- Qiu F and Dahl G (2009) A permeant regulating its permeation pore: inhibition of pannexin 1 channels by ATP. *Am J Physiol Cell Physiol* **296**:C250–C255.
- Qu Y and DUBYAK GR (2009) P2X7 receptors regulate multiple types of membrane trafficking responses and non-classical secretion pathways. *Purinergic Signal* **5**: 163–173.
- Qu Y, Misaghi S, Newton K, Gilmour LL, Louie S, Cupp JE, DUBYAK GR, Hackos D, and Dixit VM (2011) Pannexin-1 is required for ATP release during apoptosis but not for inflammasome activation. *J Immunol* **186**:6553–6561.
- Rahman OA, Sasvari-Szekely M, Szekely A, Faludi G, Guttman A, and Nemoda Z (2010) Analysis of a polymorphic microRNA target site in the purinergic receptor P2RX7 gene. *Electrophoresis* **31**:1790–1795.
- Rassendren F, Buell GN, Virginio C, Collo G, North RA, and Surprenant A (1997) The permeabilizing ATP receptor, P2X7. Cloning and expression of a human cDNA. *J Biol Chem* **272**:5482–5486.
- Riedel T, Lozinsky I, Schmalzing G, and Markwardt F (2007a) Kinetics of P2X7 receptor-operated single channels currents. *Biophys J* **92**:2377–2391.
- Riedel T, Schmalzing G, and Markwardt F (2007b) Influence of extracellular monovalent cations on pore and gating properties of P2X7 receptor-operated single-channel currents. *Biophys J* **93**:846–858.
- Riteau N, Gasse P, Fauconnier L, Gombault A, Couegnat M, Fick L, Kanellopoulos J, Quesniaux VFJ, Marchand-Adam S, and Crestani B, et al. (2010) Extracellular ATP is a danger signal activating P2X7 receptor in lung inflammation and fibrosis. *Am J Respir Crit Care Med* **182**:774–783.
- Rodriguez-Feo CL, Sexton KW, Boyer RB, Pollins AC, Cardwell NL, Nanney LB, Shack RB, Mikesch MA, McGill CH, and Driscoll CW, et al. (2013) Blocking the P2X7 receptor improves outcomes after axonal fusion. *J Surg Res* **184**:705–713.
- Roger S, Gillet L, Baroja-Mazo A, Surprenant A, and Pelegrin P (2010a) C-terminal calmodulin-binding motif differentially controls human and rat P2X7 receptor current facilitation. *J Biol Chem* **285**:17514–17524.
- Roger S, Mei ZZ, Baldwin JM, Dong L, Bradley H, Baldwin SA, Surprenant A, and Jiang LH (2010b) Single nucleotide polymorphisms that were identified in affective mood disorders affect ATP-activated P2X7 receptor functions. *J Psychiatr Res* **44**:347–355.
- Roger S, Pelegrin P, and Surprenant A (2008) Facilitation of P2X7 receptor currents and membrane blebbing via constitutive and dynamic calmodulin binding. *J Neurosci* **28**:6393–6401.
- Rokic MB and Stojilkovic SS (2013) Two open states of P2X receptor channels. *Front Cell Neurosci* **7**:215.
- Romagnoli R, Baraldi PG, Cruz-Lopez O, Lopez-Cara C, Preti D, Borea PA, and Gessi S (2008) The P2X7 receptor as a therapeutic target. *Expert Opin Ther Targets* **12**: 647–661.
- Roman S, Cusdin FS, Fonfria E, Goodwin JA, Reeves J, Lappin SC, Chambers L, Walter DS, Clay WC, and Michel AD (2009) Cloning and pharmacological characterization of the dog P2X7 receptor. *Br J Pharmacol* **158**:1513–1526.
- Ryu JK, Jantarotnait N, Serrano-Perez MC, McGeer PL, and McLarnon JG (2011) Block of purinergic P2X7R inhibits tumor growth in a C6 glioma brain tumor animal model. *J Neuropathol Exp Neurol* **70**:13–22.
- Ryu JK and McLarnon JG (2008) Block of purinergic P2X(7) receptor is neuroprotective in an animal model of Alzheimer's disease. *Neuroreport* **19**: 1715–1719.
- Sánchez-Nogueiro J, Marín-García P, and Miras-Portugal MT (2005) Characterization of a functional P2X(7)-like receptor in cerebellar granule neurons from P2X(7) knockout mice. *FEBS Lett* **579**:3783–3788.
- Sanz JM, Chiozzi P, Ferrari D, Colaianna M, Idzko M, Falzoni S, Fellin R, Trabace L, and Di Virgilio F (2009) Activation of microglia by amyloid β requires P2X7 receptor expression. *J Immunol* **182**:4378–4385.
- Savigni DL, O'Hare Doig RL, Szymanski CR, Bartlett CA, Lozić I, Smith NM, and Fitzgerald M (2013) Three Ca²⁺ channel inhibitors in combination limit chronic secondary degeneration following neurotrauma. *Neuropharmacology* **75**: 380–390.
- Sawynok J and Reid A (1997) Peripheral adenosine 5'-triphosphate enhances nociception in the formalin test via activation of a purinergic p2X receptor. *Eur J Pharmacol* **330**:115–121.
- Scheuplein F, Schwarz N, Adriouch S, Krebs C, Bannas P, Rissiek B, Seman M, Haag F, and Koch-Nolte F (2009) NAD⁺ and ATP released from injured cells induce P2X7-dependent shedding of CD62L and externalization of phosphatidylserine by murine T cells. *J Immunol* **182**:2898–2908.
- Schilling WP, Sinks WG, and Estacion M (1999) Maitotoxin activates a non-selective cation channel and a P2Z/P2X(7)-like cytolytic pore in human skin fibroblasts. *Am J Physiol* **277**:C755–C765.
- Schwarz N, Drouot L, Nicke A, Fliegerer R, Boyer O, Guse AH, Haag F, Adriouch S, and Koch-Nolte F (2012) Alternative splicing of the N-terminal cytosolic and transmembrane domains of P2X7 controls gating of the ion channel by ADP-ribosylation. *PLoS ONE* **7**:e41269.
- Seil M, Fontanils U, Etxebarria IG, Pochet S, Garcia-Marcos M, Marino A, and Dehaye J-P (2008) Pharmacological evidence for the stimulation of NADPH oxidase by P2X(7) receptors in mouse submandibular glands. *Purinergic Signal* **4**: 347–355.
- Seil M, Kabré E, Nagant C, Vandenbranden M, Fontanils U, Marino A, Pochet S, and Dehaye J-P (2010) Regulation by CRAMP of the responses of murine peritoneal macrophages to extracellular ATP. *Biochim Biophys Acta* **1798**:569–578.
- Seman M, Adriouch S, Scheuplein F, Krebs C, Freese D, Glowacki G, Deterre P, Haag F, and Koch-Nolte F (2003) NAD-induced T cell death: ADP-ribosylation of cell surface proteins by ART2 activates the cytolytic P2X7 purinoceptor. *Immunity* **19**: 571–582.
- Sesti F, Liu S, and Cai SQ (2010) Oxidation of potassium channels by ROS: a general mechanism of aging and neurodegeneration? *Trends Cell Biol* **20**:45–51.
- Sharp AJ, Polak PE, Simonini V, Lin SX, Richardson JC, Bongarzone ER, and Feinstein DL (2008) P2x7 deficiency suppresses development of experimental autoimmune encephalomyelitis. *J Neuroinflammation* **5**:33.
- Shemon AN, Slyuter R, Conigrave AD, and Wiley JS (2004) Chelerythrine and other benzophenanthridine alkaloids block the human P2X₇ receptor. *Br J Pharmacol* **142**:1015–1019.
- Shemon AN, Slyuter R, Fernando SL, Clarke AL, Dao-Ung L-P, Skarratt KK, Saunders BM, Tan KS, Gu BJ, and Fuller SJ, et al. (2006) A Thr357 to Ser polymorphism in homozygous and compound heterozygous subjects causes absent or reduced P2X7 function and impairs ATP-induced mycobacterial killing by macrophages. *J Biol Chem* **281**:2079–2086.

- Shemon AN, Sluyter R, Stokes L, Manley PW, and Wiley JS (2008) Inhibition of the human P2X₇ receptor by a novel protein tyrosine kinase antagonist. *Biochem Biophys Res Commun* **365**:515–520.
- Shemon AN, Sluyter R, and Wiley JS (2007) Rottlerin inhibits P2X₇ receptor-stimulated phospholipase D activity in chronic lymphocytic leukaemia B-lymphocytes. *Immunol Cell Biol* **85**:68–72.
- Sim JA, Young MT, Sung HY, North RA, and Surprenant A (2004) Reanalysis of P2X₇ receptor expression in rodent brain. *J Neurosci* **24**:6307–6314.
- Skarratt KK, Fuller SJ, Sluyter R, Dao-Ung LP, Gu BJ, and Wiley JS (2005) A 5' intronic splice site polymorphism leads to a null allele of the P2X₇ gene in 1–2% of the Caucasian population. *FEBS Lett* **579**:2675–2678.
- Sluyter R, Shemon AN, Hughes WE, Stevenson RO, Georgiou JG, Eslick GD, Taylor RM, and Wiley JS (2007) Canine erythrocytes express the P2X₇ receptor: greatly increased function compared with human erythrocytes. *Am J Physiol Regul Integr Comp Physiol* **293**:R2090–R2098.
- Sluyter R and Stokes L (2011) Significance of P2X₇ receptor variants to human health and disease. *Recent Pat DNA Gene Seq* **5**:41–54.
- Smart ML, Gu B, Panchal RG, Wiley J, Cromer B, Williams DA, and Petrou S (2003) P2X₇ receptor cell surface expression and cytolitic pore formation are regulated by a distal C-terminal region. *J Biol Chem* **278**:8853–8860.
- Solle M, Labasi J, Perregaux DG, Stam E, Petrushova N, Koller BH, Griffiths RJ, and Gabel CA (2001) Altered cytokine production in mice lacking P2X₇ receptors. *J Biol Chem* **276**:125–132.
- Soltoff SP, McMillian MK, and Talamo BR (1989) Coomassie Brilliant Blue G is a more potent antagonist of P₂ purinergic responses than Reactive Blue 2 (Cibacron Blue 3GA) in rat parotid acinar cells. *Biochem Biophys Res Commun* **165**:1279–1285.
- Stokes L, Trang T, Dorfman R, Smith SB, Beggs S, Ritchie J, Austin JS, Zaykin DV, Vander Meulen H, and Costigan M, et al. (2012) Genetically determined P2X₇ receptor pore formation regulates variability in chronic pain sensitivity. *Nat Med* **18**:595–599.
- Sperlágh B, Vizi ES, Wirkner K, and Illes P (2006) P2X₇ receptors in the nervous system. *Prog Neurobiol* **78**:327–346.
- Spildreorde M, Curtis SJ, Curtis BL, and Sluyter R (2014) Extracellular adenosine 5'-triphosphate and lipopolysaccharide induce interleukin-1 β release in canine blood. *Vet Immunol Immunopathol* **157**:105–110.
- Steinberg TH and Silverstein SC (1987) Extracellular ATP₄ – promotes cation fluxes in the J774 mouse macrophage cell line. *J Biol Chem* **262**:3118–3122.
- Stevenson RO, Taylor RM, Wiley JS, and Sluyter R (2009) The P2X₇ receptor mediates the uptake of organic cations in canine erythrocytes and mononuclear leukocytes: comparison to equivalent human cell types. *Purinergic Signal* **5**:385–394.
- Stock TC, Bloom BJ, Wei N, Ishaq S, Park W, Wang X, Gupta A, and Mebus CA (2012) Efficacy and safety of CE-224,535, an antagonist of P2X₇ receptor, in treatment of patients with rheumatoid arthritis inadequately controlled by methotrexate. *J Rheumatol* **39**:720–727.
- Stokes L, Fuller SJ, Sluyter R, Skarratt KK, Gu BJ, and Wiley JS (2010) Two haplotypes of the P2X₇ receptor containing the Ala-348 to Thr polymorphism exhibit a gain-of-function effect and enhanced interleukin-1 β secretion. *FASEB J* **24**:2916–2927.
- Stokes L, Jiang LH, Alcaraz L, Bent J, Bowers K, Fagura M, Furber M, Mortimore M, Lawson M, and Theaker J, et al. (2006) Characterization of a selective and potent antagonist of human P2X₇ receptors, AZ11645373. *Br J Pharmacol* **149**:880–887.
- Suadicani SO, Brosnan CF, and Scemes E (2006) P2X₇ receptors mediate ATP release and amplification of astrocytic intercellular Ca²⁺ signaling. *J Neurosci* **26**:1378–1385.
- Suadicani SO, Iglesias R, Spray DC, and Scemes E (2009) Point mutation in the mouse P2X₇ receptor affects intercellular calcium waves in astrocytes. *ASN Neuro* **1**:55–63.
- Sugiyama T, Lee SY, Horie T, Oku H, Takai S, Tanioka H, Kuriki Y, Kojima S, and Ikeda T (2013) P2X₇ receptor activation may be involved in neuronal loss in the retinal ganglion cell layer after acute elevation of intraocular pressure in rats. *Mol Vis* **19**:2080–2091.
- Surprenant A, Rassendren F, Kawashima E, North RA, and Buell G (1996) The cytolitic P_{2Z} receptor for extracellular ATP identified as a P_{2X} receptor (P2X₇). *Science* **272**:735–738.
- Syberg S, Schwarz P, Petersen S, Steinberg TH, Jensen J-EB, Teilmann J, and Jørgensen NR (2012) Association between P2X₇ receptor polymorphisms and bone status in mice. *J Osteoporos* **2012**:637986.
- Takenouchi T, Sekiyama K, Sekigawa A, Fujita M, Waragai M, Sugama S, Iwamaru Y, Kitani H, and Hashimoto M (2010) P2X₇ receptor signaling pathway as a therapeutic target for neurodegenerative diseases. *Arch Immunol Ther Exp (Warsz)* **58**:91–96.
- Taylor SRJ, Gonzalez-Begne M, Dewhurst S, Chimini G, Higgins CF, Melvin JE, and Elliott JI (2008) Sequential shrinkage and swelling underlie P2X₇-stimulated lymphocyte phosphatidylserine exposure and death. *J Immunol* **180**:300–308.
- Taylor SRJ, Gonzalez-Begne M, Sojka DK, Richardson JC, Sheardown SA, Harrison SM, Pusey CD, Tam FWK, and Elliott JI (2009a) Lymphocytes from P2X₇-deficient mice exhibit enhanced P2X₇ responses. *J Leukoc Biol* **85**:978–986.
- Taylor SRJ, Turner CM, Elliott JI, McDaid J, Hewitt R, Smith J, Pickering MC, Whitehouse DL, Cook HT, and Burnstock G, et al. (2009b) P2X₇ deficiency attenuates renal injury in experimental glomerulonephritis. *J Am Soc Nephrol* **20**:1275–1281.
- Teixeira JM, Oliveira MCG, Nociti FH Jr, Clemente-Napimoga JT, Pellegrini-da-Silva A, Parada CA, and Tambeli CH (2010a) Involvement of temporomandibular joint P2X₃ and P2X_{2/3} receptors in carrageenan-induced inflammatory hyperalgesia in rats. *Eur J Pharmacol* **645**:79–85.
- Teixeira JM, Oliveira MCG, Parada CA, and Tambeli CH (2010b) Peripheral mechanisms underlying the essential role of P2X₇ receptors in the development of inflammatory hyperalgesia. *Eur J Pharmacol* **644**:55–60.
- Théâtre E, Frederix K, Guilmain W, Delierneux C, Lecut C, Bettendorff L, Bours V, and Oury C (2012) Overexpression of CD39 in mouse airways promotes bacteria-induced inflammation. *J Immunol* **189**:1966–1974.
- Thompson RJ, Jackson MF, Olah ME, Rungta RL, Hines DJ, Beazely MA, MacDonald JF, and MacVicar BA (2008) Activation of pannexin-1 hemichannels augments aberrant bursting in the hippocampus. *Science* **322**:1555–1559.
- Tschopp J (2011) Mitochondria: Sovereign of inflammation? *Eur J Immunol* **41**:1196–1202.
- Tu G, Li G, Peng H, Hu J, Liu J, Kong F, Liu S, Gao Y, Xu C, and Xu X, et al. (2013) P2X₇ inhibition in stellate ganglia prevents the increased sympathoexcitatory reflex via sensory-sympathetic coupling induced by myocardial ischemic injury. *Brain Res Bull* **96**:71–85.
- Valdez-Morales E, Guerrero-Alba R, Liñán-Rico A, Espinosa-Luna R, Zarazua-Guzman S, Miranda-Morales M, Montaña LM, and Barajas-López C (2011) P2X₇ receptors contribute to the currents induced by ATP in guinea pig intestinal myenteric neurons. *Eur J Pharmacol* **668**:366–372.
- Vergani A, Fotino C, D'Addio F, Tezza S, Podetta M, Gatti F, Chin M, Bassi R, Molano RD, and Corradi D, et al. (2013a) Effect of the purinergic inhibitor oxidized ATP in a model of islet allograft rejection. *Diabetes* **62**:1665–1675.
- Vergani A, Tezza S, D'Addio F, Fotino C, Liu K, Niewczas M, Bassi R, Molano RD, Kleffel S, and Petrelli A, et al. (2013b) Long-term heart transplant survival by targeting the ionotropic purinergic receptor P2X₇. *Circulation* **127**:463–475.
- Verkhatsky A, Pankratov Y, Lalo U, and Nedergaard M (2012) P2X receptors in neuroglia. *Wiley Interdiscip Rev Membr Transp Signal* **1**:151–161.
- Virginio C, MacKenzie A, Rassendren FA, North RA, and Surprenant A (1999) Pore dilation of neuronal P2X receptor channels. *Nat Neurosci* **2**:315–321.
- Volonté C, Apolloni S, Skaper SD, and Burnstock G (2012) P2X₇ receptors: channels, pores and more. *CNS Neurol Disord Drug Targets* **11**:705–721.
- Wang B and Sluyter R (2013) P2X₇ receptor activation induces reactive oxygen species formation in erythroid cells. *Purinergic Signal* **9**:101–112.
- Wang J, Yang J, Liu P, Bi X, Li C, and Zhu K (2012) NAD induces astrocyte calcium flux and cell death by ART2 and P2X₇ pathway. *Am J Pathol* **181**:746–752.
- Wang X, Arcuino G, Takano T, Lin J, Peng WG, Wan P, Li P, Xu Q, Liu QS, and Goldman SA, et al. (2004) P2X₇ receptor inhibition improves recovery after spinal cord injury. *Nat Med* **10**:821–827.
- Weber FC, Esser PR, Müller T, Ganesan J, Pellegatti P, Simon MM, Zeiser R, Idzko M, Jakob T, and Martin SF (2010) Lack of the purinergic receptor P2X₇ results in resistance to contact hypersensitivity. *J Exp Med* **207**:2609–2619.
- Welter-Stahl L, da Silva CM, Schachter J, Persechini PM, Souza HS, Ojcius DM, and Coutinho-Silva R (2009) Expression of purinergic receptors and modulation of P2X₇ function by the inflammatory cytokine IFN γ in human epithelial cells. *Biochim Biophys Acta* **1788**:1176–1187.
- Westlund KN, Zhang L, Ma F, and Oz HS (2012) Chronic inflammation and pain in a tumor necrosis factor receptor (TNFR) (p55/p75^{-/-}) dual deficient murine model. *Transl Res* **160**:84–94.
- Wiley JS, Dao-Ung L-P, Li C, Shemon AN, Gu BJ, Smart ML, Fuller SJ, Barden JA, Petrou S, and Sluyter R (2003) An Ile-568 to Asn polymorphism prevents normal trafficking and function of the human P2X₇ receptor. *J Biol Chem* **278**:17108–17113.
- Wiley JS and Gu BJ (2012) A new role for the P2X₇ receptor: a scavenger receptor for bacteria and apoptotic cells in the absence of serum and extracellular ATP. *Purinergic Signal* **8**:579–586.
- Wiley JS, Sluyter R, Gu BJ, Stokes L, and Fuller SJ (2011) The human P2X₇ receptor and its role in innate immunity. *Tissue Antigens* **78**:321–332.
- Wilhelm K, Ganesan J, Müller T, Dürr C, Grimm M, Beilhack A, Krempl CD, Sorichter S, Gerlach UV, and Jüttner E, et al. (2010) Graft-versus-host disease is enhanced by extracellular ATP activating P2X₇R. *Nat Med* **16**:1434–1438.
- Williams RO (2012) What have we learned about the pathogenesis of rheumatoid arthritis from TNF-targeted therapy? *ISRN Immunol* **2012**:652739.
- Wilson HL, Francis SE, Dower SK, and Crossman DC (2004) Secretion of intracellular IL-1 receptor antagonist (type 1) is dependent on P2X₇ receptor activation. *J Immunol* **173**:1202–1208.
- Wilson HL, Varcoe RW, Stokes L, Holland KL, Francis SE, Dower SK, Surprenant A, and Crossman DC (2007) P2X receptor characterization and IL-1/IL-1Ra release from human endothelial cells. *Br J Pharmacol* **151**:115–127.
- Xie Y, Williams CD, McGill MR, Lefobsky M, Ramachandran A, and Jaeschke H (2013) Purinergic receptor antagonist A438079 protects against acetaminophen-induced liver injury by inhibiting p450 isoenzymes, not by inflammasome activation. *Toxicol Sci* **131**:325–335.
- Xu XJ, Boumehache M, Robinson LE, Marschall V, Gorecki DC, Masin M, and Murrell-Lagnado RD (2012) Splice variants of the P2X₇ receptor reveal differential agonist dependence and functional coupling with pannexin-1. *J Cell Sci* **125**:3776–3789.
- Yan Z, Khadra A, Sherman A, and Stojilkovic SS (2011) Calcium-dependent block of P2X₇ receptor channel function is allosteric. *J Gen Physiol* **138**:437–452.
- Yan Z, Li S, Liang Z, Tomić M, and Stojilkovic SS (2008) The P2X₇ receptor channel pore dilates under physiological ion conditions. *J Gen Physiol* **132**:563–573.
- Yanagisawa D, Kitamura Y, Takata K, Hide I, Nakata Y, and Taniguchi T (2008) Possible involvement of P2X₇ receptor activation in microglial neuroprotection against focal cerebral ischemia in rats. *Biol Pharm Bull* **31**:1121–1130.
- Yegutkin GG (2008) Nucleotide- and nucleoside-converting ectoenzymes: Important modulators of purinergic signalling cascade. *Biochim Biophys Acta* **1783**:673–694.
- Yiangou Y, Facer P, Durrenberger P, Chessell IP, Naylor A, Bountra C, Banati RR, and Anand P (2006) COX-2, CB2 and P2X₇-immunoreactivities are increased in activated microglial cells/macrophages of multiple sclerosis and amyotrophic lateral sclerosis spinal cord. *BMC Neurol* **6**:12.
- Yip L, Woehrlé T, Corriden R, Hirsh M, Chen Y, Inoue Y, Ferrari V, Insel PA, and Junger WG (2009) Autocrine regulation of T-cell activation by ATP release and P2X₇ receptors. *FASEB J* **23**:1685–1693.

- Young CNJ, Brutkowski W, Lien CF, Arkle S, Lochmüller H, Zablocki K, and Górecki DC (2012) P2X7 purinoceptor alterations in dystrophic mdx mouse muscles: relationship to pathology and potential target for treatment. *J Cell Mol Med* **16**:1026–1037.
- Young MT, Pelegrin P, and Surprenant A (2006) Identification of Thr283 as a key determinant of P2X7 receptor function. *Br J Pharmacol* **149**:261–268.
- Young MT, Pelegrin P, and Surprenant A (2007) Amino acid residues in the P2X7 receptor that mediate differential sensitivity to ATP and BzATP. *Mol Pharmacol* **71**:92–100.
- Zanovello P, Bronte V, Rosato A, Pizzo P, and Di Virgilio F (1990) Responses of mouse lymphocytes to extracellular ATP. II. Extracellular ATP causes cell type-dependent lysis and DNA fragmentation. *J Immunol* **145**:1545–1550.
- Zhao J, Wang H, Dai C, Wang H, Zhang H, Huang Y, Wang S, Gaskin F, Yang N, and Fu SM (2013) P2X7 blockade attenuates murine lupus nephritis by inhibiting activation of the NLRP3/ASC/caspase 1 pathway. *Arthritis Rheum* **65**:3176–3185.
- Zhou ZH, Wang JX, Liu BJ, Li M, Lu Y, and Chen HS (2012) Contribution of the spinal P2X7 receptors to bee venom-induced nociception and inflammation in conscious rats. *Neurosci Lett* **531**:145–148.
- Zhu S, Wang Y, Wang X, Li J, and Hu F (2014) Emodin inhibits ATP-induced IL-1 β secretion, ROS production and phagocytosis attenuation in rat peritoneal macrophages via antagonizing P2X₇ receptor. *Pharm Biol* **52**:51–57.