

Lawrence Berkeley National Laboratory

Recent Work

Title

The Path of Carbon in Photosynthesis V. Paper Chromatography and Radioautography of the Products (1)

Permalink

<https://escholarship.org/uc/item/7977d0dn>

Authors

Benson, A.A.
Bassham, J.A.
Calvin, M.
et al.

Publication Date

1949-06-13

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

UCRL-363
02

Unclassified Distribution

UNIVERSITY OF CALIFORNIA
Radiation Laboratory

Contract No. W-7405-eng-48

UNCLASSIFIED

THE PATH OF CARBON IN PHOTOSYNTHESIS. V. PAPER CHROMATOGRAPHY
AND RADIOAUTOGRAPHY OF THE PRODUCTS (1)

by

A. A. Benson, J. A. Bassham, M. Calvin, T. C. Goodale,
V. A. Haas and W. Stepka

June 13, 1949

Berkeley, California

THE PATH OF CARBON IN PHOTOSYNTHESIS. V. PAPER CHROMATOGRAPHY
AND RADIOAUTOGRAPHY OF THE PRODUCTS (1)

by

A. A. Benson, J. A. Bassham, M. Calvin, T. C. Goodale,

V. A. Haas and W. Stepka

Radiation Laboratory and Department of Chemistry
and Division of Plant Nutrition, University of
California, Berkeley, California

June 13, 1959

ABSTRACT

1. Paper chromatography has been employed to separate the radioactive products formed during photosynthesis in $C^{14}O_2$.
2. The method has been used for the separation and identification of carboxylic acids and phosphate esters.
3. The first observed product of carbon dioxide assimilation during photosynthesis has been isolated and shown to be phosphoglyceric acid.

For publication in The Journal of the American Chemical Society.

(1) The work described in this paper was sponsored by the Atomic Energy Commission.

THE PATH OF CARBON IN PHOTOSYNTHESIS. V. PAPER CHROMATOGRAPHY
AND RADIOAUTOGRAPHY OF THE PRODUCTS (1)

by

A. A. Benson, J. A. Bassham, M. Calvin, T. C. Goodale,

V. A. Haas and W. Stepka

Radiation Laboratory and Department of Chemistry and
Division of Plant Nutrition, University of California,
Berkeley, California

Investigation of the reduction of carbon dioxide in plants requires a rapid, general and unequivocal method for separating and identifying a large number of compounds. Paper chromatography is such a method. This paper constitutes a report of progress in the development of experimental procedures in the application of paper chromatography to the separation of compounds formed by plants during short periods of photosynthesis. The compounds involved are important intermediates of plant and animal metabolism. These methods should find application not only in other biochemical problems, but in the control of chemical synthesis (2) as well.

A number of products formed during a short period of photosynthesis with radioactive carbon dioxide have been reported (3,4,5,6,7). The classical

-
- (1) The work described in this paper was sponsored by the Atomic Energy Commission.
 - (2) R.L.M. Synge, *Biochem. J.*, 42, 99 (1948).
 - (3) A.A. Benson, et. al., "Photosynthesis in Plants" ed. by J. Franck and W. E. Loomis (Iowa State College Press, Ames, Iowa, 1949). Chapter 19.
 - (4) M. Calvin and A.A. Benson, *Science*, 107, 476 (1948).
 - (5) W. Stepka, A. A. Benson and M. Calvin, *Science*, 108, 304 (1948).
 - (6) M. Calvin and A. A. Benson, *Cold Spring Harbor Symposia in Quantitative Biology*, Vol. XIII, page 6, 1948.
 - (7) M. Calvin and A. A. Benson, *Science*, 109, 140 (1949).
-

biochemical methods and ion exchange techniques which were used in those identifications would require considerable effort to separate the large number of compounds involved. The success of paper chromatography in separating two groups of compounds important in phytosynthesis, amino acids (8) and sugars (9,10), prompted the present work on separation of polycarboxylic acids and phosphate esters. These compounds constitute the presently known intermediates in the synthesis of plant material (11).

The primary requisite for the use of this method is some means of determining the position of a compound in the chromatogram. Both chemical and physical methods have been used. The former includes color tests which depend on development of a color directly on the paper by the use of specific reagents: ninhydrin for amino acids (8), naphthoresorcinol or Tollens reagent for sugars (10), ferric ion for hydroxamic acids (12,13), pH indicators for acids (14). One physical method is the detection of fluorescence of the compound itself or of some derivative of it produced on the paper, excited by ultraviolet light (15).

A very general physical method depends on the incorporation of a radioactive isotope into the compound and its detection by a Geiger counter or a

-
- (8) R. Consden, A.H. Gordon and A.J.P. Martin, *Biochem J.*, 38, 224 (1944).
(9) A. E. Flood, E. L. Hirst and J.K.N. Jones, *Nature*, 160, 86 (1947).
(10) S. M. Partridge, *Biochem. J.*, 42, 238 (1948); *Nature*, 158, 270 (1946).
(11) A preliminary description of the general application of paper chromatography with radioisotopes has been published (7).
(12) E. Stadtman, private communication.
(13) R. M. Fink and K. Fink, Atomic Energy Commission Report MDDC-1487 (1947), "The Use of Radioactive Reagents in Filter Paper Partition Chromatography"
(14) J.W.H. Lugg and B.T. Overell, *Nature*, 160, 87 (1947); *Australian Journal Sci. Res.*, 1, 98 (1948).
(15) D.M.P. Phillips, *Nature*, 161, 53 (1948).
-

photographic film. Such a detection method finds ideal application in the problem of determining the path of carbon in photosynthesis, since it is possible to detect the number and position of all compounds containing the radioactive carbon atoms by exposing the paper to a sheet of X-ray film. Radioautography has been used to identify amino acids (16) and the extension to other compounds (7) depends only upon the ability of the method to separate the compound and the identification of the resultant spot.

Identification of those compounds for which specific color reactions exist is relatively simple. For many compounds found in plant extracts no color reactions are known. The most useful property of organic substances in the identification of tracer amounts of material has been the distribution coefficient between immiscible solvents. It is readily determined and is highly characteristic, especially when measured as a function of pH. It has proved especially useful to distinguish between a number of possibilities for an unknown spot. We have found it convenient to synthesize the suspected compound containing a radioisotope and to determine the chromatographic position by radioautography. Unequivocal identification is then accomplished by preparation of a mixed chromatogram. The isolated substance eluted (17,18) from the paper is co-chromatographed with the authentic specimens. Identity of the two samples is evidenced by the appearance of a single spot.

CHROMATOGRAPHY

Two-dimensional paper chromatograms were prepared in a conventional manner (8,19). The solvent flowed downwards through the paper suspended from

-
- (16) R.M. Fink, C.E. Dent and K. Fink, *Nature*, 160, 801 (1947).
(17) C.E. Dent, *Biochem. J.*, 41, 240 (1947).
(18) R. Consden, A.H. Gordon and A.J.P. Martin, *Biochem. J.*, 41, 590 (1947).
(19) C.E. Dent, *Biochem. J.*, 43, 169 (1948).
-

a long stainless steel trough, Fig. 1. To reduce volatilization and decomposition of compounds the paper was dried in a hood without heat.

Selection of Solvents: - Phenol, saturated with water at the room temperature, 24° C, was chosen as the first solvent and proved satisfactory in moving most of the radioactive components.

The use of lutidine or collidine as a second solvent is prohibited by the instability of many phosphate esters in these bases. Much hydrolysis and formation of immovable material occurred with lutidine and streaking of spots indicated slow decomposition.

The acidic solvent butanol-propionic acid-water was chosen as the second solvent for much of the present work since it effects an optimum separation of a majority of components. By varying the proportion of propionic acid it is possible to change the water content of the organic phase from 20 to 50%. An intermediate composition, approximately 31% water, appeared to give an optimum separation of the compounds involved. Another solvent, butanol-acetic acid-water, has been used and is especially effective in moving the phosphate esters. Such solutions must be freshly prepared, since the composition of the phases changes upon esterification. In several experiments the solvent system ethylene glycol diethyl ether-propionic acid (formic acid)-water provided a solvent with desirable characteristics and without the disadvantage of esterification.

Butanol-propionic acid-water: - Fresh solvent is prepared from equal volumes of two solutions, A (124.6 ml. n-butanol and 84 ml. water) and B (620 ml. redistilled propionic acid and 790 ml. water). A and B are adjusted to give a single phase solution which becomes cloudy upon cooling to 22° C or 2° C below the thermostatted room temperature. Stainless steel troughs 24 inches long with a semi-circular cross section 1-1/2 inches in diameter,

Fig. 1, were filled with 90-120 ml. of solvent for two papers (18 inches by 22-1/2 inches). The time required for the solvent to traverse the paper varied from fifteen to twenty hours.

Butanol-acetic acid-water: - A fresh solution is prepared from 74 ml. n-butanol, 19 ml. acetic acid and 50 ml. water.

Preparation of Radioautographs: - The dried chromatogram was folded to fit the conventional 14 inches by 17 inches Eastman No-Screen X-ray film with the origin at one corner. One or two films were then exposed to the chromatogram in a light-proof "Eastman X-Ray Exposure Holder" which can be stacked and weighted with a carton of sand to equalize pressure distribution. In the case of C^{14} - P^{32} radiograms both sides of the duplitized film are exposed by phosphorus while C^{14} exposes only the adjacent emulsion. After development of the film only the P^{32} activity is shown on one side of the film while the other is exposed by both radiations. For documentation it is advisable to use two sheets of film on one side of the chromatogram so that the second sheet shows only P^{32} activity.

The activity emerging from the paper (Whatman No. 1 filter paper) is 30% of that detectable if the sample were dried in a thin film on glass or metal and counted directly. The time of exposure of the X-ray film depends upon the intensity of the radiation coming from the paper. An activity of 15,000 disintegrations per minute per cm^2 produces satisfactory exposure in three to six days.

It has not been possible to reproduce the radiograms accurately by photography. Very faint radioactive spots are readily discerned when the film is observed by light reflected from a white background. The details of more exposed spots such as the phosphate esters are clearly seen when viewed over a standard X-ray illuminator.

Radioactive ink (20) provides an exact record of the relative position

(20) We are indebted to Mr. A. G. Hall of this laboratory for this suggestion and preparation of the stamp.

of the film and chromatogram during exposure. Opposite corners of the chromatogram are marked with stamp pad ink containing a non-volatile C¹⁴-labeled compound.

Application of Plant Extracts on Paper: - The 80 percent ethanol extract of algae (1 g. packed cells) or leaves (2 g.) is concentrated in vacuo (less than 20° C) to 2 ml. Ten to two hundred microliters of extract is applied to the 1.5 cm. origin circle. An air stream warmed with a flame, when stability of the compounds allowed it, rapidly dried the spot by flowing above and below the paper. Infra-red lamps caused decomposition and are not satisfactory heat sources.

The capacity of Whatman No. 1 filter paper varies with the type of compound and the solvent. Loads varying from a few micrograms for carboxylic acids and up to 1 or 2 mg. for the sugars have been successfully separated.

It is possible to separate simple mixtures on a larger scale by a one-dimensional technique. A mixture is applied in a 1.5 cm. wide strip near the edge of a sheet and separated in an appropriate solvent into two or more bands. As much as 20 mg. of hexoses or 4 mg. of hexose phosphates have been separated in this way.

IDENTIFICATION

The majority of compounds so far found to be involved in photosynthesis are well known metabolic intermediates whose physical and chemical properties are known. Earlier work (3) had established the presence of large fractions

of the fixed activity in alanine, malic acid, sucrose and phosphoglyceric acid. When the chromatograms were made it was a relatively straightforward matter to determine which of the forty spots detectable in a radiogram of 90 seconds photosynthesis by Scenedesmus corresponded to these compounds. Identification of the remaining compounds is begun in this work.

The positions of these radioactive compounds are reproducible for extracts of a given plant. However, absolute R_F values for compounds in such mixtures should not be applied universally (19). The presence of contiguous substances on the paper, whether they be radioactive or not, may disturb equilibrium and movement of spots. For example, inorganic phosphate is found to restrict the movement of adjacent compounds. It is necessary, then, to consider the relative location of known vicinal spots rather than the absolute position on the chromatograms. The location of a spot relative to several standards, that is, the pattern of spots, presents the most reliable method of describing chromatographic position. In this paper, the positions of the neutral amino acids, serine and alanine, are taken as reference points. A schematic plot of R_F values of compounds involved in photosynthesis is given in Fig. 2.

The Carboxylic Acids (21): - According to Consden, Gordon and Martin (8) the R_F of a solute is given by

$$R_F = \frac{A_L}{A_L + \alpha A_S}$$

where A_L is the fraction of cross-sectional area occupied by the mobile phase, A_S is the fraction of cross-sectional area occupied by the stationary

(21) Abstracted from the thesis of J. A. Bassham submitted in partial fulfillment of the requirements for the Ph.D. degree, University of California, Berkeley, California, June 1949.

phase, and g , the distribution coefficient of the solute, is the concentration of solute in the stationary phase divided by the concentration of solute in the mobile phase. This expression can be written as

$$R_F = \frac{1}{1 + gA_S/A_L}$$

The term A_S/A_L is a constant for a given paper and set of solvents while g is characteristic of the solute and determines its position on the paper.

The gross distribution of a carboxylic acid between the stationary aqueous phase and the mobile organic phase depends on the distribution coefficient of the undissociated acid and on the degree of dissociation of the acid. The degree of dissociation depends on the pH of the aqueous phase and on the dissociation constant of the acid. Consequently, the gross distribution of a given acid, and hence its R_F value, is, in general, a function of two physical properties of the acid, its distribution coefficient and its dissociation constant(s) and one experimental condition, the pH of the aqueous phase of the chromatogram.

If neutral, non-buffered solvents are employed, the pH of the aqueous phase at a given point depends only on the concentration of carboxylic acid at that point, and as the mobile phase carried the acid past this point there is a decreasing concentration and increasing degree of ionization of the acid with a resultant changing distribution of the acid between phases. The rate of movement of the acid therefore decreases when its concentration decreases and streaking is produced (14).

This effect can be avoided by maintaining the pH of the aqueous phase at a constant value. This is accomplished either by employing an acidic solvent or a buffered solvent. Lugg and Overell have used an acidic solvent

of a solution to effect the separation of carboxylic acids. By using formic acid as a "swamping" acid, they suppressed the ionization of the test acids so that the R_F values depend principally on the distribution coefficients of the free acids.

It is possible to adjust the pH of the aqueous phase by employing buffers. This can be accomplished either by applying the buffer to the origin or by dissolving the buffer in the solvent beforehand. The R_F value then depends on both the distribution coefficient and the dissociation constant of the carboxylic acid.

It is desirable in two-dimensional paper chromatography to employ a low pH solvent in one direction and a higher pH solvent in the other direction so that a different order and degree of separation is obtained in the two directions. While it is possible to obtain good separations of a number of carboxylic acids with two-dimensional chromatograms which employ acidic solvents in both directions (14), the resultant positions of the acids tend to overlap those of amino acids, sugars and other plant constituents. The use of solvents of differing pH moves the acids to a region of the paper apart from the rest of the plant constituents, as shown in Fig. 2. Such "area" separation is desirable in analysis of plant extracts in that it circumvents preliminary fractionations.

The usual method of buffering in the phenol direction has been to apply the buffer to the origin along with the mixture of acids or cell extract. Since the buffer capacity thus obtained is limited it is necessary to use very small amounts of acids. From 1 to 5 micrograms of acid gives satisfactory results. The colorimetric methods described by Lugg and Overall give best results for 20 micrograms or more, so the tracer method of detection is preferable for this purpose.

Several carboxylic acids labeled with C^{14} have been synthesized and chromatographed to determine their positions with the solvents employed. The resultant R_F values are given in Fig. 2 and they represent average values obtained from chromatograms buffered at pH 6-7 at the origin. In addition to malic, succinic, fumaric and tartaric acids, which were studied by Lugg and Overell, isocitric (22), glycollic (23) and glyceric acids have been separated.

The relation of R_F values in an acidic solvent to the distribution coefficients is shown in Table I. The R_F values and the distribution coefficients of the acids, of alanine and of glucose were measured. Using the constants of alanine as a standard, the term A_G/A_L was calculated and found to be 0.55. The R_F values of other solutes can then be calculated according to the expression

$$R_F = \frac{1}{1 + 0.55\alpha}$$

Comparison of the calculated R_F values with measured R_F values shows fairly good agreement and affords additional evidence that paper chromatograms of this type are partition chromatograms.

Inspection of radiograms of cell extracts from plants which had been exposed to C^{14} -labeled carbon dioxide for short periods indicated several compounds in the carboxylic acid region. The synthetic labeled acids were co-chromatographed, one at a time, with the plant extracts. Intensification of an unknown radioactive spot by the authentic compound demonstrated their identity. Plant acids were then eluted individually from the paper and

(22) We are indebted to Dr. R. E. Stutz for a sample of C^{14} -labeled isocitric acid.

(23) Labeled glycollic acid was obtained from Dr. B. M. Tolbert. See "Isotopic Carbon" by Calvin, Heidelberger, Reid, Tolbert and Yankwich, John Wiley and Sons, Inc., 1949.



TABLE I
Comparison of Measured and Calculated R_F Values in
Butanol-Propionic Acid-Water Solvent

Solute	α	R_F (calculated)	R_F (measured)
Alanine	3.65	-----	0.333
Fumaric	0.50	0.78	.72
Succinic	1.02	.64	.65
Glycolic	2.06	.47	.51
Malic	2.40	.43	.45
Isocitric	2.75	.40	.41
Glyceric	2.95	.38	.38
Tartaric	3.20	.36	.32
Glucose	5.75	.24	.20

co-chromatographed with the corresponding synthetic compound. Finally, if no separation appeared between unknown and synthetic acid, the unknown was subjected to suitable physical and chemical confirmatory tests.

The acids formed in short term photosynthesis and positively identified so far include malic, succinic, glycolic and fumaric. Those found in respiration experiments (dark fixation of labeled carbon dioxide without preillumination of the plant) include isocitric or citric, succinic, fumaric and malic.

Synthesis of Succinic and Malic Acids: - One gram of carboxyl-labeled sodium acetate (specific activity, 6 $\mu\text{c./mg.}$) was converted to malic and succinic acids by the following series of reactions:



The standard methods were modified when necessary for small-scale synthesis. The mixture of succinic and malic esters was obtained by using unpoisoned PtO₂ catalyst and one atmosphere of hydrogen at room temperature with rapid stirring of a 95 percent ethanol solution for two hours (24). The free acids were separated by fractional ether extraction and recrystallization. The yield of succinic acid was 28 percent of the theoretical and of malic acid 39 percent of theoretical, so that the total yield was 67 percent. The specific activity of the malic acid was 3.7 μc./mg. and that of the succinic acid was 4.2 μc./mg.

Synthesis of Fumaric Acid: - Malic acid was converted to fumaric acid by heating in nitrogen at 140° C for two hours. Fifty milligrams of carboxyl-labeled malic acid gave 30 mg. (69 percent) of fumaric acid and 13 mg. unreacted malic acid. The specific activity of the product was 4.2 μc./mg.

Synthesis of dl-Tartaric Acid: - Fumaric acid was oxidized with sodium chlorate and osmium tetroxide according to the method of Milas and Terry (25)..

Identification of Glyceric Acid: - The radioactive acid was eluted from paper with water. Hydrochloric acid was added to give a 1.0 N solution in the graduated centrifuge tube. After ether was added and the phases

(24) M. Fallebin, Ann. chim., [10] 4, 156 (1925).

(25) N. A. Milas and E. M. Terry, J. Am. Chem. Soc., 47, 1412 (1925).

mixed by forcing one phase through the other with a narrow tipped dropper, the mixture was centrifuged. In 1.0 ml. of the 7.0 ml. ether phase was found 140 c.p.m. In 0.050 ml. of the 1.8 ml. aqueous phase was found 1300 c.p.m. The resultant distribution coefficient, 0.005, approximates that of authentic glyceric acid, 0.004-0.008.

Authentic radioactive glyceric acid was prepared by hydrolysis (3) of phosphoglyceric acid eluted from a number of chromatograms. After heating two days in a sealed evacuated tube at 120° C in 1 N hydrochloric acid the solvent was evaporated in vacuo. The product was taken up in water and an aliquot part co-chromatographed with the eluted unknown material. A single spot was obtained.

Identification of Malic Acid: - The spot on a chromatogram suspected to be malic acid was eluted and its distribution coefficient between water (pH 1) and ether was found to be 0.020. The constant for low concentrations of the authentic labeled dl-malic acid is 0.017. The eluate, 20,500 c.p.m., was co-crystallized with 50.0 mg. of malic acid and found to have a specific activity of 395 c.p.m./mg. unchanged by sublimation in vacuo. This was dehydrated by heating under nitrogen for 24 hours at 145° C. The specific activity of the fumaric acid was 500 ± 50 c.p.m./mg. The distribution coefficient between water (pH 1) and ether was found to be 0.9 (authentic value, 1.0).

Phosphate Esters: - The greater part of the activity fixed in short periods of photosynthesis (26) lies in the area near the origin (Fig. 3). A characteristic of this area is the presence of three distinct spots which moved

(26) $C^{14}O_2$ was given to the plants during steady-state photosynthesis with ordinary carbon dioxide for the lengths of time indicated.

about the same small distance in the phenol but different distances in the butanol-propionic acid and thus lie one above the other in the butanol direction. As the periods of photosynthesis with $C^{14}O_2$ are shortened, the fraction of activity in this area increased. In five seconds photosynthesis only the uppermost spot in the pattern of three is radioactive. Therefore, these compounds include the first and some of the subsequent intermediates in fixation and reduction of carbon by plants (27).

It was suspected that the compounds in this area of the chromatogram were phosphate compounds. This was confirmed by chromatographing the extract of some algae which had photosynthesized for an hour with normal CO_2 in the presence of inorganic radiophosphate (P^{32}). The radiogram thus obtained (Fig. 4) was remarkably similar in this area to those obtained from radio-carbon experiments.

P^{32} -labeled known phosphorylated intermediates of glycolysis were prepared by yeast extract fermentation of glucose. A radiogram of the water extract is very similar to the P^{32} photosynthesis radiogram and the area near the origin in C^{14} photosynthesis radiogram, Fig. 3. The components of this mixture were separated by standard procedures dependent on solubility as well as by selective elution from a strong base anion exchange resin, Dowex A-1. By the latter method a series of fractions were obtained from which the radioactive compounds could be more readily separated by chromatography in the butanol-acid-water solvent, Fig. 5a,b.

Yeast Fermentation: - One hundred grams of Type 20-40 Fleischmann dry yeast was suspended in 300 ml. of M/15 potassium monohydrogen phosphate at 37° for three hours. The mixture was centrifuged and to 40 ml. of the

(27) Similar radiograms have been obtained from Chlorella, Scenedesmus, geranium and barley.

supernate was added 155 ml. of 50 percent glucose and 10 mg. sodium pyruvate. After incubation at 32° for 30 minutes vigorous fermentation had started. Then 5 ml. potassium dihydrogen phosphate containing 2 mcuries of P³² was added to the fermentation mixture. After three hours incubation, approximately 80 percent of the total phosphorus was organic. The mixture was incubated for an additional three hours after the addition of 2 ml. of 1 M sodium fluoride and 6 g. sodium pyruvate. Absolute alcohol was then added to the fermentation mixture to a final concentration of 95 percent. All of the proteins and 94 percent of the phosphorus compounds precipitated. The precipitate was washed three times with water which extracted over 95 percent of the phosphorus activity. An aliquot of this water extract gave six major spots on the radiogram.

Separation of Fermentation Products: - The water extract was absorbed on a Dowex 4-1 anion exchange resin column in the chloride form. A water wash removed 2.4 percent of the phosphate. Elution with 0.1 M sodium chloride separated the remaining phosphates into two distinct fractions. The first fraction eluted from the resin was found to contain fructose-6-phosphate, inorganic phosphate and at least three unidentified compounds. The second fraction contained only fructose-1,6-diphosphate and phosphoglyceric acid, Fig. 5a. The fractions were chromatographed in the butanol-acetic acid solvent and the individual spots of the radiograms were identified by spray tests applied to the paper or by chemical analysis of solutions of the eluted spots.

Identification of Fructose-1,6-diphosphate: - The spot having the lowest R_F value in both the phenol and butanol-acetic acid directions was eluted and identified as fructose-1,6-diphosphate by the following tests.

analysis (28,29) showed that there were two moles of phosphorus to one mole of fructose. Seven minutes hydrolysis at 100° in a 1 N hydrochloric acid converted 25 percent of the original organic phosphorus to inorganic phosphorus. When this seven-minute hydrolysate was re-chromatographed, two new radioactive spots of equal intensity appeared. One of the new spots contained only inorganic phosphate; the other spot which contained fructose and no easily hydrolyzable phosphorus was later shown to be fructose-6-phosphate.

Identification of Fructose-6-Phosphate: - In the radiograms of samples containing the first radioactive components eluted from the anion exchange column, a spot, Fig. 5b, was obtained in the butanol-acetic acid-water solvent the intensity of which, from sample to sample, corresponded directly to the fructose analysis. The eluted spots contained fructose and phosphate in a 1:1 ratio. The chromatographic position of this compound was identical with the fructose compound obtained by hydrolysis of fructose-1,6-diphosphate. Therefore, it must be an acid stable fructose monophosphate, presumably fructose-6-phosphate. A mixed chromatogram of this material with inactive glucose-1-phosphate showed considerable overlapping of the two compounds.

Identification of Phosphoglyceric Acid: - The spot labeled PGA in Fig 5a gave a negative spray test for easily hydrolyzable phosphate. A solution of the eluted spot contained no fructose and gave no inorganic phosphate when hydrolyzed for seven minutes in 1 N hydrochloric acid at 100°. When heated for one hour with α -naphthoresorcinol in concentrated sulfuric

(28) J. H. Roe, J. Biol. Chem., 107, 15 (1943).

(29) R.J.L. Allen, Biochem. J., 34, 858 (1940).

acid, a blue color (30), the same as obtained with an authentic sample of 3-phosphoglyceric acid, was observed.

Phosphoglyceric Acid in Algal Extracts: - Two early experiments indicated that the uppermost of the pattern of three spots shown in photosynthesis radiograms might be phosphoglyceric acids. The first was a chromatogram of the sodium hydroxide eluate of the weak base anion resin, Duolite A-3, which had been shown to contain largely phosphoglyceric acid (3,7) radioactivity. The position of the spot obtained corresponded with that of the uppermost of the three spots. The second experiment involved isolation of radioactive barium phosphoglycerate from a small aliquot of the extract from a one-hour photosynthesis in inorganic radiophosphate by co-crystallization with 10 mg. of the authentic barium salt. After three recrystallizations from water the salt was decationized and co-chromatographed with an algal extract (90 second photosynthesis in $C^{14}O_2$). Exact correspondence of the P^{32} activity with C^{14} activity in the uppermost spot of the phosphate area was observed. After the radiophosphorus had decayed to five percent of its initial activity, the pattern obtained was identical to that originally apparent and the uppermost spot of the pattern of three which now was made almost entirely by carbon radiation had the identical position and shape of that given by the phosphorus formerly present.

These experiments strongly indicated the identity of the upper spot as phosphoglyceric acid but did not preclude the possibility of a highly active contaminant in co-crystallizations. Synthesis of labeled phosphoglyceric acid by yeast fermentation provided authentic material for co-chromatographing with unknown mixtures.

(30) S. Rapaport, *Biochem. Z.*, 289, 406 (1937).

Carrier-Free Isolation of Phosphoglyceric Acid: - When the period of photosynthesis in radiocarbon is sufficiently shortened only one major radioactive compound appears and is in the position corresponding to the uppermost of the pattern of three. This radioactive compound was isolated from Scenedesmus which had photosynthesized for 5 seconds (7) in $C^{14}O_2$ and was identified as phosphoglyceric acid. The isolation was carried out according to the general method for phosphoglyceric acid with several modifications. It was found that the radioactive material was strongly adsorbed by inactive precipitates removed at pH 7. Thorough washing at pH 10, where the adsorbed anions are apparently displaced from the precipitate, was necessary to remove the activity. When barium chloride was added to an acid solution, the precipitate obtained was active, but repeated washing with dilute hydrochloric acid removed the adsorbed activity. When the barium-containing supernatant solution from this acid precipitation was adjusted to pH 10, no barium phosphoglycerate precipitated since the volume of the solution was large enough to retain the amount of material present. Addition of ethanol gave a radioactive precipitate. The fructose-1,6-diphosphate which would be obtained by this procedure was hydrolyzed by repeated warming in dilute acid. The amount of hexose diphosphate in algal extracts must be much smaller than the amount of phosphoglyceric acid since the radiograms of longer photosyntheses, where the reservoirs of intermediates are saturated with activity, show at least three times as much phosphoglyceric acid.

Repeated dissolution of the barium salt in 0.05 N hydrochloric acid and reprecipitation from 50 percent ethanol gave a barium salt whose phosphorus analysis and molybdate enhanced optical rotation (31) corresponded closely to those of barium 3-phosphoglycerate.

(31) O. Meyerhof and W. Schulz, Biochem. Z., 297, 60 (1938).

A mixture of the isolated material (Fraction II-a) and an equal weight of authentic barium 3-phosphoglycerate (32) was eluted from an anion exchange resin column which had previously been standardized with 3-phosphoglyceric acid. The standard and the mixed elution curves are shown in Fig. 6a,b. The somewhat low value of the optical rotation together with the barely noticeable displacement of the radioactivity elution curve might be taken to indicate the continued presence in Fraction II-a of a small amount of 2-phosphoglyceric acid of higher specific activity than the 3-phosphoglyceric acid.

We have thus isolated and obtained in a crystalline form at least one-third of the radioactivity fixed in 5 seconds as barium phosphoglycerate^W largely the 3-isomer. It is not unreasonable to suppose that the radioactivity left in the last four supernates (B,C,D,E) from the alcohol crystallizations is phosphoglyceric acid somewhat richer in the 2-isomer. The sum of these, together with that which was actually isolated, would account for over 65 percent of the total carbon fixed in 5 seconds. This closely corresponds to the distribution observed on a paper chromatogram of the same type of preparation. A small sample of phosphoglyceric acid Fraction II-a was chromatographed separately and came to the expected position.

Phosphoglyceric Acid Isolation Procedure: - Six grams of dry-old Synechococcus cells which had photosynthesized in 1.0 g. batches for 5 seconds was added to 18 g. of normal algae to provide sufficient phosphoglyceric acid for isolation. These algae were killed rapidly with acetic acid-hydrochloric acid (4:1) and all of the radioactivity was contained in the aqueous extract.

(32) Barium-3-phosphoglycerate was obtained through the courtesy of Prof. Carl Neuberg, New York University, New York.

The acid extract was concentrated (cold) to 9.7 ml. (1.6×10^6 c.p.m.) and neutralized to pH 10 with 1 N sodium hydroxide. The precipitate which formed was centrifuged and washed three times with water (12, 3.5 and 2.6 ml.) at pH 10. The combined washings and supernate (1.45×10^6 c.p.m.) were acidified with 1 N hydrochloric acid and 0.3 ml. of saturated barium chloride was added. The precipitate was washed twice with 1 N hydrochloric acid (1.5 ml.) and discarded. The combined washings and supernate was concentrated to 3.0 ml. and basified to pH 10 with 1 N sodium hydroxide. The small precipitate was discarded.

The supernate was diluted with 1.5 volumes of ethanol and centrifuged. The supernate, A, contained 0.2×10^6 c.p.m. The precipitate was washed with 1.0 ml. water (wash contained 0.05×10^6 c.p.m.) and was taken up in 4.3 ml. of 0.05 N hydrochloric acid and found to have 1.1×10^6 c.p.m. The small residue was centrifuged and discarded.

The acid solution was diluted to 8.6 ml. with ethanol and the precipitate centrifuged. The supernate, B, contained 0.23×10^6 c.p.m. The precipitate, dissolved in 2.0 ml. of 0.05 N hydrochloric acid, contained 0.78×10^6 c.p.m. Ethanol was again added to a total volume of 6.0 ml. and the precipitate was centrifuged. Supernate C contained 0.01×10^6 c.p.m. A solution of the precipitate in 1.5 ml. of 0.05 N hydrochloric acid was diluted with ethanol to 3.0 ml. and the precipitate centrifuged. The supernate, D, contained 0.2×10^6 c.p.m.

A solution of the precipitate in 0.9 ml. of 0.05 N hydrochloric acid was diluted with 0.5 ml. of ethanol to obtain Precipitate I; 3.9 mg., 0.19×10^6 c.p.m.

Anal. Calcd for $C_3H_5O_7PBa \cdot 2 H_2O$: P, 8.7%; Found, P, 8.6%.

Molar rotation (as phosphoglyceric acid) in the presence of molybdate ion: $-620^{\circ} \pm 50^{\circ}$. (Authentic value, -750°). Specific activity: 4.9×10^4 c.p.m./mg.

The supernate from Precipitate I was added to 1.5 ml. of ethanol. Precipitate II, 7 mg., 0.29×10^6 c.p.m., and supernate E, 0.08×10^6 c.p.m., were obtained. The solution of Precipitate II in 0.05 N hydrochloric acid diluted with 1.5 volumes of ethanol gave Precipitate II-a, containing 8.6 percent phosphorus and having a specific activity of 4.2×10^4 c.p.m./mg. The total activity in supernates B, C, D and E was 0.58×10^6 c.p.m. (36 percent of the total). The activity in Precipitates I and II was 0.48×10^6 c.p.m. (30 percent of the total).

Easily Hydrolyzed Phosphate Esters: - Readily hydrolyzed phosphate esters and inorganic phosphate can be recognized on the paper by spraying with suitable reagents to produce molybdenum blue. Glucose-1-phosphate, 3-phosphoglyceraldehyde and phosphopyruvic acid are easily detectable when 3 micrograms per cm^2 is present on the paper. Fructose-1,6-diphosphate is variable in its color development. This is probably caused by different degrees of hydrolysis during the spraying and drying processes.

Spray Test for Phosphates on Chromatograms: - The paper is sprayed consecutively with the following solutions (33): (a) Five grams ammonium molybdate dissolved in 100 ml. cold water and poured into 35 ml. concentrated nitric acid. Paper dried at 70° C; (b) Fifty milligrams of benzidine hydrochloride dissolved in 10 ml. concentrated acetic acid and diluted with water to 100 ml. Paper dried at room temperature; (c) Saturated sodium acetate solution. Paper dried at room temperature.

(33) Fritz Feigl, "Qualitative Analysis by Spot Tests," page 216, Norde-mann Publishing Company, New York, 1939.

Glucose-1-Phosphate: - The rate of hydrolysis of the photosynthetic intermediates in the middle one of the three phosphate spots to form radioactive glucose suggested the presence of glucose-1-phosphate. When the authentic compound containing P^{32} (34) was co-chromatographed with an algal extract, the P^{32} activity coincided with the C^{14} radioactive spot.

Phosphopyruvic Acid: - An authentic specimen of phosphopyruvic acid (35) was chromatographed and its position detected by molybdenum blue color. A photosynthetic intermediate found in this area was investigated. Acid hydrolysis formed pyruvic acid which was identified by its distribution coefficient between water and ether, solubility and volatility properties and the formation of a dinitrophenylhydrazone which co-crystallized with an authentic specimen.

A radioactive substance was eluted from a chromatogram of 5 seconds photosynthetic fixation and hydrolyzed ten minutes in 1 N hydrochloric acid. The active product was then volatile and required addition of alkali for reproducible direct counting. An aliquot part of the hydrolysate was assayed for pyruvaldehyde and the 2,4-dinitrophenylhydrazone was inactive after recrystallization from pyridine.

The distribution coefficient of the hydrolysate radioactivity between a water (pH 1) and ether was 0.15-0.18. The constant determined for low concentrations of synthetic labeled pyruvic acid is 0.16-0.18.

A solution of 135,000 c.p.m. of eluate and 10 microliters of pyruvic acid in 1 N hydrochloric acid was heated with 25 mg. 2,4-dinitrophenylhydrazine in a sealed tube for 30 minutes at 100°C. The hydrazone (theoretical yield, 39 mg.) was recrystallized from ethanol and acetic acid-water and found to have a constant specific activity of 900 c.p.m./mg. which indicated a quantitative hydrolysis to pyruvic acid.

complex and 6 mg. of 2,4-dinitrophenylhydrazine (37). The specific activity of the pure osazone (theoretical yield, 7.29 mg.) was 178 c.p.m./mg. This indicated that 1300 c.p.m., or all of the eluate activity, was triose phosphate.

Amino Acids: - Identification of radioactive amino acids by superposition of the radiogram on the chromatogram sprayed with ninhydrin has been described (5). In cases where the original separation is incomplete or where the amount of amino acid is insufficient to produce visible ninhydrin color, the radioactive spot or group of spots is eluted (17,18) and re-chromatographed with added carriers. The coincidence of shape and location of the radioactive spot with the colored spot produced by an authentic compound on the paper indicates the identity of the two.

Sugars: - Sucrose, fructose and glucose (38) are the only sugars which have been identified in the radiograms. Their chromatographic coordinates in phenol have been determined (10). The Roe test for fructose and the Tollens test for glucose (39) gave added confirmation of these identifications.

SUMMARY

1. Paper chromatography has been employed to separate the radioactive products formed during photosynthesis in $C^{14}O_2$.
2. The method has been used for the separation and identification of carboxylic acids and phosphates.esters.
3. The first observed product of carbon dioxide assimilation during photosynthesis has been isolated and shown to be phosphoglyceric acid.

(37) E. Baer and H.O.L. Fischer, J. Biol. Chem., 150, 223 (1943).

(38) Authentic specimens of these radioactive sugars were kindly supplied by Prof. W. Z. Hassid.

(39) S. M. Partridge, Nature, 158, 220 (1946).






FIG. 1 - CHROMATOGRAPHY TROUGH. FABRICATED FROM POLISHED STAINLESS STEEL, TYPE NO. 316



SCENEDESMUS
90 sec PS in 4% CO₂


CLP 784

BUTANOL - PROPIONIC ACID - WATER



PHENOL - WATER ←

FIG. 3 - RADIOGRAM OF PRODUCTS PHOTOSYNTHESIZED IN NINETY SECONDS BY SCENEDESMUS IN C¹⁴O₂. THE ALGAE WERE RAPIDLY KILLED BY POURING INTO FOUR VOLUMES OF BOILING ETHANOL. CONSIDERABLE CLARITY OF THE ORIGINAL RADIOGRAM IS LOST IN PHOTOGRAPHIC REPRODUCTION



P-LIPID

INORGANIC
P

TRIOSE - P

PGA
HMP

SCENEDESMUS
1 HOUR P³²

BUTANOL - PROPIONIC ACID - WATER

PHENOL - WATER ←

FIG. 4 - RADIOGRAM OF PRODUCTS OF ONE HOUR PHOTOSYNTHESIS BY SCENEDESMUS IN RADIOPHOSPHATE (P³²)




FIG. 5a, b - ONE DIMENSIONAL RADIOGRAMS OF P^{32} LABELED YEAST FERMENTATION INTERMEDIATES. SOLVENT: BUTANOL-ACETIC ACID-WATER. THE TWO MIXTURES WERE OBTAINED FROM ION EXCHANGE RESIN ELUATE FRACTIONS.





Fig. 6a,b.-a. Elution curve of authentic 3-phosphoglyceric acid from Dowex A-1 anion exchange of resin column. Eluant: 0.2 N sodium chloride. b. Elution curve of a mixture of equal amounts of authentic and isolated phosphoglyceric acids. ●, Total phosphorus analyses. ○, C^{14} radioactivity. The smaller peak represents inorganic phosphate and hexose monophosphates. The ordinate unit represents 100 $\mu\text{g.}$ of phosphorus/ml. of eluate and 20000 cpm./ml. of C^{14} radioactivity.