

The Physical and Molecular Properties of New Low Melting Nematics with Negative Dielectric Anisotropy

R. Dąbrowski, J. Jadżyn^a, J. Dziaduszek, Z. Stolarz, G. Czechowski^a, and M. Kasprzyk^a

Institute of Chemistry, Military University of Technology, Kaliskiego 2, 01-489 Warsaw, Poland

^a Institute of Molecular Physics, Polish Academy of Sciences, Smoluchowskiego 17, 60-179 Poznań, Poland

Reprint requests to Prof. J. J., Fax: +48-61 868-45-24

Z. Naturforsch. **54a**, 448–452 (1999); received May 8, 1999

The paper presents some basic physical properties (static electric permittivities, refraction indices, density and viscosity) of 2-chloro-4-*n*'-alkylphenyl esters of 4-*n*-alkylbicyclo[2,2,2]octane-1-carboxylic acids ($n'=7$, $n=5$ and 7) which are, at room temperature, nematics with a negative dielectric anisotropy. On the basis of temperature dependence of the principal static permittivities $\epsilon_{\parallel}(T)$ and $\epsilon_{\perp}(T)$ of the nematics, using the Maier-Meier equations, the angle between the dipole moment vector and the long axis of mesogenic molecules, the apparent molecular dipole moment square $\mu_{\text{app}}^2(T)$, and the nematic order parameter $S(T)$ were determined.

Key words: Nematics; Dielectric Anisotropy; Density; Viscosity.