



The “Plastisphere” of Biodegradable Plastics Is Characterized by Specific Microbial Taxa of Alpine and Arctic Soils

Joel Rütli^{1,2}, Damian Bölsterli¹, Lucrezia Pardi-Comensoli³, Ivano Brunner¹ and Beat Frey^{1*}

¹ Forest Soils and Biogeochemistry, Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Birmensdorf, Switzerland, ² Institute of Biogeochemistry and Pollutant Dynamics, Swiss Federal Institute of Technology, ETH Zürich, Zurich, Switzerland, ³ Laboratory for Mechanical Systems Engineering, Swiss Federal Laboratories for Materials Science and Technology (EMPA), Dübendorf, Switzerland

OPEN ACCESS

Edited by:

Montserrat Filella,
Université de Genève, Switzerland

Reviewed by:

Cristina Silva Pereira,
New University of Lisbon, Portugal
Ciro Sannino,
University of Perugia, Italy

*Correspondence:

Beat Frey
beat.frey@wsl.ch

Specialty section:

This article was submitted to
Toxicology, Pollution and the
Environment,
a section of the journal
Frontiers in Environmental Science

Received: 14 May 2020

Accepted: 28 August 2020

Published: 24 September 2020

Citation:

Rütli J, Bölsterli D,
Pardi-Comensoli L, Brunner I and
Frey B (2020) The “Plastisphere”
of Biodegradable Plastics Is
Characterized by Specific Microbial
Taxa of Alpine and Arctic Soils.
Front. Environ. Sci. 8:562263.
doi: 10.3389/fenvs.2020.562263

Plastic pollution poses a threat to terrestrial ecosystems, even impacting soils from remote alpine and arctic areas. Biodegradable plastics are a promising solution to prevent long-term accumulation of plastic litter. However, little is known about the decomposition of biodegradable plastics in soils from alpine and polar ecosystems or the microorganisms involved in the process. Plastics in aquatic environments have previously been shown to form a microbial community on the surface of the plastic distinct from that in the surrounding water, constituting the so-called “plastisphere.” Comparable studies in terrestrial environments are scarce. Here, we aimed to characterize the plastisphere microbiome of three types of plastics differing in their biodegradability in soil using DNA metabarcoding. Polylactic acid (PLA), polybutylene adipate terephthalate (PBAT), and polyethylene (PE) were buried in two different soils, from the Swiss Alps and from Northern Greenland, at 15°C for 8 weeks. While physico-chemical characteristics of the polymers only showed minor (PLA, PBAT) or no (PE) changes after incubation, a considerably lower α -diversity was observed on the plastic surfaces and prominent shifts occurred in the bacterial and fungal community structures between the plastisphere and the adjacent bulk soil not affected by the plastic. Effects on the plastisphere microbiome increased with greater biodegradability of the plastics, from PE to PLA. Copiotrophic taxa within the phyla Proteobacteria and Actinobacteria benefitted the most from plastic input. Especially taxa with a known potential to degrade xenobiotics, including Burkholderiales, Caulobacteriales, *Pseudomonas*, *Rhodococcus*, and *Streptomyces*, thrived in the plastisphere of the Alpine and Arctic soils. In addition, Saccharimonadales (superphylum Patescibacteria) was identified as a key taxon associated with PLA. The association of Saccharibacteria with plastic has not been reported before, and pursuing this finding further may shed light on the lifestyle of this obscure candidate phylum. Plastic addition affected fungal taxa to a lesser extent since only few fungal genera such as *Phlebia* and *Alternaria*

were increased on the plastisphere. Our findings suggest that the soil microbiome can be strongly influenced by plastic pollution in terrestrial cryoenvironments. Further research is required to fully understand microbial colonization on plastic surfaces and the biodegradation of plastic in soils.

Keywords: alpine soil, arctic soil, biodegradation, bioplastics, microbiome, plastisphere, *Saccharimonadales*

INTRODUCTION

Plastic pollution is a threat to ecosystems all over the globe, with clear implications for animals like birds, marine mammals and fish (Secretariat of the Convention of Biological Diversity, 2016; Thiel et al., 2018). Additionally, plastic pollution potentially causes food safety problems (Wright and Kelly, 2017; Smith et al., 2018), leads to developmental, reproductive and metabolic disorders in invertebrates through the uptake of nano-sized particles (Shen et al., 2019), and even contributes to the spread of pathogens and antibiotic resistance by serving as a raft for microorganisms (Keswani et al., 2016; Laganà et al., 2019). Most of the attention in research is paid to plastic debris in oceans, while knowledge about the occurrence, fate and potential hazards of landfill waste is scarce and its impact is likely underestimated considerably (de Souza Machado et al., 2018). Yet, some studies have examined the quantity of synthetic polymers in terrestrial ecosystems (Piehl et al., 2018; Scheurer and Bigalke, 2018; Zhang and Liu, 2018) and have documented detrimental effects of plastics in soils on earthworms (Huerta Lwanga et al., 2016) and microorganisms (Wang et al., 2016), as well as implications for agricultural productivity (Rillig et al., 2019). It is estimated that the amount of plastic released into terrestrial environments is 4–23 times higher than into marine environments (Horton et al., 2017). Plastics can reach habitats not only by direct littering and transport through rivers and oceans, but also by transport through the atmosphere and deposition in pristine regions such as alpine and Arctic environments (Allen et al., 2019). Plastic pollution has even been reported in remote areas of the Swiss Alps and the Arctic (Ambrosini et al., 2019; Bergmann et al., 2019).

A possible way to deal with the plastic pollution in the environment is to develop biodegradable plastics. However, they make up only around 1% of the total plastic currently being produced (PlasticsEurope, 2018). In addition, very little is known about the fate of biodegradable plastics in terms of microbial colonization and their degradability, in particular in soils from cold environments since new materials are usually tested for decomposition at temperatures above 20°C (Briassoulis and Dejean, 2010). In one study on the microbial degradation of plastic in temperate, marine sediment, there were no signs of biodegradation for both polyethylene (PE) and biodegradable carrier bags after a 3-month incubation at 10°C (Nauendorf et al., 2016). A second study showed a weight loss of 37% for biodegradable plastic bags incubated in agricultural soil in the lab for 3 months at 25°C (Accinelli et al., 2012). However, in the same study samples incubated in the field under natural conditions were far less prone to biodegradation.

Abbreviations: CA, contact angle; CI, carbonyl index; PAHs, polyaromatic hydrocarbons; PBAT, polybutylene adipate terephthalate; PE, polyethylene; PLA, polylactic acid.

Given that the rate of biodegradation of plastics, besides of intrinsic material parameters, largely depends on external factors like temperature, moisture, and UV radiation (O’Brine and Thompson, 2010; Wilkes and Aristilde, 2017), biodegradation in soils of cold regions is assumed to be rather slow. The resident microbiome might be another critical parameter determining the biodegradability of plastics in soils. The composition of the soil microbiome strongly depends on various climatic and soil variables (Fierer, 2017). Alpine and Arctic soils are low in carbon and nitrogen content, and knowledge of their microbial communities is limited (Frasson et al., 2015; Frey et al., 2016; Malard and Pearce, 2018; Adamczyk et al., 2019; Pontes et al., 2020; Pushkareva et al., 2020). Plastic deposition may introduce a carbon and energy source into oligotrophic soils, possibly leading to profound changes in the soil microbiome and its functions in alpine and Arctic ecosystems. However, it has not been investigated whether the enzymatic ability to degrade plastics is pervasive in all soils or whether, especially in remote soils that have only recently been affected by pollution, plastic-degrading microorganisms are lacking.

The term “plastisphere” was introduced to describe the diverse microbial community influenced by plastic surfaces, analogous to the rhizosphere surrounding plant roots (Zettler et al., 2013; Amaral-Zettler et al., 2020). Since then, researchers have analyzed the composition of this novel habitat in many studies from aquatic environments and have shown that the plastisphere selects for particular microbial communities that differ from those in the surrounding environment (De Tender et al., 2015; Oberbeckmann et al., 2018; Ogonowski et al., 2018; Kirstein et al., 2019) and from communities on other inert surfaces like glass (Kirstein et al., 2018). However, data about the plastisphere microbiome in terrestrial ecosystems are scarce, and to our knowledge plastic-colonizing microbes in soils from cold environments have not previously been studied with culture-independent approaches to cover the entire microbial community on the plastic surface.

In the present mesocosm experiment, we buried pieces of three different plastic types in two different top soils, from the Swiss Alps and from Northern Greenland, at 15°C for 8 weeks. We used polylactic acid (PLA), polybutylene adipate terephthalate (PBAT), and PE, plastics that vary in their biodegradability. We characterized the plastisphere microbiome of the three types of plastics using soil DNA metabarcoding. We aimed to determine: (1) whether different polymer types are colonized by different microbial communities; (2) whether the plastisphere microbiome is different in the two soils; and (3) whether the plastisphere microbial communities can be linked to the biodegradability of the tested plastics. We hypothesized that the plastic material would determine the associated plastisphere microbiome, and that shifts in the plastisphere microbial

communities compared to in bulk soil would be smallest in PE, which has the lowest biodegradability.

MATERIALS AND METHODS

Plastic Types

Three different types of plastic were used in the experiments. The first specimen, purchased at Coop AG (Basel, Switzerland), was a compostable plastic bag made of Ecovio, a plastic blend primarily made of PLA. The second specimen was an agricultural mulch foil, purchased at Oeremansplastic (Genderen, Netherlands), made of PBAT. While PLA is made of renewable plant sources, PBAT is fabricated from fossil fuels. The third specimen was a common non-biodegradable waste bag made of low-density PE, purchased from TopPac (Schwarzenbach, Switzerland). The exact composition of the plastics was not known, as they are commercially available products. The plastics were chosen based on their different expected biodegradabilities, ranging from high biodegradability (PLA) to medium biodegradability (PBAT) to very low biodegradability (PE). The plastic surfaces were not sterilized to avoid chemical changes of the polymers. However, we surveyed the surfaces of non-incubated plastic pieces for the presence of indigenous microorganisms by plating 4 × 4 cm pieces of plastic on R2A (Carl Roth GmbH + Co., Karlsruhe, Germany) and LB (Merck KGaA, Darmstadt, Germany) agar plates and were only able to obtain 0–2 colonies per piece of plastic.

Soils Used for Incubation

We used two different top soils (0–10 soil depth), one from an alpine and one from an Arctic environment. The alpine soil was collected on the northwestern flank of “Muot da Barba Peider” (Barba Peider) in the eastern Swiss Alps at 2979 m a.s.l. (Frey et al., 2016). The Arctic soil was sampled in Villum

at Station Nord in Northern Greenland. The coordinates, soil characteristics and climatic parameters of the two soils used are given in **Table 1**. We sampled three independent locations (field replications) for each soil type and replicated them in the mesocosm experiment (see below). Soils were transported in cold boxes to the laboratory and stored in closed plastic bags at 4°C in the dark until the start of the incubation experiment (less than 3 months).

Experimental Set-Up

Half of the soil collected from each site was autoclaved (1 × at 120°C for 20 min) to produce sterile controls for detecting possible abiotic degradation of the plastics. Sterilization of the investigated soils was confirmed by the lack of cultivable microorganisms. Incubation mesocosms were then prepared using 100 ml beaker glasses containing approximately 80 g of soil, with three replicates for each soil origin (Barba Peider, Villum), treatment (natural soil, autoclaved control) and plastic type (PLA, PBAT, PE) (36 beakers in total). Prior to incubation, the plastic pieces (4 × 4 cm) were quickly submerged in a soil suspension for 1 min to ensure contact of the indigenous microbial community of the respective soil with the plastic surface. The suspension was prepared by shaking 10 g of the corresponding soil in 40 ml of sterile dH₂O for 60 min. Plastic pieces for sterile controls were submerged in sterile dH₂O instead of the suspension. The beaker glasses were covered by parafilm and incubated at 15°C in the dark for 8 weeks. The incubation temperature represents the maximum temperature in the topsoils at both sites during snow-free season. The moisture content was determined gravimetrically and balanced once a week by adding sterile dH₂O.

At the end of the incubation experiment, samples were prepared for further processing. Bulk soil was collected from a distance of >2 cm to the incubated plastic sample. For the plastisphere samples, small soil particles adherent to the plastic piece were manually scratched off and added to the extraction tube. In addition, 4–5 small pieces of plastic (1 × 2 mm) were added to the extraction tube. For bulk soil samples approximately 250 mg fresh weight was used for DNA extraction, while for plastisphere samples only 100 mg fresh weight could be used due to the low mass of soil adherent to the plastic surfaces. Samples were added into extraction tubes and were frozen at –20°C until DNA extraction was performed. Small pieces of plastic (1 × 2 mm) were added to 5 ml Falcon tubes containing sterile tap water for cultivation of microorganisms. The remaining material from each plastic piece was rinsed with sterile dH₂O to provide a clean surface and dried at 35°C for 4 days prior to the physico-chemical analysis.

Cultivation of Microorganisms

The 5 ml Falcon tubes containing the plastic pieces were shaken with a vortex for 10 min at half speed. The resulting cell suspensions were serially diluted 1:10, 1:100, and 1:1000 in sterile tap water. All dilutions were plated in duplicate on R2A agar plates (Carl Roth GmbH + Co., Karlsruhe, Germany). The agar plates were incubated at 15°C for 2 months and growing colonies were picked once a week with a sterile tooth pick. Strains were phylogenetically characterized by sanger Sequencing as recently

TABLE 1 | Physico-chemical and site characteristics of soils used in the microcosm experiment.

Properties	Barba Peider	Villum
C [%]	0.14 ± 0.01*	2.47 ± 2.30*
N [%]	<0.02	0.25 ± 0.14*
pH [H ₂ O]	6.5 ± 0.1*	6.9 ± 0.3*
Sand [%]	80.5	88.9
Silt [%]	15.9	7.7
Clay [%]	3.6	3.4
Mean annual soil temperature [°C]	–1.8	–5.4
Minimal soil temperature [°C]	–13.5	–16.3
Maximal soil temperature [°C]	21.3	21.7
Annual precipitation [mm]	1500	188
Elevation [m a.s.l.]	2979	24
Vegetation	Sparsely vegetated (<i>Poa</i> , <i>Cerastium</i> , <i>Jacobea</i>)	Crusts and sporadic <i>Saxifraga arctica</i>
Coordinates	46° 49' 59.75" N, 9° 93' 14.30" W	81° 36' 5.26" N, 16° 39' 43.31" W

*Values represent means ± standard deviations (n = 9 for C and N, n = 3 for pH).

outlined (Lapanje et al., 2012; Brunner et al., 2018). Briefly, the 16S rDNA (bacteria) and ITS (fungi) regions of isolated organisms were amplified by colony PCR. For bacteria universal primers 27F (Heuer et al., 1997) and 907R (Muyzer et al., 1995) were used. For fungi, primers ITS1 and ITS4 (White et al., 1990) were used. PCR products were sent to Macrogen B.V. (Amsterdam, Netherlands) for sequencing. Retrieved sequences were trimmed to the high quality portion and closest related sequences were searched on the NCBI nucleotide database with the blastn algorithm and default parameters.

Light Microscopy of Plastic Surfaces

Washed plastic pieces were analyzed with a VHX-500f digital microscope (Keyence International, Mechelen, Belgium). Images were acquired with a 5× and a 50× magnification.

Physico-Chemical Analysis

Fourier-transform infrared spectroscopy (FTIR) was performed with a Tensor 27 spectrometer (Bruker, Billerica, MA, United States). Spectra were obtained with a diamond Attenuated Total Reflectance (ATR) crystal plate. For each sample, 32 scans were recorded with wavelengths in the range of 4000–650 cm^{-1} for the analysis of Barba Peider soil and 4000–320 cm^{-1} for Villum soil. The scans were recorded with a spectral resolution of 4 cm^{-1} , and the resulting spectra averaged. OPUS software (Winterthur, Switzerland) was used to collect and correct the data (baseline correction). As a standard, three measurements per replicate were performed. Carbonyl indices were used to evaluate the degradation of plastics by semi-quantitative data. The indices were calculated with the OPUS software (Winterthur, Switzerland) using the height of the peaks (K algorithm). With this technique it is possible to calculate the ratio between C = O and C-H chemical bonds ($AC = O/AC-H$). Depending on the plastic type, different wavelengths were chosen for the carbonyl index calculations ($CI_{PLA} = A_{1713}/A_{1456}$; $CI_{PBAT} = A_{1713}/A_{729}$). No carbonyl index calculations were possible for PE because no C = O bonds were detected in the incubated samples.

Contact angle analysis was used to determine the wettability of plastic surfaces by measuring the contact angle of a sitting drop. For this purpose a Drop Shape Analyzer DSA30 (Krüss, Hamburg, Germany) was used. A 2 μl drop of distilled water was placed on the plastic surface, and the contact angle was determined using the Krüss Advance software package with the Ellipse fitting method. The contact angle was measured ten times per drop at three different locations per replicate. The values obtained for the ten drops per location were averaged in advance to statistical analysis. This method was applied for all plastic types. Cleaned specimens from the mesocosm experiments were compared to plastic controls without prior incubation in the soils and to the controls incubated in sterilized soil.

DNA Extraction, PCR Amplification, and Illumina MiSeq Sequencing

DNA extraction was performed with the DNeasy PowerSoil kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol. DNA was quantified using the QubitTM high-sensitivity assay for double stranded DNA (Thermo Fisher Scientific,

Waltham, MA, United States). DNA concentrations for sterile controls were below the detection limit. The V3–V4 region of the prokaryotic small-subunit (16S) rRNA gene (bacteria and archaea) and the internal transcribed spacer region 2 (ITS2) of the eukaryotic (fungal groups and some groups of protists and green algae) ribosomal operon were PCR amplified from 5 ng of DNA template, using primers and conditions previously described (Frey et al., 2016). PCRs of individual samples were run in duplicate, pooled, and purified using Agencourt Ampure XP (Beckman Coulter). Prokaryotic and fungal amplicon pools were sent to the Génome Québec Innovation Centre at McGill University (Montreal, QC, Canada) for barcoding using the Fluidigm Access Array technology (Fluidigm) and paired-end sequencing on the Illumina MiSeq v3 platform (Illumina Inc., San Diego, CA, United States).

Sequence Quality Control, OTU Clustering, and Taxonomic Assignments

Quality filtering, OTU clustering, and assignment were conducted similarly to methods described previously (Frey et al., 2016; Luláková et al., 2019) using a customized pipeline based on UPARSE (Edgar, 2013). Briefly, paired-end reads were merged using the fastq_mergepairs algorithm (Edgar and Flyvbjerg, 2015), filtering for sequences with a minimum length of 300 bp (16SV3V4) or 200 bp (ITS2) and a minimum overlap of 50 bp. PCR primers were removed using Cutadapt (Martin, 2011) allowing for a maximum of one mismatch in the forward and reverse primer. Next, reads were quality filtered using the USEARCH fastq_filter function, discarding reads with an expected error of one or greater. De-replicated sequences were clustered into OTUs at 97% sequence identity using the cluster_otu function. The clustering step includes an “on the fly” chimera removal algorithm. The OTU centroid sequences were then filtered for the presence of ribosomal signatures using Metaxa2 (Bengtsson-Palme et al., 2015) or ITSx (Bengtsson-Palme et al., 2013). Subsequently, sequences were mapped back on the OTU centroid sequences and taxonomic classification was conducted using a naïve Bayesian classifier (Wang et al., 2007) implemented in Mothur (Schloss et al., 2009) with a minimum bootstrap support of 0.6. Prokaryotic 16SV3V4 sequences were queried against the SILVA v132 database (Quast et al., 2013), whereas eukaryotic ITS2 sequences were queried against the fungal ITS database UNITE v8.0 (Nilsson et al., 2019). Prokaryotic sequences identified as originating from organelles (chloroplast, mitochondria), as well as eukaryotic sequences identified as originating from soil animals (metazoa) or plants (viridiplantae) were not included in the analyses. Raw sequences were deposited in the NCBI Sequence Read Archive under the accession number PRJNA630025.

Statistical Analyses

Results from all statistical tests performed in this study were considered significant at $P < 0.05$ unless indicated otherwise. Unless indicated otherwise, all figures were created with the ggplot2 package (Wickham, 2016) implemented in R Development Core Team (2008). Differences in contact angle and carbonyl index data were assessed using factorial ANOVA

and Tukey's HSD *post hoc* test, implemented in R. OTU richness (S_{obs}) and Shannon index (H') were retrieved with the *estimate_richness* function implemented in the phyloseq package in R (McMurdie and Holmes, 2013). Differences in soil S_{obs} and H' data were assessed using factorial ANOVAs and paired *t*-tests implemented in StatView, 2nd edition (SAS Institute Inc., Cary, NC, United States).

Principal coordinate analysis (PCoA) of microbial β -diversity was performed using the *ordinate* function based on Bray-Curtis dissimilarities and plotted using the *plot_ordination* function, both implemented in the phyloseq package in R (McMurdie and Holmes, 2013). Differences in β -diversity were tested by permutational multivariate analysis of variance (Anderson, 2001) and pairwise PERMANOVA functions in PRIMER version 7 (Clarke and Gorley, 2006) based on 10^5 permutations (Hartmann et al., 2017). For the pairwise tests, the Monte Carlo approximated level of significance [P (MC)] was determined.

Relative abundances of the most abundant taxa were assessed in R by agglomerating OTUs on phylum and order levels using the *tax_glom* function implemented in the phyloseq package (McMurdie and Holmes, 2013), normalizing the abundances using the function *normalise_data* from the microbiomeSeq package (Ssekagiri, 2020) and using the function *aggregate_top_taxa* implemented in the microbiome package in R (Lahti and Shetty, 2019). Differences in relative abundance of the most abundant taxa were assessed using factorial ANOVAs and Tukey's HSD *post hoc* tests implemented in R. Relative abundances of taxa were plotted using the function *plot_taxa* implemented in the microbiomeSeq package (Ssekagiri, 2020).

Differential abundance analysis was performed using the *differential_abundance* function implemented in the microbiomeSeq package in R (Ssekagiri, 2020). First, the analysis was carried out using OTUs as a taxonomic classifier. In a second analysis, OTUs were aggregated to the next highest classifiable taxonomic rank for bacteria (mostly genus). For fungi, only sequences classifiable to the genus level were considered in the second analysis. In order to assess which OTUs/taxa were significantly affected by plastic type, we calculated \log_2 -fold changes relative to the bulk samples using DESeq2 (Love et al., 2014). OTUs and taxa were considered significantly different (Wald test) between plastisphere and bulk soil samples if the false discovery rate (adjusted P) was < 0.05 .

RESULTS

Physico-Chemical Measurements of Polymer Degradation

After 8 weeks of incubation in the Alpine (Barba Peider) or Arctic (Villum) soil, no clear signs of degradation were observed in any of the three plastic types (PLA, PBAT, and PE). However, PLA incubated in natural soil was covered with white stains, some of which appeared cloudy and others mycelium-like (Supplementary Figure 1).

Carbonyl indices of PLA and PBAT were significantly affected by both incubation and soil type without interactions (Table 2a). The contact angle of PBAT was significantly affected

by incubation, soil type and their interactions, whereas PE was not influenced by incubation or soil type. In contrast, PLA was affected by soil type and interactions between incubation and soil type (Table 2a).

In Barba Peider soil, the contact angle of PLA incubated in natural-soil was significantly different than that incubated in sterile soil (Table 2b). However, neither of these incubated PLAs differed from the non-incubated PLA regarding contact angle. For PLA in Villum soil, only the difference between the carbonyl index of sterile-soil incubated and non-incubated plastics was statistically significant. However, the carbonyl index of PLA incubated in natural soil was very similar to that of the sterile-soil incubated samples but lower than the non-incubated PLA ($P = 0.059$).

In Barba Peider soil, the carbonyl index and the contact angle of PBAT from the 8-week incubation in natural soil were significantly different from the non-incubated sample, as well as from PBAT incubated in sterile soil (Table 2b). In contrast, the carbonyl index of natural-soil and sterile-soil incubated samples in Villum soil was significantly different from non-incubated PBAT, but no difference was detected between natural-soil and sterile-soil incubated PBAT. No difference was detected for the contact angle of non-incubated, sterile-soil incubated and natural-soil incubated PBAT.

Polymer-Dependent Microbiome

Sequencing of Barba Peider samples yielded a total of 1,169,088 ($64,949 \pm 4,606$ per sample; mean \pm standard error) prokaryotic 16S_{V3V4} and 335,637 ($18,647 \pm 394$) eukaryotic ITS2 high-quality sequences that were clustered into 2,360 and 334 OTUs, respectively. Samples from Villum yielded a total of 412,004 ($22,889 \pm 851$ per sample) prokaryotic 16S_{V3V4} and 1,007,107 ($55,950 \pm 3,404$) eukaryotic ITS2 high-quality sequences that were clustered into 2,918 and 280 OTUs, respectively.

Bacterial and Fungal α -Diversity

We estimated the prokaryotic and fungal α -diversity of plastisphere and bulk soil samples in Barba Peider and Villum soils by analyzing the observed OTU richness (S_{obs}) and Shannon indices (H'). ANOVA of plastisphere and bulk soil samples showed significant differences for all α -diversity measures except for prokaryotic S_{obs} in Barba Peider soil (Table 3a). By pairwise comparisons of plastisphere and bulk soil of each plastic and soil type, we observed a significant reduction in prokaryotic H' in both soils in the PLA plastisphere when compared with the bulk soil (Table 3b). Similar trends were observed for S_{obs} (Figure 1). In addition, prokaryotic S_{obs} was significantly lower in the PE plastisphere when compared with the bulk soil in the Villum samples. Fungal α -diversity was significantly lower in the PLA plastisphere than in the bulk soil in samples from Barba Peider, and the same trend was true for Villum samples (Figure 2). Fungal α -diversity was lower in the plastisphere of PBAT compared with the bulk soil in the Villum samples, but no such effects of PBAT were found for Barba Peider samples. Furthermore, fungal S_{obs} of the plastisphere of PE was lower compared with the bulk soil in Villum samples.

TABLE 2 | Physico-chemical characterization of plastics before and after an 8-week incubation in autoclaved (sterile) and natural (non-sterile) soil from Barba Peider or Villum.

(a)							
Plastic type	Carbonyl index (CI)			Contact angle (CA)			
	Inc.	Soil	Inc. x Soil	Inc.	Soil	Inc. x Soil	
PLA	3.65*	30.1***	1.19 ^{ns}	3.13 ^(ns)	10.8**	8.04**	
PBAT	30.3***	38.9***	2.25 ^{ns}	9.67***	5.55*	11.2**	
PE	NA	NA	NA	1.76 ^{ns}	3.02 ^(ns)	1.63 ^{ns}	
(b)							
Plastic type	Measure	Soil	non-incubated	Barba Peider		Villum	
				Incubation	P	Incubation	P
PLA	Carbonyl index	Sterile	8.25 ± 0.14	8.41 ± 0.32	0.995	7.60 ± 0.17	0.005
		Natural	8.25 ± 0.14	8.28 ± 0.49	1.00	7.74 ± 0.18	0.059
	Contact angle	Sterile	89.6 ± 6.9	104.2 ± 9.4	0.286	80.7 ± 12.8	0.540
		Natural	89.6 ± 6.9	83.0 ± 5.8	0.846	78.9 ± 4.8	0.298
PBAT	Carbonyl index	Sterile	1.26 ± 0.01	1.26 ± 0.01	1.00	1.16 ± 0.02	<0.001
		Natural	1.26 ± 0.01	1.18 ± 0.05	<0.001	1.11 ± 0.04	<0.001
	Contact angle	Sterile	107.2 ± 9.1	108.7 ± 1.1	1.00	100.4 ± 6.2	0.883
		Natural	107.2 ± 9.1	82.8 ± 13.9	<0.001	99.3 ± 5.9	0.789
PE	Carbonyl index	Sterile	NA	NA	–	NA	–
		Natural	NA	NA	–	NA	–
	Contact angle	Sterile	87.8 ± 14.0	96.2 ± 10.7	0.907	95.9 ± 5.5	0.703
		Natural	87.8 ± 14.0	88.6 ± 8.2	1.00	97.5 ± 6.9	0.495
				0.923		1.00	

(a) Effects of the type of incubation (non-incubated, incubated in sterile soil, and incubated in natural soil), soil origin (Barba Peider and Villum), and the interaction between the two variables based on analysis of variance (ANOVA). *F*-values are given and asterisks indicate significant differences, with ****P* < 0.001, ***P* < 0.01, **P* < 0.05, ^(ns)*P* < 0.1, and ^{ns}*P* > 0.1. (b) Results of pairwise comparisons (Tukey HSD tests) between non-incubated plastics and those incubated in sterile or natural soil. *P*-values (*P* < 0.05 in bold; 0.1 > *P* > 0.05 in italics) and means ± standard deviations are given. *P*-values in the third row of each comparison indicate significance of differences between incubated sterile and natural soils. The comparison of non-incubated and sterile incubated soils indicates the abiotic effect of soils on plastics, the comparison of non-incubated and natural incubated soils indicates the combination of abiotic and microbial factors affecting the plastics, and the comparison of sterile and natural incubated soils indicates the microbial component of degradation. NA, Not applicable, Inc., incubation.

Bacterial and Fungal β-Diversity

Principal coordinate analysis and permutational analysis of variance (PERMANOVA) revealed that microbial community structures in the PLA plastisphere microbiome were significantly distinct from the microbial communities in the bulk soil from both sampling sites (Figure 3 and Table 4). In addition, PBAT plastisphere microbial community structures were significantly distinct from the microbiome in the bulk soil in Villum samples, but this was not true for samples from Barba Peider. In contrast, PE plastisphere community structures did not differ significantly from the bulk soil in samples from both sites. Bulk soil in samples with the different plastic types did not differ significantly regarding community structures (data not shown).

“Plastisphere Taxa”

Relative Abundances of Microbial Taxa

The search for taxa at the phylum and order level that are affected by the plastics revealed that most community

structural changes occurred in the PLA plastisphere at both taxonomic levels and in both soils (Figure 4, Table 5, and Supplementary Figure 2). Bacterial phyla that increased in relative abundance in the plastisphere of PLA were Actinobacteria, Proteobacteria and Patescibacteria in Barba Peider and only Patescibacteria in Villum. Mainly Acidobacteria, Chloroflexi, Gemmatimonadetes, Planctomycetes, and Nitrospirae decreased in relative abundance in the PLA plastisphere in both soils. β-Proteobacteriales, Rhizobiales, and Saccharimonadales (superphylum Patescibacteria) increased in abundance in the plastisphere of Barba Peider, whereas in Villum only the latter was enriched. Archaea were not affected by any of the plastic types. Fungal phyla and orders were largely unaffected by PLA (Table 5 and Supplementary Figures 3, 4). The only changes in the plastisphere of PLA in Barba Peider soil were a decrease in the phylum Mortierellomycota and an increase in the order Thelebolales. The only phyla enriched in the

TABLE 3 | α -diversity of prokaryotes and fungi in the plastisphere and bulk soil of Barba Peider and Villum soils.

Main test	Barba Peider				Villum			
	Richness (S_{obs})		Shannon (H')		Richness (S_{obs})		Shannon (H')	
	F	P	F	P	F	P	F	P
Prokaryotes	1.8	0.196	5.9	0.006	11.2	<0.001	35.8	<0.001
Fungi	23.3	<0.001	8.9	<0.001	30.1	<0.001	12.7	<0.001

Pairwise test	Barba Peider		Villum	
	t	P	t	P
Prokaryotes				
PLA	-3.8	0.062	-4.3	0.050
PBAT	0.3	0.786	-2.1	0.165
PE	0.2	0.838	-1.8	0.213
Fungi				
PLA	-31.2	0.001	-8.4	0.014
PBAT	-0.7	0.580	-1.7	0.24
PE	-3.0	0.097	1.4	0.289

(a) Differences between plastisphere and bulk soil samples using analysis of variance (ANOVA). F- and P-values ($P < 0.05$ in bold; $0.1 > P > 0.05$ in italics) of the observed species richness and the Shannon diversity are given. (b) Pairwise comparisons between the plastisphere and bulk soil of each plastic type using paired t-tests. P- and t-values for S_{obs} and H' are presented. Negative t-values indicate smaller means in the plastisphere compared with the bulk soil. α -diversity for prokaryotes and fungi are shown in **Figures 1, 2**, respectively.

plastisphere of PBAT were Proteobacteria in Villum soil and Mortierellomycota in Barba Peider soil (**Table 5** and **Supplementary Figures 2, 3**). In Villum, only the orders β -Proteobacteriales and Rhizobiales were significantly increased in relative abundance in the plastisphere of PBAT, whereas only Helotiales were favored in Barba Peider (**Figure 4**, **Table 5**, and **Supplementary Figure 3**). In the PE plastisphere of both soils, the only significant increase in relative abundance was observed for Actinobacteria and the associated order of Propionibacteriales (**Figure 4**, **Table 5**, and **Supplementary Figure 2**).

Specific “Plastisphere Taxa”

Differential abundance analysis was performed by comparing total abundances of OTUs in plastisphere and bulk soil samples. The number of differentially abundant OTUs for both prokaryotes and fungi in both soils was highest for PLA, followed by PBAT and lastly PE (**Supplementary Table 1**). Whereas more bacterial OTUs showed positive \log_2 -fold changes in the plastisphere compared with bulk soils, a larger portion of the fungal OTUs in Villum soil showed negative \log_2 -fold changes in the plastisphere compared with bulk soils.


A repetition of the differential abundance analysis with OTUs aggregated to the lowest classifiable taxonomic level (mostly genus) revealed the bacterial taxa most strongly affected by the different plastics, as shown in **Figures 5–7**. Most taxa with significant \log_2 -fold changes belonged to the phyla Proteobacteria and Actinobacteria, and only few exceptions were identified (Bacteroidetes, Verrucomicrobia, and

Patescibacteria). No archaeal taxa were significantly influenced by the plastics.

Bacteria

Most taxa with positive \log_2 -fold changes in the plastisphere of PLA compared with bulk soils belonged to the phyla Actinobacteria and Proteobacteria, and a few belonged to Bacteroidetes, Patescibacteria, and Verrucomicrobia. *Nocardia* and *Saccharimonadales* exhibited positive \log_2 -fold changes in the plastisphere of both soils. Several genera increased substantially in the plastisphere of PLA in Barba Peider soil, where Actinobacteria (e.g., *Streptacidiphilus*, *Catenulispora*), Proteobacteria (e.g., *Collimonas*, *Rhizobiaceae*, *Variovorax*, *Pseudomonas*) and Verrucomicrobia (*Luteolibacter*) dominated (**Figure 5A**). Furthermore, strong positive \log_2 -fold changes in the PLA plastisphere of Villum soil compared with bulk soils were observed for genera of Actinobacteria (e.g., *Nocardioides*, *Streptomyces*), Proteobacteria (e.g., *Caulobacter*, *Brevundimonas*, *Sphingorhabdus*), and Bacteroidetes (e.g., *Dyadobacter*, *Ohtaekwangia*) (**Figure 5B**).

Taxa with positive \log_2 -fold changes in the plastisphere of PBAT, as with PLA, belong to Actinobacteria and Proteobacteria, with a few belonging to Bacteroidetes and Verrucomicrobia. Positive \log_2 -fold changes on PBAT in both soils were observed for *Nocardia*. Furthermore, genera within Actinobacteria (e.g., *Rhodococcus*, *Umezawaea*), Proteobacteria (e.g., *Collimonas*, *Pseudomonas*, *Variovorax*, *Mesorhizobium*), Bacteroidetes (*Sediminibacterium*) and others had positive \log_2 -fold changes in the PBAT plastisphere of




Barba Peider soil (Figure 6A). Genera of Actinobacteria (e.g., *Actinocorallia*, *Streptomyces*) and Proteobacteria (e.g., *Aquabacterium*, *Acidovorax*, *Caulobacter*, *Brevundimonas*) showed positive \log_2 -fold changes on the PBAT surface of Villum soils (Figure 6B).

Mostly genera of the phyla Actinobacteria and Proteobacteria showed positive \log_2 -fold changes in the plastisphere of PE. The genera *Nocardia* and *Nocardioideis* (Actinobacteria) had positive \log_2 -fold changes on PE in both soils. In addition, *Rhodococcus* (Actinobacteria) showed a positive \log_2 -fold change in the plastisphere of PE in Barba Peider soil (Figure 7A), and *Streptomyces*, *Aeromicrobium* (Actinobacteria), *Aquabacterium*, *Brevundimonas*, *Sphingorhabdus* and *Solimonas* (Proteobacteria) had positive \log_2 -fold changes on PE in Villum soil (Figure 7B).

Fungi

Even though the proportion of differentially abundant fungal OTUs in Villum soil was very high (Supplementary Table 1), most of the found OTUs were non-classifiable at low taxonomic ranks. Of the fungal OTUs in Villum soil, 43% were, e.g., only classifiable to the phylum level. Only fungi classifiable to the genus level were included in the subsequent analysis (36% of OTUs in Villum soil) (Supplementary Figures 5–7). Fungi with differential abundances in the plastisphere compared with the bulk soils mostly belonged to Ascomycota, with the exception of the plastisphere of PE in Villum soil, where more fungal taxa showing an increase belonged to Basidiomycota.

Generally, observed effects were much stronger in Villum than in Barba Peider soil for PLA. In Barba Peider soil, the genera *Oidiodendron* and *Kabatiella* had the greatest positive



\log_2 -fold changes in the plastisphere of PLA (**Supplementary Figure 5A**). In Villum soil, we found positive \log_2 -fold changes for *Mycosphaerella*, *Alternaria* and *Mycoarthritis* in the plastisphere of PLA (**Supplementary Figure 5B**).


The only classifiable genus showing a positive \log_2 -fold change in the plastisphere of PBAT in Barba Peider soil was *Trichocladium* (**Supplementary Figure 6A**). Whereas the genus *Pseudogymnoascus* had the greatest positive \log_2 -fold change in the plastisphere of PBAT in Villum soil, it exhibited a negative \log_2 -fold change on the PLA surface in Barba Peider soil (**Supplementary Figure 6**). Many fungal genera were found to have negative \log_2 -fold changes especially in the plastisphere of PBAT in Villum soil (**Supplementary Figure 6B**).

The fungal genus *Mycena* was found to be strongly increased in abundance in the plastisphere of PE in Villum soil

(**Supplementary Figure 7**). However, the enrichment of *Mycena* was only observed in one of the three replicates, and the genus was additionally not detected in the bulk soil samples. Other genera with significant positive \log_2 -fold changes in the PE plastisphere in Villum soil belonged to Basidiomycota (i.e., *Phlebia* and *Tricholoma*) or Ascomycota (i.e., *Aspergillus* and *Colletotrichum*) (**Supplementary Figure 7**). No significant differences between the plastisphere of PE and the bulk soil in Barba Peider samples were found.

Cultivable “Plastisphere” Strains

Eighty bacterial and 14 fungal strains were isolated from the plastic surfaces (**Supplementary Table 2**). Bacterial isolates belonged to 28 different genera within the phyla Actinobacteria, Bacteroidetes and Proteobacteria. Fungal isolates belonged to four different genera within the phyla



Ascomycota and Mortierellomycota. A large part of the cultivated microorganisms (i.e., *Streptomyces*, *Rhodococcus*, *Pseudomonas*, *Variovorax* and *Pseudogymnoascus*) were found to be increased on the plastisphere by differential abundance analysis.

DISCUSSION

In the present study, we linked FTIR and contact angle measurements of plastics incubated in soils with shifts in the resident microbiome. To our knowledge, this is the first experimental study comparing the effects of different plastic types on the plastisphere microbiome in alpine and Arctic soils. Increasing plastic pollution in these regions due to increased human activities highlights the need for pollution control and clean-up suitable for cold regions. The biotechnological potential of terrestrial cryoenvironments is strongly linked to patterns of taxonomical and functional diversity. Microorganisms in cold environments are known to

harbor special adaptations, and particular types of genes are only in these environments present (Shi et al., 2015; da Silva et al., 2017). Our findings suggest that plastic debris provides a habitat for complex microbial assemblages that differ from those in bulk soil. The observed effects of the plastic on the microbiome were strongly dependent on the biodegradability of the plastics and were most pronounced for PLA, followed by PBAT and lastly PE.

Physico-Chemical Measurements of Polymer Degradation

Overall, analyses of the biodegradable (PLA and PBAT) and non-biodegradable (PE) plastics showed no or only weak signs of degradation after 8 weeks (initial stage of degradation) of incubation in Alpine (Barba Peider) and Arctic (Villum) soils at 15°C. The contact angle and FTIR analyses confirmed that PE is almost non-biodegradable. FTIR data allow to conclude that no oxidation of the PE occurred. Oxidation of the C-C backbone of PE is considered an important

TABLE 4 | Changes in prokaryotic and fungal community structures in the plastisphere.

Main test	Barba Peider				Villum			
	Prokaryotes		Fungi		Prokaryotes		Fungi	
	<i>Pseudo F</i>	<i>P</i>	<i>Pseudo F</i>	<i>P</i>	<i>Pseudo F</i>	<i>P</i>	<i>Pseudo F</i>	<i>P</i>
	3.665	<0.001	3.259	<0.001	4.314	<0.001	3.383	<0.001
Pairwise test	<i>t</i>	<i>P</i> (MC)	<i>t</i>	<i>P</i> (MC)	<i>t</i>	<i>P</i> (MC)	<i>t</i>	<i>P</i> (MC)
PLA	2.758	0.008	2.538	0.013	2.914	0.007	1.798	0.050
PBAT	1.731	<i>0.056</i>	1.381	0.145	1.852	0.045	2.533	0.017
PE	1.235	0.226	1.276	0.207	1.688	<i>0.058</i>	1.207	0.263

Pseudo F- and *P*-values ($P < 0.05$ in bold; $0.1 > P > 0.05$ in italics) for the differences between plastisphere and bulk soil samples and the three plastic types regarding prokaryotic and fungal β -diversity were assessed by permutational analysis of variance (PERMANOVA) in the main test. Pairwise PERMANOVA tests were performed for pairs of plastisphere and bulk soil samples of each plastic type, and the Monte Carlo approximated level of significance [*P*(MC)] was determined.

requirement for the biological degradation and assimilation of non-hydrolyzable plastics (O'Brine and Thompson, 2010; Kumar Sen and Raut, 2015).

In contrast, results for PBAT in Barba Peider soil suggested that chemical and physical changes on the polymer surface occurred due to microbial activity in this soil, as the plastic pieces incubated in natural soil differed from those incubated in sterile soil and from non-incubated pieces. On the other hand, the findings for Villum soil suggest that chemical changes occurred in the PBAT pieces as a result of unknown abiotic factors.

Notably, for the contact angle of PLA in Barba Peider soil, we observed a higher value for the sterile-soil incubated plastic relative to the non-incubated pieces and a lower angle for the natural-soil incubated plastic. Even though these results seem contradictory at first glance, the observation might be explained by opposing effects caused by abiotic degradation and microbial colonization of PLA (i.e., higher hydrophilicity of the surface due to biofilm formation). This circumstance might lead to diverging results in the physico-chemical analysis of the plastic pieces depending on the context (i.e., incubation duration). Our results demonstrate that degradation of PLA is partly abiotically driven. Abiotic degradation of PLA involves a temperature and moisture dependent chemical hydrolysis reaction (Karamanlioglu et al., 2017). It was previously shown that abiotic processes in PLA degradation (i.e., breaking down to a low molecular weight) precede microbial degradation and result in a material that is more accessible to microorganisms (Castro-Aguirre et al., 2017). The measured carbonyl indices of PLA incubated in sterile Villum soil provide further evidence that both abiotic and microbial degradation are affecting the breakdown of PLA at low temperatures.

The low level of plastic degradation observed in our study might be due to the short incubation period (8 weeks) and relatively low incubation temperature (15°C), an overall low microbial activity, or a lack of specific plastic-degrading microorganisms. Earlier soil burial experiments showed little biodegradation of PLA when incubated at 25°C for 120 days (Kamiya et al., 2007) or at 0–30°C for 1 year (Shogren et al., 2003). Another study showed little degradation of

PLA after a 1-year incubation in compost and soil at 25°C, but significant degradation in compost at 45°C after only 3 weeks of incubation (Karamanlioglu and Robson, 2013). Low temperatures and a lack of sunlight, as in our experimental set-up, potentially prevent abiotic degradation that could stimulate further microbial degradation (Singh and Sharma, 2008; Castro-Aguirre et al., 2017). Nonetheless, we detected small differences between non-incubated, sterile-soil incubated and natural-soil incubated biodegradable plastics by FTIR and contact angle measurements. It would be interesting to test different incubation temperatures and durations to investigate whether greater degrees of degradation could be achieved.

The optimal temperature for biodegradation in cold terrestrial environments remains unknown, as fast biodegradation is mostly reached at considerably higher temperatures like in industrial compost, whereas microorganisms in Arctic and alpine soils are adapted to low temperatures. Furthermore, other techniques such as respirometric measurements could be applied to better differentiate between abiotic and microbial degradation of plastics (Castro-Aguirre et al., 2017). Respirometric measurements and the analysis of more than one time-point could be possible approaches to elucidate the interplay between abiotic and biotic factors taking place in the plastisphere.

Plastics Decrease Diversity and Alter Microbial Community Structures

The two soils showed many similarities regarding the effects of plastics on microbial α -diversity. Overall, the microbial diversity was lower in the plastisphere than in bulk soil, in particular with PLA. While the PLA plastisphere clearly was most prone to changes in bacterial α -diversity in both soils, the results for fungi were less consistent. Even though effects of PLA were similar in the two soils, the fungal community in Villum soil was by far more affected by PBAT and PE. Moreover, PCoA and PERMANOVA revealed clear shifts in the microbial community structures that took place in the plastisphere of the various plastics.

TABLE 5 | Changes in the relative abundance of prokaryotic and fungal phyla and orders in the plastisphere.

Taxa	Barba Peider				Villum			
	Main test (F)	PLA	PBAT	PE	Main test (F)	PLA	PBAT	PE
Bacteria:								
Phyla								
Acidobacteria	31.5***	↓***	— ns	— ns	33.1***	↓***	↓**	↓*
Actinobacteria	39.8***	↑***	— ns	↑***	22.2***	— ns	— ns	↑***
Bacteroidetes	4.5*	— ns	— ns	— ns	0.4 ^{ns}	— ns	— ns	— ns
Chlamydiae	3.4*	— ns	— ns	— ns	2.8 ^(ns)	— ns	— ns	— ns
Chloroflexi	7.3**	↓**	— ns	— ns	13.1***	↓***	↓*	— ns
Gemmatimonadetes	31.7***	↓***	↓*	↓**	27.8***	↓***	↓**	— ns
Patescibacteria	34.7***	↑*	— ns	— ns	48.4***	↑***	— ns	— ns
Planctomycetes	22.4***	↓***	↓**	— ns	59.8***	↓***	↓**	↓***
Proteobacteria	40.0***	↑***	— ns	— ns	7.3**	— ns	↑**	— ns
Verrucomicrobia	6.0**	— ns	↓ ^(ns)	— ns	17.7***	↓***	— ns	— ns
Nitrospirae	5.4**	↓*	— ns	— ns	32.8***	↓***	↓**	↓**
Orders								
β-Proteobacteriales ¹	51.3***	↑***	— ns	— ns	8.7**	↓ ^(ns)	↑*	— ns
Chitinophagales	3.7*	— ns	— ns	— ns	4.3*	↓*	— ns	— ns
Chthoniobacteriales	4.9*	↓ ^(ns)	— ns	— ns	17.7***	↓***	— ns	— ns
Gemmatales	8.8**	↓**	↓*	— ns	54.3***	↓***	↓**	↓***
Gemmatimonadales	32.9***	↓***	↓*	↓**	29.1***	↓***	↓**	— ns
Ktedonobacteriales	5.6**	↓ ^(ns)	— ns	— ns	NA	NA	NA	NA
Parcubacteria	31.3***	↓***	— ns	— ns	6.2**	— ns	— ns	— ns
Pirellulales	12.8***	— ns	↓ ^(ns)	↓**	54.1***	↓***	↓***	↓**
Planctomycetales	9.5***	↓**	↓ ^(ns)	↓*	7.8**	— ns	— ns	— ns
Propionibacteriales	18.2***	— ns	— ns	↑***	15.9***	↑ ^(ns)	— ns	↑***
Pyrinomonadales	14.6***	↓***	— ns	↓*	29.1***	↓***	↓**	↓**
Rhizobiales	73.6***	↑***	— ns	— ns	8.7**	— ns	↑**	— ns
Saccharimonadales	26.5***	↑***	— ns	— ns	57.1***	↑***	— ns	— ns
Solibacteriales	22.1***	↓***	— ns	— ns	88.9***	↓***	↓***	↓***
Subgroup 6 ²	11.6***	↓ ^(ns)	↓**	— ns	28.0***	↓***	↓**	— ns
Tepidisphaerales	21.4***	↓***	↓ ^(ns)	— ns	20.5***	↓***	↓**	↓*
Fungi:								
Phyla								
Ascomycota	4.9*	— ns	— ns	— ns	1.4 ^{ns}	— ns	— ns	— ns
Basidiomycota	4.6*	— ns	— ns	— ns	0.9 ^{ns}	— ns	— ns	— ns
Mortierellomycota	18.3***	↓**	↑**	↑ ^(ns)	4.6*	— ns	— ns	— ns
Orders								
Cystofilobasidiales	1.2 ^{ns}	— ns	— ns	— ns	NA	NA	NA	NA
Helotiales	5.0*	— ns	↑*	— ns	2.2 ^{ns}	— ns	— ns	— ns
Pleosporales	1.2 ^{ns}	— ns	— ns	— ns	1.5 ^{ns}	— ns	— ns	— ns
Thelebolales	50.8***	↑***	— ns	— ns	17.7***	— ns	↑**	— ns

The ten most abundant bacterial phyla and orders were taken for each soil separately, resulting in a total of 11 phyla and 16 orders analyzed. The same procedure was applied to the three most abundant fungal phyla and orders, resulting in a total of three phyla and four orders. ANOVA was performed to determine the effect of plastic on the relative abundance of the selected taxa. *F*-values and statistical significances of the main effect are given in the first column for Barba Peider and for Villum. Tukey HSD tests were performed for the pairwise testing of plastisphere and bulk soil of each plastic type. The direction of change and statistical significances are given in columns two to four for each soil. ↑ represents a higher relative abundance of a taxon in the plastisphere in comparison to the respective bulk soil, while ↓ represents the opposite. No direction of change was analyzed for non-significant results. Asterisks indicate significant differences, with ****P* < 0.001, ***P* < 0.01, **P* < 0.05, ^(ns)*P* < 0.1, and ^{ns}*P* > 0.1; NA, not applicable. The relative abundances of the analyzed bacterial orders are visualized in **Figure 4**, and bacterial phyla and fungal taxa are shown in **Supplementary Figures 2–5**. ¹Silva database version v132 has changed β-Proteobacteria to the order β-Proteobacteriales (which include Burkholderiales) within the γ-Proteobacteria. ²Acidobacterial subgroup 6.

Parallel to the changes in α-diversity, we observed pronounced shifts in the microbial community structures with PLA and moderate changes with PBAT, whereas PE had no effect in either soil. Based on our hypothesis concerning the degree

of biodegradability of the different plastics, we expected to find the largest changes in microbial community structures in the plastisphere of PLA and little change for PE; this was confirmed with our DNA metabarcoding. In general,





FIGURE 4 | Relative abundance of the ten most abundant bacterial orders in the plastisphere (P) and bulk soil (S) of Barba Peider (Swiss Alps) **(A)** and Villum (Northern Greenland) **(B)** soils. The three replicates for each habitat are shown as separate columns. LCBD, local contribution to β -diversity.

it seems that the microbial community in Villum soil is more affected by plastic addition than that in Barba Peider soil. The different C and N content, and resulting higher initial microbial biomass in Villum soil could be a possible reason for these findings. In support of this explanation, Kamiya et al. (2007) found that several bioplastics were more prone to biodegradation in a soil with more organic


matter than in one with little organic matter. To improve our understanding on the biodegradation of plastics in cold environments, soils with different characteristics, e.g., varying in C, N and organic matter content, should be tested. Moreover, analyzing the plastisphere at different time-points might demonstrate if the colonization and degradation of the plastics occurs along with a succession of taxa, or if initial



microbial colonizers remain unchanged with time and at later degradation stages.


Reduced observed richness on the surface of plastics incubated in soil for 90 days was previously reported by Huang et al. (2019).

A decrease in α -diversity on plastic substrates compared with in the surroundings after only 2 weeks was previously reported in a study from a marine environment (Ogonowski et al., 2018). Most other experiments in marine environments documented higher



α -diversity on plastics than in the sea water (Zettler et al., 2013; Debroas et al., 2017; Dussud et al., 2018). De Tender et al. (2015) showed that only a very small set of OTUs in their marine study

belonged to a core plastisphere microbiome present in all samples with an abundance of at least 0.01%, providing evidence that plastisphere microbiomes are very diverse even among marine



environments. However, studies in aquatic and soil environments are difficult to compare (Bryant et al., 2016; Oberbeckmann et al., 2016; Kirstein et al., 2018). While plastic floating in water is in contact with a large number of different organisms over time and can serve as a raft for them (Oberbeckmann et al., 2016; Brunner et al., 2018), plastic substrates in soil are relatively immobile and stay permanently in contact with the inhabiting soil microbes, which are strongly adsorbed at soil surfaces (i.e., clay) (Jiang et al., 2007). There are several conceivable reasons for the reduction in the observed richness. By gaining access to a new energy and carbon source, they might outcompete other microbes not able to metabolize this carbon source in this specific niche (Dini-Andreote et al., 2015). This growth advantage can lead to the decline of other taxa, a phenomenon named “competitive exclusion” (Hibbing et al., 2010). Secondly, plastic addition can change the microenvironment of the soils by releasing potentially harmful compounds (Atuanya et al., 2016), changing the pH, or creating a barrier for water, oxygen, and nutrients (Bandopadhyay et al., 2018).

Plastic-Dependent Microbiome Relative Abundances of Microbial Taxa

Overall, the microbial communities of the plastisphere and bulk soil were characterized by taxonomic groups commonly observed in Arctic and alpine soils (Frey et al., 2016; Rime et al., 2016; Donhauser and Frey, 2018; Malard and Pearce, 2018). Only few phyla of this endemic microbiome showed an altered relative abundance in the plastisphere. Plastic addition might have little influence on the relative abundances of fungi if higher taxonomic ranks, for example phyla, are regarded. However, this interpretation might be misleading regarding the investigated soils, as the phylum Ascomycota is very dominant in these soils. Therefore, a plastic-induced selection for Ascomycota, as reported elsewhere (Muroi et al., 2016), cannot be detected by solely comparing relative abundances at the phylum level. The relative abundances of high-ranking bacterial taxa shared many similarities among the two soils. Only few phyla and orders seemed to be competitive in the plastisphere of PLA and PBAT. Especially Actinobacteria, Proteobacteria, and Patescibacteria seemed to profit from the plastisphere environment. While Proteobacteria increased in relative abundance on PLA in Barba Peider soil, it showed growth advantages on PBAT in Villum soil. Actinobacteria benefitted from PE in both soils, as well as from PLA in Barba Peider soil. Patescibacteria strongly increased in relative abundance in the PLA plastisphere in Villum soil and also profited from this plastic type in Barba Peider soil. Only little research has dealt with the colonization and biodegradation of plastics in soils so far. In one study, the effect of PBAT on the microbial community was analyzed by genetic profiling (DGGE), and Ascomycota were found to profit from the plastic, while the bacterial community did not change considerably (Muroi et al., 2016). Whereas this earlier study analyzed the long-term effects of plastic addition on a whole agricultural field, we investigated the region in proximity to single plastic pieces. A spatial limitation of the effects caused by plastic in soil might be one reason we observed more pronounced effects on the bacterial

community than reported previously. Other studies investigated the microbial community changes on PE microplastics and PLA/PHB (poly(3-hydroxybutyrate)) blend foils in laboratory incubation experiments (Jeszeová et al., 2018; Huang et al., 2019). Consistent with our findings, Huang et al. (2019) detected a significant increase in Actinobacteria on the plastic surface, and Jeszeová et al. (2018) found significant changes in the microbial community structure of PLA/PHB foil, i.e., increased numbers of *Streptomyces* and *Rhodococcus*.

Members of the Proteobacteria are physiologically and ecologically extremely diverse but are key players in C and N cycling. In particular, members of the Burkholderiales within Proteobacteria thrived in the plastisphere of both soils. Actinobacteria are ubiquitous and frequently saprophytic organisms able to degrade recalcitrant C sources. They play a vital role in the C cycle in the soil (Ventura et al., 2007; Bull, 2011; Rosenberg et al., 2014; Mohammadipanah and Wink, 2016). The ability to decompose recalcitrant compounds is an important trait in oligotrophic soil environments when more readily available substrates are rather limited. Furthermore, many members of the Actinobacteria are capable of spore formation and filamentous growth, which facilitates survival under low soil moisture (Wolf et al., 2013). With these capabilities, Actinobacteria are known to compete well under oligotrophic conditions. In contrast, Acidobacteria (i.e., subgroup Gp6) decreased in the plastisphere of all polymers. While the broad metabolic potential of Actinobacteria and Proteobacteria is widely known, it was surprising to find Patescibacteria and the associated order Saccharimonadales accumulating in the plastisphere.

Bacterial Key Taxa Associated With Plastics

Since different subgroups within a phylum can show ambiguous responses to polymer addition, we identified specific “*plastisphere taxa*” at lower taxonomic levels. Several actinobacterial taxa, such as *Rhodococcus*, *Nocardia* and *Streptomyces*, increased in the plastisphere in our study. Cultivated strains of these genera isolated from the incubated plastics yield further evidence for their presence on the plastic surfaces and complement our DNA metabarcoding data. Actinobacteria are, in general, promising candidates for biodegradation of all kinds of plastics, including PLA, PBAT, and strikingly even PE. Butbunchu and Pathom-Aree (2019) summarized the known abilities of Actinobacteria to degrade PLA. Interestingly, PLA biodegradation has been demonstrated for only a very limited number of actinobacterial families, and biodegradation by cold-adapted Actinobacteria has not yet been reported. PBAT degradation has so far only been shown for two thermophilic *Thermomonospora fusca* strains (Kleeberg et al., 1998). Other researchers reported that various isolated strains, including *Rhodococcus* spp., *Streptomyces* spp. and *Nocardia*, were associated with the biodegradation of PE (Kumar Sen and Raut, 2015; Pathak and Navneet, 2017). However, we have to consider that in most studies examining PE degradation, pretreated polymers or commercial products bearing additives have been used.

Most of the plastic-associated taxa within the Proteobacteria in our tested soils belong to a small number of orders, such as Burkholderiales, Caulobacteriales, Rhizobiales and

a few γ -Proteobacteria. Members of these orders were also found among the isolated strains obtained from the incubated plastics. Especially *Pseudomonas* is well known for its broad metabolic capabilities, and various strains of the genus are associated with the degradation of PLA, PBAT, PE, and many other plastics (Wallace et al., 2017; Wilkes and Aristilde, 2017; Bubbachat et al., 2018). Burkholderiales possess a broad range of enzymes able to degrade polyaromatic hydrocarbons (PAHs), which include important soil pollutants (Pérez-Pantoja et al., 2012). *Aquabacterium* was first isolated from biofilms of a drinking water system (Kalmbach et al., 1999), and a species of the genus was shown to degrade oil (Pham et al., 2015). Strains of the genus *Variovorax* hold highly diverse catabolic capabilities, including the degradation of 3,3'-Thiodipropionic acid (TDP), an additive widely used to stabilize polymers, as well as dimethyl terephthalate and vinyl chloride, two other chemicals used in plastic production (Satola et al., 2013; Wilson et al., 2016). *Collimonas* comprises chitinolytic species shown to degrade xenobiotics (Leveau et al., 2010); however, the genus has not been linked to plastic degradation yet. Species of the genera *Acidovorax* and *Leptothrix* are able to degrade the bioplastic poly(tetramethylene succinate)-co-adipate (PBSA) (Uchida et al., 2000; Nakajima-Kambe et al., 2009b). Additionally, the *Leptothrix* strain has been shown to degrade PLA (Nakajima-Kambe et al., 2009a). In our study, *Leptothrix* was increased in the PLA plastisphere of Barba Peider soil.

In a recent study, Rhizobiales were recognized as important taxon present on plastics floating in the North Atlantic garbage patch (Debroas et al., 2017). *Mesorhizobium* spp. have been shown to be associated with anaerobic PLA degradation (Yagi et al., 2014). *Sphingorhabdus*, a genus of the order Sphingomonadales, has been found to be able to degrade oil (Jeong et al., 2016). *Caulobacter crescentus* is a model laboratory organism that has been extensively studied concerning the process of the attachment of bacterial cells to surfaces and the formation of biofilms (Entcheva-Dimitrov and Spormann, 2004). In addition, *Caulobacter* spp. have been shown to degrade PAHs including pyrene and phenanthrene (Chang et al., 2014; Al-Thukair and Malik, 2016). A *Brevundimonas* species has been reported to degrade the biodegradable plastic poly(ϵ -caprolactone) (Nawaz et al., 2015).

We found members of the Saccharimonadales to be the major "plastisphere taxa" in both soils. To our knowledge, this is the first report of Saccharimonadales within the Patescibacteria superphylum thriving in the plastisphere. Very little is known about the ecology of Saccharimonadales, which belongs to the candidate phylum Saccharibacteria (formerly TM7), and knowledge almost exclusively stems from genomic data. Saccharibacteria have been shown to possess very small genomes and cell sizes, and they have therefore been proposed to live in symbiosis with other microorganisms depending on co-metabolism (Lemos et al., 2019). Further, Lemos et al. (2019) stated that the majority of unique genes found in nearly complete genomes of two

Saccharibacteria are of unknown function, underlining the knowledge gaps existing regarding this bacterial order. Several genes involved in the catabolism of complex C sources were identified in TM7 strain RAAC3 and other Saccharibacteria (Kantor et al., 2013; Starr et al., 2018). In addition, some Saccharibacteria were shown to take up oleic acid and to possess lipase and other exoenzymatic activities (Kindaichi et al., 2016). Interestingly, many Saccharibacteria seem to possess a D-lactate dehydrogenase-like protein acting in the conversion of pyruvate into lactate in a fermentation pathway (Lemos et al., 2019). Our data implicate that lactate, the monomer of PLA, might be used as a C source instead of being only a final waste product in the metabolism of some Saccharibacteria. Furthermore, metagenomic data indicate that representatives of the phylum are able to degrade a variety of polymers (Kantor et al., 2013; Starr et al., 2018). However, the involvement of Saccharibacteria in the degradation of PLA is speculative and needs to be addressed in future studies.

In our study, some taxa within the Bacteroidetes (i.e., *Sediminibacterium*, *Dyadobacter*, *Ohtaekwangia*) increased in the plastisphere. *Sediminibacterium* was previously shown to degrade vinylchloride (Wilson et al., 2016). This chemical is the monomer of an important plastic, namely polyvinylchloride (PVC). *Dyadobacter* and *Ohtaekwangia*, which both increased significantly in the plastisphere of PLA in Villum soil, were previously identified as potential biodegraders of xenobiotic pollutants (Willumsen et al., 2005; Tejada-Agredano et al., 2013; Bao et al., 2020).

As far as we know, the only link between Verrucomicrobia and plastics in earlier investigations was made in a study where Verrucomicrobia subdivision 1 was shown to be associated with polyethylene terephthalate (PET) in a marine environment, whereas subdivision 4 was rather found in the surrounding water (Oberbeckmann et al., 2016). Our results partly fit with this observation, as we found *Luteolibacter* (subdivision 1) to be increased in the proximity of plastic and *Lacunisphaera* (subdivision 4) increased in the bulk soil. However, *Cerasicoccus* (subdivision 4) was augmented in the plastisphere as well, which contrasts with the previous findings.

Fungal Key Taxa Associated With Plastics

Oidiodendron spp., found to be enriched in the plastisphere of PLA in Barba Peider soil, are closely related to the dominant *Pseudogymnoascus* and are found on a great variety of substrates, including wood, peat and human hair (Rice and Currah, 2005). The genus has been shown to possess the ability to degrade a broad spectrum of substrates, e.g., cellulose and lignin (Rice and Currah, 2005).

The genus *Alternaria*, found to be increased in the plastisphere of PLA in Villum soil, comprises saprobic and endophytic species and is known to cause serious disease in plants (Woudenberg et al., 2013). The genera *Alternaria* and *Pseudogymnoascus* (synonym *Geomyces*) were previously linked to polyurethane degradation (Loredo-Treviño et al., 2012). *Mycosphaerella*, another genus that increased in the plastisphere of PLA in Villum soil, likewise comprises plant pathogenic species. Species of this

genus have been shown to produce a wide variety of polymer-degrading enzymes involved in the break-down of plant cell walls (Douaiher et al., 2007).

Phlebia, a basidiomycetous genus found to be increased in the plastisphere of PE in Villum soil, is known to possess enzymes for lignin decomposition (Kantelinen et al., 1989) and has been linked to the biodegradation of PAHs (Mori et al., 2003). In our experiment, *Mycena* showed the largest growth advantage of all fungi in the plastisphere of all plastics in both soils. However, the strong increase in abundance was only observed in one replicate. Due to the lack of consistency, these results have to be interpreted with caution. Like *Phlebia*, *Mycena* has been linked to PAH degradation (Winquist et al., 2014).

CONCLUSION

As far as we know, this is the first report on the plastisphere microbiome of different plastic types (biodegradable and non-biodegradable) in terrestrial cryoenvironments using DNA metabarcoding. We found clear differences between microbial consortia on plastics and the surrounding environment, as previously reported in marine environments. Our findings suggest that plastic debris form a habitat for complex microbial assemblages with lifestyles and possibly metabolic pathways distinct from those of the microbial communities in bulk soil.

Even though degradation in our mesocosm experiment was slight, we detected pronounced shifts in bacterial and fungal α -diversity and community structures in the plastisphere compared with values in adjacent soils not affected by the plastics. Our results supported our hypothesis that the effects were larger on biodegradable (PLA, PBAT) than non-biodegradable plastics (PE).

Interestingly, members of only a few phyla (Actinobacteria, Proteobacteria, and Patescibacteria) were found to benefit from the plastics in both soils. Similarly, the key microbial taxa associated with plastics were similar in the two soils (Barba Peider and Villum). Plastic addition affected fungal taxa to a lesser extent, and we have not found any common taxa that reacted similarly in both soils. However, it is noteworthy that some genera (i.e., *Mycena* and *Plebia*) were increased on PE in the Villum. We found several candidate taxa in the plastisphere with known potential to biodegrade xenobiotics, e.g., Burkholderiales, *Pseudomonas*, *Caulobacter*, *Rhodococcus*, *Nocardia* and *Streptomyces* indicating the potential of cold-adapted microorganisms to degrade bioplastics. Plastic degraders from terrestrial cryoenvironments are thus conceivable but further investigations with single isolates or consortia of microorganisms degrading plastics needs to be performed.

In addition to identifying many organisms that are already prominently published in the context of plastics and especially their biodegradation, we found members of the Saccharimonadales to be the major “plastisphere taxa.” To our knowledge, this is the first report of Saccharimonadales within the Patescibacteria superphylum thriving in the plastisphere. Further research is needed to evaluate their

occurrence in the plastisphere of other soils from cold or temperate environments, to find traits responsible for their thriving on the PLA surface, and to analyze if they potentially even degrade PLA.

Increased human activities in the Arctic and the Alps demand environment-friendly processes for pollution control and clean-up, suitable for cold regions. Microorganisms from terrestrial cryoenvironments harbor great biotechnological potential, as they are known to possess special adaptations to cold temperatures. Major knowledge gaps still exist about the microbial life in the plastisphere, especially in terrestrial cryoenvironments. The expansive geographical dimensions of these environments point to their global importance, and we plead for further clarification of existing ambiguities. Our findings highlight that correct waste management is crucial not only for non-biodegradable but also for biodegradable plastics in order to protect environments all over the world against this serious threat.

DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found below: <https://www.ncbi.nlm.nih.gov/PRJNA630025>.

AUTHOR CONTRIBUTIONS

JR, BF, and IB designed the study. JR and DB performed the experiment. JR, LP-C, and DB participated in data analysis. JR wrote the manuscript with the help of BF and IB. All authors contributed to the article and approved the submitted version.

FUNDING

This study was partly funded by the Swiss National Science Foundation (SNSF) under the grant IZLSZ2_170941, the Swiss Polar Institute (SPI), and the BNP Paribas Swiss Foundation.

ACKNOWLEDGMENTS

We thank C. Perez-Mon for providing soils from Villum Research Station and Muot da Barba Peider; B. Stierli for laboratory assistance; R. Köchli for completing soil analyses; and S. Gunz and A. Dharmarajah for providing advice in data analysis. We acknowledge the Genetic Diversity Centre (GDC) of the ETH Zürich and the contribution of scientists at the McGill University and Génome Québec Innovation Center in Montreal, Canada for performing Illumina MiSeq sequencing. We also thank M. Dawes for her valuable contribution to the editing of this article.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fenvs.2020.562263/full#supplementary-material>

SUPPLEMENTARY DATA SHEET 1 | Supplementary Tables S1, S2.

SUPPLEMENTARY DATA SHEET 2 | Supplementary Figures S1–S7.

REFERENCES

- Accinelli, C., Saccà, M. L., Mencarelli, M., and Vicari, A. (2012). Deterioration of bioplastic carrier bags in the environment and assessment of a new recycling alternative. *Chemosphere* 89, 136–143. doi: 10.1016/j.chemosphere.2012.05.028
- Adamczyk, M., Hagedorn, F., Wipf, S., Donhauser, J., Vittoz, P., Rixen, C., et al. (2019). The soil microbiome of Gloria Mountain summits in the Swiss Alps. *Front. Microbiol.* 10:1080. doi: 10.3389/fmicb.2019.01080
- Allen, S., Allen, D., Phoenix, V. R., Le Roux, G., Durántez Jiménez, P., Simonneau, A., et al. (2019). Atmospheric transport and deposition of microplastics in a remote mountain catchment. *Nat. Geosci.* 12, 339–344. doi: 10.1038/s41561-019-0335-5
- Al-Thukair, A. A., and Malik, K. (2016). Pyrene metabolism by the novel bacterial strains *Burkholderia fungorum* (T3A13001) and *Caulobacter* sp (T2A12002) isolated from an oil-polluted site in the Arabian Gulf. *Int. Biodeter. Biodegr.* 110, 32–37. doi: 10.1016/j.ibiod.2016.02.005
- Amaral-Zettler, L. A., Zettler, E. R., and Mincer, T. J. (2020). Ecology of the plastisphere. *Nat. Rev. Microbiol.* 18, 139–151. doi: 10.1038/s41579-019-0308-0
- Ambrosini, R., Azzoni, R. S., Pittino, F., Diolaiuti, G., Franzetti, A., and Parolini, M. (2019). First evidence of microplastic contamination in the supraglacial debris of an alpine glacier. *Environ. Pollut.* 253, 297–301. doi: 10.1016/j.envpol.2019.07.005
- Anderson, M. J. (2001). Non-parametric MANOVA. *Aust. Ecol.* 26, 32–46. doi: 10.1111/j.1442-9993.2001.01070.pp.x
- Atuana, E., Udochukwu, U., and Dave-Omoregie, A. (2016). Bioavailability and toxicity of plastic contaminants to soil and soil bacteria. *Br. Microbiol. Res. J.* 13, 1–8. doi: 10.9734/bmrj/2016/25128
- Bandopadhyay, S., Martin-closas, L., and Pelacho, A. M. (2018). Biodegradable plastic mulch films: impacts on soil microbial communities and ecosystem functions. *Front. Microbiol.* 9:819. doi: 10.3389/fmicb.2018.00819
- Bao, H., Wang, J., Zhang, H., Li, J., Li, H., and Wu, F. (2020). Effects of biochar and organic substrates on biodegradation of polycyclic aromatic hydrocarbons and microbial community structure in PAHs-contaminated soils. *J. Hazard. Mater.* 385:121595. doi: 10.1016/j.jhazmat.2019.121595
- Bengtsson-Palme, J., Hartmann, M., Eriksson, K. M., Pal, C., Thorell, K., Larsson, D. G. J., et al. (2015). metaxa2: improved identification and taxonomic classification of small and large subunit rRNA in metagenomic data. *Mol. Ecol. Resour.* 15, 1403–1414. doi: 10.1111/1755-0998.12399
- Bengtsson-Palme, J., Ryberg, M., Hartmann, M., Branco, S., Wang, Z., Godhe, A., et al. (2013). Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing data. *Methods Ecol. Evol.* 4, 914–919. doi: 10.1111/2041-210X.12073
- Bergmann, M., Mützel, S., Primpke, S., Tekman, M. B., Trachsel, J., and Gerdt, G. (2019). White and wonderful? Microplastics prevail in snow from the Alps to the Arctic. *Sci. Adv.* 5, 1–11. doi: 10.1126/sciadv.aax1157
- Briassoulis, D., and Dejean, C. (2010). Critical review of norms and standards for biodegradable agricultural plastics part I. Biodegradation in soil. *J. Polym. Environ.* 18, 384–400. doi: 10.1007/s10924-010-0168-1
- Brunner, I., Fischer, M., Rüthi, J., Stierli, B., and Frey, B. (2018). Ability of fungi isolated from plastic debris floating in the shoreline of a lake to degrade plastics. *PLoS One* 13:e0202047. doi: 10.1371/journal.pone.0202047
- Bryant, J. A., Clemente, T. M., Viviani, D. A., Fong, A. A., Thomas, K. A., Kemp, P., et al. (2016). Diversity and activity of communities inhabiting plastic debris in the north Pacific gyre. *mSystems* 1, 1–19. doi: 10.1128/mSystems.0024-16
- SUPPLEMENTARY DATA SHEET 3 |** List of significant ($q > 0.05$) differently abundant (log₂-fold change) bacterial and fungal OTUs in the plastisphere compared to bulk soils.
- SUPPLEMENTARY DATA SHEET 4 |** List of significant ($q > 0.05$) differently abundant (log₂-fold change) bacterial and fungal genera in the plastisphere compared to bulk soils.
- SUPPLEMENTARY DATA SHEET 5 |** 16S rDNA sequences of isolated bacterial and fungal strains from the plastisphere.
- Bubpachat, T., Sombatsompop, N., and Prapagdee, B. (2018). Isolation and role of polylactic acid-degrading bacteria on degrading enzymes productions and PLA biodegradability at mesophilic conditions. *Polym. Degrad. Stabil.* 152, 75–85. doi: 10.1016/j.polymdegradstab.2018.03.023
- Bull, A. T. (2011). “Actinobacteria of the extremobiosphere,” in *Extremophiles Handbook*, ed. K. Horikoshi (Tokyo: Springer), 1203–1240. doi: 10.1007/978-4-431-53898-1_58
- Butbunchu, N., and Pathom-Aree, W. (2019). Actinobacteria as promising candidate for polylactic acid type bioplastic degradation. *Front. Microbiol.* 10:2834. doi: 10.3389/fmicb.2019.02834
- Castro-Aguirre, E., Auras, R., Selke, S., Rubino, M., and Marsh, T. (2017). Insights on the aerobic biodegradation of polymers by analysis of evolved carbon dioxide in simulated composting conditions. *Polym. Degrad. Stabil.* 137, 251–271. doi: 10.1016/j.polymdegradstab.2017.01.017
- Chang, Y. T., Hung, C. H., and Chou, H. L. (2014). Effects of polyethoxylylate lauryl ether (Brij 35) addition on phenanthrene biodegradation in a soil/water system. *J. Environ. Sci. Heal. A* 49, 1672–1684. doi: 10.1080/10934529.2014.951228
- Clarke, K., and Gorley, R. (2006). *PRIMER v6Q15: User Manual/Tutorial*. PRIMER-E (Plymouth: Plymouth Routines in Multivariate Ecological Research).
- da Silva, A. C., Rachid, C. T. C. C., de Jesus, H. E., Rosado, A. S., and Peixoto, R. S. (2017). Predicting the biotechnological potential of bacteria isolated from Antarctic soils, including the rhizosphere of vascular plants. *Polar Biol.* 40, 1393–1407. doi: 10.1007/s00300-016-2065-0
- de Souza Machado, A. A., Kloas, W., Zarfl, C., Hempel, S., and Rillig, M. C. (2018). Microplastics as an emerging threat to terrestrial ecosystems. *Glob. Chang. Biol.* 24, 1405–1416. doi: 10.1111/gcb.14020
- De Tender, C. A., Devriese, L. I., Haegeman, A., Maes, S., Ruttink, T., and Dawyndt, P. (2015). Bacterial community profiling of plastic litter in the Belgian part of the North Sea. *Environ. Sci. Technol.* 49, 9629–9638. doi: 10.1021/acs.est.5b01093
- Debroas, D., Mone, A., and Ter Halle, A. (2017). Plastics in the North Atlantic garbage patch: a boat-microbe for hitchhikers and plastic degraders. *Sci. Total Environ.* 599–600, 1222–1232. doi: 10.1016/j.scitotenv.2017.05.059
- Dini-Andreote, F., Stegen, J. C., Van Elsland, J. D., and Salles, J. F. (2015). Disentangling mechanisms that mediate the balance between stochastic and deterministic processes in microbial succession. *PNAS* 112, E1326–E1332. doi: 10.1073/pnas.1414261112
- Donhauser, J., and Frey, B. (2018). Alpine soil microbial ecology in a changing world. *FEMS Microbiol. Ecol.* 94, 1–31. doi: 10.1093/femsec/fiy099
- Douaiher, M. N., Nowak, E., Dumortier, V., Durand, R., Reignault, P., and Halama, P. (2007). *Mycosphaerella graminicola* produces a range of cell wall-degrading enzyme activities in vitro that vary with the carbon source. *Eur. J. Plant Pathol.* 117, 71–79. doi: 10.1007/s10658-006-9073-9
- Dussud, C., Meistertzheim, A. L., Conan, P., Pujo-Pay, M., George, M., Fabre, P., et al. (2018). Evidence of niche partitioning among bacteria living on plastics, organic particles and surrounding seawaters. *Environ. Pollut.* 236, 807–816. doi: 10.1016/j.envpol.2017.12.027
- Edgar, R. C. (2013). UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat. Methods* 10, 996–998. doi: 10.1038/nmeth.2604
- Edgar, R. C., and Flyvbjerg, H. (2015). Error filtering, pair assembly and error correction for next-generation sequencing reads. *Bioinformatics* 31, 3476–3482. doi: 10.1093/bioinformatics/btv401
- Entcheva-Dimitrov, P., and Spormann, A. M. (2004). Dynamics and control of biofilms of the oligotrophic bacterium *Caulobacter crescentus*. *J. Bacteriol.* 186, 8254–8266. doi: 10.1128/JB.186.24.8254-8266.2004

- Fierer, N. (2017). Embracing the unknown: disentangling the complexities of the soil microbiome. *Nat. Rev. Microbiol.* 15, 579–590. doi: 10.1038/nrmicro.2017.87
- Frasson, D., Udovičič, M., Frey, B., Lapanje, A., Zhang, D. C., Margesin, R., et al. (2015). *Glaciomonas alpina* sp. nov. isolated from alpine glaciers and reclassification of *glaciimonas immobilis* Cr9-12 as the type strain of *glaciimonas alpina* sp. nov. *Int. J. Syst. Evol. Microbiol.* 65, 1779–1785. doi: 10.1099/ijs.0.000174
- Frey, B., Rime, T., Phillips, M., Stierli, B., Hajdas, I., Widmer, F., et al. (2016). Bacterial diversity in European alpine permafrost and active layers. *FEMS Microbiol. Ecol.* 92:fw018. doi: 10.1093/femsec/fw018
- Hartmann, M., Brunner, I., Hagedorn, F., Bardgett, R. D., Stierli, B., Herzog, C., et al. (2017). A decade of irrigation transforms the soil microbiome of a semi-arid pine forest. *Mol. Ecol.* 26, 1190–1206. doi: 10.1111/mec.13995
- Heuer, H., Krsek, M., Baker, P., Smalla, K., and Wellington, E. M. (1997). Analysis of actinomycete communities by specific amplification of genes encoding 16S rRNA and gel-electrophoretic separation in denaturing gradients. *Appl. Environ. Microbiol.* 63, 3233–3241. doi: 10.1128/aem.63.8.3233-3241.1997
- Hibbing, M. E., Fuqua, C., Parsek, M. R., and Peterson, S. B. (2010). Bacterial competition: surviving and thriving in the microbial jungle. *Nat. Rev. Microbiol.* 8, 15–25. doi: 10.1038/nrmicro2259
- Horton, A. A., Walton, A., Spurgeon, D. J., Lahive, E., and Svendsen, C. (2017). Microplastics in freshwater and terrestrial environments: evaluating the current understanding to identify the knowledge gaps and future research priorities. *Sci. Total Environ.* 586, 127–141. doi: 10.1016/j.scitotenv.2017.01.190
- Huang, Y., Zhao, Y., Wang, J., Zhang, M., Jia, W., and Qin, X. (2019). LDPE microplastic films alter microbial community composition and enzymatic activities in soil. *Environ. Pollut.* 254:112983. doi: 10.1016/j.envpol.2019.112983
- Huerta Lwanga, E., Gertsen, H., Gooren, H., Peters, P., Salánki, T., Van Der Ploeg, M., et al. (2016). Microplastics in the terrestrial ecosystem: implications for *Lumbricus terrestris* (Oligochaeta, Lumbricidae). *Environ. Sci. Technol.* 50, 2685–2691. doi: 10.1021/acs.est.5b05478
- Jeong, H. I., Jin, H. M., and Jeon, C. O. (2016). Complete genome sequence of *Sphingorhabdus* sp. M41, a versatile hydrocarbon degrader, isolated from crude oil-contaminated coastal sediment. *J. Biotechnol.* 227, 41–42. doi: 10.1016/j.jbiotec.2016.04.016
- Jeszeová, L., Puškárová, A., Bučková, M., Kraková, L., Grivalský, T., Danko, M., et al. (2018). Microbial communities responsible for the degradation of poly(lactic acid)/poly(3-hydroxybutyrate) blend mulches in soil burial respirometric tests. *World J. Microbiol. Biotechnol.* 34:101. doi: 10.1007/s11274-018-2483-y
- Jiang, D., Huang, Q., Cai, P., Rong, X., and Chen, W. (2007). Adsorption of *Pseudomonas putida* on clay minerals and iron oxide. *Colloids Surf. B* 54, 217–221. doi: 10.1016/j.colsurfb.2006.10.030
- Kalmbach, S., Manz, W., Wecke, J., and Szewzyk, U. (1999). *Aquabacterium* gen. nov., with description of *Aquabacterium citratiphilum* sp. nov., *Aquabacterium parvum* sp. nov. and *Aquabacterium commune* sp. nov., three in situ dominant bacterial species from the Berlin drinking water system. *Int. J. Syst. Bacteriol.* 49, 769–777. doi: 10.1099/00207713-49-2-769
- Kamiya, M., Asakawa, S., and Kimura, M. (2007). Molecular analysis of fungal communities of biodegradable plastics in two Japanese soils. *Soil Sci. Plant Nutr.* 53, 568–574. doi: 10.1111/j.1747-0765.2007.00169.x
- Kantelinen, A., Hatakka, A., and Viikari, L. (1989). Production of lignin peroxidase and laccase by *Phlebia radiata*. *Appl. Microbiol. Biotechnol.* 31, 234–239.
- Kantor, R. S., Wrighton, K. C., Handley, K. M., Sharon, I., Hug, L. A., Castelle, C. J., et al. (2013). Small genomes and sparse metabolisms of sediment-associated bacteria from four candidate phyla. *mBio* 4:e00708-13. doi: 10.1128/mBio.00708-13
- Karamanlioglu, M., Preziosi, R., and Geoffrey D. R. (2017). Abiotic and biotic environmental degradation of the bioplastic polymer poly(lactic acid): a review. *Polym. Degrad. Stabil.* 137, 122–130. doi: 10.1016/j.polymdegradstab.2017.01.009
- Karamanlioglu, M., and Robson, G. D. (2013). The influence of biotic and abiotic factors on the rate of degradation of poly(lactic acid) (PLA) coupons buried in compost and soil. *Polym. Degrad. Stabil.* 98, 2063–2071. doi: 10.1016/j.polymdegradstab.2013.07.004
- Keswani, A., Oliver, D. M., Gutierrez, T., and Quilliam, R. S. (2016). Microbial hitchhikers on marine plastic debris: human exposure risks at bathing waters and beach environments. *Mar. Environ. Res.* 118, 10–19. doi: 10.1016/j.marenvres.2016.04.006
- Kindaichi, T., Yamaoka, S., Uehara, R., Ozaki, N., Ohashi, A., Albertsen, M., et al. (2016). Phylogenetic diversity and ecophysiology of Candidate phylum Saccharibacteria in activated sludge. *FEMS Microbiol. Ecol.* 92, 1–11. doi: 10.1093/femsec/fiw078
- Kirstein, I. V., Wichels, A., Gullans, E., Krohne, G., and Gerdt, G. (2019). The plastisphere – uncovering tightly attached plastic “specific” microorganisms. *PLoS One* 14:e215859. doi: 10.1371/journal.pone.0215859
- Kirstein, I. V., Wichels, A., Krohne, G., and Gerdt, G. (2018). Mature biofilm communities on synthetic polymers in seawater – specific or general? *Mar. Environ. Res.* 142, 147–154. doi: 10.1016/j.marenvres.2018.09.028
- Kleeberg, I., Hetz, C., Kroppenstedt, R. M., Müller, R. J., and Deckwer, W. D. (1998). Biodegradation of aliphatic-aromatic copolyesters by *Thermomonospora fusca* and other thermophilic compost isolates. *Appl. Environ. Microbiol.* 64, 1731–1735. doi: 10.1128/aem.64.5.1731-1735.1998
- Kumar Sen, S., and Raut, S. (2015). Microbial degradation of low density polyethylene (LDPE): a review. *J. Environ. Chem. Eng.* 3, 462–473. doi: 10.1016/j.jece.2015.01.003
- Laganà, P., Caruso, G., Corsi, I., Bergami, E., Venuti, V., Majolino, D., et al. (2019). Do plastics serve as a possible vector for the spread of antibiotic resistance? First insights from bacteria associated to a polystyrene piece from King George Island (Antarctica). *Int. J. Hyg. Environ. Health* 222, 89–100. doi: 10.1016/j.ijheh.2018.08.009
- Lahti, L., and Shetty, S. (2019). *Microbiome R Package (1.8.0)*. Available online at: <http://microbiome.github.io> (accessed March 4, 2020).
- Lapanje, A., Wimmersberger, C., Furrer, G., Brunner, I., and Frey, B. (2012). Pattern of elemental release during the granite dissolution can be changed by aerobic heterotrophic bacterial strains isolated from *Damma glacier* (Central Alps) deglaciated granite. *Microbiol. Ecol.* 63, 865–882. doi: 10.1007/s00248-011-9976-7
- Lemos, L. N., Medeiros, J. D., Dini-Andreote, F., Fernandes, G. R., Varani, A. M., Oliveira, G., et al. (2019). Genomic signatures and co-occurrence patterns of the ultra-small Saccharimonadia (phylum CPR/Patescibacteria) suggest a symbiotic lifestyle. *Mol. Ecol.* 28, 4259–4271. doi: 10.1111/mec.15208
- Leveau, J. H. J., Uroz, S., and de Boer, W. (2010). The bacterial genus *Collimonas*: mycophagy, weathering and other adaptive solutions to life in oligotrophic soil environments. *Environ. Microbiol.* 12, 281–292. doi: 10.1111/j.1462-2920.2009.02010.x
- Loredo-Treviño, A., Gutiérrez-Sánchez, G., Rodríguez-Herrera, R., and Aguilar, C. N. (2012). Microbial enzymes involved in polyurethane biodegradation: a review. *J. Polym. Environ.* 20, 258–265. doi: 10.1007/s10924-011-0390-5
- Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 1–21. doi: 10.1186/s13059-014-0550-8
- Luláková, P., Perez-Mon, C., Šantrůčková, H., Ruethi, J., and Frey, B. (2019). High-alpine permafrost and active-layer soil microbiomes differ in their response to elevated temperatures. *Front. Microbiol.* 10:668. doi: 10.3389/fmicb.2019.00668
- Malard, L. A., and Pearce, D. A. (2018). Microbial diversity and biogeography in Arctic soils. *Environ. Microbiol. Rep.* 10, 611–625. doi: 10.1111/1758-2229.12680
- Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.* 17, 10. doi: 10.14806/ej.17.1.200
- McMurdie, P. J., and Holmes, S. (2013). Phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* 8:e0061217. doi: 10.1371/journal.pone.0061217
- Mohammadipanah, F., and Wink, J. (2016). Actinobacteria from arid and desert habitats: diversity and biological activity. *Front. Microbiol.* 6:1541. doi: 10.3389/fmicb.2015.01541
- Mori, T., Kitano, S., and Kondo, R. (2003). Biodegradation of chloronaphthalenes and polycyclic aromatic hydrocarbons by the white-rot fungus *Phlebia lindtneri*. *Appl. Microbiol. Biotechnol.* 61, 380–383. doi: 10.1007/s00253-003-1253-3
- Muroi, F., Tachibana, Y., Kobayashi, Y., Sakurai, T., and Kasuya, K. I. (2016). Influences of poly(butylene adipate-co-terephthalate) on soil microbiota and plant growth. *Polym. Degrad. Stabil.* 129, 338–346. doi: 10.1016/j.polymdegradstab.2016.05.018
- Muyzer, G., Teske, A., Wiersen, C. O., and Jannasch, H. W. (1995). Phylogenetic relationships of Thiomicrospiraspecies and their identification in deep-sea

- hydrothermal vent samples by denaturing gradient gel electrophoresis of 16S rDNA fragments. *Arch. Microbiol.* 164, 165–172. doi: 10.1007/BF02529967
- Nakajima-Kambe, T., Ichihashi, F., Matsuzoe, R., Kato, S., and Shintani, N. (2009a). Degradation of aliphatic-aromatic copolyesters by bacteria that can degrade aliphatic polyesters. *Polym. Degrad. Stabil.* 94, 1901–1905. doi: 10.1016/j.polymdegradstab.2009.08.006
- Nakajima-Kambe, T., Toyoshima, K., Saito, C., Takaguchi, H., Akutsu-Shigeno, Y., Sato, M., et al. (2009b). Rapid monomerization of poly(butylene succinate)-co-(butylene adipate) by *Leptothrix* sp. *J. Biosci. Bioeng.* 108, 513–516. doi: 10.1016/j.jbiosc.2009.05.018
- Nauendorf, A., Krause, S., Bigalke, N. K., Gorb, E. V., Gorb, S. N., Haeckel, M., et al. (2016). Microbial colonization and degradation of polyethylene and biodegradable plastic bags in temperate fine-grained organic-rich marine sediments. *Mar. Pollut. Bull.* 103, 168–178. doi: 10.1016/j.marpolbul.2015.12.024
- Nawaz, A., Hasan, F., and Shah, A. A. (2015). Degradation of poly(ϵ -caprolactone) (PCL) by a newly isolated *Brevundimonas* sp. strain MRL-AN1 from soil. *FEMS Microbiol. Lett.* 362, 1–7. doi: 10.1093/femsle/fnu004
- Nilsson, R. H., Larsson, K. H., Taylor, A. F. S., Bengtsson-Palme, J., Jeppesen, T. S., Schigel, D., et al. (2019). The UNITE database for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Res.* 47, D259–D264. doi: 10.1093/nar/gky1022
- Oberbeckmann, S., Kreikemeyer, B., and Labrenz, M. (2018). Environmental factors support the formation of specific bacterial assemblages on microplastics. *Front. Microbiol.* 8:2709. doi: 10.3389/fmicb.2017.02709
- Oberbeckmann, S., Osborn, A. M., and Duhaime, M. B. (2016). Microbes on a bottle: substrate, season and geography influence community composition of microbes colonizing marine plastic debris. *PLoS One* 11:e159289. doi: 10.1371/journal.pone.0159289
- O’Brine, T., and Thompson, R. C. (2010). Degradation of plastic carrier bags in the marine environment. *Mar. Pollut. Bull.* 60, 2279–2283. doi: 10.1016/j.marpolbul.2010.08.005
- Ogonowski, M., Motiei, A., Ininbergs, K., Hell, E., Gerdes, Z., Udekwi, K. I., et al. (2018). Evidence for selective bacterial community structuring on microplastics. *Environ. Microbiol.* 20, 2796–2808. doi: 10.1111/1462-2920.14120
- Pathak, V. M., and Navneet. (2017). Review on the current status of polymer degradation: a microbial approach. *Bioresour. Bioprocess* 4:15. doi: 10.1186/s40643-017-0145-9
- Pérez-Pantoja, D., Donoso, R., Agulló, L., Córdova, M., Seeger, M., Pieper, D. H., et al. (2012). Genomic analysis of the potential for aromatic compounds biodegradation in Burkholderiales. *Environ. Microbiol.* 14, 1091–1117. doi: 10.1111/j.1462-2920.2011.02613.x
- Pham, V. H. T., Jeong, S. W., and Kim, J. (2015). *Aquabacterium olei* sp. nov., an oil-degrading bacterium isolated from oil-contaminated soil. *Int. J. Syst. Evol. Microbiol.* 65, 3597–3602. doi: 10.1099/ijsem.0.000458
- Piehl, S., Leibner, A., Löder, M. G. J., Dris, R., Bogner, C., and Laforsch, C. (2018). Identification and quantification of macro- and microplastics on an agricultural farmland. *Sci. Rep.* 8:17950. doi: 10.1038/s41598-018-36172-y
- PlasticsEurope. (2018). *Plastics - The facts 2017*. Brussels: PlasticsEurope.
- Pontes, A., Ruethi, J., Frey, B., Aires, A., Thomas, A., Overy, D., et al. (2020). *Cryolevonia* gen. nov. and *Cryolevonia schaffbergensis* sp. nov., a cryophilic yeast from ancient permafrost and melted sea ice. *Int. J. Syst. Evol. Microbiol.* 70, 2334–2338. doi: 10.1099/ijsem.0.004040
- Pushkareva, E., Eckhardt, K. U., Hotter, V., Frossard, A., Leinweber, P., Frey, B., et al. (2020). Chemical composition of soil organic matter and potential enzyme activity in the topsoil along a moisture gradient in the High Arctic (Svalbard). *Geoderma* 368, 114304. doi: 10.1016/j.geoderma.2020.114304
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., et al. (2013). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* 41, 590–596. doi: 10.1093/nar/gks1219
- R Development Core Team (2008). *R: A Language and Environment for Statistical Computing*. (3.6.2). Vienna: R Foundation for Statistical Computing.
- Rice, A. V., and Currah, R. S. (2005). Oidiodendron: a survey of the named species and related anamorphs of *Myxotrichum*. *Stud. Mycol.* 53, 83–120. doi: 10.3114/sim.53.1.83
- Rillig, M. C., Lehmann, A., de Souza Machado, A. A., and Yang, G. (2019). Microplastic effects on plants. *New Phytol.* 223, 1066–1070. doi: 10.1111/nph.15794
- Rime, T., Hartmann, M., and Frey, B. (2016). Potential sources of microbial colonizers in an initial soil ecosystem after retreat of an alpine glacier. *ISME J.* 10, 1625–1641. doi: 10.1038/ismej.2015.238
- Rosenberg, E., DeLong, E. F., Lory, S., Stackebrandt, E., and Thompson, F. (2014). *The Prokaryotes: Actinobacteria*. Berlin: Springer.
- Satola, B., Wübbeler, J. H., and Steinbüchel, A. (2013). Metabolic characteristics of the species *Variovorax paradoxus*. *Appl. Microbiol. Biot.* 97, 541–560. doi: 10.1007/s00253-012-4585-z
- Scheurer, M., and Bigalke, M. (2018). Microplastics in Swiss Floodplain soils. *Environ. Sci. Technol.* 52, 3591–3598. doi: 10.1021/acs.est.7b06003
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., et al. (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.* 75, 7537–7541. doi: 10.1128/AEM.01541-09
- Secretariat of the Convention of Biological Diversity (2016). *Marine Debris: Understanding, Preventing and Mitigating the Significant Adverse Impacts on Marine and Coastal Biodiversity*. CBD Technical Series (Vol. 83). Montreal, QC: Convention of Biological Diversity, doi: 10.1080/14888386.2007.9712830
- Shen, M., Zhang, Y., Zhu, Y., Song, B., Zeng, G., Hu, D., et al. (2019). Recent advances in toxicological research of nanoplastics in the environment: a review. *Environ. Pollut.* 252, 511–521. doi: 10.1016/j.envpol.2019.05.102
- Shi, Y., Grogan, P., Sun, H., Xiong, J., Yang, Y., Zhou, J., et al. (2015). Multi-scale variability analysis reveals the importance of spatial distance in shaping Arctic soil microbial functional communities. *Soil Biol. Biochem.* 86, 126–134. doi: 10.1016/j.soilbio.2015.03.028
- Shogren, R. L., Doane, W. M., Garlotta, D., Lawton, J. W., and Willett, J. L. (2003). Biodegradation of starch/poly(lactic acid)/poly(hydroxyester-ether) composite bars in soil. *Polym. Degrad. Stabil.* 79, 405–411. doi: 10.1016/S0141-3910(02)00356-7
- Singh, B., and Sharma, N. (2008). Mechanistic implications of plastic degradation. *Polym. Degrad. Stabil.* 93, 561–584. doi: 10.1016/j.polymdegradstab.2007.11.008
- Smith, M., Love, D. C., Rochman, C. M., and Neff, R. A. (2018). Microplastics in seafood and the implications for human health. *Curr. Environ. Health Rep.* 5, 375–386. doi: 10.1007/s40572-018-0206-z
- Ssekagiri, A. (2020). *microbiomeSeq: Microbial Community Analysis in an Environmental Context (0.1)*. Available online at: <http://www.github.com/umerijaz/microbiomeSeq> (accessed March 4, 2020).
- Starr, E. P., Shi, S., Blazewicz, S. J., Probst, A. J., Herman, D. J., Firestone, M. K., et al. (2018). Stable isotope informed genome-resolved metagenomics reveals that *Saccharibacteria* carbon. *Microbiome* 6:122.
- Tejeda-Agredano, M. C., Gallego, S., Vila, J., Grifoll, M., Ortega-Calvo, J. J., and Cantos, M. (2013). Influence of the sunflower rhizosphere on the biodegradation of PAHs in soil. *Soil Biol. Biochem.* 57, 830–840. doi: 10.1016/j.soilbio.2012.08.008
- Thiel, M., Luna-Jorquera, G., Álvarez-Varas, R., Gallardo, C., Hinojosa, I. A., Luna, N., et al. (2018). Impacts of marine plastic pollution from continental coasts to subtropical gyres-fish, seabirds, and other vertebrates in the SE Pacific. *Front. Mar. Sci.* 5:238. doi: 10.3389/fmars.2018.00238
- Uchida, H., Nakajima-Kambe, T., Shigeno-Akutsu, Y., Nomura, N., Tokiwa, Y., and Nakahara, T. (2000). Properties of a bacterium which degrades solid poly(tetramethylene succinate)-co-adipate, a biodegradable plastic. *FEMS Microbiol. Lett.* 189, 25–29. doi: 10.1016/S0378-1097(00)00246-9
- Ventura, M., Canchaya, C., Tauch, A., Chandra, G., Fitzgerald, G. F., Chater, K. F., et al. (2007). Genomics of actinobacteria: tracing the evolutionary history of an ancient phylum. *Microbiol. Mol. Biol. Rev.* 71, 495–548. doi: 10.1128/mmb.00005-07
- Wallace, P. W., Haernvall, K., Ribitsch, D., Zitzenbacher, S., Schittmayer, M., Steinkellner, G., et al. (2017). PpEst is a novel PBAT degrading polyesterase identified by proteomic screening of *Pseudomonas pseudoalcaligenes*. *Appl. Microbiol. Biotechnol.* 101, 2291–2303. doi: 10.1007/s00253-016-7992-8
- Wang, J., Lv, S., Zhang, M., Chen, G., Zhu, T., Zhang, S., et al. (2016). Effects of plastic film residues on occurrence of phthalates and microbial activity in soils. *Chemosphere* 151, 171–177. doi: 10.1016/j.chemosphere.2016.02.076

- Wang, Q., Garrity, G. M., Tiedje, J. M., and Cole, J. R. (2007). Naïve Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.* 73, 5261–5267. doi: 10.1128/AEM.00062-07
- White, T., Bruns, T., Lee, S., and Taylor, J. (1990). “Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics,” in *PCR Protocols: A Guide to Meth- Ods and Applications*, eds M. Innis, D. Gelfand, J. Sninsky, and T. White (San Diego, CA: Academic Press), 315–322.
- Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis* (3.2.1). New York, NY: Springer-Verlag.
- Wilkes, R. A., and Aristilde, L. (2017). Degradation and metabolism of synthetic plastics and associated products by *Pseudomonas* sp.: capabilities and challenges. *J. Appl. Microbiol.* 123, 582–593. doi: 10.1111/jam.13472
- Willumsen, P. A., Johansen, J. E., Karlson, U., and Hansen, B. M. (2005). Isolation and taxonomic affiliation of N-heterocyclic aromatic hydrocarbon-transforming bacteria. *Appl. Microbiol. Biotechnol.* 67, 420–428. doi: 10.1007/s00253-004-1799-8
- Wilson, F. P., Liu, X., Mattes, T. E., and Cupples, A. M. (2016). *Nocardioidea*, *Sediminibacterium*, *Aquabacterium*, *Variovorax*, and *Pseudomonas* linked to carbon uptake during aerobic vinyl chloride biodegradation. *Environ. Sci. Pollut. Res. Int.* 23, 19062–19070. doi: 10.1007/s11356-016-7099-x
- Winqvist, E., Björklöf, K., Schultz, E., Räsänen, M., Salonen, K., Anasonye, F., et al. (2014). Bioremediation of PAH-contaminated soil with fungi - From laboratory to field scale. *Int. Biodeter. Biodegr.* 86, 238–247. doi: 10.1016/j.ibiod.2013.09.012
- Wolf, A. B., Vos, M., De Boer, W., and Kowalchuk, G. A. (2013). Impact of matric potential and pore size distribution on growth dynamics of filamentous and non-filamentous soil bacteria. *PLoS One* 8:e83661. doi: 10.1371/journal.pone.0083661
- Woudenberg, J. H. C., Groenewald, J. Z., Binder, M., and Crous, P. W. (2013). *Alternaria* redefined. *Stud. Mycol.* 75, 171–212. doi: 10.3114/sim0015
- Wright, S. L., and Kelly, F. J. (2017). Plastic and human health: a micro issue? *Environ. Sci. Technol.* 51, 6634–6647. doi: 10.1021/acs.est.7b00423
- Yagi, H., Ninomiya, F., Funabashi, M., and Kunioka, M. (2014). Mesophilic anaerobic biodegradation test and analysis of eubacteria and archaea involved in anaerobic biodegradation of four specified biodegradable polyesters. *Polym. Degrad. Stabil.* 110, 278–283. doi: 10.1016/j.polymdegradstab.2014.08.031
- Zettler, E. R., Mincer, T. J., and Amaral-Zettler, L. A. (2013). Life in the “plastisphere”: microbial communities on plastic marine debris. *Environ. Sci. Technol.* 47, 7137–7146. doi: 10.1021/es401288x
- Zhang, G. S., and Liu, Y. F. (2018). The distribution of microplastics in soil aggregate fractions in southwestern China. *Sci. Total Environ.* 642, 12–20. doi: 10.1016/j.scitotenv.2018.06.004

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Rüthi, Bölsterli, Pardi-Comensoli, Brunner and Frey. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.