

NANO EXPRESS

Open Access



The Potential Liver, Brain, and Embryo Toxicity of Titanium Dioxide Nanoparticles on Mice

Xiaochuan Jia^{1,2}, Shuo Wang^{1*}, Lei Zhou² and Li Sun²

Abstract

Nanoscale titanium dioxide (nano-TiO₂) has been widely used in industry and medicine. However, the safety of nano-TiO₂ exposure remains unclear. In this study, we evaluated the liver, brain, and embryo toxicity and the underlying mechanism of nano-TiO₂ using mice models. The results showed that titanium was distributed to and accumulated in the heart, brain, spleen, lung, and kidney of mice after intraperitoneal (i.p.) nano-TiO₂ exposure, in a dose-dependent manner. The organ/body weight ratios of the heart, spleen, and kidney were significantly increased, and those of the brain and lung were decreased. High doses of nano-TiO₂ significantly damaged the functions of liver and kidney and glucose and lipid metabolism, as showed in the blood biochemistry tests. Nano-TiO₂ caused damages in mitochondria and apoptosis of hepatocytes, generation of reactive oxygen species, and expression disorders of protective genes in the liver of mice. We found ruptured and cracked nerve cells and inflammatory cell infiltration in the brain. We also found that the activities of constitutive nitric oxide synthases (cNOS), inducible NOS (iNOS), and acetylcholinesterase, and the levels of nitrous oxide and glutamic acid were changed in the brain after nano-TiO₂ exposure. Ex vivo mouse embryo models exhibited developmental and genetic toxicity after high doses of nano-TiO₂. The size of nano-TiO₂ particles may affect toxicity, larger particles producing higher toxicity. In summary, nano-TiO₂ exhibited toxicity in multiple organs in mice after exposure through i.p. injection and gavage. Our study may provide data for the assessment of the risk of nano-TiO₂ exposure on human health.

Keywords: Titanium dioxide, Nanoparticle, Liver toxicity, Brain toxicity, Fetus toxicity

Background

Nanoscale titanium dioxide (nano-TiO₂) is widely used in the food industry. It has been used for the production of coated candy, preserved fruits, chewing gum, carbonated drinks, powdered drinks (in unsweetened dosage form or concentrated), milk and dairy products, and other food categories [1, 2]. The concentration of nano-TiO₂ in food reaches as high as 0.5–9 g/kg [1, 3], and many food products that are claimed nano-TiO₂-free contain nano-TiO₂ [2]. Nano-TiO₂ has also been widely used in biomedicine, organic pollutant treatment, materials engineering, and cosmetics [4–6]. However, the safety of nano-TiO₂ exposure remains unclear.

Studies have shown that nano-TiO₂ can become enriched and toxic in multiple organs after entering the

body through several methods, such as administration via the abdominal cavity or inhalation [7, 8]. Nano-TiO₂ may be toxic to several types of cell, such as human lymphoblastoid cells and hepatoma cells [9, 10]. It can induce an acute stress reaction in glial cells of mouse brains, leading to neuron damage and dysfunction [11]. The survival rate of neuron cell lines exposed to nano-TiO₂ particles is significantly decreased in a typical time- and dose-dependent manner [12].

Studies have revealed several mechanisms by which these nanoparticles cause toxicity. Nano-TiO₂ particles may cause genetic toxicity through changing the structure of molecular complex and the permeability of cell membrane [13–15]. Nano-TiO₂ may produce oxidative stress. During oxidative stress, reactive oxygen species (ROS), such as hydroxyl radicals, are generated and cause DNA oxidation, generating 8-OHG, leading to errors and mutations in DNA replication [16, 17].

* Correspondence: jwy7412@126.com

¹Tianjin University of Science and Technology, Tianjin 300457, China
Full list of author information is available at the end of the article

Moreover, ROS may induce inflammation and mutual feed-forward interaction between oxidative stress and inflammation, resulting in DNA damage and cell apoptosis [18, 19]. However, the comprehensive systematic data on the toxicity of nano-TiO₂ remains limited. Our objective was to reveal the effect and the underlying mechanism of nano-TiO₂ exposure on human health.

In this study, we evaluated the effect and the underlying mechanism of nano-TiO₂ exposure using mice models. Our findings showed that nano-TiO₂ may be enriched and cause toxicity in several organs such as the liver, kidney, spleen, heart, lung, and brain through generating an oxidation-reduction imbalance and disorders of gene expression. This may also cause damage to embryonic development. Our study may provide data to assess the potential risk to human health of nano-TiO₂ exposure.

Methods

Chemicals and Reagents

Micro-scale TiO₂ (micro-TiO₂) and 5 nm of TiO₂ in the form of anatase were purchased from Sigma-Aldrich (Shanghai, China), and 10, 60, and 90 nm of TiO₂ (anatase) were purchased from Run He Ltd. (Shanghai, China). The formaldehyde, nitric acid, hydrogen peroxide, and heparin sodium were reagent grade and were purchased from Sigma-Aldrich (Shanghai, China). Phosphate buffer (PBS), penicillin, and streptomycin were purchased from Gibco (San Diego, USA). Total RNA extraction kits were purchased from Takara (Dalian, China). Reactive oxygen species assay kits were purchased from Jianchen Ltd. (Nanjing, China). Stock TiO₂ suspension (1%) in Hank's solution was sterilized at 121 °C for 30 min. The suspension was sonicated and diluted to the desired concentration just before use.

Animals and Models

For the study of liver and brain toxicity, ICR (imprinting control region) mice (22 ± 3 g, half male and half female) were purchased from the animal center of China Medical University. All experimental procedures involving animals were pre-approved by the Institutional Ethics Committee Tianjin University of Science and Technology and were conducted in accordance with the international guidelines for care and use of laboratory animals. For the study of mouse embryo toxicity, ICR mice (45 females, 20–35 g; 15 males, 35–40 g) were purchased from Beijing Weitong Lihua Ltd. (Beijing, China). All mice were healthy and sexually mature. Five days before treatment, mice were reared in separate cages in a house with good ventilation, a 12-h light/dark cycle, 20 ± 2 °C, 60 ± 10% relative humidity, and ad libitum access to food and water.

Dosing regimen 1 was designed for a general toxicity and brain toxicity test. The mice were randomly divided into six groups and an additional control group, with 10 mice/group.

A nanoscale TiO₂ (nano-TiO₂) suspension was injected (intraperitoneal (i.p.), 5, 10, 50, 100, 150, and 200 mg/kg) once a day for 14 days. Saline solution was injected into mice of the control group. The mice were observed every day, and no animal died during the study. On the 15th day, blood samples were collected from the orbital sinus. All mice were individually weighed, were anesthetized with 2% Phenobarbital (60 ml/kg, i.p.), and then were sacrificed through cervical dislocation. All tissue samples were collected (brain tissue isolated from cortex and hippocampus) and were stored at –80 °C. Each heart, liver, spleen, lung, and kidney was cut into two portions. One portion was soaked in formalin (10%) solution at 4 °C for pathological examination. The other portion was stored at –20 °C for the determination of the titanium content.

Dosing regimen 2 was designed for the liver toxicity test. Mice were divided into three experimental groups and one control group. Nano-TiO₂ (5, 10, 50 mg/kg) was administered once per day through gavage for 60 days. The mice in the control group received 0.5% CMC (carboxymethyl cellulose). The mice were observed every day, and no animal died during the study. On the 60th day, the mice were anesthetized with 2% Phenobarbital (60 ml/kg, i.p.), and then were sacrificed through cervical dislocation, the livers were immediately collected and were processed for examination using electron microscopy, determination of ROS and lipid oxidation, and analysis of gene expression.

Determination of Titanium Content in Target Tissues

A piece of 0.1–0.5-g frozen tissue sample was cut and thawed at room temperature, and then was digested in HNO₃ (0.5 mL) and H₂O₂ (0.5 mL) at 160 °C. After being diluted to 3 mL with 3% nitric acid, the concentration of titanium in the solution was determined using inductively coupled plasma mass spectrometry (ICP-MS). Titanium content in target tissues was then calculated.

Blood Biochemistry Tests and Calculation of Organ/Body Weight Ratio

The levels of the enzymes in serum samples were analyzed by an automatic biochemistry analyzer (TBA-2000FR, Toshiba, Tokyo, Japan). These enzymes are biomarkers related to the function of the liver and kidney.

The organ/body weight ratio was calculated based on the weights of the organ and the body. The body weights were measured before anesthesia and sacrifice. Organs were weighted after isolation from anesthetized and sacrificed mice.

Pathological Examination and Transmission Electron Microscopy

Pathological examination of the liver or brain tissues soaked in formalin was performed under a light microscope after

hematoxylin staining. For transmission electron microscopy (TEM), the liver tissues were embedded in epoxy resin EPON 812 and were cut into sections as thin as <500 μm after glutaraldehyde and osmic acid fixation. The sections were stained with saturated acetic acid uranium solution (pH 3.5) and lead citrate (pH 12) for 1–2 h. The stained sections were examined using TEM.

Determination of the Levels of Reactive Oxygen Species, the Activities of Their Metabolic Enzymes, and the Levels of Neurotransmitters

For the liver tissues, the superoxide anion (O_2^-) levels were determined using XTT. The activity of catalase (CAT) was determined using OD values at 240 nm, following published procedures [20]. The levels of lipid peroxidation were determined by the content of malondialdehyde (MDA) following published procedures [21].

Brain tissues were homogenized with pre-cooled 1% polyvinylpyrrolidone solution (50 mM in pH 7.6 PBS) after isolation. Supernatants were collected after centrifugation at 15,000 rpm for 20 min (Eppendorf 5418, Hamburg, Germany) and were used for subsequent analysis of the activities of superoxide enzyme (SOD), CAT, ascorbate peroxidase (APX), and glutathione peroxidase (GSHPx). SOD activity was determined using NBT (nitro-tetrazolium chloride blue test). Catalase activity was determined using a kit (CAT assay kit, A007-2, Nanjing Jiancheng Bioengineering Institute, Nanjing, China). The activity of APX was measured using a kit (APX assay kit, A123, Nanjing Jiancheng Bioengineering Institute, Nanjing, China). GSHPx activity was determined using a kit (GSHPx assay kit, A005, Nanjing Jiancheng Bioengineering Institute, Nanjing, China). The activities of constitutive nitric oxide synthase (cNOS), inducible nitric oxide synthase (iNOS), and acetylcholinesterase (AChE) were determined using commercial kits (AChE assay kit, A024, Nanjing Jiancheng Bioengineering Institute, Nanjing, China).

The levels of ROS in brain tissues were determined by adding 2', 7' dichlorofluorescein diacetate to a final concentration of 10 μM in brain tissue homogenate and by incubating at 37 $^\circ\text{C}$ for 30 min; the tissues were then subjected to analysis using flow cytometry.

Determination of Relative mRNA Levels

Total RNA was extracted from liver tissue samples using a commercial kit (TaKaRa MiniBEST Universal RNA Extraction Kit, 9767, Takara, Dalian, China). Complementary DNA was synthesized using reverse transcription with random primer. The relative mRNA levels of SOD, CAT, GSHPx, MT, HSP70, CYP1A, P53, GST, and TF were determined using a real-time quantitative PCR (qPCR) kit (One Step SYBR[®] PrimeScript[™] RT-PCR Kit, PR066A, Takara, Dalian, China). All the primers (Table 1)

Table 1 Primers for the gene expression analysis using real-time PCR

Genes	Sequences (5'–3')	Amplicon (bp)
CAT	Forward: AGCGACCAGATGAAGCAGTG Reverse: GGGTGACCTCAAAGTATCCAAA	241
CYP1A	Forward: CGTCGCAGAGTATCCAGAGG Reverse: TTAACCGGGTAGCCGTCAAT	204
GSHPx	Forward: GGGACTACACCGAGATGAACG Reverse: TCCGCAGGAAGGTAAGAGC	231
GST	Forward: CCCTCTTTGGGGCTTTAT Reverse: GGTTCTGGGACAGCAGGGT	191
HSP 70	Forward: CATCGCCAACGACCAGG Reverse: ACCGCATCGCCGAACCT	162
MT	Forward: CTCCTGCACTTGACCCAGC Reverse: CACATTTGGAGCAGCCAC	100
P53	Forward: GCTGGTTCATCACTCCTCCC Reverse: GCTTCCCCATTTCACTCTGG	216
SOD	Forward: CTGGACAAACCTGAGCCCTAA Reverse: TCCCCAGCAGCGGAATAA	242
TF	Forward: TGTCAGAGCACGAGAATACCAA Reverse: ATAAAACCTCCGCCCCAC	224

were synthesized and purchased from Shanghai Sangon Ltd.

Ex Vivo Embryo Toxicity Test

Embryos of 8.5 embryonic days were isolated from female mice after cervical dislocation and then were cultured in 50.0-mL Hank's solution containing 3 mL immediately centrifuged serum (ICS) from rats, micro-TiO₂, or nano-TiO₂ (0.0, 50.0, 100.0, and 200.0 $\mu\text{g}/\text{mL}$) with 3 embryos in each bottle for 48 h.

To determine the effect of micro-TiO₂ or nano-TiO₂ exposure time on embryos, the embryos were cultured in 50.0-mL Hank's solution containing 3 mL ICS from rats, micro-TiO₂, or nano-TiO₂ (200.0 $\mu\text{g}/\text{mL}$) with 3 embryos in each bottle for 16, 26, and 48 h, and then were washed for 48 h with pre-warmed 37 $^\circ\text{C}$ Hank's solution and cultured in 50.0-mL Hank's solution containing 3 mL ICS from rats.

Embryonic development was evaluated using the Maele-Fabry Van score [22]. The yolk sac diameter, embryonic crown–rump length, head length, and numbers of body sections were examined under a dissecting microscope. The malformation rate of developing embryos was evaluated based on the scores of morphological changes of the forebrain, midbrain, hindbrain, the forelimb bud, the hindlimb bud, the auditory and visual systems, and the heart. More than 10 embryos from 2 ICR mice at embryonic day 10.5 were isolated for control.

Statistical Analysis

Data was analyzed using SPSS 13 (IBM, Illinois, USA). The difference between the treatment group and the control group was analyzed using a Dunnett's *t* test. The difference among groups was analyzed using ANOVA. The comparison between two of multiple samples were analyzed using the LSD and SNK tests. Categorical data were analyzed using the chi-square test and rank sum test. If $P < 0.05$, the difference was considered significant.

Results

Tissue Distribution of Titanium in Mice After Nanoscale Titanium Dioxide Exposure

We treated mice with nano-TiO₂ (i.p., 5, 10, 50, 100, 150, and 200 mg/kg) for 14 days and determined titanium contents in the organs of mice. The results revealed that titanium was accumulated in the organs of mice treated with different doses of nano-TiO₂ (Fig. 1). The magnitude of accumulation was dose-dependent (Fig. 1). The liver was the organ where titanium was enriched most followed by the kidney. The magnitude of accumulation of titanium was approximately the same in the spleen, lung, brain, and heart (Fig. 1). The results suggest that nano-TiO₂ can be absorbed through GI track and distributed to tissues through the circulatory system and deposited in the organs liver, kidney, spleen, lung, brain, and heart.

General Toxicity of Nanoscale Titanium Dioxide in Mice

We treated mice with different doses of nano-TiO₂ for 14 days and found that there was no difference in body weight gains among groups of mice treated with different doses (data not shown). Low doses of nano-

TiO₂ (5 and 10 mg/kg) did not change the organ/body weight ratio of the liver, kidney, spleen, lung, heart, and brain in mice after i.p. exposure for 14 days (Fig. 2). However, the high doses of nano-TiO₂ (50, 100, 150, and 200 mg/kg) significantly increased the organ/body weight ratio of the liver, kidney, spleen, and heart and decreased those of lung and brain in mice in a dose-dependent manner (Fig. 2).

Lower doses (5, 10, 50, and 100 mg/kg) of nano-TiO₂ did not change any blood biochemistry index (Fig. 3). High doses of nano-TiO₂ (150 to 200 mg/kg) elevated liver function biomarkers alkaline phosphatase (ALP) and alanine aminotransferase (ALT), albumin (ALB), leucine aminopeptidase (LAP), butyrylcholinesterase (PChe), total bilirubin (TBIL), and total protein (TP) levels (Fig. 3). High doses decreased serum uric acid (UA) and blood urea nitrogen (BUN) levels, which are biomarkers for kidney function. They increased serum aspartate aminotransferase (AST), creatine kinase (CK), lactate dehydrogenase (LDH), and alpha hydroxybutyrate dehydrogenase (HBDH) levels, which are indices for myocardial damage (Fig. 3).

These results suggest that high doses of TiO₂ can cause severe damage to the liver, kidney, heart, and other organs in a dose-dependent manner.

Liver Toxicity of Nano-TiO₂ in Mice

We further evaluated the liver toxicity of nano-TiO₂. Using light microscopy, we found that there was no significant change in the livers of mice exposed to low dose (i.p. for 14 days, 5 mg/kg) nano-TiO₂ (Fig. 4a, b). We observed marked vascular obstruction and dilation (Fig. 4c, 50 mg/kg), increases in basophils (Fig. 4d,






Fig. 1 Titanium was accumulated in organs of mice exposed to nano-TiO₂. Mice were treated with nano-TiO₂ suspension or saline solution as indicated through intraperitoneal injection once a day for 14 days. * compared to control, $P < 0.05$, # compared to control, $P < 0.01$






Fig. 4 Histology of livers in mice treated with nano-TiO₂ exposed to nano-TiO₂. Mice were treated with nano-TiO₂ suspension or saline solution as indicated through intraperitoneal injection once a day for 14 days. **a** Control. **b** TiO₂, 5 mg/kg. **c** TiO₂, 50 mg/kg. **d** TiO₂, 100 mg/kg. **e** TiO₂, 150 mg/kg. **f** TiO₂, 200 mg/kg

100 mg/kg), partial ischemia in the liver (Fig. 4e, 150 mg/kg), and obstruction of central veins (Fig. 4f, 200 mg/kg) in mice exposed to nano-TiO₂ (i.p.).

However, using TEM, we found a slight swelling of the mitochondria in hepatocytes and presence of condensed chromatin and apoptotic cells in the liver tissues in mice exposed to low dose of nano-TiO₂ (gavage for 60 days, 5 mg/kg) (Fig. 5a, b). We observed nano-TiO₂ in the mitochondria of hepatocytes, swelling mitochondria, and vacuoles in mitochondria of the liver cells of mice treated with 10 mg/kg nano-TiO₂ (gavage for 60 days, Fig. 5c). We further observed nucleolus collapse, scattered chromatin, obvious apoptosis, and/or apoptotic bodies in the liver cells of mice treated with 50 mg/kg nano-TiO₂ (gavage for 60 days, Fig. 5d). The results indicated that nano-TiO₂ can lead to pathological damage in liver cells at the subcellular and cellular levels.

The treatment of mice with 5 mg/kg nano-TiO₂ for 60 days did not change the levels of ROS such as O²⁻, H₂O₂, nitric oxide (NO), and MDA (Fig. 6), or the mRNA levels of SOD, CAT, GSHPx, MT, GST, HSP70, P53, and TF genes in liver tissues (Fig. 7). Treatment of mice with 10 or 50 mg/kg nano-TiO₂ for 60 days resulted in significant increases in the levels of O²⁻, H₂O₂,

NO, and MDA (Fig. 6), decreases in the mRNA levels of SOD, CAT, MT, GST, HSP70, P53, TF, and GSHPx genes, and increases in the mRNA levels of CYP1A genes in the livers of mice (Fig. 7). The results showed that high doses of nano-TiO₂-induced oxidative stress and changes in the expression of protective genes in the livers of exposed mice.

Brain Toxicity of Nanoscale Titanium Dioxide in Mice

We further evaluated brain toxicity of nano-TiO₂. We first examined the ratios of brain/body weights in the mice exposed to nano-TiO₂ (i.p. for 14 days). Low doses (5, 10, 50 mg/kg) did not change the ratios of brain/body weights, and higher doses (100, 150, 200 mg/kg) significantly decreased the ratios of brain/body weights in a dose-dependent manner (Fig. 2). The concentration of Ti in the brain tissues was significantly increased in a dose-dependent manner (Fig. 1).

We also examined the histological changes in the brain of mice exposed to nano-TiO₂ (i.p. for 14 days) using hematoxylin staining. We observed that low doses of nano-TiO₂ (50 mg/kg) did not change the histology of brain tissue in mice after i.p. exposure for 14 days (Fig. 8a, b). Treatment of mice with 100 mg/kg of nano-TiO₂ resulted




Fig. 5 Ultramicroscopic structure of hepatocytes in mice exposed to nano-TiO₂. Mice were treated with nano-TiO₂ as indicated through gavage once per day for 60 days. The mice in the control group received 0.5% CMC (carboxymethyl cellulose). **a** Control (x8000). **b** TiO₂ (5 mg/kg) (x8000). **c** TiO₂ (10 mg/kg) (x10,000). Arrows indicates mitochondria and vacuoles in mitochondria. **d** TiO₂ (50 mg/kg) (x 10,000)

in ruptured and cracked nerve cells in the brain tissue (Fig. 8c). Treatment of mice with 150 mg/kg of nano-TiO₂ resulted in invasion of inflammatory cells in the brain tissue (Fig. 8d). The results showed that high doses of nano-TiO₂ can cause morphological damage to brain tissue, resulting in an inflammatory reaction.

We determined the effects of nano-TiO₂ on the redox state and signal molecules in brain tissue of mice exposed to nano-TiO₂ (i.p. for 14 days). We observed that a low dose (5 mg/kg) nano-TiO₂ did not change O²⁻, H₂O₂ and MDA levels, did not change the activities of the antioxidant enzymes APX, CAT, GSHPx, and SOD,

or the levels of non-enzymatic antioxidants ASA/DASA and GSH/GSSG. Nor did it change nitric oxide synthase (NOS) activity and NO levels in brain tissues (Figs. 9 and 10). Higher doses of nano-TiO₂ increased O²⁻, H₂O₂, and MDA levels, decreased the activities of the antioxidant enzymes APX, CAT, GSHPx, and SOD, decreased the levels of the non-enzymatic antioxidants ASA/DASA and GSH/GSSG, increased the levels of NO and activities of NOS, and decreased the levels of AChE and blood glucose (GLU) in the brain tissues (Figs. 9 and 10). These results suggest that nano-TiO₂ may cause damage in the brain in mice after i.p. exposure.





Fig. 6 ROS production rates and lipid peroxidation levels in livers of mice exposed to nano-TiO₂. Mice were treated with nano-TiO₂ as indicated through gavage once per day for 60 days. The mice in the control group received 0.5% CMC (carboxymethyl cellulose). * compared to control, *P* < 0.05, normalized to total protein




Toxic Effect of Nano-TiO₂ on Ex Vivo Mouse Embryos

To evaluate the developmental toxicity of nano-TiO₂, we first compared the growth and development of in vivo embryos and ex vivo embryos. The results showed that the growth and development of ex vivo embryos was similar to those of in vivo embryos (data not shown). Therefore, we used ex vivo embryos to study the toxic effects of nano-TiO₂ on embryos.

We investigated the effects of different doses (final concentrations 0.0, 50.0, 100.0, and 200.0 μg/mL) and different exposure times of micro-TiO₂/nano-TiO₂ on embryonic growth and development as well as the

morphology of tissues and organs through examining the embryonic VXY diameter, crown-rump length, head length, and the number of body sections. The results showed that the micro-TiO₂ did not change these indicators at any dose (Table 2).

For different sizes of nano-TiO₂, treatment of embryos of 5–10 nm, 60 nm, 90 nm with 50.0 μg/mL TiO₂ had no effect on embryo VXY diameter, crown-rump length, head length, and the number of body sections (Table 2). Higher doses (100.0 and 200.0 μg/mL) decreased VXY diameter, crown-rump length, head length, and number of body sections, and increased malformation rate (Table 2).







Table 2 Effects of TiO₂ of different sizes at different concentration on the growth and development of mouse in vitro embryos (N = 15)

TiO ₂ concentration	VXY diameter	CRL (mm)	Head length (mm)	The number of body sections	Malformation rate (%)	Death rate (%)
micro-TiO ₂						
0.0	4.37 ± 0.08	3.35 ± 0.06	1.45 ± 0.04	30.68 ± 1.65	0	0
50.0	4.36 ± 0.07	3.34 ± 0.13	1.49 ± 0.12	30.46 ± 1.27	20.0*	0
100.0	4.46 ± 0.09	3.39 ± 0.08	1.46 ± 0.11	30.82 ± 1.49	33.3*	0
200.0	4.51 ± 0.11	3.38 ± 0.09	1.46 ± 0.08	30.68 ± 1.29		
5–10 nm						
0.0	4.36 ± 0.07	3.34 ± 0.05	1.46 ± 0.05	30.78 ± 1.65	0	0
50.0	4.37 ± 0.06	3.37 ± 0.8	1.47 ± 0.8	30.63 ± 1.47	0	0
100.0	4.01 ± 0.24*	2.59 ± 0.28*	1.16 ± 0.08#	22.24 ± 3.47*	20.0*	0
200.0	3.21 ± 0.28 [#]	2.36 ± 0.49 [#]	1.06 ± 0.12 [#]	17.68 ± 4.22 [#]	33.3*	0
60 nm						
0.0	4.38 ± 0.07	3.35 ± 0.06	1.46 ± 0.06	30.81 ± 1.72	0	0
50.0	4.37 ± 0.06	3.36 ± 0.05	1.47 ± 0.08	30.63 ± 2.14	0	0
100.0	3.41 ± 0.34*	2.51 ± 0.32*	1.22 ± 0.18#	24.17 ± 4.61 [#]	6.67	0
200.0	4.32 ± 0.26	2.29 ± 0.51*	0.96 ± 0.21 [#]	15.36 ± 4.71 [#]	60.0*	0
90 nm						
0.0	4.37 ± 0.07	3.35 ± 0.05	1.46 ± 0.06	30.91 ± 1.67	0	0
50.0	4.38 ± 0.06	3.38 ± 0.11	1.47 ± 0.08	30.13 ± 1.61	0	0
100.0	4.09 ± 0.14*	2.89 ± 0.02*	1.19 ± 0.11 [#]	19.17 ± 4.56 [#]	52.7 [#]	6.67
200.0	3.72 ± 0.21 [#]	2.52 ± 0.21 [#]	1.08 ± 0.15 [#]	14.67 ± 3.57 [#]	74.1 [#]	26.67*

Compared to control; *P < 0.05, [#]P < 0.01

For the same dose, there was no obvious difference among groups treated with different sizes of nano-TiO₂, 50.0, or 100 µg/mL. Treatment of embryos with 200 µg/mL nano-TiO₂ significantly decreased VXY diameter, crown–rump length, head length, and the number of body sections of mice embryos with the increasing the size of nano-TiO₂ particles (Table 2).

Treatment of mice embryos with micro-TiO₂ (200.0 µg/mL) for 16, 24, and 48 h did not change VXY diameter, crown–rump length, head length, and the number of body sections (Table 3). Treatment of mice embryos with nano-TiO₂ (5–10 nm and 60 nm, 90 nm, 200.0 µg/mL) for 16 h also did not change VXY diameter, crown–rump length, head length, and the number of body sections (Table 3). However, the treatment of mice embryos with nano-TiO₂ (5–10 nm and 60 nm, 90 nm, 200.0 µg/mL) for 24 and 48 h decreased VXY diameter, crown–rump length, head length, the number of body sections, and increased the malformation rate (Table 3). For the same exposure time, there was no difference in VXY diameter, crown–rump length, head length, the number of body sections, or malformation rate among groups of different sizes of nano-TiO₂ particles (Table 3).

In summary, these results indicate that nano-TiO₂ had toxic effects on the growth and development of mouse

embryos in dose-dependent and time-dependent manners. The sizes of the nano-TiO₂ particles may affect toxicity with a trend of increasing toxicity associated with larger nano-TiO₂ particles.

Discussion

Nano-TiO₂ has been widely used in industry and medicine. However, the safety of nano-TiO₂ exposure remains unclear. In the present study, we investigated the potential toxicity of nano-TiO₂, using mice models. We find that nano-TiO₂ accumulates in the heart, liver, kidney, spleen, lung, and brain of mice after exposure (i.p. injection) in a dose-dependent manner. High doses of nano-TiO₂ significantly increase the organ/body weight ratios of the liver, kidney, spleen, and heart, and decrease those of the lung and brain in a dose-dependent manner. Moreover, high doses of nano-TiO₂ significantly increase the levels of ALT, ALP, LAP, PChE, TP, ALB, and TBIL, which are indices for liver function. They decrease the levels of UA and BUN, which are renal function indicators. Further, high doses significantly increase the activities of CK, LDH, AST, and HBDH, and significantly increase the levels of GLU, triglyceride, total cholesterol, and high-density lipoprotein. Low doses of nano-TiO₂ do not change these biochemical parameters. Our data support that nano-TiO₂ may be toxic and may affect the

Table 3 Effects of TiO₂ of different sizes at different exposure time on the growth and development of mouse in vitro embryos

DOE (h)	Embryos (n)	VXY diameter	CRL (mm)	Head length (mm)	The number of body sections	Neubert scores	Malformation rate (%)
Micro-TiO ₂							
0	15	4.38 ± 0.08	3.35 ± 0.09	1.45 ± 0.04	30.76 ± 1.21	54.36 ± 1.32	0
16	12	4.39 ± 0.11	3.35 ± 0.13	1.47 ± 0.12	30.81 ± 1.13	54.14 ± 1.16	0
26	12	4.36 ± 0.15	3.36 ± 0.12	1.49 ± 0.11	30.62 ± 1.26	53.28 ± 1.08	0
48	15	4.41 ± 0.13	3.38 ± 0.11	1.49 ± 0.08	30.42 ± 1.22	53.45 ± 1.22	0
5–10 nm							
0	15	4.37 ± 0.06	3.35 ± 0.06	1.46 ± 0.06	30.86 ± 1.61	54.43 ± 1.38	0
16	12	4.35 ± 0.04	3.03 ± 0.07	1.47 ± 0.12	30.43 ± 1.32	53.12 ± 1.26	13.33
26	12	3.93 ± 0.07*	2.76 ± 0.05*	1.14 ± 0.07*	25.26 ± 1.31*	41.32 ± 1.38*	75.00 [#]
48	15	3.01 ± 0.06 [#]	2.08 ± 0.08 [#]	0.94 ± 0.08 [#]	18.34 ± 1.12 [#]	26.24 ± 1.43 [#]	94.44 [#]
60 nm							
0	15	4.39 ± 0.06	3.36 ± 0.07	1.47 ± 0.08	30.90 ± 1.62	54.13 ± 1.31	0
16	12	4.36 ± 0.04	3.13 ± 0.13	1.22 ± 0.17*	30.63 ± 1.45	52.31 ± 1.26	8.33
24	12	4.05 ± 0.09*	2.77 ± 0.08 [#]	1.15 ± 0.06*	27.68 ± 1.30*	41.32 ± 1.48*	66.67*
48	15	3.04 ± 0.11 [#]	2.13 ± 0.16 [#]	0.94 ± 0.78 [#]	19.04 ± 1.07 [#]	27.24 ± 1.15 [#]	100.00 [#]
90 nm							
0	15	4.39 ± 0.06	3.36 ± 0.06	1.47 ± 0.08	30.90 ± 1.62	54.13 ± 1.31	0
16	12	4.37 ± 0.04	3.35 ± 0.08	1.22 ± 0.17*	30.63 ± 1.45	52.31 ± 1.26	0
24	12	4.25 ± 0.06	3.08 ± 0.07 [#]	1.15 ± 0.06*	27.68 ± 1.30*	41.32 ± 1.48*	56.67*
48	15	3.34 ± 0.09 [#]	2.29 ± 0.05 [#]	0.94 ± 0.78 [#]	19.04 ± 1.07 [#]	27.24 ± 1.15 [#]	86.67 [#]

DOE duration of exposure;

Compared to controls, **P* < 0.05; [#]*P* < 0.01

liver, kidney, heart, GLU, and lipid metabolism at high doses in a dose-dependent manner.

In the present study, we investigated the mechanism of liver toxicity of nano-TiO₂. We find that high doses of nano-TiO₂ may cause swelling of hepatocytes with obvious vacuoles in cells, and nuclear condensation in hepatocytes, and apoptosis and necrosis of hepatocytes in liver tissues. This is consistent with previous studies [7, 23, 24]. After the treatment of mice with high doses of nano-TiO₂, we find that the levels of CAT, GSHPx, and SOD are significantly decreased, and there is nano-TiO₂ in the mitochondria of hepatocytes, revealed by TEM. This is consistent with previous studies [7, 25–28] suggesting that nano-TiO₂ generates excess ROS and reduces the antioxidant capacity of the cells through damaging the mitochondria. This is further supported by observation that nano-TiO₂ can significantly decrease the mRNA levels of SOD, CAT, GSHPx, MT and HSP70, CYP1A1, p53, GST, and TF genes in the mouse liver. SOD, CAT, GSH PX, and MT are involved in liver cell detoxification, CYP1A1 is involved in toxic-substance metabolism and defense against invasion from harmful substances, and HSP70 and p53 are involved in repairing liver cell DNA damage [10, 29–39]. These findings

support that the mechanisms for nano-TiO₂ liver toxicity are damaging mitochondria, generating ROS, and causing expression disorders of protective genes.

In the present study, we investigated the mechanism of brain neurotoxicity of nano-TiO₂. We find that high doses of nano-TiO₂ can produce lipid peroxidation and decrease antioxidant capacity, including SOD, CAT, APX, and GSHPx activities, resulting in oxidative stress, which may damage unsaturated fatty acids and brain tissue [24, 26, 37, 40]. We observed rupture and cracking in nerve cells and the infiltration of inflammatory cells in the brain. We further found that the activities of cNOS and iNOS are increased, and NO is excessively released. Glutamic acid levels and AChE activity are decreased in the brain. This is consistent with the effect of Fe₂O₃ nanoparticles on olfactory bulb cells [40] and the effect of nano-TiO₂ on mouse hippocampal neurons [31, 41]. Glutamate is the most abundant amino acid in excitatory neurotransmitters of the nervous system. It is critical for the brain's development and function [42]. Acetylcholinesterase is a key enzyme for levels of acetylcholine, which is critical for the function of the peripheral and central nervous systems. Nitric oxide regulates many central nervous functions, such as synaptic plasticity,

the sleep–wake cycle, and hormone secretion [43]. Therefore, nano-TiO₂ may cause oxidative stress and may disrupt orders of neurochemical metabolism in brain tissue and therefore have neurotoxicity in the central nervous system.

We find that the micro-TiO₂ and low doses of nano-TiO₂ (5–10 nm and 60 nm and 90 nm) do not exhibit toxicity on ex vivo mouse embryos, while high doses of nano-TiO₂ (100–200.0 µg/ml) exhibit toxicity on ex vivo mouse embryos, as revealed by evaluation of morphology of exposed embryos. Whole embryo culture is a useful tool to assess the developmental toxicity of chemicals [44, 45]. Previous studies show that exposure of 14-day pregnant mice to a single dose of nano-TiO₂ in the nasal cavity increase the sensitivity of inflammatory response in F1 generation [46, 47]. Nano-TiO₂ does not affect white pregnant Kunming mice but inhibits growth, increases the rate of stillbirth, and exhibits developmental toxicity [48]. These studies indicate the presence of the developmental and genetic toxicity of nano-TiO₂. This is further supported by studies that show cleavage and oxidative damage of DNA by nano-TiO₂, for example, in Zebra fish [16, 49, 50]. Additionally, another shows an increase in the sister chromatid exchange rates in Chinese hamster ovary cells [51]. Nano-TiO₂ may also prevent chromosome formation during metaphase in the ovary when TiO₂ concentration is high [51]. These studies consistently show that exposure to high doses of nano-TiO₂ is linked with developmental and genetic toxicity. Furthermore, our data indicated that the size of nano-TiO₂ particles may affect its toxicity, with the trend of increasing toxicity being associated with larger nano-TiO₂ particles (Table 2).

In the current study, we found that titanium accumulates in a dose-dependent manner in the heart, liver, kidney, spleen, lung, and brain of mice after i.p. injection of nano-TiO₂. This is consistent with published reports that absorption and distribution of nano-TiO₂ is dependent on blood circulation. Nanoparticles can be absorbed in mesenchymal cells through being ingested by airway epithelial cells; they can then penetrate into the blood or lymph, thus gradually being distributed to the whole body [52, 53]. It is worth noting that nano-TiO₂ in the abdomen cavity can be absorbed and transported to the brain by the circulatory system, and nano-TiO₂ can enter directly into the central nervous system without crossing the blood–brain barrier. This is consistent with previous studies [41, 54]. Nanoparticles can also be absorbed by the terminal nerve cell in the respiratory tract and then be transferred to the ganglion through the axon, eventually entering central nervous cells [8, 55]. Nano-TiO₂ can be absorbed in the nasal cavity through the olfactory epithelium, and then be transported to other parts of the brain, such as the hippocampus, through the olfactory nerve [41, 54].

Therefore, the brain may be directly exposed to nano-TiO₂. Damage in the brain may be caused directly or indirectly by nano-TiO₂.

Conclusions

Ingested nano-TiO₂ can be distributed to and accumulated in the heart, brain, spleen, lung, and kidney. It exhibits toxicity and causes disorders of the GLU and lipid metabolism. Nano-TiO₂ causes liver and brain toxicity mainly through increasing oxidative stress, decreasing antioxidant levels, and changing the expression of the protective genes in the liver. In addition, nano-TiO₂ has adverse effects on the growth and development of mouse embryos and the morphology of the tissues and organs. The size of nano-TiO₂ particles may affect their toxicity, with a trend of increasing toxicity being associated with larger nano-TiO₂ particles. These toxic effects are dose-dependent. Our study may provide data for the assessment of the risk of nano-TiO₂ exposure on human health.

Abbreviations

ALB: Albumin; ALP: Alkaline phosphatase; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; BUN: Blood urea nitrogen; CK: Creatine kinase; cNOS: Constitutive nitric oxide synthases; HBDH: Hydroxybutyrate dehydrogenase; ICR: Imprinting control region; iNOS: Inducible NOS; LAP: Qleucine aminopeptidase; LDH: Lactate dehydrogenase; Nano-TiO₂: Nanoscale titanium dioxide; PChe: Butyrylcholinesterase; ROS: Reactive oxygen species; TBIL: Total bilirubin; TP: Total protein; UA: Uric acid

Acknowledgements

None.

Availability of Data and Materials

The datasets supporting the conclusions of this article are included within the article.

Authors' Contributions

XCJ and SW designed the study. XCJ, SW, LZ, and LS collected and analyzed the data. XCJ, SW, LZ, and LS contributed samples collection and intellectual input. XCJ and SW drafted and wrote the manuscript. SW revised the manuscript critically for intellectual content. All authors gave intellectual input to the study and approved the final version of the manuscript.

Competing Interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Tianjin University of Science and Technology, Tianjin 300457, China. ²Technical Center for Safety of Industrial Products of Tianjin Entry-Exit Inspection and Quarantine Bureau, Tianjin 300308, China.

Received: 10 October 2016 Accepted: 30 November 2016

Published online: 02 August 2017

References

1. Weir A, Westerhoff P, Fabricius L, Hristovski K, von Goetz N (2012) Titanium dioxide nanoparticles in food and personal care products. *Environ Sci Technol* 46:2242–2250
2. Chen XX, Cheng B, Yang YX, Cao A, Liu JH, Du LJ et al (2013) Characterization and preliminary toxicity assay of nano-titanium dioxide additive in sugar-coated chewing gum. *Small* 9:1765–1774

3. Peters RJ, van Bommel G, Herrera-Rivera Z, Helsper HP, Marvin HJ, Weigel S et al (2014) Characterization of titanium dioxide nanoparticles in food products: analytical methods to define nanoparticles. *J Agric Food Chem* 62:6285–6293
4. Riehemann K, Schneider SW, Luger TA, Godin B, Ferrari M, Fuchs H (2009) Nanomedicine—challenge and perspectives. *Angew Chem Int Ed Eng* 48:872–897
5. Laurencin CT, Kumbar SG, Nukavarapu SP (2009) Nanotechnology and orthopedics: a personal perspective. *Wiley Interdiscip Rev Nanomed Nanobiotechnol* 1:6–10
6. Karaca G, Tasdemir Y (2014) Migration of PAHs in food industry sludge to the air during removal by UV and TiO₂. *Sci Total Environ* 488–489:356–361
7. Cui Y, Gong X, Duan Y, Li N, Hu R, Liu H et al (2010) Hepatocyte apoptosis and its molecular mechanisms in mice caused by titanium dioxide nanoparticles. *J Hazard Mater* 183:874–880
8. Hu R, Gong X, Duan Y, Li N, Che Y, Cui Y et al (2010) Neurotoxicological effects and the impairment of spatial recognition memory in mice caused by exposure to TiO₂ nanoparticles. *Biomaterials* 31:8043–8050
9. Wang JJ, Sanderson BJ, Wang H (2007) Cyto- and genotoxicity of ultrafine TiO₂ particles in cultured human lymphoblastoid cells. *Mutat Res* 628:99–106
10. Wang L, Mao J, Zhang GH, Tu MJ (2007) Nano-cerium-element-doped titanium dioxide induces apoptosis of Bel 7402 human hepatoma cells in the presence of visible light. *World J Gastroenterol* 13:4011–4014
11. Pogue AI, Jones BM, Bhattacharjee S, Percy ME, Zhao Y, Lukiw WJ (2012) Metal-sulfate induced generation of ROS in human brain cells: detection using an isomeric mixture of 5- and 6-Carboxy-2',7'-Dichlorofluorescein Diacetate (Carboxy-DCFDA) as a cell Permeant tracer. *Int J Mol Sci* 13:9615–9626
12. Wu J, Sun J, Xue Y (2010) Involvement of JNK and P53 activation in G2/M cell cycle arrest and apoptosis induced by titanium dioxide nanoparticles in neuron cells. *Toxicol Lett* 199:269–276
13. Geiser M, Rothen-Rutishauser B, Kapp N, Schurch S, Kreyling W, Schulz H et al (2005) Ultrafine particles cross cellular membranes by nonphagocytic mechanisms in lungs and in cultured cells. *Environ Health Perspect* 113:1555–1560
14. Ma L, Ze Y, Liu J, Liu H, Liu C, Li Z et al (2009) Direct evidence for interaction between nano-anatase and superoxide dismutase from rat erythrocytes. *Spectrochim Acta A Mol Biomol Spectrosc* 73:330–335
15. Ma L, Zhao J, Wang J, Liu J, Duan Y, Liu H et al (2009) The acute liver injury in mice caused by Nano-Anatase TiO₂. *Nanoscale Res Lett* 4:1275–1285
16. Hirakawa K, Mori M, Yoshida M, Oikawa S, Kawanishi S (2004) Photo-irradiated titanium dioxide catalyzes site specific DNA damage via generation of hydrogen peroxide. *Free Radic Res* 38:439–447
17. Reeves JF, Davies SJ, Dodd NJ, Jha AN (2008) Hydroxyl radicals (*OH) are associated with titanium dioxide (TiO₂) nanoparticle-induced cytotoxicity and oxidative DNA damage in fish cells. *Mutat Res* 640:113–122
18. Goncalves DM, Chiasson S, Girard D (2010) Activation of human neutrophils by titanium dioxide (TiO₂) nanoparticles. *Toxicol in Vitro* 24:1002–1008
19. Kang SJ, Kim BM, Lee YJ, Hong SH, Chung HW (2009) Titanium dioxide nanoparticles induce apoptosis through the JNK/p38-caspase-8-bid pathway in phytohemagglutinin-stimulated human lymphocytes. *Biochem Biophys Res Commun* 386:682–687
20. Jambunathan N (2010) Determination and detection of reactive oxygen species (ROS), lipid peroxidation, and electrolyte leakage in plants. *Methods Mol Biol* 639:292–298
21. Auer T, Khoschsorur GA, Rabl H, Iberer F, Petutschnigg B, Wasler A et al (1995) Detection of lipid peroxidation products by malondialdehyde (MDA-TBA reaction) in organ transplantation. *Transplant Proc* 27:2749–2751
22. Van Maele-Fabry G, Delhaise F, Picard JJ (1990) Morphogenesis and quantification of the development of post-implantation mouse embryos. *Toxicol in Vitro* 4:149–156
23. Duan Y, Liu J, Ma L, Li N, Liu H, Wang J et al (2010) Toxicological characteristics of nanoparticulate anatase titanium dioxide in mice. *Biomaterials* 31:894–899
24. Cui Y, Liu H, Ze Y, Zengli Z, Hu Y, Cheng Z et al (2012) Gene expression in liver injury caused by long-term exposure to titanium dioxide nanoparticles in mice. *Toxicol Sci* 128:171–185
25. Paunesku T, Vogt S, Lai B, Maser J, Stojicevic N, Thurn KT et al (2007) Intracellular distribution of TiO₂-DNA oligonucleotide nanoconjugates directed to nucleolus and mitochondria indicates sequence specificity. *Nano Lett* 7:596–601
26. Zhang L, Liang Z, Yang K, Xia S, Wu Q, Zhang L et al (2012) Mesoporous TiO₂ aerogel for selective enrichment of phosphopeptides in rat liver mitochondria. *Anal Chim Acta* 729:26–35
27. Zhang R, Niu Y, Du H, Cao X, Shi D, Hao Q et al (2009) A stable and sensitive testing system for potential carcinogens based on DNA damage-induced gene expression in human HepG2 cell. *Toxicol in Vitro* 23:158–165
28. Yang Z, Zhang Y, Yang Y, Sun L, Han D, Li H et al (2010) Pharmacological and toxicological target organelles and safe use of single-walled carbon nanotubes as drug carriers in treating Alzheimer disease. *Nanomedicine* 6:427–441
29. Rahman Q, Lohani M, Dopp E, Pemsel H, Jonas L, Weiss DG et al (2002) Evidence that ultrafine titanium dioxide induces micronuclei and apoptosis in Syrian hamster embryo fibroblasts. *Environ Health Perspect* 110:797–800
30. Gretzer C, Gisselalt K, Liljensten E, Ryden L, Thomsen P (2003) Adhesion, apoptosis and cytokine release of human mononuclear cells cultured on degradable poly(urethane urea), polystyrene and titanium in vitro. *Biomaterials* 24:2843–2852
31. Wang ML, Tuli R, Manner PA, Sharkey PF, Hall DJ, Tuan RS (2003) Direct and indirect induction of apoptosis in human mesenchymal stem cells in response to titanium particles. *J Orthop Res* 21:697–707
32. Larsson J, Persson C, Tengvall P, Lundqvist-Gustafsson H (2004) Anti-inflammatory effects of a titanium-peroxy gel: role of oxygen metabolites and apoptosis. *J Biomed Mater Res A* 68:448–457
33. Park EJ, Yi J, Chung KH, Ryu DY, Choi J, Park K (2008) Oxidative stress and apoptosis induced by titanium dioxide nanoparticles in cultured BEAS-2B cells. *Toxicol Lett* 180:222–229
34. Liu S, Xu L, Zhang T, Ren G, Yang Z (2010) Oxidative stress and apoptosis induced by nanosized titanium dioxide in PC12 cells. *Toxicology* 267:172–177
35. Shi Y, Wang F, He J, Yadav S, Wang H (2010) Titanium dioxide nanoparticles cause apoptosis in BEAS-2B cells through the caspase 8/t-bid-independent mitochondrial pathway. *Toxicol Lett* 196:21–27
36. Shukla RK, Kumar A, Pandey AK, Singh SS, Dhawan A (2011) Titanium dioxide nanoparticles induce oxidative stress-mediated apoptosis in human keratinocyte cells. *J Biomed Nanotechnol* 7:100–101
37. Xue C, Liu W, Wu J, Yang X, Xu H (2011) Chemoprotective effect of N-acetylcysteine (NAC) on cellular oxidative damages and apoptosis induced by nano titanium dioxide under UVA irradiation. *Toxicol in Vitro* 25:110–116
38. Marquez-Ramirez SG, Delgado-Buenrostro NL, Chirino YI, Iglesias GG, Lopez-Marure R (2012) Titanium dioxide nanoparticles inhibit proliferation and induce morphological changes and apoptosis in glial cells. *Toxicology* 302:146–156
39. Srivastava RK, Rahman Q, Kashyap MP, Singh AK, Jain G, Jahan S et al (2013) Nano-titanium dioxide induces genotoxicity and apoptosis in human lung cancer cell line, A549. *Hum Exp Toxicol* 32:153–166
40. Wang L, Yang L, Yang F, Li X, Song Y, Wang X et al (2010) Involvements of H₂O₂ and metallothionein in NO-mediated tomato tolerance to copper toxicity. *J Plant Physiol* 167:1298–1306
41. Wang J, Zhou G, Chen C, Yu H, Wang T, Ma Y et al (2007) Acute toxicity and biodistribution of different sized titanium dioxide particles in mice after oral administration. *Toxicol Lett* 168:176–185
42. Zhou Y, Danbolt NC (2014) Glutamate as a neurotransmitter in the healthy brain. *J Neural Transm* 121:799–817
43. Calabrese V, Mancuso C, Calvani M, Rizzarelli E, Butterfield DA, Stella AM (2007) Nitric oxide in the central nervous system: neuroprotection versus neurotoxicity. *Nat Rev Neurosci* 8:766–775
44. Klug S, Neubert D (1993) The use of whole embryo culture to elucidate teratogenic mechanisms. *Toxicol in Vitro* 7:727–734
45. Stahlmann R, Klug S, Foerster M, Neubert D (1993) Significance of embryo culture methods for studying the prenatal toxicity of virustatic agents. *Reprod Toxicol* 7(Suppl 1):129–143
46. Mohammadipour A, Fazel A, Haghiri H, Motejaded F, Rafatpanah H, Zabihi H et al (2014) Maternal exposure to titanium dioxide nanoparticles during pregnancy; impaired memory and decreased hippocampal cell proliferation in rat offspring. *Environ Toxicol Pharmacol* 37:617–625
47. Yamashita K, Yoshioka Y, Higashisaka K, Mimura K, Morishita Y, Nozaki M et al (2011) Silica and titanium dioxide nanoparticles cause pregnancy complications in mice. *Nat Nanotechnol* 6:321–328
48. Shimizu M, Tainaka H, Oba T, Mizuo K, Umezawa M, Takeda K (2009) Maternal exposure to nanoparticulate titanium dioxide during the prenatal period alters gene expression related to brain development in the mouse. *Part Fibre Toxicol* 6:20

49. Singh N, Manshian B, Jenkins GJ, Griffiths SM, Williams PM, Maffei TG et al (2009) NanoGenotoxicology: the DNA damaging potential of engineered nanomaterials. *Biomaterials* 30:3891–3914
50. Ze Y, Hu R, Wang X, Sang X, Ze X, Li B et al (2014) Neurotoxicity and gene-expressed profile in brain-injured mice caused by exposure to titanium dioxide nanoparticles. *J Biomed Mater Res A* 102:470–478
51. Gao G, Ze Y, Zhao X, Sang X, Zheng L, Ze X et al (2013) Titanium dioxide nanoparticle-induced testicular damage, spermatogenesis suppression, and gene expression alterations in male mice. *J Hazard Mater* 258-259:133–143
52. Huang H, Liu X, Qu C, Liu C, Chen L, Hong F (2008) Influences of calcium deficiency and cerium on the conversion efficiency of light energy of spinach. *Biometals* 21:553–561
53. Zhang W, Zhang Z, Zhang Y (2011) The application of carbon nanotubes in target drug delivery systems for cancer therapies. *Nanoscale Res Lett* 6:555
54. Baggs RB, Ferin J, Oberdorster G (1997) Regression of pulmonary lesions produced by inhaled titanium dioxide in rats. *Vet Pathol* 34:592–597
55. Zhao H, Cheng Z, Cheng J, Hu R, Che Y, Cui Y et al (2011) The toxicological effects in brain of mice following exposure to cerium chloride. *Biol Trace Elem Res* 144:872–884

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
