



# The potential of anti-malarial compounds derived from African medicinal plants. Part I: A pharmacological evaluation of alkaloids and terpenoids

Amoa Onguéné *et al.*

REVIEW

Open Access

# The potential of anti-malarial compounds derived from African medicinal plants. Part I: A pharmacological evaluation of alkaloids and terpenoids

Pascal Amoa Onguéné<sup>1†</sup>, Fidele Ntie-Kang<sup>2†</sup>, Lydia Likowo Lifongo<sup>2</sup>, Jean Claude Ndom<sup>1</sup>, Wolfgang Sippl<sup>3</sup> and Luc Meva'a Mbaze<sup>1\*</sup>

## Abstract

Traditional medicine caters for about 80% of the health care needs of many rural populations around the world, especially in developing countries. In addition, plant-derived compounds have played key roles in drug discovery. Malaria is currently a public health concern in many countries in the world due to factors such as chemotherapy faced by resistance, poor hygienic conditions, poorly managed vector control programmes and no approved vaccines. In this review, an attempt has been made to assess the value of African medicinal plants for drug discovery by discussing the anti-malarial virtue of the derived phytochemicals that have been tested by *in vitro* and *in vivo* assays. This survey was focused on pure compounds derived from African flora which have exhibited anti-malarial properties with activities ranging from "very active" to "weakly active". However, only the compounds which showed anti-malarial activities from "very active" to "moderately active" are discussed in this review. The activity of 278 compounds, mainly alkaloids, terpenoids, flavonoids, coumarines, phenolics, polyacetylenes, xanthenes, quinones, steroids, and lignans have been discussed. The first part of this review series covers the activity of 171 compounds belonging to the alkaloid and terpenoid classes. Data available in the literature indicated that African flora hold an enormous potential for the development of phytomedicines for malaria.

**Keywords:** Africa, Malaria, Medicinal plants, Natural products, Traditional medicine

## Background

Malaria is an infectious disease with ravaging effects in the world. The World Health Organization (WHO) has published statistics which reveal that half the world's population is at risk of malaria and that one to two million annual deaths can be attributed to malaria alone [1,2]. Four protozoan species of the genus *Plasmodium* (*Plasmodium falciparum*, *Plasmodium malariae*, *Plasmodium ovale*, and *Plasmodium vivax*) are responsible for this infection, although the majority of fatal cases are caused by *P. falciparum* [3]. Malaria has been treated

with quinine, chloroquine, mefloquine, and artemisinin (Figure 1), among other drugs. However, the protozoans have developed resistance against many of the current treatment regimens [4]. In the quest to identify new anti-malarial chemotherapeutic agents, many research groups have resorted to plant sources [3,5,6]. This is because of the use of many of these plant materials in the treatment of malaria and fevers in African traditional medicine (ATM) [7]. There has been a general call for the use of natural products as drugs for malaria or as sources of inspiration for the development of novel anti-malarials [8-11] in order to possibly avoid problems related to drug resistance [12].


The African continent is very rich in floral biodiversity and its plant materials are endowed with natural products (NPs) with intriguing chemical structures and promising

\* Correspondence: lmbazze@yahoo.fr

†Equal contributors

<sup>1</sup>Department of Chemistry, Faculty of Science, University of Douala, PO Box 24157, Douala, Cameroon

Full list of author information is available at the end of the article



biological activities. Therefore, the next generation anti-malarials or the scaffolds necessary for their synthesis may be found in plants currently used in ATM [13,14]. It should also be mentioned that malaria mostly affects the populations of Africa, Asia and Latin America. Asia has offered artemisinin to humanity while Latin America has offered quinine. Many researchers are therefore of the opinion that it is Africa's turn to offer a new anti-malarial drug to humanity. Why do we not yet find a (real) anti-malarial drug from Africa? This brings us to the need to have an overview of the anti-malarial/anti-plasmodial activity of compounds from bitter African plants (alkaloids and terpenoids). Several research groups in Africa have been involved in the bioassay-guided fractionation of plant extracts, leading to the isolation, purification and characterization of a significant number of NPs, some with remarkable anti-malarial activities. The literature survey reported in this work has led to the identification of several vast screening efforts of crude extracts derived from plants used in ATM, harvested from the following countries, just to mention a few: the Democratic Republic of Congo [15,16], Nigeria [17-19], Mozambique, Cape Verde, Guinea-Bissau, São Tomé and Príncipe and Angola [20], Mali and São Tomé and Príncipe [21], Madagascar [22-24], Congo [25], Benin [26], Burkina Faso [27], South Africa [28], Ivory Coast [29], West African countries [30], Tanzania [31], Kenya [32], and East African countries [33,34].

The potential of plant-derived NPs for anti-malarial drug discovery has been examined in a number of review papers [3,35-41]. Other review articles have concentrated on anti-malarials from specific countries/regions in Africa [19,20,42-46]. However, there has been no review offering coverage of promising anti-malarials from the entire African continent in the last ten years [13]. In this review series, the potential of plant-derived NPs that could be developed into drugs have been discussed, by giving an overview of the most pertinent *in vitro* and *in vivo* screening results reported in the literature.

### Promising anti-malarial alkaloids and terpenoids derived from African flora

#### Alkaloids

Previous studies have shown that plant-derived alkaloids have a great potential for anti-malarial drug development [33-36,39,40,42-44]. Tables 1 and 2 summarize the most promising alkaloids derived from African medicinal plants with significant anti-malarial properties. The chemical structures are shown in Figures 2, 3, 4, 5, 6 and 7, according to the alkaloid subclasses.

#### Indole alkaloids

Several indole alkaloids, derived from African medicinal plants, have shown interesting *in vitro* anti-malarial activities, among them compounds **1**, **2** and **10** to **19**. Nkunya *et al.* have isolated prenylated indole alkaloids from *Monodora* and *Isolona* species (Annonaceae) growing in Tanzania [47]. According to their report, 6-(3-methyl-but-2-enyl)-1,3-dihydro-indol-2-one (**1**), 3-[6-(3-methyl-but-2-enyl)-1H-indolyl]-6-(3-methyl-but-2-enyl)-1H-indole or annonidine F (**2**), 1H-indole-5-carbaldehyde (**3**), 6-(3-methyl-2-butenyl)-1H-indole (**4**), 6-(3-methylbuta-1,3-dienyl)-1H-indole (**5**), 6-(4-oxo-but-2-enyl)-1H-indole (**6**) and 3-geranylindole (**7**) were isolated from *Monodora angolensis* while 3-(1,1-dimethyl-but-2-enyl)-5-(3-methyl-but-2-enyl)-1H-indole or caulidine A (**8**), 4-[3-(1,1-dimethyl-but-2-enyl)-1H-indol-5-yl]-but-3-en-2-one or caulidine B (**9**), 5-(3-methyl-2-butenyl)-1H-indole and 5-(3-methylbuta-1,3-dienyl)-1H-indole were obtained from *Isolona cauliflora*. The compounds with the most promising, measured, anti-malarial activities were **1** and **2**, both having *in vitro* anti-malarial activities against the multidrug resistant strain K1 of *P. falciparum* (IC<sub>50</sub> = 21 µg mL<sup>-1</sup> for each compound). Moreover, their measured cytotoxicities, in the brine shrimp test were IC<sub>50</sub> = 4.08 and 5.28 µg mL<sup>-1</sup>, respectively.

The compound 17-O-acetyl,10-hydroxycorynantheol (**10**) was isolated from *Strychnos usambarensis* (harvested in Rwanda), along with isostrychnopentamine (**18**), the main alkaloid responsible for the anti-plasmodial activity of the plant, by Cao *et al.* [48]. The study showed that

**Table 1 Summary of anti-malarial alkaloids derived from the African flora – indoles, naphthoisoquinolines and furoquinolines**

| Compound subclass            | Isolated metabolites          | Plant species (Family)   | Part of plant studied                                   | Place of harvest (locality, country)   | Author, reference  |
|------------------------------|-------------------------------|--|---|--|--|
| Indole alkaloids             | <b>1, 2, 3, 4, 5, 6 and 7</b> | <i>Monodora angolensis</i> (Annonaceae)  | Stem and root bark                                      | Kiwanda, Tanzania  | Nkunya et al. [47]   |
|                              | <b>8 and 9</b>                | <i>Isolona cauliflora</i> (Annonaceae)   | Stem and root bark, and flower stalks                   | Namikwe Island, Tanzania   | Nkunya et al. [47]   |
|                              | <b>10</b>                     | <i>Strychnos usambarensis</i> (Loganiaceae)  | Leaves  | Akagera National Park, Rwanda  | Cao et al. [48]  |
|                              | <b>11 and 12</b>              | <i>Penianthus longifolius</i> (Menispermaceae)   | Stem bark   | Cameroon   | Bidla et al. [49]  |
|                              | <b>13</b>                     | <i>Glossocalyx brevipes</i> (Siparunaceae)   | Leaves  | Kumba, Cameroon  | Mbah et al. [50]   |
|                              | <b>14</b>                     | <i>Fagara zanthoxyloides</i> (Rutaceae)  | Roots   | Nigeria  | Odebiyi et al. [51]  |
|                              | <b>15</b>                     | <i>Picralima nitida</i> (Apocynaceae)  | Fruits  | Nnewi, Nigeria   | Okunji et al. [52]   |
|                              | <b>16</b>                     | <i>Strychnos usambarensis</i> (Loganiaceae)  | Roots   | Akagera National Park, Rwanda  | Frédérich et al. [55]  |
|                              | <b>17, 18 and 19</b>          | <i>Strychnos usambarensis</i> (Loganiaceae)  | Leaves  | Akagera National Park, Rwanda  | Frédérich et al. [56]  |
|                              | Naphthoisoquinolines          | <b>20, 21, 22, 23 and 24</b>   | <i>Ancistrocladus robertsoniorum</i> (Acistrocladaceae) | Stems and leaves   | Buda Mafisini Forest, Kenya  |
| <b>25, 26, 27, 28 and 29</b> |                               | <i>Ancistrocladus tanzaniensis</i> (Acistrocladaceae)  | Leaves  | Uzungwa Mountains, Tanzania  | Bringmann et al. [58]  |
| <b>30</b>                    |                               | <i>Triphyophyllum peltatum</i> (Dioncophyllaceae)  | Roots   | Parc de Tai, West Ivory Coast  | Bringmann et al. [59]  |
| <b>31</b>                    |                               | <i>Triphyophyllum peltatum</i> (Dioncophyllaceae)  | Root bark   | West Ivory Coast   | Bringmann et al. [60]  |
| <b>32</b>                    |                               | Mixture of <i>Triphyophyllum peltatum</i> , <sup>a</sup> <i>Dioncophyllum thollonii</i> <sup>b</sup> and <i>Habropetalum dawefi</i> (Dioncophyllaceae) | Root and bark <sup>a</sup><br>Twigs <sup>b, c</sup>     | West Ivory Coast <sup>a</sup><br>Gabon <sup>b</sup><br>Sierra Leone <sup>c</sup> | Bringmann et al. [61]  |
| <b>33, 34, 35 and 36</b>     |                               | <i>Triphyophyllum peltatum</i> (Dioncophyllaceae)  | Leaves and twigs  | Mt. Nabemba, Congo Republic <sup>a</sup> and West Ivory Coast <sup>b</sup>       | Bringmann et al. [62-65] <sup>a</sup><br>François et al. [66] <sup>b</sup> |
| Furoquinolines               |                               | <b>37 and 38</b>   | <i>Vepris uguenensis</i> (Rutaceae)                     | Roots  | Baringo District, Kenya  |
|                              | <b>39 and 40</b>              | <i>Toddalia asiatica</i> (Rutaceae)  | Roots   | Ol Ari Nyiro Ranch, Kenya  | Gakunju et al. [69]  |
|                              | <b>41</b>                     | <i>Teclea gerrardii</i> (Rutaceae)   | Root bark   | Durban, South Africa   | Waffo et al. [70]  |

<sup>a,b</sup> and <sup>c</sup> correspond to the respective references.

compound **10** is one of the most promising, monomeric indole alkaloids known to date, showing an *in vitro* activity against *P. falciparum* close to 5  $\mu\text{M}$  and a high selectivity.

Indoles with interesting anti-malarial properties have also been derived from two plant species growing in Cameroon: *Penianthus longifolius* and *Glossocalyx brevipes* [49,50]. Bidla et al. have isolated palmitine (**11**) and jatrorrhizine (**12**) from the stem bark of *Penianthus longifolius*. Compounds **11** and **12** showed promising *in vitro* activities on various strains of *P. falciparum* with  $\text{IC}_{50}$  values ranging from 0.28 to 0.35  $\mu\text{g mL}^{-1}$  [49], meanwhile Mbah et al. isolated liriodenine (**13**) from *Glossocalyx brevipes* (Siparunaceae), which exhibited anti-malarial activity against the D-6 drug sensitive

strain from Sierra Leone and the NF54 strain with  $\text{IC}_{50}$  values of 2.37  $\mu\text{M}$  and 1.32  $\mu\text{M}$ , respectively [50].

From the plant species growing in Nigeria, fagaronine (**14**) and alstonine (**15**) were derived from *Fagara zanthoxyloides* (Rutaceae) and *Picralima nitida* (Apocynaceae) respectively [51,52]. While fagaronine (**14**) inhibited *P. falciparum* growth *in vitro* at  $\text{IC}_{50} = 0.018 \mu\text{g mL}^{-1}$ , alstonine (**15**) has been noted to be the most active indole alkaloid derived from *Picralima nitida* [53]. It is noteworthy that indole and dihydroindole alkaloids are common in *Picralima nitida* growing in Nigeria, the major constituents including akuammiline, akuammidine, akuammine, akuammigine, akuammicine, picraline, and alstonine [54]. Some of the aforementioned alkaloids have exhibited *in vitro* anti-malarial activity against *P. falciparum*

**Table 2 Summary of anti-malarial alkaloids derived from the African flora – acridones, amides and cryptolepines**

| Compound subclass | Isolated metabolites                                | Plant species (Family)                               | Part of plant studied  | Place of harvest (locality, country)                    | Author, reference  |
|-------------------|---|--|------------------------|---|--|
| Acridones         | 42 <sup>a, b, c</sup> and 43 <sup>b</sup>           | <i>Teclea gerrardii</i> <sup>a</sup>                 | Root bark <sup>a</sup> | Durban, South Africa <sup>a</sup>                       | Waffo et al. [70] <sup>a</sup>   |
|                   |   | <i>Zanthoxylum leprieurii</i> <sup>b</sup>           | Fruits <sup>b</sup>    | Yaoundé, Cameroon <sup>b</sup>                          | Tchinda et al. [71] <sup>b</sup>   |
|                   |   | <i>Teclea trichocarpa</i> <sup>c</sup><br>(Rutaceae) | Leaves <sup>c</sup>    |   |  |
|                   | 44, 45, 46, 47, 48 and 49                           | <i>Teclea trichocarpa</i> (Rutaceae)                 | Leaves                 | Nairobi, Kenya  | Wurithi et al. [72]  |
|                   | 50  | <i>Vepris uguenensis</i> (Rutaceae)                  | Roots                  | Baringo District, Kenya                                 | Cheplogoi et al. [68], Kiplimo [73]  |
| Amides            | 51  | <i>Hugonia castaneifolia</i> (Linaceae)              | Root bark              | Pugu forest, Tanzania                                   | Baraza et al. [74]   |
|                   | 52  | <i>Beilschmiedia zenkeri</i> (Lauraceae)             | Bark                   | Yaoundé, Cameroon                                       | Lenta et al. [75]  |
| Cryptolepines     | 53  | <i>Sida acuta</i> (Malvaceae)                        | Aerial parts           | Ivory Coast   | Banzounzi et al. [76]  |
|                   | 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, and 63' | <i>Cryptolepis sanguinolenta</i> (Periplocaceae)     | Stems <sup>a</sup>     | Mampong-Akwapim, Ghana, Guinea Bissau and other regions | Barku et al. [77] <sup>a</sup>   |
|                   |   |  | Root bark <sup>b</sup> |   | Cimanga et al. [78,79] <sup>a, b</sup>   |
|                   |   |  | Roots <sup>c</sup>     |   | Ablordeppey et al. [80] <sup>a</sup><br>Paulo et al. [81] <sup>c</sup><br>Hadden et al. [82] |

<sup>a,b</sup> and <sup>c</sup> correspond to the respective references.


comparable to chloroquine and quinine [52], the IC<sub>50</sub> values varying from 0.01 to 0.9 µg mL<sup>-1</sup> [53].

Frédérich et al. examined the roots and leaves of *Strychnos usambarensis* (Loganiaceae) growing in Rwanda [55,56]. Four potent anti-malarial bisindole alkaloids; 10'-hydroxyusambarensine (**16**), strychnopentamine (**17**), isostrychnopentamine (**18**) and chrysopentamine (**19**) have been isolated. Compounds **17** to **19** showed interesting anti-malarial activities against chloroquine-sensitive line (FCA 20) from Ghana (IC<sub>50</sub> values from 117 to 579 nM), against moderately chloroquine-resistant line (FCB1-R) from Colombia (IC<sub>50</sub> values from 107 to 550 nM) and against chloroquine-resistant line (W2) from Indochina (Laos) (IC<sub>50</sub> values from 145 to 507 nM). The results of anti-plasmodial activities were comparable to those of the anti-malarial drugs quinine and chloroquine, which is indicative of the absence of cross-resistance with chloroquine. Meanwhile, compound **16** had moderate *in vitro* activity against two strains of *P. falciparum*.


#### Naphthoisoquinolines

These compounds are characterized by the C5/C8' linkage between the naphthalene and the isoquinoline portions of these alkaloids (Figure 3). They have been isolated from *Ancistrocladus* (Acistrocladaceae), *Triphyophyllum*, *Dioncophyllum*, and *Habropetalum* (Dioncophyllaceae) species. The chemical significance of naphthylisoquinoline alkaloids rests on their unique structure and their biological activities [45,46].



The anti-malarial properties of some of these species have been investigated by Bringmann et al. [57-66]. Regarding the Acistrocladaceae-derived naphthoisoquinolines, compounds **20** to **24**, derived from the stems and leaves of *Ancistrocladus robertsoniorum* growing in Kenya, exhibited moderate anti-malarial activities (IC<sub>50</sub> values from 2.0 to 15.9 µM) against the K-1 and NF54 strains of *P. falciparum* [57], meanwhile the Tanzanian species, *Ancistrocladus tanzaniensis*, gave compounds **25** to **29** with IC<sub>50</sub> values ranging from 0.1 to 3.6 µg mL<sup>-1</sup> against the K1 strain and between 1.9 and 34.1 µg mL<sup>-1</sup> against the 3D7 strain [58]. Habropetaline A (**30**) and 5'-O-demethyl-dioncophylline A (**31**) were derived from the roots of *Triphyophyllum peltatum*, harvested in the Parc de Tai, in west Ivory Coast [59,60]. Both naphthoisoquinolines exhibited interesting anti-plasmodial activities against drug-sensitive and drug-resistant strains of the parasite. Habropetaline A (**30**) showed very good effect against *P. falciparum*, without cytotoxicity, with respective IC<sub>50</sub> values of 5.0 and 2.3 ng mL<sup>-1</sup> for the strains K1 (chloroquine and pyrimethamine resistant) and NF54 (sensitive to all known drugs). Compound **30** was almost as active as artemisinin (K1: 1.2 ng mL<sup>-1</sup>, NF54: 1.2 ng mL<sup>-1</sup>) and is known to be one of the most potent NPs used against *P. falciparum* [59]. On the other hand, 5'-O-demethyl-dioncophylline A (**31**) showed improved *in vitro* anti-malarial activity (IC<sub>50</sub> = 0.340 µg mL<sup>-1</sup>) against the erythrocytic forms of *P. falciparum* [60]. Jozi-peltine A (**32**), the dimer of the highly hydroxylated



**Figure 2** Anti-malarial indole alkaloids derived from the African flora.




**Figure 3** Naphthoisoquinolines with anti-plasmodial activity, derived from plants used in African traditional medicine.



naphthylisoquinoline alkaloid dioncopeltine A (36), was derived from a mixture of root and bark of *Triphyophyllum peltatum* and *Dioncophyllum thollonii*, along with twigs of *Habropetalum dawei* (Dioncophyllaceae), harvested from different regions on the continent [61]. Although this compound showed some *in vitro* anti-plasmodial activity against *P. falciparum* ( $K_1 = 875 \text{ ng mL}^{-1}$ ,  $\text{NF54} = 2530 \text{ ng mL}^{-1}$ ), it is significantly less active than its monomeric precursor, dioncopeltine A (36) ( $K_1 = 4.8 \text{ ng mL}^{-1}$ ,  $\text{NF54} = 3.3 \text{ ng mL}^{-1}$ ). This observation could lead to the conclusion that only naphthoisoquinolines containing one phenolic OH group each (such as dioncophylline A (36) and ancistrocladine (28)), could easily undergo the required dimerization reaction, implying that doubling of the number of free OH groups would increase the anti-plasmodial activity [61]. Dioncophyllines A (33), B (34) and C (35) and dioncopeltine A (36) were also active in the *in vivo* rodent model [66], with dioncophylline C (35) exhibiting a 50% effective dosage ( $\text{ED}_{50}$ ) of  $10.71 \text{ mg kg}^{-1} \text{ day}^{-1}$ . Four daily treatments with  $50 \text{ mg kg}^{-1} \text{ day}^{-1}$  were needed to achieve radical cure, one oral dose being sufficient to kill 99.6% of the parasites. Intravenous application of dioncophylline C was shown to be even more effective, with an  $\text{ED}_{50}$  of  $1.90 \text{ mg kg}^{-1} \text{ day}^{-1}$  and no noticeable toxic effects. Compound 35 also suppressed more established *Plasmodium berghei* infections when orally applied at day 3 after infection. It should be mentioned that rodent malaria is a well-known animal model for testing new

compounds and plant extracts. However, trial in human being is decisive to identify a “hit” as “a real hit”; and this is a good way to assess toxicity and safety. Both dioncopeltine A (36) and dioncophylline C (35) were active against the chloroquine-resistant *P. berghei* Anka CRS parasites. The naphthoisoquinolines are also known to exhibit other biological activities, e.g. dioncophylline A (33), is the main cytotoxin in *Ancistrocladus letestui* [67]. The above observations all point to the fact that naphthylisoquinoline alkaloids are promising lead compounds for the development of anti-malarial drugs.






### Furoquinolines

This subclass of alkaloids is easily identified with the *Vepris*, *Toddalia* and *Teclea* genera of the Rutaceae family. From the roots of *Vepris uguenensis*, Cheplogoi *et al.* isolated flindersiamine (37) and maculosidine (38) [68]. Although compound 37 lacked anti-malarial efficacy against all tested strains, maculosidine (38) exhibited moderate anti-malarial activity against two strains of *P. falciparum*, with  $IC_{50}$  values of 29.2 and 40.4  $\mu\text{g mL}^{-1}$  against the chloroquine-susceptible 3D7 and the chloroquine-resistant FCM29 strains respectively. Nitidine (39) has been derived from the roots of *Toddalia asiatica* harvested in Kenya and modified to yield the reduced derivative 5,6-dihydroneitidine (40) [69]. Even though nitidine is mostly known for its potential anticancer properties, the investigations of Gakunju *et al.* showed the alkaloidal extract of the roots of this plant to have high activity against the chloroquine-resistant K39 strain of *P. falciparum*, with an  $IC_{50}$  value of 0.04  $\mu\text{g mL}^{-1}$ . Further phytochemical analysis on the extract by these authors yielded nitidine as a major compound. *In vitro* screening against the K39 strain of *P. falciparum* revealed that nitidine exhibited high anti-plasmodial activity, with an  $IC_{50}$  of 0.045  $\mu\text{g mL}^{-1}$ , in addition to its known cytotoxic property. In order to remove toxicity, synthetic modification led to 5,6-dihydroneitidine (40), with a much weaker anti-malarial activity ( $IC_{50}$  of 1.03  $\mu\text{g mL}^{-1}$ , 23 times weaker than nitidine). Evoxine (41), derived from *Teclea gerrardii* (Rutaceae) harvested from Durban, South Africa, displayed moderate anti-plasmodial activity against the CQS D10 strain of *P. falciparum*, with  $IC_{50}$  value 24.5  $\mu\text{M}$  [70].

### Acridones

The most promising anti-plasmodial acridones derived from the African flora include arborinine (42), xanthoxoline (43), normelicopicine (44), skimmianine (45), dictamnine (46), melicopicine (47), tecleanthine (48) and 6-methoxytecleanthine (49), shown in Figure 5, isolated



from *Teclea* and *Zanthoxylum* species. Compound 42 was derived from *Teclea gerrardii*, *Zanthoxylum leprieurii* and *Teclea trichocarpa* (Rutaceae) and has shown anti-plasmodial activity against 3D7 strains ( $IC_{50} = 4.5 \mu\text{g mL}^{-1}$ ),

almost equally active as compound **43** ( $IC_{50} = 4.6 \mu\text{g mL}^{-1}$ ) [70,71]. Compounds **44** to **49** showed moderate activity against the chloroquine-sensitive HB3 and the chloroquine-resistant K-1 strains of *P. falciparum*, with respective anti-plasmodial  $IC_{50}$  values of 14.7, 9.3, 59.0, 53.0, and  $56.9 \mu\text{M}$  [72]. Compound **50** (maculosidine), derived from *Vepris uguenensis* (Rutaceae), exhibited anti-malarial activities at 13.8 and  $40.4 \mu\text{g mL}^{-1}$  against the 3D7 (chloroquine susceptible, CQS) and FCM29 (chloroquine resistant, CQR) strains of *P. falciparum*, respectively [73].

#### Amides

*N*-isobutyldeca-2,4-dienamide (**51**) and pipyahyne (**52**), Figure 6, are two amides respectively derived from *Hugonia castaneifolia* (Linaceae) and *Beilschmiedia zenkeri* (Lauraceae) [74,75]. It has been shown that compound **51** had moderate anti-plasmodial activity against the K-1 strain of *P. falciparum*, with an  $IC_{50}$  value of  $5.4 \mu\text{g mL}^{-1}$  [74], while compound **52** showed activity against the chloroquine-resistant W2 strain of *P. falciparum*, with an  $IC_{50}$  value of  $3.7 \mu\text{M}$  [75].

#### Cryptolepines

Cryptolepine (**53**), derived from *Sida acuta* (Malvaceae), growing in Ivory Coast, has shown very good anti-malarial activity [76]. According to Banzouzi *et al.*, the  $IC_{50}$  values obtained for the extracts from this plant ranged from 3.9 to  $5.4 \mu\text{g mL}^{-1}$ . Purification of this active fraction led to the identification of cryptolepine (**53**) as the active anti-plasmodial constituent of the plant. Compound **53** exhibited  $IC_{50}$  values against the chloroquine-sensitive strain (respectively 0.13 and  $0.17 \mu\text{g mL}^{-1}$  after 24 and 72 hours) from Nigeria and the Fcm29 chloroquine-resistant strain (respectively 0.17 and  $0.17 \mu\text{g mL}^{-1}$  after 24 and 72 hours) from Cameroon. Cryptolepine derivatives (**54** to **63**), Figure 7, isolated from the stems, roots and root bark of *Cryptolepis sanguinolenta* (Periplocaceae) growing in diverse regions in Africa, have also exhibited potent anti-malarial properties [77-80].

Cimanga *et al.* assessed three different extracts and four alkaloids from the root bark of *Cryptolepis sanguinolenta* *in vitro* against *P. falciparum* D-6 (chloroquine-sensitive strain), K-1, and W-2 (chloroquine-resistant strains). Cryptolepine (**53**) and its hydrochloride salt, 11-hydroxycryptolepine (**56**), and neocryptolepine (**61**) showed strong anti-plasmodial activity against *P. falciparum* chloroquine-resistant strains. Quindoline (**54**) was less active. The highest activity was obtained with cryptolepine (**53**). *In vivo* tests on infected mice showed that cryptolepine exhibited a significant chemosuppressive effect against *Plasmodium yoelii* and *Plasmodium berghei*, while cryptolepine (**53**) had the same effect against *P. yoelii* only. Compounds **54** and **56** did not show activity in this *in vivo* test system [79].

Another study by Paulo *et al.* on the roots of *Cryptolepis sanguinolenta* harvested from Guinea-Bissau led to the isolation of cryptolepinoic acid (**62**) and methyl cryptolepinoate (**63**) in addition to **53**, **54** and **56** from the ethanol and chlorophorm extracts of the leaves [81]. The isolated compounds and extracts were tested *in vitro* against *P. falciparum* K1 (multidrug-resistant strain) and T996 (chloroquine-sensitive clone). All extracts had 90% inhibition of *P. falciparum* K1 growth at concentrations  $<23 \mu\text{g mL}^{-1}$ . Cryptolepine (**53**) was the most active alkaloid tested with  $IC_{50}$  values ( $0.23 \mu\text{M}$  to K1;  $0.059 \mu\text{M}$  to T996), compared to chloroquine ( $0.26 \mu\text{M}$  to K1;  $0.019 \mu\text{M}$  to T996). The indolobenzazepine alkaloid cryptoleptine (**57**) was the second most active with  $IC_{50}$  values of  $0.8 \mu\text{M}$  (K1) and  $1.2 \mu\text{M}$  (T996). Cryptolepinoic acid (**62**) showed no significant activity while its ethyl ester derivative (**63'**) was active against *P. falciparum* K1 ( $IC_{50} = 3.7 \mu\text{M}$ ). All the indoloquinoline alkaloids showed cross-resistance with chloroquine but not the indolobenzazepine cryptoleptine (**57**). It was noticed that alkaloids with weakly basic characteristics were active whereas other structurally related alkaloids with different acid-base profiles were inactive. These observations are in agreement with the anti-malarial mechanism of action for quinolines. According to Hadden *et al.*, the unusual incorporation of the isopropyl group at the 11-position of the indolo [3,2-*b*] quinoline nucleus in 11-isopropylcryptolepine (**56**) is suggestive of a mixed biosynthetic origin for the alkaloid [82].

#### Terpenoids

Terpenoids with most promising anti-malarial properties are summarized in Tables 3, 4 and 5 (according to their subclasses), while the chemical structures are shown in Figures 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18 and 19.

#### Clerodane and labdane diterpenoids

Mambu *et al.* isolated the clerodanes (13*S*)-ent-7 $\beta$ -hydroxy-3-cleroden-15-oic acid (**64**), ent-7 $\beta$ -hydroxy-2-oxo-3-cleroden-15-oic acid (**65**), ent-2,7-dioxo-3-cleroden-15-oic acid (**66**) and ent-18-(*E*)-caffeoyloxy-7 $\beta$ -hydroxy-3-cleroden-15-oic acid (**67**) and the labdanes (13*S*)-ent-18-(*E*)-coumaroyloxy-8(17)-labden-15-oic acid (**68**), ent-18-(*E*)-caffeoyloxy-8(17)-labden-15-oic acid (**69**) and ent-15-*E*-caffeoyloxy-8(17)-labden-18-oic acid (**70**) from *Nuxia sphaerocephala*, growing in Ankazobe, 100 km from Antananarivo, Madagascar [83]. The compounds showed *in vitro* anti-plasmodial activities against the FcB1 *P. falciparum* strain, with respective  $IC_{50}$  values 14.6, 4.3, 8.0, 7.3, 11.4, 21.0,  $16.0 \mu\text{g mL}^{-1}$ . Duker-Eshun *et al.* obtained aframodial (**71**), (*E*)-8(17), 12-labddiene-15,16-dial (**72**), (*E*)-15,15-diethoxylabda-8(17),12-dien-16-al (**73**) and coronarin B (**74**) from the fruits and leaves of *Aframomum latifolium* or *Aframomum sceptrum* (Zingiberaceae) harvested in

**Table 3 Summary of anti-malarial terpenoids derived from the African flora, part 1: clerodanes, labdanes, limonoids, bisnorterpenes and acyclic triterpenes**

| Compound subclass                  | Isolated metabolites   | Plant species (Family)  | Part of plant studied                                | Place of harvest (locality, country) | Author, reference                        |
|------------------------------------|--|---|--|--------------------------------------|--|
| Clerodane and labdane diterpenoids | <b>64 to 70</b>  | <i>Nuxia sphaerocephala</i> (Loganiaceae)                                   | Leaves   | Ankazobe, Madagascar                 | Mambu et al. [83]                        |
|                                    | <b>71 to 74</b>  | <i>Aframomum latifolium</i> or <i>sceptrum</i> (Zingiberaceae)              | Fruits and Leaves                                    | Accra, Ghana                         | Duker-Eshun et al. [84]                  |
|                                    | <b>75, 76</b> and <b>77</b>  | <i>Turreanthus africanus</i> (Meliaceae)                                    | Seeds  | Mt. Cameroon, Cameroon               | Ngemenya et al. [85]                     |
|                                    | <b>78, 79, 80,</b> and <b>81</b>                                   | <i>Aframomum zambesiicum</i> (Zingiberaceae)                                | Seeds  | Nyasoso, Cameroon                    | Kenmogne et al. [86]                     |
|                                    | <b>74<sup>a, b</sup>, 82<sup>a</sup></b> and <b>83<sup>b</sup></b> | <i>Aframomum escapum<sup>a</sup></i> (Zingiberaceae)                        | Seeds <sup>a</sup><br>Fruits and leaves <sup>b</sup> | Mogbi, Cameroon <sup>a</sup>         | Ayimele et al. [87] <sup>a</sup>         |
|                                    |  | <i>Aframomum latifolium</i> and <i>sceptrum<sup>b</sup></i> (Zingiberaceae) |  | Accra, Ghana <sup>b</sup>            | Duker-Eshun et al. [84] <sup>b</sup>     |
| Limonoids                          | <b>84, 85, 86,</b> and <b>87</b>                                   | <i>Aframomum arundinaceum</i> (Zingiberaceae)                               | Seeds  | Maha, Cameroon                       | Wabo et al. [88]                         |
|                                    | <b>88</b>  | <i>Vepris uguenensis</i> (Rutaceae)   | Roots  | Baringo District, Kenya              | Cheplogoi et al. [73], Kiplimo [73]      |
|                                    | <b>89, 90, 91, 92,</b> and <b>93</b>                               | <i>Khaya grandifoliola</i> (Meliaceae)                                      | Bark and seeds                                       | Foumban, Cameroon                    | Bickii et al. [89]                       |
|                                    | <b>94</b> and <b>95</b>  | <i>Entandrophragma angolense</i> (Meliaceae)                                | Stem bark  | Awae forest reserve, Cameroon        | Bickii et al. [90]                       |
| Bisnorterpenes                     | <b>96, 97, 98,</b> and <b>99</b>                                   | <i>Ekebergia capensis</i> (Zingiberaceae)                                   | Stem bark  | Mt Kenya, Kenya                      | Murata et al. [91]                       |
| Bisnorterpenes                     | <b>100, 101, 102,</b> and <b>103</b>                               | <i>Salacia madagascariensis</i> (Celastraceae)                              | Roots  | Tanzania                             | Thiem et al. [92]                        |
| Acyclic triterpenes                | <b>104, 105, 106,</b> and <b>107</b>                               | <i>Ekebergia capensis</i> (Zingiberaceae)                                   | Stem bark  | Mt Kenya, Kenya                      | Murata et al. [91]                       |
|                                    | <b>108</b>   | <i>Aframomum escapum</i> (Zingiberaceae)                                    | Seeds  | Mogbi, Cameroon                      | Ayimele et al. [87]<br>Lopez et al. [93] |

<sup>a,b</sup> and <sup>c</sup> correspond to the respective references.

Accra, Ghana. Respective IC<sub>50</sub> values of 25, 48, 24, and 26 μM against the chloroquine-sensitive strain (3D7) were obtained for these compounds [84]. Anti-plasmodial activities were also obtained for labdanes 16-oxolabda-8(17),12(*E*)-dien-15-oic acid (**75**), methyl-14,15-epoxy-labda-8(17), 12(*E*)-diene-16-oate (**76**) and turraeanin A (**77**), from the seeds of *Turreanthus africanus* (Meliaceae), a plant generally used in preparations against fevers and malaria in ATM [85]. Compound **75** showed the highest anti-plasmodial activity (IC<sub>50</sub> of 26 μg mL<sup>-1</sup>) on chloroquine-sensitive *P. falciparum* F 32, *in vitro*, while compounds **76** and **77** rather had moderate activities [85]. Other anti-malarial labdanes of *Aframomum* sp. include 3-deoxyaulacocarpin A (**78**), zambesiacolactones A (**79**) and B (**80**) and aulacocarpin A (**81**) from seeds of *Aframomum zambesiicum* [86]; coranarin B (**82**) from the seeds of *Aframomum escapum* [87]; galanal A (**83**) from the leaves of *Aframomum sceptrum* [84]; galanal B

(**84**), galanolactone (**85**), (*E*)-8β,17-epoxylabd-12-ene-5,16 dial (**86**) and (*E*) labda-8,12-diene-15,16 dial (**87**) from the seeds of *Aframomum arundinaceum* [88]. Among these compounds, compound **76** (3-deoxyaulacocarpin A), derived from *Aframomum zambesiicum*, was both the least polar and the most active compound, with an IC<sub>50</sub> of 4.97 μM (1.73 μg/ml) [86].

#### Limonoids

Limonoids with good anti-plasmodial activities have been isolated from *Vepris uguenensis* (Rutaceae), harvested in Kenya [68,73], as well as from *Khaya grandifoliola* (Meliaceae) and *Entandrophragma angolense* (Meliaceae) harvested in Cameroon [89,90]. The chemical structures are shown in Figure 10. Methyl uguenesonate (**88**) from *Vepris uguenensis* displayed mild activity, with IC<sub>50</sub> values of 10.4 and 13.8 μg mL<sup>-1</sup>, against the CQS and CQR strains of *P. falciparum*, respectively [68,73]. The bark

**Table 4 Summary of anti-malarial terpenoids derived from the African flora, part 2: cassane furanoditerpenes, abietane diterpenes and sesquiterpenes**

| Compound subclass                         | Isolated metabolites                   | Plant species (Family)                                    | Part of plant studied     | Place of harvest (locality, country)      | Author, reference         |
|---|--|---|---------------------------|---|---------------------------|
| Cassane furanoditerpenes                  | 109 and 110                            | <i>Caesalpinia volkensii</i> (Leguminosae)                | Root bark                 | Gatamaiyo forest, Kenya                   | Ochieng et al. [94]       |
| Abietane diterpenes                       | 111 and 112                            | <i>Plectranthus hadiensis</i> (Lamiaceae)                 | Leaves                    | South Africa                              | van Zyla et al. [95]      |
|   | 113                                    | <i>Plectranthus lucidus</i> (Lamiaceae)                   | Leaves                    | South Africa                              | van Zyla et al. [95]      |
|   | 114                                    | <i>Plectranthus ecklonii</i> (Lamiaceae)                  | Leaves                    | South Africa                              | van Zyla et al. [95]      |
|   | 115, 116 and 117                       | <i>Plectranthus purpuratus</i> (Lamiaceae)                | Leaves                    | South Africa                              | van Zyla et al. [95]      |
|   | 118                                    | <i>Fuerstia africana</i> (Lamiaceae)                      | Aerial parts              | Ngong Hills, Kenya                        | Koch et al. [96]          |
|   | 119                                    | <i>Hoslundia opposita</i> (Lamiaceae)                     | Root bark                 | Tanzania                                  | Achenbach et al. [97]     |
|   | 120                                    | <i>Hyptis suaveolens</i> (Lamiaceae)                      | Leaves                    | Southeastern Nigeria                      | Chukwujekwu et al. [98]   |
| Sesquiterpenes and sesquiterpene lactones | 121, 122, 123, and 124                 | <i>Vernonia amygdalina</i> (Asteraceae)                   | Leaves                    | Mahale National Mountains Park, Tanzania  | Ohigashi et al. [99]      |
|   | 125                                    | <i>Vernonia brachycalyx</i> (Asteraceae)                  | Leaves                    | Machakos District, Kenya                  | Oketch-Rabah et al. [100] |
|   | 126                                    | <i>Ajuga remota</i> (Lamiaceae)                           | Aerial parts              | Nairobi, Kenya                            | Kuria et al. [101]        |
|   | 127, 128 and 129                       | <i>Reneilmia cincinnata</i> (Zingiberaceae)               | Fruits                    | Bafut, Cameroon                           | Tchuendem et al. [102]    |
|   | 130 and 131                            | <i>Acanthospermum hispidum</i> (Asteraceae)               | Flowers, leaves and stems | Danto/Porto-Novo, Benin                   | Ganfon et al. [103]       |
|   | 132, 133, 134, and 135                 | <i>Vernonia angulifolia</i> (Asteraceae)                  | Aerial parts              | University of KwaZulu-Natal, South Africa | Pedersen et al. [104]     |
|   | 136                                    | <i>Dicoma tomentosa</i> (Asteraceae)                      | Whole plant               | Poun, Burkina Faso                        | Jansen et al. [105]       |
|   | 137                                    | <i>Artemisia annua</i> (Asteraceae)                       | Seeds                     | Kjenzi (Bugarama), Burundi                | Reale et al. [106]        |
|   | 138                                    | <i>Dicoma anomala</i> subsp. <i>gerardii</i> (Asteraceae) | Root stocks               | Brits region, South Africa                | Becker et al. [107]       |
|   | 139                                    | <i>Tithonia diversifolia</i> (Asteraceae)                 | Aerial parts              | São Tomé and Príncipe islands             | Goffin et al. [108]       |
| 140                                       | <i>Scleria striatinux</i> (Cyperaceae) | Roots   | Oku, Cameroon             | Efange et al. [109]                       |                           |
| Coloratane sesquiterpenes                 | 141 to 148                             | <i>Warburgia ugandensis</i> (Canellaceae)                 | Stem bark                 | Dello Menna, Ethiopia                     | Wube et al. [110]         |

and seed extracts of *Khaya grandifoliola* (Meliaceae), a plant species widely used in the Central African subregion to treat various ailments, including malaria have also been investigated. Seven limonoids were isolated, among which five were significantly active (with  $IC_{50}$  values ranging between 1.25 and 9.63  $\mu\text{g mL}^{-1}$ ). These include: methylangolensate (89); gedunin (1.25  $\mu\text{g mL}^{-1}$ ) (90); 7-deacetylkhivorin (91); 1-deacetylkhivorin (92) and 6-acetylswietenolide (93). The same authors also investigated the stem bark of *Entandrophragma angolense* (Meliaceae) and isolated known limonoids with anti-malarial activity; 7 $\alpha$ -acetoxydihydronomilin (94), 7 $\alpha$ -obacunylacetate (95) and methylangolensate (89). Compounds 89 and 95 were considerably active against *P. falciparum* W2 with respective  $IC_{50}$  values of 2.0 and 19.5  $\mu\text{g mL}^{-1}$ .

Among the four limonoids derived from the stem bark of *Ekebergia capensis*, namely 7-deacetoxy-7-oxogedunin (96), ekeberins C1 (97), C2 (98) and C3 (99), only compound 96 exhibited significant activity against the chloroquine-susceptible FCR-3 strain of *P. falciparum*, with an  $IC_{50}$  of 6  $\mu\text{M}$ , but it lacked efficacy against the chloroquine-resistant K-1 strain [91].

#### Bisnorterpenes

Bisnorterpenes with interesting anti-plasmodial properties were purified from the roots of *Salacia madagascariensis* (Celastraceae), a shrub found in East Africa whose roots are used in the treatment of malaria, fever and menorrhagia specifically in Tanzania [92]. This plant is a rich source of bisnortriterpenes with potent antiprotozoal activity [45]. Four bisnortriterpenes; isoiguesterin

**Table 5 Summary of anti-malarial triterpenoids derived from the African flora, part 3: Beilshmiedic acid derivatives and pentacyclic triterpenes**

| Compound subclass             | Isolated metabolites                        | Plant species (Family)                           | Part of plant studied | Place of harvest (locality, country) | Author, reference      |
|-------------------------------|---|--|-----------------------|--------------------------------------|------------------------|
| Beilshmiedic acid derivatives | <b>149, 150, 151, 152, and 153</b>          | <i>Beilschmiedia cryptocaryoides</i> (Lauraceae) | Bark                  | Ranomafana-Ifanadiana, Madagascar    | Talontsi et al. [111]  |
| Pentacyclic triterpenes       | <b>154</b>                                  | <i>Schefflera umbellifera</i> (Araliaceae)       | Leaves                | South Africa                         | Mthembu [112]          |
|                               | <b>155</b>                                  | <i>Maytenus senegalensis</i> (Celastraceae)      | Root bark             | Eastern region of Sudan              | Khalid et al. [113]    |
|                               | <b>156, 157, 158, 159, and 160</b>          | <i>Nuxia sphaerocephala</i> (Loganiaceae)        | Leaves                | Ankazobe, Madagascar                 | Mambu et al. [83]      |
|                               | <b>161 and 162</b>                          | <i>Hymenocardia acida</i> (Phyllanthaceae)       | Bark                  | Chad                                 | Mahmout et al. [114]   |
|                               | <b>161</b>                                  | <i>Cassia siamea</i> (Fabaceae)                  | Stems                 | Otu (Oyo State), Nigeria             | Ajaiyeoba et al. [115] |
|                               | <b>163 and 164</b>                          | <i>Entandrophragma angolense</i> (Meliaceae)     | Stem bark             | Awae forest reserve, Cameroon        | Bickii et al. [90]     |
|                               | <b>165</b>                                  | <i>Hypericum lanceolatum</i> (Hypericaceae)      | Stem bark             | Mt. Bamboutos, Cameroon              | Zofou et al. [116]     |
|                               | <b>166</b>                                  | <i>Psorospermum glaberrimum</i> (Hypericaceae)   | Stem bark             | Ekombitié, Cameroon                  | Lenta et al. [117]     |
|                               | <b>167</b>                                  | <i>Baillonella toxisperma</i> (Sapotaceae)       | Stem bark             | Korup forest reserve, Cameroon       | Mbah et al. [118]      |
|                               | <b>168</b>                                  | <i>Kigelia africana</i> (Bignoniaceae)           | Stem bark             | Bandjoun, Cameroon                   | Zofou et al. [119]     |
| <b>169, 170 and 171</b>       | <i>Cogniauxia podolaena</i> (Cucurbitaceae) | Stem bark  | Congo                 | Banzouzi et al. [120]                |                        |

(**100**), 20-*epi*-isoiguesterinol (**101**), isoiguesterinol (**102**) and 6-oxoisoiguesterinol (**103**), were reported from the roots of this plant [92], Figure 11. However, only the first two showed high activity, with respective IC<sub>50</sub> values of 200 and 68 ng mL<sup>-1</sup> against the D6 strain of *P. falciparum*, and 170 and 68 ng mL<sup>-1</sup> (against the W2 strain of *P. falciparum*), respectively.


#### Acyclic triterpenes

The most active acyclic triterpenes have been found in the stem bark of *Ekebergia capensis* (Zingiberaceae) by Murata et al. [91]. Four triterpenes from the stem bark of this species, comprising two new acyclic triterpenoids, namely ekeberin D4 (**104**) and D5 (**105**) and two known ones (3*R*,2*R*)-2,3,22,23-tetrahydroxy-2,6,10,15,19,23-hexamethyl-6,10,14,18-tetracosatetraene (**106**) and (2*R*,3*R*,2*R*)-2-hydroxymethyl-2,3,22,23-tetrahydroxy-2,6,10,15,19,23-hexamethyl-6,10,14,18-tetracosatetraene (**107**) have been identified, Figure 12. Compounds **106** and **107** exhibited moderate anti-malarial activity against the FCR-3 strain of *P. falciparum*, with IC<sub>50</sub> values of 55 and 18 μM respectively, in addition to the good activities against the chloroquine-resistant K-1 strain, against which they had respective IC<sub>50</sub> values of 7 and 59 μM. The triterpene **104** lacked efficacy, while **105** had an IC<sub>50</sub> of 137 μM against the same parasite [91]. (+) *S*-nerolidol (**108**) isolated from the seeds of *Aframomum escapum* [87], is an important constituent of essential oils used in the treatment of

malaria. This compound is also found in *Artemisia herba alba* and in lemon grass, and is able to arrest development of the intraerythrocytic stages of the parasite. Compound **108** was identified as the active constituent leading to 100% growth inhibition at the schizont stage [93].

#### Cassane furanoditerpenes


Ochieng et al. isolated the cassane furanoditerpenes; deoxycasaldekarin C (**109**) and casaldekarin C (**110**) from the the chloroform and ethyl acetate extracts of the root bark of *Caesalpinia volkensii* from Kenya (Figure 13). These two compounds have exhibited antinociceptive and anti-plasmodial activities [94]. The anti-plasmodial activities were evaluated against chloroquine-sensitive (D6) and chloroquine-resistant (W2), with respective IC<sub>50</sub> values of 25.67 and 30.33 μg mL<sup>-1</sup> for compound **109** and respective IC<sub>50</sub> values of 34.44 and 30.69 μg mL<sup>-1</sup> for compound **110**. The results however demonstrated that *Caesalpinia volkensii* and other members of this genus contain cassane furanoditerpenes, which play a role in the medicinal properties of their plant root barks. The antinociceptive action in chemical models of nociception in mice suggests that the root bark extract and the active principles (furanoditerpenes) represent potential therapeutic options for the management of pain related ailments and not malaria [94].



### Abietane diterpenes

The anti-malarial properties of *Plectranthus* sp. (Lamiaceae), harvested in South Africa, have been determined by van Zyla *et al.* [95]. Seven abietane diterpenes (see Figure 14) were isolated from *Plectranthus hadiensis*, *Plectranthus lucidus*, *Plectranthus ecklonii*, *Plectranthus purpuratus* subsp. *purpuratus* and *Plectranthus purpuratus* subsp. *tongaensis*; 7 $\alpha$ -formyloxy-6,12-dihydroxy-abieta- 8,12-diene-11,14-dione (**111**), 7 $\alpha$ -acetoxy-6,12-dihydroxy-abieta-8,12-diene-11,14-dione (**112**), 11-hydroxy-2 $\alpha$ -(4-hydroxybenzoyloxy)-abieta-5,7,9(11),13-tetraene-12-one (**113**), 11-hydroxy-2 $\alpha$ -(3,4-dihydroxybenzoyloxy)abieta-5,7,9(11),13-tetraene-12-one (**114**), 11-hydroxy-19-(methyl-buten-

2-oyloxy)-abieta-5,7,9(11),13-tetraene-12-one (**115**), 11-hydroxy-19-(4-hydroxybenzoyloxy)-abieta-5,7,9(11),13-tetraene-12-one (**116**) and 11-hydroxy-19-(3,4-dihydroxybenzoyloxy)-abieta-5,7,9(11),13-tetraene-12-one (**117**). These compounds were tested for their antiplasmodial activity and for their ability to inhibit  $\beta$ -haematin formation. Overall, they showed good activity (IC<sub>50</sub> values ranging from 3.11 to 14.65  $\mu$ M), with compound **114** being 62% as effective as chloroquine in inhibiting  $\beta$ -haematin formation. Compounds **111**, **114** and **117** were more active than quinine. However, the cytotoxicity profile indicated a low degree of specificity towards the malaria parasite. When combined with quinine, three




compounds (**114**, **115** and **117**) interacted in an additive manner whereas compound **111** interacted synergistically [95].

Two other abietane diterpenes with anti-plasmodial activities; ferruginol (**118**) and 3-*O*-benzoylhosloppone (**119**), were respectively isolated by Koch *et al.* [96] and Achenbach *et al.* [97]. Compound **118** was isolated from the methanol extract of dried aerial parts (leaves and stems) of *Fuerstia africana* (Lamiaceae), a low-growing herb endemic to tropical East Africa [96], while compound **119** was isolated from the root bark of *Hoslundia opposita* (Lamiaceae), harvested in Tanzania [97]. The anti-malarial activity of compound **118**, determined using the D6 (chloroquine-sensitive derived from CDC Sierra Leone) clone of *P. falciparum*, showed a strong activity with an IC<sub>50</sub> of 1.95 μg mL<sup>-1</sup> compared to chloroquine IC<sub>50</sub> = 1.94 μg mL<sup>-1</sup> [96]. Meanwhile, 3-*O*-benzoylhosloppone (**119**) inhibited the growth of the multidrug resistant strain K1 of *P. falciparum* *in vitro* with an IC<sub>50</sub> value of 0.4 μg mL<sup>-1</sup> [97]. A bioactivity-guided fractionation of the petroleum ether extract of the leaves of *Hyptis suaveolens*, from Nigeria, led to the isolation of the abietane-type diterpenoid endoperoxide, 13α-*epi*-dioxiabiet-8(14)-en-18-ol (**120**), a molecule with high anti-plasmodial activity (IC<sub>50</sub> = 0.1 μg mL<sup>-1</sup>) [98].

#### Sesquiterpenes and sesquiterpene lactones

Sesquiterpenes derived from *Vernonia* sp. are known to have interesting anti-plasmodial activities. The compounds include vernodalin (**121**), vernodalol (**122**), vernolide (**123**), hydroxyvernolide (**124**), derived from the leaves of *Vernonia amygdalina* by Ohigashi *et al.* [99], in addition to 16,17-dihydrobrachycalixolide (**125**) isolated from the leaves of the sister species, *Vernonia brachycalyx*, as a major anti-plasmodial compound, by Oketch-Rabah *et al.*, Figures 15 and 16 [100]. These compounds exhibited moderate anti-plasmodial activity against the multidrug-resistant K-1 strain of *P. falciparum*, vernodalin (**121**) being the most active compound with an IC<sub>50</sub> value of 4 μg mL<sup>-1</sup>. Meanwhile, compounds **122**, **123** and **124** had IC<sub>50</sub> values of 4.2, 8.4 and 11.4 μg mL<sup>-1</sup>, respectively [99]. The measured activities of the compounds correlates with the uses of the plants in ATM (the leaves of *Vernonia amygdalina* are used in the treatment of various diseases, including malaria). Quantitative analysis showed that young leaves of this species have a higher concentration of compound **121** than the other derived compounds, suggesting that the anti-malarial efficacy of the leaf extracts of this species may be partly due to the high content of this NP. It has also been reported that dry leaves of *Vernonia brachycalyx* contain 0.2-0.4% of the sesquiterpene dilactone **125**. This compound exhibited moderate to high anti-plasmodial activity against the K39, 3D7, V1/S and Dd2 *P. falciparum*




strains, with IC<sub>50</sub> values of 4.2, 13.7, 3.0, and 16  $\mu\text{g mL}^{-1}$ , respectively [100]. In spite of the anti-plasmodial activity of this compound, it also had higher toxicity against human lymphocytes, indicating that the anti-plasmodial activity may have been due to the general toxicity the compound had on cells. Despite these observations, the leaves of this species are still used in the treatment of malaria and parasitic infections in East Africa [45]. Ajuagarin-1 (126) is another sesquiterpene, which has been reported from aerial parts of *Ajuga remota*, harvested in Kenya [101]. The compound has exhibited moderate anti-malarial properties against the chloroquine-sensitive FCA20/GHA strain of *P. falciparum*, with an IC<sub>50</sub> of 23  $\mu\text{M}$  [101].

The sesquiterpenoids oplodiol (127), 5*E*,10(14)-germacradien-1 $\beta$ ,4 $\beta$ -diol (128) and 1(10) *E*,5*E*-germacradien-4 $\alpha$ -ol (129), derived from *Reneilmia cincinnata*, with respective IC<sub>50</sub> values of 4.17, 1.63 and 1.54  $\mu\text{M}$ , were


used to validate the use of this plant in ATM to cure malaria and other fevers in Cameroon [102]. In addition, Ganfon *et al.* investigated the antiparasitic activities of two sesquiterpenic lactones isolated from *Acanthospermum hispidum* harvested in Benin Republic [103]. From their results, two known sesquiterpenic lactones were isolated: 15-acetoxy-8 $\beta$ -[(2-methylbutyryloxy)]-14-oxo-4,5-*cis*-acanthospermolide), 130 and 9 $\alpha$ -acetoxy-15-hydroxy-8 $\beta$ -[(2-methylbutyryloxy)]-14-oxo-4,5-*trans*-acanthospermolide), 131. Compounds 130 and 131 showed *in vitro* anti-plasmodial activity against the chloroquine-sensitive strain (3D7) with IC<sub>50</sub> values of 2.9 and 2.23  $\mu\text{M}$ , respectively. Only 131 showed a high selectivity index (SI: 18.4) on *Plasmodium* compared to cytotoxicity against human fibroblasts cell line (WI38). Furthermore, the crude acidic water extract and fractions containing one of the two isolated compounds displayed a weak *in vivo* anti-malarial activity against *P. berghei berghei* with a long half-life






causing a delayed effect. *In vivo* acute (2000 mg kg<sup>-1</sup>) and sub-acute (1000 mg kg<sup>-1</sup>) toxicity tests of the crude acidic water extract did not show toxicity. Moreover, the crude acidic water extract, fractions and pure isolated compounds from *Acanthospermum hispidum* showed promising *in vitro* anti-plasmodial activity. Despite the fact that this study did not show *in vivo* acute and subacute toxicities of the crude acidic water extract, its weak *in vivo* anti-malarial activity and the *in vitro* cytotoxicity of pure compounds and enriched extracts containing **130** and **131** indicate that the aerial parts of this plant should be used with caution for malaria treatments [103].

The combined use of bioassay-guided fractionation based on *in vitro* anti-plasmodial assay and dereplication based on HPLC-PDA-MS-SPE-NMR by Pederson *et al.* [104], led to isolation of (6*S*,7*R*,8*S*)-14-acetoxy-8-[2-hydroxymethylacrylat]-15-helianga-1(10),4,11(13)-trien-15-*al*-6,12-olid or vernangulide A (**132**) and (5*R*,6*R*,7*R*,8*S*,10*S*)-14-acetoxy-8-[2-hydroxymethylacrylat]-elema-1,3,11(13)-trien-15-*al*-6,12-olid or vernangulide B (**133**), along with vernodalol (**134**), vernodalin (**135**) and 11,13β-dihydroxyvernodalin (**136**) from the dichloromethane/methanol 1:1 and methanol extracts of the aerial parts







of *Distephanus angulifolius*. The isolated compounds showed  $IC_{50}$  values in the range 1.6 to 3.8  $\mu$ M and 2.1 to 4.9  $\mu$ M against chloroquine-sensitive D10 and chloroquine-resistant W2 *P. falciparum* strains, respectively. Janson *et al.* identified urospermal A-15-O-acetate (**136**) as the main active compound responsible for the anti-plasmodial activity of *Dicoma tomentosa* (Asteraceae) from Burkina Faso [105]. Based on their results, the  $IC_{50}$  of the

compound was  $<1 \mu\text{g mL}^{-1}$  against both 3D7 and W2 strains. Compound **136** was found to be the main cytotoxic compound ( $SI = 3.3$ ). A rapid quantification of the anti-malarial drug, artemisinin (**137**) in *Artemisia annua* plants cultivated for the first time in Burundi by Reale *et al.*, revealed the prospect of cultivating *Artemisia* and eventually using the active principle to offer the population of Burundi a fundamental resource in a country where malaria is endemic [106].

Standard phytochemical analysis techniques, including solvent-solvent extraction, thin-layer- and column chromatography, were used by Becker *et al.* to isolate a eudesmanolide-type sesquiterpene lactone, dehydrobrachylaenolide (**138**), as the main active constituent of *Dicoma anomala* subsp. *gerrardii* from the Brits region of North West Province of South Africa [107]. The compound demonstrated an *in vitro*  $IC_{50}$  of 1.865  $\mu$ M against a chloroquine-sensitive strain (D10) of *P. falciparum*. The biological activities of synthetic analogues of compound **138** showed that a methylene lactone group must be present in the eudesmanolide before any significant anti-malarial activity could be observed. This feature is absent





in the artemisinins and suggests that eudesmanolide-type sesquiterpene lactones have a different mode of action from artemisinins. This hypothesis was further confirmed by microarray gene ontology analysis [107]. The ether extract from aerial parts of *Tithonia diversifolia* collected in São Tomé and Príncipe demonstrated good anti-plasmodial activity ( $IC_{50}$  of  $0.75 \mu\text{g mL}^{-1}$  against the FCA strain) and fractionation of this extract yielded the sesquiterpene lactone tagitinin C (**139**) as an active compound against *P. falciparum* ( $IC_{50}$  of  $0.33 \mu\text{g mL}^{-1}$  against the FCA strain) [108].

Okundoperoxide (**140**), a new compound with a cyclic endoperoxide moiety, was isolated by Efangé *et al.* from *Scleria striatinux* (Cyperaceae), a spice commonly used in Cameroonian folk medicine to treat malaria and other fevers. This molecule exhibited significant anti-plasmodial activity, with  $IC_{50}$  values of 0.47, 0.48, 1.49, and  $1.30 \mu\text{g mL}^{-1}$ , on *P. falciparum* W2, D6, K1, and NF54, respectively. Moreover, the molecule showed no significant toxicity against mammalian cells [109].

#### Coloratane sesquiterpenes


Wube *et al.* demonstrated the antiprotozoal activity of *Warburgia ugandensis* (Canellaceae) from Ethiopia towards *Trypanosoma brucei rhodesiense* and *P. falciparum* *in vitro* and attributed the anti-plasmodial activity to the presence of drimane and coloratane sesquiterpenes. These include 4(13),7-coloratadiene-12,11-olide (**141**),  $11\alpha$ -hydroxymuzigadiolide (**142**), muzigadiol (**143**),  $6\alpha$ ,  $9\alpha$ -dihydroxy-4(13),7-coloratadiene-11,12-dial (**144**), cinnamolide (**145**), cinnamolide- $3\beta$ -acetate (**146**), mukaadial (**147**) and ugandensidial (**148**), Figure 17. The anti-plasmodial assays also revealed that the six coloratane and six drimane sesquiterpenes isolated from this extract exhibited significant antitrypanosomal activity with  $IC_{50}$  values ranging from 0.45 to  $>114 \mu\text{M}$ . Among the compounds tested against the malaria parasite *P. falciparum*  $11\alpha$ -hydroxymuzigadiolide (**142**) was most active with an  $IC_{50}$  value of  $6.40 \mu\text{M}$  [110].

#### Beilshmiedic acid derivatives


Beilshmiedic acid derivatives exhibiting antibacterial and anti-plasmodial activities were obtained from *Beilshmiedia cryptocaryoides* (Lauraceae) collected from Madagascar (Table 5). The work of Talontsi *et al.* [111] led to the isolation of four new beilshmiedic acid derivatives, cryptobeilic acids A – D (**149** to **152**), and tsangibeilin B (**153**), Figure 19. Compounds **149** to **153** exhibited anti-plasmodial activity against erythrocytic stages of chloroquine-resistant *P. falciparum* strain NF54 (with  $IC_{50}$  values ranging from 5.35 to  $17.70 \mu\text{M}$ ) and weak cytotoxicity against L6 cell lines (with  $IC_{50}$  values ranging from 20.4 to  $61.0 \mu\text{M}$ ), the most promising antiplasmodial activity being shown by compound **150**.

#### Pentacyclic triterpenes


The crude organic (methanol/dichloromethane (1:1)) extract of the leaves of *Schefflera umbellifera* (Araliaceae) exhibits promising anti-malarial activity. Bioassay-guided fractionation of this extract yielded the active compound, 3-hydroxy-20(29)-lupen-28-ol (**154**), Figure 19, which exhibited good anti-plasmodial activity ( $IC_{50}$  of  $3.2 \mu\text{g mL}^{-1}$ ), when tested against a chloroquine-susceptible malarial strain (D10). The reference compound (chloroquine) gave an  $IC_{50}$  of  $27.2 \text{ ng mL}^{-1}$  [112]. The quinonemethide




15-Acetoxy-8-[(2-methylbutyryloxy)]-14-oxo-4,5-cis-acanthospermolide (130)




9-Acetoxy-15-hydroxy-8-(2-methylbutyryloxy)-14-oxo-4,5-trans-acanthospermolide (131)




Vernangulide A (132)




Vernangulide B (133)




Vernodalol (134)




Vernodalin (135)




Urospermal A-15-O-acetate (136)




Artemisinin (137)



Dehydrobrachylaenolide (138)




Tagitinin C (139)




Okundoperoxide (140)


**Figure 16** Some anti-malarial sesquiterpenes and sesquiterpene lactones from African medicinal plants - II.




4(13),7-Coloratadiene-12,11-olide (141)




11 $\alpha$ -Hydroxymuzigadiolide (142)




Muzigadiol (143)




6 $\alpha$ ,9 $\alpha$ -Dihydroxy-4(13),7-coloratadiene-11,12-dial (144)




Cinnamolide (145)



Cinnamolide-3 $\beta$ -acetate (146)




Mukaadial (147)




Ugandensidial (148)

**Figure 17** Coloratane sesquiterpenes from African medicinal plants with anti-malarial activity.




triterpene, pristimerin or (20 $\alpha$ )-3-hydroxy-2-oxo-24-norfriedela-1(10),3,5,7-tetraen-carboxylic acid-(29)-methyl ester (**155**) was obtained by a bioactivity directed fractionation of the chloroform extract of the root bark of *Maytenus senegalensis* (Celastraceae) harvested from Sudan by Khalid *et al.* [113]. The *in vitro* anti-plasmodial activity of the isolated compound against chloroquine-resistant strain (Dd2) of *P. falciparum* was  $IC_{50} = 0.5 \mu\text{g mL}^{-1}$ , while the cytotoxicity on lymphocyte proliferation model was detected at  $IC_{50} = 6.8 \mu\text{g mL}^{-1}$ . The lupane-type triterpenoids 3-oxolupenal (3-oxolup-20(29)-en-30-al) (**156**), 3 $\beta$ -hydroxylupenal (3 $\beta$ -hydroxylup-20(29)-en-30-al) (**157**) and 3-oxolupenol (30-hydroxylup-20(29)-en-3-one) (**158**) were obtained from the leaves of *Nuxia sphaerocephala* (Loganiaceae), along with oleanolic acid (**160**), its acetylated ester (**159**), lupeol, uvaol, ursolic acid, and 3 $\beta$ -acetylursolic acid [83]. Among the compounds isolated from this study, **156** and **157** showed the best inhibitory activity against *P. falciparum* with the  $IC_{50}$  values between 1.55 and 4.67  $\mu\text{g mL}^{-1}$  *in vitro*, respectively.


Another lupane-type triterpene, lupeyl docosanoate (**162**), was isolated from the bark extract of *Hymenocardia acida* (Phyllanthaceae) collected in Chad, along with lupeol (**161**) and  $\beta$ -sitosterol by Mahmoud *et al.* [114]. The anti-malarial property of compound **162** justifies the ethnobotanic use of the plant in the treatment of malaria. *Cassia siamea* (Fabaceae) was identified from an ethnobotanical survey of southwest Nigeria as a remedy for febrile illness. Bioassay-guided fractionation of stem bark of the plant extract, using the parasite lactate dehydrogenase assay and multi-resistant strain of *P. falciparum* (K1) for assessing the *in vitro* anti-malarial activity led to the isolation of emodin and lupeol (**161**) from the ethyl acetate extract [115]. Both compounds were found to be the active principles responsible for the anti-plasmodial property with  $IC_{50}$  values of 5  $\mu\text{g mL}^{-1}$ , for each compound. The compounds 22-hydroxyhopan-3-one (**163**) and 24-methylene cycloartenol (**164**) from the stem bark of *Entandrophragma angolense* (Meliaceae) had moderate activities against *P. falciparum* W2 [90]. Zofou *et al.* evaluated the anti-plasmodial activity of betulinic acid (**165**) from the stem bark of the African St John's wort, *Hypericum lanceolatum* (Hypericaceae). The compound had an  $IC_{50}$  of 2.05  $\mu\text{g mL}^{-1}$  [116]. The *n*-hexane extract of *Psorospermum glaberrimum* from Cameroon showed good anti-plasmodial activity against the *P. falciparum* W2 strain, with  $IC_{50}$  of 0.87  $\mu\text{g mL}^{-1}$  [117]. Lenta *et al.* isolated betulinic acid (**165**) and friedelan-3-ol (**166**) from this extract. The measured *in vitro* activity of compound **165** against the *P. falciparum* W2 strain gave an  $IC_{50}$  of 2.33  $\mu\text{g mL}^{-1}$ . Mbah *et al.* isolated 3-*O*-betulinic acid *p*-coumarate (**167**) from *Baillonella toxisperma*, with an  $IC_{50}$  of 1.65  $\mu\text{M}$  [118]. The triterpenoid 2 $\beta$ ,3 $\beta$ ,19 $\alpha$ -trihydroxy-urs-12-20-en-28-




3-Hydroxy-20(29)-lupen-28-ol (154)




Pristimerin (155)




3-Oxolupenal (3-oxolup-20(29)-en-30-al) (156)




3β-Hydroxylupenal (3β-hydroxylup-20(29)-en-30-al) (157)




3-Oxolupenal (30-hydroxylup-20(29)-en-3-one) (158)




3β-Acetyloleanolic acid (159)




Oleanolic acid (160)




Lupeol (161): R = H




Lupeyl docosanoate (162): R =  $\text{C}_{22}\text{H}_{43}\text{O}_2$




22-Hydroxyhopan-3-one (163)




24-Methylene cycloartenol (164)




Betulinic acid (165)




3-Friedelanone (166)



3-O-betulinic acid *p*-coumarate (167)




2β,3β,19α-Trihydroxy-urs-12-20-en-28-oic acid (168)



Cucurbitacin B (169): R = H

Cucurbitacin D (170): R = H



20-Epibryonolic acid (171)

**Figure 19** Pentacyclic triterpenes from African medicinal plants with antiplasmodial activity.

oic acid (**168**) was isolated by Zofou *et al.* [119] from the stem bark of *Kigelia africana* (Bignoniaceae). This compound exhibited an  $IC_{50}$  of  $0.90 \mu\text{g mL}^{-1}$  against the W2 strain of *P. falciparum*.

*Cogniauxia podolaena* (Cucurbitaceae) is traditionally used in Congo Brazzaville for the treatment of malaria. The anti-plasmodial activity of the plant and some of the isolated compounds responsible for its activity were assessed by Banzouzi *et al.* [120]. Cucurbitacin B (**169**), cucurbitacin D (**170**) and 20-*epibryonolic* acid (**171**) were assayed for anti-plasmodial activity (on FcM29, a chloroquine-resistant strain of *P. falciparum*) and cytotoxicity (on KB and Vero cell lines). The compounds showed respective  $IC_{50}$  values of 1.6, 4.0 and  $2.0 \mu\text{g mL}^{-1}$  on FcM29. Compounds **169** and **170** both showed high cytotoxicity whereas **171** showed a better selectivity index.

## Conclusions

In this review an attempt has been made to document anti-malarial activities of NPs derived from African medicinal plants. It covers results published until the time of submission of the article. The first part of the review involves naturally occurring, anti-plasmodial/anti-malarial alkaloids and terpenoids while the second part of the review focuses on the remaining classes of compounds. Some of the compounds have been isolated from plants reputed to have a long history of usage in ATM, inferring that knowledge from ATM could be very useful in drug discovery efforts from African medicinal plants. From every indication, recent research efforts on new anti-malarial agents should focus on two main areas: the search for new chemical entities (NCEs) of natural/semi-synthetic origin, and the development of phytomedicines [37]. It should be mentioned that African researchers have, knowingly or unknowingly, blown the former avenue out of proportion. This is basically as a result of the fact that most of the research activities on medicinal plants going on in Africa are carried out by academic research groups and the focus is on publications, not application. This calls for the need to develop the necessary applications required to turn acquired knowledge on NPs derived from African medicinal plants into concrete applications in phytomedicine, within an industrial setting. It has been noticed that among the anti-malarials mentioned in this review, most have never been tested for cytotoxicity and very few have been tested for *in vivo* antoplasmodial activity. Another limitation is the, often small, quantities of compounds isolated from the plants which frustrate ambitions of large-scale screening efforts. Since some complex anti-malarial mixtures derived from plant extracts sometimes lose their anti-malarial properties when pure compounds are isolated, due to synergism of molecules in mixture, the trend towards the development of

total extracts into phytomedicines or improved traditional preparations is to be encouraged. Moreover, the isolation and characterization of NPs is an expensive endeavour, not within the reach of the average African research group. However, the attempt to validate ATM remedies as drugs will also face a number of limitations, among which are dosage determinations, variations of the concentration of the active ingredients in the plants with seasonal variations, the rapid loss of tropical forests and the extinction of key species, intellectual property rights management, the intervariability of plant species, quality control, and the conservation of biodiversity. The reconciliation between academic-oriented research and the development of phytomedicines could be feasible with the establishment of African centres of excellence in drug discovery [121], an initiative of the African Network for Drugs and Diagnostics Innovation (ANDI) [122], ATM being a major hub in this endeavour. In order to enhance modern drug discovery efforts from phytochemicals derived from the African flora, a recent effort by the authors of this paper has been to develop virtual libraries including NPs derived from African medicinal plants that have been reported in the literature, for computer-aided drug discovery (CADD). These include the CamMedNP database, containing three-dimensional structures of NPs derived from Cameroonian medicinal plants [123], the ConMedNP database, which covers ten countries in the Central African geographical region, converging the Congo Basin [124] and the AfroDb database, which is a select highly potent dataset, covering compounds with remarkable activities derived from plants across the entire continent [125]. Such databases could serve as starting points for virtual screening (VS) and CADD, leading to the identification of *in silico* hits, followed by validation by biological assays. These efforts have been in line with the prediction of DMPK profiles of the NPs, with a view to prioritizing hit selection during VS campaigns [125-127].

## Abbreviations

AfroDb: African medicinal plants active compound database; ATM: African traditional medicine; ADME/T: Absorption, distribution, metabolism, excretion, and toxicology; ANDI: African network for drugs and diagnostics innovation; CADD: Computer-aided drug design; CamMedNP: Cameroonian medicinal plant and natural products database; ConMedNP: Congo basin medicinal plant and natural products database; DMPK: Drug metabolism and pharmacokinetics; NP: Natural product; VS: Virtual screening; WHO: World Health Organization.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

FNK, LLL, JCN, and LMM conceived the idea. FNK, LLL and PAO participated in the data collection. FNK and PAO contributed in the data analysis, the discussion of results and the conception of the paper under the supervision of LMM, WS, LLL, and JCN. FNK and PAO wrote the first draft of the paper and all authors agreed on the final version before submission.

## Acknowledgements

Financial support is acknowledged from Lhasa Ltd, Leeds, UK through the Chemical and Bioactivity Information Centre (CBIC), University of Buea, Cameroon.

## Author details

<sup>1</sup>Department of Chemistry, Faculty of Science, University of Douala, PO Box 24157, Douala, Cameroon. <sup>2</sup>Chemical and Bioactivity Information Centre, Department of Chemistry, Faculty of Science, University of Buea, PO Box 63, Buea, Cameroon. <sup>3</sup>Department of Pharmaceutical Sciences, Martin-Luther University of Halle-Wittenberg, Wolfgang-Langenbeck Str. 4, Halle, Saale 06120, Germany.

Received: 24 October 2013 Accepted: 10 December 2013

Published: 13 December 2013

## References

- WHO: *World malaria report 2012*. Geneva: World Health Organization; 2012. Available from [http://who.int/malaria/publications/world\\_malaria\\_report\\_2012/wmr2012\\_no\\_profiles.pdf](http://who.int/malaria/publications/world_malaria_report_2012/wmr2012_no_profiles.pdf) (accessed on August 02, 2013).
- Vogel G: **Infectious disease - New map illustrates risk from the 'other' malaria.** *Science* 2010, **329**:618–618.
- Nogueira CR, Lopes LMX: **Antiplasmodial natural products.** *Molecules* 2011, **16**:2146–2190.
- White NJ: **Antimalarial drug resistance.** *J Clin Invest* 2004, **113**:1084–1092.
- Chin YW, Balunas MJ, Chai HB, Kinghorn AD: **Drug discovery from natural sources.** *AAPS J* 2006, **8**(2):E239–E253.
- Fabricant DS, Farnsworth NR: **The value of plants used in traditional medicine for drug discovery.** *Environ Health Perspect* 2001, **109**:69–75.
- Addae-Mensah I, Fakorede F, Holtel A, Nwaka S: **Traditional medicines as a mechanism for driving research innovation in Africa.** *Malar J* 2011, **10**(Suppl 1):S9.
- Ginsburg H, Deharo E: **A call for using natural compounds in the development of new antimalarial treatments – an introduction.** *Malar J* 2011, **10**(Suppl 1):S1.
- Guantai E, Chibale K: **How can natural products serve as a viable source of lead compounds for the development of new/novel anti-malarials?** *Malar J* 2011, **10**(Suppl 1):S2.
- Cruz LR, Spangenberg T, Lacerda MVG, Wells TNC: **Malaria in South America: a drug discovery perspective.** *Malar J* 2013, **12**:168.
- Wells TNC: **Natural products as starting points for future anti-malarial therapies: going back to our roots?** *Malar J* 2011, **10**(Suppl 1):S3.
- Anthony MP, Burrows JN, Duparc S, JMoehrl J, Wells TNC: **The global pipeline of new medicines for the control and elimination of malaria.** *Malar J* 2012, **11**:316.
- Hostettmann K, Marston A, Ndjoko K, Wolfender JL: **The potential of African plants as a source of drugs.** *Curr Org Chem* 2000, **4**:973–1010.
- Efange SMN: **Natural products: a continuing source of inspiration for the medicinal chemist.** In *Advances in phytomedicine*. Edited by Iwu MM, Wootton JC. Amsterdam: Elsevier Science; 2002:61–69.
- Tona L, Ngimbi NP, Tsakala M, Mesia K, Cimanga K, Apers S, De Bruyne T, Pieters L, Totte J, Vlietinck AJ: **Antimalarial activity of 20 crude extracts from nine African medicinal plants used in Kinshasa, Congo.** *J Ethnopharmacol* 1999, **68**:193–203.
- Tona L, Cimanga RK, Mesia K, Musuamba CT, De Bruyne Apers TS, Hernans N, Van Miert S, Pieters L, Totté J, Vlietinck AJ: **In vitro antiplasmodial activity of extracts and fractions from seven medicinal plants used in the Democratic Republic of Congo.** *J Ethnopharmacol* 2004, **93**:27–32.
- Dike IP, Obembe OO, Adebiji FE: **Ethnobotanical survey for potential anti-malarial plants in south-western Nigeria.** *J Ethnopharmacol* 2012, **144**:618–626.
- Gbadamosi IT, Moody JO, Lawal AM: **Phytochemical screening and proximate analysis of eight ethnobotanicals used as antimalaria remedies in Ibadan, Nigeria.** *J Appl Biosci* 2011, **44**:2967–2971.
- Adebayo JO, Krettli AU: **Potential antimalarials from Nigerian plants: a review.** *J Ethnopharmacol* 2011, **133**:289–302.
- Silva JR DA, Ramos A DS, Machado M, De Moura DF, Neto Z, Canto-Cavaleiro MM, Figueiredo P, Do Rosário VE, Amaral ACF, Lopes D: **A review of antimalarial plants used in traditional medicine in communities in Portuguese-speaking countries: Brazil, Mozambique, Cape Verde, Guinea-Bissau, São Tomé and Príncipe and Angola.** *Mem Inst Oswaldo Cruz* 2011, **106**(Suppl. 1):142–158.
- Ancolio C, Azas N, Mahiou V, Ollivier E, Di Giorgio C, Keita A, Timon-David P, Balansard G: **Antimalarial activity of extracts and alkaloids isolated from six plants used in traditional medicine in Mali and Sao Tome.** *Phytother Res* 2002, **16**:646–649.
- Rasoanaivo P, Ratsimamanga-Urverg S, Ramanitrahasimbola D, Rafatro H, Rakoto-Ratsimamanga A: **Criblage d'extraits de plantes de Madagascar pour recherche d'activité antipaludique et d'effet potentialisateur de la chloroquine.** *J Ethnopharmacol* 1999, **64**:117–126.
- Puri M, Masum H, Heys J, Singer PA: **Harnessing biodiversity: the Malagasy Institute of Applied Research (IMRA).** *BMC Int Health Hum Rights* 2010, **10**(Suppl 1):S9.
- Randrianarivojosia M, Rasidimanana VT, Rabarison H, Cheplogoi PK, Ratsimbason M, Mulholland DA, Maucière P: **Plants traditionally prescribed to treat tazo (malaria) in the eastern region of Madagascar.** *Malar J* 2003, **2**:25.
- Mbatchi SF, Mbatchi B, Banzouzi JT, Bansimba T, Nsonde Ntandou GF, Ouamba JM, Berry A, Benoit-Vical F: **In vitro antiplasmodial activity of 18 plants used in Congo Brazzaville traditional medicine.** *J Ethnopharmacol* 2006, **104**:168–174.
- Bero J, Ganfon H, Jonville MC, Frédéric M, Gbaguidi F, DeMol P, Moudachirou M, Quetin-Leclercq J: **In vitro antiplasmodial activity of plants used in Benin in traditional medicine to treat malaria.** *J Ethnopharmacol* 2009, **122**:439–444.
- Kumulungui BS, Ondo-Azi AS, Mintsu NA, Fumoux F, Traore A: **In vitro antiplasmodial activity of seven plants commonly used against malaria in Burkina Faso.** *J Med Plant Res* 2012, **6**:2284–2288.
- Pillay P, Maharaj VJ, Smith PJ: **Investigation South African plants as a source of antimalarial drugs.** *J Ethnopharmacol* 2008, **129**:438–454.
- Vonthron-Sénécheau C, Weniger B, Ouattara M, Bi FT, Alphonse K, Lobstein A, Brun R, Anton R: **In vitro antiplasmodial activity and cytotoxicity of ethnobotanically selected Ivorian plants.** *J Ethnopharmacol* 2003, **87**:221–225.
- Soh PN, Benoit-Vical F: **Are West African plants a source of future antimalarial drugs?** *J Ethnopharmacol* 2007, **114**:130–140.
- Maregesi S, Van Miert S, Pannecouque C, Haddad MHF, Hermans N, Wright CW, Vlietinck AJ, Apers S, Pieters L: **Screening of Tanzanian medicinal plants against *Plasmodium falciparum* and human immunodeficiency virus.** *Planta Med* 2010, **76**:195–201.
- Muthaura CN, Rukunga GM, Chhabra SC, Mungai GM, Njagi ENM: **Traditional antimalarial phytotherapy remedies used by the Kwale community of the Kenyan Coast.** *J Ethnopharmacol* 2007, **114**:377–386.
- Waako PJ, Katuura E, Smith P, Folb P: **East African medicinal plants as a source of lead compounds for the development of new antimalarial drugs.** *Afr J Ecol* 2007, **45**(Suppl. 1):102–106.
- Ayuko TA, Njau RN, Cornelius W, Leah N, Ndieye IO: **In vitro antiplasmodial activity and toxicity assessment of plant extracts used in traditional malaria therapy in the Lake Victoria Region.** *Mem Inst Oswaldo Cruz* 2009, **104**(5):689–694.
- Bhatnagara S, Dasa P: **Antimalarial activity in tropical plants: a review.** *J Herbs Spices Med Plants* 2007, **13**. 10.1300/J044v13n01\_09.
- Schwikkard S, Van Heerden FR: **Antimalarial activity of plant metabolites.** *Nat Prod Rep* 2002, **19**:675–692.
- Oliviera AB, Dolabela MF, Braga FC, Jacome RLRP, Varotti FP, Povoia MM: **Plant-derived antimalarial agents: new leads and efficient phytomedicines. Part I. Alkaloids.** *Anais da Academia Brasileira de Ciencias* 2009, **81**:715–740.
- Batista R, Junior AJS, Oliveira AB: **Plant-derived antimalarial agents: new leads and efficient phytomedicines. Part II. Non-alkaloidal natural products.** *Molecules* 2009, **14**:3037–3072.
- Bero J, Frédéric M, Quetin-Leclercq J: **Antimalarial compounds isolated from plants used in traditional medicine.** *J Pharm Pharmacol* 2009, **61**:1401–1433.
- Saxena S, Pant N, Jain DC, Bhakuni RS: **Antimalarial agents from plant sources.** *Curr Sci* 2003, **85**:1314–1329.
- Frédéric M, Tits M, Angenot L: **Potential antimalarial activity of indole alkaloids.** *Trans R Soc Trop Med Hyg* 2008, **102**:11–19.
- Kuete V, Efferth T: **Cameroonian medicinal plants: pharmacology and derived natural products.** *Front Pharmacol* 2010, **1**:123.
- Ntie-Kang F, Lifongo LL, Mbaze LM, Ekwelle N, Owono Owono LC, Megnassan E, Judson PN, Sippl W, Efange SMN: **Cameroonian medicinal plants: a bioactivity versus ethnobotanical survey and chemotaxonomic classification.** *BMC Complement Altern Med* 2013, **13**:147.
- Titanji VPK, Zofou D, Ngemenya MN: **The antimalarial potential of medicinal plants used for the treatment of malaria in Cameroonian folk medicine.** *Afr J Trad CAM* 2008, **5**:302–321.



45. Magadula JJ, Erasto P: **Bioactive natural products derived from the East African flora.** *Nat Prod Rep* 2009, **26**:1535–1554.
46. Zofou D, Ntie-Kang F, Sippl W, Efange SMN: **Bioactive natural products derived from the Central African flora against neglected tropical diseases and HIV.** *Nat Prod Rep* 2013, **30**:1098–1120.
47. Nkunya MH, Makangara JJ, Jonker SA: **Prenylindoles from Tanzanian *Monodora* and *Isolona* species.** *Nat Prod Res* 2004, **18**:253–258.
48. Cao MR, Tits M, Angenot LM, Frédéric M: **17-O-acetyl,10-hydroxycorynanthol, a selective antiplasmodial alkaloid isolated from *Strychnos usambarensis* leaves.** *Planta Med* 2011, **77**:2050–2053.
49. Bidla G, Titanji VP, Joko B, Ghazali GE, Bolad A, Berzins K: **Antiplasmodial activity of seven plants used in African folk medicine.** *Ind J Pharmacol* 2004, **36**:245–246.
50. Mbah JA, Tane P, Ngadjui BT, Connolly JD, Okunji CC, Iwu MM, Schuster BM: **Antiplasmodial agents from the leaves of *Glossocalyx brevipes*.** *Planta Med* 2004, **70**:437–440.
51. Odebiyi OO, Sofowora ES: **Antimicrobial alkaloids from a Nigerian chewing stick (*Fagara zanthoxyloides*).** *Planta Med* 1979, **36**:204–207.
52. Okunji CO, Iwu MM, Ito Y, Smith PL: **Preparative separation of indole alkaloids from the rind of *Picralima nitida* (Stapf) T. Durand & H. Durand by pH zone refining countercurrent chromatography.** *J Liquid Chromatogr Relat Technol* 2005, **28**:775–783.
53. Kassim OO, Loyevsky M, Elliott B, Geall A, Amonoo H, Gordeuk VR: **Effects of root extracts of *Fagara zanthoxyloides* on the *in vitro* growth and stage distribution of *Plasmodium falciparum*.** *Antimicrob Agents Chemother* 2005, **49**:264–268.
54. Ansa-Asamoah R, Kapadia GJ, Lloyd HA, Sokoloski EA: **Picratidine, a new indole alkaloid from *Picralima nitida* seeds.** *J Nat Prod* 1990, **53**:975–977.
55. Frédéric M, Tits Hayette MP, Brandt V, Penelle J, De Mol PG, Lablès G, Angenot L: **10'-hydroxyusambarensine, a new antimalarial bisindole alkaloid from the roots of *Strychnos usambarensis*.** *J Nat Prod* 1999, **62**:619–621.
56. Frédéric M, Cristino A, Choi YH, Verpoorte R, Tits M, Angenot L, Prost E, Nuzillard JM, Zèches-Hanrot M: **Chrysopentamine, an antiplasmodial anhydronium base from *Strychnos usambarensis* leaves.** *Planta Med* 2004, **70**:72–76.
57. Bringmann G, Teltschik F, Michael M, Busemann S, Rückert M, Haller R, Bär S, Robertson SA, Kaminsky R: **Ancistrobertsonines B, C, and D as well as 1,2-didehydroancistrobertsonine D from *Ancistrocladus robertsonianus*.** *Phytochemistry* 1999, **52**:321–332.
58. Bringmann G, Dreyer M, Faber JH, Dalsgaard PW, Jaroszewski JW, Ndangalasi H, Mbago F, Brun R, Christensen SB: **Ancistrotanzanine C and related 5,1'- and 7,3'-coupled naphthylisoquinoline alkaloids from *Ancistrocladus tanzaniensis*.** *J Nat Prod* 2004, **67**:743–748.
59. Bringmann G, Messer K, Schwöbel B, Brun R, Aké Assi L: **Habropetaline A, an antimalarial naphthylisoquinoline alkaloid from *Triphophyllum peltatum*.** *Phytochemistry* 2003, **62**:345–349.
60. Bringmann G, Saeb W, God R, Schäffer M, François G, Peters K, Peters EM, Proksch P, Hostettmann K, Aké Assi L: **5'-O-demethylioncophylline A, a new antimalarial alkaloid from *Triphophyllum peltatum*.** *Phytochemistry* 1998, **49**:1667–1673.
61. Bringmann G, Saeb W, Wohlfarth M, Messer K, Brun R: **Jozipeltine A, a novel, unnatural dimer of the highly hydroxylated naphthylisoquinoline alkaloid dioncopenine A.** *Phytochemistry* 2000, **56**:5871–5875.
62. Bringmann G, Rübenacker M, Jansen JR, Scheutzw D: **On the structure of the Dioncophyllaceae alkaloids dioncophylline A ("triphophylline") and "O-methyltriphophylline."** *Tetrahedron Lett* 1990, **31**:639–642.
63. Bringmann G, Rübenacker M, Geuder T, Aké Assi L: **Dioncophylline B, a naphthylisoquinoline alkaloid with a new coupling type from *Triphophyllum peltatum*.** *Phytochemistry* 1991, **30**:3845–3847.
64. Bringmann G, Rübenacker M, Vogt P, Busse H, Aké Assi L, Peters K, Von Schnering HG: **Dioncopenine A and dioncolactone A: alkaloids from *Triphophyllum peltatum*.** *Phytochemistry* 1991, **30**:1691–1696.
65. Bringmann G, Rübenacker M, Weirich R, Aké Assi L: **Dioncophylline C from the roots of *Triphophyllum peltatum*, the first 5,19-coupled Dioncophyllaceae alkaloid.** *Phytochemistry* 1992, **31**:4019–4024.
66. François G, Timperman G, Eling W, Aké Assi L, Holenz J, Bringmann G: **Naphthylisoquinoline alkaloids against malaria: evaluation of the curative potentials of dioncophylline C and dioncopenine A against *Plasmodium berghei* *in vivo*.** *Antimicrob Agents Chemother* 1997, **41**:2533–2539.
67. Hallock YF, Hughes CB, Cardellina JH II, Schäffer M, Gulden KP, Bringmann G, Boyd MR: **Dioncophylline A, the principal cytotoxin from *Ancistrocladus letestui*.** *Nat Prod Lett* 1995, **6**:315–320.
68. Cheplogoi PK, Mulholland DA, Coombes PH, Randrianarivelosia M: **An azole, an amide and a limonoid from *Vepris uguenensis* (Rutaceae).** *Phytochemistry* 2008, **69**:1384–1388.
69. Gakunju DMN, Mberu EK, Dossaji SF, Gray AI, Waigh RD, Waterman PG, Watkins WM: **Potent antimalarial activity of the alkaloid nitidine, isolated from a Kenyan herbal remedy.** *Antimicrob Agents Chemother* 1995, **39**:2606–2609.
70. Waffo AFK, Coombes PH, Crouch NR, Mulholland DA, El Amin SMM, Smith PJ: **Acridone and furoquinoline alkaloids from *Teclea gerrardii* (Rutaceae: Toddalioideae) of southern Africa.** *Phytochemistry* 2007, **68**:663–667.
71. Tchinda AT, Fuendjiep V, Sajjad A, Matchawe C, Wafu P, Khan S, Tane P, Choudhary MI: **Bioactive compounds from the fruits of *Zanthoxylum lepreurii*.** *Pharmacologyonline* 2009, **1**:406–415.
72. Muriithi MW, Abraham WR, Addae-Kyereme J, Scowen I, Croft SL, Gitu PM, Kendrick H, Njagi ENM, Wright CW: **Isolation and *in vitro* antiplasmodial activities of alkaloids from *Teclea trichocarpa*: *in vivo* antimalarial activity and X-ray crystal structure of normelicopine.** *J Nat Prod* 2002, **65**:956–959.
73. Kiplimo JJ: **The phytochemical and biological activity of secondary metabolites from Kenyan and *Vernonia* and *Vepris* species.** PhD thesis. University of Kwazulu-Natal, Department of Chemistry; 2012.
74. Baraza LD, Joseph CC, Munisi JJE, Nkunya MHH, Arnold N, Porzel A, Wessjohann L: **Antifungal rosane diterpenes and other constituents of *Hugonia castaneifolia*.** *Phytochemistry* 2008, **69**:200–205.
75. Lenta BN, Tantangmo F, Devkota KP, Wansi JD, Chouna JR, Soh RC R, Neumann B, Stammer HG, Tsamo E, Sewald N: **Bioactive constituents of the stem bark of *Beilschmiedia zenkeri*.** *J Nat Prod* 2009, **72**:2130–2134.
76. Banzouzi JT, Prado R, Menan H, Valentin A, Roumestan C, Mallié M, Pelissier Y, Blache Y: **Studies on medicinal plants of Ivory Coast: Investigation of *Sida acuta* for *in vitro* antiplasmodial activities and identification of an active constituent.** *Phytomedicine* 2004, **11**:338–341.
77. Barku VYA, Opoku-Boahen Y, Dzotsi EY: **Isolation and pharmacological activities of alkaloids from *Cryptolepis sanguinolenta* (Lindl) Schl.** *Int Res J Biochem Bioinform* 2012, **2**:58–61.
78. Cimanga K, De Bruyne T, Peters L, Claeys M, Vlietinck A: **New alkaloids from *Cryptolepis sanguinolenta*.** *Tetrahedron Lett* 1996, **37**:1703–1706.
79. Cimanga K, De Bruyne T, Peters L, Vlietinck AJ: ***In vitro* and *in vivo* antiplasmodial activity of cryptolepine and related alkaloids from *Cryptolepis sanguinolenta*.** *J Nat Prod* 1997, **60**:688–691.
80. Ablordepey SY, Hufford CD, Bourne RF, Dwuma-Badu D: **<sup>1</sup>H-NMR and <sup>13</sup>C-NMR assignments of Cryptolepine, a 3:4 benzo-6-carboline derivative isolated from *Cryptolepis sanguinolenta*.** *Planta Med* 1990, **56**:416.
81. Paulo A, Gomes ET, Steele J, Warhurst DC, Houghton PJ: **Antiplasmodial activity of *Cryptolepis sanguinolenta* alkaloids from leaves and roots.** *Planta Med* 2000, **66**:30–34.
82. Hadden CE, Sharaf MHM, Guido JE, Robins RH, Tackie AN, Phoebe CH Jr, Schiff PL Jr, Martin GE: **11-Isopropylcryptolepine: A novel alkaloid isolated from *Cryptolepis sanguinolenta* characterized using submicro NMR techniques.** *J Nat Prod* 1999, **62**:238–240.
83. Mambu L, Grellet P, Florent L, Joyeau R, Ramanitrahasimbola D, Rasoanaivo P, Frappier F: **Clerodane and labdane diterpenoids from *Nuxia sphaerocephala*.** *Phytochemistry* 2006, **67**:444–451.
84. Duker-Eshun G, Jaroszewski JW, Asomaning WA, Oppong-Boachie F, Olsen CE, Christensen SB: **Antiplasmodial activity of labdanes from *Aframomum latifolium* and *Aframomum sceptrum*.** *Planta Med* 2002, **68**:642–644.
85. Ngemanya MN, Akam TM, Yong JN, Tane P, Fanzo- Free SMN, Berzins K, Titanji VPK: **Antiplasmodial activities of some products from *Turreanthus africanus* (Meliaceae).** *Afr J Health Sci* 2006, **13**:1–2.
86. Kenmogne M, Prost E, Harakat D, Jacquier MJ, Frederich M, Sondengam LB, Zeches M, Waffo-Teguio P: **Five labdane diterpenoids from the seeds of *Aframomum zambesiaceum*.** *Phytochemistry* 2006, **67**:433–438.
87. Ayimele GA, Tane P, Connolly JD: **Aulacocarpin A and B, nerolidol and  $\beta$ -sitosterol glucoside from *Aframomum escapum*.** *Biochem Syst Ecol* 2004, **32**:1205–1207.
88. Wabo HK, Tane P, Connolly JD: **Diterpenoids and sesquiterpenoids from *Aframomum arundinaceum*.** *Biochem Syst Ecol* 2006, **34**:603–605.
89. Bickii J, Njifutue N, Foyere JA, Basco LK, Ringwald P: ***In vitro* antimalarial activity of limonoids from *Khaya grandifoliola* C.D.C. (Meliaceae).** *J Ethnopharmacol* 2000, **69**:27–33.

90. Bickii J, Tchouya GR, Tchouankeu JC, Tsamo E: **The antiplasmodial agents of the stem bark of *Entandrophragma angolense* (Meliaceae).** *Afr J Tradit Complement Altern Med* 2007, **4**:135–139.
91. Thiem DA, Sneden AT, Khan SI, Tekwani LB: **Bisnortriterpenes from *Salacia madagascariensis*.** *J Nat Prod* 2005, **68**:251–254.
92. Murata T, Miyase T, Muregi FW, Naashima-Ishibashi Y, Umehara K, Warashina T, Kanou S, Mkoji GM, Terada M, Ishih A: **Antiplasmodial triterpenoids from *Ekebergia capensis*.** *J Nat Prod* 2008, **71**:167–174.
93. Lopes NP, Kato MJ, Andrade EHA, Maia JGS, Yoshida M, Plancharat AR, Katzinet AM: **Antimalarial use of volatile oil from leaves of *Virola surinamensis* (Rol.) Warb. by Waiãpi Amazon Indians.** *J Ethnopharmacol* 1999, **67**:313–319.
94. Ochieng' CO, Owuor PO, Mang'uro LAO, Akala H, Ishola IO: **Antinociceptive and antiplasmodial activities of cassane furanoditerpenes from *Caesalpinia volkensii* H. root bark.** *Fytoterapia* 2012, **83**:74–80.
95. Van Zyla RL, Khan F, Edwards TJ, Drewesc SE: **Antiplasmodial activities of some abietane diterpenes from the leaves of five *Plectranthus* species.** *S Afr J Sci* 2008, **104**:62–65.
96. Koch TCEFA, Orjala J, Mutiso PC, Soejarto DD: **An antimalarial abietane diterpene from *Fuerstia africana*.** *Biochem Syst Ecol* 2006, **34**:270–272.
97. Achenbach H, Waibel R, Nkunya MHM, Weenen H: **Antimalarial compounds from *Hoslundia opposita*.** *Phytochemistry* 1992, **31**:3781–3784.
98. Chukwujekwu JC, Smith P, Coombes PH, Mulholland DA, Van Staden J: **Antiplasmodial diterpenoid from the leaves of *Hyptis suaveolens*.** *J Ethnopharmacol* 2005, **102**:295–297.
99. Ohigashi H, Hoffman MA, Izutsu D, Koshimizu K, Kawanakan M, Sugiyama H, Kirby GC, Warhurst DC, Allen D, Wright CW, Phillipson JD, Timon-David P, Delmas F, Elias R, Balansard G: **Toward the chemical ecology of medicinal plant use in chimpanzees: The case of *Vernonia amygdalina*, a plant used by wild chimpanzees possibly for parasite-related diseases.** *J Chem Ecol* 1994, **20**:541–553.
100. Oketch-Rabah HA, Lemmich E, Dossaji SF, Theander TG, Olsen CE, Cornett C, Kharazmi A, Christensen SB: **Two new antiprotozoal 5-methylcoumarins from *Vernonia brachycalyx*.** *J Nat Prod* 1997, **60**:458–461.
101. Kuria KAM, Chepkwony H, Govaerts C, Roets E, Busson R, De Witte P, Zupko I, Hoornaert G, Quiryren L, Maes L, Janssens L, Hoogmartens J, Laekeman G: **The antiplasmodial activity of isolates from *Ajuga remota*.** *J Nat Prod* 2002, **65**:789–793.
102. Tchuendem MHK, Mbah JA, Tsopmo A, Ayafor JF, Sterner O, Okunji CC, Iwu MM, Schuster BM: **Anti-plasmodial sesquiterpenoids from the African *Reneilimia cinninata*.** *Phytochemistry* 1999, **52**:1095–1099.
103. Ganfon H, Bero J, Tchinda AT, Gbaguidi F, Gbenou J, Moudachirou M, Frédéric M, Quetin-Leclercq J: **Antiparasitic activities of two sesquiterpene lactones isolated from *Acanthospermum hispidum* D.C.** *J Ethnopharmacol* 2012, **141**:411–417.
104. Pedersen MM, Chukwujekwu JC, Lategan CA, Van Staden J, Smith PJ, Staerk D: **Antimalarial sesquiterpene lactones from *Distephanus angulifolius*.** *Phytochemistry* 2009, **70**:601–607.
105. Jansen O, Tits M, Angenot L, Nicolas JP, De Mol P, Nikiema JB, Frédéric M: **Anti-plasmodial activity of *Dicoma tomentosa* (Asteraceae) and identification of urospermal A-15-O-acetate as the main active compound.** *Malar J* 2012, **11**:289.
106. Reale S, Pace L, Monti P, De Angelis F, Marcozzi G: **A rapid method for the quantification of artemisinin in *Artemisia annua* L. plants cultivated for the first time in Burundi.** *Nat Prod Res* 2008, **22**:360–364.
107. Becker JWV, Van der Merwe M, Van Brummelen AC, Pillay P, Crampton BG, Mmutlane EM, Parkinson C, Van Heerden FR, Crouch NR, Smith PJ, Mancama DT, Maharaj VJ: **In vitro anti-plasmodial activity of *Dicoma anomala* subsp. *gerrardii* (Asteraceae): identification of its main active constituent, structure-activity relationship studies and gene expression profiling.** *Malar J* 2011, **10**:295.
108. Goffin E, Da Cunha AP, Ziemons E, Tits M, Angenot L, Frederich M: **Quantification of tagitinin C in *Tithonia diversifolia* by reversed-phase highperformance liquid chromatography.** *Phytochem Anal* 2003, **14**:378–380.
109. Efange SMN, Brun R, Wittlin S, Connolly JD, Hoye TR, McAkam T, Makolo FL, Mbah JA, Nelson DP, Nyongbela KD, Wirumum CK: **Okundoperoxide, a bicyclic cyclofarnesylsesquiterpene endoperoxide from *Scleria striatinux* with antiplasmodial activity.** *J Nat Prod* 2009, **72**:280–283.
110. Wube AA, Bucar F, Gibbons S, Asres K, Rattray L, Croft SL: **Antiprotozoal activity of drimane and coloratane sesquiterpenes towards *Trypanosoma brucei rhodesiense* and *Plasmodium falciparum* in vitro.** *Phytother Res* 2010, **24**:1468–1472.
111. Talontsi FM, Lamshöft M, Bauer JO, Razakarivony AA, Andriamihaja B, Strohmann C, Spiteller M: **Antibacterial and antiplasmodial constituents of *Beilschmiedia cryptocaryoides*.** *J Nat Prod* 2013, **76**:97–102.
112. Mthembu XS: **A phytochemical study of *Schefflera umbellifera* and *Elephantorrhiza elephantina*.** *MSc thesis, School of Chemistry, Pietermaritzburg, South Africa: University of KwaZulu-Natal*; 2007.
113. Khalid SA, Friedrichsen GM, Christensen SB, Tahir AE, Sattic GM: **Isolation and characterization of pristimerin as the antiplasmodial and antileishmanial agent of *Maytenus senegalensis* (Lam.) Exell.** *Arkivoc* 2007, **ix**:129–134.
114. Mahmoud Y, Mianpeurem T, Dolmazon R, Bouchu D, Fenet B: **15ème colloque sur la Pharmacopée et la Médecine Traditionnelles Africaines.** Libreville: Conseil Africain et Malgache pour l'Enseignement Supérieur (CAMES); 2008:1.
115. Ajaiyeoba EO, Ashidi JS, Okpako LC, Houghton PJ, Wright CW: **Antiplasmodial compounds from *Cassia siamea* stem bark extract.** *Phytother Res* 2008, **22**:254–255.
116. Zofou D, Kowa TK, Wabo HK, Ngemenya MN, Tane P, Titanji VPK: ***Hypericum lanceolatum* (Hypericaceae) as a potential source of new anti-malarial agents: a bioassay-guided fractionation of the stem bark.** *Malar J* 2011, **10**:167.
117. Lenta BN, Devkota PK, Ngouela S, Boyom FF, Naz Q, Choudhary MI, Tsamo E, Rosenthal PJ, Sewald N: **Anti-plasmodial and cholinesterase inhibiting activities of some constituents of *Psorospermum glaberrimum*.** *Chem Pharm Bull* 2008, **56**:222–226.
118. Mbah JA, Ndikum G, Zofou D, Ngemenya MN, Efange SMN: **Antiplasmodial triterpenes from the stem bark of *Baillonella toxisperma*.** *ISESCO J Sci Tech* 2011, **7**:84–87.
119. Zofou D, Tene M, Tane P, Titanji VPK: **Antimalarial drug interactions of compounds isolated from *Kigelia africana* (Bignoniaceae) and their synergism with artemether, against the multidrug-resistant W2mf *Plasmodium falciparum* strain.** *Parasitol Res* 2012, **110**:539–544.
120. Banzouzi JT, Soh PN, Mbachi B, Cavé A, Ramos S, Retailleau P, Rakotonandrasana O, Berry A, Benoit-Vical F: ***Cogniauxia podolaena*: bioassay-guided fractionation of defoliated stems, isolation of active compounds, antiplasmodial activity and cytotoxicity.** *Planta Med* 2008, **74**:1453–1456.
121. Nwaka S, Ochem A, Besson D, Ramirez B, Fakorede F, Botros S, Inyang U, Mgone C, Aday-Mensah I, Konde V, Nyasse B, Okole B, Guantai A, Loots G, Atadja P, Ndumbe P, Sanou I, Olesen O, Ridley R, Ilunga T: **Analysis of pan-African Centres of excellence in health innovation highlights opportunities and challenges for local innovation and financing in the continent.** *BMC Int Health Hum Rights* 2012, **12**:11.
122. Nwaka S, Ilunga TB, Da Silva JS, Verde ER, Hackley D, Vré RD, Mboya-Okeyo T, Ridley RG: **Developing ANDI: A novel approach to health product R & D in Africa.** *PLoS Med* 2010, **7**:e1000293.
123. Ntie-Kang F, Mbah JA, Mbaze LM, Lifongo LL, Scharfe M, Ngo Hanna J, Cho-Ngwaf, Onguéné PA, Owono LCO, Megnassan E, Sippl W, Efange SMN: **CamMedNP: Building the Cameroonian 3D structural natural products database for virtual screening.** *BMC Complement Altern Med* 2013, **13**:88.
124. Ntie-Kang F, Onguéné PA, Scharfe M, Owono LCO, Megnassan E, Mbaze LM, Sippl W, Efange SMN: **ConMedNP: a natural product library from Central African medicinal plants for drug discovery.** *RSC Adv* 2014, **4**:409–419.
125. Ntie-Kang F, Zofou D, Babiaka SB, Meudom R, Scharfe M, Lifongo LL, Mbah JA, Mbaze LM, Sippl W, Efange SMN: **AfroDb: A select highly potent and diverse natural product library from African medicinal plants.** *PLOS One* 2013, **8**:e78085.
126. Ntie-Kang F, Mbah JA, Lifongo LL, Owono LCO, Megnassan E, Mbaze LM, Judson PN, Sippl W, Efange SMN: **Assessing the pharmacokinetic profile of the CamMedNP natural products database: an in silico approach.** *Org Med Chem Lett* 2013, **3**:10.
127. Ntie-Kang F, Lifongo LL, Mbah JA, Owono LCO, Megnassan E, Mbaze LM, Judson PN, Sippl W, Efange SMN: **In silico drug metabolism and pharmacokinetic profiles of natural products from medicinal plants in the Congo basin.** *In Silico Pharmacol* 2013, **1**:12.

doi:10.1186/1475-2875-12-449

**Cite this article as:** Amoa Onguéné et al: The potential of anti-malarial compounds derived from African medicinal plants. Part I: A pharmacological evaluation of alkaloids and terpenoids. *Malaria Journal* 2013 **12**:449.