



The Power of a Positive Human–Animal Relationship for Animal Welfare

Jean-Loup Rault^{1*}, Susanne Waiblinger¹, Xavier Boivin² and Paul Hemsworth³

¹ Institute of Animal Welfare Science, University of Veterinary Medicine, Vienna, Austria, ² Unité Mixte de Recherche sur les Herbivores, Université Clermont Auvergne, INRA, VetAgro Sup, UMR Herbivores, Saint-Genès-Champagnelle, France, ³ Animal Welfare Science Centre, Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Parkville, VIC, Australia

OPEN ACCESS

Edited by:

Luigi Baciadonna,
Queen Mary University of London,
United Kingdom

Reviewed by:

Helen Gray,
Newcastle University, United Kingdom
Monica Padilla De La Torre,
Norwegian University of Life
Sciences, Norway

*Correspondence:

Jean-Loup Rault
jean-loup.rault@vetmeduni.ac.at

Specialty section:

This article was submitted to
Animal Behavior and Welfare,
a section of the journal
Frontiers in Veterinary Science

Received: 03 August 2020

Accepted: 29 September 2020

Published: 09 November 2020

Citation:

Rault J-L, Waiblinger S, Boivin X and Hemsworth P (2020) The Power of a Positive Human–Animal Relationship for Animal Welfare. *Front. Vet. Sci.* 7:590867. doi: 10.3389/fvets.2020.590867

Domestic animals often seek and enjoy interacting with humans. Positive human–animal relationships can elicit positive emotions and other positive welfare outcomes. Nevertheless, our understanding of the underlying processes that govern the positive perception of humans by animals is incomplete. We cover the potential mechanisms involved in the development and maintenance of positive human–animal relationships from the perspective of the animal. This encompasses habituation, associative learning, and possibly attachment or bonding based on communication and social cognition. We review the indicators from the literature to assess a positive human–animal relationship. We operationally define this positive relationship as the animal showing voluntary approach and spatial proximity (seeking) and signs of anticipation, pleasure, relaxation, or other indicators of a rewarding experience from interacting with the human. For research, we recommend accounting for the baseline human–animal relationship in the animal's everyday life, and incorporating a control treatment rather than only comparing positive to negative interaction treatments. Furthermore, animal characteristics, such as previous experience, genetics, and individual predisposition, as well as contextual characteristics related to the social and physical environment, may modulate the perception of humans by animals. The human–animal relationship is also influenced by human characteristics, such as the person's familiarity to the animal, attitudes, skills, and knowledge. We highlight implications for current practices and suggest simple solutions, such as paying attention to the animal's behavioral response to humans and providing choice and control to the animal in terms of when and how to interact with humans. Practical applications to achieve a positive perception of humans could be better utilized, such as by incorporating training principles, while keeping in mind trust and safety of both partners. Overall, there is growing evidence in the scientific literature that a positive human–animal relationship can bring intrinsic rewards to the animals and thereby benefit animal welfare. Further research is needed on the underlying processes to establish an effective positive human–animal relationship, especially in regard to the type, frequency, and length of human interaction necessary. In particular, the importance of providing animals with a sense of agency over their interactions with humans remains poorly understood.

Keywords: agency, domestic, interaction, inter-species, perception, positive welfare, welfare assessment, well-being

INTRODUCTION

The human–animal relationship (HAR) is an important determinant of animal welfare (1–3). Numerous studies have demonstrated the detrimental effects of a negative HAR on animal and human welfare, that is, productivity, companionship, health (4). A negative HAR can impair animal welfare with negative consequences on the animal's productivity, health, and welfare, primarily through fear as an underlying mechanism (1, 5). In comparison, the benefits of a positive HAR for animal welfare are poorly understood and appreciated. Domestic animals often seek and enjoy interacting with humans, beyond depending on humans for food (6–9). Animals may perceive interacting with humans *per se* as rewarding (5, 10, 11).

This review compiles the recent knowledge of the welfare benefits for animals of interacting positively with humans and provides recommendations to assess and utilize a positive HAR. We focus on the HAR from the perspective of the non-human animal (hereby referred to as “animal”) unless stated otherwise. For the HAR from the human's perspective, we refer the readers to other reviews (5, 12, 13). We restrict the scope of this article to domesticated species, primarily farm and companion animals, because they have been (and still are) selected over thousands of years with a major influence on their response to humans (9, 14), and most domestic animals experience frequent interactions with humans. Notwithstanding, animals from other species are also able to develop positive relationships with humans, for instance, animals kept in zoos (1, 15, 16) or laboratories (17, 18), and therefore examples on these species are included where relevant.

MECHANISMS FOR THE FORMATION OF A POSITIVE HAR

Definitions

A positive HAR can be defined conceptually based on a positive perception by the animal of the human. Because perception is challenging to assess practically, a positive HAR can be defined operationally in that the animal shows voluntary approach and spatial proximity (seeking) and signs of anticipation, pleasure, relaxation, or other indicators of a rewarding experience arising from interacting with the human. Fear of humans prevents a positive perception of humans, but low or no fear is in itself not a sufficient condition. A positive HAR brings beneficial short-term [e.g., positive emotions (19)] and long-term [e.g., stress resilience (20)] welfare outcomes for the animal when or after interacting with the human (see section Implications for Practice).

Habituation

HARs are most often referred to in the context of fear of humans (21), although positive HARs have received increased attention recently (1–3, 22, 23). This questions whether a positive HAR can be understood, as for negative HAR, solely as a consequence of a reduction in the fear response to humans or an absence of fear. When a stimulus is unfamiliar, fear is usually the default response. Fear of humans can be reduced through habituation, defined as a reduction in response resulting from repeated exposure to a stimulus (24). Although it can reduce

fear of humans by leading to a neutral response, habituation is insufficient to reach a positive HAR. This non-associative learning process can occur by direct exposure, but also be facilitated or inhibited by social learning or transmission from the dam or other animals (25, 26).

Interactions vs. Relationship

The formation of a relationship is a progressive process, reinforced upon subsequent interactions. This highlights the difference between an interaction and a relationship based on a single vs. multiple events between two individuals, respectively (27), with a relationship developing on the basis that animals are able to memorize and predict future interactions with humans (28, 29). We focus on the relationship rather than interactions because the HAR is more relevant for welfare because of its long-lasting and integrative nature (i.e., comprising past interactions, present, and predicting future ones). Of course, there is a link between interactions and the resulting relationship. In particular, the formation of a positive HAR may be jeopardized by negative interactions, even when the occurrence of positive interactions far outweighs negative interactions (30). However, a strong or high-quality HAR may endure deviation from positive interactions or be more resilient to aversive events (31, 32). The time at which a relationship is formed remains difficult to determine, but it can be defined as the time at which the animal forms expectations of its interaction with humans.

Associative Learning

Associative learning can accelerate the formation of a relationship, by the animal associating humans with positive aspects either through classical conditioning (the human presence itself or its concurrent association with a positive event) or operant conditioning (interacting with the human leads to positive consequences). A positive HAR can be established by human contact that is inherently rewarding such as through stroking or brushing [dog (33), sheep (6, 7), cattle (11, 34, 35), pig (36, 37)] or play interactions [dog (38), cat (39)]. However, not all individuals react in the same manner to putative positive interactions. For example, previous interactions affect the way animals perceived human contact [pig (32, 40)], supporting a role for ontogeny. Furthermore, animals from different genetic origins can also perceive stroking by humans differently [dog (9), sheep (41)], supporting a role for phylogeny. The role of potential modulating factors such as individual differences (e.g., personality) and affective states should be investigated further. It should be noted that a positive HAR cannot simply be explained by food or other resources provided by humans, although food can facilitate the development of a positive HAR [sheep (7), cow (42), pig (8), cat (39)].

Bonding

In addition to associative learning processes, a number of phenomena have been proposed to explain the formation of a positive HAR, in particular aspects relevant to social bonding and related constructs. Familiarity with a human does not necessarily equate to a positive HAR, although it may be conducive to it given that repeated non-aversive exposure can facilitate positive

appraisal [“the mere exposure” effect (43)]. The attachment theory has been used in the context of the HAR (26), originating from the study of infant–parent relationships and defined as an affectional bond binding the individuals together in space and enduring over time (44). Without any obvious reinforcement, and because animals need to feel safe and have a basis from which they can explore their world, attachment can occur with familiar individuals such as the mother, peers, other conspecifics, and even individuals from other species such as humans (26). These animals calm quickly after a short period of social isolation when in the presence of a familiar human [dog (33, 45), cat (46), hand-reared lamb (26, 47), pig (48)]. The socialization process (49, 50) may also play a role in the context of the HAR, through learning how to behave toward others. Indeed, a successful relationship encompasses both the intent by the animal and the human to interact, as well as competent social skills relying on sociocognitive and communicative abilities (see other contributions in this Special Issue).

ASSESSMENT OF A POSITIVE HUMAN-ANIMAL RELATIONSHIP

The HAR can be observed either through observations of spontaneous interactions (i.e., without interference) or through stimulus-evoked situations and tests that investigate the HAR in a more systematic way.

Indicators

A number of biological changes can occur before, during, and/or after interactions with humans (Table 1). Most of these indicators are based on features of the interactions, indirectly reflecting the HAR. Some indicators can distinguish different qualities of the relationship (e.g., evaluate which animals have a better relationship than others), but it is generally difficult to set a threshold where a positive HAR starts, apart from some indicators that clearly reflect a positive HAR. The assessment of a positive HAR requires a holistic analysis, given that several indicators need to be considered together for a full understanding. Care is required in assessing a positive HAR because, for example, the motivation to interact with humans may at the time be conflicting with other motivations, and some indicators of a positive HAR are species-specific.

Behavioral Changes

We describe here in a typical chronological order the behavioral changes associated with a positive HAR and their specificity to a positive HAR.

The animal can show signs of anticipation before the interaction takes place in cases when the human interaction is predictable or environmental cues signal the arrival of a human. These could be considered “appetitive” signs, such as pacing, vocalizations, or increased behavioral transitions (83). For example, captive Bottlenose dolphins anticipated interactions with humans, through increased surface looking and spy hopping, and these anticipatory behaviors correlated with their subsequent level of engagement in the interaction (84). These anticipatory signals can nevertheless be ambiguous indicators, as

either indicative of positive (excitement) or negative (frustration, for instance, if the delay is too long) states depending on the situation (83).

The first reaction of an animal to the arrival of a human in its environment is an orientation response. The animal typically turns its attention toward the human, possibly using various senses other than vision. The orientation response indicates that the animal notices the presence of the human but is not in itself an indicator of the quality of the HAR because of its potential ambiguous underlying motivations reflecting either a positive (e.g., interest) or negative (e.g., vigilance) state. However, head, ear, and body posture or movement and accompanying behaviors may help to distinguish at least between a negative HAR and a neutral to positive HAR; for example, in cattle, head stays in normal position and ears not erected or even hanging loosely while looking toward the person and ongoing rumination.

Once the human enters the animal’s environment, the latency to approach, in the form of voluntary seeking behavior of the animal, is generally an indicator of a positive HAR and/or curiosity. Approach is context-specific (e.g., novelty of the situation and stimulus) and species-specific and therefore should be used with other indicators. However, a lack of approach does not preclude a positive HAR but may just indicate low motivation for (physical) interaction at this time (68); this especially accounts for situations in the home environment where many distractions or competing motivations may occur (e.g., feeding, resting). In many cases, spatial proximity is also a sensitive indicator of a positive HAR, for instance, reflected by the duration of time spent near a human (30, 51).

The number or duration of interactions initiated by the animal is often used as an indicator of the quality of the HAR. Although a quantifiable metric, it does not necessarily reflect the relationship because the animal may modulate the interaction with the human according to its needs; for instance, the animal may want to interact more if it is distressed, may not have interacted for some time, or conversely may not be interested at that time in interacting (68). In this regard, further research is required on refined indicators of interactions (85), such as by studying the complementarity, reciprocity, and synchrony of behavioral exchanges that have been shown to be important in the quality of parent–infant interactions (70). For instance, behavioral synchrony has been shown to be linked to affiliation in humans (86), and locomotor synchrony has been observed between dogs and their owners (69). Further, dogs with lower initial oxytocin levels received more stroking from their owner (87), demonstrating the dynamic interplay of the HAR.

The type of behaviors and body posture displayed during approach and contact with humans, reflecting the animal’s level of engagement in the interaction, can provide information regarding the perception and motivation of the animal. In particular, solicitation behaviors such as species-specific grooming solicitation postures and other types of physical solicitation for contact such as touching, nudging, scratching the human with the paw, or vocalizations are indicators of the animal’s motivation to engage and can be interpreted as clear signs of a positive perception of the human. Animals may also expose body areas where they wish to be stroked, for example,

TABLE 1 | List of indicators of a positive human–animal relationship.

Category	Indicator	Examples of measures	Direction of change	Specificity ¹	Key references
Movement/location	Approach ²	Latency to approach/touch human	↓	Conditional	(22)
		Direction of movement relative to human	↔	Conditional	(22)
Expressive behaviors	Spatial proximity	Time in proximity of human	↑	Conditional	(30, 51)
	Vocalizations ²	Yapping, purring, chirping, grunting, other low-frequency vocalizations	↑	Conditional	(46, 48)
	Ear posture ²	Relaxed ears	↑	Yes	(52–58)
	Tail posture	Tail low, or wagging	↑	Conditional	(59)
	Body posture	Relaxed, laying down near the human	↑	Yes	(48)
	Facial expression	Muscle movement, change in facial display, closed/half-closed eyes	?	?	(60, 61)
	Qualitative behavior assessment		Positively valenced factors	↑	Yes
Preference for human ²		Choice or motivation for specific human(s) over other stimuli, memory of humans	↑	Conditional	(28, 32, 36, 48, 63, 64)
Characteristics of the interaction	Initiating physical interaction ²	Exposing a body area	↑	Conditional	(35, 37, 52, 65)
		Solicitation behaviors like nudging, scratching, play bow	↑	Yes	(66)
	Interaction features	Frequency or duration of physical contact, eye gaze	↑	Conditional	(67)
	Reaction to human contact	Acceptance of stroking or touch	↑	Conditional	(68)
	Behavioral synchrony	Temporal synchrony of behavioral exchanges/movement	↑	?	(69)
	Behavioral matching	Complementarity, reciprocity of behavioral exchanges	↑	?	(70)
During or postinteraction effects	Relaxation ²	Duration or shorter latency to rest or sleep, rumination	↑	Yes	(55, 71–74)
	Exploration ²	Secure base exploration	↑	Conditional	(26, 50)
Postinteraction behavioral changes	Separation distress ²	Searching behavior, distress vocalizations, or contact calls	↑	Conditional	(47, 51, 75, 76)
Physiological indicators	Oxytocin ²	Hormone concentration or change	↑	Conditional	(77)
	Heart rate ²	Heart rate beat per min	↓	Conditional	(54, 78, 79)
	Parasympathetic activity ²	Heart rate variability: high-frequency band, RMSSD	↑	Conditional	(54, 78, 79)
Cognitive and neurobiological measures	Cognitive bias	Positive judgment of ambiguous cues	↑	Conditional	(40, 80)
	Neurobiology	EEG, fNIRS, neuroimaging (e.g., MRI, PET scan), postmortem measures	↑ or ↓	Conditional	(37, 81, 82)

These are general indicators, and their expression may be species-specific. This table aims to stimulate discussion and consideration of the complexity of positive HAR assessment and our current state of knowledge on its assessment.

¹“Conditional”: the presence of this sign could indicate a positive HAR, but may need to be interpreted in conjunction with other indicators or the context (e.g., conflicting motivations).

²The presence of this sign could indicate a positive HAR, but its absence does not necessarily rule out a positive HAR.

the ventral neck area in cattle (35, 52), the abdominal area in pigs (37), or the back rather than the head region in dogs (65). These types of behavioral responses, exposing often vulnerable body region, may be interpreted as involving a level of trust reflecting a positive HAR, although some behaviors such as lying with the belly exposed may also indicate submission in dogs, for instance, and therefore do not necessarily indicate a positive HAR. In most cases, these behaviors are similar to those shown during intraspecific sociopositive interactions, although there are some interspecies specific behaviors [e.g., dog vs. wolf (88)].

The response of the animal in the presence of the human is obviously a key indicator of a positive HAR. A lack of avoidance

response to humans is usually indicative of low fear of humans (22). Ear position changes or positions (forward vs. side or backward, or erected vs. hanging) have been used to interpret the valence of human contact [sheep (53, 54), cow (52, 55, 56), dog (57), horse (58)], and recent work investigated more subtle changes in facial expression [cat (60), parrot (61)]. Tail wagging in dogs is often cited as an indicator of enjoyment, but it may be a sign of arousal rather than specifically positive valence (59). In some species, some vocalizations are often associated with positive interactions, for instance, purring in cats (46). A rapid reduction in distress vocalizations and increased proximity seeking toward humans can also be interpreted as a positive

perception of human presence [goat (75, 76), hand-reared sheep (47, 51)]. Redirected or displacement behaviors [e.g., in dogs yawning, lip- or muzzle-licking, and looking away or toward the ceiling (89)] may be negative indicators, reflecting a reluctance to interact or conflicting motivations. Similarly, pigs that were used to stroking or scratching expressed more high-pitched vocalizations when the handler did not provide gentle tactile contact, which the authors interpreted as indicators of stress possibly resulting from frustration due to the fact that the previously-handled piglets expected positive human contact (90).

Finally, qualitative behavior assessment in which human observers rank the bodily expression of the animals using word descriptors (91) seems promising as a holistic approach for differentiating HAR (62).

Behavioral changes to assess a positive HAR may be species-, individual-, and context-specific (see section Implications for Practice). In particular, the way the human and the animal initiate the contact or interact appear as important modulators of these changes [pig (92), dog (65, 66, 78, 87)].

Physiological Changes

In addition to behavioral changes, studies have also shown a wide array of physiological changes linked to human–animal interactions (93).

Oxytocin, in particular, has attracted a lot of attention for its link to social processes. Positive interactions, in particular with familiar humans, generally raise oxytocin concentration [reviewed in (77)]. The relationship between positive HAR and changes in oxytocin concentrations is nevertheless complex and not fully understood [dog (66, 67, 79, 87); sheep (94); domestic species (77); dairy cattle, pig, goat (95)].

Cortisol concentration also changes following positive interactions, with the direction of change reflecting either excitement [dog (79)] or conversely relaxation [dog (66)], and change in cortisol concentration is time- and context-dependent.

Heart rate and heart rate variability measures provide dynamic information on activation of the autonomic response. In general, studies show a reduced heart rate and an increase in measures of parasympathetic activation (e.g., high frequency, or greater root mean square of successive differences) during or after interacting positively with a human [sheep (54); dog (78, 79)], partly dependent on the body region of grooming [horse (96); cow (52)] or the type of interactions (97).

Finally, the involvement of other physiological changes, especially neurotransmitters such as opioids and dopamine and immune parameters, such as immunoglobulin-A require further research. In fact, positive interactions induce an array of physiological and immune changes in both humans and animals (93), and -omics approaches [e.g., transcriptomics, proteomics, metabolomics (98)] could be useful to decipher the biological pathways modulated by positive HAR and its effects on health. For instance, rabbits that received regular positive human contact showed lower incidence of atherosclerosis (99). Gently handled chickens had a higher immune response and disease resistance (100), and mere regular visual contact with humans increased the antibody response to Newcastle disease vaccine and reduced

heterophil-to-lymphocyte ratio following capture and restraint later in life (101).

Cognitive and Neurobiological Effects

Few studies have focused on the cognitive and neurobiological changes induced by a positive HAR. Cognitive bias tasks have recently been popular as an indirect assessment of emotional states by studying affect-related cognitive changes (102). In rats, tickling by a human induces a more positive judgment of ambiguous cues, suggesting that it induces a positive emotional state (80). Similarly, piglets that experienced positive human contact judged ambiguous cues more positively (40). Conversely, dogs show a more negative judgment of ambiguous cues after being left alone (103). Whether a positive HAR leads to positive emotional states requires further research.

Other approaches have relied on the animal's memory of humans. Pigs can remember positive interactions with humans for at least 5 weeks (32). Horses that were trained using positive reinforcement training with positive human interactions remembered the human 6 months later and spent more time close to the familiar human (63). Sheep can be trained to discriminate sheep and human faces (104) and remember those faces for over 2 years (28), and sheep also recognize their familiar caretaker without any pretraining (64). These findings support that a positive HAR can be long-lasting.

Finally, neurobiological studies of positive HAR are still in the early stage with the use of, for instance, functional near-infrared spectroscopy [dog (81)], electroencephalography [pig (37)], or postmortem brain measures [sheep (82)]. Our understanding of the neuroscience of human–animal interactions could progress with new techniques such as neuroimaging [dog (105)], allowing non-invasive longitudinal neurobiological studies.

Postinteraction Changes

Most studies have focused on studying those biological changes when or around the time a human is present. There are also a number of changes that can occur following positive human–animal interactions, that is, at other times than when the animal and human interact. These can be indicative of positive (relaxation or “postconsummatory”) or negative (e.g., separation distress, searching behavior) effects. These effects that outlast the interaction *per se* are often overlooked as compared to the changes occurring during the interaction. Indices of relaxation include hanging ear posture [cattle (55)], lower heart rate [dog (71)], greater parasympathetic activity [various species (72)], elevated brain oxytocin concentration [pig (73)], and shorter latency to rest or better sleep quality [dog (74)].

There can also be indicators of attempts to restore contact, for example, after interruption of an interaction, as evidenced by signs of separation distress or searching behavior [dog (45); hand-reared sheep (47)]. Although these may be signs of distress and negative emotional states, searching behavior and separation distress when an interaction is disrupted are nevertheless signs of a positive HAR.

Further research is warranted on whether a positive HAR can induce baseline biological changes on an animal, for example, changes to its time-budget outside of the interactive sessions

with humans. For instance, gentle human interactions during milking or rearing can lead to fewer aggressive interactions between dairy cows once they return to the herd (106) and lower adrenocortical activity in calves (107). Similarly, flocks of gently handled chickens showed fewer agonistic interactions (108).

Epistemological Considerations for the Investigation of a Positive Human–Animal Relationship

Motivation and preference tests can be used to assess the HAR (22). They can provide insight into the animal's perception (109), by testing animals on what they find positively and negatively reinforcing; what they want or do not want and how much they value the stimulus. Nevertheless, preferences and motivation may vary with the time of day, environmental conditions, the animal's previous experience, and the current condition and familiarity with the options under study (110), requiring careful interpretation.

The most common tests used for HAR assessment have been the stationary/passive human test, approaching/active human test, and tests involving separation from a human [reviewed in (22)]. We cover below various aspects of the development and use of tests to specifically assess a positive HAR.

An animal voluntarily approaching and interacting (non-aggressively) with a human is a prime indicator of a positive HAR. This is nonetheless not sufficient to qualify as a specific positive HAR indicator because the animal may approach and interact because of curiosity or a motivation to explore. The motivation to explore may also be initially affected by how fear-provoking the situation is. Conversely, the lack of approach is not sufficiently conclusive to reject a positive HAR due to potentially conflicting motivations and a momentary lack of motivation for interaction.

Tests based on avoidance responses (e.g., distance of withdrawal by the animal from an approaching human) are often used to measure the fear dimension of the HAR. However, acceptance of approach and subsequent touch and stroking by a human are clear indicators of a positive HAR and can be more sensitive in differentiating the quality of HAR than approach behavior toward a stationary person [pig (36, 92), cattle (35, 68, 111)]. The sensitivity of the tests nevertheless depends on the species tested and contextual features (22), as well as phenomena such as generalization of the response toward unfamiliar humans.

Situations where there is a lack of control offered to the animal because the animal is restrained or limited to a constrained space or when contact is imposed on the animal without the possibility for the animal to avoid or withdraw may influence the validity of the HAR assessment. Nevertheless, the few studies to date comparing restrained and unrestrained animals showed relatively similar responses to humans (68, 112, 113). Standardized interactions by the humans, such as imposing contact on the animal or using highly standardized interactive features (e.g., predetermined interaction in terms of bout frequency or duration) are commonly used in research settings as they provide experimental control. However, free-choice interactions may replicate real-life situations more faithfully because control over the situation may be linked to the perception

of the situation, although this hypothesis remains to be tested. It may be important that the animal is provided with a sense of control or agency (114) by free-choice approach about when and how to interact (61, 78). This is similar to the case for second-person neuroscience (85) that emphasizes the need to look at situations of active social engagement and reciprocal behaviors, rather than passive observation or being subjected to a situation with a lack of agency. This argument is based on the fact that an interaction typically involves active participation from both agents.

The test should be conducted in an appropriate environment. Animals have been most often tested individually, which may not reflect their typical reaction when in their social group. Furthermore, testing environments have most often been barren, offering few choices other than interacting with the human. Hence, this questions the specificity and validity of the animal's response toward the human as an indicator of a positive HAR in cases where there is a lack of choice (110).

The experience of the animal with humans is obviously crucial to consider, as additional positive interaction treatments may fail to show additional effects if the HAR is already positive (107, 115). Hence, it is important to assess the “baseline” HAR in the animal's real-life environment (i.e., outside of the experimental treatments) and take into account the ratio of negative and positive human contact (106).

Many studies, to date, compared positive and negative human interaction treatments, but lacked a control treatment [e.g., (73, 74, 116)]. This control treatment usually consists of minimal human contact involved in routine care and management (117), or human present with no active interactions (52). It is crucial to demonstrate that the HAR is specifically positive, rather than neutral. If comparing only positive and negative interactions without a control treatment, a potential difference may be induced by negative treatment effects without being able to distinguish them from the positive treatment effects.

As mentioned earlier, more detailed analysis of the interaction could assist in assessing the quality of the HAR, for instance, based on the synchrony between partners (69), or the functional complementarity of the exchange and/or responsiveness using similar approaches to those used in humans (70).

IMPLICATIONS FOR PRACTICE

Developing and Cultivating a Positive Human–Animal Relationship: How?

The HAR is a dynamic and reciprocal process modulated by individual and contextual features. An understanding of its development and regulatory mechanisms provides practical opportunities to develop and maintain a positive HAR for animal caretakers.

Gentle handling is particularly effective [sheep (6), pig (32, 92), ostrich (118)], although passive human presence may be required initially to habituate the animal [pig (36)]. Note that some species may not need physical contact, and visual contact may be sufficient [e.g., poultry (119, 120)], although the need for and type of contact are strongly species-dependent. Positive

interactions involve several species-specific sensory channels: tactile, visual, auditory, and olfactory, and are often multimodal [dairy cow (121), sheep (82), pig (23)]. In many species, brief (from 15 s to a few minutes) opportunities to interact with humans over days or several weeks are sufficient to reduce the animal's fear of humans and encourage approach and interaction [dog (33), horse (122), cattle (123), pig (92, 124), poultry (120)], suggesting that a positive HAR from the perspective of the animal can develop rapidly. Studies examining tickling of rats on positive affective states have demonstrated the importance of the dosage and characteristics of this technique (125). Further research is needed to determine the minimal “dose” of human contact required to form a positive HAR in terms of type, frequency, and duration of interaction.

Incorporating training principles, primarily through the use of positive reinforcement, has been broadly and successfully used in practice for zoo and companion animals to improve handling by reducing the aversiveness of some procedures [dog (126), cat (127), horse (63), primates (128)]. Training is not yet commonly used in farm settings despite proof of its effectiveness in research settings [pig (129, 130), sheep (131), cattle]. Given that human contact *per se* can be perceived as inherently rewarding, it could be used as a reward during training (e.g., stroking, brushing, playing), although food rewards may facilitate this process.

It may be easier to develop a HAR with young animals [dog (132), pig (124)] because they may have had fewer negative experiences with humans, have greater learning ability (133) and higher levels of curiosity and exploration (26, 50) than adult animals. In particular, the development of a HAR may be most effective during sensitive periods for socialization such as during early life (49) or socially stressful periods such as after weaning (134). Social facilitation, building on the transmission of the HAR with the dam or other conspecifics, can also be effective [horse (135), sheep (26), pig (136)]. There is even evidence of transgenerational transmission of positive HAR, as human contact altered mother quails' egg physiological environment and led to less emotionally reactive offspring (137).

Familiarity and previous experience with humans can influence the HAR. Nevertheless, if the animal's experience with familiar humans is mainly positive, domestic animals can generalize their positive response toward unfamiliar humans [sheep (138–140), dog (141, 142), pig (8, 10, 36), horse (143), cattle (34)], although the animal may still prefer familiar over unfamiliar humans [sheep (116)]. Generalization of the HAR to unfamiliar humans depends not only on past experiences but also familiarity of the context such as the behavior or other characteristics of the human and the location [cattle (42, 144, 145), pig (32, 36, 90, 146)], which may affect the motivation to approach and remain near an unfamiliar human. As such, a positive HAR is not necessarily limited to a personalized one, that is, toward a specific human.

It is important to keep in mind the potential modulating effects on the HAR due to genetics and species differences [fox (147), dog (9), sheep (41)], individual differences [dog (78)], previous experience and age [pig (32, 36)], social context [cattle (148), sheep (26), pig (136)], and other context-specific aspects.

In addition, the attitudes, skills, and knowledge of humans influence their behavior toward animals and in turn the animal's perception of humans (1, 5). Although beyond the scope of this article, these human factors should be considered when thinking of the HAR. There is also increasing evidence that animals can recognize human facial expression of emotions [dog (149–151), horse (152), goat (153)] or human bodily expression [cat (154)] and prefer positive human emotional expressions.

The predictability of the interaction can strongly affect the animal's response to humans [pig (30), beef cattle (144), sheep (116), dog (155)], because as mentioned previously the relationship is based on the animal's expectation of its interaction with humans. The HAR concept implies the predictability of human–animal interactions. In addition, the provision of choice and control available to the animal in terms of when and how to interact appears to be important [dog (79), pig (92), cattle (11)].


A key aspect for the human is to pay attention to the animal's response to humans. A positive HAR can be assessed based on behavioral observations as highlighted previously (see section Assessment of a Positive Human–Animal Relationship), and as such it is feasible to cultivate a positive HAR in practice based on this knowledge and without the need for specific equipment.

The Benefits of a Positive HAR: Why?

The HAR can have important and long-lasting effects on the welfare of animals, and this relationship is often critical to the domestic animals' role, for example, animal productivity and ease of handling and management, as well as companionship and satisfaction for the human. Evidence is accumulating on the potential welfare benefits of a positive HAR (**Figure 1**).

There are benefits of a positive HAR on stress resilience. For example, offering positive interactions to shelter dogs can reduce their cortisol level (156) and combined with training increases adoptability (157). Walking and stroking shelter dogs for 15 min once a week for 6 weeks increased the time they spent visible from the front of the pen and tail wagging (158). Only three 10-min bouts of handling was sufficient for shelter dogs to show a preference for the handler (33). Five minutes of weekly brushing dairy heifers facilitated their acclimation to the milking routine (123). Five seconds of back scratching of sows for 1 week prior to farrowing reduced piglet mortality in sows (159), although this stroking treatment was confounded with music from a radio. These examples indicate that brief positive interactions with humans can benefit animal management and animal welfare.

A positive HAR can also buffer aversive procedures where humans are involved such as veterinary inspections or management interventions [sheep (131), pig (129), cow (160), ostrich (118)], presumably by removing human-related stress-eliciting components. In addition, humans can provide social support to animals during stressful times, especially for animals kept in suboptimal social environment [pig (161), sheep (47), chimpanzee (17)]. Stroking by the owner calms the behavioral and heart rate responses of dogs to subsequent separation (71). The effectiveness of providing social support can be modulated by the quality of the HAR [dog (66, 162)].



As an enrichment strategy, positive interactions with humans present several advantages as they usually occur daily and can be combined with routine checks, can be manipulated for their predictability to minimize habituation, and do not require additional resources (*e.g.*, material). For example, orangutans preferred to stay in the part of their zoo enclosure where they can be close to and observe visitors (163), suggesting that interactions with humans may be enriching for them.

There is limited direct evidence to date that a positive HAR stimulates positive affective states in domestic animals (19). Tickling of rats (80) or gentle contact of pigs (40) by humans induces more positive judgment of ambiguous cues, suggestive of a positive emotional state. Positive or negative human interactions influence the sleeping patterns of dogs (74), although in the absence of a control treatment it remains to be determined whether this was the result of the positive or negative interactions.

Developing a positive HAR provides benefits in the long term. The persistence of the effect of early positive human contacts [5–16 weeks, pig (32, 124); 6 months, dairy cattle (11); 6–8 months, beef cattle (134); 24 months (164); 8 months, goat (76); 25 months, (165); 6–8 months, horse (63)] makes it an intervention with potentially long-lasting effects. Nevertheless, there may also be risks or disadvantages of a positive HAR. For example, pigs that experienced positive human interactions can be difficult to handle in familiar locations because of low fear of humans (166); however, pigs that are fearful of both humans and the unfamiliar handling location take longer to move and balk more than pigs that have experienced positive human interactions (146), suggesting an interplay between the

HAR and the familiarity of the environment. It is also important to keep in mind trust and safety of both partners, because animals with low fear of humans can be dangerous, especially in case of inappropriate human behavior as it is often the case in dog bites of children (155) or during risky or potentially aversive procedures that involve close contact or handling [horse (25), dairy cow (121, 167)]. In order to be able to both manage the animals in a practicable manner and minimize the risks of aggression or injuries, a positive HAR may benefit from settings boundaries such as respecting a safe distance and avoiding potentially dangerous interactions. Social animals usually learn to distinguish acceptable from unacceptable social behaviors during their development as part of the socialization process and the development of their social skills, and this socialization process may also affect the animal's behavior toward humans.

Hence, a positive HAR can provide animals with positive welfare outcomes (20, 168), such as greater stress resilience, social support, environmental enrichment, possibly positive affective states, as well as benefits to their role for humans.

CONCLUSIONS

Positive experiences with humans lead to domestic animals seeking and interacting with humans. Consequently, a positive HAR can bring intrinsic rewards to the animal. It can be used to elicit positive emotions and other positive welfare outcomes. Nevertheless, our understanding of the underlying processes that govern the positive perception of humans by animals is incomplete and will benefit from further research, especially in regard to the type, frequency, and length of human interaction

necessary to establish an effective positive HAR. In particular, the importance of providing animals with a sense of agency and its effect on the HAR remains poorly understood. Further research is needed to identify how much changes in features of interaction reflect the quality of the relationship.

AUTHOR CONTRIBUTIONS

J-LR wrote the first draft of the manuscript and drafted **Table 1**. XB drafted **Figure 1**. SW, XB, and PH edited the manuscript. All authors reviewed and approved the final manuscript.

REFERENCES

- Hemsworth PH, Sherwen SL, Coleman GJ. Human contact. In: Appleby M, editor. *Animal Welfare*. Wallingford: CAB International (2018). p. 294–314.
- Waiblinger S. Agricultural animals. In: Hosey G, Melfi V, editors. *Anthrozoology: Human-Animal Interactions in Domesticated and Wild Animals*. Oxford: Oxford University Press (2019), p. 32–58.
- Boivin X. Animal experience of domestication. In: Butterworth A, editor. *Animal Welfare in a Changing World*. Wallingford: CABI (2018). p. 154–61.
- Pinillos RG, Appleby MC, Manteca X, Scott-Park F, Smith C, Velarde A. One welfare—a platform for improving human and animal welfare. *Vet Rec*. (2016) 179:412–13. doi: 10.1136/vr.i5470
- Hemsworth PH, Coleman GJ. *Human-Livestock Interactions: The Stockperson and The Productivity and Welfare of Intensively Farmed Animals*. Wallingford: CABI. (2011).
- Tallet C, Veissier I, Boivin X. Human contact and feeding as rewards for the lamb's affinity to their stockperson. *Appl Anim Behav Sci*. (2005) 94:59–73. doi: 10.1016/j.applanim.2005.02.007
- Tallet C, Veissier I, Boivin X. How does the method used to feed lambs modulate their affinity to their human caregiver? *Appl Anim Behav Sci*. (2009) 119:56–65. doi: 10.1016/j.applanim.2009.03.012
- Hemsworth PH, Verge J, Coleman GJ. Conditioned approach-avoidance responses to humans: the ability of pigs to associate feeding and aversive social experiences in the presence of humans with humans. *Appl Anim Behav Sci*. (1996) 50:71–82. doi: 10.1016/0168-1591(96)01065-9
- Lazzaroni M, Range F, Backes J, Portele K, Scheck K, Marshall-Pescini S. The effect of domestication and experience on the social interaction of dogs and wolves with a human companion. *Front Psychol*. (2020) 11:785. doi: 10.3389/fpsyg.2020.00785
- Hemsworth PH, Barnett JL, Campbell RG. A study of the relative aversiveness of a new daily injection procedure for pigs. *Appl Anim Behav Sci*. (1996) 49:389–401. doi: 10.1016/0168-1591(96)01060-X
- Bertenshaw CE, Rowlinson P. Exploring heifers' perception of 'positive' treatment through their motivation to pursue a retreated human. *Anim Welfare*. (2008) 17:313–9.
- Friedman E, Krause-Parello CA. Companion animals and human health: benefits, challenges, and the road ahead for human-animal interaction. *Rev Sci Tech*. (2018) 37:71–82. doi: 10.20506/rst.37.1.2741
- McCune S, McCardle P, Griffin JA, Esposito L, Hurley K, Bures R, et al. Editorial: human-animal interaction (hai) research: a decade of progress. *Front Vet Sci*. (2020) 7:44. doi: 10.3389/fvets.2020.00044
- Price EO. Behavioral development in animals undergoing domestication. *Appl Anim Behav Sci*. (1999) 65:245–71. doi: 10.1016/S0168-1591(99)00087-8
- Claxton AM. The potential of the human-animal relationship as an environmental enrichment for the welfare of zoo-housed animals. *Appl Anim Behav Sci*. (2011) 133:1–10. doi: 10.1016/j.applanim.2011.03.002
- Hosey G, Melfi V. Human-animal interactions, relationships and bonds: a review and analysis of the literature. *Int J Comp Psychol*. (2014) 27:117–42.
- Baker KC. Benefits of positive human interaction for socially-housed chimpanzees. *Anim Welfare*. (2004) 13:239–45.
- LaFollette MR, Riley MC, Cloutier S, Brady CM, O'Haire ME, Gaskill BN. Laboratory animal welfare meets human welfare: a cross-sectional study of professional quality of life, including compassion fatigue in laboratory animal personnel. *Front Vet Sci*. (2020) 7:114. doi: 10.3389/fvets.2020.00114
- Boissy A, Manteuffel G, Jensen MB, Moe RO, Spruijt B, Keeling LJ, et al. Assessment of positive emotions in animals to improve their welfare. *Physiol Behav*. (2007) 92:375–97. doi: 10.1016/j.physbeh.2007.02.003
- Rault J-L, Hintze S, Camerlink I, Yee JR. Positive welfare and the Like: distinct views and a proposed framework. *Front Vet Sci*. (2020) 7:370. doi: 10.3389/fvets.2020.00370
- Rushen J, Taylor AA, de Passillé AM. Domestic animals' fear of humans and its effect on their welfare. *Appl Anim Behav Sci*. (1999) 65:285–303. doi: 10.1016/S0168-1591(99)00089-1
- Waiblinger S, Boivin X, Pedersen V, Tosi M-V, Janczak AM, Visser EK, et al. Assessing the human-animal relationship in farmed species: a critical review. *Appl Anim Behav Sci*. (2006) 101:185–242. doi: 10.1016/j.applanim.2006.02.001
- Tallet C, Brajon S, Devillers N, Lensink J. Pig-human interactions: creating a positive perception of humans to ensure pig welfare. In: Spinka M, editor. *Advances in Pig Welfare*. Duxford: Woodhead Publishing (2018). p. 381–93.
- Rankin CH, Abrams T, Barry RJ, Bhatnagar S, Clayton DF, Colombo J, et al. Habituation revisited: an updated and revised description of the behavioral characteristics of habituation. *Neurobiol Learning Mem*. (2009) 92:135–38. doi: 10.1016/j.nlm.2008.09.012
- Hausberger M, Roche H, Henry S, Visser EK. A review of the human-horse relationship. *Appl Anim Behav Sci*. (2008) 109:1–24. doi: 10.1016/j.applanim.2007.04.015
- Nowak R, Boivin X. Filial attachment in sheep: similarities and differences between ewe-lamb and human-lamb relationships. *Appl Anim Behav Sci*. (2015) 164:12–28. doi: 10.1016/j.applanim.2014.09.013
- Hinde RA. Interactions, relationships and social structure. *Man*. (1976) 11:1–17. doi: 10.2307/2800384
- Kendrick KM, da Costa AP, Leigh AE, Hinton MR, Peirce JW. Sheep don't forget a face. *Nature*. (2001) 414:165–6. doi: 10.1038/35102669
- Boivin X, Lensink J, Tallet C, Veissier I. Stockmanship and farm animal welfare. *Anim Welfare*. (2003) 12:479–92.
- Hemsworth PH, Barnett JL, Hansen C. The influence of inconsistent handling by humans on the behaviour, growth and corticosteroids of young pigs. *Appl Anim Behav Sci*. (1987) 17:245–52. doi: 10.1016/0168-1591(87)90149-3
- Schmid C, Boivin X, Scala S, Waiblinger S. Effect of previous stroking on reactions to a veterinary procedure: behaviour and heart rate of dairy cows. *Interact Stud*. (2010) 11:467–81. doi: 10.1075/is.11.3.08sch
- Brajon S, Laforest J-P, Bergeron R, Tallet C, Hötzel M-J, Devillers N. Persistency of the piglet's reactivity to the handler following a previous positive or negative experience. *Appl Anim Behav Sci*. (2015) 162:9–19. doi: 10.1016/j.applanim.2014.11.009
- Gácsi M, Topál J, Miklósi Á, Dóka A, Csányi V. Attachment behavior of adult dogs (*Canis familiaris*) living at rescue centers: forming new bonds. *J Comparat Psychol*. (2001) 115:423–31. doi: 10.1037/0735-7036.115.4.423
- Lensink BJ, Boivin X, Pradel P, Le Neindre P, Veissier I. Reducing veal calves' reactivity to people by providing additional human contact. *J Anim Sci*. (2000) 78:1213–8. doi: 10.2527/2000.7851213x
- Schmid C, Boivin X, Waiblinger S. Stroking different body regions of dairy cows: effects on avoidance and approach behavior toward humans. *J Dairy Sci*. (2008) 91:596–05. doi: 10.3168/jds.2007-0360
- Brajon S, Laforest J-P, Bergeron R, Tallet C, Devillers N. The perception of humans by piglets: recognition of familiar handlers and generalisation to unfamiliar humans. *Anim Cogn*. (2015) 18:1299–316. doi: 10.1007/s10071-015-0900-2
- Rault J-L, Truong S, Hemsworth L, Le Chevoir M, Bauquier S, Lai A. Gentle abdominal stroking (belly rubbing) of pigs by a human reduces EEG total power and increases EEG frequencies. *Behav Brain Res*. (2019) 374:111892. doi: 10.1016/j.bbr.2019.04.006

38. Jerolmack C. Humans, animals, and play: theorizing interaction when intersubjectivity is problematic. *Sociol Theor.* (2009) 27:371–89. doi: 10.1111/j.1467-9558.2009.01353.x
39. Karsh EB, Turner DC. The human-cat relationship. In: Turner DC, Bateson P, editors. *The Domestic Cat, The Biology of its Behaviour*. Cambridge: Cambridge University Press (1988). p. 159–77.
40. Brajon S, Laforest J-P, Schmitt O, Devillers N. The way humans behave modulates the emotional state of piglets. *PLoS ONE.* (2015) 10:e0133408. doi: 10.1371/journal.pone.0133408
41. Tamioso PR, Boissy A, Boivin X, Chandèze H, Andanson S, Delval É, et al. Spatial distance and reactivity traits alter the positive perception of brushing by ewes. *Animal.* (2019) 14:150–60. doi: 10.1017/S1751731119001435
42. Jago JG, Krohn CC, Matthews LR. The influence of feeding and handling on the development of the human-animal interactions in young cattle. *Appl Anim Behav Sci.* (1999) 62:137–51. doi: 10.1016/S0168-1591(98)00219-6
43. Montoya RM, Horton RS, Vevea JL, Citkovic M, Lauber EA. A re-examination of the mere exposure effect: the influence of repeated exposure on recognition, familiarity, and liking. *Psychol Bull.* (2017) 143:459–98. doi: 10.1037/bul0000085
44. Bowlby J. *Attachment and Loss*, Vol. 1. New York, NY: Basic Books (1969).
45. Prato-Previde E, Cusance DM, Spiezio C, Sabatini F. Is the dog-human relationship an attachment bond? An observational study using Ainsworth's strange situation. *Behaviour.* (2003) 140:225–54. doi: 10.1163/156853903321671514
46. Eriksson M, Keeling LJ, Rehn T. Cats and owners interact more with each other after a longer duration of separation. *PLoS ONE.* (2017) 12:e0185599. doi: 10.1371/journal.pone.0185599
47. Rault J-L, Boissy A, Boivin X. Separation distress in artificially-reared lambs depends on human presence and the number of conspecifics. *Appl Anim Behav Sci.* (2011) 132:42–50. doi: 10.1016/j.applanim.2011.02.011
48. Villain A, Lanthony M, Guérin C, Nous C, Tallet C. Manipulable object and human contact: preferences and modulation of emotional states in weaned piglets. *Front Vet Sci.* (this issue). doi: 10.3389/fvets.2020.577433
49. Bateson P. How do sensitive periods arise and what are they for? *Anim Behav.* (1979) 27:470–86. doi: 10.1016/0003-3472(79)90184-2
50. Scott JP. The phenomenon of attachment in human-nonhuman relationships. In: Davis H, Balfour D, editors. *The Inevitable Bond: Examining Scientist-Animal Interactions*. Cambridge: Cambridge University Press (1992).
51. Boivin X, Tournadre H, Le Neindre P. Hand-feeding and gentling influence early-weaned lambs' attachment responses to their stockperson. *J Anim Sci.* (2000) 78:879–84. doi: 10.2527/2000.784879x
52. Schmied C, Waiblinger S, Scharl T, Leisch F, Boivin X. Stroking of different body regions by a human: effects on behaviour and heart rate of dairy cows. *Appl Anim Behav Sci.* (2008) 109:25–38. doi: 10.1016/j.applanim.2007.01.013
53. Reefmann N, Wechsler B, Gyax L. Behavioural and physiological assessment of positive and negative emotion in sheep. *Anim Behav.* (2009) 78:651–9. doi: 10.1016/j.anbehav.2009.06.015
54. Coulon M, Nowak R, Peyrat J, Chandèze H, Boissy A, Boivin X. Do Lambs perceive regular human stroking as pleasant? behavior and heart rate variability analyses. *PLoS ONE.* (2015) 10:e0118617. doi: 10.1371/journal.pone.0118617
55. Proctor HS, Carder G. Can ear postures reliably measure the positive emotional state of cows? *Appl Anim Behav Sci.* (2014) 161:20–7. doi: 10.1016/j.applanim.2014.09.015
56. Lange A, Franzmayr S, Wisenöcker V, Futschik A, Waiblinger S, Lürzel S. Effects of different stroking styles on behaviour and cardiac parameters in heifers. *Animals.* (2020) 10:426. doi: 10.3390/ani10030426
57. Tami G, Gallagher A. Description of the behaviour of domestic dog (*Canis familiaris*) by experienced and inexperienced people. *Appl Anim Behav Sci.* (2009) 120:159–69. doi: 10.1016/j.applanim.2009.06.009
58. Hausberger M, Muller C. A brief note on some possible factors involved in the reactions of horses to humans. *Appl Anim Behav Sci.* (2002) 76:339–44. doi: 10.1016/S0168-1591(02)00016-3
59. McGowan RTS, Rehn T, Norling Y, Keeling LJ. Positive affect and learning: exploring the "Eureka Effect" in dogs. *Anim Cogn.* (2014) 17:577–87. doi: 10.1007/s10071-013-0688-x
60. Caeiro CC, Burrows AM, Waller BM. Development and application of CatFACS: are human cat adopters influenced by cat facial expressions? *Appl Anim Behav Sci.* (2017) 189:66–78. doi: 10.1016/j.applanim.2017.01.005
61. Bertin A, Beraud A, Lansade L, Blache M-C, Diot A, Mulot B, et al. Facial display and blushing: means of visual communication in blue-and-yellow macaws (*Ara Ararauna*)? *PLoS ONE.* (2018) 13:e0201762. doi: 10.1371/journal.pone.0201762
62. Minero M, Tosi MV, Canali E, Wemelsfelder F. Quantitative and qualitative assessment of the response of foals to the presence of an unfamiliar human. *Appl Anim Behav Sci.* (2009) 116:74–81. doi: 10.1016/j.applanim.2008.07.001
63. Sankey C, Richard-Yris M-A, Leroy H, Henry S, Hausberger M. Positive interactions lead to lasting positive memories in horses, *Equus caballus*. *Anim Behav.* (2010) 79:869–75. doi: 10.1016/j.anbehav.2009.12.037
64. Knolle F, Goncalves RP, Morton AJ. Sheep recognize familiar and unfamiliar human faces from two-dimensional images. *Roy Soc Open Sci.* (2017) 4:171228. doi: 10.1098/rsos.171228
65. Kuhne F, Höfler JC, Struwe R. Behavioral and cardiac responses by dogs to physical human-dog contact. *J Vet Behav.* (2014) 9:93–7. doi: 10.1016/j.jveb.2014.02.006
66. Rehn T, Handlin L, Uvnäs-Moberg K, Keeling LJ. Dogs' endocrine and behavioural responses at reunion are affected by how the human initiates contact. *Physiol Behav.* (2014) 124:45–53. doi: 10.1016/j.physbeh.2013.10.009
67. Nagasawa M, Mitsui S, En S, Ohtani N, Ohta M, Sakuma Y, et al. Oxytocin-gaze positive loop and the coevolution of human-dog bonds. *Science.* (2015) 348:333–6. doi: 10.1126/science.1261022
68. Waiblinger S, Menke C, Fölsch DW. Influences on the avoidance and approach behaviour of dairy cows towards humans on 35 farms. *Appl Anim Behav Sci.* (2003) 84:23–39. doi: 10.1016/S0168-1591(03)00148-5
69. Duranton C, Bedossa T, Gaunet F. Interspecific behavioural synchronization: dogs exhibit locomotor synchrony with humans. *Sci Rep.* (2017) 7:12384. doi: 10.1038/s41598-017-12577-z
70. Feldman R. From biological rhythms to social rhythms: physiological precursors of mother-infant synchrony. *Dev Psychol.* (2006) 42:175–88. doi: 10.1037/0012-1649.42.1.175
71. Mariti C, Carlone B, Protti M, Diverio S, Gazzano A. Effects of petting before a brief separation from the owner on dog behavior and physiology: a pilot study. *J Vet Behav.* (2018) 27:41–6. doi: 10.1016/j.jveb.2018.07.003
72. von Borell E, Langbein J, Després G, Hansen S, Leterrier C, Marchant-Forde J, et al. Heart rate variability as a measure of autonomic regulation of cardiac activity for assessing stress and welfare in farm animals—a review. *Physiol Behav.* (2007) 92:293–316. doi: 10.1016/j.physbeh.2007.01.007
73. Rault J-L. Effects of positive and negative human contacts and intranasal oxytocin on cerebrospinal fluid oxytocin. *Psychoneuroendocrino.* (2016) 69:60–6. doi: 10.1016/j.psyneuen.2016.03.015
74. Kis A, Gergely A, Galambos Á, Abdai J, Gombos F, Bódizs R, et al. Sleep macrostructure is modulated by positive and negative social experience in adult pet dogs. *P Roy Soc B: Biol Sci.* (2017) 284:20171883. doi: 10.1098/rspb.2017.1883
75. Lyons DM, Price EO. Relationships between heart rates and behavior of goats in encounters with people. *Appl Anim Behav Sci.* (1987) 18:363–9. doi: 10.1016/0168-1591(87)90230-9
76. Boivin X, Braastad BO. Effects of handling during temporary isolation after early weaning on goat kids' later response to humans. *Appl Anim Behav Sci.* (1996) 48:61–71. doi: 10.1016/0168-1591(95)01019-X
77. Rault J-L, van den Munkhof M, Buisman-Pijlman FTA. Oxytocin as an Indicator of psychological and social well-being in domesticated animals: a critical review. *Front Psychol.* (2017) 8:1521. doi: 10.3389/fpsyg.2017.01521
78. McGowan RTS, Bolte C, Barnett HR, Perez-Camargo G, Martin F. Can you spare 15 min? The measurable positive impact of a 15-min petting session on shelter dog well-being. *Appl Anim Behav Sci.* (2018) 203:42–54. doi: 10.1016/j.applanim.2018.02.011
79. Handlin L, Hydbring-Sandberg E, Nilsson A, Ejdebäck M, Jansson A, Uvnäs-Moberg K. Short-term interaction between dogs and their owners: effects on oxytocin, cortisol, insulin and heart rate—an exploratory study. *Anthrozoös.* (2011) 24:301–15. doi: 10.2752/175303711X13045914865385
80. Rygula R, Pluta H, Popik P. Laughing rats are optimistic. *PLoS ONE.* (2012) 7:e51959. doi: 10.1371/journal.pone.0051959

81. Gygas L, Reefmann N, Pilheden T, Scholkmann F, Keeling L. Dog behavior but not frontal brain reaction changes in repeated positive interactions with a human: a non-invasive pilot study using functional near-infrared spectroscopy (fNIRS). *Behav Brain Res.* (2015) 281:172–6. doi: 10.1016/j.bbr.2014.11.044
82. Guesdon V, Nowak R, Meurisse M, Boivin X, Cornilleau F, Chaillou E, et al. Behavioral evidence of heterospecific bonding between the lamb and the human caregiver and mapping of associated brain network. *Psychoneuroendocrino.* (2016) 71:159–69. doi: 10.1016/j.psyneuen.2016.05.020
83. Spruijt BM, R. van den Bos, Pijlman FTA. A concept of welfare based on reward evaluating mechanisms in the brain: anticipatory behaviour as an indicator for the state of reward systems. *Appl Anim Behav Sci.* (2001) 72:145–71. doi: 10.1016/S0168-1591(00)00204-5
84. Clegg ILK, Rödel HG, Boivin X, Delfour F. Looking forward to interacting with their caretakers: dolphins' anticipatory behaviour indicates motivation to participate in specific events. *Appl Anim Behav Sci.* (2018) 202:85–93. doi: 10.1016/j.applanim.2018.01.015
85. Redcay E, Schilbach L. Using second-person neuroscience to elucidate the mechanisms of social interaction. *Nat Rev Neurosci.* (2019) 20:495–505. doi: 10.1038/s41583-019-0179-4
86. Chartrand TL, Bargh JA. The chameleon effect: the perception–behavior link and social interaction. *J Pers Soc Psychol.* (1999) 76:893–910. doi: 10.1037/0022-3514.76.6.893
87. Petersson M, Uvnäs-Moberg K, Nilsson A, Gustafson L-L, Hydbring-Sandberg E, Handlin L. Oxytocin and cortisol levels in dog owners and their dogs are associated with behavioral patterns: an exploratory study. *Front Psychol.* (2017) 8:1796. doi: 10.3389/fpsyg.2017.01796
88. Gácsi M, Gyori B, Miklósi Á, Virányi Z, Kubinyi E, Topál J, et al. Species-specific differences and similarities in the behavior of hand-raised dog and wolf pups in social situations with humans. *Dev Psychobiol.* (2005) 47:111–22. doi: 10.1002/dev.20082
89. Luescher AU, Reisner IR. Canine aggression toward familiar people: a new look at an old problem. *Vet Clin N Am Small Anim Pract.* (2008) 38:1107–30. doi: 10.1016/j.cvsm.2008.04.008
90. Tallet C, Sy K, Prunier A, Nowak R, Boissy A, Boivin X. Behavioural and physiological reactions of piglets to gentle tactile interactions vary according to their previous experience with humans. *Livest Sci.* (2014) 167:331–41. doi: 10.1016/j.livsci.2014.06.025
91. Wemelsfelder F, Hunter TEA, Mendl MT, Lawrence AB. Assessing the 'whole animal': a free choice profiling approach. *Anim Behav.* (2001) 62:209–20. doi: 10.1006/anbe.2001.1741
92. Hemsworth PH, Barnett JL, Hansen C. The influence of handling by humans on the behaviour, reproduction and corticosteroids of male and female pigs. *Appl Anim Behav Sci.* (1986) 15:303–14. doi: 10.1016/0168-1591(86)90123-1
93. Odendaal JSJ, Meintjes RA. Neurophysiological correlates of affiliative behaviour between humans and dogs. *Vet J.* (2003) 165:296–301. doi: 10.1016/S1090-0233(02)00237-X
94. Coulon M, Nowak R, Andanson S, Ravel C, Marnet PG, Boissy A, et al. Human–lamb bonding: oxytocin, cortisol and behavioural responses of lambs to human contacts and social separation. *Psychoneuroendocrino.* (2013) 38:499–508. doi: 10.1016/j.psyneuen.2012.07.008
95. Lürzel S, Bückendorf L, Waiblinger S, Rault J-L. Salivary oxytocin in pigs, cattle, and goats during positive human-animal interactions. *Psychoneuroendocrino.* (2020) 115:104636. doi: 10.1016/j.psyneuen.2020.104636
96. Feh C, de Mazières J. Grooming at a preferred site reduces heart rate in horses. *Anim Behav.* (1993) 46:1191–4. doi: 10.1006/anbe.1993.1309
97. Lange A, Bauer L, Futschik A, Waiblinger S and Lürzel S. Talking to Cows: Reactions to Different Auditory Stimuli During Gentle Human-Animal Interactions. *Front. Psychol.* (2020) 11:579346. doi: 10.3389/fpsyg.2020.579346
98. Hasin Y, Seldin M, Lusis A. Multi-omics approaches to disease. *Genome Biol.* (2017) 18:83. doi: 10.1186/s13059-017-1215-1
99. Nerem R, Levesque M, Cornhill J. Social environment as a factor in diet-induced atherosclerosis. *Science.* (1980) 208:1475–6. doi: 10.1126/science.7384790
100. Gross WB, Siegel PB. Effects of early environmental stresses on chicken body weight, antibody response to RBC antigens, feed efficiency and response to handling. *Avian Dis.* (1979) 24:569–79. doi: 10.2307/1589792
101. Zulkifli I, Gilbert J, Liew PK, Ginsos J. The effects of regular visual contact with human beings on fear, stress, antibody and growth responses in broiler chickens. *Appl Anim Behav Sci.* (2002) 79:103–12. doi: 10.1016/S0168-1591(02)00135-1
102. Mendl M, Burman OHP, Parker RMA, Paul ES. Cognitive bias as an indicator of animal emotion and welfare: Emerging evidence and underlying mechanisms. *Appl Anim Behav Sci.* (2009) 118:161–81. doi: 10.1016/j.applanim.2009.02.023
103. Mendl M, Brooks J, Basse C, Burman O, Paul E, Blackwell E, et al. Dogs showing separation-related behaviour exhibit a 'pessimistic' cognitive bias. *Chor Biol.* (2010) 20:R839–40. doi: 10.1016/j.cub.2010.08.030
104. Kendrick KM, Atkins K, Hinton MR, Broad KD, Fabre-Nys C, Keverne B. Facial and vocal discrimination in sheep. *Anim Behav.* (1995) 49:1665–76. doi: 10.1016/0003-3472(95)90088-8
105. Andics A, Miklósi Á. Neural processes of vocal social perception: Dog-human comparative fMRI studies. *Neurosci Biobehav Rev.* (2018) 85:54–64. doi: 10.1016/j.neubiorev.2017.11.017
106. Reiter D. *The Influence of the Milker's Behaviour on Social Behaviour of Dairy Cows.* Vienna: University of Veterinary Medicine, Vienna (2018).
107. Lürzel S, Windschnurer I, Futschik A, Palme R, Waiblinger S. Effects of gentle interactions on the relationship with humans and on stress-related parameters in group-housed calves. *Anim Welfare.* (2015) 24:475–84. doi: 10.7120/09627286.24.4.475
108. Collins JW, Siegel PB. Human handling, flock size and responses to an E. coli challenge in young chickens. *Appl Anim Behav Sci.* (1987) 19:183–8. doi: 10.1016/0168-1591(87)90215-2
109. Dawkins MS. The science of animal suffering. *Ethology.* (2008) 114:937–45. doi: 10.1111/j.1439-0310.2008.01557.x
110. Kirkden RD, Pajor EA. Using preference, motivation and aversion tests to ask scientific questions about animals' feelings. *Appl Anim Behav Sci.* (2006) 100:29–47. doi: 10.1016/j.applanim.2006.04.009
111. Waiblinger S, Wagner K, Hillmann E, Barth K. Short- and long-term effects of rearing dairy calves with contact to their mother on their reactions towards humans. *J Dairy Res.* (2020) 87:148–53. doi: 10.1017/S0022029920000576
112. Hemsworth PH, Brand A, Willems P. The behavioural response of sows to the presence of human beings and its relation to productivity. *Livest Prod Sci.* (1981) 8:67–74. doi: 10.1016/0301-6226(81)90031-2
113. Windschnurer I, Schmied C, Boivin X, Waiblinger S. Reliability and inter-test relationship of tests for on-farm assessment of dairy cows' relationship to humans. *Appl Anim Behav Sci.* (2008) 114:37–53. doi: 10.1016/j.applanim.2008.01.017
114. Spinka M. Animal agency, animal awareness and animal welfare. *Anim Welfare.* (2019) 28:11–20. doi: 10.7120/09627286.28.1.011
115. Baciadonna L, Nawroth C, McElligott A. Judgement bias in goats (*Capra hircus*): investigating the effects of human grooming. *PeerJ.* (2016) 4:e2485. doi: 10.7717/peerj.2485
116. Destrez A, Coulon M, Deiss V, Delval E, Boissy A, Boivin X. The valence of the long-lasting emotional experiences with various handlers modulates discrimination and generalization of individual humans in sheep. *J Anim Sci.* (2013) 91:5418–26. doi: 10.2527/jas.2012-5654
117. Chijiwa H, Kuroshima H, Hori Y, Anderson JR, Fujita K. Dogs avoid people who behave negatively to their owner: third-party affective evaluation. *Anim Behav.* (2015) 106:123–7. doi: 10.1016/j.anbehav.2015.05.018
118. Muvhali PT. Improving ostrich welfare by developing positive human-animal interactions (thesis). University of Stellenbosch, Stellenbosch, South Africa (2018).
119. Jones RB. Reduction of the domestic chick's fear of human beings by regular handling and related treatments. *Anim Behav.* (1993) 46:991–8. doi: 10.1006/anbe.1993.1280
120. Barnett JL, Hemsworth PH, Hennessy DP, McCallum TH, Newman EA. The effects of modifying the amount of human contact on behavioural, physiological and production responses of laying hens. *Appl Anim Behav Sci.* (1994) 41:87–100. doi: 10.1016/0168-1591(94)90054-X
121. Hemsworth PH, Barnett JL, Tilbrook AJ, Hansen C. The effects of handling by humans at calving and during milking on the behaviour and milk cortisol

- concentrations of primiparous dairy cows. *Appl Anim Behav Sci.* (1989) 22:313–26. doi: 10.1016/0168-1591(89)90026-9
122. Heird JC, Whitaker DD, Bell RW, Ramsey CB, Lokey CE. The effects of handling at different ages on the subsequent learning ability of 2-year-old horses. *Appl Anim Behav Sci.* (1986) 15:15–25. doi: 10.1016/0168-1591(86)90018-3
 123. Bertenshaw C, Rowlinson P, Edge H, Douglas S, Shiel R. The effect of different degrees of 'positive' human-animal interaction during rearing on the welfare and subsequent production of commercial dairy heifers. *Appl Anim Behav Sci.* (2008) 114:65–75. doi: 10.1016/j.applanim.2007.12.002
 124. Hemsworth PH, Barnett JL. The effects of early contact with humans on the subsequent level of fear of humans in pigs. *Appl Anim Behav Sci.* (1992) 35:83–90. doi: 10.1016/0168-1591(92)90018-7
 125. LaFollette MR, O'Haire ME, Cloutier S, Blankenberger WB, Gaskill BN. Rat tickling: a systematic review of applications, outcomes, and moderators. *PLoS ONE.* (2017) 12:e0175320. doi: 10.1371/journal.pone.0175320
 126. Rooney NJ, Cowan S. Training methods and owner-dog interactions: links with dog behaviour and learning ability. *Appl Anim Behav Sci.* (2011) 132:169–77. doi: 10.1016/j.applanim.2011.03.007
 127. Pratsch L, Mohr N, Palme R, Rost J, Troxler J, Arhant C. Carrier training cats reduces stress on transport to a veterinary practice. *Appl Anim Behav Sci.* (2018) 206:64–74. doi: 10.1016/j.applanim.2018.05.025
 128. Laule GE, Bloomsmith MA, Schapiro SJ. The use of positive reinforcement training techniques to enhance the care, management, and welfare of primates in the laboratory. *J Appl Anim Welfare Sci.* (2003) 6:163–73. doi: 10.1207/S15327604JAWS0603_02
 129. Muns R, Rault J-L, Hemsworth P. Positive human contact on the first day of life alters the piglet's behavioural response to humans and husbandry practices. *Physiol Behav.* (2015) 151:162–7. doi: 10.1016/j.physbeh.2015.06.030
 130. Tallet C, Rakotomahandy M, Guérin C, Lemasson A, Hausberger M. Postnatal auditory preferences in piglets differ according to maternal emotional experience with the same sounds during gestation. *Sci Rep.* (2016) 6:37238. doi: 10.1038/srep37238
 131. Tosi MV, Hemsworth PH. Stockperson-and husbandry interactions animal welfare in the extensive livestock industries. *Int Soc Appl Ethol.* (2002) 129. Available online at: https://www.applied-ethology.org/res/2002%20isae%20in%20egmond%20aan%20zee_%20netherlands.pdf
 132. Scott JP. The process of primary socialization in canine and human infants. *Monogr Soc Res Child Dev.* (1963) 28:1–47. doi: 10.2307/1165508
 133. Foster T, DeFazio R, Bizon J. Characterizing cognitive aging of spatial and contextual memory in animal models. *Front Aging Neurosci.* (2012) 4:12. doi: 10.3389/fnagi.2012.00012
 134. Boivin X, Neindre PL, Chupin JM. Establishment of cattle-human relationships. *Appl Anim Behav Sci.* (1992) 32:325–35. doi: 10.1016/S0168-1591(05)80025-5
 135. Henry S, Hemery D, Richard MA, Hausberger M. Human-mare relationships and behaviour of foals toward humans. *Appl Anim Behav Sci.* (2005) 93:341–62. doi: 10.1016/j.applanim.2005.01.008
 136. Tallet C, Leribillard O, Rault J-L, Meunier-Salaün M-C. Social transmission of approach behaviour toward humans in pigs [In French]. *Conference Proceedings of the annual congress of the French society for the study of animal behaviour (SFECA).* Rennes, France (2018).
 137. Bertin A, Richard-Yris M-A, Houdelier C, Lumineau S, Möstl E, Kuchar A, et al. Habituation to humans affects yolk steroid levels and offspring phenotype in quail. *Horm Behav.* (2008) 54:396–402. doi: 10.1016/j.yhbeh.2008.04.012
 138. Price EO, Thos J. Behavioral responses to short-term social isolation in sheep and goats. *Appl Anim Ethol.* (1980) 6:331–9. doi: 10.1016/0304-3762(80)90133-9
 139. Munksgaard L, De Passillé AM, Rushen J, Thodberg K, Jensen MB. Discrimination of people by dairy cows based on handling. *J Dairy Sci.* (1997) 80:1106–12. doi: 10.3168/jds.S0022-0302(97)76036-3
 140. Markowitz TM, Dally MR, Gursky K, Price EO. Early handling increases lamb affinity for humans. *Anim Behav.* (1998) 55:573–87. doi: 10.1006/anbe.1997.0640
 141. Hubrecht RC. A comparison of social and environmental enrichment methods for laboratory housed dogs. *Appl Anim Behav Sci.* (1993) 37:345–61. doi: 10.1016/0168-1591(93)90123-7
 142. Hubrecht RC. Enrichment in puppyhood and its effects on later behavior of dogs. *Lab Anim Sci.* (1995) 45:70–5.
 143. Fureix C, Jégo P, Sankey C, Hausberger M. How horses (Equus caballus) see the world: humans as significant objects. *Anim Cogn.* (2009) 12:643–54. doi: 10.1007/s10071-009-0223-2
 144. Boivin X, Garel JP, Mante A, Le Neindre P. Beef calves react differently to different handlers according to the test situation and their previous interactions with their caretaker. *Appl Anim Behav Sci.* (1998) 55:245–57. doi: 10.1016/S0168-1591(97)00050-6
 145. de Passillé AM, Rushen J, Ladewig J, Petherick C. Dairy calves' discrimination of people based on previous handling. *J Anim Sci.* (1996) 74:969–74. doi: 10.2527/1996.745969x
 146. Hemsworth PH, Coleman GJ, Cox M, Barnett JL. Stimulus generalization: the inability of pigs to discriminate between humans on the basis of their previous handling experience. *Appl Anim Behav Sci.* (1994) 40:129–42. doi: 10.1016/0168-1591(94)90077-9
 147. Trut L, Oskina I, Kharlamova A. Animal evolution during domestication: the domesticated fox as a model. *BioEssays.* (2009) 31:349–60. doi: 10.1002/bies.200800070
 148. Boivin X, Gilard F, Egal D. The effect of early human contact and the separation method from the dam on responses of beef calves to humans. *Appl Anim Behav Sci.* (2009) 120:132–9. doi: 10.1016/j.applanim.2009.05.011
 149. Nagasawa M, Murai K, Mogi K, Kikusui T. Dogs can discriminate human smiling faces from blank expressions. *Anim Cogn.* (2011) 14:525–33. doi: 10.1007/s10071-011-0386-5
 150. Corsin Müller A, Schmitt K, Anjuli Barber LA, Huber L. Dogs can discriminate emotional expressions of human faces. *Curr Biol.* (2015) 25:601–5. doi: 10.1016/j.cub.2014.12.055
 151. Albuquerque N, Guo K, Wilkinson A, Savalli C, Otta E, Mills D. Dogs recognize dog and human emotions. *Biol Lett.* (2016) 12:20150883. doi: 10.1098/rsbl.2015.0883
 152. Smith AV, Proops L, Grounds K, Wathan J, McComb K. Functionally relevant responses to human facial expressions of emotion in the domestic horse (Equus caballus). *Biol Lett.* (2016) 12:20150907. doi: 10.1098/rsbl.2015.0907
 153. Nawroth C, Albuquerque N, Savalli C, Single M-S, McElligott AG. Goats prefer positive human emotional facial expressions. *Roy Soc Open Sci.* (2018) 5:180491. doi: 10.1098/rsos.180491
 154. Merola I, Lazzaroni M, Marshall-Pescini S, Prato-Previde E. Social referencing and cat-human communication. *Anim Cogn.* (2015) 18:639–48. doi: 10.1007/s10071-014-0832-2
 155. Arhant C, Beetz AM, Troxler J. Caregiver reports of interactions between children up to 6 years and their family dog—implications for dog bite prevention. *Front Vet Sci.* (2017) 4:130. doi: 10.3389/fvets.2017.00130
 156. Coppola CL, Grandin T, Enns RM. Human interaction and cortisol: can human contact reduce stress for shelter dogs? *Physiol Behav.* (2006) 87:537–41. doi: 10.1016/j.physbeh.2005.12.001
 157. Luescher AU, Tyson Medlock R. The effects of training and environmental alterations on adoption success of shelter dogs. *Appl Anim Behav Sci.* (2009) 117:63–8. doi: 10.1016/j.applanim.2008.11.001
 158. Normando S, Corain L, Salvadoretti M, Meers L, Valsecchi P. Effects of an Enhanced Human Interaction Program on shelter dogs' behaviour analysed using a novel nonparametric test. *Appl Anim Behav Sci.* (2009) 116:211–9. doi: 10.1016/j.applanim.2008.10.005
 159. De Meyer D, Amalraj A, Van Limbergen T, Fockedekey M, Edwards S, Moustsen VA, et al. Short Communication: effect of positive handling of sows on litter performance and pre-weaning piglet mortality. *Animal.* (2020) 14:1733–9. doi: 10.1017/S1751731120000427
 160. Waiblinger S, Menke C, Korff J, Bucher A. Previous handling and gentle interactions affect behaviour and heart rate of dairy cows during a veterinary procedure. *Appl Anim Behav Sci.* (2004) 85:31–42. doi: 10.1016/j.applanim.2003.07.002
 161. Pedersen V, Barnett JL, Hemsworth PH, Newman EA, Schirmer B. The effects of handling on behavioural and physiological responses to housing in tether-stalls among pregnant pigs. *Anim Welfare.* (1998) 7:137–50.

162. Cimarelli G, Turcsán B, Bánlaki Z, Range F, Virányi Z. Dog owners' interaction styles: their components and associations with reactions of pet dogs to a social threat. *Front Psychol.* (2016) 7:1979. doi: 10.3389/fpsyg.2016.01979
163. Bloomfield RC, Gillespie GR, Kerswell KJ, Butler KL, Hemsworth PH. Effect of partial covering of the visitor viewing area window on positioning and orientation of zoo orangutans: a preference test. *Zoo Biol.* (2015) 34:223–9. doi: 10.1002/zoo.21207
164. Boivin X, Le Neindre P, Chupin JM, Garel JP, Trillat G. Influence of breed and early management on ease of handling and open-field behaviour of cattle. *Appl Anim Behav Sci* (1992) 32:313–23. doi: 10.1016/S0168-1591(05)80024-3
165. Lyons DM. Individual differences in temperament of dairy goats and the inhibition of milk ejection. *Appl Anim Behav Sci.* (1989) 22:269–82. doi: 10.1016/0168-1591(89)90022-1
166. Day JEL, Spoolder HAM, Burfoot A, Chamberlain HL, Edwards SA. The separate and interactive effects of handling and environmental enrichment on the behaviour and welfare of growing pigs. *Appl Anim Behav Sci.* (2002) 75:177–92. doi: 10.1016/S0168-1591(01)00199-X
167. Lindahl C, Pinzke S, Herlin A, Keeling LJ. Human-animal interactions and safety during dairy cattle handling—comparing moving cows to milking and hoof trimming. *J Dairy Sci.* (2016) 99:2131–41. doi: 10.3168/jds.2014-9210
168. Lawrence AB, Vigers B, Sandøe P. What is so positive about positive animal welfare?—a critical review of the literature. *Animals.* (2019) 9:783. doi: 10.3390/ani9100783

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Rault, Waiblinger, Boivin and Hemsworth. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Rault, J-L;Waiblinger, S;Boivin, X;Hemsworth, P

Title:

The Power of a Positive Human-Animal Relationship for Animal Welfare

Date:

2020-11-09

Citation:

Rault, J. -L., Waiblinger, S., Boivin, X. & Hemsworth, P. (2020). The Power of a Positive Human-Animal Relationship for Animal Welfare. FRONTIERS IN VETERINARY SCIENCE, 7, <https://doi.org/10.3389/fvets.2020.590867>.

Persistent Link:

<http://hdl.handle.net/11343/253054>

License:

[CC BY](#)