

RESEARCH ARTICLE

The prehistoric roots of Chinese cuisines: Mapping staple food systems of China, 6000 BC–220 AD

Xinyi Liu^{1*}, Rachel E. B. Reid²

1 Department of Anthropology, Washington University in St. Louis, St. Louis, Missouri, United States of America, **2** Department of Geosciences, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, United States of America

* liuxinyi@wustl.edu**OPEN ACCESS**

Citation: Liu X, Reid REB (2020) The prehistoric roots of Chinese cuisines: Mapping staple food systems of China, 6000 BC–220 AD. PLoS ONE 15(11): e0240930. <https://doi.org/10.1371/journal.pone.0240930>

Editor: Peter F. Biehl, University at Buffalo - The State University of New York, UNITED STATES

Received: June 24, 2020

Accepted: October 5, 2020

Published: November 4, 2020

Copyright: © 2020 Liu, Reid. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: National Science Foundation, under grant 1826727, "The origins and spread of millet cultivation", PI: X. Liu; the European Research Council, under grant 249642, "Food Globalisation in Prehistory", PI, M. Jones. Sponsors did not play any role in research, decision to publish, or preparation of the manuscript.

Competing interests: No authors have competing interests.

Abstract

We conducted a meta-analysis of published carbon and nitrogen isotope data from archaeological human skeletal remains ($n = 2448$) from 128 sites across China in order to investigate broad spatial and temporal patterns in the formation of staple cuisines. Between 6000–5000 cal BC we found evidence for an already distinct north versus south divide in the use of main crop staples (namely millet vs. a broad spectrum of C_3 plant based diet including rice) that became more pronounced between 5000–2000 cal BC. We infer that this pattern can be understood as a difference in the spectrum of subsistence activities employed in the Loess Plateau and the Yangtze-Huai regions, which can be partly explained by differences in environmental conditions. We argue that regional differentiation in dietary tradition are not driven by differences in the conventional "stages" of shifting modes of subsistence (hunting-foraging-pastoralism-farming), but rather by myriad subsistence choices that combined and discarded modes in a number of innovative ways over thousands of years. The introduction of wheat and barley from southwestern Asia after 2000 cal BC resulted in the development of an additional east to west gradient in the degree of incorporation of the different staple products into human diets. Wheat and barley were rapidly adopted as staple foods in the Continental Interior contra the very gradual pace of adoption of these western crops in the Loess Plateau. While environmental and social factors likely contributed to their slow adoption, we explored local cooking practice as a third explanation; wheat and barley may have been more readily folded into grinding-and-baking cooking traditions than into steaming-and-boiling traditions. Changes in these culinary practices may have begun in the female sector of society.

Introduction

Staple foods pass through a long transformative process as they are acquired, prepared, and distributed by human societies, and the performances of staple food preparation and presentation are intimately connected with social relationships [1]. Recent investigations have shown that between 5000 and 1500 cal BC, the Eurasian and African landmass underpinned a

continental-scale process of food ‘globalisation’ of staple crops [2, 3]. By 1500 cal BC, the process brought together previously isolated agricultural systems to form a new kind of management system that enabled multi-cropping and fundamentally transformed Eurasian diets. China plays an important role in this narrative for its diverse forms of food products in the Neolithic but also as both the source of eastern domesticates (e.g., rice, broomcorn and foxtail millet) that moved from China to the West, and the recipient of southwest Asian grains (i.e. wheat and barley) that moved east-wards. Understanding the prehistoric roots of Chinese staple cuisines provides perspectives that not only transform our knowledge of the past but also raise awareness of the present and future utilities of these cereals.

Cultivation of staple cereals has played a vital role in the development of many aspects of Chinese culture from prehistory to today. Globally, the process employs millions of people and presently feeds 20% of the world’s population [4]. Cereals are the most important food source in the world, contributing as much as 70% of energy intake in developing countries [5]. In China, cereal foods such as rice and wheat products contributed 75–85% towards the daily dietary intake for average low/medium-income individuals in the 1980s [6]. Regional variations in cereal management and choice of staple products provide a key to understanding food production and consumption in China. These variations (e.g., rice in the lower Yangtze, wheat in the northwest, barley in Tibet) have been well documented historically [4, 7].

The diversity in staple choices has often been linked to origins of the diversity of regional cooking techniques. Early communities in East and West Asia, for example, were characterized by differences in food processing technologies: culinary traditions based on boiling and steaming of grain in the East, and by grinding grain and baking the resulting flour in the West. While the Pre-Pottery Neolithic cultures of Southwest Asia made extensive use of querns for flour production and constructed clay ovens for baking bread and roasting foods, cultures in Neolithic China elaborated forms of ceramic vessels for boiling, steaming, and serving [8]. Current evidence places pottery in south China *c.* 18,000 years ago, associated with hunter-gathers [13]. By contrast, in Southwest Asia, ceramics developed relatively late, dated *c.* 8,500 years ago [8]. This contrast has led to the hypothesis that these distinct East-West cooking technologies are deep seated in cultural differences between peoples that predate domestication [8]. In the context of early globalization of staple crops, the dispersal of cereals into new areas was not necessarily accompanied by the dispersal of culinary traditions. Novel grains could sometimes be incorporated into existing local traditions of food processing or sometimes lose the status of being staple grain.

Here, we integrate a large body of isotopic data from both English and Chinese publications to explore broad spatial and temporal patterns in the prehistoric roots of Chinese staple cuisines and assess possible gender distinctions in the context of staple consumption. Other recent reviews of this literature, though are not as broad in geographic scope, have shed important light on this topic [55, 9]. We additionally explore the nature of regional differences in staple traditions and consider the context in which culinary innovation arose.

Isotopic values from archaeological skeletons provide direct proxies for long-term consumption practices of individuals in the past. Due to the resolution of this technique—which doesn’t enable assessment of the contribution of minor dietary components—we focus our discussion on the consumption of staple foods. Carbon isotope values ($\delta^{13}\text{C}$) vary primarily according to the photosynthetic pathways employed by plants at the base of the food chain [10, 11]. The potential for using $\delta^{13}\text{C}$ values to differentiate between different types of cereal diets was first realized in detecting the C_4 domesticate maize (*Zea mays*) in Americas [10]. Nitrogen isotope values ($\delta^{15}\text{N}$) provide further information about past diets by situating the consumers on the trophic food chain; $\delta^{15}\text{N}$ values increase by 3–5 ‰ with each step in a trophic chain [12]. Nonetheless, without site specific faunal baseline data, dietary reconstruction with $\delta^{15}\text{N}$

values at this broad geographic scope is challenging. There are now over 90 publications presenting isotopic results from >120 sites and >2000 human individuals from prehistoric China. We compiled these data to investigate the historic geography of staple cuisines between 6000 cal BC and 220 cal AD, capitalizing on the contrasting isotopic signatures among major crop domesticates, including rice, wheat and barley (C_3 plants), and broomcorn and foxtail millets (C_4 plants). We focus on three regions featuring differing environmental characteristics and distinct agricultural and culinary traditions: (1) the broad Loess Plateau including the Yellow, Wei and Xiliao Rivers, (2) the Yangtze and Huai Rivers, and (3) the Continental Interior bordering the Loess Plateau and Eurasian steppe including the Tibetan Plateau (Fig 1).

Setting up the geography

China's vast landmass ranges across contrasting ecological extremes, from tropical in the south, to sub-Arctic in the north, alpine in the west and marshy lowlands in the east [13]. A key dynamic climatic element is the monsoonal system, comprising a warm, wet summer monsoon, and a cold, dry winter monsoon. The summer monsoon brings water from the Pacific and Indian Oceans onto much of the east and south of China while the winter monsoon drives the movement of Aeolian dust from the Gobi desert to the Loess Plateau. The sensitivity of the monsoonal system to fluctuations in the relative temperatures of land and ocean has rendered it the most variable part of the physical environment, critically affecting water availability in many parts of China, particularly towards the south and east [13]. These features have led to an agriculture that is diverse in its crop ecology, elaborate in its management of water, and with its most intense sedentary cultivations in the east of the country, including the broad Loess Plateau and the Yangtze and Huai Rivers, which is geographically divided by the Qinling Mountains and Huai River [14, 15].

The central/eastern parts of China host the most productive soils and have an enduring association with important staple cereals: the Yangtze and Huai Rivers with rice (*Oryza sativa*), and the Loess Plateau with broomcorn and foxtail millet (*Panicum miliaceum* and *Seteria italica*) [13, 16]. The oldest archaeological sites preserving broomcorn and foxtail millet remains do not, however, lie in direct proximity to the great rivers. Sites with millet remains are instead located along the foothills of the eastern edge of the Loess Plateau at a considerable distance from the rivers themselves [17]. The earliest sites with rice are situated in the middle and lower Yangtze and Huai River valleys [18] at locations associated with minor tributaries and intermountain plains where cultivation could be easily managed [19]. In the archaeobotanical record dating to before 5000 cal. BC, a north-south divide is observable on either side of the Huai River–Qinling Mountain line, a topographical reference used by modern geographers to distinguish between north and south China. North of this line, millet cultivation was predominant in the Loess Plateau, while south of the line, subsistence was based on a diverse spectrum of food resources including cultivation of rice and managing free-living plants prevalent in the Yangtze-Huai Region [15, 16].

The same general area of central/eastern China came into contact with Central Asia and possibly South Asia in the Bronze Age, facilitating the adoption of a variety of cereal crops originating in the west [2]. To the west and north of this principal area of Chinese agriculture, the Loess Plateau and the upper Yangtze is flanked by the mountainous Continental Interior. This includes the Mongolian plateau, the Gobi desert, the Hexi Corridor, western Sichuan and northern Yunnan bordering the eastern Tibetan Plateau, as well as the northern and eastern parts of the Tibetan Plateau itself. In the context of a trans-Eurasian exchange of cereal crops, the founder crops from the Fertile Crescent (modern-day Iran, Iraq, Syria and southern Turkey)—notably free-threshing wheat (*Triticum aestivum*) and naked barley (*Hordeum vulgare*)—




Fig 1. Site maps. (a) Site locations with isotope (white circles) and archaeobotanical (black triangles) data (see Tables 1–3, S1 for isotope studies and S2 for archaeobotanical). (b) Proposed culinary traditions in China after 2000 cal BC as described in the discussion. Regional difference in cooking follows the hypothesis proposed by Fuller and Rowlands (2011). Map generated using ArcMap v. 10.2 and NASA Blue Marble with data set obtained from NASA Earth Observatory (public domain). See: <http://earthobservatory.nasa.gov/Features/BlueMarble/>.

<https://doi.org/10.1371/journal.pone.0240930.g001>

were introduced to China between 3000 and 1500 cal BC possibly along multiple routes through the Continental Interior [20]. The introduction of the Fertile Crescent grains in the Bronze Age significantly transformed the staple food system in China.

Prehistoric people did not subsist on cereals alone. Archaeobotanical evidence shows that, since the terminal Pleistocene, a variety of plants including acorns, beans, tubers and grasses (*Triticeae* and *Paniceae*) were used in the Loess Plateau [21]. Over the course of the Neolithic (c. 8000–1500 BC), additional plant and animal domesticates were introduced into human and animal diets, including pigs (*Sus scrofa*), soybean (*Glycine max*), adzuki bean (*Vigna angularis*), buckwheat (*Fagopyrum esculentum*), and hemp seed (*Cannabis sativa*) [13, 22]. Recent research shows that in the Yangtze-Huai region, rice cultivation emerged in the context of broad spectrum foraging focused on the collection of tree nuts, especially acorns (*Quercus* spp.), fruits such as peaches and apricots (*Prunus* spp.), and wetland nuts and tubers, including water chestnuts (*Trapa natans*), foxnuts (*Euryale ferox*), lotus root (*Nelumbo nucifera*), job's tears (*Coix lachrymal-jobi*) and barnyard grasses (*Echinochloa* spp.) [23–25]. With the exception of job's tears and some members of *Triticeae* and *Paniceae*, all the fruits, nuts, tubers and beans identified employ the C₃ photosynthetic pathway.

Materials and methods

To locate published archaeological isotopic studies from China, we searched Web of Science and Google Scholar using combinations of the following keywords: stable isotopes, China, human diet, bone collagen, and apatite. To include data published in Chinese, we searched the China Academic Journals Database using the same set of keywords. We restricted our search to articles concerned with post-Paleolithic archaeological sites and specimens dating to before 220 AD, the ending point of the Han Dynasty. Our search yielded isotopic data from 128 sites in over 90 articles published in English and Chinese between 1984 and 2018 (Tables 1–3, S1 Table). The articles are primarily concerned with $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values from archaeological human bone collagen (n = 83, including 7 review articles), although a subset includes carbon and oxygen isotope data ($\delta^{13}\text{C}$ and $\delta^{18}\text{O}$) from bone apatite and/or tooth enamel (n = 6). Several recent articles also feature sulphur isotope data ($\delta^{34}\text{S}$), but these are presently few in numbers (n = 6). We did not consider articles focusing on strontium (Sr) isotope ratios, which are commonly used as a geographical fingerprinting tool. As we only use published data for the meta-analysis, no permits were required for this study, which compiled with all relevant regulations. All published data compiled in this study is presented in S1 Table and summarized in Tables 1–3 with references to the original studies. Specimen IDs (where are available) as given in the original isotopic studies are also presented in S1 Table.

We performed all statistical analyses in R [26]. Before beginning analyses, we filtered out samples with poor C:N ratios (< 2.9 or > 3.6) suggesting that they were contaminated or poorly preserved. We described the basic structure of the data by calculating the mean and standard

Table 1. Published human C and N isotope values from before 5000 cal BC.

Site	Site number	Region	Province	Cultural group	Estimated dates (cal. BC/AD)	Mean $\delta^{15}\text{N}$	SD $\delta^{15}\text{N}$	Mean $\delta^{13}\text{C}$	SD $\delta^{13}\text{C}$	n	Reference number
Xinglonggou	1	A	Inner Mongolia	Xinglongwa	6200–5300	9.8	0.8	-9.9	1.1	30	[43, 59]
Xinglongwa	2	A	Inner Mongolia	Xinglongwa	6200–5300	-	-	-8.9	1.7	7	[59]
Xiaojingshan	3	A	Shandong	Houli	6200–5500	9.0	0.6	-17.8	0.3	10	[60]
Baijia	4	A	Shaanxi	Laoguantai	5700–5300	10.9	1.7	-13.7	1.9	3	[61]
Beiliu	5	A	Shaanxi	Laoguantai	6000–5000	8.5	0.1	-12.0	0.6	6	[62, 63]
Jiahu	6	B	Henan	Jiahu	7000–5800	8.9	0.9	-20.3	0.5	14	[64]

Summary of published carbon and nitrogen isotope values measured in human bone collagen from sites dating to before 5000 cal BC.

<https://doi.org/10.1371/journal.pone.0240930.t001>

Table 2. Published human C and N isotope values from between 5000 and 2000 cal BC.

Site	Site number	Region	Province	Cultural group	Estimated dates (cal. BC/AD)	Mean $\delta^{15}\text{N}$	SD $\delta^{15}\text{N}$	Mean $\delta^{13}\text{C}$	SD $\delta^{13}\text{C}$	n	Reference number
Baiyinchanghan	7	A	Inner Mongolia	Hongshan	4300–3900	8.6	0.3	-8.8	0.4	3	[43]
Caomaoshan	8	A	Inner Mongolia	Hongshan	3400–3100	9.1	0.4	-9.3	0.6	7	[43]
Dakou	9	A	Inner Mongolia	Dakou	2300–1900	7.5	1.0	-8.9	1.8	2	[65]
Miaozigou	10	A	Inner Mongolia	Miaozigou	3500–3000	9.2	0.2	-7.2	0.2	9	[66]
Xinglongwa	11	A	Inner Mongolia	Hongshan	4500–3000	8.7	-	-5.4	-	1	[59]
Xishan	12	A	Inner Mongolia	Xiaoheyuan	2900–2400	8.8	0.4	-7.5	0.5	16	[42]
Qingliangsi	13	A	Shanxi	Yangshao/Longshan	5000–2000	8.5	1.1	-8.1	0.9	27	[67]
Taosi	14	A	Shanxi	Longshan	2500–2000	8.9	1.3	-7.3	2.4	15	[68, 69]
Xinhuaacun	15	A	Shanxi	Longshan	2300–2000	6.5	1.4	-7.3	0.7	2	[65]
Beiqian	16	A	Shandong	Dawenkou	4000–3000	8.8	1.0	-9.6	0.8	38	[44]
Beizhuang	17	A	Shandong	Beizhuang	4500–2500	13.2	-	-7.9	-	1	[59]
Guzhendu	18	A	Shandong	Dawenkou	4000–3000	9.6	-	-8.5	0.7	4	[59]
Xigongqiao	19	A	Shandong	Dawenkou	4000–3000	8.1	2.1	-15.3	3.8	10	[70]
Banpo	20	A	Shaanxi	Yangshao	4800–4300	9.1	-	-15.0	-	1	[42]
Beiliu	21	A	Shaanxi	Yangshao	4000–3500	8.7	0.9	-12.0	1.4	6	[62, 63]
Beishouling	22	A	Shaanxi	Yangshao	5000–3500	-	-	-13.8	0.9	3	[68]
Dongying II	23	A	Shaanxi	Longshan	2600–2000	9.4	0.3	-8.0	1.3	5	[71]
Jiangzhai	24	A	Shaanxi	Yangshao	4900–4000	8.6	0.6	-9.9	1.2	20	[42, 72]
Quanhucun	25	A	Shaanxi	Yangshao	3500–3000	11.5	-	-11.2	-	1	[73]
Shengedaliang	26	A	Shaanxi	Longshan	2500–2000	8.8	1.4	-8.5	1.8	28	[74]
Shijia	27	A	Shaanxi	Yangshao	4300–4000	8.1	0.5	-10.0	0.7	9	[42]
Xipo	28	A	Shaanxi	Yangshao	4000–3300	9.4	1.0	-9.7	1.1	31	[75]
Xunyi	29	A	Shaanxi	Longshan	2400–2000	8.2	0.1	-7.1	0.1	3	[65]
Yuhuazhai	30	A	Shaanxi	Yangshao	5000–3000	8.4	1.9	-10.9	4.4	35	[65, 75]
Zhouyuan	31	A	Shaanxi	Longshan	2500–2000	6.7	3.9	-8.0	0.7	5	[76]
Guanjia	32	A	Henan	Yangshao	4000–3500	6.2	0.7	-8.0	0.6	21	[50]
Wadian	33	A	Henan	Wangwan III	2400–2000	8.2	1.3	-11.0	2.1	12	[77]
Xiaowu	34	A	Henan	Yangshao	5000–4000	7.8	0.8	-10.3	1.2	74	[78]
Xishan	35	A	Henan	Yangshao	5000–3000	9.0	0.8	-8.2	1.5	39	[75]
Gouwan	36	B	Henan	Yangshao/Qujialing	5000–2600	8.3	1.1	-14.3	1.9	41	[79]
Haojiatai	37	B	Henan	Longshan	2500–2000	9.2	1.1	-13.1	4.9	11	[130]
Jiazhuang	38	B	Henan	Longshan	2500–2000	12.7	-	-19.1	-	1	[130]
Meishan	39	B	Henan	Longshan	2500–2000	10.2	1.5	-15.0	2.8	4	[130]
Pingliangtai	40	B	Henan	Longshan	2500–2000	9.0	1.0	-8.7	1.2	8	[130]
Xiazhai	41	B	Henan	Longshan	2500–2000	8.2	0.7	-10.2	1.9	22	[130]
Sanxingcun	42	B	Jiangsu	Majiabang	4500–3500	9.7	0.3	-20.0	0.2	19	[81]
Qinglongquan	43	B	Hubei	Qujialing/Shijiahe	3000–2200	9.0	1.1	-14.6	1.4	27	[82, 83]
Hemudu	44	B	Zhejiang	Hemudu	5000–4000	11.4	0.3	-18.2	2.2	4	[59]
Songze	45	B	Zhejiang	Songze	4000–3300	10.9	1.6	-19.9	0.4	2	[59]
Tashan	46	B	Zhejiang	Hemudu-Majiabang	3900–2200	6.9	4.6	-20.7	4.5	4	[84]

(Continued)

Table 2. (Continued)

Site	Site number	Region	Province	Cultural group	Estimated dates (cal. BC/AD)	Mean $\delta^{15}\text{N}$	SD $\delta^{15}\text{N}$	Mean $\delta^{13}\text{C}$	SD $\delta^{13}\text{C}$	n	Reference number
Tanshishan	47	B	Fujian	Tanshishan	3000–4300	10.8	1.6	-18.4	1.1	17	[46]
Liyudun	48	B	Guangdong		c. 5000	13.8	1.4	-17.0	1.3	2	[44]
Hupo	49	C	Qinghai	Banshan/Machang	2300–2000	7.6	0.3	-8.7	0.4	6	[85]
Mozuizi	50	C	Gansu	Machang	2400–2000	8.3	0.4	-7.2	0.4	14	[47]
Wuba	51	C	Gansu	Banshan/Machang	2500–1900	9.2	1.1	-7.5	1.3	55	[47]
Zongri	52	C	Qinghai	Zongri	3700–2300	8.3	0.5	-10.1	1.1	24	[86]
Dadiwan	53	C	Gansu	Yangshao	4550–2950	9.7	0.8	-9.8	3.0	6	[87]

Summary of published carbon and nitrogen isotope values measured in human bone collagen from sites dating to between 5000 and 2000 cal BC.

<https://doi.org/10.1371/journal.pone.0240930.t002>

deviations of the isotope data by time period, region, province and/or sex. We recognize that present-day provinces are artificial borders, however for the sake of simplicity, we compare isotopic data among provinces as a way to examine north to south and east to west geographic gradients. Although these data did not always conform to the assumptions of parametric statistics (specifically, residuals were not always normally distributed and groups did not necessarily have equal variance), we nonetheless chose to use ANOVA with posthoc Tukey's test for multigroup comparisons and were cautious about rejecting the null-hypothesis when p -values were close to 0.05. In the case of highly heteroscedastic groups, we used the more conservative Welch's ANOVA for multi-group comparisons (see [S3A–S3D Table](#) and [S4A–S4H Table](#) for results).

Mixing model

To estimate the proportional contributions of potential plant and animal food resources to past human diets at Xinglonggou—one of the oldest sites at which humans were using millet as a staple food—we used the Bayesian stable isotope mixing model MixSIAR [27, 28] following the best practices for stable isotope mixing models outlined by Phillips et al. [29]. We grouped dietary items into ecologically relevant isotopically distinct source groups by assessing whether sources had significantly different means using MANOVA followed by Tukey tests; dietary items that were isotopically indistinct ($p > 0.05$) were grouped and averaged over all of the samples. We accounted for concentration dependence by including the digestible [C] and [N] of the potential dietary resources, which we calculated from data available in the United States Department of Agriculture (USDA) Nutrient Database following Koch and Phillips [30]. To account for human diet-to-collagen isotope discrimination, we used a nitrogen isotope dietary discrimination factor of 3.5 ± 1 ‰ and a carbon isotope discrimination factor of 5 ± 1 ‰ [12, 31]. We initially tried nitrogen isotope discrimination factors of between 4.6–6 ‰ [32], but found that these values placed the human collagen samples well outside the dietary mixing space. To evaluate the sensitivity of the model to the nitrogen isotope discrimination factor, we ran the model using several discrimination factors ([S5 Table](#)); the estimated mean proportional contribution of C_4 plants to human diet varies by just 4% among the models. We conducted Markov Chain Monte Carlo (MCMC) sampling within MixSIAR using the “very long” chain length, which includes running three replicate chains, each with 1,000,000 draws, a burn-in of 500,000, and a thinning rate of 500. We used Gelman-Rubin diagnostics to confirm model convergence [33]. Although the relative abundance of various dietary resources

Table 3. Published human C and N isotope values from between 2000 cal BC and 212 cal AD.

Site	Site number	Region	Province	Cultural group	Estimated dates (cal. BC/AD)	Mean $\delta^{15}\text{N}$	SD $\delta^{15}\text{N}$	Mean $\delta^{13}\text{C}$	SD $\delta^{13}\text{C}$	n	Reference number
Beiqian	54	A	Shandong	Zhou Dynasty	1046–256	10.5	0.5	-9.2	0.8	4	[44]
Qianzhangda	55	A	Shandong	Shang/Western Zhou	1200–771	10.0	1.2	-8.9	1.4	49	[88]
Xiyasi	56	A	Shandong	Eastern Zhou	770–221	8.1	0.9	-11.9	2.2	30	[50]
Liulihe	57	A	Hebei	Western Zhou	1046–771	-	-	-8.2	1.4	19	[59]
Nancheng	58	A	Hebei	Proto Shang	2000–1600	9.4	0.0	-6.9	0.1	75	[89]
Neiyangyuan	59	A	Shanxi	Spring and Autumn	770–476	9.6	1.0	-8.6	1.6	23	[90]
Niedian	60	A	Shanxi	Xia	1900–1500	10.5	0.7	-7.1	0.3	60	[91]
Xiaonanzhuang	61	A	Shanxi	Eastern Zhou	770–221	10.5	0.9	-8.0	0.4	16	[92]
Xiaoshuangqiao	62	A	Shanxi	Shang	1600–1046	8.5	1.6	-10.0	1.7	10	[93]
Anyang	63	A	Henan	Shang	1400–1046	-	-	-8.2	2.5	39	[59]
Changxinyua	64	A	Henan	Eastern Zhou	770–221	7.7	1.0	-10.3	1.4	15	[50]
Erlitou	65	A	Henan	Erlitou	1900–1500	11.9	4.2	-9.4	2.1	31	[59, 69]
Handeng	66	A	Henan	Proto Shang	1750–1600	9.3	-	-6.7	-	1	[94]
Liuzhuang	67	A	Henan	Proto Shang	2000–1600	9.7	1.5	-8.2	1.9	21	[95]
Xiaomintun	68	A	Henan	Shang	1400–1046	9.5	0.6	-11.5	2.7	4	[96]
Xinzhai	69	A	Henan	Erlitou	1900–1500	9.0	1.0	-9.6	1.4	8	[97]
Xinzheng City	70	A	Henan	Eastern Zhou	770–221	8.8	0.8	-11.0	1.6	75	[80]
Xuecun	71	A	Henan	Han	202BC–220AD	10.6	1.3	-13.7	1.2	53	[80]
Yanshi	72	A	Henan	Shang	1600–1046	-	-	-7.6	0.8	3	[59]
Yinxu	73	A	Henan	Shang	1250–1046	9.1	1.2	-8.5	1.0	130	[98–100, 104]
Fenggeling	74	A	Shaanxi		500–300	9.1	0.4	-9.2	0.5	4	[65]
Guandao	75	A	Shaanxi	Han	202BC–220AD	10.4	0.3	-10.7	0.8	5	[101]
Guangming	76	A	Shaanxi	Western Han	202BC–8AD	11.0	0.8	-9.8	0.9	7	[101]
Jianhe	77	A	Shaanxi	Warring States	476–221	8.7	0.5	-9.2	0.7	14	[102]
Jichang	78	A	Shaanxi	Eastern Han	8–220AD	9.0	0.9	-12.0	1.2	30	[101]
Lintong	79	A	Shaanxi		300BC–700AD	9.4	1.3	-13.8	3.8	3	[65]
Muzhuzhuliang	80	A	Shaanxi	Longshan/Xia	1900–1700	8.8	0.6	-8.2	1.5	8	[103]
Shimao	81	A	Shaanxi	Longshan/Xia	2100–1600	6.9	0.9	-8.4	0.1	4	[65]
Shigushan	82	A	Shaanxi	Western Zhou	1200–900	9.4		-9.8		1	[104]
Sunjianantou	83	A	Shaanxi	Eastern Zhou	770–221	8.5	1.0	-10.8	1.3	25	[105]
Xinhua	84	A	Shaanxi		2000–1700	8.2	-	-8.7	-	1	[65]
Zhanguo	85	A	Shaanxi		450–350	8.8	-	-14.8	-	1	[65]
Zhouyuan	86	A	Shaanxi	Western Zhou	1200–900	9.8	1.3	-10.9	2.3	9	[104]
Dabaoshan	87	A	In. Mongolia	Warring States	410–180	9.6	0.9	-9.0	1.4	40	[106]
Dashanqian	88	A	In. Mongolia	Upper Xiajiadian	900–200	9.3	0.6	-7.0	0.4	9	[107]
Huhewusu	89	A	In. Mongolia	Western Han	202BC–8AD	9.1	0.6	-9.1	0.7	5	[108]
Jinggouzi	90	A	In. Mongolia	Eastern Zhou	770–221	9.8	0.6	-12.4	0.7	10	[109]
Nalintaohai	91	A	In. Mongolia	Western Han	202BC–8AD	13.3	1.2	-10.0	0.8	7	[110]
Tuoba Xianbei	92	A	In. Mongolia	Tuoba Xianbei	100BC–557AD	10.4	1.3	-11.4	2.8	65	[111]
Xindianzi	93	A	In. Mongolia	Eastern Zhou	770–221	10.3	0.8	-11.5	0.9	20	[112]
Xinglongwa	94	A	In. Mongolia	Lower Xiajiadian	1500–1300	-	-	-4.2	0.9	2	[59]

(Continued)

Table 3. (Continued)

Site	Site number	Region	Province	Cultural group	Estimated dates (cal. BC/AD)	Mean $\delta^{15}\text{N}$	SD $\delta^{15}\text{N}$	Mean $\delta^{13}\text{C}$	SD $\delta^{13}\text{C}$	n	Reference number
Xinglongwa III	95	A	In. Mongolia	Lower Xiajiadian	1500–1300	9.8	0.9	-7.0	0.6	9	[43, 59]
Zhukaigou	96	A	In. Mongolia	Zhukaigou	2200–1600	9.0	1.1	-8.2	0.4	2	[65]
Buziping	97	C	Gansu	Qijia	2100–1700	8.1	-	-7.3	-	1	[48]
Buzishan	98	C	Gansu	Qijia	2100–1700	8.3	-	-7.3	-	1	[48]
Ganguai	99	C	Gansu	Siba	1400–900	11.6	0.9	-15.3	1.5	30	[83]
Huoshagou	100	C	Gansu	Siba	2000–1300	12.0	1.3	-12.0	1.9	30	[83]
Huoshiliang	101	C	Gansu		2135–1690	8.0	2.6	-8.8	0.1	2	[58]
Lianhuatai	102	C	Gansu	Xindian	1500–1000	8.6	0.3	-10.0	0.3	6	[104]
Lixian	103	C	Gansu	Spring and Autumn	750–500	8.8	0.3	-13.1	4.1	3	[62]
Mogou	104	C	Gansu	Qijia/Siwa	1800–1100	10.2	1.2	-13.9	1.5	85	[83, 87]
Mozuizi	105	C	Gansu	Han	202BC–220AD	10.5	0.8	-15.7	1.4	6	[83]
Qijiaping	106	C	Gansu	Qijia	1500–1200	9.8	0.9	-8.9	1.1	42	[111]
Xiahaishi	107	C	Gansu	Qijia	2200–1900	8.6	1.0	-7.5	0.3	13	[48, 104]
Xichengyi	108	C	Gansu	Siba	2000–1000	11.7	2.1	-9.0	0.6	4	[113]
Xishan	109	C	Gansu	Spring and Autumn	770–403	7.9	1.8	-13.4	4.0	41	[85]
Zhanqi	110	C	Gansu	Siwa	1100–900	10.4	0.6	-15.5	1.0	31	[47, 104]
Lajia	111	C	Qinghai	Qijia	2000–1200	10.0	0.2	-7.9	0.4	4	[114]
Lajigai	112	C	Qinghai	Kayue	1400–1000	9.0	0.5	-14.9	1.8	5	[85]
Sanheyi	113	C	Qinghai	Qijia	2000–1800	8.1	1.5	-9.1	0.5	5	[85]
Shangsunjia	114	C	Qinghai	Kayue	1500–600	9.8	1.4	-16.2	1.3	21	[59]
Donghuigou	115	C	Xinjiang	Hongshankou	900–0	13.3	0.6	-18.4	0.4	13	[115]
Duogang	116	C	Xinjiang	Qunbake	900–500	12.6	0.6	-14.5	1.0	39	[116]
Gumugou	117	C	Xinjiang	Xiaohe	c. 1800	14.6	0.6	-18.2	0.2	10	[117, 118]
Heigouliang	118	C	Xinjiang	Western Han	500BC–8AD	12.5	0.6	-18.5	0.4	36	[119, 120]
Qiongkeke	119	C	Xinjiang	Early Iron Age	500–202	12.7	0.4	-16.2	0.2	8	[121]
Kelasu	120	C	Xinjiang	Early Iron Age/ Han	500BC–220AD	11.8	0.6	-16.6	0.4	7	[122]
Tianshanbeilu	121	C	Xinjiang	Tianshanbeilu	2000–1300	14.7	0.9	-15.4	1.3	124	[123, 124]
Xiabandi	122	C	Xinjiang	Andronovo	1500–600	12.3	1.0	-18.2	0.8	27	[125]
Yanbulake	123	C	Xinjiang	Early Iron Age	1000–500	-	-	-14.6	1.7	2	[59]
Yanghai	124	C	Xinjiang		1200–100	12.1	1.0	-16.3	1.1	22	[126]
Shenmingpu	125	B	Henan	Warring States/ Han	475BC–220AD	8.5	1.1	-14.6	2.2	32	[94]
Boyangcheng	126	B	Anhui	Spring and Autumn	1122–771	10.9	1.0	-18.8	1.5	38	[127]
Jinlianshan	127	B	Yunnan		700	9.8	0.9	-18.8	0.4	9	[128]
Shilinggang	128	B	Yunnan		850–250	10.0	1.0	-18.8	0.7	16	[129]

Summary of published carbon and nitrogen isotope values measured in human bone collagen from sites dating to between 2000 cal BC and 212 cal AD.

<https://doi.org/10.1371/journal.pone.0240930.t003>

found at the archaeological sites could arguably be used to construct informative priors, we chose to use uninformative priors (i.e., flat) for past human diet because of the potential for differences in preservation and/or sampling effort between floral and faunal material.

Kellner and Schoeninger's approach

Partitioning the relative contributions of plant and animal resources to human diet is difficult to accomplish using stable isotope values of bulk collagen alone because collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values reflect both dietary protein and dietary non-protein disproportionately (approximately 60% of the carbon atoms in collagen come from dietary protein [10, 34–38]). One approach to addressing this issue is to use $\delta^{13}\text{C}$ values in collagen and apatite from the same individual to model the regression lines of energy and protein sources, as collagen and bioapatite reflect dietary protein and the whole diet disproportionately [39]. Using the limited available data, we additionally followed [39] approach of plotting collagen $\delta^{13}\text{C}$ against apatite $\delta^{13}\text{C}$ with their modern-calibrated C_3 and C_4 protein regression lines [40, 41]. We identified three populations between 5000–2000 cal BC to be included in the analysis: Jiangzhai, Shijia and Banpo [42]. We also included Jiahu, a site that predates 5000 cal BC.

Results

Mapping staple cuisines in prehistoric China

We considered temporal and spatial patterns by organizing the results in three successive periods: 6000–5000 cal BC, 5000–2000 cal BC, and post 2000 cal BC (Liu et al. 2019), and within the geographic framework of the three regions described above: the broad Loess Plateau, the Yangtze-Huai Region, and the Continental Interior (Fig 1).

6000–5000 cal BC. Carbon and nitrogen isotope values measured in human bones are reported from five sites dating to the period between 6000–5000 cal BC (Fig 2A–2C). With the exception of Jiahu from the Yangtze-Huai Region, the sites are located in the Loess Plateau. Carbon isotope ratios from Jiahu are consistent with a predominantly- C_3 diet (mean $\delta^{13}\text{C} < -17\text{‰}$). In the Loess Plateau, human values from Xiaojingshan, Baijia and Beiliu are consistent with a mixed C_3 - C_4 diet ($\delta^{13}\text{C}$ values from -17 to -12‰), while at Xinglonggou and Xinglongwa, people have carbon isotope values indicative of a C_4 -plant dominated diet (mean $\delta^{13}\text{C} > -12\text{‰}$). The two regions have significantly different $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values ($\delta^{13}\text{C}$: $F_{1,68} = 93.4$, $p = 2.16 \text{ e-}14$, $\delta^{15}\text{N}$: $F_{1,66} = 4.8$, $p = 0.0319$).

Xinglonggou I (c. 6000 cal BC) provides a unique case study, as there are additionally data available from a range of both plant and animal dietary sources. At Xinglonggou, $\delta^{13}\text{C}$ values measured in human bone collagen are consistent with a C_4 diet with relatively high $\delta^{15}\text{N}$ values (Fig 3B; mean $\delta^{13}\text{C} = -9.9 \pm 1.1 \text{‰}$; $\delta^{15}\text{N} = 9.8 \pm 0.8 \text{‰}$, $n = 32$) [43]. The majority of animals (except dogs) from the same site demonstrate consistency with a C_3 diet and relatively low nitrogen isotope values (Fig 3A and 3B; mean $\delta^{13}\text{C} = -19.0 \pm 2.4 \text{‰}$; $\delta^{15}\text{N} = 5 \pm 1.4 \text{‰}$, $n = 50$). Carbon isotope ratios in humans seemingly suggest that humans directly consumed millet as a staple food, perhaps on a daily basis. Nitrogen isotope ratios on the other hand, suggest that the animal protein consumption at Xinglonggou was also significant, with a human-animal collagen offset of about 5 ‰ in $\delta^{15}\text{N}$ values.

To further explore these localized dietary patterns observed in bulk collagen data, we used an isotope mixing model to quantify the importance of C_4 plants to human diets at this site. The results suggest that the proportional contribution of C_4 plants (likely millet) to human diet at Xinglonggou was between 52–62% (95% CI; Fig 3C, S5 Table). Herbivores formed the second most important human dietary item, accounting for approximately 33–46%. These results confirm that humans in the Xinglonggou community relied on C_4 plants as a staple food. Nonetheless, when dietary reconstruction is based on bulk collagen isotopic determinations, informative variation at the molecular level is masked. Future research at the compound specific level that separates essential and non-essential amino acid isotope values could be




Fig 2. Box- and scatterplots of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values measured in human bone collagen. Data are from sites occupied pre-5000 cal BC (a-c), 5000–2000 cal BC (d-f), and post-2000 cal BC (g-i). Regions are differentiated by color and provinces by shape. Boxplots illustrate minimum, first quartile, median, third quartile, and maximum; means are depicted as hollow black diamonds and outliers as black dots. See Tables 1–3 for data citations and S1 Table for original data.

<https://doi.org/10.1371/journal.pone.0240930.g002>

undertaken to confirm or refute the validity of these interpretations derived from bulk collagen isotope data.

5000–2000 cal BC. Between 5000 and 2000 cal BC, isotope data from the Loess Plateau and the Yangtze-Huai Region reveal a more pronounced north-south distinction in human diets. Limited data are available from the Continental Interior. Humans from the Yangtze-Huai Region preserve isotopic signatures consistent with C_3 -dominated diets, while humans from the Loess Plateau present isotopic values suggesting they consumed a varying degree of C_4 plant foods (Fig 2D–2F). There is a statistically significant difference in the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values from these two regions (Welch’s ANOVA, $\delta^{13}\text{C}$: $F_{1,586} = 291.1, p < 2.2e-16$; $\delta^{15}\text{N}$: $F_{1,569} = 20.9, p = 5.785e-06$). Twenty-seven out of thirty populations from the Loess Plateau show significant consumption of C_4 plants ($\delta^{13}\text{C} > -12\text{‰}$, see Table 3). High $\delta^{13}\text{C}$ values can also be




Fig 3. Xinglonggou mixing model results. (a) Animal bone collagen and plant $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values from Xinglonggou; standard ellipse areas are depicted as ovals (C₃ plant—green, omnivore—pink, C₄ plant—orange, and herbivore—purple). (b) $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for grouped dietary sources (mean \pm 1 std dev) plotted with trophic corrected $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values from human bone collagen from Xinglonggou. The dashed line and the gray area illustrate the minimum and maximum convex hulls for the dietary mixing space, respectively. And (c) median (lines in center of boxes = median, box boundaries = 50% CI, error bars = 50% CI) proportional contributions of each dietary resource to humans from Xinglonggou.

<https://doi.org/10.1371/journal.pone.0240930.g003>

caused by significant consumption of marine resources, making it difficult to distinguish between C₄ and marine dietary inputs for coastal sites as in Shandong and the Lower Yangtze, where marine resources were abundant in the archaeological record [44]. The three Loess Plateau sites that do not exhibit dominant C₄ consumption at this time are all located in more southerly provinces that border on the Yangtze-Huai Region (Shandong and Shaanxi), suggesting that there is probably northward expansion of rice cultivation at this time (Fig 4A). In the Yangtze-Huai Region, seven out of thirteen populations have isotope values consistent with a predominantly-C₃ diet ($\delta^{13}\text{C} < -17\text{‰}$, see Table 3). The other six populations are consistent with a predominantly-C₄ diet. The latter group comes from the southern Henan and Hubei provinces, and likely reflects the southern expansion of millet cultivation in this period. Several individuals with extremely high $\delta^{15}\text{N}$ values in the Yangtze-Huai region (Fig 2F) come from coastal sites at which marine resources were likely being consumed [45, 46].

Some interesting patterns emerged in the collagen versus apatite $\delta^{13}\text{C}$ plot, using Kellner and Schoeninger's approach as described in the methods (Fig 5). At Jiahu, humans plot along the C₃ protein line, but their position on the y-axis ($\sim -10\text{‰}$) suggests that their energy comes from a mixture of C₃ and C₄ resources. Humans from Jiangzhai and Shijia, on the other hand, plot more closely to the C₄ protein line and their position along the y-axis, with apatite $\delta^{13}\text{C}$ values $> -5\text{‰}$, suggests their dietary energy is derived primarily from C₄ energy sources. Both of these sites are located on the Loess Plateau and these results help to clarify that some humans from this time period and region were likely consuming fully C₄ diets. The individual from Banpo, another Loess Plateau site, tells a slightly different story because they fall between the C₃ and C₄ protein lines, suggesting a mixed protein diet; their apatite $\delta^{13}\text{C}$ value is similarly suggestive of mixed C₃ and C₄ energy sources. Although this method is not quantitative, it nonetheless allows for energy and protein resources to be evaluated separately, allowing for a deeper understanding of past human diet than bulk collagen isotope values provide. Indeed, these data suggest that later in the period of 5000–2000 cal BC, some humans on the Loess Plateau were consuming millet directly as well as animals provisioned with millet (Jiangzhai and Shijia).

2000 cal BC– 220 cal AD. Between 2000 cal BC and 220 cal AD, China's staple food system experienced a major shift resulting from the introduction of wheat and barley (both are C₃ plants) [47–49]. The compiled isotopic data reflect distinct dietary choices between the prehistoric communities in the Loess Plateau and the Continental Interior (Fig 2G–2I). In the Loess




Fig 4. Isotope data by province, 5000–2000 cal BC. Boxplots of human bone collagen $\delta^{13}\text{C}$ (a) and $\delta^{15}\text{N}$ values (b) by province from sites that date to between 5000–2000 cal BC arranged from north (left) to south (right). Regions are differentiated by color: Loess Plateau (red) and Yangtze-Huai (gray). Provinces sharing a letter are not significantly different (Tukey's HSD). Statistics are summarized in [S3A and S3B Table](#).

<https://doi.org/10.1371/journal.pone.0240930.g004>

Plateau, other than a few exceptional individuals from Henan Province, humans show isotopic signatures consistent with predominantly- C_4 or mixed C_3 - C_4 consumption. Conversely, humans from the Continental Interior exhibit a broader spectrum of dietary habits including predominantly- C_3 , mixed C_3 - C_4 , and predominantly- C_4 diets. The two regions show statistically significant differences in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values ($\delta^{13}\text{C}$: Welch's ANOVA, $F_{1,1548} = 1113.8$, $p < 2.2e-16$; $\delta^{15}\text{N}$: Welch's ANOVA, $F_{1,1454} = 335.13$, $p < 2.2e-16$). Human data from 39 out of 43 sites from the Loess Plateau suggest that millet consumption was very significant ($\delta^{13}\text{C} > -12\text{‰}$), while human data from 19 out of 28 sites from the Continental Interior are consistent with C_3 or C_3 - C_4 mixed diets ($\delta^{13}\text{C} < -12\text{‰}$). The significantly different $\delta^{15}\text{N}$ values between the two regions could be the result of a combination of several factors, including variable animal protein input, differences in crop $\delta^{15}\text{N}$ values caused by variable soil ^{15}N enrichment, or aridity in the Continental Interior. The earlier north to south divide in staple crop use is accompanied by a new divide between the east and the west (see [Fig 6](#)).

Gendered consumption

We next consider the differences in staple consumption between males and females between 5000 and 2000 cal BC. In the Loess Plateau, females and males do not have significantly different $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values ([Fig 7A and 7B](#)). In the Yangtze-Huai Region, no significant




Fig 5. Kellner and Schoeninger’s approach. $\delta^{13}C_{\text{apatite}}$ and $\delta^{13}C_{\text{collagen}}$ measured in archaeological humans from one site dating to >5000 cal BC in the Yangtze-Huai region (gray, Jiahu) and three sites from the Loess Plateau dating to between 5000 and 2000 cal BC (red). Humans plotting along the C₃ regression line are interpreted to consume primarily C₃ protein, while those along the C₄ regression line consume C₄ protein. The position on the line along the y-axis is further indicative of the energy source, with low apatite $\delta^{13}C$ values corresponding with C₃ energy sources and high apatite $\delta^{13}C$ values corresponding with C₄ energy sources.

<https://doi.org/10.1371/journal.pone.0240930.g005>




Fig 6. Isotope data by province, post-2000 cal BC. Boxplots of human bone collagen $\delta^{13}C$ (a) and $\delta^{15}N$ values (b) by province from sites that date to post-2000 cal BC arranged from east (left) to west (right). Regions are differentiated by color: Loess Plateau (red) and Continental Interior (blue). Provinces sharing a letter do not have significantly different means (Tukey’s HSD). Statistics are summarized in S3C and S3D Table.

<https://doi.org/10.1371/journal.pone.0240930.g006>




Fig 7. Isotope data by sex. Boxplots and scatterplots of human bone collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values by sex from sites occupied between 5000–2000 cal BC (a-c), and post-2000 cal BC (d-f). Regions are differentiated by color and sex by shape and shade (female = lighter shaded, open triangles, male = darker shaded, open circles). Boxplot components are described in Fig 2. N-values listed at the base of the boxplots are divided by female (gray) and male (black). In panels a, b, d, and e, groups sharing a letter do not have significantly different means (Tukey's HSD). Statistics are summarized in S4A and S4B Table.

<https://doi.org/10.1371/journal.pone.0240930.g007>

difference in $\delta^{15}\text{N}$ is observed, but significant differences are observed in $\delta^{13}\text{C}$ values ($p = 0.0014$, Fig 7A and 7B, S4A and S4B Table), with males exhibiting higher carbon isotope values. This difference is primarily driven by regional variations within the Yangtze-Huai Region; when sexed individuals are compared within provinces, no significant differences are observed (Fig A in S1 File, S4C and S4D Table). It does not appear that social customs prohibited the consumption of C_4 -plants or animals consuming C_4 products by females in either the Loess Plateau or Yangtze-Huai Region during 5000–2000 cal BC, despite the fact that males consumed these foods to a higher degree than females in both regions.

After 2000 cal BC, males exhibited higher $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values than females in all three regions (Fig 7D and 7E), although the Loess Plateau is the only region where male and female $\delta^{13}\text{C}$ values differ significantly ($p = 0.01$, S4E and S4F Table). Because there is a risk of conflating gender differences with differences in social status, particularly when sample sizes are small, we focus our discussion on the Loess Plateau, where sample sizes are greatest. Lower $\delta^{13}\text{C}$ values in females from the Loess Plateau ($n = 218$) could indicate that females had greater access to newly introduced C_3 crops than males ($n = 293$). When gendered differences are considered at the provincial level, differences (although not significant) are evident in several provinces where males display higher access to C_4 resources and protein products (e.g., Shandong, Henan, Shaanxi, Inner Mongolia and Gansu; Fig B in S1 File, S4G and S4H Table). Only in Henan do males have significantly higher $\delta^{13}\text{C}$ values than females, which has been clearly documented at sites in the region [50].

Discussion

Our results suggest that both environmental and cultural-culinary conditions contributed significantly to the formation of staple cuisines in China. We shall next consider the observed isotopic patterns in two temporal and spatial dimensions. Before 2000 cal BC, the north-and-south dietary division will be considered in the context of regional variations of subsistence activities, which are partly driven by differences in environmental conditions. After 2000 cal BC, the introduction of crops originating from southwestern Asia resulted in an additional east-to-west gradient in the degree of incorporation of wheat and barley in human diets. We shall explore this pattern in relation to culinary traditions and emphasize the incompatibility of novel exotic grains with local culinary practice.

It is no exaggeration to say the millennium between 6000 and 5000 cal BC is crucial to understanding the origins of farming activities in East Asia [14, 16]. The north and south dietary divergence observed in this period is better understood as a difference in the spectrum of subsistence activities, rather than as separated peoples. In the Yangtze-Huai Region, human carbon isotope values are consistent with a C_3 -plant dominated diet, which likely consisted of C_3 resources identified in the archaeobotanical record (i.e. rice, fruits, tubers, nuts) [24]. In the Loess Plateau, however, humans relied on C_4 foods. Broomcorn and/or foxtail millet were documented at all four northern sites (or associated cultural sites) in high quantities [13, 51]. No other C_4 cereal has been identified in the plant macrofossil assemblages from this time period. There is microbotanical evidence for job's tears (a C_4 plant) at Xinglonggou [52], however, given the tropical adaptation of genus *Coix*, job's tears were unlikely to be cultivated on a large enough scale to become a staple cereal 7500 years ago. The tradition of consumption of C_4 crops as staple foods emerged in this period and was particularly pronounced among the Xinglonggou cultural communities. At Xinglonggou, we estimated the proportional contribution of C_4 plants to human diet to be greater than 50%, nearly two-times more significant than herbivores, the second most important dietary resource.

Above all, the distinct subsistence modes between north and south in Neolithic China are driven by regional environmental differences. The lower catchment of the Yangtze and Huai Rivers was an intricate deltaic wetland crisscrossed by hundreds of distributaries, merging and diverging with seasonal flooding. People in this region relied overwhelmingly on wetland resources, including rice—an aquatic plant—for their subsistence. In contrast, landscapes in the north form a single relatively uniform semi-arid zone across the Loess Plateau. From early on, millet cultivation became the key component of agrarian based subsistence in the Loess Plateau. Within this perspective, the broad spectrum of subsistence activities in the Yangtze-Huai within an environmental mosaic consisting of swamps, marshes, fens and wetlands can be seen as the mirror image of unified agrarian practices based on millet grain in the northern Loess Plateau. That is, both of these highly sustainable systems in the north and south took advantage of the subsistence options their landscape setting provided. And this arrangement seems to have persisted for another 3000 years (5000–2000 cal BC). The regional difference in dietary tradition between north and south, along with the variation within each region, challenges the conventional “stages” of shifting modes of subsistence—hunting, foraging, pastoralism, and farming—in an evolutionary framework. Both historical and archaeological evidence shows that peoples moved fairly readily between distinctive modes of subsistence and the same people might have practiced more than one subsistence mode in a single lifetime [53, 54]. In China as elsewhere, it seems early peoples combined subsistence modes in a number of innovative hybrids that co-existed over thousands of years. The north-south dietary distinction in China resonates with the conceptual distinction in southwest Asia between the northern “Hilly Flanks” and the southern Mesopotamian alluvium highlighted by James Scott [54].

The rapid adoption of wheat and barley as staple foods in the Continental Interior by 2000 cal BC contrasts the very gradual pace of the adoption of these western crops in the Loess Plateau. In a recent review focused on northern China, the authors noted that the shift from a C₄-dominated diet to a mixed C₃-C₄ diet at this time was concurrent with “Holocene Event 3” at 4200 BP (2,250 BC) [55]. A global aridification event may well be part of the explanation of the readiness of communities in the Continental Interior to accept wheat and barley as new staples. Nonetheless, the question remains—what delayed the adoption of wheat and barley in the Loess Plateau? As discussed elsewhere, one plausible social explanation is that in the early stages of their adoption in the Loess Plateau, these crops were exclusively used by the few—such as elites, ritual specialists or others—rather than the many [56]. But this is not the only explanation.

An alternative interpretation lies in the deep-seated East and West culinary distinction. As established, boiling and steaming of grains and other foods appear to have been and remained the predominant East Asian methods for preparing foods. By contrast, cereals in southwestern and Central Asia such as wheat and barley were processed for a flour-focused food system. Such an East-West culinary distinction can be traced back to the pre-agricultural Palaeolithic [57]. These culinary preferences had consequences for the selection of grain quality and features, with gluten protein being the target of selection in west Eurasia for making bread, and starch properties being selected in East Asia for the function of boiling-steaming. It has been hypothesized that the western boundary of the boiling-steaming culinary tradition appears to correspond approximately to the geographic range of the summer monsoon [57]. In other words, the people of the Loess Plateau and Continental Interior each belonged to two distinct culinary systems: the boiling-and-steaming cultures in the East and grinding-and-baking cultures in the West. The gradual adoption of western grains (wheat and barley) and the isotopic evidence associated with it could be understood in this context. The dispersal of crops into new areas was not necessarily accompanied by the spread of the culinary traditions from their regions of origin. Novel grains could instead be incorporated into existing local practices of food processing. In the case of wheat, it has been illustrated this incorporation may have exerted selection on the crops for phenotypic traits adapted to the eastern cooking traditions [56]. In southeast Asia, the preference for cultivation of cereals that show within-species variation for stickiness of the cooked grain are typified by the eastern boiling-and-steaming cultures [58]. In both cases, it is plausible that the novel grains from the West (i.e. wheat and barley) might be initially “rejected” as a staple grain because of their incompatibility with local culinary practice, and this is consistent with the isotopic results showing a significant delay in human consumption of wheat and barley as a staple food in the Loess Plateau. Within the context of symbolism and social use of food [1], culinary traditions are often related to kinship and family structure, and that was the case in southeast Asia with the sticky food culture [58]. In the post-2000 BC Loess Plateau, newly introduced staple cereals from the West were consumed by females to a greater degree than males. This hints at the gender roles in the context of social status of grains and food processing with the female sector of the society being the primary agent of the process, pioneering innovations in culinary practice.

Conclusion

Modern Chinese cuisine formed over thousands of years through the development of diverse regional subsistence systems and cuisines, which were further influenced by food traditions from other parts of the world. Our results help to illustrate the ways in which both environment and culture contributed to shaping the Chinese staple food system over the past 8000 years. A distinct north versus south divide in Chinese ancient staple cuisines was already evident isotopically between 6000–5000 cal BC and became more pronounced between 5000–

2000 cal BC. We infer that this pattern is better understood as a difference in the spectrum of subsistence activities, which was partly driven by environmental differences between the Loess Plateau and the Yangtze-Huai region. The introduction of wheat and barley from southwestern Asia after 2000 cal BC resulted in the development of an additional east to west gradient in the degree of incorporation of the different staple products into human diets. We argue the regional differences in dietary tradition between and within the three broad regions throughout the Neolithic and the Bronze Age could not be understood in the conventional “stages” of shifting modes of subsistence: hunting-foraging-pastoralism-farming. Instead the same people might have practiced more than one subsistence mode and combined them in a number of innovative hybrids that co-existed over thousands of years. The rapid adoption of wheat and barley as staple foods in the Continental Interior by 2000 cal BC contrasts the very gradual pace of the adoption of these western crops in the Loess Plateau. Apart from the possible environmental and social drivers, we explored a third explanation that these novel grains may have at first been ignored as a staple grain because of their incompatibility with local culinary practice; people of the Loess Plateau belonged to a boiling-and-steaming culture while those in the Continental Interior belonged to a grinding-and-baking culture into which wheat and barley were more readily folded. Finally, in some cases in the Loess Plateau, newly introduced staple cereals from the West were consumed by females to a greater extent than males, suggesting that the female sector of society may have pioneered the innovations in culinary practice.

Supporting information

S1 File. Supplementary material: Gender differences at the provincial level.

(DOCX)

S2 File. R scripts used in this study.

(DOCX)

S1 Table. Published carbon and nitrogen isotope data used in this study. Data are from archaeological human skeletal remains ($n = 2448$) from 128 sites across China.

(XLSX)

S2 Table. Summary of key archaeobotanical studies from selected sites cross China. See [2] for detailed site/assemblage information.

(XLSX)

S3 Table. A-D. ANOVA results by province. Results of comparisons of human $\delta^{13}\text{C}$ (Table A in S3 Table) values and $\delta^{15}\text{N}$ (Table B in S3 Table) by province in time period II (5000–2000 cal BC), and results of comparisons of human $\delta^{13}\text{C}$ (Table C in S3 Table) values and $\delta^{15}\text{N}$ (Table D in S3 Table) by province in time period III (2000 cal BC– 220 cal AD).

(XLSX)

S4 Table. A-H. ANOVA results by sex. Results of comparisons of male and female $\delta^{13}\text{C}$ (Table A in S4 Table) and $\delta^{15}\text{N}$ (Table B in S4 Table) values by region in time period II (5000–2000 cal BC); ANOVA comparisons of human $\delta^{13}\text{C}$ (Table C in S4 Table) and $\delta^{15}\text{N}$ (Table D in S4 Table) values by sex and province in time period II (5000–2000 cal BC); ANOVA comparisons of male and female $\delta^{13}\text{C}$ (Table E in S4 Table) and $\delta^{15}\text{N}$ (Table F in S4 Table) values by region in time period III (2000 cal BC– 220 cal AD); and ANOVA comparisons of human $\delta^{13}\text{C}$ (Table G in S4 Table) and $\delta^{15}\text{N}$ (Table H in S4 Table) values by sex and province in time period III (2000 cal BC– 220 cal AD).

(XLSX)

S5 Table. Mixing model results for Xinglonggou humans. Results from the model run highlighted in green are reported in the text. (XLSX)

Acknowledgments

We acknowledge David Redhouse at University of Cambridge for assistance to generate the map. We are also grateful to Professor Fiona Marshall and Dr Petra Vaiglova for their feedback on an earlier draft of the manuscript.

Author Contributions

Conceptualization: Xinyi Liu.

Data curation: Xinyi Liu, Rachel E. B. Reid.

Formal analysis: Xinyi Liu, Rachel E. B. Reid.

Funding acquisition: Xinyi Liu.

Investigation: Xinyi Liu.

Methodology: Xinyi Liu, Rachel E. B. Reid.

Software: Rachel E. B. Reid.

Visualization: Xinyi Liu, Rachel E. B. Reid.

Writing – original draft: Xinyi Liu, Rachel E. B. Reid.

Writing – review & editing: Xinyi Liu, Rachel E. B. Reid.

References

1. Douglas M. Standard Social Uses of Food: Introduction. In: Douglas M, editor. *Food in the Social Order: Studies of Food and Festivities in Three American Communities*. New York: Russell Sage Foundation; 1984. p. 1–39.
2. Liu X, Jones PJ, Motuzaite Matuzeviciute G, Hunt HV, Lister DL, An T, et al. From ecological opportunism to multi-cropping: mapping food globalisation in prehistory. *Quaternary Science Reviews*. 2019; 206(15):21–8.
3. Jones MK, Hunt HV, Lightfoot E, Lister D, Liu X, Motuzaite-Matuziviciute G. Food globalization in prehistory. *World Archaeology*. 2011; 43(4):665–75.
4. Bray F. *Science and Civilisation in China—Agriculture*. Needham J, editor. Cambridge: Cambridge University Press; 1984.
5. Alexandratos N, editor. *World Agriculture: towards 2030/50, interim report. An FAO perspective*. London/Rome: Earthscan/FAO; 2006.
6. Kearney J. Food consumption trends and drivers. *Philosophical Transactions of the Royal Society B*. 2010; 365:2793–807.
7. Han M. *Historical Agricultural Geography of China*. Beijing: Peking University Press; 2012.
8. Fuller D, Rowlands M. Ingestion and Food Technologies: Maintaining Differences over the Long-term in West, South and East Asia. In: Wilkinson TC, Sherratt S, Bennet J, editors. *Intervening Worlds: systemic interactions in Eurasia, 7th to 1st millennia BC*. Oxford: Oxbow Books; 2011.
9. Hu Y. Thirty-four years of stable isotopic analyses of ancient skeletons in China: An overview, progress and prospects. *Archaeometry*. 2018; 60(1):144–56.
10. van der Merwe NJ, Vogel JC. C-13 Content of human collagen as a measure of prehistoric diet in woodland North-America. *Nature*. 1978; 276(5690):815–6. ISI:A1978GA88900044. <https://doi.org/10.1038/276815a0> PMID: 364321
11. Ambrose SH, Norr L. Experimental evidence for the relationship of the carbon isotope ratios of whole diet and dietary protein to those of bone collagen and carbonate. In: Lambert JB, Grupe G, editors. *Prehistoric Human Bone: Archaeology at the Molecular Level*. Berlin: Springer-Verlag; 1993. p. 1–37.

12. Hedges REM, Reynard LM. Nitrogen isotopes and the trophic level of humans in archaeology. *Journal of Archaeological Science*. 2007; 34(8):1240–51.
13. Liu X, Fuller D, Jones MK. Early agriculture in China. In: Barker G, Goucher C, editors. *The Cambridge World History—Volume II: A world with agriculture, 12,000 BCE–500CE*. Cambridge: Cambridge University Press; 2015. p. 310–34.
14. Crawford GW. East Asian plant domestication. In: Stark MT, editor. *Archaeology of Asia*. Malden: Blackwell Publishing; 2006. p. 77–95.
15. Crawford G. Domestication and the origins of agriculture in China. In: Feinman G, Bekke D, Niziolek L, editors. *Visions Through the Ages*. Chicago: University of Chicago Press; 2018. p. 45–63.
16. Zhao Z. New archaeobotanic data for the study of the origins of agriculture in China. *Current Anthropology*. 2011; 52:S295–S304.
17. Liu X, Hunt HV, Jones MK. River valleys and foothills: changing archaeological perceptions of north China's earliest farms. *Antiquity*. 2009; 83(319):82–95.
18. Gross BL, Zhao Z. Archaeological and genetic insights into the origins of domesticated rice. *Proceedings of National Academy of Science of the United States of America*. 2013; 111(17):6190–7.
19. Ren X, Lemoine X, Mo D, Kidder TR, Guo Y, Qin Z, et al. Foothills and intermountain basins: Does China's Fertile Arc have a 'Hilly Flanks'? *Quaternary International*. 2016; 426(28):86–96.
20. Liu X, Lister DL, Zhao Z, Petrie CA, Zeng X, Jones PJ, et al. Journey to the East: diverse routes and variable flowering times for wheat and barley *en route* to prehistoric China. *PLOS ONE*. 2017; 12(11): e0209518.
21. Liu L, Bestel S, Shi J, Song Y, Chen X. Paleolithic human exploitation of plant foods during the last glacial maximum in North China. *Proceedings of National Academy of Science of the United States of America*. 2013; 110(14):5380–5.
22. Lee GA, Crawford GW, Liu L, Sasaki Y, Chen X. Archaeological soybean (*Glycine max*) in East Asia: Does size matter? *PLoS ONE*. 2011; 6(11):e26720. <https://doi.org/10.1371/journal.pone.0026720> PMID: 22073186
23. Zheng Y, Crawford G, Chen X. Archaeological evidence for peach (*Prunus persica*) cultivation and domestication in China. *PLoS ONE*. 2014; 9(9):e106595. <https://doi.org/10.1371/journal.pone.0106595> PMID: 25192436
24. Zhao Z. Flotation results from the Jiahu, Wuyang county, Henan. In: Zhao Z, editor. *Paleoethnobotany: Theories, Methods and Practice*. Beijing: Academy Press; 2010. p. 108–18.
25. Yang X, Fuller D, Huan X, Perry L, Li Q, Li Z, et al. Barnyard grasses were processed with rice around 10,000 years ago. *Scientific Reports*. 2015; 5: <https://doi.org/10.1038/srep16251>
26. R-Core-Team. R: A language and environment for statistical computing Vienna, Austria: R Foundation for Statistical Computing; 2017. Available from: <https://www.R-project.org>.
27. Stock BC, Semmens BX. MixSIAR GUI User Manual. Version 3.1.: <https://github.com/brianstock/MixSIAR>; 2016. <https://doi.org/10.5281/zenodo.1209993> p.
28. Parnell AC, Phillips DL, Bearhop S, Semmens BX, Ward EJ, Moore JW, et al. Bayesian stable isotope mixing models. *Environmetrics*. 2013; 24(6):387–99.
29. Phillips DL, Inger R, Bearhop S, Jackson AL, Jonathan WM, Andrew CP, et al. Best practices for use in stable isotope mixing models in food-web studies. *Canadian Journal of Zoology*. 2014; 92:823–35.
30. Koch PL, Phillips DL. Incorporating concentration dependence in stable isotope mixing models: a reply to Robbins, Hilderbrand and Farley (2002). *Oecologia*. 2002; 133(1):14–8. <https://doi.org/10.1007/s00442-002-0977-6> ISI:000178224300003. PMID: 24599364
31. Newsome SD, Phillips DL, Culleton BJ, Guilderson TP, Koch PL. Dietary reconstruction of an early to middle Holocene human population from the central California coast: insights from advanced stable isotope mixing models. *Journal of Archaeological Science*. 2004; 25:1051–62.
32. O'Connell TC, Kneale CJ, Tasevska N, Kuhnle GG. The diet-body offset in human nitrogen isotopic values: A controlled dietary study. *American Journal of Physical Anthropology*. 2012; 149(3):426–34. <https://doi.org/10.1002/ajpa.22140> PMID: 23042579
33. Gelman A, Rubin DB. Inference from iterative simulation using multiple sequences. *Statistical Science*. 1992; 7(4):457–72.
34. Corr LT, Sealy JC, Horton MC, Evershed RP. A novel marine dietary indicator utilising compound-specific bone collagen amino acid $\delta^{13}\text{C}$ values of ancient humans. *Journal of Archaeological Science*. 2005; 32:321–30.
35. Fogel ML, Tuross N. Extending the limits of paleodietary studies of humans with compound specific carbon isotope analysis of amino acids. *Journal of Archaeological Science*. 2003; 30(5):535–45. [https://doi.org/10.1016/s0305-4403\(02\)00199-1](https://doi.org/10.1016/s0305-4403(02)00199-1) ISI:000182001300003.

36. Jim S, Jones V, Ambrose SH, Evershed RP. Quantifying dietary macronutrient sources of carbon for bone collagen biosynthesis using natural abundance stable carbon isotope analysis. *British Journal of Nutrition*. 2006; 95:1055–62. <https://doi.org/10.1079/bjn20051685> PMID: [16768826](https://pubmed.ncbi.nlm.nih.gov/16768826/)
37. Jim S, Ambrose SH, Evershed RP. Stable carbon isotopic evidence for differences in the dietary origin of bone cholesterol, collagen and apatite: implications for their use in palaeodietary reconstruction. *Geochimica et Cosmochimica Acta*. 2004; 68:61–72.
38. Tieszen LL, Fagre T. Effect of diet quality and composition on the isotopic composition of respiratory CO₂, bone collagen, bioapatite and soft tissues. In: Lambert JB, Grupe G, editors. *Prehistoric Human Bone: Archaeology at the Molecular Level*. Berlin: Springer-Verlag; 1993. p. 121–55.
39. Kellner CM, Schoeninger MJ. A simple carbon isotope model for reconstructing prehistoric human diet. *American Journal of Physical Anthropology*. 2007; 133:1112–27. <https://doi.org/10.1002/ajpa.20618> PMID: [17530667](https://pubmed.ncbi.nlm.nih.gov/17530667/)
40. Froehle AW, Kellner CM, Schoeninger MJ. FOCUS: effect of diet and protein source on carbon stable isotope ratios in collagen: follow up to Warinner and Tuross. *Journal of Archaeological Science*. 2010; 37:2662–70.
41. Warinner C, Tuross N. Alkaline cooking and stable isotope tissue-diet spacing in swine: archaeological implications. *Journal of Archaeological Science*. 2009; 36:1690–7.
42. Pechenkina EA, Ambrose SH, Xiaolin M, Benfer JRA. Reconstructing northern Chinese Neolithic subsistence practices by isotopic analysis. *Journal of Archaeological Science*. 2005; 32(8):1176–89.
43. Liu X, Jones MK, Zhao Z, Liu G, O'Connell TC. The earliest evidence of millet as a staple crop: New light on Neolithic foodways in North China. *American Journal of Physical Anthropology*. 2012; 149(2):238–90.
44. Wang F, Fan R, Kang H, Jin G, Luan F, Fang H, et al. Reconstructing the food structure of ancient coastal inhabitants from Beiqian village: Stable isotopic analysis of fossil human bone. *Chinese Science Bulletin*. 2012; 57(17):2148–54.
45. Hu Y, Li F, Wang C, Michael R. Guangdong Zhanjiang Liyudun yizhi rengu de C/N wending tongweisu fenxi (Carbon and Nitrogen stable isotope analysis of the human bones from the Liyudun site, Zhanjiang, Guangdong: a preliminary exploration of the Neolithic human lifestyle in South China). *Renlei Xue Xuebao (Acta Anthropologica Sinica)*. 2010; 29:264–9.
46. Wu M, Ge W, Chen Z. Haiyang xing juluo xianmin de shiwu jiegou: Tanshishan yizhi xinshiqi shidai wanqi rengu de tan dan wending tongweisu fenxi (Diets of a late Neolithic maritime settlement: Carbon and nitrogen isotope analysis of human skeletons recovered from Tanshishan). *Acta Anthropologica Sinica*. 2016; 35(2):246–56.
47. Liu X, Lightfoot E, O'Connell TC, Wang H, Li S, Zhou L, et al. From necessity to choice: dietary revolutions in west China in the second millennium BC. *World Archaeology*. 2014; 46(5):661–80.
48. Ma MM, Dong GH, Lightfoot E, Wang H, Liu XY, Jia X, et al. Stable isotope analysis of human and faunal remains in the western loess plateau, approximately 2000 cal. BC. *Archaeometry*. 2014; 56(S1):237–55.
49. Ma M, Dong G, Liu X, Lightfoot E, Chen F, Wang H, et al. Stable isotope analysis of human and animal remains at the Qijiaping site in middle Gansu, China. *International Journal of Osteoarchaeology*. 2015; 25(6):923–34.
50. Dong Y, Morgan C, Chinenov Y, Zhou L, Fan W, Ma X, et al. Shifting diets and the rise of male-biased inequality on the central plains of China during Eastern Zhou. *Proceedings of National Academy of Science of the United States of America*. 2017; 114(5):932–7.
51. Crawford G, Chen X, Wang J. Houli culture rice from the Yuezhuang site, Jinan. *Dongfang Kaogu (East Asia Archaeology)*. 2006; 3:247–51.
52. Liu L, Duncan NA, Chen X, Liu G, Zhao H. Plant domestication, cultivation, and foraging by the first farmers in early Neolithic Northeast China: Evidence from microbotanical remains. *The Holocene*. 2015; 25(12):1965–78.
53. Zeder MA. The origins of agriculture in the Near East. *Current Anthropology*. 2011; 52(S4):S221–S35.
54. Scott JC. *Against the Grain: A Deep History of the Earliest States*. New Haven and London: Yale University Press; 2017.
55. Cheung C, Zhang H, Hepburn J, Yang DY, Richards MP. Stable isotope and dental caries data reveal abrupt changes in subsistence economy in ancient China in response to global climate change. *PLOS ONE*. 2019; Published: July 22, 2019; <https://doi.org/10.1371/journal.pone.0218943>.
56. Liu X, Lister DL, Zhao Z-Z, Staff RA, Jones PJ, Zhou L-P, et al. The virtues of small grain size: Potential pathways to a distinguishing feature of Asian wheats. *Quaternary International*. 2016; 426:107–9.

57. Fuller D, Rowland M. Ingestion and food technologies: maintaining differences over the long-term in West, South and East Asia. In: Wilkinson TC, editor. *Interweaving Worlds: Systematic Interactions in Eurasia, 7th to the 1st millennia BC*. 37–60. Oxford: Oxbow; 2011.
58. Fuller DQ, Castillo C. Diversification and cultural construction of a crop: the case of rice and waxy cereals in the food cultures of eastern Asia. In: Lee-Thorp JA, Katzenberg MA, editors. *The Oxford Handbook of the Archaeology of Diet*. Oxford: Oxford University Press; 2016. p. published online: <https://doi.org/10.1093/oxfordhb/9780199694013.013.8>
59. Zhang X, Wang J, Xian Z, Qiu S. Gu renlei shiwu jiegou yanjiu (Studies on ancient human diet). *Kaogu (Archaeology)*. 2003; 425:62–75.
60. Hu Y, Wang SG, Luan FS, Wang CS, Richards MP. Stable isotope analysis of humans from Xiaojingshan site: implications for understanding the origin of millet agriculture in China. *Journal of Archaeological Science*. 2008; 35(11):2960–5. <https://doi.org/10.1016/j.jas.2008.06.002> ISI:000259760600009.
61. Atahan P, Dodson J, Li X, Zhou X, Hu S, Chen L, et al. Early Neolithic diets at Baijia, Wei River valley, China: stable carbon and nitrogen isotope analysis of human and faunal remains. *Journal of Archaeological Science*. 2011; 38:2811–7.
62. Guo Y, Xia Y, Dong Y, Yu B, Fan Y, Wen F, et al. Beiliu yizhi rengu de wending tongweisu fenxi (Stable isotope analysis of human remains from Beiliu). *Kaogu yu Wenwu (Archaeology and Cultural Relics)*. 2016;(1):115–20.
63. Guo Y, Yu Y, Xia Y, Dong Y, Fan Y, Wen F, et al. Shiqian shiqi shehui xingzhi chutan—Yi Beiliu yizhi xianmin shiwu jiegou wending tongweisu fenxi weili (Investigating social structure in prehistory—A case of isotope analysis of the ancient diet at Beiliu). *Huaxia Kaogu (Huaxia Archaeology)*. 2017; (1):45–53.
64. Hu Y, Ambrose SH, Wang C. Stable isotopic analysis of human bones from Jiahu site, Henan, China: implications for the transition to agriculture. *Journal of Archaeological Science*. 2006; 33:1319–30.
65. Atahan P, Dodson J, Li X, Zhou X, Chen L, Barry L, et al. Temporal trends in millet consumption in northern China. *Journal of Archaeological Science*. 2014; 50:171–7.
66. Zhang Q, Eng JT, Wei J, Zhu H. Neimenggu Chayouqian Qi Miaoizigou yizhi xinshiqi shidai regu de wending tongweisu fenxi (Paleodietary studies using stable Carbon and Nitrogen isotopes from human bone: an example from the Miaoizigou site). *Renlei Xue Xuebao (Acta Anthropologica Sinica)*. 2010; 29:270–5.
67. Ling X, Chen Q, Xue X, Zhao C. Shanxi Ruicheng Qingliangsi modi chutu rengu de tongweisu fenxi (Stable isotope analysis on human bones from Qingliangsi site, Ruicheng, Shanxi Province). *Quaternary Sciences*. 2010; 30:415–21.
68. Cai L, Qiu S. Tan shisan ceding he gudai shipu yanjiu (Carbon-13 values and palaeodiet in China). *Kaogu (Archaeology)*. 1984; 205:945–55.
69. Zhang X, Qiu S, Bo G, Wang J, Zhong J. Erlitou yizhi Taosi yizhi bofen rengu tan shisan dan shiwu fenxi (Carbon and nitrogen isotope analysis to skeleton remains recovered from Erlitou and Taosi). In: Yuan J, editor. *Keji Kaogu—II (Archaeological Science—II)*. Beijing: Kexue Chubanshe (China Science Publishing House); 2007. p. 41–8.
70. Hu Y, He D, Dong Y, Wang C, Gao M, Lan Y. Shandong Tengzhou Xigongqiao yizhi rengu de wending tongweisu fenxi (Stable isotopic analysis on human remains from Xigongqiao, Tengzhou, Shandong). *Disiji Yanjiu (Quaternary Sciences)*. 2006; 25:561–7.
71. Chen X-L, Hu S-M, Hu Y-W, Wang W-L, Ma Y-Y, Lü P, et al. Raising practices of Neolithic livestock evidenced by stable isotope analysis in the Wei River valley, north China. *International Journal of Osteoarchaeology*. 2014; 26:42–52.
72. Guo Y, Hu Y, Gao Q, Wang C, Richard MP. Jiangzhai yizhi xianmin shipu fenxi (Stable Carbon and Nitrogen isotope evidence in human diets based on evidence from the Jiangzhai site, China). *Renlei Xue Xuebao (Acta Anthropologica Sinica)*. 2011; 30:149–57.
73. Hu Y, Hu S, Wang W, Wu X, Marshall FB, Chen X, et al. Earliest evidence for commensal processes of cat domestication. *Proc Natl Acad Sci USA*. 2014; 111(1):116–20. <https://doi.org/10.1073/pnas.1311439110> PMID: 24344279
74. Chen X, Guo X, Wang W, Hu S, Yang M, Wu Y, et al. The subsistence patterns of the Shengedaliang site (~4,000 yr BP) revealed by stable carbon and nitrogen isotopes in northern Shaanxi, China. *Science China*. 2017; 60(2):268–76.
75. Zhang X, Qiu S, Zhong J, Zhao X, Sun F, Cheng L, et al. Zhongyuan diqu jichu Yangshao wenhua shiqi kaogu yizhi de renlei shiwu zhuangkuang fenxi (Studies on diet of the ancient people of the Yangshao cultural sites in the Central Plains). *Renlei Xue Xuebao (Acta Anthropologica Sinica)*. 2010; 29:197–207.

76. Cheng P, Zhou W, Gong W, Zhu Y, Yang Y, Burr GS, et al. Paleodietary analysis of humans in Guanzhong basin, Shaanxi province since 8000 BP. *Radiocarbon*. 2017; 59(5):1435–46.
77. Chen X-L, Fang Y-M, Hu Y-W, Hou Y-F, Lü P, Yuan J, et al. Isotopic reconstruction of the late Longshan period (ca. 4200–3900 BP) dietary complexity before the onset of state-level societies at the Wadian site in the Ying River valley, central plains, China. *International Journal of Osteoarchaeology*. 2016; 26:808–17.
78. Shu T, Wei X, Wu X. Wuxiao yizhi rengu de tan dan wending tongweisu fenxi. *Huaxia Kaogu (Huaxia Archaeology)*. 2016;(1):48–55.
79. Fu Q, Jin S, Hu Y, Zhao M, Pan J, Wang C. Agricultural development and human diets in Gouwan site, Xichuan, Henan. *Chinese Science Bulletin*. 2010; 55(7):614–20.
80. Zhou L, Garvie-Lok SJ, Fan W, Chu X. Human diets during the social transition from territorial states to empire: Stable isotope analysis of human and animal remains from 770 BCE to 220 CE on the Central Plains of China. *Journal of Archaeological Science: Reports*. 2017; 11:211–23.
81. Hu Y, Wang G, Cui Y, Dong Y, Guan L, Wang C. Jiangsu Jintan Sanxingcun yizhi xianmin de shipu yanjiu (Palaeodietary study of Sanxingcun site, Jintan, Jiangsu). *Chinese Science Bulletin*. 2007; 52:660–4.
82. Guo Y, Hu Y, Zhu J, Zhou M, Wang C, Richards MP. Qinglongquan yizhi ren he zhu gu de C/N wending tongweisu fenxi (Carbon and Nitrogen isotopic analysis on human and pig bones from Qinglongquan site). *Zhongguo Kexue (Science of China)*. 2011; 41:52–60.
83. Guo Y, Fan Y, Hu Y, Zhu J, Richards MP. Diet transition or human migration in the Chinese Neolithic? Dietary and migration evidence from the stable isotope analysis of humans and animals from the Qinglongquan site, China. *International Journal of Osteoarchaeology*. 2015; 28(2):85–94.
84. Zhang G, Jiang L, Hu Y, Si Y, Lü P, Song G, et al. Zhejiang Tanshan yizhi ren he dongwu gu de C/N wending tongweisu fenxi (Carbon and nitrogen isotope analysis of human and animal skeletal remains from Tanshan, Zhejiang). *Huaxia Kaogu (Huaxia Archaeology)*. 2015;(2):138–46.
85. Ma M, Dong G, Jia X, Wang H, Cui Y, Chen F. Dietary shift after 3500 cal yr BP and its influencing factors in northwestern China: Evidence from stable isotopes. *Quaternary Science Reviews*. 2016; 145:57–70.
86. Cui Y, Hu Y, Chen H, Dong Y, Guan L, Weng Y, et al. Zongri yizhi rengu de wending tongweisu fenxi (Stable isotopic analysis on human bones from Zongri site). *Disiji Yanjiu (Quaternary Sciences)*. 2006; 26:604–11.
87. Barton L, Newsome SD, Chen F-H, Wang H, Guilderson TP, Bettinger RL. Agricultural origins and the isotopic identity of domestication in northern China. *Proceedings of National Academy of Sciences of the United States of America*. 2009; 106:5523–8.
88. Zhang X, Qiu S, Zhong J, Liang Z. Shandong Tengzhou shi Qianzhangda mudi chutu rengu de tan dan wending tongweisu fenxi (Carbon and Nitrogen isotopic analysis on human bones from Qianzhangda site, Tengzhou in Shandong Province). *Kaogu (Archaeology)*. 2012; 9(9):851–64.
89. Ma Y, Fuller BT, Wei D, Shi L, Zhang X, Hu Y, et al. Isotopic perspectives ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{34}\text{S}$) of diet, social complexity, and animal husbandry during the proto-Shang period (ca. 2000–1600 BC) of China. *American Journal of Physical Anthropology*. 2016; 160:433–45. <https://doi.org/10.1002/ajpa.22980> PMID: 26972928
90. Pei D, Hu Y, Wang Y, Zhang Q, Zhang G, Tian J, et al. Shanxi Xiangning Neiyangyuan yizhi xianmin shiwu jigou fenxi (Palaeodietary analysis of humans from the Neiyangyuan site of Xiangning, Shanxi Province). *Renlei Xue Xuebao (Acta Anthropologica Sinica)*. 2008; 27:379–84.
91. Wang Y, Nan P, Wang X, Wei D, Hu Y, Wang C. Xiangjin shehui dengji xianmin de shiwu jigou chayi —yi Shanxi Niedian yizhi weili (Dietary differences in humans with similar social hierarchies: Example from the Niedian site, Shanxi). *Acta Anthropologica Sinica*. 2014; 33(1):82–9.
92. Tang M, Wang X, Hou K, Hou L. Shanxi Jinzhong Xiaonanzhuang mudi rengu de C, N wending tongweisu: Shi lun xiaomai zai Shanxi de tuiguang (Carbon and nitrogen stable isotope of the human bones from the Xiaonanzhuang cemetery, Jinzhong, Shanxi: A preliminary study on the expansion of wheat in ancient Shanxi, China). *Acta Anthropologica Sinica*. 2018; 37(2):318–30.
93. Wang N, Li S, Li H, Hu Y, Song G. Oxygen isotope analysis of ancient bone collagen and its application in the study of human migration. *Science China Press*. 2015; 60(9):838–46.
94. Hou L, Wang N, Lv P, Hu Y, Song G, Wang C. Shenmingpu yizhi Zhanguo zhi lianghan xianmin shiwu jigou he nongye jingji de zhuanbian (Palaeodiet and the development of agricultural economy in Shenmingpu site). *Zhongguo Kexue (Science of China)*. 2012; 42:1018–25.
95. Hou L, Hu Y, Zhao X, Li S, Wei D, Hou Y, et al. Human subsistence strategy at Liuzhuang site, Henan, China during the proto-Shang culture (~2000–1600 BC) by stable isotopic analysis. *Journal of Archaeological Science*. 2013; 40:2344–51.

96. Si Y, Li Z. Xiaomintun yizhi wan Shang xianmin de dongwu danbai xiaofei ji xiangguan wenti chutan (Investigating). *Yindu Xuekan (Yindu Studies)*. 2017; 3:18–23.
97. Wu X, Xiao H, Wei C, Pan Y, Huang Y, Zhao C, et al. Henan Xinzhai yizhi ren, zhu shiwu jiegou yu nongye xingtai he jiazhu xunyang de wending tongweisu (Dietary reconstruction of human and pig uncovered from Xinzhai site, Henan Province: isotopic evidences of agriculture and pig domestication). *Keji Kaogu (Science for Archaeology)*. Beijing: Science Press; 2007. p. 49–58.
98. Cheung C, Jing Z, Tang J, Yue Z, Richards MP. Examining social and cultural differentiation in early Bronze Age China using stable isotope analysis and mortuary patterning of human remains at Xin'anzhuang, Yinxu. *Archaeological and Anthropological Science*. 2015; 9(5):799–816.
99. Cheung C, Jing Z, Tang J, Weston DA, Richards MP. Diets, social roles, and geographical origins of sacrificial victims at the royal cemetery at YinXu, Shang China: New evidence from stable carbon, nitrogen, and sulfur isotope analysis. *Journal of Anthropological Archaeology*. 2017; 48:28–45.
100. Zhang X, Xu G, He Y, Qiu S. Yinxu 54 hao mu chutu rengu de tan dan wending tongweisu fenxi (Carbon and nitrogen isotopic analysis of human skeletons recovered from tomb 54 at Yinxu). *Kaogu (Archaeology)*. 2017;(3):340–9.
101. Zhang G, Hu Y, Nehlich O, Yang W, Liu D, Song G, et al. Guanzhong lianghan xianmin shengye moshi ji yu beifang youmu minzu jian chayi de wending tongweisu fenxi (Exploring the subsistence differences between farming communities in Guanzhong and the northern pastoralists during the Han period—Stable isotopic analysis). *Huaxia Kaogu (Huaxia Archaeology)*. 2013;(3):131–41.
102. Ling X, Wang W, Chen Q, Sun L, Hu Y. Baoji Jianhe mudi chutu Zhanguo shiqi qin rengu de wending tongweisu fenxi (Stable isotopic analysis on human bones from the Warring States period of Jianhe cemetery in Baoji). *Archaeology and Cultural Relics*. 2010; 1(1):95–8.
103. Chen X, Guo X, Hu Y, Wang W, Wang C. Shaanxi Shenmu Muzhuzhuliang yizhi xianmin de shipu fenxi (Paleodietary analysis of Muzhuzhuliang, Shenmu, Shaanxi). *Kaogu yu Wenwu (Archaeology and Cultural Relics)*. 2015;(5):112–7.
104. Cheung C, Jing Z, Tang J, Richards MP. Social dynamics in early Bronze Age China: A multi-isotope approach. *Journal of Archaeological Science: Reports*. 2017; 16:90–101.
105. Ling X, Chen Q, Tian Y, Li Y, Zhao C, Hu Y. Shanxi Fengxiang Sunjianantou Qin mu chutu rengu zhong de C he N tongweisu fenxi (Carbon and Nitrogen stable isotopic analysis on human bones from the Qin tomb of Sunjianantou site, Fengxiang, Shaanxi Province). *Renlei Xue Xuebao (Acta Anthropologica Sinica)*. 2010; 29:54–61.
106. Zhang X, Zhang X, Sou M, Wei D, Hu Y. The influence of agriculture in the process of population integration and cultural interaction during the Eastern Zhou period in central-south Inner Mongolia: Carbon and nitrogen stable isotope analysis of human bones from the Dabaoshan cemetery, Helingeer county. *Science China*. 2018; 61(2):205–14.
107. Zhang C, Zhang Q, Peng S, Wang L, Zhu Y, Guo Z. Neimenggu Chifeng shi Dashanqian yizhi xiajiadian shangceng wenhua "jisi keng" chutu rengu wending tongweisu fenxi (Stable isotope analysis of human skeletons recovered from "sacrificial pit" at Dashanqian, a upper Xiajiadian cultural site, Chifeng, Inner Mongolia). *Kaogu yu Wenwu (Archaeology and Cultural Relics)*. 2015;(4):107–10.
108. Zhang Q, Guo L, Zhu H. Neimenggu Chayouqian Qi Huhewusu Handai mudi chutu rengu de wending tongweisu fenxi (Stable isotopic analysis on human bones from Huhewusu Han Dynasty tomb in Chayouqian Qi, Inner Mongolia). *Caoyuan Wenwu (Cultural Relics of Grassland)*. 2012; 2(2):99–101.
109. Zhang Q, Eng JT, Wang L, Ta L. Neimenggu Linxi xian Jinggouzi xiqu mudi rengu de wending tongweisu fenxi (Paleodietary studies using stable carbon isotopes from human bones: example from Jinggouzi cemetery, Inner Mongolia). *Bianjiang Kaogu Yanjiu (Research of China's Frontier Archaeology)*. 2008; 7:322–7.
110. Zhang Q, Hu Y, Wei J, Zhu H. Neimenggu Bayannaouer shi Nalintaohai Han mu chutu rengu de wending tongweisu fenxi (Stable isotope analysis of human skeletons recovered from Nalintaohai, a Han dynasty cemetery in Bayannour city, Inner Mongolia). *Acta Anthropologica Sinica*. 2012; 31(4):408–14.
111. Zhang G, Hu Y, Wang L, Cao C, Li X, Wu X, et al. A paleodietary and subsistence strategy investigation of the Iron Age Tuoba Xianbei site by stable isotopic analysis: A preliminary study of the role of agriculture played in pastoral nomad societies in northern China. *Journal of Archaeological Science: Reports*. 2014; 2:699–707.
112. Zhang Q, Zhu H, Hu Y, Li Y, Cao J. Neimenggu Helingeer xian Xindianzi mudi gudai jumin de shipu fenxi (Dietary analysis to the human population from Xindianzi cemetery, Helinge'er county, Inner Mongolia). *Wenwu (Cultural Relics)*. 2006; 596:87–91.
113. Zhang X, Zhang J, Li Z, Zhang L, Chen G, Wang P, et al. Gansu Zhangye shi Xichengyi yizhi xianmin shiwu zhuangkuang de chubu fenxi (Preliminary isotopic analysis of ancient diets at Xinchengyi, Zhangye, Gansu). *Kaogu (Archaeology)*. 2015;(7):830–40.

114. Zhang X, Ye M. Lajia yizhi xianmin shiwu de chubu tansuo—Lajia yizhi zainan xianchang chutu rengu de tan dan wending tongweisu fenxi (Preliminary investigation to ancient diets of Lajia community—Carbon and nitrogen isotopic analysis to human remains recovered from Lajia). *Nanfang Wenwu* (Southern Cultural Relics). 2016;(4):197–202.
115. Ling X, Chen X, Wang J, Che L, Ma J, Ren M, et al. Xinjiang Balikun Dongheigou yizhi chutu rengu de tan dan tongweisu fenxi (Carbon and nitrogen isotope analysis of human skeletons recovered from Dongheigou, Balikun, Xinjiang). *Acta Anthropologica Sinica*. 2013; 32(2):219–25.
116. Zhang X, Qiu S, Zhang J, Guo W. Xinjiang Duogang mudi chutu rengu de tandan wending tongweisu fenxi (Carbon and nitrogen isotope analysis of human remains recovered from Duogang cemetery, Xinjiang). *Nanfang Wenwu* (Southern Cultural Relics). 2014;(3):79–59.
117. Qu Y, Yang Y, Hu Y, Wang C. The extraction of hair keratin at the Gumugou cemetery, Xinjiang and stable isotope analysis (C, N). *Geochimica*. 2013; 42(5):448–54.
118. Zhang Q, Zhu H. Xinjiang Gumugou mudi rengu de wending tongweisu fenxi: zaoqi Luobubo xianmin yinshi jieyou chutan (Stable isotopic analysis on human bones from Gumugou cemetery, Xinjiang: paleodietary reconstruction on Luobubo population). *Xiyu Yanjiu* (the Western Regions Studies). 2011; 3(3):91–6.
119. Wang T, Fuller BT, Wei D, Chang XE, Hu YW. Investigating dietary patterns with stable isotope ratios of collagen and starch grain analysis of dental calculus at the Iron Age cemetery site of Heiguoliang, Xinjiang, China. *International Journal of Osteoarchaeology*. 2016; 26:693–704.
120. Zhang Q, Chang X, Liu G. Xinjiang Balikun xian Heiguoliang mudi chutu rengu de shixing fenxi (Stable isotopic analysis on human bones from Heiguoliang cemetery in Barkol, Xinjiang). *Xiyu Yanjiu* (the Western Regions Studies). 2009; 3:45–9.
121. Zhang Q, Li S. Xinjiang Nileke Xian Kongkeke yihao mudi gudai jumin de shiwu jieyou fenxi (Analysis on paleodiet of ancient inhabitants in no. 1 cemetery of Qiongkeke at Nika County, Xinjiang). *Xiyu Yanjiu* (the Western Regions Studies). 2006; 4:78–82.
122. Chen X, Yu J, You Y. Tan dan wending tongweisu suojian Xinjiang Kalasu mudi de zang ma xisu (Horse burial customs in Kalasu cemetery of Xinjiang: Carbon and nitrogen isotopic perspectives). *Xiyu Yanjiu* (Studies of Western Regions). 2017;(4):89–98.
123. Wang T, Wei D, Chang X, Yu Z, Zhang X, Wang C, et al. Tianshanbeilu and the isotopic millet road: Reviewing the late Neolithic/Bronze Age radiation of human millet consumption from north China to Europe. *National Science Review*. 2017: <https://doi.org/10.1093/nsr/nwx015>
124. Zhang Q, Chang X, Liu G. Xinjiang Hami Tianshanbeilu mudi chutu rengu de wending tongweisu fenxi (Stable isotopic analysis of human bones unearthed from Tianshan Beilu cemetery in Hami, Xinjiang). *Xiyu Yanjiu* (the Western Regions Studies). 2010; 2:38–43.
125. Zhang X, Wei D, Yu Y, Nie Y, Hu Y. Carbon and nitrogen stable isotope ratio analysis of Bronze Age humans from the Xiabandi cemetery, Xinjiang, China: Implications for cultural interactions between the East and West. *Science China Press*. 2016; 61(32):3509–19.
126. Si Y, Lü E, Li X, Jiang H, Hu Y, Wang C. Exploration of human diets and populations from the Yanghai Tombs, Xinjiang. *Science China Press*. 2013; 58(15):1422–9.
127. Xia Y, Zhang J, Yu F, Zhang H, Wang T, Hu Y, et al. Breastfeeding, weaning, and dietary practices during the Western Zhou Dynasty (1122–771 BC) at Boyangcheng, Anhui Province, China. *American Journal of Physical Anthropology*. 2017; 165(2):343–52. <https://doi.org/10.1002/ajpa.23358> PMID: [29131307](https://pubmed.ncbi.nlm.nih.gov/29131307/)
128. Zhang Q. Yunnan Chengjiang xian Jinlianshan mudi chutu rengu wending tongweisu de chubu fenxi (Preliminary isotopic analysis of human skeletons recovered from Jinlianshan cemetery, Chengjiang county, Yunnan). *Kaogu* (Archaeology). 2011;(1):30–3.
129. Ren L, Li X, Kang L, Brunson K, Liu H, Dong W, et al. Human paleodiet and animal utilization strategies during the Bronze Age in northwest Yunnan Province, southwest China. *PLOS ONE*. 2017; 12(5): <https://doi.org/10.1371/journal.pone.0177867>
130. Zhou L. Wending tongweisu shijiao xia de Henan Longshan muzang yu shehui (Mortuary practice and social structure of Longshan culture in Henan—an isotopic perspective). *Huaxia Kaogu* (Huaxia Archaeology). 2017;(3):145–52.