



University of
Salford
MANCHESTER

The prevalence of *Neospora caninum* and co-infection with *Toxoplasma gondii* by PCR analysis in naturally occurring mammal populations

Hughes, JM, Williams, RH, Morley, EK, Cook, DAN, Terry, RS, Murphy, RG, Smith, JE and Hide, G

<http://dx.doi.org/10.1017/S0031182005008784>

Title	The prevalence of <i>Neospora caninum</i> and co-infection with <i>Toxoplasma gondii</i> by PCR analysis in naturally occurring mammal populations
Authors	Hughes, JM, Williams, RH, Morley, EK, Cook, DAN, Terry, RS, Murphy, RG, Smith, JE and Hide, G
Publication title	Parasitology
Publisher	Cambridge University Press
Type	Article
USIR URL	This version is available at: http://usir.salford.ac.uk/id/eprint/658/
Published Date	2006

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: library-research@salford.ac.uk.

The prevalence of *Neospora caninum* and co-infection with *Toxoplasma gondii* by PCR analysis in naturally occurring mammal populations

J. M. HUGHES¹, R. H. WILLIAMS¹, E. K. MORLEY¹, D. A. N. COOK¹, R. S. TERRY²,
R. G. MURPHY³, J. E. SMITH² and G. HIDE^{1*}

¹Centre for Parasitology, Molecular Epidemiology and Ecology, Biosciences Research Institute, School of Environment and Life Sciences, University of Salford, Salford M5 4WT

²School of Biology, University of Leeds, Leeds LS2 9JT

³Research Institute for the Built and Human Environment, University of Salford, Salford M5 4WT

(Received 1 June 2005; revised 26 July 2005; accepted 27 July 2005; first published online 15 September 2005)

SUMMARY

Neospora caninum and *Toxoplasma gondii* are closely related intracellular protozoan parasites associated with bovine and ovine abortion respectively. Little is known about the extent of *Neospora/Toxoplasma* co-infection in naturally infected populations of animals. Using nested PCR techniques, based on primers from the Nc5 region of *N. caninum* and SAG1 for *T. gondii*, the prevalence of *N. caninum* and its co-infection with *T. gondii* were investigated in populations of *Mus domesticus*, *Rattus norvegicus* and aborted lambs (*Ovis aries*). A low frequency of infection with *N. caninum* was detected in the *Mus domesticus* (3%) and *Rattus norvegicus* (4.4%) populations. A relatively high frequency of infection with *N. caninum* was detected in the brains of aborted lambs (18.9%). There was no significant relationship between *N. caninum* and *T. gondii* co-infection. Investigation of the tissue distribution of *Neospora*, in aborted lambs, showed that *Neospora* could not be detected in tissues other than brain and this was in contrast to *Toxoplasma* where the parasite could be frequently detected in a range of tissues.

Key words: *Neospora caninum*, *Toxoplasma gondii*, PCR, co-infection, naturally acquired infection, mouse, rat, sheep.

INTRODUCTION

Toxoplasma gondii and *Neospora caninum* are closely related Apicomplexan parasites which are morphologically similar but possess some structural, molecular and antigenic differences (Ellis *et al.* 1998; Speer *et al.* 1999; Howe and Sibley, 1999). They have similar life-cycles with different definitive hosts, the felids (Hutchinson, 1965) and canids (McAlister *et al.* 1998; Gondim *et al.* 2004) respectively, and have similar intermediate hosts including a wide range of mammals (Dubey, 1999; Tenter, Heckerth and Weiss, 2000). The ingestion of infective oocysts from the environment and tissue cysts from raw meat is considered to be a major route of infection of *T. gondii* (Tenter *et al.* 2000) and vertical transmission from mother to offspring a major route of infection of *N. caninum* (Davison, Otter and Trees, 1999). Both parasites cause significant disease and economic loss in the farming industry with *T. gondii* primarily causing abortion

and foetal abnormality in sheep, and *N. caninum* causing abortion and foetal abnormality in cattle. *T. gondii* is also an important pathogen of humans while, in contrast, there is currently little evidence of human infection caused by *N. caninum* (Graham *et al.* 1999).

The balance of similarities and differences between these parasites raises the question as to whether frequent co-infection of individual hosts occurs and the degree to which these parasites may interact by acting antagonistically or synergistically during infection and disease. For example, does infection with one parasite prevent infection with the other or is the outcome of infection influenced by co-infection? Previous serological investigations have shown that antibodies to both parasites were present in several species including coyotes (Lindsay *et al.* 1996) and non-domestic felids from southern Africa (Cheadle, Spencer and Blagburn, 1999) but few studies have investigated this question in other species. Previous *in vitro* studies (Sunderman and Estridge, 1999) have demonstrated that *Toxoplasma* and *Neospora* can co-exist in tissue culture, and both species could invade the same host cell and undergo endodyogeny. Immunological studies (Innes *et al.* 2001) showed that, although there was evidence of induction of cross-reactive immune recognition following

* Corresponding author: Centre for Parasitology, Molecular Epidemiology and Ecology, Biosciences Research Institute, School of Environment and Life Sciences, University of Salford, Salford M5 4WT. Tel: +161 295 3371. Fax: +161 295 5015. E-mail: g.hide@salford.ac.uk

immunization with *T. gondii* in sheep, this was not sufficient to prevent foetal loss following subsequent *N. caninum* oocyst challenge.

The main objectives of this study were to develop a sensitive nested PCR technique for the detection of *N. caninum* DNA in host tissue, to use this to determine the prevalence of *N. caninum* in 3 naturally infected mammalian populations in Britain (rats, mice and sheep), and to determine the levels of *N. caninum* co-infection with *T. gondii* in these species.

MATERIALS AND METHODS

Collection of samples and DNA extraction

House mice (*Mus domesticus*) used in this study were captured within domestic dwellings in the Cheetham Hill area of Manchester, UK as part of a pest control programme (Marshall *et al.* 2004), brown rats (*Rattus norvegicus*) were captured in urban areas of Manchester. Tissue was obtained from lambs (*Ovis aries*) which were either aborted, dead at birth or died shortly after birth from 2 farms and contained individuals from 3 breeds, Suffolk cross, Charolais and Lleyns as described by Duncanson *et al.* (2001), Williams *et al.* (2005) and Morley *et al.* (2005). Approximately 1 g of brain tissue was dissected from mice, rats and lambs and, in the case of lambs, tissue was from the midbrain area along with samples of umbilical cord, tongue and heart. Cord tissue was not available for all lambs due to farming constraints during sampling and this was especially the case for the *N. caninum*-infected aborted lambs of which very few were obtained. All tissue samples were aseptically collected, using sterile instruments with great care being taken to exclude contamination either across tissues or from the surrounding environment, and stored frozen until DNA could be extracted under clean laboratory conditions as described by Williams *et al.* (2005).

DNA was extracted using a proteinase K lysis followed by phenol chloroform, ethanol precipitation method as described previously (Duncanson *et al.* 2001). DNA was resuspended in 100 μ l of T.E. buffer, pH 8.0 and stored at 4 °C.

PCR detection of Toxoplasma gondii

Detection of *T. gondii* was carried out using nested PCR amplification of the Surface Antigen Gene 1 (Savva *et al.* 1990) as modified by Morley *et al.* (2005). The PCR product from one positive example of each species was extracted from the gel using the GENECLEAN 2 kit (Q-BIOgene) and sequenced in both directions (Lark Technologies Inc.) to confirm the detection of *Toxoplasma* (Accession nos. DQ077665 (sheep), DQ077664 (rat) and DQ077666 (mouse)).

PCR detection of Neospora caninum

The Nc5 region of *N. caninum* has been cloned and shown to be highly specific to *N. caninum* by DNA hybridization or PCR and is clearly distinguishable from *T. gondii* (Kaufmann *et al.* 1996; Yamage, Flechner and Gottstein, 1996). This gene was selected as the target for PCR amplification and detection using primers Np21 PLUS (5' GGG TGT GCG TCC AAT CCT GTA AC 3') and Np6 PLUS (5' CTC GCC AGT CAA CCT ACG TCT TCT 3') (Liddell, Jenkins and Dubey, 1999) both alone and as a nested technique combined with primers Np6 (5' CAG TCA ACC TAC GTC TTC T 3') and Np7 (5' GGG TGA ACC GAG GGA GTT G 3') (Yamage *et al.* 1996; Baszler *et al.* 1999) in the 2nd round. Both primer sets had previously been shown to be *N. caninum* specific, we further ensured the specificity of the technique by confirming there was no amplification using our nested technique, from 4 different isolates (RH, Martin, 17695 and COR), representing the 3 lineages of *T. gondii*. The 1st round reaction was performed as described by Liddell *et al.* (1999) with minor modifications as described below, 50 μ l reactions contained 5 μ l of 10 \times PCR buffer (HT biotechnology) (100 mM Tris-HCl, pH 9.0, 15 mM MgCl₂, 500 mM KCl, 1% Triton X-100, 0.1% (w/v) stabilizer), 0.5 mM MgCl₂, 12.5 nM each dNTP, 10 pM each primer, 2.5 units Taq polymerase (Bioline) and 1 μ l of test DNA or water. After an initial denaturation step of 5 min at 95 °C, 40 cycles of PCR were performed with cycle times of 40 sec at 94 °C, 40 sec at 63 °C, 1 min 10 sec at 72 °C, followed by a final step of 10 min at 72 °C. The second round PCR was carried out as described by Baszler *et al.* (1999), reaction conditions were the same as the 1st round except 25 pM of primers Np7 and Np6 were used. Two microlitres of 1st round product was used as the template and cycling conditions were as the 1st round except that an annealing temperature of 56 °C was used. These PCR methods were compared to the semi-nested technique of Baszler *et al.* (1999) to calculate the performance of the test. All PCR reactions were performed using a Stratagene ROBOCYCLER™. PCR products were run on a 2% Agarose TBE gel containing ethidium bromide and visualized on an Alpha imager™ 1220. All PCR reactions were performed a minimum of 3 times to ensure comprehensive sampling of the test DNA. Due to the sensitive nature of the nested PCR technique negative controls were included in all PCR reactions and contamination could easily be tracked since many samples were naturally negative and therefore most positive reactions were interspaced between negative reactions. Positive PCR products from each host species were extracted from the gel using the GENECLEAN 2 kit (Q-BIOgene) cloned using the TOPO TA cloning® kit (Invitrogen) and sequenced (Lark technologies Inc.) to confirm the




Fig. 1. Comparison of the detection of *Neospora caninum* DNA, using single and nested PCR techniques. Rat DNA was spiked with dilutions of *N. caninum* DNA and PCR amplification was carried out using single round PCR using primers Np21PLUS and Np6PLUS (Panel A) and nested PCR 1st round primers Np21PLUS and Np6PLUS, 2nd round primers Np6 and Np7 (Panel B). Lane 1 Undiluted *N. caninum* DNA, Lane 2 1/5 dilution of *N. caninum* DNA, Lane 3 1/25 dilution of *N. caninum* DNA, Lane 4 1/125 dilution of *N. caninum* DNA, Lane 5 1/625 dilution of *N. caninum* DNA Lane 6 negative control, Lane M Hyper ladder1 marker (Bioline).

presence of DNA corresponding to the *N. caninum* Nc5 region. All negative PCR samples underwent a Tubulin PCR (Terry *et al.* 2001), targeting host DNA, to check the sample was suitable for PCR.

RESULTS

Evaluation of a nested PCR detection system for Neospora caninum

Previous work on *T. gondii* detection (Duncanson *et al.* 2001) suggested that the use of a nested PCR reaction would be required for the accurate detection of apicomplexan parasites directly from mammalian tissues. Although single-round PCR techniques (e.g. Liddell *et al.* 1999) and a semi-nested technique (Baszler *et al.* 1999) are available for the detection of *N. caninum*, we decided to develop a more sensitive detection method which can detect small quantities of *Neospora* DNA, in the presence of huge excesses of mammalian DNA. A series of dilution experiments were carried out using the NP21PLUS, NP6PLUS, NP7 and NP6 primers to determine the sensitivity of this nested PCR approach and compared it with the semi nested technique of Baszler *et al.* (1999). A series of dilutions of *N. caninum* DNA in genomic rat DNA was used as the template in nested PCR reactions using primers NP21PLUS and NP6PLUS in the first round followed by NP7 and NP6 in the second round. The results presented in Fig. 1 show that in the first round *Neospora* DNA could be detected at a dilution of 1 in 25 (lane 3) and following the nested PCR, detection was increased 5-fold to a 1 in 125 dilution (lane 4). We found the semi-nested technique of Baszler *et al.* (1999) was 4-fold less sensitive than our nested technique when used in the presence of host DNA (data not shown). An initial comparison of the single-round and nested PCR approaches on actual tissue samples from sheep revealed that the single-round approach underestimated the number of positive samples by 36% when compared to nested PCR. Even after many repeated tests of the same samples we were still unable to detect 14% of positive

tissue samples using the single round alone, when compared with the nested technique. Furthermore when testing DNA extracted from infected tissue samples, bands from the single round PCR were always faint and usually difficult to distinguish clearly. Tissues positive for *N. caninum* usually produced a clear strong band following the second round, allowing easier interpretation of gels. Sequencing of the *N. caninum* PCR product amplified from an example of each host species (Accession nos. DQ077661 (sheep), DQ077662 (rat) and DQ077663 (mouse)) confirmed the specificity of the PCR with greater than 97% homology with the published Nc5 sequence (Accession no. X84238) (Yamaga *et al.* 1996) (Fig. 2).

Neospora caninum and Toxoplasma gondii co-infection

To compare rates of co-infection of *T. gondii* and *N. caninum*, nested PCR was carried out on DNA extracted from brain tissue from the mice, rat and lamb populations. Three (3.0%) of the 100 mice tested positive by PCR for *N. caninum* and 53 (53.0%) were positive for *T. gondii*. Two (4.4%) of the 43 rats were positive for *N. caninum* and 19 (42.2%) were *T. gondii* positive. Of the 74 aborted lambs 14 (18.9%) tested *N. caninum* positive and 52 (70.3%) *T. gondii* positive (Table 1). As shown in Table 1, both co-infection and the absence of co-infection were present in the mice with 2 mice (2.0%) testing positive for *T. gondii* and *N. caninum*, and 1 (1.0%) testing positive for *N. caninum* alone. Neither of the 2 *N. caninum*-positive rats tested positive for Toxoplasmosis. The numbers of *N. caninum*-positive samples in these 2 species was too small to enable further statistical analysis.

Of the 74 lambs 9 (12.2%) were positive for *N. caninum* and *T. gondii*, 5 (6.7%) for *N. caninum* alone, 43 (58.1%) for *T. gondii* alone and 17 (23%) were negative for both parasites. Statistical analysis using the χ^2 contingency table test for association and Yates' correction for continuity demonstrated there

```

NCSSDNAP      GGGTGAACCGAGGGAGTTGGTAGCGGTGAGAGGTGGGATACGTGGTTTGTGGTTAGTCAT
NCMus         GGGTGAACCGAGGGAGTTGGTAGCGGTGAGAGGTGGGATACGTGGTTTGTGGTTAGTCAT
NCOvis        GGGTGAACCGAGGGAGTTGGTAGCGGTGAGAGGTGGGATACGTGGTTTGTGGTTAGTCAT
NCRattus      GGGTGAACCGAGGGAGTTGGTAGCGGTGAGAGGTGGGATACGTGGTTTGTGGTTAGTCAT
*****

NCSSDNAP      TCGTCACGTTGAAATCAGCCTGCGTCAGGGTGAGGACAGTGTGTCAATGATACTTATCGA
NCMus         TCGTCACGTTGAAATCAGCCTGCGTCAGGGTGTGGACAGTGTGTCAATGATACTTATCGA
NCOvis        TCGTCACGTTGAAATCAGCCTGCGTCAGGGTGTGGACAGTGTGTCAATGATACTTATCCA
NCRattus      TCGTCACGTTGAAATGAGCCTGCGTCAGGGTGTGGACAGTGTGTCAATGATACTTATCCA
*****

NCSSDNAP      GAGTTCAGTGTCTGTGTTGAGGCAACACCGCGGCCTGATGACGGGGGAGATTATTCGA
NCMus         GAGTTCAGTGTCTGTGTTGAGGCAACACCGCGGCCTGATGACGGGGGAGATTATTCGA
NCOvis        GAGTTCAGTGTCTGTGTTGAGGCAACACCGCGGCCTGATGACGGGGGAGATTATTCGA
NCRattus      GAGTTCAGTGTCTGTGTTGAGGCAACACCGCGGCCTGATGACGGGGGAGATTATTCGA
*****

NCSSDNAP      TAGGGAGCAAGCGGACGAGGGAAGGGGCAGAAGACGTAGGTTGACTG
NCMus         TAGGGAGCAAGCGGACGAGGGAAGGGGCAGAAGACGTAGGTTGACTG
NCOvis        TAGGGAGCAAGCGGACGAGGGAAGGGGCAGAAGACGTAGGTTGACTG
NCRattus      TGGGGAGCAAGCGGACGAGGGAAGGGGCAGAAGACGTAGGTTGACTG
*****
    
```

Fig. 2. Sequence alignment of PCR fragments amplified from naturally infected mammalian brains using *Neospora caninum*-specific primers compared with part of the Nc5 region of *N. caninum* (Yamaga *et al.* 1996) NCSSDNAP (Accession no. X84238), NCMus, PCR product amplified from a naturally infected mouse (Accession no. DQ077663), NCOvis, PCR product amplified from a naturally infected sheep (Accession no. DQ077661), NCRattus, PCR product amplified from a naturally infected rat (Accession no. DQ077662). ★ Denotes exact match between all sequences, ● denotes a mismatch with at least 1 sequence.

was neither positive or negative association between *N. caninum* and *T. gondii* infection in this population of lambs ($P=0.85$). Within our sample of 3 breeds, Suffolk cross, Charolais and Lleyens, we found individuals from all 3 breeds infected with *N. caninum* showing that there was no association with breed.

Tissue distribution of Neospora caninum in lambs

As differences have been reported in the tissue distribution of *T. gondii* and *N. caninum* (Duncanson *et al.* 2001; Pereira-Bueno *et al.* 2002) we investigated this aspect using the aborted lamb tissue

samples. Samples for brain, heart and cord, where available, were collected from *N. caninum*-positive lambs and analysed for *N. caninum* presence, a similar number of *T. gondii*-positive lambs tissues were compared. Although *T. gondii* could be detected reliably in a number of tissues, *N. caninum* was only detectable in the DNA extracted from brain tissue (Table 2). Fig. 3 shows PCR results of *N. caninum* and *T. gondii* specific PCRs from 1 lamb that was positive for both *N. caninum* and *T. gondii*. The results presented clearly show that *N. caninum* was only detectable in DNA extracted from brain tissue but *T. gondii* was detectable in DNA extracted from brain, heart, cord and tongue tissues.

Table 1. Frequency of the levels of infection and co-infection of *Neospora caninum* and *Toxoplasma gondii* in the brain tissue of mice (A), rats (B) and aborted lambs (C)

	<i>Neospora</i> -positive	<i>Neospora</i> -negative	Total
(A) Mice			
<i>Toxoplasma</i> -positive	2 (2%)	51 (51%)	53 (53%)
<i>Toxoplasma</i> -negative	1 (1%)	46 (46%)	47 (47%)
Total	3 (3%)	97 (97%)	100
(B) Rats			
<i>Toxoplasma</i> -positive	0 (0%)	19 (42.2%)	19 (42.2%)
<i>Toxoplasma</i> -negative	2 (4.4%)	24 (53.3%)	26 (57.8%)
Total	2 (4.4%)	43 (95.6%)	45
(C) Aborted lambs			
<i>Toxoplasma</i> -positive	9 (12.2%)	43 (58.1%)	52 (70.3%)
<i>Toxoplasma</i> -negative	5 (6.7%)	17 (23%)	22 (29.7%)
Total	14 (18.9%)	60 (81.1%)	74

Table 2. Comparison of sites of successful detection of *Toxoplasma gondii* and *Neospora caninum* in different tissues of *Toxoplasma* and *Neospora* positive lambs respectively

(“+” Denotes successful SAG1 or Nc5 PCR, “-” denotes unsuccessful SAG1 or Nc5 PCR, “nd” denotes no sample available.)

<i>Toxoplasma gondii</i> -positive lambs				<i>Neospora caninum</i> -positive lambs			
Lamb	Brain	Heart	Cord	Lamb	Brain	Heart	Cord
A	+	+	+	1	+	-	nd
B	+	+	+	2	+	-	nd
C	+	+	+	3	+	-	nd
D	+	+	-	4	+	-	-
E	+	+	+	5	+	nd	nd
F	+	+	+	6	+	-	nd
G	+	-	+	7	+	-	nd
H	+	+	+	8	+	-	-
I	+	nd	+	9	+	nd	nd
J	+	+	+	10	+	nd	nd
K	+	+	+	11	+	nd	nd
L	+	+	+	12	+	nd	nd
M	+	-	+	13	+	-	-
N	+	+	nd	14	+	-	-




Fig. 3. Detection of *Toxoplasma gondii* and *Neospora caninum* DNA in different lamb tissues. Lamb DNA samples were amplified by PCR to detect *N. caninum* DNA (Panel A) and *T. gondii* DNA (Panel B). DNA was extracted from tongue (Lane 1), cord (Lane 2), heart (Lane 3), brain (Lane 4), Lane 5 negative control, Lane 6 positive control, Lane M Hyper ladder1 marker (Bioline).

DISCUSSION

Since *N. caninum* was identified as a separate species a number of methods of identification have been developed, including serological, immunohistochemical labelling and molecular methods including PCR. PCR is recognized as a specific sensitive technique used to identify species-specific parasite DNA sequences (Wastling, Nicoll and Buxton, 1993). A number of PCR methods have been used to assess *Neospora* infection e.g. Payne and Ellis (1996), Lally, Jenkins and Dubey (1996), Ho *et al.* (1997) and Gottstein *et al.* (1998). We further developed a nested PCR technique based on the highly repeated Nc5 region specific to *N. caninum* (Kaufmann *et al.* 1996) using primers NP21PLUS, NP6PLUS (Liddell *et al.* 1999) combined with NP7, NP6 (Yamaga *et al.* 1996; Baszler *et al.* 1999). This increased the sensitivity of detection 5-fold and also allowed the use of greater quantities of host DNA as template without masking the final result. In common with previous authors (Ho *et al.* 1997; Almeria *et al.* 2002) we found that the same samples did

not always test positive and a minimum of 3 PCR reactions were carried out on separate occasions. Examination of field samples demonstrated that, despite repeated testing, the single-round PCR underestimated the number of *Neospora*-infected individuals by 14.3%, when compared to the nested technique. In contrast Baszler *et al.* (1999), in a study of *N. caninum* prevalence in foetal tissues from spontaneous bovine abortions, found that although semi-nested PCR increased the sensitivity of detection, it was unnecessary for the detection of infection. Two possible explanations for this discrepancy exist. Firstly, different host species (i.e. bovine *vs* ovine) may have a different parasite density and may therefore require different test conditions. Secondly, Baszler *et al.* (1999) compared only single-round and semi-nested PCR techniques. We have shown that semi-nested PCR is 4-fold less sensitive than our technique, clearly demonstrating that a nested PCR technique is required for diagnosis in sheep. Previous studies using primers Np21PLUS and NP6PLUS have shown a detection limit of 1 tachyzoite in 0.5 g brain tissue (Almeria *et al.* 2002) by adding the

second round of PCR using primers NP7 and NP6, we increased the sensitivity by 5-fold to the equivalence of 0.2 tachyzoites in 0.5 g brain tissue.

In our samples we found very low levels of *N. caninum* infection in both mice (3/100, 3%) and rats (2/45, 4.4%). It is interesting to note that a recent study of rats from cattle farms with *N. caninum* associated abortion found serological evidence of exposure to *N. caninum* in 9/55 rats, with *N. caninum* DNA detected by a single-round PCR in 2 of these serologically positive individuals (Huang *et al.* 2004). In both host species levels of *T. gondii* infection were significantly higher (53/100, 53% and 19/45, 42.2% respectively). This suggests that there is no particular linkage in the epidemiological factors that determine prevalence levels of *T. gondii* and *N. caninum* in mice and rats. The rats and mice in our study were collected from an urban environment, investigation of similar samples from a rural area might provide insight into the importance of different modes of transmission, for example, it might be expected in an urban area with a greater density of dogs that infection by the ingestion of oocysts may increase levels of infection when combined with vertical transmission whereas in rural areas with fewer dogs vertical transmission may be the more important transmission route. Further work is needed to address this question.

In our studies of aborted lambs we found high levels of infection with *Neospora* (14/74, 18.9%). Naturally occurring *Neospora* infection of ovines detected either by serological survey or PCR have been reported as rare (Hemphill and Gottstein, 2000). However, Hassig *et al.* (2003) detected *N. caninum* using a single-round PCR method in the brain tissue of 4 out of 20 (20%) aborted lambs and our data are in accordance with these figures. Further work may be required to investigate this variability in reported prevalence.

A number of serological investigations into co-infection of *Neospora* and *Toxoplasma* have been reported in the literature including co-infection rates of 4/53 (7.7%) of coyotes (Lindsay *et al.* 1996), 4/68 (5.9%) of non-domestic felids from southern Africa (Cheadle *et al.* 1999), 0/66 camels (Hilali *et al.* 1998), 0/221 red foxes (Jakubek *et al.* 2001), 1/549 red foxes (Hamilton *et al.* 2005), 12/144 (10.5%) sheep (Hassig *et al.* 2003) and 3.5% of 597 sheep (Figliuolo *et al.* 2004). In our experiments we found co-infection with DNA from both parasites in 2% of mice but not in any of the 43 rats tested, however the prevalence of *Neospora* was low and, therefore, statistical analysis was inappropriate. Of the aborted lambs 9/74 (12.2%) were co-infected, confirming previous serological results (Hassig *et al.* 2003; Figliuolo *et al.* 2004) that co-infection can occur in sheep. However, our results were in contrast to a PCR based study in Switzerland (Hassig *et al.* 2003) where there was no observed co-infection with toxoplasmosis in the 4/20

(20%) aborted lambs infected with *Neospora*. This is probably due to the small numbers of animals that exhibited either *N. caninum* or *T. gondii* ($n=7$). On the other hand our larger data set of infected animals ($n=57$) allowed us to address the question of co-infection and we demonstrated there was no statistical association between infection with *Neospora* and *Toxoplasma* ($P=0.85$). Our results support the *in vitro*, immunological and serological evidence that there is no exclusivity of infection and that co-infection would appear to be a random event.

We investigated tissue distribution in aborted lambs to answer a number of questions including whether or not *Neospora* could be detected in cord tissue. Lamb cord is foetally derived and if *Neospora* could be reliably detected in it we could use this as a means of screening live births for *Neospora* infection. *Neospora* has been amplified from many experimentally and naturally infected bovine tissues again with the most successful amplification from brain tissue (Gottstein *et al.* 1998; Baszler *et al.* 1999). Ho *et al.* (1997) suggested that experimentally infected bovines appear to have a wider tissue distribution of *Neospora* sp. DNA than naturally infected animals, with natural infection being detected in brain and spinal cord and rarely in other tissue. Our studies support this hypothesis in the case of aborted lambs.

Our investigations demonstrated that naturally occurring infection of *N. caninum* occurs in all three species of mammals investigated but at a much lower prevalence than *T. gondii*. We also found evidence that co-infection with *T. gondii* was occurring but that there was no specific association. The incidence of *N. caninum*-infected mice and rats has implications for the translocation of Neosporosis across wider geographical areas due to the spread of infection through rodent and other pest populations. Furthermore, the apparent broad host range of *Neospora* raises the question as to if or when it might appear as a human parasite. Further research is needed to investigate these questions.

We would like to acknowledge The University of Salford, Perry Foundation and The Wellcome Trust for providing funding for the work conducted in this study. Thanks go to A. M. Tenter and J. P. Dubey for the control *Neospora caninum* DNA. We gratefully acknowledge the Cheetham and Broughton SRB, Peter Marshall and Allan Maiden and family for their involvement in this research.

REFERENCES

- Almeria, S., Ferrer, D., Pabon, M, Castella, J. and Manas, S.** (2002). Red foxes (*Vulpes vulpes*) are a natural intermediate host of *Neospora caninum*. *Veterinary Parasitology* **107**, 287–294.
- Baszler, T. V., Gay, L. J. C., Long, M. T. and Mathison, B. A.** (1999). Detection by PCR of *Neospora caninum* in fetal tissues from spontaneous bovine abortions. *Journal of Clinical Microbiology* **37**, 4059–4064.

- Cheadle, M. A., Spencer, J. A. and Blagburn, B. L.** (1999). Seroprevalences of *Neospora caninum* and *Toxoplasma gondii* in nondomestic felids from southern Africa. *Journal of Zoo and Wildlife Medicine* **30**, 248–251.
- Davison, H. C., Otter, A. and Trees, A. J.** (1999). Estimation of vertical and horizontal transmission parameters of *Neospora caninum* in dairy cattle. *International Journal for Parasitology* **29**, 1683–1689.
- Dubey, J. P.** (1999). Recent advances in *Neospora* and Neosporosis. *Veterinary Parasitology* **84**, 349–367.
- Duncanson, P., Terry, R. S., Smith, J. E. and Hide, G.** (2001). High levels of congenital transmission of *Toxoplasma gondii* in a commercial sheep flock. *International Journal for Parasitology* **31**, 1699–1703.
- Ellis, J. T., Amoyl, G., Ryce, C., Harper, P. A. W., Clough, K. A., Homan, W. L. and Brindley, P. J.** (1998). Comparison of the large subunit ribosomal DNA of *Neospora* and *Toxoplasma* and development of a new genetic marker for their differentiation based on the D2 domain. *Molecular and Cellular Probes* **12**, 1–13.
- Figliuolo, L. P. C., Kasai, N., Ragozo, A. M. A., de Paula, V. S. O., Dias, R. A., Souza, S. L. A., Gennari, S. M.** (2004). Prevalence of anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in ovine from Sao Paulo State, Brazil. *Veterinary Parasitology* **123** 161–166.
- Gondim, L. F. P., McAllister, M. M., Pitt, W. C. and Zemlicka, D. E.** (2004). Coyotes (*Canis latrans*) are definitive hosts of *Neospora caninum*. *International Journal for Parasitology* **34**, 159–161.
- Gottstein, B., Hentrich, B., Wyss, R., Thur, B., Busato, A., Stark, K. D. C. and Muller, N.** (1998). Molecular and immunodiagnostic investigations on bovine neosporosis in Switzerland. *International Journal for Parasitology* **28**, 679–691.
- Graham, D. A., Calvert, V., Whyte, M., Marks, J.** (1999). Absence of serological evidence for human *Neospora caninum* infection. *Veterinary Record* **144**, 672–673.
- Hamilton, C. M., Gray, R., Wright, S. E., Gangadharan, B., Laurenson, K., Innes, E. A.** (2005). Prevalence of antibodies to *Toxoplasma gondii* and *Neospora caninum* in red foxes (*Vulpes vulpes*) from around the UK. *Veterinary Parasitology* **130**, 169–173.
- Hassig, M., Sager, H., Reitt, K., Ziegler, D., Strabel, D. and Gottstein, B.** (2003). *Neospora caninum* in sheep: a herd case report. *Veterinary Parasitology* **117**, 213–220.
- Hilali, M., Romand, S., Thulliez, P., Kwok, O. C. and Dubey, J. P.** (1998). Prevalence of *Neospora caninum* and *Toxoplasma gondii* antibodies in sera from camels from Egypt. *Veterinary Parasitology* **75**, 269–271.
- Hemphill, A. and Gottstein, B.** (2000). A European perspective on *Neospora caninum*. *International Journal for Parasitology* **30**, 877–924.
- Ho, M. S. Y., Barr, B. C., Rowe, J. D., Anderson, M. L., Sverlow, K. W., Packham, A., Marsh, A. E. and Conrad, P. A.** (1997). Detection of *Neospora* sp. From infected bovine tissues by PCR and probe hybridization. *Journal of Parasitology* **83**, 508–514.
- Howe, D. K. and Sibley, D.** (1999). Comparison of the major antigens of *Neospora caninum* and *Toxoplasma gondii*. *International Journal for Parasitology* **29** 1489–1496.
- Huang, C. C., Yang, C. H., Watanabe, Y., Laio, Y. K. and Ooi, H. K.** (2004). Finding of *Neospora caninum* in the wild brown rat (*Rattus norvegicus*). *Veterinary Research* **35**, 283–290.
- Hutchinson W. M.** (1965). Experimental transmission of *Toxoplasma gondii*. *Nature, London* **206**, 961–962.
- Innes, E. A., Lunden, A., Esteban, I., Marks, J., Maley, S., Wright, S., Rae, A., Harkins, D., Vermeulen, A., McKendrick, I. J. and Buxton, D.** (2001). A previous infection with *Toxoplasma gondii* does not protect against a challenge with *Neospora caninum* in pregnant sheep. *Parasite Immunology* **23**, 121–132.
- Jakubek, E. B., Brojer, C., Regnersen, C., Uggl, A., Schares, G. and Bjorkman, C.** (2001). Seroprevalences of *Toxoplasma gondii* and *Neospora caninum* in Swedish red foxes (*Vulpes vulpes*). *Veterinary Parasitology* **102**, 167–172.
- Kaufmann, H., Yamage, M., Roditi, I., Dobbelaere, D., Dubey, J. P., Holmdahl, O. J. M., Trees, A. and Gottstein, B.** (1996). Discrimination of *Neospora caninum* from *Toxoplasma gondii* and other apicomplexan parasites by hybridization and PCR. *Molecular and Cellular Probes* **10**, 289–297.
- Lally, N. C., Jenkins, M. C. and Dubey, J. P.** (1996). Development of a polymerase chain reaction assay for the diagnosis of neosporosis using the *Neospora caninum* 14-3-3 gene. *Molecular and Biochemical Parasitology* **75**, 169–178.
- Liddell, S., Jenkins, M. C. and Dubey, J. P.** (1999). A competitive PCR assay for the quantitative detection of *Neospora caninum*. *International Journal for Parasitology* **29**, 1583–1587.
- Lindsay, D. S., Kelly, E. J., McKown, R. D., Stein, F. J., Plozer, J., Herman, J., Blagburn, B. L. and Dubey, J. P.** (1996). Prevalence of *Neospora caninum* and *Toxoplasma gondii* antibodies in coyotes (*Canis latrans*) and experimental infection of coyotes with *Neospora caninum*. *Journal of Parasitology* **82**, 657–659.
- Marshall, P. A., Hughes, J. M., Williams, R. H., Smith, J. E., Murphy, R. G. and Hide, G.** (2004). Detection of high levels of congenital transmission of *Toxoplasma gondii* in natural urban populations of *Mus domesticus*. *Parasitology* **128**, 39–42.
- McAllister, M. M., Dubey, J. P., Lindsay, D. S., Jolley, W. R., Wills, R. A. and McGuire, A. M.** (1998). Dogs are the definitive hosts of *Neospora caninum*. *International Journal of Parasitology* **28**, 1473–1478.
- Morley, E. K., Williams, R. H., Hughes, J. M., Terry, R. S., Duncanson, P., Smith, R. S. and Hide, G.** (2005). Significant familial differences in the frequency of abortion and *Toxoplasma gondii* infection within a flock of Charollais sheep. *Parasitology* **131**, 181–185.
- Payne, S. and Ellis, J.** (1996). Detection of *Neospora caninum* DNA by the polymerase chain reaction. *International Journal for Parasitology* **26**, 347–351.
- Pereira-Bueno, J., Quintanilla-Gozalo, A., Perez-Perez, V., Espi-Felgueroso, A., Alvarez-Garcia, G., Collantes-Fernandez, E. and Ortega-Mora, L. M.** (2002). Evaluation by different diagnostic techniques of bovine abortion associated with *Neospora caninum* in Spain. *Veterinary Parasitology* **111**, 143–152.

- Savva, D., Morris, J. C., Johnson, J. D. and Holliman, R. E.** (1990). Polymerase chain reaction for detection of *Toxoplasma gondii*. *Journal of Medical Microbiology* **32**, 25–31.
- Speer, C. A., Dubey, J., McAllister, M. M. and Blixt, J. A.** (1999). Comparative ultrastructure of tachyzoites, bradyzoites and tissue cysts of *Neospora caninum* and *Toxoplasma gondii*. *International Journal for Parasitology* **29**, 1509–1519.
- Sundermann, C. A. and Estridge, B. H.** (1999). Growth and competition between *Neospora caninum* and *Toxoplasma gondii* in vitro. *International Journal for Parasitology* **29**, 1725–1732.
- Tenter, A. M., Heckeroth, A. R. and Weiss, L. M.** (2000). *Toxoplasma gondii*: from animals to humans. *International Journal for Parasitology* **30**, 1217–1258.
- Terry, R. S., Smith, J. E., Duncanson, P. and Hide, G.** (2001). MGE-PCR: a novel approach to the analysis of *Toxoplasma gondii* strain differentiation using mobile genetic elements. *International Journal for Parasitology* **31**, 155–161.
- Wastling, J. M., Nicoll, S. and Buxton, D.** (1993). Comparison of 2 gene Amplification methods for the detection of *Toxoplasma gondii* in experimentally infected sheep. *Journal of Medical Microbiology* **38**, 360–365.
- Williams, R. H., Morley, E. K., Hughes, J. M., Duncanson, P., Terry, R. S., Smith, J. E. and Hide, G.** (2005). High levels of congenital transmission of *Toxoplasma gondii* in longitudinal and cross-sectional studies on sheep farms provides evidence of vertical transmission in ovine hosts. *Parasitology* **130**, 301–307.
- Yamage, M., Flechner, O. and Gottstein, B.** (1996). *Neospora caninum*: specific oligonucleotide primers for the detection of brain “cyst” DNA of experimentally-infected nude mice by the polymerase chain reaction (PCR). *Journal of Parasitology* **82**, 272–279.