

THE PRODUCT OF OPERATORS WITH CLOSED RANGE AND AN EXTENSION OF THE REVERSE ORDER LAW

SAICHI IZUMINO

(Received March 7, 1981)

1. Introduction. Let A be a (bounded linear) operator on a Hilbert space H . If A has closed range, then there is a unique operator A^\dagger called the Moore-Penrose inverse or generalized inverse of A , which satisfies the following four identities [2, p. 321]:

$$AA^\dagger A = A, \quad A^\dagger AA^\dagger = A^\dagger, \quad (A^\dagger A)^* = A^\dagger A \quad \text{and} \\ (AA^\dagger)^* = AA^\dagger.$$

We denote by (CR) the set of all operators on H with closed range (or equivalently, that of all operators with Moore-Penrose inverses). For two operators A and B in (CR) , one problem is to find the condition under which the product AB is in (CR) . Bouldin [3] [5] gave a geometric characterization of the condition in terms of the angle between two linear subspaces, and recently Nikaido [16] showed a topological characterization of it (for Banach space operators). Another problem is to represent the Moore-Penrose inverse $(AB)^\dagger$ in a reasonable form, that is, to generalize the reverse order law $(AB)^{-1} = B^{-1}A^{-1}$ for invertible operators. Many authors [1], [4], [6], [9], [10], [18]-[20], etc. (some of them in the setting of matrices) studied this problem. Barwick and Gilbert [1], Bouldin [4] [6], Galperin and Waksman [9], etc. proved some necessary and sufficient conditions which guarantee the "generalized" reverse order law $(AB)^\dagger = B^\dagger A^\dagger$ holds.

In this paper we shall treat the product of two operators with closed range. In Section 2 we shall show some norm inequalities for the product to have closed range, which enable us to refine the results in [3] and [16]. In Section 3, using our result in [12], we shall present an extension of the (generalized) reverse order law, and extend some main results in [1], [4], [6] and [9].

Throughout this paper all operators are bounded linear. A projection is a selfadjoint idempotent operator, and it is an orthogonal projection onto a closed linear subspace of H . For projections P and Q onto the closed linear subspaces M and N , we write, in lattice theoretic notations, P^\perp , $P \vee Q$ and $P \wedge Q$ the projections onto the orthocomplement M^\perp of

M , the norm closure $(M + N)^-$ of $M + N$ and the intersection $M \cap N$, respectively. For an operator A we shall denote by $\ker A$ and $\text{ran } A$ the kernel and the range, respectively. The lower bound $\gamma(A)$ of A ($A \neq 0$) is defined by

$$\gamma(A) = \inf \{ \|Ax\| : x \in (\ker A)^\perp, \|x\| = 1 \}.$$

It is well-known [2, p. 311] that $A \in (CR)$ if and only if $\gamma(A) > 0$, and in this case [2, p. 325]

$$(1.1) \quad \gamma(A) = \|A^\dagger\|^{-1}.$$

If $A \in (CR)$, then $A^* \in (CR)$ and $A^{*(\dagger)} = A^{\dagger(*)}$ [2, p. 320] ($A^{\alpha(\beta)}$ means $(A^\alpha)^\beta$). Moreover, $A^\dagger A (= A^* A^{*(\dagger)})$ and $AA^\dagger (= A^{*(\dagger)} A^*)$ are the projections onto $(\ker A)^\perp (= \text{ran } A^*)$ and $\text{ran } A (= (\ker A^*)^\perp)$, respectively. For further basic properties of Moore-Penrose inverses we refer to [2], [11] or [15].

We would like to express our thanks to the referee for his kind advice.

2. The closedness of range of the product operator. An operator $A \in (CR)$ is easily characterized as an operator satisfying $AXA = A$ for some operator X (cf. $AA^\dagger A = A$ for such an A), i.e., a relatively regular element of the operator algebra on H . Hence by [17, Result 3.1] (cf. [14, Theorem 1]) on relative regularity we, at once, have the following proposition, which shows that the problem on the closedness of $\text{ran } AB$ is reduced to that of $\text{ran } A^\dagger ABB^\dagger$, the range of the product of two projections.

PROPOSITION 2.1. *Let $A, B \in (CR)$. Then $AB \in (CR)$ if and only if $A^\dagger ABB^\dagger \in (CR)$.*

The following result shows a norm characterization of the closedness of $\text{ran } PQ$ for two projections P and Q .

PROPOSITION 2.2. *Let P and Q be projections. If $PQ \neq 0$, then*

$$(2.1) \quad \gamma(PQ)^2 + \|P^\perp Q(P \vee Q^\perp)\|^2 = 1.$$

Hence (even if $PQ = 0$) $PQ \in (CR)$ if and only if

$$(2.2) \quad \|P^\perp Q(P \vee Q^\perp)\| < 1.$$

PROOF. Since $\text{ran } QP \subset \text{ran } Q(Q^\perp \vee P) \subset (\text{ran } QP)^- = (\ker PQ)^\perp$, we have

$$(2.3) \quad (\ker PQ)^\perp = \text{ran } Q(Q^\perp \vee P).$$

Let $x \in (\ker PQ)^\perp$ and $\|x\| = 1$. Then, since $x = Q(Q^\perp \vee P)x = Qx$, we

have $\|PQx\|^2 + \|P^\perp Q(P \vee Q^\perp)\|^2 \geq \|PQx\|^2 + \|P^\perp Qx\|^2 = \|Qx\|^2 = 1$. By definition, the infimum of $\|PQx\|$ is $\gamma(PQ)$. Hence we have $\gamma(PQ)^2 + \|P^\perp Q(P \vee Q^\perp)\|^2 \geq 1$. To show the converse inequality, note $\gamma(PQ) \leq \|PQx\|$ ($x \in (\ker PQ)^\perp$, $\|x\| = 1$). Hence $\gamma(PQ)^2 + \|P^\perp Qx\|^2 \leq \|PQx\|^2 + \|P^\perp Qx\|^2 = 1$. Since the supremum of $\|P^\perp Qx\|$ is $\|P^\perp Q(P \vee Q^\perp)\|$, we obtain $\gamma(PQ)^2 + \|P^\perp Q(P \vee Q^\perp)\|^2 \leq 1$. Now, the equivalence $PQ \in (CR) \Leftrightarrow (2.2)$ (between $PQ \in (CR)$ and (2.2)) is clear if $PQ \neq 0$. If $PQ = 0$, then $Q(P \vee Q^\perp) = 0$ (say, by (2.3)), so that (2.2) is clear. q.e.d.

By (1.1) we easily see $\gamma(A) = \gamma(A^*)$ for an operator $A \neq 0$, in particular, $\gamma(PQ) = \gamma(QP)$ ($PQ \neq 0$). Hence by (2.1) we have (even if $PQ = 0$)

$$(2.4) \quad \|P^\perp Q(P \vee Q^\perp)\| = \|Q^\perp P(Q \vee P^\perp)\| .$$

Between two closed linear subspaces M and N we define the angle $\alpha(M, N)$ ($0 \leq \alpha(M, N) \leq \pi/2$) as the arccosine of

$$\sup \{ |(x, y)| : \|x\| = \|y\| = 1, x \in M, y \in N \} ,$$

and $\alpha(M, N) = \pi/2$ when either M or N is $\{0\}$.

Suppose $A, B \in (CR)$, and write $P = A^\dagger A$, $Q = BB^\dagger$. Then $P^\perp(P \vee Q^\perp) = P^\perp \wedge (P^\perp \wedge Q)^\perp$ is the projection onto $L := \ker A \cap (\ker A \cap \text{ran } B)^\perp$. If neither L nor $\text{ran } B$ is $\{0\}$, then

$$\begin{aligned} \|P^\perp Q(P \vee Q^\perp)\| &= \|QP^\perp(P \vee Q^\perp)\| \\ &= \sup \{ |(P^\perp(P \vee Q^\perp)x, Qy)| : \|x\| = \|y\| = 1 \} \\ &= \sup \{ |(x, y)| : \|x\| = \|y\| = 1, x \in L, y \in \text{ran } B \} . \end{aligned}$$

Hence, by Propositions 2.1 and 2.2 we have the following result due to Bouldin [3] (cf. [5]).

COROLLARY 2.3 [3, Theorem]. *Let $A, B \in (CR)$. Then $AB \in (CR)$ if and only if $\alpha(\ker A \cap (\ker A \cap \text{ran } B)^\perp, \text{ran } B) > 0$.*

For another characterization of the closedness of $\text{ran } PQ$, we have

PROPOSITION 2.4. *Let P and Q be projections. If $PQ \neq 0$, then*

$$(2.5) \quad \gamma(PQ) \geq \gamma(P^\perp + Q) \geq (1 - \|P^\perp Q(P \vee Q^\perp)\|)^2 .$$

Hence (even if $PQ = 0$) $PQ \in (CR)$ if and only if $P^\perp + Q \in (CR)$.

PROOF. Note first that $(\ker QP)^\perp = \text{ran } P(P^\perp \vee Q)$, $(\ker (P^\perp + Q))^\perp = \text{ran } (P^\perp \vee Q)$, and that both the subspaces are not $\{0\}$. Let $x \in (\ker QP)^\perp$ and $\|x\| = 1$. Then $x = P(P^\perp \vee Q)x = Px$ and

$$\|QPx\| = \|(P^\perp + Q)Px\| = \|(P^\perp + Q)x\| \geq \gamma(P^\perp + Q) .$$

The last inequality follows from the fact $x = (P^\perp \vee Q)x \in (\ker (P^\perp + Q))^\perp$.

Hence we have $\gamma(QP) \geq \gamma(P^\perp + Q)$. Since $\gamma(PQ) = \gamma(QP)$, we have the left hand side inequality of (2.5). Next, note $\langle (P^\perp + Q)x, x \rangle \geq \langle (P^\perp Q^\perp P^\perp + Q)x, x \rangle$ for any $x \in H$ and $P^\perp Q^\perp P^\perp + Q = 1 - (Q^\perp P + PQ^\perp) + PQ^\perp P$. Hence, if $x = (P^\perp \vee Q)x$ and $\|x\| = 1$ then

$$\begin{aligned} \|(P^\perp + Q)x\| &\geq \langle (P^\perp + Q)x, x \rangle \geq 1 - 2 \operatorname{Re} \langle Q^\perp P x, x \rangle + \langle PQ^\perp P x, x \rangle \\ &\geq 1 - 2\|Q^\perp P x\| + \|Q^\perp P x\|^2 = (1 - \|Q^\perp P x\|)^2 \\ &\geq (1 - \|Q^\perp P(P^\perp \vee Q)\|)^2. \end{aligned}$$

Hence $\gamma(P^\perp + Q) \geq (1 - \|Q^\perp P(Q \vee P^\perp)\|)^2$. By (2.4) this implies the right hand side inequality of (2.5). Now the equivalence $PQ \in (CR) \Leftrightarrow P^\perp + Q \in (CR)$ is clear by (2.5) and (2.2) if $PQ \neq 0$. If $PQ = 0$, then $\operatorname{ran}(P^\perp + Q) = \operatorname{ran} P^\perp(1 + Q) = \operatorname{ran} P^\perp$, so that $P^\perp + Q \in (CR)$. q.e.d.

Before an application we remark that $A \in (CR)$ if and only if $AA^* \in (CR)$. This is seen by the facts $\operatorname{ran} AA^* \subset \operatorname{ran} A \subset (\operatorname{ran} AA^*)^-$, and $\operatorname{ran} A = \operatorname{ran} A \cdot (A^\dagger A)^* = \operatorname{ran} AA^* A^{\dagger(*)} \subset \operatorname{ran} AA^* \subset \operatorname{ran} A$ for $A \in (CR)$.

The equivalence (1) \Leftrightarrow (3) of the following corollary was shown by Nikaido [16, Corollary 1].

COROLLARY 2.5. *Let $A, B \in (CR)$. Write $P = A^\dagger A$ and $Q = BB^\dagger$. Then the following conditions are equivalent.*

- (1) $AB \in (CR)$.
- (2) $P^\perp + Q \in (CR)$.
- (3) $\ker A + \operatorname{ran} B$ is closed.

PROOF. (1) \Leftrightarrow (2) By Propositions 2.1 and 2.4.

(2) \Leftrightarrow (3) We employ a technique in [7, Theorem 2.2]. Let $T = \begin{Bmatrix} P^\perp & Q \\ 0 & 0 \end{Bmatrix}$ be a operator matrix on the product Hilbert space $H \oplus H$. Then $\operatorname{ran} T = (\operatorname{ran} P^\perp + \operatorname{ran} Q) \oplus \{0\}$ and $\operatorname{ran} TT^* = \operatorname{ran}(P^\perp + Q) \oplus \{0\}$. Hence by the above remark we have the desired equivalence. q.e.d.

COROLLARY 2.6. *Let P and Q be projections. Then $\operatorname{ran} P + \operatorname{ran} Q$ is closed if and only if $\|PQ(P^\perp \vee Q^\perp)\| < 1$.*

PROOF. By Corollary 2.5 and Proposition 2.4. q.e.d.

For a pair of two closed linear subspaces M and N , the gap $g(M, N)$ is defined (cf. [13, p. 219]) by

$$g(M, N) = \inf \{d(x, N)/d(x, M \cap N) : x \in M \setminus N\},$$

where $d(x, L)$ is the distance from x to L . We set $g(M, N) = 1$ when $M \subset N$. Let P and Q be the projections onto M and N , respectively. Then by a simple calculation we have $g(M, N) = \gamma(Q^\perp P)$ ($M \not\subset N$), or by

(2.1) (even if $M \subset N$)

$$g(M, N) = (1 - \|PQ(P^\perp \vee Q^\perp)\|^2)^{1/2}.$$

Clearly, Corollary 2.6 says that $g(M, N) > 0$ if and only if $M + N$ is closed, which is a well-known result [13, IV, Theorem 4.3] (on a Banach space).

3. The reverse order law. We state a result which we proved in [12].

LEMMA 3.1 [12, Lemmas 2.1 and 3.2]. *Let $A \in (CR)$, and let R be a projection commuting with $A^\dagger A$. Then $AR \in (CR)$, $C := 1 - ARA^\dagger + ARA^*$ is invertible and*

$$(3.1) \quad (AR)(AR)^\dagger = C^{-1}ARA^*.$$

Using the above lemma we have

LEMMA 3.2. *Let $A, B, AB \in (CR)$. Write $P = A^\dagger A$ and $Q = BB^\dagger$. Then $C := 1 - A(P^\perp \vee Q)A^\dagger + A(P^\perp \vee Q)A^*$ is invertible, and*

$$(3.2) \quad (AB)(AB)^\dagger = C^{-1}A(P^\perp \vee Q)A^*.$$

PROOF. Put $R = P^\perp \vee Q$. Since $\text{ran } AB = \text{ran } AQ \subset \text{ran } AR \subset (\text{ran } AR)^\perp = (\text{ran } AQ)^\perp = \text{ran } AB$, we have $\text{ran } AB = \text{ran } AR$, i.e., $(AB)(AB)^\dagger = (AR)(AR)^\dagger$. Since R commutes with $P = A^\dagger A$, we have, by Lemma 3.1, the required assertions. q.e.d.

COROLLARY 3.3. *Let P and Q be projections. If $PQ \in (CR)$, then*

$$(3.3) \quad (PQ)(PQ)^\dagger = P(P^\perp \vee Q), \quad (PQ)^\dagger(PQ) = Q(Q^\perp \vee P).$$

We remark that the second identity of (3.3) can be also obtained from (2.3).

For the Moore-Penrose inverse of $(PQ)^\dagger$, we have the following result which is considered as an extension of [10, Theorem 3].

LEMMA 3.4. *Let P and Q be projections. If $PQ \in (CR)$, then $R := 1 - (P \vee Q^\perp)Q + PQ$ is invertible and*

$$(3.4) \quad (PQ)^\dagger = R^{-1}P(P^\perp \vee Q).$$

PROOF. Since $R = 1 - (P \vee Q^\perp - P)Q = 1 - (P \vee Q^\perp)P^\perp Q$ and since $\|(P \vee Q^\perp)P^\perp Q\| < 1$ by (2.1), we see that R is invertible. By (3.3) we see $(P \vee Q^\perp)Q(PQ)^\dagger = (PQ)^\dagger(PQ)(PQ)^\dagger = (PQ)^\dagger$. Hence we have

$$R(PQ)^\dagger = (PQ)^\dagger - (P \vee Q^\perp)Q(PQ)^\dagger + PQ(PQ)^\dagger = (PQ)(PQ)^\dagger = P(P^\perp \vee Q).$$

This implies the desired identity.

q.e.d.

Now we state the main theorem of this section.

THEOREM 3.5. *Let $A, B \in (CR)$. If $AB \in (CR)$, then*

$$\begin{aligned} (AB)^\dagger &= (AB)^\dagger(AB) \cdot B^\dagger \cdot (PQ)^\dagger \cdot A^\dagger \cdot (AB)(AB)^\dagger \\ &= f(B^*, Q^\perp \vee P) \cdot B^\dagger \cdot \{1 - (P \vee Q^\perp)Q + PQ\}^{-1} \\ &\quad \times (P^\perp \vee Q) \cdot A^\dagger \cdot f(A, P^\perp \vee Q), \end{aligned}$$

where $P = A^\dagger A$, $Q = BB^\dagger$ and $f(S, T) = (1 - STS^\dagger + STS^*)^{-1}STS^*$.

PROOF. Note $PQ \in (CR)$ by Proposition 2.1. The first identity is obtained from the fact:

$$\begin{aligned} (AB)B^\dagger(PQ)^\dagger A^\dagger(AB) &= A(A^\dagger ABB^\dagger)(A^\dagger ABB^\dagger)^\dagger(A^\dagger ABB^\dagger)B \\ &= A(A^\dagger ABB^\dagger)B = AB. \end{aligned}$$

The second identity is shown by (3.2), (3.4) and the identity $(AB)^\dagger(AB) = (B^*A^*)(B^*A^*)^\dagger$. q.e.d.

In each of the following two corollaries, $(AB)^\dagger$ is represented by a rational function in $A, A^\dagger, B, B^\dagger$ and their adjoints under a certain condition which is satisfied for invertible operators. Hence our theorem is, in a sense, a reasonable extension of the reverse order law.

COROLLARY 3.6. *Let $A, B, AB \in (CR)$. If $P := A^\dagger A$ and $Q := BB^\dagger$ commute, then*

$$(AB)^\dagger = f(B^*, P)B^\dagger A^\dagger f(A, Q) \quad (f \text{ is defined in Theorem 3.5}).$$

PROOF. Since P and Q commute, we see that PQ is a projection. Hence $(PQ)^\dagger = PQ (= QP)$, because $R^\dagger = R$ for a projection R . Since $(AB)(AB)^\dagger = (AQ)(AQ)^\dagger$, and since Q commutes with $A^\dagger A = P$, we have, by (3.1), $(AB)(AB)^\dagger = f(A, Q)$. Similarly we have $(AB)^\dagger(AB) = f(B^*, P)$. Hence by the first identity of Theorem 3.5 we have the desired representation of $(AB)^\dagger$. q.e.d.

We remark that the assumption $AB \in (CR)$ is not needed in Corollary 3.6. For, if P and Q commute then PQ is a projection and $PQ \in (CR)$, so that $AB \in (CR)$ (say, by Proposition 2.1).

COROLLARY 3.7. *Let $A, B, AB \in (CR)$. If $P^\perp \vee Q = P \vee Q^\perp = 1$, i.e., $\ker A$ and $\text{ran } B$ are complementary, then*

$$(AB)^\dagger = B^\dagger(1 - Q + PQ)^{-1}A^\dagger.$$

PROOF. By assumption $f(A, P^\perp \vee Q) = f(A, 1) = (1 - AA^\dagger + AA^*)^{-1}AA^*$. Since $(1 - AA^\dagger + AA^*)AA^\dagger = AA^*$ (cf. $A^*AA^\dagger = A^*$), we have $f(A, 1) = AA^\dagger$. Similarly we have $f(B^*, Q^\perp \vee P) = B^\dagger B$. Hence by the second

identity of Theorem 3.5 we have the required equation. q.e.d.

The following result was essentially shown in [16, Proposition 1] (for Banach space operators).

COROLLARY 3.8. *Let $A, B \in (CR)$ and let $AB \neq 0$. Then*

$$(3.5) \quad \gamma(AB) \geq \gamma(A)\gamma(B)\gamma(PQ) .$$

PROOF. If $AB \in (CR)$, then by Theorem 3.5 $\|(AB)^\dagger\| \leq \|B^\dagger\| \|(PQ)^\dagger\| \|A^\dagger\|$. Hence by (1.1) we obtain (3.5). If $AB \notin (CR)$, then $PQ \notin (CR)$. Hence (3.5) is clear. q.e.d.

The next two propositions extend (or refine) Bouldin [4, Theorem 3.1] [6, Theorem 3.3], Barwick and Gilbert [1, Theorems 1 and 2], Shinozaki and Sibuya [18, Propositions 3.2 and 4.3].

First we state a useful lemma for our discussion.

LEMMA 3.9 [8, Theorem 2]. *Let T be an idempotent operator with $\|T\| \leq 1$. Then T is a projection.*

PROPOSITION 3.10. *Let $A, B, AB \in (CR)$. Then the following conditions are equivalent.*

- (1) $A^\dagger A$ commutes with BB^* .
- (2) $(AB)^\dagger(AB) = B^\dagger A^\dagger AB$.
- (3) $C := 1 - A^{*(\dagger)}BB^\dagger A^* + ABB^\dagger A^*$ is invertible, and

$$(AB)^\dagger = B^\dagger A^* C^{-1} .$$

PROOF. (1) \Rightarrow (2) Since $A^* = A^\dagger A A^*$ (and $B^* = B^\dagger B B^*$), we have

$$\begin{aligned} (AB)^\dagger(AB) &= (AB)^*(AB)^{\dagger(*)} = B^* A^*(AB)^{\dagger(*)} = B^\dagger B B^* \cdot A^\dagger A A^* \cdot (AB)^{\dagger(*)} \\ &= B^\dagger A^\dagger A B B^* A^* (AB)^{\dagger(*)} = B^\dagger A^\dagger (AB) (AB)^* (AB)^{\dagger(*)} \\ &= B^\dagger A^\dagger (AB) (AB)^\dagger (AB) = B^\dagger A^\dagger AB . \end{aligned}$$

(2) \Rightarrow (3) We first show that $P := A^\dagger A$ and $Q := BB^\dagger$ commute. Since $AB = (AB)(AB)^\dagger(AB) = AB \cdot B^\dagger A^\dagger AB$, we have $PQ = A^\dagger A B B^\dagger = A^\dagger \cdot A B B^\dagger A^\dagger A B \cdot B^\dagger = (PQ)^2$. Besides, clearly $\|PQ\| \leq 1$. Hence by Lemma 3.9 PQ is a projection, so that P and Q commute. Now by Corollary 3.6 we see $(AB)^\dagger = f(B^*, P)B^\dagger A^\dagger f(A, Q)$. Since $f(B^*, P) = (AB)^\dagger(AB) = B^\dagger A^\dagger AB$, and since $f(A, Q) = f(A, Q)^* = AQA^*C^{-1}$, we have $(AB)^\dagger = B^\dagger A^\dagger A B B^\dagger A^\dagger A Q A^* C^{-1} = B^\dagger A^* C^{-1}$.

(3) \Rightarrow (1) Let $(AB)^\dagger = B^\dagger A^* C^{-1}$. Then $(AB)^\dagger C = B^\dagger A^*$ or

$$(3.6) \quad (AB)^\dagger - (AB)^\dagger A^{\dagger(*)} Q A^* + (AB)^\dagger A Q A^* = B^\dagger A^* .$$

Since $(AB)(AB)^\dagger A Q A^* = A B B^\dagger A^*$, multiplying (3.6) by AB from the left, we have $(AB)(AB)^\dagger - (AB)(AB)^\dagger A^{\dagger(*)} Q A^* + A B B^\dagger A^* = A B B^\dagger A^*$. Hence

$$(3.7) \quad (AB)(AB)^\dagger = (AB)(AB)^\dagger A^{\dagger(*)}QA^* .$$

If we multiply (3.7) by $(AB)^\dagger$ from the left, then we obtain $(AB)^\dagger = (AB)^\dagger A^{\dagger(*)}QA^*$. Hence by (3.6) we see

$$(3.8) \quad (AB)^\dagger AQA^* = B^\dagger A^* .$$

Now, if we assume that P and Q commute, then by (3.8)

$$\begin{aligned} PBB^* &= PQBB^* = QPBB^* = BB^\dagger A^\dagger ABB^* = BB^\dagger A^* A^{\dagger(*)} BB^* \\ &= B \cdot (AB)^\dagger AQA^* \cdot A^{\dagger(*)} BB^* = B(AB)^\dagger AQA^\dagger ABB^* \\ &= B(AB)^\dagger (AB)B^* . \end{aligned}$$

This shows that PBB^* is selfadjoint. Hence P and BB^* commute, which is the assertion (1). To see that P and Q commute, take the adjoints in (3.7). Then we have $(AB)(AB)^\dagger = AQA^\dagger(AB)(AB)^\dagger$. Multiplying by AB from the right, we have $AB = AQA^\dagger AB$. By this identity we easily see $PQ = (PQ)^\dagger$, so that P and Q commute (cf. Proof of (2) \Rightarrow (3)). q.e.d.

Similarly to Proposition 3.10 we have:

PROPOSITION 3.10'. *Let $A, B, AB \in (CR)$. Then the following conditions are equivalent.*

- (1) BB^\dagger commutes with A^*A .
- (2) $(AB)(AB)^\dagger = ABB^\dagger A^\dagger$.
- (3) $D := 1 - B^*A^\dagger AB^{*(\dagger)} + B^*A^\dagger AB$ is invertible, and

$$(AB)^\dagger = D^{-1}B^*A^\dagger .$$

PROOF. Replace, in Proposition 3.10, A and B by B^* and A^* respectively, and take the adjoints. q.e.d.

COROLLARY 3.11 [6, Theorem 3.3]. *Let $A, B, AB \in (CR)$. Then the following conditions are equivalent.*

- (1) $A^\dagger A$ commutes with BB^* and BB^\dagger commutes with A^*A .
- (2) $(AB)^\dagger(AB) = B^\dagger A^\dagger AB$ and $(AB)(AB)^\dagger = ABB^\dagger A^\dagger$.
- (3) $(AB)^\dagger = B^\dagger A^\dagger$.

PROOF. The equivalence (1) \Leftrightarrow (2) is clear by Propositions 3.10 and 3.10'. If (2) is assumed, then $A^\dagger A$ and BB^\dagger commute (cf. Proof of Proposition 3.10 (2) \Rightarrow (3)). Hence $(AB)^\dagger = (AB)^\dagger(AB)(AB)^\dagger = B^\dagger A^\dagger AB(AB)^\dagger = B^\dagger A^\dagger ABB^\dagger A^\dagger = B^\dagger A^\dagger$, which is the assertion (3). The implication (3) \Rightarrow (2) is clear. q.e.d.

The following proposition is a Hilbert space version of a result due to Galperin and Waksman ([9, Theorem 2]).

PROPOSITION 3.12. *Let $A, B, AB \in (CR)$. Then the following conditions*

are equivalent.

- (1) $\text{ran } B^*A^* = \text{ran } B^*A^*$ and $\text{ran } A^{*(\dagger)}B = \text{ran } AB$.
- (2) $(AB)^\dagger = B^\dagger(A^*ABB^\dagger)^\dagger A^\dagger$.

PROOF. Note first that B^*A^* , $B^\dagger A^\dagger$, $A^{*(\dagger)}B \in (CR)$, say, by Proposition 2.1. Write $P = A^\dagger A$ and $Q = BB^\dagger$, and let $X = B^\dagger(PQ)^\dagger A^\dagger$. Then clearly $XABX = X$, so that $X \in (CR)$. Next we want to show

$$(3.9) \quad \text{ran } X = \text{ran } B^\dagger A^\dagger \quad \text{and} \quad \text{ran } X^* = \text{ran } A^{*(\dagger)}B .$$

Since $(PQ)^\dagger P = (PQ)^\dagger$ by (3.4), and since $\text{ran } B^\dagger(Q^\perp \vee P) = \text{ran } B^\dagger P$ (cf. Proof of Lemma 3.2), we have

$$\begin{aligned} \text{ran } X &= \text{ran } B^\dagger(PQ)^\dagger A^\dagger = \text{ran } B^\dagger(PQ)^\dagger P = \text{ran } B^\dagger(PQ)^\dagger = \text{ran } B^\dagger(PQ)^\dagger(PQ) \\ &= \text{ran } B^\dagger(Q^\perp \vee P) = \text{ran } B^\dagger P = \text{ran } B^\dagger A^\dagger . \end{aligned}$$

Similarly we have the other identity of (3.9). Now, if we assume (1), then by (3.9) we obtain

$$(3.10) \quad \text{ran } X = \text{ran } B^*A^* \quad \text{and} \quad \text{ran } X^* = \text{ran } AB ,$$

or equivalently

$$(3.11) \quad XX^\dagger = (AB)^\dagger(AB) \quad \text{and} \quad X^\dagger X = (AB)(AB)^\dagger .$$

Hence $X = XX^\dagger XX^\dagger X = (AB)^\dagger(AB) \cdot X \cdot (AB)(AB)^\dagger = (AB)^\dagger(AB)(AB)^\dagger = (AB)^\dagger$, which is the assertion (2). Conversely, if we assume (2), i.e., $X = (AB)^\dagger$, then clearly (3.11) and hence (3.10) are valid. Hence by (3.9) we have the assertion (1). q.e.d.

We remark that the condition $P^\perp \vee Q = P \vee Q^\perp = 1$ ($P = A^\dagger A$, $Q = BB^\dagger$) taken in Corollary 3.7 implies the assertion (2) (hence also (1)) of the above proposition.

The following result adds to Corollary 3.11 another condition in order that $(AB)^\dagger = B^\dagger A^\dagger$ holds.

COROLLARY 3.13 (cf. [9, Theorem 3]). *Let $A, B, AB \in (CR)$. Then $(AB)^\dagger = B^\dagger A^\dagger$ if and only if*

$$(3.12) \quad A^\dagger A \text{ and } BB^\dagger \text{ commute, and (1) (or equivalently (2)) of Proposition 3.12 holds.}$$

PROOF. If $(AB)^\dagger = B^\dagger A^\dagger$ then $(AB)^\dagger AB = B^\dagger A^\dagger AB$, so that $A^\dagger A$ and BB^\dagger commute. Since $(A^*ABB^\dagger)^\dagger = A^*ABB^\dagger = BB^\dagger A^\dagger A$, we have $B^\dagger(A^*ABB^\dagger)^\dagger A^\dagger = B^\dagger A^\dagger = (AB)^\dagger$, the assertion (2) of Proposition 3.12. Conversely, if (3.12) is assumed then $(AB)^\dagger = B^\dagger(A^*ABB^\dagger)^\dagger A^\dagger = B^\dagger BB^\dagger A^\dagger AA^\dagger = B^\dagger A^\dagger$, as desired. q.e.d.

REFERENCES

- [1] D. T. BARWICK AND J. D. GILBERT, On generalizations of reverse order law, *SIAM J. Appl. Math.* 27 (1974), 326-330.
- [2] A. BEN-ISRAEL AND T. N. E. GREVILLE, *Generalized Inverses: Theory and Applications*, Wiley, New York, 1974.
- [3] R. BOULDIN, The product of operators with closed range, *Tôhoku Math. J.* 25 (1973), 359-363.
- [4] R. BOULDIN, The pseudo-inverse of a product, *SIAM J. Appl. Math.* 24 (1973), 489-495.
- [5] R. BOULDIN, Closed range and relativity for products, *J. Math. Anal. Appl.* 61 (1977), 397-403.
- [6] R. BOULDIN, Generalized inverses and factorizations, to appear in *Recent Application of Generalized Inverses* edited by M. Z. Nashed.
- [7] P. A. FILLMORE AND J. P. WILLIAMS, On operator ranges, *Advances in Math.* 7 (1971), 254-281.
- [8] T. FURUTA AND R. NAKAMOTO, Some theorems on certain contraction operators, *Proc. Japan Acad.* 45 (1969), 565-566.
- [9] A. M. GALPERIN AND Z. WAKSMAN, On pseudoinverses of operator products, *Linear Algebra and Appl.* 33 (1980), 123-131.
- [10] T. N. E. GREVILLE, Solutions of the matrix equation $XAX = X$, and relations between oblique and orthogonal projections, *SIAM J. Appl. Math.* 26 (1974), 823-832.
- [11] C. W. GROETSCH, *Generalized Inverses of Linear Operators: Representation and Approximation*, Marcel Dekker, New York and Basel, 1977.
- [12] S. IZUMINO, Inequalities on operators with closed range, *Math. Japon.* 25 (1980), 423-429.
- [13] T. KATO, *Perturbation Theory for Linear Operators*, Springer-Verlag, Berlin, Heidelberg and New York, 1966.
- [14] J. J. KOLIHA, The product of relatively regular operators, *Commentationes Math. Univ. Carolinae* 16 (1975), 531-539.
- [15] M. Z. NASHED, *Generalized Inverses and Applications*, Academic Press, New York, San Francisco and London, 1976.
- [16] G. NIKAIKO, Remarks on the lower bound of linear operators, *Proc. Japan Acad.* 56 (1980), 321-323.
- [17] S. V. PHADKE AND N. T. TAKARE, Generalized inverses and operator equations, *Linear Algebra and Appl.* 23 (1979), 191-199.
- [18] N. SHINOZAKI AND M. SIBUYA, The reverse order law $(AB)^- = B^-A^-$, *Linear Algebra and Appl.* 9 (1974), 29-40.
- [19] N. SHINOZAKI AND M. SIBUYA, Further results on the reverse order law, *Linear Algebra and Appl.* 27 (1979), 9-16.
- [20] E. A. WIBKER, R. B. HOWE AND J. D. GILBERT, Explicit solutions to the reverse order law, $(AB)^+ = B_m^-A_l^-$, *Linear Algebra and Appl.* 25 (1979), 107-114.

FACULTY OF EDUCATION
 TOYAMA UNIVERSITY
 TOYAMA, 930
 JAPAN