

The property of universality for some monoid algebras over non-commutative rings

Elena P. Cojuhari

Abstract. We define on an arbitrary ring A a family of mappings $(\sigma_{x,y})$ subscripted with elements of a multiplicative monoid G . The assigned properties allow to call these mappings derivations of the ring A . A monoid algebra of G over A is constructed explicitly, and the universality property of it is shown.

Mathematics subject classification: Primary 16S36; Secondary 13N15, 16S10.

Keywords and phrases: Derivations, monoid algebras, free algebras.

In this note we consider monoid algebras over non-commutative rings. First, we introduce axiomatically a family of mappings $\sigma = (\sigma_{x,y})$ defined on a ring A and subscripted with elements of a multiplicative monoid G . Due to their assigned properties these mappings can be called derivations of A . Next, we construct a monoid algebra $A\langle G \rangle$ by means of the family σ , and the universality of it is shown.

1. Let A be a ring (in general non-commutative) and G a multiplicative monoid. Throughout the paper we consider $1 \neq 0$ (where 0 is the null element of A , and 1 is the unit element for multiplication), the unit element of G is denoted by e . We introduce a family of mappings of A into itself by the following assumption.

(A) For each $x \in G$ there exists a unique family $\sigma_x = (\sigma_{x,y})_{y \in G}$ of mappings $\sigma_{x,y} : A \rightarrow A$ such that $\sigma_{x,y} = 0$ for almost all $y \in G$ (here and thereafter, almost all will mean all but a finite number, that is, $\sigma_{x,y} \neq 0$ only for a finite set of $y \in G$) and for which the following properties are fulfilled:

- (i) $\sigma_{x,y}(a + b) = \sigma_{x,y}(a) + \sigma_{x,y}(b)$ ($a, b \in A; x, y \in G$);
- (ii) $\sigma_{x,y}(ab) = \sum_{z \in G} \sigma_{x,z}(a)\sigma_{z,y}(b)$ ($a, b \in A; x, y \in G$);
- (iii) $\sigma_{xy,z} = \sum_{uv=z} \sigma_{x,u} \circ \sigma_{y,v}$ ($x, y, z \in G$);
- (iv₁) $\sigma_{x,y}(1) = 0$ ($x \neq y; x, y \in G$);
- (iv₂) $\sigma_{x,x}(1) = 1$ ($x \in G$);
- (iv₃) $\sigma_{e,x}(a) = 0$ ($x \neq e; x \in G$);
- (iv₄) $\sigma_{e,e}(a) = a$ ($a \in A$).

In (ii) the elements are multiplied as in the ring A , but in (iii) the symbol \circ means the composition of maps.

Examples. 1. Let A be a ring and let G be a multiplicative monoid, and let σ be a monoid-homomorphism of G into $End(A)$, i.e. $\sigma(xy) = \sigma(x) \circ \sigma(y)$ ($x, y \in G$) and $\sigma(e) = 1_A$. We define $\sigma_{x,y} : A \rightarrow A$ such that $\sigma_{x,x} = \sigma(x)$ for $x \in G$ and $\sigma_{x,y} = 0$ for $y \neq x$. The properties (i) – (iv₄) of (A) are verified at once.

2. Let A be a ring, and let α be an endomorphism of A and δ be an α -differentiation of A , i.e.

$$\delta(a + b) = \delta(a) + \delta(b), \delta(ab) = \delta(a)b + \alpha(a)\delta(b)$$

for every $a, b \in A$. Denote by G the monoid of elements x_n ($n = 0, 1, \dots$) endowed with the law of composition defined by $x_n x_m = x_{n+m}$ ($n, m = 0, 1, \dots; x_0 := e$). We write σ_{nm} instead of σ_{x_n, x_m} by defining $\sigma_{nm} : A \rightarrow A$ as the following mappings $\sigma_{00} = 1_A, \sigma_{10} = \delta, \sigma_{11} = \alpha, \sigma_{nm} = 0$ for $m > n$ and $\sigma_{nm} = \sum_{j_1 + \dots + j_n = m} \sigma_{1j_1} \circ \dots \circ \sigma_{1j_n}$ ($m = 0, 1, \dots, n; n = 1, 2, \dots$), where $j_k = 0, 1$ ($k = 1, \dots, n$). The family $\sigma = (\sigma_{nm})$ satisfies the axioms (i) – (iv₄) of (A).

2. Next, we consider an algebra $A\langle G \rangle$ connected with the structure of differentiation $\sigma = (\sigma_{x,y})$. Let $A\langle G \rangle$ be the set of all mappings $\alpha : G \rightarrow A$ such that $\alpha(x) = 0$ for almost all $x \in G$. We define the addition in $A\langle G \rangle$ to be the ordinary addition of mappings into the additive group of A and define the operation of A on $A\langle G \rangle$ by the map $(a, \alpha) \rightarrow a\alpha$ ($a \in A$), where $(a\alpha)(x) = a\alpha(x)$ ($x \in G$). Note that, in respect to these operations, $A\langle G \rangle$ forms a left module over A . Following notations made in [1] we write an element $\alpha \in A\langle G \rangle$ as a sum $\alpha = \sum_{x \in G} a_x \cdot x$, where by $a \cdot x$ ($a \in A, x \in G$) is denoted the mapping whose value at x is a and 0 at elements different from x . Certainly, the above sum is taken over almost all $x \in G$. $A\langle G \rangle$ becomes a ring if for elements of the form $a \cdot x$ ($a \in A; x \in G$) we define their product by the rule

$$(a \cdot x)(b \cdot y) = \sum_{z \in G} a\sigma_{x,z}(b) \cdot zy \quad (a, b \in A; x, y \in G)$$

and then extend for $\alpha, \beta \in A\langle G \rangle$ by the property of distributivity. We let

$$\alpha\alpha = \sum_{x \in G} \left(\sum_{y \in G} a_y \sigma_{y,x}(a) \right) \cdot x, \quad (a \in A, \alpha \in A\langle G \rangle)$$

for $a \in A$ and $\alpha \in A\langle G \rangle$, and thus we obtain an operation of A on $A\langle G \rangle$ and in such a way we make $A\langle G \rangle$ into a right A -module. Thus, we may view $A\langle G \rangle$ as an algebra over A .

Remark. Let us consider the situation described in Example 1. Then the law of multiplication in $A\langle G \rangle$ is given as follows

$$\left(\sum_{x \in G} a_x \cdot x \right) \left(\sum_{x \in G} b_x \cdot x \right) = \sum_{x \in G} \sum_{y \in G} a_x \sigma_{x,x}(b_y) \cdot xy.$$

In this case, the monoid algebra $A\langle G \rangle$ represents a crossed product [2, 3] of the multiplicative monoid G over the ring A with respect to the factors $\rho_{x,y} = 1$ ($x, y \in G$). If G is a group, and $\sigma : G \rightarrow \text{End}(A)$ is such that $\sigma(x) = 1_A$ for all $x \in G$, we evidently obtain an ordinary group ring [4] (the commutative case see also [5]).

3. In this subsection we show that $A\langle G \rangle$ is a free G -algebra over A . Let B be another ring. Given a ring-homomorphism $f : A \rightarrow B$ it can be defined on the ring B a structure of A -module, defining the operation of A on B by the map $(a, b) \rightarrow f(a)b$ for all $a \in A$ and $b \in B$. We denote this operation by $a * b$. The axioms for a module are trivially verified. Let now $\varphi : G \rightarrow B$ be a multiplicative monoid-homomorphism. Denote by $\langle B; f, \varphi \rangle$ the module formed by all linear combinations of elements $\varphi(x)$ ($x \in G$) over A in respect to the operation $*$. The axioms for a left A -module are trivially verified.

We assume that the homomorphisms f and φ satisfy the following assumption.

$$(B) \quad \varphi(G)f(A) \subset \langle B; f, \varphi \rangle.$$

Thus, it is postulated that an element $\varphi(x)f(a)$ ($a \in A, x \in G$) can be written as a linear combination of the form $\sum_{b \in B, y \in G} b \varphi(y)$. The coefficients b depend on $\varphi(x), \varphi(y)$ and $f(a)$. To designate this fact we denote the corresponding coefficients by $\sigma_{\varphi(x), \varphi(y)}(f(a))$. Therefore, it can be considered that there are defined a family of mappings $\sigma_{\varphi(x), \varphi(y)} : B \rightarrow B$ such that

$$\varphi(x)f(a) = \sum_{y \in G} \sigma_{\varphi(x), \varphi(y)}(f(a))\varphi(y) \quad (a \in A, x \in G).$$

By these considerations, we may view $\langle B; f, \varphi \rangle$ as a right A -module. In order to make the module $\langle B; f, \varphi \rangle$ to be a ring we require the following additional assumption.

(C) The homomorphisms f and φ are such that the following diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \sigma_{x,y} \uparrow & & \uparrow \sigma_{\varphi(x), \varphi(y)} \\ A & \xrightarrow{f} & B \end{array}$$

is commutative for every $x, y \in G$, i.e. $\sigma_{\varphi(x), \varphi(y)} \circ f = f \circ \sigma_{x,y}$ ($x, y \in G$).

We define multiplication in $\langle B; f, \varphi \rangle$ by the rules

$$\begin{aligned} \left(\sum_{x \in G} a_x * \varphi(x) \right) \left(\sum_{x \in G} b_x * \varphi(x) \right) &= \sum_{x \in G} \sum_{y \in G} (a_x * \varphi(x))(b_y * \varphi(y)), \\ (a_x * \varphi(x))(b_y * \varphi(y)) &= f(a_x) \sum_{z \in G} \sigma_{\varphi(x), \varphi(z)}(f(b_y))\varphi(zy). \end{aligned}$$

The verification that $\langle B; f, \varphi \rangle$ is a ring under the above laws of composition is direct. Thus, we have made $\langle B; f, \varphi \rangle$ into an algebra over A (in general, non-commutative).

Next, we define a category \mathcal{C} whose objects are algebras $\langle B; f, \varphi \rangle$ constructed as above, and whose morphisms between two objects $\langle B; f, \varphi \rangle$ and $\langle B'; f', \varphi' \rangle$ are ring-homomorphisms $h : B \rightarrow B'$ making the diagrams commutative:

$$\begin{array}{ccc} G & \equiv \equiv & G \\ \varphi \downarrow & & \downarrow \varphi' \\ B & \xrightarrow{h} & B' \\ f \uparrow & & \uparrow f' \\ A & \equiv \equiv & A \end{array}$$

The axioms for a category are trivially satisfied. We call a universal object in the category \mathcal{C} a free G -algebra over A , or a free (A, G) -algebra. It turns out that the monoid algebra $A\langle G \rangle$ represents a free (A, G) -algebra. To this end, we observe that the mapping $\varphi_0 : G \rightarrow A\langle G \rangle$ given by $\varphi_0(x) = 1 \cdot x$ ($x \in G$) is a monoid-homomorphism. The mapping φ_0 is embedding of G into $A\langle G \rangle$. In addition, we have a ring-homomorphism $f_0 : A \rightarrow A\langle G \rangle$ given by $f_0(a) = a \cdot e$ ($a \in A$). Obviously, f_0 is also an embedding. We identify $A\langle G \rangle$ with the triple $\langle A\langle G \rangle; f_0, \varphi_0 \rangle$ and in this sense we treat $A\langle G \rangle$ as an object of the category \mathcal{C} . The property of the universality of $A\langle G \rangle$ is formulated by the following assertion.

Theorem 1. *Let A be a ring, and G a multiplicative monoid for which the assumptions (A), (B) and (C) are satisfied. Then for every object $\langle B; f, \varphi \rangle$ of the category \mathcal{C} there exists a unique ring-homomorphism $h : A\langle G \rangle \rightarrow B$ making the following diagram commutative*

$$\begin{array}{ccc} G & \text{====} & G \\ \varphi_0 \downarrow & & \downarrow \varphi \\ A\langle G \rangle & \xrightarrow{h} & B \\ f_0 \uparrow & & \uparrow f \\ A & \text{====} & A \end{array}$$

The relation with the theory of skew polynomial rings [6–8] and with those obtained by Yu. M. Ryabukhin [9] (see also [10]), and further properties of the general derivation mappings $\sigma_{x,y}$ ($x, y \in G$) will be given in a subsequent publication.

References

- [1] LANG S. *Algebra*. Addison Wesley, Reading, Massachusetts, 1970.
- [2] BOVDI A.A. *Crossed products of a semigroup and a ring*. Sibirsk. Mat. Zh., 1963, **4**, p. 481–499.
- [3] PASSMAN D.S. *Infinite crossed products*. Academic Press, Boston, 1989.
- [4] BOVDI A.A. *Group rings*. Kiev UMK VO, 1988 (in Russian).
- [5] KARPILOVSKY G. *Commutative group algebras*. New-York, 1983.
- [6] COHN P.M. *Free rings and their relations*. Academic Press, London, New-York, 1971.
- [7] SMITS T.H.M. *Skew polynomial rings*. Indag. Math., 1968, **30**, p. 209–224.
- [8] SMITS T.H.M. *The free product of a quadratic number field and semifield*. Indag. Math., 1969, **31**, p. 145–159.
- [9] RYABUKHIN YU.M. *Quasi-regular algebras, modules, groups and varieties*. Buletinul A.S.R.M., Matematica, 1997, N 1(23), p. 6–62 (in Russian).
- [10] ANDRUNAKIEVICH V.A., RYABUKHIN YU.M. *Radicals of algebras and structure theory*. Moscow, Nauka, 1979 (in Russian).

State University of Moldova
A. Mateevici str. 60
Chisinau MD-2009, Moldova
E-mail: *cojuhari@usm.md*

Received August 28, 2006