

 Open access • Journal Article • DOI:10.1136/EBNURS-2011-100294

## The reciprocal relationship between physical activity and depression in older European adults — [Source link](#)

John A Owiti, Kamaldeep Bhui

**Institutions:** Queen Mary University of London

**Published on:** 01 Oct 2012 - Evidence-Based Nursing (Evid Based Nurs)

**Topics:** Child and adolescent psychiatry, Population ageing, Depression (differential diagnoses) and Life expectancy

Related papers:

- [Depression and physical disability](#)
- [Study of a cohort of patients newly diagnosed with depression in general practice: prevalence, incidence, comorbidity, and treatment patterns.](#)
- [Depression Care Across the Lifespan](#)
- [Relationship between depression symptom and family factors in children.](#)
- [Anxiety and Depression in Older People: Diagnostic Challenges](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/the-reciprocal-relationship-between-physical-activity-and-gtbr4fa42p>



## Quantitative - other

# The reciprocal relationship between physical activity and depression in older European adults

John Owiti, Kamaldeep Singh Bhui

10.1136/ebnurs-2011-100294

Department of Psychiatry,  
Wolfson Institute of Preventive  
Medicine, Queen Mary  
University of London, UK

Correspondence to:

**Kamaldeep Singh Bhui**  
Department of Psychiatry,  
Wolfson Institute of Preventive  
Medicine, Queen Mary  
University, Charterhouse  
Square, Old Anatomy building,  
London, EC1M 6BQ, UK;  
k.s.bhui@qmul.ac.uk

## Implications for practice and research

- The study highlights the mental health benefits of regular physical activity in the prevention of future depressive illness among older adults.
- Physical activity appears to be effective for promoting mental health and preventing depression, although further study is needed of different types of depression, for example, recurrent or chronic depression.
- Future studies, specifically randomised trials, should also determine the relationship between dose, and type, of physical activity and depression, and unexpected events in a prospective design if physical activity is to be routinely recommended.

## Context

Depression is approximately twice as prevalent among women compared with men and is associated with functional disability, co-morbid medical conditions and social deprivation.<sup>1</sup> Longer life expectancy is leading to a growing ageing population for whom depression is a major public mental health problem. Greater physical activity can prevent the onset of depression and reduce symptoms of depression in older adults<sup>2</sup> and children and adolescents.<sup>3-4</sup> Regular physical activity may be an adjunct treatment for depression.<sup>5</sup> However, reverse causality is often a possible explanation for previous findings, and even in prospective designed studies, the interactive confounding effects of age, gender, physical activity and depression have not been fully investigated.

## Methods

The two-wave cross-lagged panel design with a 2-year follow-up used structural equation modelling to investigate the reciprocal nature of the physical activity and depression in a community sample of older adults. The study evaluates the moderating effects of gender and age. The study tested two effects: physical activity causing depression and depression causing physical activity. The cross-lagged modelling technique is widely used to assess causal associations in data derived from non-experimental, longitudinal research designs.

## Findings

The study found that higher levels of physical activity were associated with lower levels of affective suffering and motivation in cross-sectional analyses at baseline and at follow-up. Lower baseline physical activity was associated with more depressive symptoms at follow-up 2 years later. The trends seemed strongest for men, but there were no significant differences between men and women. There was not an association between baseline depressive symptoms and later physical activity at follow-up in the overall sample. Depression to physical activity relationships were

found primarily in younger age groups (<65), whereas physical activity to depression relationships were primarily supported in older age groups (>65).

## Commentary

Lindwall and colleagues' study provides additional evidence on the benefits of physical activity preventing incident depression in older adults. This study supports other cross-sectional and longitudinal studies. The promotion of physical activity in the prevention of depression may be cost-effective and physical activity as an adjunct treatment may avoid the adverse effects of antidepressant medications. However, more evidence is needed from high-quality studies with adequate power and sample size. These studies should help understand the mechanisms that link physical activity and its preventive effects, whether these are mediated by specific types of activity or other factors like leisure time, living near appropriate facilities, social support and levels of co-morbid health problems.

Future studies should include additional covariates including culture and ethnicity, as moderating or precipitating factors, rather than control for them; furthermore studies are needed to show a dose-response relation between activity (type and intensity) and phenotypes for clinical depression.<sup>6</sup> The new mental health policy calls for a focus on cost-effective preventive interventions in promoting both physical and mental well-being.<sup>7</sup> Physical and mental healthcare professionals might prioritise promotion of physical activity in their interventions. However, benefits must be weighted against injuries, accidents and unexpected events; these need to be assessed in randomised controlled trials and other study types as appropriate.

Competing interests None.

## References

1. McDougall FA, Kvaal K, Matthews FE, *et al.* Prevalence of depression in older people in England and Wales: the MRC CFA Study. *Psychol Med* 2007;37:1787-95.
2. Strawbridge WJ, Deleger S, Roberts RE, *et al.* Physical activity reduces the risk of subsequent depression for older adults. *Am J Epidemiol* 2002;156:328-34.
3. Larun L, Nordheim LV, Ekland E, *et al.* Exercise in prevention and treatment of anxiety and depression among young children and young people. *Cochrane Database Syst Rev* 2006;3.
4. Rothern C, Edwards P, Bhui K, *et al.* Physical activity and depressive symptoms in adolescents: a prospective study. *BMC Med* 2010;8:32.
5. Mead GE, Morley W, Campbell P, *et al.* Exercise for depression. *Cochrane Database Syst Rev* 2009;3.
6. Paffenbarger RS Jr, Lee IM, Leung R. Physical activity and personal characteristics associated with depression and suicide in American college men. *Acta Psychiatr Scand Suppl* 1994;377:16-22.
7. Department of Health, No Health Without Mental Health: A Cross-government Mental Health Outcomes Strategy for People of all Ages, 2010.