



The Relationship between Saccades and Locomotion

Anshul Srivastava,¹ Omar F. Ahmad,¹
Christopher Pham Pacia,² Mark Hallett,¹
Codrin Lungu³

¹Human Motor Control Section, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD, USA

²Department of Biomedical Engineering, Washington University in St. Louis, Saint Louis, MO, USA

³Division of Clinical Research, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD, USA

ABSTRACT

Human locomotion involves a complex interplay among multiple brain regions and depends on constant feedback from the visual system. We summarize here the current understanding of the relationship among fixations, saccades, and gait as observed in studies sampling eye movements during locomotion, through a review of the literature and a synthesis of the relevant knowledge on the topic. A significant overlap in locomotor and saccadic neural circuitry exists that may support this relationship. Several animal studies have identified potential integration nodes between these overlapping circuitries. Behavioral studies that explored the relationship of saccadic and gait-related impairments in normal conditions and in various disease states are also discussed. Eye movements and locomotion share many underlying neural circuits, and further studies can leverage this interplay for diagnostic and therapeutic purposes.

Key Words

Gait; posture; saccade; fixation; locomotion; deep brain stimulation

Received: March 21, 2018 Accepted: April 26, 2018

Corresponding author: Codrin Lungu, MD

<https://orcid.org/0000-0001-6253-3281>

Division of Clinical Research, National Institute of Neurological Disorders and Stroke, National Institutes of Health, 6001 Executive Blvd, #2188, Rockville, MD 20852, USA
Tel: +1-301-496-9135 Fax: +1-301-480-1080 E-mail: lunguci@ninds.nih.gov

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Visual information from the environment is gathered through quick eye movements, which consist of a series of saccades and fixations. Saccades align the fovea with an object of interest.¹ Once an object is foveated, it is held stationary during a fixation, allowing time for the visual information to be collected.¹ Efficient locomotion is dependent upon visual information that is gathered by these quick eye movements. Understanding the relationship among fixations, saccades and locomotion may provide insight into how these seemingly parallel and potentially integrated systems work together.

When studied independently, saccadic and locomotor parameters (Table 1) can be measured precisely. It is difficult to reach the same level of precision when measuring both parameters simultaneously. To get around this, most studies in the literature have correlated saccadic eye movement or gait-related parameters^{2,3} with a given disease state or functional impairment.

In this review, we explore the literature for correlations made between saccades and locomotion. We present the neural circuitry of saccadic and gait-related circuitry and the similarities between them. We highlight brain regions that have been found in animal studies that potentially integrate these two networks. Lastly, we review neurodegenerative diseases

that manifest saccadic and gait-related impairments.

NEURAL COMPONENTS OF SACCADES, FIXATIONS AND LOCOMOTION

Fixation, saccades and locomotion are served by specific areas and networks of the brain. It is particularly interesting to compare the neural components of saccades and locomotion because there are many overlapping brain areas, suggesting a potential integrated neural network between them.

The most relevant areas that support neural integration between saccades and locomotion would likely be at the level of the brainstem and the cerebellum (Figure 1). Afferent inputs between these two parallel networks differ greatly, in that spinal cord pathways provide the majority of sensory information for locomotion, while the geniculate and extrageniculate pathways are important for saccades. On the other hand, modulating structures such as the cerebral cortex, basal ganglia and thalamus are common to all sensorimotor networks and are non-specific to locomotion and saccades. Saccades and locomotion are primitive functions, are well-developed in lower species,^{4,5} and are more likely to be preserved in primitive integrating brain areas, such as the brainstem and cerebellum, more specifically, the mesopontine tegmentum and the cerebellar vermis.

Table 1. Eye movement/fixation parameters and gait/balance parameters

Saccadic/fixation parameters ²	
Fixation duration	Duration of time that the eyes remain fixated. Measured in milliseconds to seconds.
Saccadic duration	Duration of time between saccadic initiation and the saccadic endpoint.
Saccadic latency	Time taken for the eyes to move (saccade) after the target appears. Measured in milliseconds or seconds.
Saccadic amplitude	Arc distance of rotational movement made during a saccade. Sometimes called saccadic size. Measured in degrees or minutes.
Saccadic peak velocity	During a saccade, it is the highest velocity attained. Measured in degrees/seconds.
Saccadic intrusions	Series of irregular interruptions by fast eye movements during primary fixation.
Saccadic gain	Ratio of the actual saccadic amplitude over the intended saccadic amplitude.
Main sequence	Relationship among saccadic peak velocity, duration and amplitude.
Gait/balance parameters ³	
Step length	Distance between initial ground contact of one foot and initial ground contact of the opposite foot.
Step time	Time in seconds between initial ground contact of one foot and initial ground contact of the opposite foot.
Step width	Lateral distance between the centers of the heels when both feet are on the ground (i.e., double stance).
Stride length	Distance between initial ground contact of one foot and initial ground contact of the same foot, constituting the distance of one gait cycle.
Stride time	Time between initial ground contact of one foot and initial ground contact of the same foot, constituting the time of one gait cycle.
Postural sway	Horizontal movement of the center of gravity while standing still.
Swing phase	Remaining 40% of the gait cycle, when the foot no longer is in contact with the ground, spanning from initial swing phase to initial contact.
Cadence	Steps per minute.
Stance phase	Initial 60% of the gait cycle, when the foot is in contact with the ground, spanning from initial contact to terminal double stance.




Figure 1. Brain areas involved in saccades/fixations and locomotor activities. Possible integration areas are shaded in orange. PFC: prefrontal cortex, PMA: premotor cortex, SMA: supplementary motor cortex, PMC: primary motor cortex, PPC: posterior parietal cortex, PT: putamen, CN: caudate nucleus, GP: globus pallidus, STN: subthalamic nucleus, SN: substantia nigra, SC: superior colliculus, MLR: mesencephalic locomotor region, PMRF: pontomedullary reticular formation, MRF: mesencephalic reticular formation, FN: fastigial nucleus, NPH: nucleus prepositus hypoglossi, MedRF: medullary reticular formation, PPRF: paramedian pontine reticular formation, VC: visual cortex, VA: ventral anterior, VL: ventrolateral nucleus, PPN: pedunculo-pontine nucleus.

Mesencephalic Locomotor Region/ Mesencephalic Reticular Formation

The pedunculo-pontine nucleus (PPN) and the cuneiform nucleus (CNF) make up the mesencephalic locomotor region (MLR),⁶ also known as the mesencephalic reticular formation (MRF). The MLR/MRF is involved in eye movement-related activity.⁷⁻⁹ In addition, it promotes locomotion through the reticulo-spinal pathways¹⁰ and influences postural tone and locomotor rhythmicity.¹¹⁻¹³ In animal studies, stimulation of the CNF has been found to be associated with locomotor initiation, while stimulation of the PPN was associated with locomotor suppression.¹⁴ The PPN contains cholinergic, glutaminergic and GABAergic neurons; the cholinergic neurons are those closely associated with locomotion.¹⁵ PPN cholinergic neurons are also associated with rapid eye movements in sleep.¹⁵ The PPN directly innervates the motor neurons involved in eye movements and receives direct projections from the frontal and supplementary eye fields in the cortex.¹⁶⁻²⁰ Neuronal recordings of the PPN in primates have shown different firing patterns during fixations and saccades.^{21,22}

The PPN receives input from the cerebral cortex and has reciprocal connections with components of the basal ganglia, namely, the substantia nigra [both the substantia nigra pars reticulata (SNr) and the substantia nigra pars compacta (SNc)], globus pallidus and subthalamic nucleus (STN).²³⁻²⁸

Superior colliculus

The superior colliculus (SC) receives inputs from the retina and visual cortex (VC).²⁹⁻³² Neurons in the SC have projections to saccade generators in the brainstem.³³ The SC has been reported to be associated with fixation- and saccade-related activity.³⁴⁻³⁶ There is no evidence for locomotor function related to the SC; however, the SC does receive afferents from various subcortical structures common to the locomotor network, such as the SNr, pretectum, and other nuclei in the pons and medulla. SC efferents project to the thalamus, MLR/MRF, paramedian pontine reticular formation (PPRF), cerebellar locomotor region and cerebellar vermis.³⁷ The PPRF is important for coordinating horizontal saccadic eye movements, but its role in locomotion has not yet been ex-

ploded. The PPRF receives input from the frontal eye fields (FEF) through the contralateral SC³⁸ and contains burst neurons that generate horizontal saccades.³⁹⁻⁴¹

Pontomesencephalic reticular formation

Reticulospinal neurons in the pontomesencephalic reticular formation are involved in controlling and maintaining head movements and in generating the quick phase of vestibular and optokinetic head nystagmus toward the same side.⁴² Omnidirectional pause neurons (OPNs) are inhibitory interneurons in the pontomesencephalic reticular formation that are thought to stabilize fixations and saccades in the horizontal, vertical and oblique directions. OPNs are tonically active during fixations and are silent (i.e., “paused”) during saccades.⁴³ Dysfunction in OPNs is thought to result in fixational instability, with reports of macrosaccadic oscillations, saccadic dysmetria, ocular flutter, and opsoclonus.^{44,45} The pontomesencephalic reticular formation is also involved in transmitting locomotor signals to central pattern generators in the spinal cord⁴⁶ and in controlling balance, locomotion and posture.^{47,48}

Cerebellar vermis

The cerebellum is involved in both locomotion⁴⁹⁻⁵⁴ and saccades.⁵⁵⁻⁶⁵ The fastigial nucleus (FN) of the cerebellum receives input from the vermis, which in turn receives input from the SC through the nucleus reticularis tegmenti pontis.^{55,66,67} Brainstem saccade generators are driven by the FN and the vermis.⁴¹ Studies of transcranial magnetic stimulation directed toward the cerebellar vermis have demonstrated that this area coordinates saccades ipsilateral to the side of stimulation.⁶⁸ Neuronal discharge in the FN, also known as the cerebellar locomotor region, is linked to coding of proximal movement during locomotion.^{55,69} The FN is thought to act as a pacemaker during locomotion⁷⁰ and projects to the pontomedullary reticular formation in the brainstem.

Thalamus

The thalamus serves as the major relay between cortical and subcortical saccadic generators.⁷¹⁻⁷³ The internal medullary lamina, a myelinated area that divides the thalamus into the anterior, medial and lateral masses, contains nuclei that relay information among multiple areas that control saccades, namely,

the frontal and parietal eye fields, SC, PPRF, striatum, cerebellum and the lateral geniculate nuclei.⁷¹

The lateral geniculate nuclei and pulvinar are two thalamic nuclei in the ventrolateral area that specifically process visual input. The lateral geniculate nucleus projects information from the retina to the VC. Connections between the SC and the lateral geniculate nucleus contribute to saccades that are involved in foveating objects of interest with a high degree of resolution (e.g., facial recognition).⁷⁴ The pulvinar has connections between the SC and visual cortices and is involved in visuospatial attention to areas in the visual field.⁷⁵ The pulvinar is an important relay for generating saccades toward visual targets or reflexive saccades toward or away from stimuli, and this nucleus influences visually guided behavior, including locomotion. It has been speculated that visual and motor information may be integrated in the pulvinar, allowing a distinction between changes in the visual environment caused by external sources versus self-generated visual motion (caused by eye movements or locomotion).⁷⁴

The ventrolateral nucleus (VL) receives all major saccade-generating afferents in the brainstem and cerebellum and projects to the frontal eye field and the supplementary eye field.⁷⁶ Similar to the pulvinar, the VL is closely involved in visually guided saccades.⁷⁷ The VL is also a major afferent to the primary motor cortex, and it is not surprising that this region is important for locomotion.^{78,79}

The thalamic reticular nucleus is a thin capsule of inhibitory GABAergic neurons that surrounds the dorsolateral thalamus and functions to modulate thalamocortical and corticothalamic signals for a multitude of functions.⁸⁰ In terms of saccadic and locomotor networks, this region functions as an inhibitory modulator. The thalamic reticular nucleus sends reciprocal inhibitory signals to the lateral geniculate nucleus in response to saccade-related visual perturbations to maintain a stable image.⁸¹ Recordings have revealed phasic bursts of activity in reticular neurons within the receptive field of distal limbs during walking tasks that are thought to fine tune ongoing locomotor activity.⁸²

Basal ganglia

The basal ganglia refers mainly to the caudate and the putamen, which consist of the striatum, globus pallidus, substantia nigra and STN. The nigrostri-

tal pathway modulates the striatum, affecting all motor output, and is not specific to saccadic or locomotor control, though its influence over these functions is considerable.⁸³⁻⁸⁷ The STN receives inputs from the cortex via the striatum and the globus pallidus externa (GPe) through the indirect pathway and direct connections from the cortex through the hyperdirect pathway.⁸⁸ The STN receives inputs from the brainstem, thalamus and cortex. Efferents from the STN travel mainly to the GPi and SNr.⁸⁹⁻⁹¹ There is evidence that patients with Parkinson disease (PD) who receive deep brain stimulation (DBS) of the STN experience a significant improvement in both saccadic performance^{92,93} and locomotion⁹³⁻⁹⁵ compared to patients that receive other DBS targets, such as the globus pallidus interna (GPi). GPi DBS has been shown to improve locomotion,⁹⁶ but there is less evidence supporting an improvement in saccadic performance,⁹⁷ though one study found improvement in antisaccades.⁹⁸

ANIMAL STUDIES EXPLORING THE INTEGRATION BETWEEN EYE MOVEMENT AND LOCOMOTOR CIRCUITRY

Thus far, we have identified brain areas that are common to both saccades and gait in humans. Animal studies have provided much of the direct evidence for the integration of networks controlling saccades and gait.

Semi-intact experiments in lampreys undergoing electrical stimulation of the optic tectum have demonstrated a stimulus-dependent coordination of eye movements with steering and goal-directed behavior. Saitoh et al.⁹⁹ showed that, with increasing stimulation of the lateral optic tectum, there is a stepwise recruitment of eye movements, followed by a coordinated lateral bending of the body, and then by coordinated locomotor movements. Stimulating other areas, such as the caudomedial tectum, elicits different behaviors, such as struggling behavior, characterized by undulating body movements with anti-phasic eye movements. These experiments have lent support for the role of the optic tectum (SC in primates) as a stepwise integrating interface for patterned visuomotor and locomotor behavior.⁹⁹

The coordination between eye movements and spinal locomotor patterns is also preserved and adapt-

able at different stages of development. Uckerman et al.¹⁰⁰ demonstrated how the *Xenopus laevis* (XL) frog adapts visuomotor control to maintain image stabilization when swimming as it transitions from a tadpole to an adult frog. In the tadpole, propulsion is achieved with undulating tail movements, requiring conjugate left-right eye rotations to maintain a stable binocular image. In the frog, forward acceleration is achieved with rhythmic bilateral leg kicking that requires nonconjugate, convergent-divergent, eye movements. In fixed-head preparations, a strict 1:1 relationship was found between eye movements and spontaneous fictive swimming movements. Vestibular and visual input were controlled for by transecting the optic nerves and ablating the vestibular end organs. In tadpoles, the eyes rotate laterally, countering each lateral tail movement, while in frogs, the eyes converge or diverge in phase with the kick cycle. This experiment provided evidence for multimodal integration between spinal central pattern generators and eye movements during locomotion in XL. More importantly, the ability of visuomotor and locomotor networks to change in a coordinated fashion at different stages of development in XL suggested that they are integrated. This adaptability is probably evolutionarily preserved in other forms of locomotion, such as quadrupedal and bipedal ambulation. The OPN, as mentioned earlier, coordinates horizontal, vertical and oblique fixations and saccades. It is possible that the omnidirectional stabilizing capability of these interneurons provides a mechanism for the adaptability of reflexive saccades to different locomotive head perturbations across species.

Schwarz et al.¹⁰¹ performed microelectrode recordings of nondopaminergic SNr neurons in cats as they received different sensory stimuli, such as mechanical skin stimulation, passive and active limb movement, and visual and vestibular stimuli. Neurons within the receptive field of each limb showed regular discharge patterns that were in phase with the step cycle during locomotion. Avoiding or navigating around an obstacle had the greatest effect on neuronal firing rates. Objects moving within the contralateral visual field modulated the firing rates of a small population of neurons related to saccades. Similar findings of saccades and neuronal discharge in the SNr have been found in monkeys.¹⁰² The authors concluded that the SNr functions as an output

station that processes convergent multimodal sensory input (e.g., joint position, limb movement, direction and amplitude of saccades) and fine tunes spinal motor output to adequately address changing environments.

The PPN has also been suggested to serve as a multimodal integrative interface.¹⁰³ Suppression of spontaneous locomotion and rhythmic eye movements was observed with stimulation of the ventral PPN in anesthetized and acutely decerebrated cats.¹⁴

Saccade-related¹⁰⁴⁻¹⁰⁶ and locomotion-related¹⁰⁷⁻¹⁰⁹ neuronal activity has been reported in Purkinje cells in the cerebellar vermis in various studies using microstimulation and optogenetic techniques in non-human primates and other mammals.

The SC and PPRF^{110,111} have been shown in rhesus monkeys to influence coordinated head-eye movements, an important component of steering during locomotion.¹¹²

Saleem et al.¹¹³ showed that, in order for mice to accurately gauge their speed when navigating their environment, visual speed, derived from optic input, and running speed, derived from proprioceptive input, are integrated and encoded with weighted sums within the neurons of the V1 area of the occipital cortex. While this does not pertain to eye movements per se, it at least provides more evidence linking visual sampling (which requires adequate saccades and fixations) and locomotion.

While numerous studies have suggested a multimodal integration between saccades and locomotion, the challenge of establishing a neural basis for this interaction, especially in humans, is hindered by the technical limitations related to studying the circuitry of eye movements during the act of walking. Therefore, the level at which these circuits interact with each other in real time and how activating or inactivating various nodes within one neural circuit may affect the functions of the other are not yet known.

BEHAVIORAL STUDIES IN HEALTHY INDIVIDUALS EXPLORING THE RELATIONSHIP AMONG SACCADES, FIXATIONS AND LOCOMOTION

During ambulation, the limbs, body, head and eyes move in a coordinated manner.^{112,114,115} Saccadic eye movements allow the fovea to maintain fixations

on relevant objects in the environment in a dynamic manner to allow guidance of locomotion. Any problems in this fixation-saccade strategy may lead to visual and gait impairments.

The visuomotor and locomotor systems influence each other via a continuous feedback loop, though the exact network is not well delineated.¹¹⁶⁻¹¹⁸ Several studies have focused on gaze fixations and saccadic eye movements during stepping¹¹⁹⁻¹²⁵ to describe how eye movements influence gait parameters. In one study, visual information gathered during the latter half of the preceding step was shown to influence the step length of the following step.¹²⁶ It has also been suggested that, while walking on uneven ground or terrain, visual information from two steps is required to direct foot placement.¹²⁷

Marigold and Patla¹²⁸ found that, when walking on a varying terrain, participants visually fixated on areas of the ground where they eventually stepped. Additionally, fixations were frequently guided to the transition zones between the varying surfaces (e.g., solid to compliant, rocky to slippery, tilted to irregular, etc). Hollands and Marple-Horvat¹²⁹ studied the eye movements of healthy participants who were made to walk in different conditions that varied in terms of the amount visual information available to the participants as they stepped onto stepping stones. The time interval between saccadic onset and foot-lift was similar in all conditions, but the interval between saccadic onset and footfall onto the stepping target differed significantly depending on the amount of visual information present. Patla and Vickers¹³⁰ found that healthy participants fixated on footfall targets that were an average of two steps ahead. Elderly participants with a history of falls tended not to look two steps ahead but instead fixated more on the imminent footfall target.¹³¹ This finding may be the result of impaired central processing of visually guided information in that group, as suggested in another study, in which elderly participants with a high risk of falling had longer latencies from saccadic initiation to foot-lift than elderly individuals with a low risk of falling.¹³²

Saccades were also studied in individuals during turning maneuvers. These studies supported a “top-down” model, in which saccadic initiation precedes, and possibly influences, turning of the head, trunk and legs.^{112,114} Imai et al.¹¹⁴ observed that when participants were asked to move in a straight line and

turn 90 degrees, a saccade was made in the direction of the turn. A similar observation was made by Hollands et al.,¹³³ in which healthy participants made saccades in order to position their gaze in line with the endpoint of the required travel path.

Anxiety can influence the interplay between gait and saccades. It has been suggested that early gaze transfer due to anxiety over impending obstacles is correlated with stepping inaccuracies. Investigators observed the visual and stepping behavior of an 87-year-old female when she was directed to walk along a stepping path before and after an obstacle. At the beginning of the experiment, she fixated on the stepping path before the obstacle. After falling twice, she stopped fixating on the stepping path, and instead fixated on the obstacle itself.¹³⁴ In a similar study, elderly participants with a high risk of falling were more likely to transfer their gaze early from a stepping target along a path to an impending obstacle.¹³⁵ One study indirectly showed a relationship between saccades and gait during an episode of anxiety/fear, in which participants with a fear of heights made more vertical than horizontal saccades when walking on a fire escape 20 meters above ground compared to the saccades of the controls.¹³⁶ The amygdala plays an important role in anxiety and has been found to be involved in saliency coding when scanning a visual scene.¹³⁷ States of increased anxiety may disrupt fixations and saccades through this pathway.

The relationship between saccades and gait was observed in healthy participants as they moved along a pathway with irregularly placed stepping stones, both with and without an alcohol dose. Gait

impairments were observed in terms of increased step cycle durations and missed footfall targets. In terms of saccadic impairments, a large proportion of the saccades of the successive stepping stones were inaccurate and were accompanied by corrective saccades.¹³⁸ Alcohol has been shown to cause saccadic dysmetria.¹³⁹ The combination of impaired saccadic control and stepping accuracy implicates the cerebellum [See Supplementary Table 1 (in the online only Data Supplement) for summary of the studies of this section].

SACCADES AND GAIT IN NEURODEGENERATIVE DISEASES

While saccadic and gait abnormalities have been studied separately in various neurodegenerative disorders (Table 2),¹⁴⁰⁻¹⁷⁴ simultaneous recordings of eye movements and gait in these disorders have rarely been reported.

PD is well known as having both saccadic¹⁵¹ and gait abnormalities.¹⁷⁵ In PD, both saccades and step length can be hypometric. Side-to-side asymmetry, in terms of step length and saccadic amplitude, is often seen in PD. Nemanich and Earhart reported that, in PD, freezing of gait is associated with increased saccadic latency and variability.¹⁷⁶ The researchers found that PD patients with freezing of gait were slower in initiating pro- and antisaccades. Saccadic velocity and gain variability were also increased in PD with freezing of gait. Performance of antisaccades was impaired in PD patients with freezing of gait compared to patients without freezing.¹⁷⁷ In an-

Table 2. Separate studies showing saccadic abnormalities or gait abnormalities in essential tremor, PD, PSP, Huntington disease and cerebellar ataxia

Disorder	Saccadic abnormalities	Gait abnormalities
Essential tremor	Slow saccades and increased square-wave jerks ¹⁴⁰	Tandem gait difficulty ¹⁴¹⁻¹⁴⁹
PD	Hypometric saccades and prolonged saccadic latency ^{150,151}	Freezing of gait, falls, turning impairment, and decreased stride length ^{152,153}
PSP	Fixational saccades that are abnormally large. Square wave jerks more frequent, larger, and markedly more horizontal ¹⁵⁴ Vertical saccades (slow and hypometric, both up and down) ¹⁵⁵	Hypokinetic gait characteristics: decreased velocity and step length ¹⁵⁶ Interstep variability and asymmetry during gait. Slower cadence. Freezing of gait and frequent falls ¹⁵⁷
Huntington disease	Slow saccades ¹⁵⁸⁻¹⁶¹ Increased variability in saccadic reaction times and occurrence of errors ^{162,163} Hypometric primary saccades ¹⁶⁴	Gait characteristic variation in each walk, with mean decreases in velocity, stride length, and cadence. Decreased gait velocity ¹⁶⁵⁻¹⁶⁷ Disordered regulation of footstep timing; reduced stride length ¹⁶⁸
Cerebellar ataxia	Square-wave jerks, saccadic dysmetria, and reduced saccadic velocity ¹⁶⁹⁻¹⁷¹	Decreased step length, stride length, and gait speed ¹⁷²⁻¹⁷⁴

PD: Parkinson disease, PSP: progressive supranuclear palsy

other study, saccadic frequency was found to increase in both patients with PD and their age-matched controls when approaching a turn, but the PD patients made fewer preparatory saccades than the controls before the turn.^{178,179} During the turn, the PD patients made more saccades, and the saccadic velocity was slower than that of the controls.¹⁸⁰

The likely neural components affecting both saccades and locomotion in PD include the STN, the SNr, and the MLR/MFR.¹⁷⁵ In PD, degeneration of dopaminergic neurons in the SNc affects the direct and indirect pathways, resulting in bradykinetic movements that affect locomotion and saccades. More specifically, there is increased excitation of the STN, causing an increased inhibitory effect of the GPi and SNr through the indirect pathway. As mentioned earlier in the current review, DBS of the STN affects both saccadic and locomotor performance when compared to DBS of the GPi.⁹²⁻⁹⁵ In terms of eye movements, the effect on these pathways in PD results in increased excitation of the SNr, which leads to abnormal saccade generation in the SC. There is also increased excitation of the PPN, which, as mentioned previously, has projections that are related to saccades and locomotion. In a recent imaging study, PPN alterations were suggested to be related to both saccadic and postural impairments in patients with PD.¹⁸¹ It was observed that functional connectivity involving the PPN and FEF correlated with antisaccadic latencies in healthy participants but not in PD patients with postural instability. Additionally, saccadic impairment correlated with gait initiation im-

pairment in patients.

Additional examples of neurological disorders with abnormal saccades and postural instability other than PD^{182,183} include progressive supranuclear palsy,¹⁸⁴ cerebellar ataxia,¹⁸⁵ essential tremor,¹⁸⁶ and Huntington disease.^{187,188} Some studies have reported that abnormalities in saccadic eye movements are correlated with body sway, even in healthy individuals.^{189,190} These findings of these studies reflect an integration between postural dynamics and eye movements.

Patients with cerebellar ataxia have ataxic gait and dysmetric saccades. Dysmetric saccades consist of hypometric or hypermetric initial saccades, followed by a corrective saccade. TMS studies have implicated the ipsilateral cerebellar vermis in saccadic dysmetria.⁶⁸ Studies of visual fixation in patients with cerebellar ataxia have discovered the presence of dysmetric saccades. During locomotor tasks with visually guided stepping, both dysmetric saccades and ataxic gait were detected.^{191,192} Other studies have found correlations between efficient footfalls and oculomotor function^{127,129,130} in healthy subjects.

Studies of saccadic performance in patients with gait impairment could provide insight into how eye movements affect motor abnormalities such as freezing of gait, imbalance, turning difficulties and falls. Beyond that, saccadic eye movement training as a gait rehabilitation strategy could be an important therapeutic option. Some studies have reported saccadic eye movement training as a strategy for alleviating gait abnormalities in terms of improvement in

Table 3. Eye movement training and gait

Authors	Year	Participants	Method	Main findings
Eye movement training and gait				
Zampieri and Di Fabio ¹⁹³	2008	19 moderately affected progressive supranuclear pals patients	Balance training and eye movement exercises Eye movement training: eye movement practice on the computer screen with randomly appearing arrows on the screen	Improvements in stance time and walking speed in the treatment group
Crowdy et al ¹⁹⁴	2002	2 cerebellar patients	Foot placement (stepping task) Eye movement training: rehearsal of saccades for footfall targets in a stationary standing condition	Improvements in oculomotor and locomotor performance following eye-movement rehearsal
Kang and Yu ¹⁹⁵	2016	14 stroke patients	Foot placement (stepping task) Eye movement training: visual scanning of the picture cards, fixating gaze on a moving baton	Improvements in walking speed, step length and cadence

stance time and accuracy in stepping in patients (Table 3).¹⁹³⁻¹⁹⁵

CONCLUSIONS AND FUTURE DIRECTIONS

Eye movements and locomotion share common neural substrates and potentially have interlinked neural circuitries. The mesopontine tegmentum and cerebellar vermis are the most likely areas to have specific neural connections between these parallel networks. Physiological studies in animals and behavioral studies in healthy individuals have supported the hypothesis that these connections are preserved and adaptable across species. Many neurodegenerative disorders demonstrate coexisting eye movement and gait abnormalities. Correlations have been made in these disease states, further providing evidence of interlinked neural circuitry. As the technology of mobile eye-tracking improves, future studies exploring eye movement abnormalities in real time with simultaneous gait recording will further elucidate the interplay between these two networks. In addition, such studies may potentially serve to develop new diagnostic or disease severity markers.

Supplementary Materials

The online-only Data Supplement is available with this article at <https://doi.org/10.14802/jmd.18018>.

Conflicts of Interest

The authors have no financial conflicts of interest.

REFERENCES

- Purves D, Augustine GJ, Fitzpatrick D, Hall WC, LaMantia AS, McNamara JO, et al. Eye movements and sensory motor integration. In: Purves D, Augustine GJ, Fitzpatrick D, Hall WC, LaMantia AS, McNamara JO, et al. editors. *Neuroscience*. 3rd ed. Sunderland, MA: Sinauer Associates, Inc., 2004;453-467.
- Holmqvist K, Nyström M, Andersson R, Dewhurst R, Jarodzka H, van de Weijer J. *Eye Tracking: A Comprehensive Guide to Methods and Measures*. 1st ed. Oxford: Oxford University Press, 2011;537.
- Hollman JH, McDade EM, Petersen RC. Normative spatio-temporal gait parameters in older adults. *Gait Posture* 2011; 34:111-118.
- Harcourt-Smith WE, Aiello LC. Fossils, feet and the evolution of human bipedal locomotion. *J Anat* 2004;204:403-416.
- Walls GL. The evolutionary history of eye movements. *Vision Res* 1962;2:69-80.
- Ryczko D, Dubuc R. The multifunctional mesencephalic locomotor region. *Curr Pharm Des* 2013;19:4448-4470.
- Graf WM, Ugolini G. The central mesencephalic reticular formation: its role in space-time coordinated saccadic eye movements. *J Physiol* 2006;570:433-434.
- Waitzman DM, Silakov VL, Cohen B. Central mesencephalic reticular formation (cMRF) neurons discharging before and during eye movements. *J Neurophysiol* 1996;75: 1546-1572.
- Perkins E, May PJ, Warren S. Feed-forward and feedback projections of midbrain reticular formation neurons in the cat. *Front Neuroanat* 2014;7:55.
- Skinner RD, Kinjo N, Henderson V, Garcia-Rill E. Locomotor projections from the pedunculopontine nucleus to the spinal cord. *Neuroreport* 1990;1:183-186.
- Garcia-Rill E, Skinner RD, Fitzgerald JA. Activity in the mesencephalic locomotor region during locomotion. *Exp Neurol* 1983;82:609-622.
- Garcia-Rill E, Houser CR, Skinner RD, Smith W, Woodward DJ. Locomotion-inducing sites in the vicinity of the pedunculopontine nucleus. *Brain Res Bull* 1987;18:731-738.
- Sherman D, Fuller PM, Marcus J, Yu J, Zhang P, Chamberlin NL, et al. Anatomical location of the mesencephalic locomotor region and its possible role in locomotion, posture, cataplexy, and parkinsonism. *Front Neurol* 2015;6:140.
- Takakusaki K, Saitoh K, Harada H, Okumura T, Sakamoto T. Evidence for a role of basal ganglia in the regulation of rapid eye movement sleep by electrical and chemical stimulation for the pedunculopontine tegmental nucleus and the substantia nigra pars reticulata in decerebrate cats. *Neuroscience* 2004;124:207-220.
- Garcia-Rill E, Hyde J, Kezunovic N, Urbano FJ, Petersen E. The physiology of the pedunculopontine nucleus: implications for deep brain stimulation. *J Neural Transm (Vienna)* 2015;122:225-235.
- Cohen B, Waitzman DM, Büttner-Ennever JA, Matsuo V. Horizontal saccades and the central mesencephalic reticular formation. *Prog Brain Res* 1986;64:243-256.
- Huerta MF, Krubitzer LA, Kaas JH. Frontal eye field as defined by intracortical microstimulation in squirrel monkeys, owl monkeys, and macaque monkeys: I. Subcortical connections. *J Comp Neurol* 1986;253:415-439.
- Stanton GB, Goldberg ME, Bruce CJ. Frontal eye field efferents in the macaque monkey: I. Subcortical pathways and topography of striatal and thalamic terminal fields. *J Comp Neurol* 1988;271:473-492.
- Huerta MF, Kaas JH. Supplementary eye field as defined by intracortical microstimulation: connections in macaques. *J Comp Neurol* 1990;293:299-330.
- Shook BL, Schlag-Rey M, Schlag J. Primate supplementary eye field: I. Comparative aspects of mesencephalic and pontine connections. *J Comp Neurol* 1990;301:618-642.
- Okada K, Kobayashi Y. Fixational saccade-related activity of pedunculopontine tegmental nucleus neurons in behaving monkeys. *Eur J Neurosci* 2014;40:2641-2651.
- Okada K, Kobayashi Y. Rhythmic firing of pedunculopontine tegmental nucleus neurons in monkeys during eye movement task. *PLoS One* 2015;10:e0128147.
- Lavoie B, Parent A. Pedunculopontine nucleus in the squirrel monkey: projections to the basal ganglia as revealed by anterograde tract-tracing methods. *J Comp Neurol* 1994; 344:210-231.
- Matsumura M, Nambu A, Yamaji Y, Watanabe K, Imai H, Inase M, et al. Organization of somatic motor inputs from the frontal lobe to the pedunculopontine tegmental nucleus in the macaque monkey. *Neuroscience* 2000;98:97-110.
- Martinez-Gonzalez C, Bolam JP, Mena-Segovia J. Topo-

- graphical organization of the pedunculopontine nucleus. *Front Neuroanat* 2011;5:22.
26. Martinez-Gonzalez C, van Andel J, Bolam JP, Mena-Segovia J. Divergent motor projections from the pedunculopontine nucleus are differentially regulated in Parkinsonism. *Brain Struct Funct* 2014;219:1451-1462.
 27. Lau B, Welter ML, Belaid H, Fernandez Vidal S, Bardinet E, Grabli D, et al. The integrative role of the pedunculopontine nucleus in human gait. *Brain* 2015;138:1284-1296.
 28. Strumpf H, Noesselt T, Schoenfeld MA, Voges J, Panther P, Kaufmann J, et al. Deep brain stimulation of the pedunculopontine tegmental nucleus (PPN) influences visual contrast sensitivity in human observers. *PLoS One* 2016;11:e0155206.
 29. Perry VH, Cowey A. Retinal ganglion cells that project to the superior colliculus and pretectum in the macaque monkey. *Neuroscience* 1984;12:1125-1137.
 30. Harting JK, Glendenning KK, Diamond IT, Hall WC. Evolution of the primate visual system: anterograde degeneration studies of the tecto-pulvinar system. *Am J Phys Anthropol* 1973;38:383-392.
 31. Fries W. Cortical projections to the superior colliculus in the macaque monkey: a retrograde study using horseradish peroxidase. *J Comp Neurol* 1984;230:55-76.
 32. Lock TM, Baizer JS, Bender DB. Distribution of corticotectal cells in macaque. *Exp Brain Res* 2003;151:455-470.
 33. Izawa Y, Sugiuchi Y, Shinoda Y. Neural organization from the superior colliculus to motoneurons in the horizontal oculomotor system of the cat. *J Neurophysiol* 1999;81:2597-2611.
 34. Munoz DP, Guitton D. Control of orienting gaze shifts by the tectoreticulospinal system in the head-free cat. II. Sustained discharges during motor preparation and fixation. *J Neurophysiol* 1991;66:1624-1641.
 35. Munoz DP, Wurtz RH. Fixation cells in monkey superior colliculus. I. Characteristics of cell discharge. *J Neurophysiol* 1993;70:559-575.
 36. Munoz DP, Wurtz RH. Fixation cells in monkey superior colliculus. II. Reversible activation and deactivation. *J Neurophysiol* 1993;70:576-589.
 37. May PJ. The mammalian superior colliculus: laminar structure and connections. *Prog Brain Res* 2006;151:321-378.
 38. Leichnetz GR, Gonzalo-Ruiz A, DeSalles AA, Hayes RL. The origin of brainstem afferents of the paramedian pontine reticular formation in the cat. *Brain Res* 1987;422:389-397.
 39. Sparks DL. The brainstem control of saccadic eye movements. *Nat Rev Neurosci* 2002;3:952-964.
 40. Cohen B, Komatsuzaki A. Eye movements induced by stimulation of the pontine reticular formation: evidence for integration in oculomotor pathways. *Exp Neurol* 1972;36:101-117.
 41. Scudder CA, Kaneko CS, Fuchs AF. The brainstem burst generator for saccadic eye movements: a modern synthesis. *Exp Brain Res* 2002;142:439-462.
 42. Suzuki SS, Siegel JM, Wu MF. Role of pontomedullary reticular formation neurons in horizontal head movements: an ibotenic acid lesion study in the cat. *Brain Res* 1989;484:78-93.
 43. Optican LM. The role of omnipause neurons: why glycine? *Prog Brain Res* 2008;171:115-121.
 44. Averbuch-Heller L, Kori AA, Rottach KG, Dell'osso LF, Remler BF, Leigh RJ. Dysfunction of pontine omnipause neurons causes impaired fixation: macrosaccadic oscillations with a unilateral pontine lesion. *Neuro-ophthalmology* (Aeolus Press) 1996;16:99-106.
 45. Thurtell MJ, Tomsak RL, Leigh RJ. Disorders of saccades. *Curr Neurol Neurosci Rep* 2007;7:407-416.
 46. Sakai ST, Davidson AG, Buford JA. Reticulospinal neurons in the pontomedullary reticular formation of the monkey (*Macaca fascicularis*). *Neuroscience* 2009;163:1158-1170.
 47. Schepens B, Stapley P, Drew T. Neurons in the pontomedullary reticular formation signal posture and movement both as an integrated behavior and independently. *J Neurophysiol* 2008;100:2235-2253.
 48. Stapley PJ, Drew T. The pontomedullary reticular formation contributes to the compensatory postural responses observed following removal of the support surface in the standing cat. *J Neurophysiol* 2009;101:1334-1350.
 49. He YC, Wu GY, Li D, Tang B, Li B, Ding Y, et al. Histamine promotes rat motor performances by activation of H(2) receptors in the cerebellar fastigial nucleus. *Behav Brain Res* 2012;228:44-52.
 50. Chambers WW, Sprague JM. Functional localization in the cerebellum. I. Organization in longitudinal cortico-nuclear zones and their contribution to the control of posture, both extrapyramidal and pyramidal. *J Comp Neurol* 1955;103:105-129.
 51. Chambers WW, Sprague JM. Functional localization in the cerebellum. II. Somatotopic organization in cortex and nuclei. *AMA Arch Neurol Psychiatry* 1955;74:653-680.
 52. Sprague JM, Chambers WW. Regulation of posture in intact and decerebrate cat. I. Cerebellum, reticular formation, vestibular nuclei. *J Neurophysiol* 1953;16:451-463.
 53. Thach WT, Goodkin HP, Keating JG. The cerebellum and the adaptive coordination of movement. *Annu Rev Neurosci* 1992;15:403-442.
 54. Dichgans J, Diener HC. Clinical evidence for functional compartmentalization of the cerebellum. In: Bloedel JR, Dichgans J, Precht W, editors. *Cerebellar Functions*. Berlin Heidelberg: Springer-Verlag, 1984;126-147.
 55. Zhang XY, Wang JJ, Zhu JN. Cerebellar fastigial nucleus: from anatomic construction to physiological functions. *Cerebellum Ataxias* 2016;3:9.
 56. Joshi AC, Das VE. Muscimol inactivation of caudal fastigial nucleus and posterior interposed nucleus in monkeys with strabismus. *J Neurophysiol* 2013;110:1882-1891.
 57. Kleine JE, Guan Y, Büttner U. Saccade-related neurons in the primate fastigial nucleus: what do they encode? *J Neurophysiol* 2003;90:3137-3154.
 58. Helmchen C, Straube A, Büttner U. Saccade-related activity in the fastigial oculomotor region of the macaque monkey during spontaneous eye movements in light and darkness. *Exp Brain Res* 1994;98:474-482.
 59. Fuchs AF, Robinson FR, Straube A. Role of the caudal fastigial nucleus in saccade generation. I. Neuronal discharge pattern. *J Neurophysiol* 1993;70:1723-1740.
 60. Robinson FR, Straube A, Fuchs AF. Role of the caudal fastigial nucleus in saccade generation. II. Effects of muscimol inactivation. *J Neurophysiol* 1993;70:1741-1758.
 61. Ohtsuka K, Noda H. Saccadic burst neurons in the oculomotor region of the fastigial nucleus of macaque monkeys. *J Neurophysiol* 1991;65:1422-1434.
 62. Helmchen C, Büttner U. Saccade-related Purkinje cell activity in the oculomotor vermis during spontaneous eye movements in light and darkness. *Exp Brain Res* 1995;103:198-208.
 63. Takagi M, Zee DS, Tamargo RJ. Effects of lesions of the oculomotor vermis on eye movements in primate: saccades. *J Neurophysiol* 1998;80:1911-1931.

64. Thier P, Dicke PW, Haas R, Thielert CD, Catz N. The role of the oculomotor vermis in the control of saccadic eye movements. *Ann N Y Acad Sci* 2002;978:50-62.
65. Kojima Y, Soetedjo R, Fuchs AF. Effect of inactivation and disinhibition of the oculomotor vermis on saccade adaptation. *Brain Res* 2011;1401:30-39.
66. Giolli RA, Gregory KM, Suzuki DA, Blanks RH, Lui F, Betelak KF. Cortical and subcortical afferents to the nucleus reticularis tegmenti pontis and basal pontine nuclei in the macaque monkey. *Vis Neurosci* 2001;18:725-740.
67. Matsuzaki R, Kyuhou S. Pontine neurons which relay projections from the superior colliculus to the posterior vermis of the cerebellum in the cat: distribution and visual properties. *Neurosci Lett* 1997;236:99-102.
68. Hashimoto M, Ohtsuka K. Transcranial magnetic stimulation over the posterior cerebellum during visually guided saccades in man. *Brain* 1995;118:1185-1193.
69. Ilg W, Giese MA, Gizewski ER, Schoch B, Timmann D. The influence of focal cerebellar lesions on the control and adaptation of gait. *Brain* 2008;131:2913-2927.
70. Zwergal A, la Fougère C, Lorenzl S, Rominger A, Xiong G, Deutschenbaur L, et al. Functional disturbance of the locomotor network in progressive supranuclear palsy. *Neurology* 2013;80:634-641.
71. Schlag J, Schlag-Rey M. Role of the central thalamus in gaze control. *Prog Brain Res* 1986;64:191-201.
72. Watanabe Y, Funahashi S. Neuronal activity throughout the primate mediodorsal nucleus of the thalamus during oculomotor delayed-responses. II. Activity encoding visual versus motor signal. *J Neurophysiol* 2004;92:1756-1769.
73. Kunimatsu J, Tanaka M. Roles of the primate motor thalamus in the generation of antisaccades. *J Neurosci* 2010;30:5108-5117.
74. Roth MM, Dahmen JC, Muir DR, Imhof F, Martini FJ, Hofer SB. Thalamic nuclei convey diverse contextual information to layer 1 of visual cortex. *Nat Neurosci* 2016;19:299-307.
75. Robinson DL, McClurkin JW. The visual superior colliculus and pulvinar. *Rev Oculomot Res* 1989;3:337-360.
76. Tanaka M, Kunimatsu J. Contribution of the central thalamus to the generation of volitional saccades. *Eur J Neurosci* 2011;33:2046-2057.
77. Kronenbueger M, González EG, Liu LD, Moro E, Steinbach MJ, Lozano AM, et al. Involvement of the human ventrolateral thalamus in the control of visually guided saccades. *Brain Stimul* 2010;3:226-229.
78. Armstrong DM, Drew T. Discharges of pyramidal tract and other motor cortical neurones during locomotion in the cat. *J Physiol* 1984;346:471-495.
79. Drew T. Motor cortical activity during voluntary gait modifications in the cat. I. Cells related to the forelimbs. *J Neurophysiol* 1993;70:179-199.
80. Pinault D. The thalamic reticular nucleus: structure, function and concept. *Brain Res Brain Res Rev* 2004;46:1-31.
81. Wurtz RH, McAlonan K, Cavanaugh J, Berman RA. Thalamic pathways for active vision. *Trends Cogn Sci* 2011;15:177-184.
82. Marlinski V, Beloozerova IN. Burst firing of neurons in the thalamic reticular nucleus during locomotion. *J Neurophysiol* 2014;112:181-192.
83. Watanabe M, Munoz DP. Probing basal ganglia functions by saccade eye movements. *Eur J Neurosci* 2011;33:2070-2090.
84. Watanabe M, Munoz DP. Saccade reaction times are influenced by caudate microstimulation following and prior to visual stimulus appearance. *J Cogn Neurosci* 2011;23:1794-1807.
85. Ding L, Gold JJ. Separate, causal roles of the caudate in saccadic choice and execution in a perceptual decision task. *Neuron* 2012;75:865-874.
86. Phillips JM, Everling S. Neural activity in the macaque putamen associated with saccades and behavioral outcome. *PLoS One* 2012;7:e51596.
87. Kravitz AV, Freeze BS, Parker PR, Kay K, Thwin MT, Deisseroth K, et al. Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. *Nature* 2010;466:622-626.
88. Jahanshahi M, Obeso I, Rothwell JC, Obeso JA. A fronto-striato-subthalamic-pallidal network for goal-directed and habitual inhibition. *Nat Rev Neurosci* 2015;16:719-732.
89. Van Der Kooy D, Hattori T. Single subthalamic nucleus neurons project to both the globus pallidus and substantia nigra in rat. *J Comp Neurol* 1980;192:751-768.
90. Canteras NS, Shammah-Lagnado SJ, Silva BA, Ricardo JA. Afferent connections of the subthalamic nucleus: a combined retrograde and anterograde horseradish peroxidase study in the rat. *Brain Res* 1990;513:43-59.
91. Kita T, Kita H. The subthalamic nucleus is one of multiple innervation sites for long-range corticofugal axons: a single-axon tracing study in the rat. *J Neurosci* 2012;32:5990-5999.
92. Nilsson MH, Patel M, Rehncrona S, Magnusson M, Fransson PA. Subthalamic deep brain stimulation improves smooth pursuit and saccade performance in patients with Parkinson's disease. *J Neuroeng Rehabil* 2013;10:33.
93. Lohnes CA, Earhart GM. Effect of subthalamic deep brain stimulation on turning kinematics and related saccadic eye movements in Parkinson disease. *Exp Neurol* 2012;236:389-394.
94. Ferrarin M, Carpinella I, Rabuffetti M, Rizzone M, Lopiano L, Crenna P. Unilateral and bilateral subthalamic nucleus stimulation in Parkinson's disease: effects on EMG signals of lower limb muscles during walking. *IEEE Trans Neural Syst Rehabil Eng* 2007;15:182-189.
95. Chang JY, Shi LH, Luo F, Woodward DJ. High frequency stimulation of the subthalamic nucleus improves treadmill locomotion in unilateral 6-hydroxydopamine lesioned rats. *Brain Res* 2003;983:174-184.
96. Pötter-Nerger M, Volkmann J. Deep brain stimulation for gait and postural symptoms in Parkinson's disease. *Mov Disord* 2013;28:1609-1615.
97. Fridley J, Adams G, Sun P, York M, Atassi F, Lai E, et al. Effect of subthalamic nucleus or globus pallidus interna stimulation on oculomotor function in patients with Parkinson's disease. *Stereotact Funct Neurosurg* 2013;91:113-121.
98. Antoniadou CA, Rebelo P, Kennard C, Aziz TZ, Green AL, FitzGerald JJ. Pallidal deep brain stimulation improves higher control of the oculomotor system in Parkinson's disease. *J Neurosci* 2015;35:13043-13052.
99. Saitoh K, Ménard A, Grillner S. Tectal control of locomotion, steering, and eye movements in lamprey. *J Neurophysiol* 2007;97:3093-3108.
100. von Uckeremann G, Lambert FM, Combes D, Straka H, Simmers J. Adaptive plasticity of spino-extraocular motor coupling during locomotion in metamorphosing *Xenopus laevis*. *J Exp Biol* 2016;219:1110-1121.
101. Schwarz M, Sontag KH, Wand P. Sensory-motor processing in substantia nigra pars reticulata in conscious cats. *J Physiol* 1984;347:129-147.
102. Hikosaka O, Wurtz RH. Visual and oculomotor functions

- of monkey substantia nigra pars reticulata. I. Relation of visual and auditory responses to saccades. *J Neurophysiol* 1983;49:1230-1253.
103. Winn P. Experimental studies of pedunculopontine functions: are they motor, sensory or integrative? *Parkinsonism Relat Disord* 2008;14 Suppl 2:S194-S198.
 104. Ohtsuka K, Edamura M, Kawahara K, Aoki M. The properties of goal-directed eye movements evoked by microstimulation of the cerebellar vermis in the cat. *Neurosci Lett* 1987;76:173-178.
 105. Godschalk M, Van der Burg J, Van Duin B, De Zeeuw CI. Topography of saccadic eye movements evoked by microstimulation in rabbit cerebellar vermis. *J Physiol* 1994;480:147-153.
 106. El-Shamayleh Y, Kojima Y, Soetedjo R, Horwitz GD. Selective optogenetic control of Purkinje cells in monkey cerebellum. *Neuron* 2017;95:51-62.e4.
 107. Sauerbrei BA, Lubenov EV, Siapas AG. Structured variability in Purkinje cell activity during locomotion. *Neuron* 2015;87:840-852.
 108. Hoogland TM, De Groot JR, Witter L, Canto CB, De Zeeuw CI. Role of synchronous activation of cerebellar Purkinje cell ensembles in multi-joint movement control. *Curr Biol* 2015;25:1157-1165.
 109. Andersson G, Armstrong DM. Complex spikes in Purkinje cells in the lateral vermis (b zone) of the cat cerebellum during locomotion. *J Physiol* 1987;385:107-134.
 110. Gandhi NJ, Barton EJ, Sparks DL. Coordination of eye and head components of movements evoked by stimulation of the paramedian pontine reticular formation. *Exp Brain Res* 2008;189:35-47.
 111. Gandhi NJ, Katnani HA. Motor functions of the superior colliculus. *Annu Rev Neurosci* 2011;34:205-231.
 112. Grasso R, Prévost P, Ivanenko YP, Berthoz A. Eye-head coordination for the steering of locomotion in humans: an anticipatory synergy. *Neurosci Lett* 1998;253:115-118.
 113. Saleem AB, Ayaz A, Jeffery KJ, Harris KD, Carandini M. Integration of visual motion and locomotion in mouse visual cortex. *Nat Neurosci* 2013;16:1864-1869.
 114. Imai T, Moore ST, Raphan T, Cohen B. Interaction of the body, head, and eyes during walking and turning. *Exp Brain Res* 2001;136:1-18.
 115. Wilkie RM, Wann JP. Judgments of path, not heading, guide locomotion. *J Exp Psychol Hum Percept Perform* 2006;32:88-96.
 116. Rietdyk S, Rhea CK. Control of adaptive locomotion: effect of visual obstruction and visual cues in the environment. *Exp Brain Res* 2006;169:272-278.
 117. Vitória R, Lirani-Silva E, Barbieri FA, Raile V, Stella F, Gobbi LT. Influence of visual feedback sampling on obstacle crossing behavior in people with Parkinson's disease. *Gait Posture* 2013;38:330-334.
 118. Stuart S, Alcock L, Galna B, Lord S, Rochester L. The measurement of visual sampling during real-world activity in Parkinson's disease and healthy controls: a structured literature review. *J Neurosci Methods* 2014;222:175-188.
 119. Land MF, Lee DN. Where we look when we steer. *Nature* 1994;369:742-744.
 120. Land MF. Motion and vision: why animals move their eyes. *J Comp Physiol A* 1999;185:341-352.
 121. Land MF. Eye movements of vertebrates and their relation to eye form and function. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 2015;201:195-214.
 122. Patla AE, Tomescu SS, Greig M, Novak A. Gaze fixation patterns during goal-directed locomotion while navigating around obstacles and a new route-selection model. In: van Gompel RPG, Fischer MH, Murray WS, Hill RL, editors. *Eye Movements: A Window on Mind and Brain*. 1st ed. Amsterdam: Elsevier, 2007;677-696.
 123. Higuchi T. Visuomotor control of human adaptive locomotion: understanding the anticipatory nature. *Front Psychol* 2013;4:277.
 124. Rivers TJ, Sirota MG, Guttentag AI, Ogorodnikov DA, Shah NA, Beloozerova IN. Gaze shifts and fixations dominate gaze behavior of walking cats. *Neuroscience* 2014;275:477-499.
 125. Foulsham T. Eye movements and their functions in everyday tasks. *Eye (Lond)* 2015;29:196-199.
 126. Matthis JS, Barton SL, Fajen BR. The biomechanics of walking shape the use of visual information during locomotion over complex terrain. *J Vis* 2015;15:10.
 127. Matthis JS, Fajen BR. Visual control of foot placement when walking over complex terrain. *J Exp Psychol Hum Percept Perform* 2014;40:106-115.
 128. Marigold DS, Patla AE. Gaze fixation patterns for negotiating complex ground terrain. *Neuroscience* 2007;144:302-313.
 129. Hollands MA, Marple-Horvat DE. Coordination of eye and leg movements during visually guided stepping. *J Mot Behav* 2001;33:205-216.
 130. Patla AE, Vickers JN. How far ahead do we look when required to step on specific locations in the travel path during locomotion? *Exp Brain Res* 2003;148:133-138.
 131. Yamada M, Higuchi T, Mori S, Uemura K, Nagai K, Aoyama T, et al. Maladaptive turning and gaze behavior induces impaired stepping on multiple footfall targets during gait in older individuals who are at high risk of falling. *Arch Gerontol Geriatr* 2012;54:e102-e108.
 132. Greany JF, Di Fabio RP. Saccade to stepping delays in elders at high risk for falling. *Aging Clin Exp Res* 2008;20:428-433.
 133. Hollands MA, Patla AE, Vickers JN. "Look where you're going!": gaze behaviour associated with maintaining and changing the direction of locomotion. *Exp Brain Res* 2002;143:221-230.
 134. Young WR, Hollands MA. Newly acquired fear of falling leads to altered eye movement patterns and reduced stepping safety: a case study. *PLoS One* 2012;7:e49765.
 135. Young WR, Wing AM, Hollands MA. Influences of state anxiety on gaze behavior and stepping accuracy in older adults during adaptive locomotion. *J Gerontol B Psychol Sci Soc Sci* 2012;67:43-51.
 136. Kugler G, Huppert D, Eckl M, Schneider E, Brandt T. Visual exploration during locomotion limited by fear of heights. *PLoS One* 2014;9:e105906.
 137. Gonzalez Andino SL, Grave de Peralta Menendez R. Coding of saliency by ensemble bursting in the amygdala of primates. *Front Behav Neurosci* 2012;6:38.
 138. Crowdy KA, Marple-Horvat DE. Alcohol affects eye movements essential for visually guided stepping. *Alcohol Clin Exp Res* 2004;28:402-407.
 139. Jäntti V, Lang AH, Keskinen E, Lehtinen I, Pakkanen A. Acute effects of intravenously given alcohol on saccadic eye movements and subjective evaluations of intoxication. *Psychopharmacology (Berl)* 1983;79:251-255.
 140. Gitchel GT, Wetzel PA, Baron MS. Slowed saccades and increased square wave jerks in essential tremor. *Tremor Other Hyperkinet Mov (N Y)* 2013;3:tre-03-178-4116-2.
 141. Singer C, Sanchez-Ramos J, Weiner WJ. Gait abnormality in essential tremor. *Mov Disord* 1994;9:193-196.

142. Hubble JP, Busenbark KL, Pahwa R, Lyons K, Koller WC. Clinical expression of essential tremor: effects of gender and age. *Mov Disord* 1997;12:969-972.
143. Stolze H, Petersen G, Raethjen J, Wenzelburger R, Deuschl G. The gait disorder of advanced essential tremor. *Brain* 2001;124:2278-2286.
144. Lim ES, Seo MW, Woo SR, Jeong SY, Jeong SK. Relationship between essential tremor and cerebellar dysfunction according to age. *J Clin Neurol* 2005;1:76-80.
145. Kronenbuerger M, Konczak J, Ziegler W, Buderath P, Frank B, Coenen VA, et al. Balance and motor speech impairment in essential tremor. *Cerebellum* 2009;8:389-398.
146. Fasano A, Herzog J, Raethjen J, Rose FE, Muthuraman M, Volkmann J, et al. Gait ataxia in essential tremor is differentially modulated by thalamic stimulation. *Brain* 2010;133:3635-3648.
147. Rao AK, Gillman A, Louis ED. Quantitative gait analysis in essential tremor reveals impairments that are maintained into advanced age. *Gait Posture* 2011;34:65-70.
148. Hoskocová M, Ulmanová O, Sprdlík O, Sieger T, Nováková J, Jech R, et al. Disorders of balance and gait in essential tremor are associated with midline tremor and age. *Cerebellum* 2013;12:27-34.
149. Rao AK, Louis ED. Timing control of gait: a study of essential tremor patients vs. age-matched controls. *Cerebellum Ataxias* 2016;3:5.
150. MacAskill MR, Anderson TJ. Eye movements in neurodegenerative diseases. *Curr Opin Neurol* 2016;29:61-68.
151. Srivastava A, Sharma R, Sood SK, Shukla G, Goyal V, Behari M. Saccadic eye movements in Parkinson's disease. *Indian J Ophthalmol* 2014;62:538-544.
152. Bloem BR, Hausdorff JM, Visser JE, Giladi N. Falls and freezing of gait in Parkinson's disease: a review of two interconnected, episodic phenomena. *Mov Disord* 2004;19:871-884.
153. Svehlík M, Zwick EB, Steinwender G, Linhart WE, Schwingenschuh P, Katschnig P, et al. Gait analysis in patients with Parkinson's disease off dopaminergic therapy. *Arch Phys Med Rehabil* 2009;90:1880-1886.
154. Otero-Millan J, Serra A, Leigh RJ, Troncoso XG, Macknik SL, Martinez-Conde S. Distinctive features of saccadic intrusions and microsaccades in progressive supranuclear palsy. *J Neurosci* 2011;31:4379-4387.
155. Chen AL, Riley DE, King SA, Joshi AC, Serra A, Liao K, et al. The disturbance of gaze in progressive supranuclear palsy: implications for pathogenesis. *Front Neurol* 2010;1:147.
156. Hatanaka N, Sato K, Hishikawa N, Takemoto M, Ohta Y, Yamashita T, et al. Comparative gait analysis in progressive supranuclear palsy and Parkinson's disease. *Eur Neurol* 2016;75:282-289.
157. Amano S, Skinner JW, Lee HK, Stegemöller EL, Hack N, Akbar U, et al. Discriminating features of gait performance in progressive supranuclear palsy. *Parkinsonism Relat Disord* 2015;21:888-893.
158. Lasker AG, Zee DS. Ocular motor abnormalities in Huntington's disease. *Vision Res* 1997;37:3639-3645.
159. Golding CV, Danchaivijitr C, Hodgson TL, Tabrizi SJ, Kennard C. Identification of an oculomotor biomarker of preclinical Huntington disease. *Neurology* 2006;67:485-487.
160. Hicks SL, Robert MP, Golding CV, Tabrizi SJ, Kennard C. Oculomotor deficits indicate the progression of Huntington's disease. *Prog Brain Res* 2008;171:555-558.
161. Patel SS, Jankovic J, Hood AJ, Jeter CB, Sereno AB. Reflexive and volitional saccades: biomarkers of Huntington disease severity and progression. *J Neurol Sci* 2012;313:35-41.
162. Blekher T, Johnson SA, Marshall J, White K, Hui S, Weaver M, et al. Saccades in presymptomatic and early stages of Huntington disease. *Neurology* 2006;67:394-399.
163. Peltsch A, Hoffman A, Armstrong I, Pari G, Munoz DP. Saccadic impairments in Huntington's disease. *Exp Brain Res* 2008;186:457-469.
164. Winograd-Gurvich CT, Georgiou-Karistianis N, Evans A, Millist L, Bradshaw JL, Churchyard A, et al. Hypometric primary saccades and increased variability in visually-guided saccades in Huntington's disease. *Neuropsychologia* 2003;41:1683-1692.
165. Koller WC, Trimble J. The gait abnormality of Huntington's disease. *Neurology* 1985;35:1450-1454.
166. Rao AK, Muratori L, Louis ED, Moskowitz CB, Marder KS. Spectrum of gait impairments in presymptomatic and symptomatic Huntington's disease. *Mov Disord* 2008;23:1100-1107.
167. Thaut MH, Miltner R, Lange HW, Hurt CP, Hoemberg V. Velocity modulation and rhythmic synchronization of gait in Huntington's disease. *Mov Disord* 1999;14:808-819.
168. Bilney B, Morris ME, Churchyard A, Chiu E, Georgiou-Karistianis N. Evidence for a disorder of locomotor timing in Huntington's disease. *Mov Disord* 2005;20:51-57.
169. Termsarasab P, Thammongkolchai T, Rucker JC, Frucht SJ. The diagnostic value of saccades in movement disorder patients: a practical guide and review. *J Clin Mov Disord* 2015;2:14.
170. Christova P, Anderson JH, Gomez CM. Impaired eye movements in presymptomatic spinocerebellar ataxia type 6. *Arch Neurol* 2008;65:530-536.
171. Federighi P, Cevenini G, Dotti MT, Rosini F, Pretelegiani E, Federico A, et al. Differences in saccade dynamics between spinocerebellar ataxia 2 and late-onset cerebellar ataxias. *Brain* 2011;134:879-891.
172. Stolze H, Klebe S, Petersen G, Raethjen J, Wenzelburger R, Witt K, et al. Typical features of cerebellar ataxic gait. *J Neurol Neurosurg Psychiatry* 2002;73:310-312.
173. Buckley E, Mazzà C, McNeill A. A systematic review of the gait characteristics associated with cerebellar ataxia. *Gait Posture* 2018;60:154-163.
174. Palliyath S, Hallett M, Thomas SL, Lebedowska MK. Gait in patients with cerebellar ataxia. *Mov Disord* 1998;13:958-964.
175. Boonstra TA, van der Kooij H, Munneke M, Bloem BR. Gait disorders and balance disturbances in Parkinson's disease: clinical update and pathophysiology. *Curr Opin Neurol* 2008;21:461-471.
176. Nemanich ST, Earhart GM. Freezing of gait is associated with increased saccade latency and variability in Parkinson's disease. *Clin Neurophysiol* 2016;127:2394-2401.
177. Walton CC, O'Callaghan C, Hall JM, Gilat M, Mowszowski L, Naismith SL, et al. Antisaccade errors reveal cognitive control deficits in Parkinson's disease with freezing of gait. *J Neurol* 2015;262:2745-2754.
178. Ambati VN, Saucedo F, Murray NG, Powell DW, Reed-Jones RJ. Constraining eye movement in individuals with Parkinson's disease during walking turns. *Exp Brain Res* 2016;234:2957-2965.
179. Galna B, Lord S, Daud D, Archibald N, Burn D, Rochester L. Visual sampling during walking in people with Parkinson's disease and the influence of environment and dual-task. *Brain Res* 2012;1473:35-43.
180. Lohnes CA, Earhart GM. Saccadic eye movements are re-

- lated to turning performance in Parkinson disease. *J Parkinsons Dis* 2011;1:109-118.
181. Ewenczyk C, Mesmoudi S, Gallea C, Welter ML, Gaymard B, Demail A, et al. Antisaccades in Parkinson disease: a new marker of postural control? *Neurology* 2017; 88:853-861.
 182. Mancini M, Carlson-Kuhta P, Zampieri C, Nutt JG, Chiari L, Horak FB. Postural sway as a marker of progression in Parkinson's disease: a pilot longitudinal study. *Gait Posture* 2012;36:471-476.
 183. Stylianou AP, McVey MA, Lyons KE, Pahwa R, Luchies CW. Postural sway in patients with mild to moderate Parkinson's disease. *Int J Neurosci* 2011;121:614-621.
 184. Takamatsu Y, Matsuda N, Aiba I. Body sway during static standing in patients with progressive supranuclear palsy. *J Neurol Sci* 2017;381:836.
 185. Umemura K, Ishizaki H, Matsuoka I, Hoshino T, Nozue M. Analysis of body sway in patients with cerebellar lesions. *Acta Otolaryngol Suppl* 1989;468:253-261.
 186. Arkadir D, Louis ED. The balance and gait disorder of essential tremor: what does this mean for patients? *Ther Adv Neurol Disord* 2013;6:229-236.
 187. Panzera R, Salomonczyk D, Pirogovosky E, Simmons R, Goldstein J, Corey-Bloom J, et al. Postural deficits in Huntington's disease when performing motor skills involved in daily living. *Gait Posture* 2011;33:457-461.
 188. Grimbergen YAM, Knol MJ, Bloem BR, Kremer BPH, Roos RAC, Munneke M. Falls and gait disturbances in Huntington's disease. *Mov Disord* 2008;23:970-976.
 189. Legrand A, Mazars KD, Lazzareschi J, Lemoine C, Olivier I, Barra J, et al. Differing effects of prosaccades and antisaccades on postural stability. *Exp Brain Res* 2013;227: 397-405.
 190. Rodrigues ST, Aguiar SA, Polastri PF, Godoi D, Moraes R, Barela JA. Effects of saccadic eye movements on postural control stabilization. *Motriz: Revista de Educação Física* 2013;19:614-619.
 191. Crowdy KA, Hollands MA, Ferguson IT, Marple-Horvat DE. Evidence for interactive locomotor and oculomotor deficits in cerebellar patients during visually guided stepping. *Exp Brain Res* 2000;135:437-454.
 192. Marple-Horvat DE, Crowdy KA. Direct visualisation of gaze and hypometric saccades in cerebellar patients during visually guided stepping. *Gait Posture* 2005;21:39-47.
 193. Zampieri C, Di Fabio RP. Balance and eye movement training to improve gait in people with progressive supranuclear palsy: quasi-randomized clinical trial. *Phys Ther* 2008;88:1460-1473.
 194. Crowdy KA, Kaur-Mann D, Cooper HL, Mansfield AG, Offord JL, Marple-Horvat DE. Rehearsal by eye movement improves visuomotor performance in cerebellar patients. *Exp Brain Res* 2002;146:244-247.
 195. Kang KY, Yu KH. The effects of eye movement training on gait function in patients with stroke. *J Phys Ther Sci* 2016; 28:1816-1818.
 196. Hollands MA, Marple-Horvat DE, Henkes S, Rowan AK. Human eye movements during visually guided stepping. *J Mot Behav* 1995;27:155-163.
 197. Stuart S, Galna B, Delicato LS, Lord S, Rochester L. Direct and indirect effects of attention and visual function on gait impairment in Parkinson's disease: influence of task and turning. *Eur J Neurosci* 2017;46:1703-1716.
 198. Paquette C, Fung J. Old age affects gaze and postural coordination. *Gait Posture* 2011;33:227-232.
 199. Di Fabio RP, Greany JF, Zampieri C. Saccade-stepping interactions revise the motor plan for obstacle avoidance. *J Mot Behav* 2003;35:383-397.
 200. Di Fabio RP, Zampieri C, Greany JF. Aging and saccade-stepping interactions in humans. *Neurosci Lett* 2003;339: 179-182.