

The Remodeling of Connexins Localized at Pulmonary Vein – Left Atria in Triggering and Maintenance of Atrial Fibrillation

Guo-qiang Zhong¹, Ri-xin Xiong², Hong-xing Song,
Yun Ling, Jing-chang Zhang and Zhe Wei

¹*The Department of Cardiology, The First Affiliated Hospital
of Guangxi Medical University,*

²*The Department of Cardiology, The People's Hospital
of Guangxi Zhuang Autonomous Region,
China*

1. Introduction

Atrial fibrillation is the most common sustained arrhythmia and the major cardiac cause of stroke (Sellers & Newby, 2011). Recent studies in patients and animals with paroxysmal atrial fibrillation had shown that the arrhythmia was triggered by focal sources from muscular sleeves originated in the left atria extending into pulmonary veins (Date et al., 2007; Chen et al., 2001; Patterson et al., 2007; Honjo et al., 2003). What is more, the autonomic nervous system has a crucial role in the genesis, maintenance and abruption of atrial fibrillation (Duffy & Wit, 2008; Sandres et al., 2004), and there is a fat pad localized in superior vena cava and the root of aorta as the origin of cardiac nerve, being called SVC-Ao fat pad (Kapa et al., 2010; Volders, 2010; Ji et al., 2010). The mechanism of atrial fibrillation involves multiple effects (Chiou et al., 1997; Tsuboi et al., 2000; Hoffmann et al., 2006; Haissaguerre et al., 1998; Wit & Boyden, 2007), the expression of connexins is changed and is associated with increased propensity for arrhythmias (Kanagaratnam et al., 2002; Herve et al., 2007; Stergiopoulos et al., 1999; Verheule S et al., 2002; Kanagaratnam et al., 2007; Valiunas et al., 2001; Valiunas et al., 2000; Cottrell et al., 2002; Elenes et al., 1999). Pulmonary veins and left atria are the primary structure of genesis, persistence of atrial fibrillation. Vagus nerve plays a key role in the initiation and maintenance of atrial fibrillation, it can mediate the electrical remodeling of atria and pulmonary vein, enhance the maintenance and stability of atrial fibrillation. However, these effects can be inhibited by avianizing vagus nerve.

2. Overview of gap junction structure

The name “gap junction” was coined from their appearance in electron micrographs in the 60s (Revel & Karnovsky, 1967). They are specialized at cell-cell contact regions that contain tens to thousands of intercellular channels that link two apposed cells. These channels facilitate a form of intercellular communication by permitting the regulated passage of ions and small molecules from one cell to another (Bennett & Goodenough, 1978). The gap junction

membrane channel is composed of macular aggregations of intercellular channels permitting the direct intercellular transfer of ions and small molecules. Each intercellular channel is formed by the apposition of two hexameric transmembrane channels (connexons), one from each cell (Yeager & Gilula, 1992). Each intercellular channel is composed of two oligomers with each of two adjacent tissue cells contributing one oligomer. Each connexon is built from six copies of one or more members of a protein family called the connexins (Unger et al., 1999; Perkins et al., 1998). The functional cell-cell channel is formed by the end-to-end docking of the extracellular domains of the two connexons. Therefore, the gap junction membrane channel can be thought of as a dimer of two hexamers joined together in the gap region. In this manner, the membrane channel extends across both cell membranes.

Vertebrate gap junction channels are assembled to form multimers of one or more different proteins from a multigene family of homologous protein, which are composed of an entirely different gene family but the sequences predict a similar folding pattern. Up to now, more than 20 different connexin genes have been identified in the mouse genome and at least 6 others have been identified in other vertebrates. Connexins are highly homologous proteins with 50%-80% identity between amino acid sequences and display considerable amino acid sequences conservation between species. The distribution and developmental regulation of connexins are tissue specific. The predicted molecular mass of the connexin protein family ranges from ~26 to ~60KDa, and the proteins are named "connexin" followed by their predicted molecular mass. The connexin family can be subdivided into two classifications called α and β based on similarities in certain regions of the primary sequence. The x-ray diffraction analysis by Tibbitts et al. (1990) indicated that there was more α -helical content that could be accounted for by four transmembrane helices. The E1 and E2 loops are thought to be as rigid as the transmembrane domain (Hoh et al., 1993; Sosinsky, 1992). Hence, the extracellular region is visible with the crystallographic averaging used. Each extracellular protrusion may therefore include an extension of the intramembrane α -helical structure (Tibbitts et al., 1990). Mutagenesis studies have suggested that the extracellular loops contain disulfide-bonded β -sheet conformation (Foote & Nicholson, 1997), which would be expected to act as a rigid domain, for instance, connexin43 (isolated from heart tissue, Yeager & Gilula, 1992), connexin32 and connexin 26 (isolated from liver tissue, Fallon & Goodenough, 1981; Hertzberg, 1984) would be seen for the same structure. In the human heart, 4 main isoforms are expressed. Connexin43 is expressed in all chambers of the heart, but predominantly in the ventricles. Connexin45 is found in the conduction system of the heart and at low levels in the atrial and ventricular working myocardium and connexin37 is located in the endothelial gap junctions in many vessels. Finally, connexin40 is expressed mainly in the atrial working myocardium, the conduction system, and the vasculature. Connexin40 was first described in a range of animal species and subsequently mapped to human chromosome. It became apparent that connexin40 was expressed in the atrioventricular conduction system and abundantly expressed in the atrial but not in the ventricular gap junctions. Recently, a new connexin was described in the mouse heart, i.e. connexin30.2 (the human equivalent is connexin31.9), which in mice seems responsible for slowing of impulse conduction in the atrioventricular node. However, the role of connexin31.9 in the human heart is unclear, for it is not detectable in the human cardiac conduction system. Connexons formed the functional intercellular channel, which is defined as homotypic when the connexin composition of the contributing connexons is identical or heterotypic when different. The ability of connexins to form homomeric/ heterotypic channels has been examined in the *Xenopus* oocyte and HeLa

cell expression systems as well as in other settings. Connexin43 and connexin40 are co-expressed in several tissues, including cardiac atrial and ventricular myocytes and vascular smooth muscle. It has been shown that these connexins form functional homomeric/homotopic channels with distinct permeability and gating properties.

3. Connexon and the heart development

In the vertebrate embryo, a single outflow vessel initially develops from the common ventricles undergo septation to form the four-chamber heart (Kirby & Waldo, 1995). The four-chambered heart develops from a single straight tube composed of three layers: an inner endocardium and outer myocardium separated by a thick extracellular matrix called cardiac jelly. The growing tube loops, with the convexity of the loop demarcating a functional inflow from an outflow portion of the looped tube. This original convexity forms the two ventricles by expansion and septation, and as the ventricular septum forms, the inflow and outflow must be redefined. Initially, all of the inflow is through the atrioventricular canal connecting a common atrium to the presumptive left ventricle; the entire outflow is from the presumptive right ventricle into the aortic sac via the conotruncus. Several septa are formed simultaneously, making a rearrangement of inflow and outflow critical. The atrioventricular canal is divided into left and right channels, the nascent right and left ventricles are separated by the ventricular septum, the conotruncus is converted into aortic vestibule and semilunar valve continuous with the left ventricle, and the pulmonary infundibulum and semilunar valve originating from the right ventricle (see Fig1.). Concomitantly, the outflow tract septates and rotates to generate the pulmonary and




Fig. 1. Major events in development of the heart from a single straight tube. Early looping is probably a function of the myocardium. Convergence of the outflow and inflow tracts occurs during late looping and is critical for normal wedging. Wedging produces alignment of the three components that complete outflow septation. A, aorta; V, ventricle; P, pulmonary trunk; RA, right atrium; LA, left atrium; RV, right ventricle; LV, left ventricle.

aortic outflow tracts, becoming connected to the right and left ventricular chambers, respectively. This morphogenetic sequence is significantly affected by the activity of neural crest cells. With the development of heart, the co-expression of multiple connexins in diverse tissues support the idea that communication between cells does not depend on just one type of connexin, which results in forming heterotypic channels with properties different from those homotypic parental connexons.

Pluripotent mammalian embryonic stem cells, which are derived from the inner cell mass of preimplantation blastocysts, have the capacity to differentiate into cells of all three germ layers. Under suitable conditions, embryonic stem cells remain pluripotent through repeated rounds of cell division in culture. The recent isolation of human embryonic stem cells has spurred great interest in their potential use for therapeutic tissue repair because appropriate manipulation of the culture environment can induce both mouse and human embryonic stem cells to differentiate in vitro into specific somatic cell types, and the result shows human embryonic stem cells express RNA encoding most of the known human connexin genes. The mRNA of Cx25, Cx26, Cx30.3, Cx31, Cx31.1, Cx31.9, Cx37, Cx40, Cx43, Cx45, and Cx46 for all of the undifferentiated human embryonic stem cultures that were detected. The remaining connexins (Cx30, Cx30.2, Cx32, Cx36, Cx47, Cx59, and Cx62) were also reliably detected (Rackauskas et al., 2010).

4. Cardiac electrical properties of connexins and the methods for detection

For effective cardiac output, it is essential that electrical excitation spread rapidly throughout the atria and ventricles. This is effected by electrical coupling through gap junction channels at contact sites between myocytes (Danik te al., 2008; Harris, 2001; Söhl & Willecke, 2004; Moreno, 2004). These channels form a low-resistance pathway between adjacent myocytes and consist of connexin proteins. The connexin family is a large multigene family, and the channels formed by different members of this family have distinct electrical and regulatory properties (Reisner et al., 2009; Gros & Jongsma, 1996; Verheijck et al., 2001; Valiunas et al., 2002). Voltage sensitivity is particularly important in regulating the intercellular coupling between excitable cells. Cx43 channels are relatively insensitive to changes in transjunctional voltage compared with channels composed of Cx45 (Rackauskas et al., 2007; Bruzzone et al., 1993; Haubrich et al., 1996). Each gap junction composing hemichannel contains two V_j sensitive gates. The fast gate is located at the cytoplasmic entrance of hemichannels and operates from open to residual state. The slow, or “loop” gate is located toward extracellular ends of hemichannels and exhibits slow gating transition to the fully closed state. Cx26, Cx30, Cx50 close at positive voltages, and Cx31, Cx32, Cx37, Cx40, Cx43, Cx45, Cx57 at negative. Interestingly, Cx46 hemichannels close at both, positive and negative voltages. Besides, each type of connexon has a certain sensitivity to H⁺, it is showed the order of decreasing sensitivity to PH: Cx50>Cx46>Cx45>Cx26>Cx37>Cx43>Cx40>Cx32, anyway, it is not completely clear whether H⁺ acts directly on GJ channels. Cytoplasmic C-tail of connexins contains multiple serine, threonine, and tyrosine residues that may be phosphorylated by various protein kinases (Valiunas et al., 2000; Haubrich et al., 1996). Phosphorylation modifies electrical and metabolic communication between contiguous cells by changing channel molecular structure that affects channel unitary conductance, mean open time, or open probability. Moreover, phosphorylation alters the net charge of C-terminus that in turn may modulate voltage or pH sensitivity of the connexins (Kirchhoff et al., 2000, 1998; Bukauskas et al., 2004).

From the gene level to protein expression, novel methods of detection of gap junctions can lead us to realize the therapeutic potential, including morphological diversities, ultrastructure and spatial heterogeneity. The authors induced rapid persistent atrial pacing dogs and did a comparison between removal and retention of cardiac vagus nerve in dogs (superior vena cava and aorta root fat pad, SVC-Ao fat pad), then recorded the effective refractory period and the dispersion of effective refractory period, analyzed the expression of atrial and pulmonary vein connexin40, connexin43 and the heterogeneity of its spatial distribution and the changes of atrial collagen volume fraction, tried to illuminate the correlation of atrial fibrillation with connexin40, connexin43 remodeling and the function of vagus nerve. There was obvious shortness of effective refractory period in atrial fibrillation animal models, while the removal of vagus nerve could attenuate this effect, meanwhile, the quantity and distribution of connexin40 and connexin43 changed at different locations and the deposit of collagen fibers, which might explain the mechanism of electrical heterogeneity and the structural remodeling originated from the pulmonary vein and left atria for the atrial fibrillation.

5. The distribution of connexins in heart and the relationship with arrhythmia

Gap junction remodeling is a common response to many forms of heart disease (Strom et al.,2010; Burstein et al.,2009; Kieken et al.,2009; Qu et al., 2009; Yamada et al.,2008). Previous studies have demonstrated that loss of cell-cell coupling is highly arrhythmic (Chaldoupi et al., 2009), however, a detailed understanding of arrhythmia dynamics has been lacking. With respect to mechanistically link a primary perturbation of gap junction function with arrhythmia, conduction properties, arrhythmia dynamics, and cellular electrophysiological characteristics have been detected (Zhou et al., 2008; Severs et al., 2008; Gutstein et al., 2001). Accumulative evidences showed that ventricular gap junctions contain at least 20 times more connexin43 than connexin40, while atrial gap junctions contain much more connexin40 than connexin43 (Kontogeorgis et al., 2008; Lin et al., 2010). In the ventricle, the heterogeneous loss of connexin43 gap junctions in a murine conditional cardiac connexin43 knockout model best exemplified how the focal loss of cardiac gap junctions lead to significant dispersion of conduction, increased incidence of spontaneous arrhythmias, and loss of ventricular systolic function with only minor reductions in overall connexin43 expression. The gating of connexin43-containing ventricular gap junctions during the action potential is also proposed to promote cardiac arrhythmias via inactivation and recovery that depends on transjunctional voltage (V_j) and contributes to conduction slowing or block and the formation of reentrant arrhythmias. In the atria, targeted gene deletion of connexin40 in mice produced multiple aberrations: P wave and PQ interval prolongation, prolonged sinus-node-recovery time, prolonged Wenckebach period, burst-pacing induced atrial tachyarrhythmias, reduced atrial, A-V node, and left bundle branch conduction velocity, right bundle branch block, and reduced interatrial conduction heterogeneity(Leaf et al., 2008; Gutstein et al., 2005). There are close correlation between electrical properties and the anatomy of pulmonary veins and left atria. In animal studies, data showed that segmental muscle disconnection and differential muscle narrowing at pulmonary vein and left atria junctions and complex fiber orientations within the pulmonary vein provide robust anatomical bases for conduction disturbance at the pulmonary vein and left atria junction and complex intra pulmonary vein conduction patterns (Date et al., 2007).

6. The anatomy and electrical activation in pulmonary vein and left atria

Catheter radiofrequency ablation of triggers originated from the pulmonary veins may successfully terminate paroxysmal atrial fibrillation (Jais et al., 1994). Because ablation within the pulmonary veins may result in stenosis of the pulmonary veins, the junctions between the pulmonary veins and the left atria have become new ablation targets for achieving electrical pulmonary vein isolation (Haissaguerre et al., 1998, 2000). Segmental ablation can successfully isolate the pulmonary vein by placing radiofrequency lesions in only 21% to 59% of the circumference of the pulmonary veins ostia (Pappone et al., 2000; Oral et al., 2002). Previous studies showed that ectopic beats originated from the pulmonary vein propagated to the left atria with characteristically long conduction time, often with conduction delay or block within the pulmonary vein or at the pulmonary vein and left atria junction (Nathan & Eliakim, 1966). The complex arrangement of myocardial fibers in the pulmonary vein and/or in the pulmonary vein and left atria junction is a possible reason for conduction delay or block in the pulmonary vein and left atria junction and within the pulmonary vein (Saito et al., 2000). Ho et al reported that a differential thickness of muscle sleeves could also account for a variable safety factor of propagation across the pulmonary vein and left atria junction (Ho et al., 2001). Hocini reported that zones of activation delay were observed in canine pulmonary veins and correlated with abrupt changes in fascicle orientation (Hocini et al., 2002). Hamabe identified that segmental muscle disconnection and differential muscle narrowing at pulmonary vein and left atria junctions and complex fiber orientations within the pulmonary vein provide robust anatomical bases for conduction disturbance at the pulmonary vein and left atria junction and complex intra pulmonary vein conduction patterns (Hamabe et al., 2003, see Fig2. and Fig3).




Fig. 2. Ectopic excitement existing in pulmonary veins and left atria junction, propagating into atria and forming reentry, which is so called "Atrial fibrillation begets Atrial fibrillation". SVC, superior vena cava; IVC, inferior vena cava; LIPV, left inferior pulmonary vein; RIPV, right inferior pulmonary vein; LSPV: left superior pulmonary vein; RSPV: right superior pulmonary vein. LA: left atria; RA: right atria.




Fig. 3. Electrical remodeling of atria, Acetylcholine injecting at RAA activates ganglion to output impulse and effects on the excited foci in pulmonary veins, therefore, the arrhythmia perpetuates. Ach, acetylcholine; RAA: right atrial appendage; RSPV: right superior pulmonary vein; LSPV: left superior pulmonary vein. (Lu et al. 2008)

The autopsied heart showed that myocardial cells were localized to pulmonary vein between 9 and 38 mm from the pulmonary vein and left atria junction (Roux et al., 2004). The sleeve was composed of circularly and longitudinally oriented bundles of cardiomyocytes. The peripheral end of the myocardial sleeve was irregular. The longest myocardial sleeves were found in the superior veins and were longitudinally oriented. At the pulmonary vein and left atria junction, the circular bundles were not often circumferential. Pulmonary vein myocardial architecture confirmed the possibility of initiating atrial fibrillation. Tan (Tan et al., 2006) examined 192 sections in 32 veins obtained from 8 healthy human heart, each segment included between 7 and 20 mm of the adjoining left atria, and found 1) pulmonary vein and left atria muscular discontinuities and abrupt 90° changes in myofiber orientation were present in the pulmonary vein and left atria junction in over half of all segments examined; 2) These anisotropic features were found more frequently in the anterosuperior than posteroinferior junctions; 3) Adrenergic and cholinergic nerve densities were highest in the left atria within 5 mm from the pulmonary vein and left atria junction, higher in the superior aspect of left superior pulmonary vein, anterosuperior aspect of right superior pulmonary vein, and inferior aspects of the both inferior pulmonary veins than diametrically opposite, and higher in the epicardial than endocardial half of the tissue; 4) adrenergic and cholinergic nerves were highly co-located at tissue and cellular levels of spatial organization. Tan suggest that: 1) the pulmonary vein and left atria junction contains anatomical substrates (muscular discontinuities and abrupt fiber orientation changes) to support reentry; 2) there are no good empiric targets for segmental pulmonary vein isolation because of the widespread distributions of pulmonary vein and left atria muscular discontinuities; 3) the left atria region close to the pulmonary vein and left atria junction rather than farther away in the left atria or pulmonary vein would be the most appropriate target for autonomic modulation procedures; 4) it is not possible to selectively ablate either adrenergic or cholinergic nerves in this location because

both nerve types are highly co-located in this region. Steiner found that pulmonary vein myocardial sleeves frequently harbour pathological lesions, particularly senile atrial amyloid, and scarring in atrial fibrillation (Steiner et al., 2005). The degree of scarring of the sleeves did not correlate with the degree of coronary atherosclerosis, and inferred the genesis of the scarring is not post necrotic but degenerative, due to diffuse hypoxia of the sleeve myocardium. Amyloidosis and particularly scarring of pulmonary vein myocardial sleeves appear generally in the elderly population as an arrhythmogenic substrate for atrial fibrillation.

In animal studies, pacing had different effects on connexin40 and connexin43 gap junctions, collagen content increased as well (Yeh et al., 2006). They found there was a 98% increase in connexin43 in pacing 2 weeks, and a 74% increase in pacing 6-8weeks animals. In contrast, connexin40 decreased 47% in pacing 2 weeks but increased 44% in pacing 6-8weeks animals. Our studies also found the moderate to severe deposit of collagen fibers in canine atria after persistent rapid atrial pacing (XIONG et al., 2010, see Fig4. and Fig5.), connexin40 mRNA expression decreased in left atria and right atria, but increased in left atria appendage, right atria appendage and atrial septum; connexin43 mRNA expression was reduced in left atria, right atria, left atrial appendage and right atrial appendage while increased in atrial septum (see Fig6., Fig7. and Fig8.). The pacing induced collagen remodeling and modulation on connexin40 mRNA and connexin43 mRNA expressions could be partially attenuated by removing SVC-Ao fat pad suggesting vagal nervation plays a key role in the initiation and preservation of atrial fibrillation .




Fig. 4. Different degree of collagen fibers deposits at intercellular substance.1A to 1E: normal distribution of collagen fibers in the whole atria; 2A to 2E: severe degree of collagen fibers deposits at different locations of atria after persistent atrial pacing; 3A to 3E: moderate degree of collagen fibers deposits at different locations of atria after persistent atrial pacing without SVC-Ao fat pad.A: left atria; B: right atria; C: left atrial appendage; D: right atrial appendage; E: atrial septum; SVC-Ao fat pad: superior vena cava and aorto root fat pad.




Fig. 5. Collagen fibers wrap the cardiac muscle cells and interrupt the normal intercellular electrical and signal conduction.




Fig. 6. Collagen Volume Fraction compared among three different groups. There was no significant spatial difference on collagen distribution in sham operated group and SVC-Ao fat pad removal group, while significant spatial difference on collagen distribution in SVC-Ao fat pad reserved group. CVF: collagen volume fraction; SVC-Ao fat pad: superior vena cava and aorto root fat pad; LA: left atria; RA: right atria; LAA: left atrial appendage; RAA: right atrial appendage; AS: atrial septum.




Fig. 7. Expression of Cx40 mRNA compared among three different groups. There was significant difference of increased Cx40 mRNA expression after persistent atrial pacing, the expression at atria in SVC-Ao fat pad removal group but that at atrial appendage in SVC-Ao fat pad reserved group are the significant difference. SVC-Ao fat pad: superior vena cava and aorto root fat pad; LA: left atria; RA: right atria; LAA: left atrial appendage; RAA: right atrial appendage; AS: atrial septum; Cx40: connexin 40.




Fig. 8. Expression of Cx43 mRNA compared among three different groups. There was significant difference of increased Cx43 mRNA expression in SVC-Ao fat pad removal group but decreased Cx43 mRNA expression in SVC-Ao fat pad reserved group. The most interesting locations are left and right atria. SVC-Ao fat pad: superior vena cava and aorto root fat pad; LA: left atria; RA: right atria; LAA: left atrial appendage; RAA: right atrial appendage; AS: atrial septum; Cx43: connexin 43.

7. Acknowledgment

This work was supported by grants from the Science Foundation for Returnees of Guangxi Zhuang Autonomous Region of China (No. 0575014) and the Key Scientific Research Subject of Medical Treatment and Public Health of Guangxi Zhuang Autonomous Region of China (No. 200633).

8. References

- Bennett, MV., Goodenough, DA. (1978). Gap junctions, electrotonic coupling, and intercellular communication. *Neurosci Res Program Bull*, Vol.6, No.3, pp.1-486, ISSN
- Bruzzo, R., Haefliger, JA., Gimlich, RL. (1993). Connexin40, a component of gap junctions in vascular endothelium, is restricted in its ability to interact with other connexins. *Mol Biol Cell*, Vol.4, No.1, pp.7-20, ISSN 1059-1524
- Bukauskas, FF., Verselis, VK.(2004) . Gap junction channel gating. *Biochim Biophys Acta*, Vol.1662, No.1-2, pp.42-60, ISSN 1388-1981
- Burstein, B., Comtois, P., Michael, G., Nishida, K., Villeneuve, L., Yeh, YH., Nattel, S.(2009). Changes in connexin expression and the atrial fibrillation substrate in congestive heart failure. *Circ Res*, Vol.105, No.12, pp.1213-1222, ISSN 0009-7330
- Chiou, CW., Eble, JN., Zipes, DP.(1997). Efferent vagal innervation of the canine atria and sinus and atrioventricular nodes. The third fat pad. *Circulation*, Vol.95, No.11, pp.2573-2584, ISSN 0009-7322
- Chen, YJ., Chen, SA., Chen, YC., Yeh, HI., Chan, P., Chang, MS., Lin, CI.(2001). Effects of Rapid Atrial Pacing on the Arrhythmogenic Activity of Single Cardiomyocytes From Pulmonary Veins: Implication in Initiation of Atrial Fibrillation. *Circulation*, Vol.104, No.11, pp.2849-2854, ISSN 0009-7322
- Cottrell, GT., Wu, Y., Burt, JM.(2002). Cx40 and Cx43 expression ratio influences heteromeric/heterotypic gap junction channel properties. *Am J Physiol Cell Physiol*, Vol.282, No.6, pp. C1469–C1482, ISSN 0363-6143
- Chaldoupi, SM., Loh, P., Hauer, RN.(2009). The role of connexin40 in atrial fibrillation. *Cardiovasc Res*, Vol.84, No.1, pp. 15-23, ISSN 0008-6363
- Date, T., Yamane, T., Inada, K. (2007). The effects of pulmonary vein isolation on the morphology of p waves: the contribution of pulmonary vein muscle excitation to the formation of p waves. *Pacing Clin Electrophysiol*, Vol.30, No.1, pp. 93-101, ISSN 0147-8389
- Duffy, HS., Wit, AL.(2008). Is there a role for remodeled connexins in AF? No simple answers. *J Mol Cell Cardiol*, Vol.44, No.1, pp.4-13, ISSN 0022-2828
- Danik, SB., Rosner, G., Lader, J.(2008). Electrical remodeling contributes to complex tachyarrhythmias in connexin43-deficient mouse hearts. *FASEB J*, Vol.22, No.4, pp. 1204-1212, ISSN 0892-6638
- Date, T., Yamane, T., Inada, K.(2007). The effects of pulmonary vein isolation on the morphology of p waves: the contribution of pulmonary vein muscle excitation to the formation of p waves. *Pacing Clin Electrophysiol*, Vol.30, No.1, pp. 93-101, ISSN 0147-8389


- Elenes, S., Rubart, M., Moreno AP.(1999). Junctional communication between isolated pairs of canine atrial cells is mediated by homoheneous and heterogeneous gap junction channels. *J Cardiovasc Electrophysiol*, Vol.10, No.7, pp. 990–1004, ISSN 1045-3873
- Gros, DB., Jongasma, HJ. (1996). Connexins in mammalian heart function. *Bioessays*, Vol.18, No.9, pp. 719-730, ISSN 0265-9247
- Gutstein, DE., Danik, SB., Lewitton, S.(2005). Focal gap junction uncoupling and spontaneous ventricular ectopy. *Am J Physiol Heart Circ Physiol*, Vol.289, No.3, pp. H1091-H1098, ISSN 0363-6135
- Gutstein, DE., Morley, GE., Vaidya, D.(2001). Heterogeneous expression of Gap junction channels in the heart leads to conduction defects and ventricular dysfunction. *Circulation*, Vol.104, No.10, pp. 1194-1199, ISSN 0009-7322
- Hoffmann, E., Sulke, N., Edvardsson, N.(2006). on behalf of the Atrial Fibrillation Therapy (AFT) Trial Investigators Initiation of AF; New Insights Into the Initiation of Atrial Fibrillation:A Detailed Intraindividual and Interindividual Analysis of the Spontaneous Onset of Atrial Fibrillation Using New Diagnostic Pacemaker Features. *Circulation*, Vol.113, No.16, pp. 1933–1941, ISSN 0009-7322
- Haissaguerre, M., Jais, P., Shah, DC.(1998). Spontaneous Initiation of Atrial Fibrillation by Ectopic Beats Originating in the Pulmonary Veins. *N Engl J Med*, Vol.339, No.10, pp. 659–666, ISSN 0028-4793
- Honjo, H., Boyett, MR., Niwa, R.(2003). Pacing-Induced Spontaneous Activity in Myocardial Sleeves of Pulmonary Veins After Treatment With Ryanodine. *Circulation*, Vol.107, No.14, pp. 1937–1943, ISSN 0009-7322
- Herve, JC., Bourmeyster, N., Sarrouilhe, D.(2007). Gap junctional complexes:From partners to functions. *Prog Biophys Mol Biol*, Vol.94, No.1-2, pp. 29–65, ISSN 0079-6107
- Harris, AL. (2001).Emerging issues of connexin channels: biophysics fills the gap. *Q Rev Biophys*, Vol.34, No.3, pp. 325-472, ISSN 0033-5835
- Haubrich, S., Schwarz, HJ., Bukauskas, F.(1996). Incompatibility of connexin 40 and 43 Hemichannels in gap junctions between mammalian cells is determined by intracellular domains. *Mol Biol Cell*, Vol.7, No.12, pp. 1995-2006, ISSN 1059-1524
- Haissaguerre, M., Shah, DC., Jais, P.(2000). Electrophysiological breakthroughs from the left atrium to the pulmonary veins. *Circulation*, Vol.102, No.20, pp. 2463–2465, ISSN 0009-7322
- Ho, SY., Cabrera, JA., Tran, VH.(2001). Architecture of the pulmonary veins: relevance to radiofrequency ablation. *Heart*, Vol.86, No.3, pp. 265–270, ISSN 1355-6037
- Hocini, M., Ho, SY., Kawara, T.(2002). Electrical conduction in canine pulmonary veins: electrophysiological and anatomic correlation. *Circulation*, Vol.105, No.20, pp. 2442–2448, ISSN 0009-7322
- Hamabe, A., Okuyama, Y., Miyauchi, Y.(2003). Correlation between anatomy and electrical activation in canine pulmonary veins. *Circulation*, Vol.107, No.11, pp. 1550-1555, ISSN 0009-7322
- Ji, S., Gupta, N., Weiss, JN.(2010). The heart and its nerves: a nervous bond. *Heart Rhythm*, Vol.7, No.4, pp. 504-505, ISSN 1547-5271
- Jais, P., Haissaguerre, M., Shah, DC.(1997). A focal source of atrial fibrillation treated by discrete radiofrequency ablation. *Circulation*, Vol.95, No.3, pp. 572–576, ISSN 0009-7322

- Kapa, S., Venkatachalam, KL., Asirvatham, SJ.(2010). The autonomic nervous system in cardiac electrophysiology: an elegant interaction and emerging concepts. *Cardiol Rev*, Vol.18, No.6, pp. 275-284, ISSN 1061-5377
- Kanagaratnam, P., Rothery, SR., Patel, P.(2002). Relative Expression of Immunolocalized Connexins 40 and 43 Correlates With Human Atrial Conduction Properties. *J Am Coll Cardiol*, Vol.39, No.1, pp. 116-123, ISSN 0735-1097
- Kanagaratnam, P., Cherian, A., Stanbridge, RDL.(2004). Relationship between connexins and atrial activation during human atrial fibrillation. *J Cardiovasc Electrophys*, Vol.15, No.2, pp. 206-213, ISSN 1045-3873
- Kirby, ML., Waldo, KL.(1995). Neural crest and cardiovascular patterning. *Circ Res*, Vol.77, No.2, pp. 211-215, ISSN 0009-7330
- Kirchhoff, S., Kim, JS., Hagedorff, A.(2000). Abnormal cardiac conduction and morphogenesis in connexin40 and connexin43 double-deficient mice. *Circ Res*, Vol.87, No.5, pp. 399-405, ISSN 0009-7330
- Kirchhoff, S., Nelles, E., Hagedorff, A.(1998). Reduced cardiac conduction velocity and predisposition to arrhythmias in connexin40-deficient mice. *Curr Biol*, Vol.8, No.5, pp. 299-302, ISSN 0960-9822
- Kieken, F., Mutsaers, N., Dolmatova, E.(2009). Structural and molecular mechanisms of gap junction remodeling in epicardial border zone myocytes following myocardial infarction. *Circ Res*, Vol.104, No.9, pp. 1103-1112, ISSN 0009-7330
- Kontogeorgis, A., Kaba, RA., Kang, E.(2008). Short-term pacing in the mouse alters cardiac expression of connexin43. *BMC Physiol*, Vol.8, p. 8,
- Lin, X., Gemel, J., Glass, A.(2010). Connexin40 and connexin43 determine gating properties of atrial gap junction channels. *J Mol Cell Cardiol*, Vol.48, No.1, pp. 238-245, ISSN 0022-2828
- Leaf, DE., Feig, JE., Vasquez, C.(2008). Connexin40 imparts conduction heterogeneity to atrial tissue. *Circ Res*, Vol.103, No.9, pp. 1001-1008, ISSN 0009-7330
- Lu Z, Scherlag BJ, Lin J, Niu G, Ghias M, Jackman WM, Lazzara R, Jiang H, Po SS.(2008). Autonomic mechanism for complex fractionated atrial electrograms: evidence by fast fourier transform analysis. *J Cardiovasc Electrophysiol*, Vol.19, No.8, pp. 835-842, ISSN 1045-3873
- Moreno, AP.(2004). Biophysical properties of homomeric and heteromultimeric channels formed by cardiac connexins. *Cardiovasc Res*, Vol.62, No.2, pp. 276-286, ISSN 0008-6363
- Nathan, H., Eliakim, M.(1966). The junction between the left atrium and the pulmonary veins: an anatomic study of human hearts. *Circulation*, Vol.34, No.3, pp. 412- 422, ISSN 0009-7322
- Oral, H., Knight, BP., Tada, H.(2002). Pulmonary vein isolation for paroxysmal and persistent atrial fibrillation. *Circulation*, Vol.105, No.9, pp. 1077-1081, ISSN 0009-7322
- Patterson, E., Lazzara, R., Szabo, B.(2006). Sodium-Calcium Exchange Initiated by the Ca²⁺ Transient: An Arrhythmia Trigger Within Pulmonary Veins. *Journal of the American College of Cardiology*, Vol.47, No.6, pp. 1196-1206, ISSN 0735-1097

- Page, RL., Wilkinson, WE., Clair, WK.(1994). Asymptomatic arrhythmias in patients with symptomatic paroxysmal atrial fibrillation and paroxysmal supraventricular tachycardia. *Circulation*, Vol.89, No.1, pp. 224-227, ISSN 0009-7322
- Perkins, GA., Goodenough, DA., Sosinsky, GE.(1998). Formation of the gap junction intercellular channel requires a 30 degree rotation for interdigitating two apposing connexons. *J Mol Biol*, Vol.277, No.2, pp. 171-177, ISSN 0022-2836
- Pappone, C., Rosanio, S., Oreto, G.(2000). Circumferential radiofrequency ablation of pulmonary vein ostia: a new anatomic approach for curing atrial fibrillation. *Circulation*, Vol.102, No.21, pp. 2619-2628, ISSN 0009-7322
- Qu, J., Volpicelli, FM., Garcia, LI.(2009). Gap junction remodeling and spironolactone-dependent reverse remodeling in the hypertrophied heart. *Circ Res*, Vol.104, No.3, pp. 365-371, ISSN 0009-7330
- Revel, JP., Karnovsky, MJ.(1967). Hexagonal array of subunits in intercellular junctions of the mouse heart and liver. *J Cell Biol*, Vol.33, No.3, pp. C7-C12, ISSN 0021-9525
- Rackauskas, M., Neverauskas, V., Skeberdis, VA.(2010). Diversity and properties of connexin gap junction channels. *Medicina*, Vol.46, No.1, pp. 1-12, ISSN 1010-660X
- Reisner, Y., Meiry, G., Levin, ZN.(2009). Impulse conduction and gap junctional remodelling by endothelin-1 in cultured neonatal rat ventricular myocytes. *J Cell Mol Med*, Vol.13, No.3, pp. 562-573, ISSN 1582-1838
- Rackauskas, M., Kreuzberg, MM., Pranevicius, M.(2007). Gating properties of heterotypic gap junction channels formed of connexins 40, 43, and 45. *Biophys J*, Vol.92, No.6, pp. 1952-1965, ISSN 0006-3495
- Roux, N., Havet, E., Mertl, P. The myocardial sleeves of the pulmonary veins: potential implications for atrial fibrillation. *Surg Radiol Anat*, Vol.26, No.4, pp. 285-289, ISSN 0930-1038
- Sellers, MB., Newby, LK.(2011). Atrial fibrillation, anticoagulation, fall risk, and outcomes in elderly patients. *Am Heart J*, Vol.161, No.2, pp. 241-246, ISSN 0002-8703
- Sandres, P., Jais, P., Hocini, M. Electrical disconnection of the coronary sinus by radiofrequency catheter ablation to isolate a trigger of atrial fibrillation. *J Cardiovasc Electr*, Vol.15, No.3, pp. 364-368, ISSN 1045-3873
- Stergiopoulos, K., Alvarado, JL., Mastroianni, M.(1999). Heterodomain interactions as a mechanism for the regulation of connexin channels. *Circ Res*, Vol.84, No.10, pp. 1144-1155, ISSN 0009-7330
- Söhl, G., Willecke, K. Gap junctions and the connexin protein family. *Cardiovasc Res*, Vol.62, No.2, pp. 228-232, ISSN 0008-6363
- Strom, M., Wan, X., Poelzing, S.(2010). Gap junction heterogeneity as mechanism for electrophysiologically distinct properties across the ventricular wall. *Am J Physiol Heart Circ Physiol*, Vol.298, No.3, pp. H787-H794, ISSN 0363-6135
- Severs, NJ., Bruce, AF., Dupont, E.(2008). Remodelling of gap junctions and connexin expression in diseased myocardium. *Cardiovasc Res*, Vol.80, No.1, pp. 9-19, ISSN 0008-6363
- Saito, T., Waki, K., Becker, AE.(2000). Left atrial myocardial extension onto pulmonary veins in humans: anatomic observations relevant for atrial arrhythmias. *J Cardiovasc Electrophysiol*, Vol.11, No.8, pp. 888-894, ISSN 1045-3873

- Steiner, I., Hájková, P., Kvasnicka, J.(2005). Pulmonary veins and atrial fibrillation: a pathological study of 100 hearts. *Cesk Patol*, Vol.41, No.4, pp. 124-131
- Tsuboi, M., Furukawa, Y., Nakajima, K.(2000). Inotropic, chronotropic, and dromotropic effects mediated via parasympathetic ganglia in the dog heart. *Am J Physiol Heart Circ Physiol*, Vol.279, No.3, pp. H1201-1207, ISSN 0363-6135
- Tan, AY., Li, H., Hogiu, WS.(2006). Autonomic innervation and segmental muscular disconnections at the human pulmonary vein-atrial junction: implications for catheter ablation of atrial-pulmonary vein junction. *J Am Coll Cardiol*, Vol.48, No.1, pp. 132-143, ISSN 0735-1097
- Unger, VM., Kumar, NM., Gilula, NB.(1999). Three-dimensional structure of a recombinant gap junction membrane channel. *Science*, Vol.283, No.5405, pp. 1176-1180, ISSN 0036-8075
- Volders, PG.(2010). Novel insights into the role of the sympathetic nervous system in cardiac arrhythmogenesis. *Heart Rhythm*, Vol.7, No.12, pp. 1900-1906, ISSN 1547-5271
- Verheule, S., Wilson, EE., Arora, R.(2002). Tissue structure and connexin expression of canine pulmonary veins. *Cardiovasc Res*, Vol.55, No.4, pp. 727-738, ISSN 0008-6363
- Valiunas, V., Gemel, J., Brink, PR.(2001). Gap junction channels formed by coexpressed connexin40 and connexin43. *Am J Physiol Heart Circ Physiol*, Vol.281, No.4, pp. H1675-H1689, ISSN 0363-6135
- Valiunas, V., Weingart, R., Brink, PR.(2000). Formation of heterotypic gap junction channels by connexins 40 and 43. *Circ Res*, Vol.86, No.2, pp. E42-E49, ISSN 0009-7330
- Verheijck, EE., Kempen, MJ., Veereschild, M.(2001). Electrophysiological features of the mouse sinoatrial node in relation to connexin distribution. *Cardiovasc Res*, Vol.52, No.1, pp. 40-50, ISSN 0008-6363
- Valiunas, V., Weingart, R., Brink, PR.(2000). Formation of heterotypic gap junction channels by connexins 40 and 43. *Circ Res*, Vol.86, No.2, pp. E42-E49, ISSN 0009-7330
- Valiunas, V., Beyer, EC., Brink, PR.(2002). Cardiac gap junction channels show quantitative differences in selectivity. *Circ Res*, Vol.91, No.2, pp. 104-111, ISSN 0009-7330
- Valiunas, V., Polosina, YY., Miller, H.(2005). Connexin-specific cell-to-cell transfer of short interfering RNA by gap junctions. *J Physiol*, Vol.568, No.Pt2, pp. 459-468, ISSN 0022-3751
- Wit, AL., Boyden, PA.(2007). Triggered activity and atrial fibrillation. *Heart Rhythm*, Vol.4, No.3, pp. S17-S23, ISSN 1547-5271
- Wijffels MC, Kirchhof CJ, Dorland R, Allessie MA.(1995). Atrial fibrillation begets atrial fibrillation. A study in awake chronically instrumented goats. *Circulation*, Vol.92, No.7, pp. 1954-1968, ISSN 0009-7322
- Xiong RX, Zhong GQ, Song HX, Zhang JC, Ling Y.(2010). Collagen spatial distribution in rapid atrial pacing dogs with or without superior vena cava and aortic root fat. *Zhonghua Xin Xue Guan Bing Za Zhi*, Vol.38, No.2, pp. 171-174, ISSN 0253-3758
- Yeager, M., Gilula, NB.(1992). Membrane topology and quaternary structure of cardiac gap junction ion channels. *J Mol Biol*, Vol.223, No.4, pp. 929-948, ISSN 0022-2836
- Yamada, KA.(2008). A change of heart: heterogeneous remodeling in heart failure. *Heart Rhythm*, Vol.5, No.8, pp. 1186-1188, ISSN 1547-5271

- Yeh HI, Lai YJ, Lee SH, Chen ST, Ko YS, Chen SA, Severs NJ, Tsai CH.(2006). Remodeling of myocardial sleeve and gap junctions in canine superior vena cava after rapid pacing. *Basic Res Cardiol*, Vol.101, No.4, pp. 269-280, ISSN 0300-8428
- Zhou SH, He XZ, Liu QM, Du WH, Li XP, Zhou T, Tang JQ, Li RS.(2008). Study on the spatial distribution pattern of Cx40 gap junctions in the atria of patients with coronary heart disease. *Cardiol J*, Vol.15, No.1, pp. 50-56, ISSN 1897-5593



Advances in Electrocardiograms - Clinical Applications

Edited by PhD. Richard Millis

ISBN 978-953-307-902-8

Hard cover, 328 pages

Publisher InTech

Published online 25, January, 2012

Published in print edition January, 2012

Electrocardiograms have become one of the most important, and widely used medical tools for diagnosing diseases such as cardiac arrhythmias, conduction disorders, electrolyte imbalances, hypertension, coronary artery disease and myocardial infarction. This book reviews recent advancements in electrocardiography. The four sections of this volume, Cardiac Arrhythmias, Myocardial Infarction, Autonomic Dysregulation and Cardiotoxicology, provide comprehensive reviews of advancements in the clinical applications of electrocardiograms. This book is replete with diagrams, recordings, flow diagrams and algorithms which demonstrate the possible future direction for applying electrocardiography to evaluating the development and progression of cardiac diseases. The chapters in this book describe a number of unique features of electrocardiograms in adult and pediatric patient populations with predilections for cardiac arrhythmias and other electrical abnormalities associated with hypertension, coronary artery disease, myocardial infarction, sleep apnea syndromes, pericarditides, cardiomyopathies and cardiotoxicities, as well as innovative interpretations of electrocardiograms during exercise testing and electrical pacing.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Guo-qiang Zhong, Ri-xin Xiong, Hong-xing Song, Yun Ling, Jing-chang Zhang and Zhe Wei (2012). The Remodeling of Connexins Localized at Pulmonary Vein – Left Atria in Triggering and Maintenance of Atrial Fibrillation, *Advances in Electrocardiograms - Clinical Applications*, PhD. Richard Millis (Ed.), ISBN: 978-953-307-902-8, InTech, Available from: <http://www.intechopen.com/books/advances-in-electrocardiograms-clinical-applications/the-remodeling-of-connexins-localized-at-pulmonary-vein-left-atria-in-triggering-and-maintenance-of->

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.