



The Role of Fractional Time-Derivative Operators on Anomalous Diffusion

Angel A. Tateishi¹, Haroldo V. Ribeiro^{2*} and Ervin K. Lenzi³

¹ Departamento de Física, Universidade Tecnológica Federal do Paraná, Pato Branco, Brazil, ² Departamento de Física, Universidade Estadual de Maringá, Maringá, Brazil, ³ Departamento de Física, Universidade Estadual de Ponta Grossa, Ponta Grossa, Brazil

OPEN ACCESS

Edited by:

Matjaž Perc,
University of Maribor, Slovenia

Reviewed by:

Jordan Yankov Hristov,
University of Chemical Technology
and Metallurgy, Bulgaria
Angel R. Plastino,
National University of Northwestern
Buenos Aires, Argentina

*Correspondence:

Haroldo V. Ribeiro
hvr@dfi.uem.br

Specialty section:

This article was submitted to
Interdisciplinary Physics,
a section of the journal
Frontiers in Physics

Received: 11 September 2017

Accepted: 05 October 2017

Published: 25 October 2017

Citation:

Tateishi AA, Ribeiro HV and Lenzi EK
(2017) The Role of Fractional
Time-Derivative Operators on
Anomalous Diffusion.
Front. Phys. 5:52.
doi: 10.3389/fphy.2017.00052

The generalized diffusion equations with fractional order derivatives have shown to be quite efficient to describe the diffusion in complex systems, with the advantage of producing exact expressions for the underlying diffusive properties. Recently, researchers have proposed different fractional-time operators (namely: the Caputo-Fabrizio and Atangana-Baleanu) which, differently from the well-known Riemann-Liouville operator, are defined by non-singular memory kernels. Here we proposed to use these new operators to generalize the usual diffusion equation. By analyzing the corresponding fractional diffusion equations within the continuous time random walk framework, we obtained waiting time distributions characterized by exponential, stretched exponential, and power-law functions, as well as a crossover between two behaviors. For the mean square displacement, we found crossovers between usual and confined diffusion, and between usual and sub-diffusion. We obtained the exact expressions for the probability distributions, where non-Gaussian and stationary distributions emerged. This former feature is remarkable because the fractional diffusion equation is solved without external forces and subjected to the free diffusion boundary conditions. We have further shown that these new fractional diffusion equations are related to diffusive processes with stochastic resetting, and to fractional diffusion equations with derivatives of distributed order. Thus, our results suggest that these new operators may be a simple and efficient way for incorporating different structural aspects into the system, opening new possibilities for modeling and investigating anomalous diffusive processes.

Keywords: fractional-time operators, anomalous diffusion, continuous time random walk, fractional order derivatives, fractional calculus

1. INTRODUCTION

The *random walk* concept is one of the broadest and versatile paradigms to deal with statistical fluctuations. The term “random walk” was coined in 1905 by Pearson [1], but the fundamental relationship between this concept and the usual diffusion equation was reported earlier in the seminal works of Rayleigh [2–4] in sound theory, Bachelier [5] in economics, Einstein [6], and von Smoluchowski [7] in the Brownian motion theory. Due to this intrinsic relation, the usual random walk is characterized by Gaussian, Markovian, and ergodic properties, which lead to a linear time dependence of the mean square displacement, $(\Delta x)^2 \sim t$. The versatility of this

concept relies on the possibility of generalizations and extensions to describe systems with one or more characteristics of anomalous diffusion: non-Gaussian distributions; long-range memory effects (non-Markovian); non-ergodicity; divergent mean square displacement (Lévy walks); and nonlinear mean square displacement, $(\Delta x)^2 \sim t^\alpha$ (sub-diffusion: $\alpha < 1$, superdiffusion: $\alpha > 1$, confined or saturated diffusion: $\alpha = 0$).

In the context of generalizations, the first landmark is the work proposed in 1965 by Montroll and Weiss [8], in which they introduce the *continuous time random walk* concept (see [9] for a general overview). This framework is characterized by a joint distribution of jump length and waiting time $\psi(x, t)$, where $\lambda(x) = \int_0^\infty dt \psi(x, t)$ is the jump length distribution and $\omega(t) = \int_{-\infty}^\infty dx \psi(x, t)$ is the waiting time distribution. Subsequently, a connection with a generalized master equation is proposed to discuss memory effects in the continuous time random walk [10–12]; the waiting time distribution $\omega(t)$ is strictly related to the memory kernel of the generalized master equation. Moreover, natural extensions of both random walk and continuous time random walk were proposed to study transport properties in systems with structural complexity, such as disordered, random, and fractal environments [13].

The second landmark, “a modern era of the continuous time random walk” according to Kutner and Masoliver [9], is the development of the intrinsic relationship between this formalism and the fractional diffusion equations. Among the seminal works, we have found a simple mention in the work of Klafter et al. [14] on the possibility of having a fractional diffusion equation to describe anomalous transport. However, were Hilfer et al. [15] that, 90 years after Einstein’s work [6], established a rigorous and precise connection between the continuous time random walk and the fractional master equation as well as with the fractional diffusion equation (a special case of the former). This result was later extended by Compte [16] in the long-time limit, where it is shown that any decoupled continuous time random walk having no characteristic scale of time or space (power-law memories) corresponds to a time- or space-fractional diffusion equation, respectively with Riemann-Liouville time derivative or Riesz space derivative. The subsequent success and development of the fractional approach are well documented in two review articles by Metzler and Klafter [17, 18], and in several articles by Barkai [19–22].

Since memory effects underlying a continuous time random walk are implicitly considered by the differential operators, the versatility of the fractional formalism is mainly related to two remarkable features. First, this formalism handles very well the physical requirements of a system by dealing boundary conditions and external forces in a simple manner. Second, it takes advantage of traditional tools from mathematical physics and statistics for obtaining exact expressions to describe complex systems with anomalous behaviors. For instance, in the following fractional differential equation

$$\frac{\partial}{\partial t} \rho(x, t) = {}_0\mathcal{D}_t^{1-\alpha} (\mathcal{L}\{\rho(x, t)\}), \quad (1)$$

where

$$\mathcal{L}\{\rho(x, t)\} = D \frac{\partial^2}{\partial x^2} \rho(x, t) - \frac{\partial}{\partial x} [F(x, t) \rho(x, t)],$$

the unusual relaxation can be associated with a continuous time random walk where the waiting time distribution is a power-law. This process is also strictly related to the Riemann-Liouville fractional operator [23]

$${}_0\mathcal{D}_t^{1-\alpha} \rho(x, t) = \frac{1}{\Gamma(\alpha)} \frac{d}{dt} \int_0^t \frac{\rho(x, t')}{(t-t')^{1-\alpha}} dt', \quad (2)$$

where $0 < \alpha < 1$ is the *fractional order exponent* (or the *anomalous exponent*), a quantity that can be interpreted as an index of memory in empirical systems [24]. The fractional operator is also responsible for introducing a nonlinear time dependence in the mean square displacement of the system [17]. Thus, a large class of complex phenomena can be effectively described by extending the standard differential operator to a non-integer order [25–34]; indeed, as pointed out by West [35], the fractional calculus provides a suitable framework to deal with complex systems.

Recently, researchers have made and promoted remarkable progress toward improving experimental techniques for investigating diffusive processes, mainly illustrated by the developments in the single-particle tracking technique [36–39]. Such improvements yield novel insights into transport properties of biological systems [40–42] and nanomaterials [43–45], where the high-resolution of the experiments has found different diffusive behaviors depending on the time scale. In this context, an important question is whether other forms of fractional differential operators (replacing the Riemann-Liouville one) such as those recently-proposed with non-singular kernels [46–51] are suitable to describe the aforementioned situations. To answer this question, we investigate an one-dimensional diffusive process described by the fractional diffusion equation

$$\frac{\partial}{\partial t} \rho(x, t) = D \mathcal{F}_t^\alpha \left(\frac{\partial^2}{\partial x^2} \rho(x, t) \right), \quad (3)$$

where D is the generalized diffusion coefficient. This equation is also subjected to the free diffusion boundary conditions $\rho(\pm\infty, t) = 0$ and to the initial condition $\rho(x, 0) = \varphi(x)$.

The fractional operator in Equation (3) is defined as

$$\mathcal{F}_t^\alpha \rho(x, t) = \frac{\partial}{\partial t} \int_0^t \rho(x, t') \mathcal{K}(t-t') dt', \quad (4)$$

in order to consider situations with singular and non-singular kernels in a unified way. It is worth noting that $\mathcal{K}(t) = \delta(t)$ recovers the usual diffusion equation. Here we consider three different forms for the kernel $\mathcal{K}(t)$. The first one is

$$\mathcal{K}(t) = t^{\alpha-1} / \Gamma(\alpha), \quad (5)$$

which corresponds to the well-known Riemann-Liouville fractional operator [52] for $0 < \alpha < 1$. The second one is

$$\mathcal{K}(t) = b \exp\left(-\frac{\alpha t}{1-\alpha}\right), \quad (6)$$

which corresponds to the fractional operator of Caputo-Fabrizio [46, 53, 54]. As discussed in Caputo and Fabrizio [46], the ratio $\alpha/(1-\alpha)$ maps the time from the range $[0, \infty]$ to the range $[0, 1]$ (see also [54] for details on the role of the fractional order α).

Finally, the third one is

$$\mathcal{K}(t) = b E_{\alpha} \left(-\frac{\alpha t^{\alpha}}{1-\alpha} \right), \quad (7)$$

where $E_{\alpha}(\dots)$ is the Mittag-Leffler function [52]. This kernel corresponds to the fractional operator of Atangana and Baleanu [47]. Further possibilities for the kernel $\mathcal{K}(t)$ are discussed by Gómez-Aguilar et al. [48]. We observe that the Riemann-Liouville operator have a singularity at the origin ($t = 0$), while the recently-proposed Caputo-Fabrizio and Atangana-Baleanu are non-singular operators [46–51]. In the previous definitions, the parameter b is a normalization constant and α is the fractional order exponent.

Our main goal here is to verify how these different fractional operators modify the fractional diffusion Equation (1) and what are the effect of these choices on the underlying diffusive properties of a system modeled by this equation. The rest of this manuscript is organized as follows. In Section II, we investigate general solutions and processes related with Equation (3) when considering different choices (singular and non-singular) for the kernel $\mathcal{K}(t)$. In Section 3, we present a summary of our results and some concluding remarks.

2. DIFFUSION AND FRACTIONAL OPERATORS

We start by noting that the solution of the fractional diffusion Equation (3) in the Fourier-Laplace space is

$$\rho(k, s) = \frac{\varphi(k)}{s + s\mathcal{D}\mathcal{K}(s)k^2}, \quad (8)$$

where $\rho(k, s)$ is the Fourier-Laplace transformation of the probability distribution $\rho(x, t)$. This result can be related to different situations depending on the choice of the kernel $\mathcal{K}(s)$.

Within the continuous time random walk formalism and by following the works of Metzler and Klafter [17], we can show that the waiting time $\omega(t)$ and the jump $\lambda(x)$ probability distribution associated with Equation (3) are (in the Laplace and Fourier spaces)

$$\omega(s) = \frac{\mathcal{K}(s)/\tau_c}{1 + \mathcal{K}(s)/\tau_c} \quad (9)$$

and $\lambda(k) = 1 - k^2 D \tau_c$, where τ_c is a characteristic waiting time of the underlying continuous time random walk. We observe that the jump probability distribution is characterized by a Gaussian asymptotic behavior [$\lambda(x) \sim e^{-x^2/(2D\tau_c)^2}$] and thus has a finite characteristic jump length, regardless of the choice for the kernel

$\mathcal{K}(t)$. On the other hand, the inverse Laplace transform of the waiting time distribution is given by

$$\begin{aligned} \omega(t) &= \frac{1}{\tau_c} \int_0^t dt' \mathcal{K}(t') + \sum_{n=1}^{\infty} \left(-\frac{1}{\tau_c} \right)^n \int_0^t dt_n \mathcal{K}(t - t_n) \cdots \\ &\times \int_0^{t_4} dt_3 \mathcal{K}(t_4 - t_3) \int_0^{t_2} dt_1 \mathcal{K}(t_2 - t_1) \mathcal{K}(t_1), \end{aligned} \quad (10)$$

yielding different situations that depends on $\mathcal{K}(t)$.

The choice $\mathcal{K}(t) = \delta(t)$ leads to usual diffusion and an exponential distribution for the waiting times

$$\omega(t) = \frac{1}{\tau_c} e^{-t/\tau_c}. \quad (11)$$

For the fractional operator of Riemann-Liouville, we find

$$\omega(t) = \frac{1}{\tau_c} t^{\alpha-1} E_{\alpha, \alpha} \left(-\frac{1}{\tau_c} t^{\alpha} \right), \quad (12)$$

where $E_{\alpha, \alpha}(\dots)$ is the generalized Mittag-Leffler function [52] whose asymptotic behavior is described by a power-law, $\omega(t) \sim 1/t^{1+\alpha}$ for $t \rightarrow \infty$. For the Atangana-Baleanu operator (Equation 7), the waiting time distribution is given by

$$\omega(t) = \frac{\xi b \gamma}{\pi \tau_c} \sin(\pi \gamma) \int_0^{\infty} d\eta \frac{\eta^{\gamma} e^{-\eta b(1-\alpha)t/\tau_c}}{(1-\eta)^2 + 2\xi(1-\eta)\eta^{\gamma} \cos(\pi \gamma) + \eta^{2\gamma}} \quad (13)$$

where $\gamma = 1 - \alpha$ and $\xi = \alpha b/\tau_c$. This expression is very interesting because for small times we have a stretched exponential, that is,

$$\omega(t) \sim E_{\alpha} \left(-\frac{\alpha t^{\alpha}}{1-\alpha} \right) \sim \exp \left(-\frac{\alpha t^{\alpha}}{(1-\alpha)\Gamma(\alpha)} \right), \quad (14)$$

while for long times we have the same power-law behavior of the Riemann-Liouville operator. Thus, the Atangana-Baleanu operator yields a crossover between a stretched exponential and a power-law distribution.

In the case of the Caputo-Fabrizio operator, the connection with the continuous time random walk is more complex and not compatible with its standard interpretation. As we shall discuss later on, the diffusion equation associated with this operator is connected to a diffusive process with stochastic resetting [55, 56], where the waiting time distribution is exponential.

Figure 1 depicts the behavior of the waiting time distribution $\omega(t)$ for the different kernels previously-discussed. For long times, we confirm that the operators of Riemann-Liouville and Atangana-Baleanu yield the same power-law decay for $\omega(t)$. We further note that the Atangana-Baleanu operator yields a non-divergent $\omega(t)$, an interesting feature that is not observed for the singular kernel of Riemann-Liouville.

We now focus on finding the formal solutions for the fractional diffusion Equation (1) when considering the three different fractional operators. These solutions are obtained by performing the inverse of Fourier and Laplace transforms of the




FIGURE 1 | Changes in the waiting time distribution $\omega(t)$ caused by the different forms of the kernel $\mathcal{K}(t)$ defining the fractional operator of Equation (4). The different curves correspond the $\omega(t)$ when choosing the kernels of Riemann-Liouville (Equation 5, blue), Atangana-Baleanu (Equation 7, red), and the usual (Brownian motion) case [$\mathcal{K}(t) = \delta(t)$, green]. For simplicity, we have considered $\tau_c = 1$ and $\alpha = 1/2$. We note that the asymptotic behavior of $\omega(t)$ is a power-law for the kernels of Riemann-Liouville and Atangana-Baleanu, that is, $\omega(t) \sim 1/t^{1+\alpha}$. In the usual, we have an exponential behavior.

$\rho(k, s)$ expressed in Equation (8), where the Laplace transform of the kernel $\mathcal{K}(s)$ appears. In the well-known case of the Riemann-Liouville operator [17], we have

$$\mathcal{K}(s) = s^{-\alpha} \tag{15}$$

and consequently

$$\rho(x, t) = \int_{-\infty}^{\infty} dx' \mathcal{G}(x - x', t) \varphi(x'), \tag{16}$$

where the Green function is

$$\mathcal{G}(x, t) = \frac{1}{2|x|} \mathbf{H}_{1,1}^{1,0} \left[\frac{|x|}{\sqrt{D} t^\alpha} \middle| \begin{matrix} (1, \frac{\alpha}{2}) \\ (1, 1) \end{matrix} \right]. \tag{17}$$

Here $\mathbf{H}(\dots)$ stands for the Fox H-function [57]. Having found the probability distribution, we can show that the mean square displacement is

$$\langle (\Delta x)^2 \rangle = \langle (x - \langle x \rangle)^2 \rangle = \frac{2D t^\alpha}{\Gamma(1 + \alpha)}, \tag{18}$$

which corresponds to the typical case of anomalous diffusion, where $\alpha < 1$ represents sub-diffusion and $\alpha \rightarrow 1$ recovers the usual diffusion. The time-dependent behavior of a typical probability distribution $\rho(x, t)$ (with $\alpha = 1/2$) is shown in **Figure 2A**. We observe that the Riemann-Liouville operator leads to a tent-shaped distribution, whose tails are longer than the Gaussian distribution of the usual diffusion (**Figure 2D**). **Figure 3** shows the corresponding behavior for mean square

displacement of Equation (18), which is a power-law function of the time t with an exponent α .

For the Caputo-Fabrizio operator, the Laplace transform of the kernel in Equation (6) is

$$\mathcal{K}(s) = \frac{b}{\left(s + \frac{\alpha}{1-\alpha}\right)}, \tag{19}$$

which substituted into Equation (8) yields

$$\rho(k, s) = \frac{\left(s + \frac{\alpha}{1-\alpha}\right) \varphi(k)}{s \left(s + \frac{\alpha}{1-\alpha} + D b k^2\right)}. \tag{20}$$

By performing the inverse Fourier and Laplace transforms, we have

$$\begin{aligned} \rho(x, t) = & \int_{-\infty}^{\infty} dx' \mathcal{G}(x - x', t) \varphi(x') \\ & + \frac{\alpha}{1-\alpha} \int_0^t dt' \int_{-\infty}^{\infty} dx' \mathcal{G}(x - x', t') \varphi(x') \end{aligned} \tag{21}$$

where the Green function is

$$\mathcal{G}(x, t) = \frac{e^{-\frac{\alpha t}{1-\alpha}}}{\sqrt{4\pi D b t}} e^{-\frac{x^2}{4D b t}}, \tag{22}$$

A typical shape of this distribution is shown in **Figure 2B**. We observe that this distribution is very similar to a Gaussian for small times, and exhibits a tent-shape to long times. However, differently from the distribution obtained for the Riemann-Liouville operator (Equations 16 and 17), the distribution obtained from Equations (21) and (22) displays a stationary behavior for $t \rightarrow \infty$, that is,


$$\rho(x, t \rightarrow \infty) \sim \int_{-\infty}^{\infty} dx' \varphi(x') e^{-\frac{\alpha|x-x'|}{(1-\alpha)D b}}, \tag{23}$$

a result that corresponds to confined diffusion. **Figure 2B** also shows this stationary solution (dashed line); in particular, we observe that the shape of $\rho(x, t)$ is practically constant for $t \gtrsim 5$ in that case. This behavior also appears in the mean square displacement

$$\langle (\Delta x)^2 \rangle = \frac{2D b (1 - \alpha)}{\alpha} \left(1 - e^{-\frac{\alpha t}{1-\alpha}}\right), \tag{24}$$

which behaves linearly in time for small times and saturates in $2D b (1 - \alpha)/\alpha$ for long times. **Figure 3** illustrates this crossover, a common feature of systems where diffusion is confined or hindered [41, 58]. In particular, the same crossover between usual and confined diffusion is observed in simulations of diffusion with immobile obstacles or obstacles moving according to an Ornstein-Uhlenbeck process [59, 60].

An intriguing feature of the diffusion equation with the Caputo-Fabrizio fractional operator is that it can be related to a diffusion with stochastic resetting [61]. Indeed, we find out



that the fractional diffusion Equation (3) with the kernel of Equation (6) can be rewritten as

$$\frac{\partial}{\partial t} \rho(x, t) = Db \frac{\partial^2}{\partial x^2} \rho(x, t) - \frac{\alpha}{1-\alpha} Db \int_0^t dt' e^{-\frac{\alpha}{1-\alpha}(t-t')} \frac{\partial^2}{\partial x^2} \rho(x, t'). \quad (25)$$

This equation is essentially the same obtained by Hristov [49] when by analyzing a heat diffusion equation with non-singular memory. Also, by integrating both sides of the fractional diffusion Equation (3), we obtain

$$Db \int_0^t dt' e^{-\frac{\alpha}{1-\alpha}(t-t')} \frac{\partial^2}{\partial x^2} \rho(x, t') = \rho(x, t) - \varphi(x), \quad (26)$$

which after substituting into Equation (25) yields

$$\frac{\partial}{\partial t} \rho(x, t) = Db \frac{\partial^2}{\partial x^2} \rho(x, t) - \frac{\alpha}{1-\alpha} [\rho(x, t) - \varphi(x)]. \quad (27)$$

Equation (27) with $\varphi(x) = \delta(x - x_0)$ is the same obtained by Evans and Majumdar [61] when studying a random walker whose position is redefined to the position x_0 with a rate $r = \alpha/(1 - \alpha)$. Thus, the fractional exponent α in the fractional diffusion

equation of Caputo-Fabrizio can be related to a well-defined physical quantity (resetting rate).

Also, the mean square displacement of Equation (24) is analogous to results obtained from a random walk description of a diffusive process with stochastic resetting, subjected to an exponential waiting time distribution [55, 56]. As discussed in these works, a suitable continuous time random walk formulation is established by considering a density of particles $\mathcal{J}(x, t)$ whose dynamics is governed by

$$\begin{aligned} \mathcal{J}(x, t) = & \delta(t)\delta(x) + r\delta(x) \int_0^t dt' \omega(t') \mathcal{J}(x, t-t') \\ & + (1-r) \int_0^t dt' \psi(x', t') \mathcal{J}(x-x', t-t'), \quad (28) \end{aligned}$$

when particles start the random walk at the origin ($x = 0$) with $\rho(x, t) = \int_0^t \Phi(t') \mathcal{J}(x, t-t') dt'$ and $\Phi(t) = \int_t^\infty \omega(t') dt'$. In Equation (28), r is a resetting rate, $\psi(x, t)$ is joint distribution of jump length and waiting time, $\lambda(x) = \int_0^\infty dt \psi(x, t)$ is the jump length distribution, and $\omega(t) = \int_{-\infty}^\infty dx \psi(x, t)$ is waiting time distribution. By considering $\lambda(x)$ Gaussian and $\omega(t)$ exponentially distributed, we can show that this formalism leads to Equation (27). It is worth remarking that by comparison with this framework, we can infer that the diffusion equation with




FIGURE 3 | Changes in the evolving behavior of the mean square displacement $(\Delta x)^2$ caused by the different fractional time operators. The curves show $(\Delta x)^2$ vs. t when considering the operators of Riemann-Liouville (blue, Equation 18), Caputo-Fabrizio (purple, Equation 24), Atangana-Baleanu (red, Equation 35), and the usual case [green, $(\Delta x)^2 \propto t$]. For simplicity, we have considered $\alpha = 1/2$ and $D b = 1$. It is worth noting the Atangana-Baleanu operator shows a crossover between usual (for small times) and sub-diffusion (for long time). In the Caputo-Fabrizio case, the diffusion is usual for small times and saturates for large times.

the Caputo-Fabrizio operator leads to the same waiting time distribution of the usual diffusion, that is, an exponential.

Finally, for the Atangana-Baleanu operator, the Laplace transform of the kernel in Equation (7) is

$$\mathcal{K}(s) = \frac{b s^{\alpha-1}}{\left(s^\alpha + \frac{\alpha}{1-\alpha}\right)}, \quad (29)$$

which substituted into Equation (8) yields

$$\rho(k, s) = \frac{\left(s^\alpha + \frac{\alpha}{1-\alpha}\right) \varphi(k)}{s^\alpha \left(s + \frac{\alpha}{1-\alpha} s^{1-\alpha} + D b k^2\right)}, \quad (30)$$

the solution for the fractional diffusion Equation (3) in the Fourier-Laplace space. By evaluating the inverse Fourier and Laplace transforms, we obtain

$$\begin{aligned} \rho(x, t) &= \int_{-\infty}^{\infty} dx' \mathcal{G}(x-x', t) \varphi(x') \\ &+ \frac{\alpha}{\Gamma(\alpha)(1-\alpha)} \int_0^t \frac{dt'}{(t-t')^{1-\alpha}} \int_{-\infty}^{\infty} dx' \varphi(x') \mathcal{G}(x-x', t'), \end{aligned} \quad (31)$$

where the Green function is

$$\begin{aligned} \mathcal{G}(x, t) &= \frac{e^{-\frac{x^2}{4D b t}}}{\sqrt{4\pi D b t}} \\ &+ \frac{1}{|x|} \sum_{n=1}^{\infty} \frac{\left(-\frac{\alpha}{1-\alpha}\right)^n}{\Gamma(1+n)} t^{n\alpha} \mathbf{H}_{2,2}^{2,0} \left[\frac{x^2}{D b t} \begin{matrix} (1+\alpha n, 1), (1, 1) \\ (1, 2), (1+n, 1) \end{matrix} \right]. \end{aligned} \quad (32)$$

Once again, $\mathbf{H}(\dots)$ stands for the Fox H-function [57]. We can also show that for $|x| \rightarrow \infty$, Equation (32) is approximated by

$$\mathcal{G}(x, t) \approx \frac{1}{\sqrt{4\pi D b t}} e^{-f(x,t)}, \quad (33)$$

where

$$f(x, t) = \frac{x^2}{4D b t} + \frac{\alpha}{1-\alpha} t^\alpha \left(\frac{x^2}{4D b t}\right)^{1-\alpha}. \quad (34)$$

A typical behavior for the distribution $\rho(x, t)$ for this operator is shown in **Figure 2C**. Similarly to the Caputo-Fabrizio operator, the profile of $\rho(x, t)$ resembles a Gaussian for small times, while exhibits a tent-shape for long times. However, the distribution does not have a stationary solution for the Atangana-Baleanu operator. This crossover between two behaviors for $\rho(x, t)$ is also present in Equation (34), and can be better quantified by analyzing the mean square displacement. For this operator, we have

$$(\Delta x)^2 = 2D b t E_{\alpha,2} \left(-\frac{\alpha}{1-\alpha} t^\alpha\right), \quad (35)$$

where $E_{\alpha,2}(\dots)$ is the generalized Mittag-Leffler function [52]. By considering the asymptotic limits of this function, we can show that $(\Delta x)^2 \sim t$ for small times, and $(\Delta x)^2 \sim t^{1-\alpha}$ for long times.

This crossover between usual and sub-diffusion is present in several biological systems [62–66] and is also illustrated in **Figure 3** for $\alpha = 1/2$. A similar situation appears in simulations of diffusion with obstacles moving according to a usual random walk [59, 60], where the same crossover between usual and sub-diffusion with $\alpha = 1/2$ is observed. It is worth mentioning that crossovers between diffusive regimes can also be described by generalized Langevin equations [67] and fractional (with the Riemann-Liouville operator) Kramers equations [19], among other approaches [68, 69]. In particular, the usual Langevin equation [70] predicts a crossover between ballistic and usual diffusion, which has been experimentally observed only in 2011 [71]. However, the diffusion equation in terms of these new operators lead to these crossovers without explicitly considering external forces, inertial effects, and reaction terms.

The fractional diffusion equation with the Atangana-Baleanu operator can be further related to fractional derivatives of distributed order as proposed by Caputo [72, 73] and worked out in Chechkin [68] and Lenzi et al. [69], that is,

$$\int_0^1 dv w(v) \frac{\partial^v}{\partial t^v} \rho(x, t) = D \frac{\partial^2}{\partial x^2} \rho(x, t), \quad (36)$$

TABLE 1 | Summary of the changes caused by the different fractional operators on the diffusion Equation (1).

Fractional operator	Waiting time distribution	Mean square displacement	Probability distribution
Riemann- Liouville	Power-law	Power-law and scale-invariant	Non-Gaussian
Caputo- Fabrizio	Exponential	Crossover from usual to confined diffusion	Crossover from Gaussian to non-Gaussian with steady state
Atangana- Baleanu	Crossover from stretched exp. to power-law	Crossover from usual to sub-diffusion	Crossover from Gaussian to non-Gaussian

where $w(\nu)$ is the distribution of the fractional order exponent ν and

$$\frac{\partial^\nu}{\partial t^\nu} \rho(x, t) = \frac{1}{\Gamma(1-\nu)} \int_0^t \frac{dt'}{(t-t')^\nu} \frac{\partial}{\partial t'} \rho(x, t') \quad (37)$$

is the fractional time derivative of Caputo. Indeed, by substituting the kernel of Equation (7) into Equation (3) and taking the Laplace transform, we have

$$s\rho(x, s) - \rho(x, 0) = Db \frac{s^{1-\alpha}}{s^{1-\alpha} + \frac{\alpha}{1-\alpha}} \frac{\partial^2}{\partial x^2} \rho(x, s), \quad (38)$$

which can be rewritten as

$$\begin{aligned} s\rho(x, s) - \rho(x, 0) + \frac{\alpha}{1-\alpha} s^{-\alpha} [s\rho(x, s) - \rho(x, 0)] \\ = Db \frac{\partial^2}{\partial x^2} \rho(x, s). \end{aligned} \quad (39)$$

By calculating the inverse Laplace transform of the previous equation, we find

$$\begin{aligned} \frac{\partial}{\partial t} \rho(x, t) + \frac{\alpha}{\Gamma(\alpha)(1-\alpha)} \int_0^t \frac{dt'}{(t-t')^{1-\alpha}} \frac{\partial}{\partial t'} \rho(x, t') \\ = Db \frac{\partial^2}{\partial x^2} \rho(x, t), \end{aligned} \quad (40)$$

which can also be written as

$$\frac{\partial}{\partial t} \rho(x, t) + \frac{\alpha}{1-\alpha} \frac{\partial^{1-\alpha}}{\partial t^{1-\alpha}} \rho(x, t) = Db \frac{\partial^2}{\partial x^2} \rho(x, t). \quad (41)$$

We note that Equation (41) is a special case of Equation (36) with $w(\nu) = \delta(\nu-1) + \frac{\alpha}{1-\alpha} \delta(\nu+\alpha-1)$. Analogously to results reported here, the solutions of Equation (41) are also characterized by two diffusive regimes [68, 69].

3. DISCUSSION AND CONCLUSIONS

We presented a detailed investigation of the changes in the fractional diffusion equation when the well-established Riemann-Liouville operator is replaced by the recently-proposed operators of Caputo-Fabrizio and Atangana-Baleanu. These changes are summarized in **Table 1**. Within the context of the continuous time random walk, we verified that these new fractional operators modify the behavior of the waiting time distribution. In the

Caputo-Fabrizio case, we found that the waiting time distribution is described by an exponential distribution; while the Atangana-Baleanu operator yields a distribution that decays as a stretched exponential for small times and as a power-law (with the same exponent of the Riemann-Liouville operator) for long times.

We obtained the exact solutions of the fractional diffusion equation and the time dependence of the mean square displacement when considering these different fractional operators. Our results reveal that these new operators lead to non-Gaussian distributions and different diffusive regimes depending on the time scale. For the Caputo-Fabrizio operator, the probability distribution $\rho(x, t)$ displays a stationary state as well as saturated diffusion for long times. This is a remarkable feature because the fractional diffusion equation is solved without external forces and subjected to the free diffusion boundary conditions. For the Atangana-Baleanu operator, we found a crossover between two diffusive regimes: a usual for small times and a sub-diffusive for long times, a feature observed in several empirical systems.

By properly manipulating the fractional diffusion equations, we demonstrated that the results obtained with these new fractional operators could be connected with other diffusive models. The fractional diffusion equation with the Caputo-Fabrizio operator recovers a diffusive process with stochastic resetting, where the fractional order exponent is directly related to the resetting rate. Also, the equation with the Atangana-Baleanu operator can be associated with a fractional diffusion equation with derivatives of distributed order. Our results thus suggest that these new fractional operators may be a simple and efficient way for incorporating different memory effects, opening new possibilities for modeling and investigating the anomalous diffusive processes.

AUTHOR CONTRIBUTIONS

All authors listed, have made substantial, direct and intellectual contribution to the work, and approved it for publication.

FUNDING

AT thanks the financial support of the CNPq under Grant No. 462067/2014-9. HR thanks the financial support of the CNPq under Grant No. 440650/2014-3 and CAPES. EL thanks the financial support of the CNPq under Grant No. 303642/2014-9.

REFERENCES

- Pearson K. The problem of the random walk. *Nature* (1905) 72:294. doi: 10.1038/072294b0
- Rayleigh L. On the resultant of a large number of vibrations of the same pitch and arbitrary phase. *Philos Mag.* (1880) 10:73. doi: 10.1080/14786448008626893
- Rayleigh L. *The Theory of Sound, 2nd Edn.*, Vol. 1. London: Macmillan (1894).
- Rayleigh, L. The problem of the random walk. *Nature* (1905) 72:318. doi: 10.1038/072318a0
- Bachelier L. Théorie de la Spéculation. *Ann Sci.* (1900) 17:21.
- Einstein A. Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen. *Ann Phys.* (1905) 17:549–60. doi: 10.1002/andp.19053220806
- von Smoluchowski M. Zur kinetischen theorie der brownischen molecularbewegung und der suspensionen. *Ann Phys.* (1906) 21:756–80. doi: 10.1002/andp.19063261405
- Montroll EW, Weiss GH. Random walks on lattices: II. *J Math Phys.* (1965) 6:167–81. doi: 10.1063/1.1704269
- Kutner R, Masoliver J. The continuous time random walk, still trendy: fifty-year history, state of art and outlook. *Eur Phys J B* (2017) 50:90. doi: 10.1140/epjb/e2016-70578-3
- Montroll EW, Scher H. Random walks on lattices. IV. Continuous-time walks and influence of absorbing boundaries. *J Stat Phys.* (1973) 9:101–35. doi: 10.1007/BF01016843
- Kenkre VM, Montroll EW, Shlesinger MF. Generalized master equations for continuous-time random walks. *J Stat Phys.* (1973) 9:45–50. doi: 10.1007/BF01016796
- Shlesinger MF. Asymptotic solutions of continuous-time random walks. *J Stat Phys.* (1974) 10:421–34. doi: 10.1007/BF01008803
- Havlin S, Avraham YB. *Diffusion and Reactions in Fractals and Disordered Systems.* Cambridge: Cambridge University Press (2000).
- Klafter J, Blumen A, Shlesinger MF. Stochastic path to anomalous diffusion. *Phys Rev A* (1987) 35:3081–5. doi: 10.1103/PhysRevA.35.3081
- Hilfer R, Anton L. Fractional master equations and fractal time random walks. *Phys Rev E* (1995) 51:R848–51. doi: 10.1103/PhysRevE.51.R848
- Compte A. Stochastic foundations of fractional dynamics. *Phys Rev E* (1996) 53:4191–3. doi: 10.1103/PhysRevE.53.4191
- Meztlar R, Klafter J. The random walk's guide to anomalous diffusion: a fractional dynamics approach. *Phys Rep.* (2000) 339:1–77. doi: 10.1016/S0370-1573(00)00070-3
- Meztlar R, Klafter, J. The restaurant at the end of the random walk: recent developments in the descriptions of anomalous transport by fractional dynamics. *J Phys A* (2004) 37:R161–R208. doi: 10.1088/0305-4470/37/31/R01
- Barkai E, Silbey RJ. Fractional Kramers equation. *J Phys Chem B* (2000) 104:3866–74. doi: 10.1021/jp993491m
- Barkai E. Fractional Fokker-Planck equation, solution, and application. *Phys Rev E* (2001) 63:046118. doi: 10.1103/PhysRevE.63.046118
- Barkai E, CTRW pathways to the fractional diffusion equation. *Chem Phys.* (2002) 284:13–27. doi: 10.1016/S0301-0104(02)00533-5
- Sung J, Barkai E, Silbey RJ, Lee S. Fractional dynamics approach to diffusion-assisted reactions in disordered media. *J Chem Phys.* (2002) 116, 2338–41. doi: 10.1063/1.1448294
- Barkai E, Metzler R, Klafter J. From continuous time random walks to the fractional Fokker-Planck equation. *Phys Rev E* (2000) 61:132–8. doi: 10.1103/PhysRevE.61.132
- Du M, Wang Z, Zu H. Measuring memory with the order of fractional derivative. *Sci Rep.* (2013) 3:431. doi: 10.1038/srep03431
- Zahran MA. On the derivation of fractional diffusion equation with an absorbent term and a linear external force. *Appl Math Model.* (2009) 33:3088–92. doi: 10.1016/j.apm.2008.10.013
- Zahran MA, Abulwafa EM, El-Wakil SA. The fractional Fokker Planck equation on comb-like model. *Phys A* (2003) 323:237–48. doi: 10.1016/S0378-4371(02)02026-5
- El-Wakil SA, Zahran MA, Abulwafa EM. Fractional (space time) diffusion equation on comb-like model. *Chaos Solit Fract.* (2004) 20:1113–20. doi: 10.1016/j.chaos.2003.09.032
- Villamaina D, Sarracino A, Gradenigo G, Puglisi A, Vulpiani A. On anomalous diffusion and the out of equilibrium response function in one-dimensional models. *J Stat Mech.* (2011) L01002. doi: 10.1088/1742-5468/2011/01/L01002
- Burioni R, Cassi D, Giusiano G, Regina S. Anomalous diffusion and Hall effect on comb lattices. *Phys Rev E* (2003) 67:016116. doi: 10.1103/PhysRevE.67.016116
- da Silva LR, Tateishi AA, Lenzi MK, Lenzi EK, da Silva PC. Green function for a non-Markovian Fokker-Planck equation: comb-model and anomalous diffusion. *Braz J Phys.* (2009) 39:483–7. doi: 10.1590/S0103-97332009000400025
- Iomin A. Subdiffusion on a fractal comb. *Phys Rev E* (2011) 83:052106. doi: 10.1103/PhysRevE.83.052106
- Baskin E, Iomin A. Superdiffusion on a comb structure. *Phys Rev Lett.* (2004) 93:120603. doi: 10.1103/PhysRevLett.93.120603
- Tateishi AA, Lenzi EK, Ribeiro HV, Evangelista LR, Mendes RS, da Silva LR. Solutions for a diffusion equation with a backbone term. *J Stat Mech.* (2011) P02022. doi: 10.1088/1742-5468/2011/02/P02022
- Metzler R, Nonnenmacher TF. Space- and time-fractional diffusion and wave equations, fractional Fokker Planck equations, and physical motivation. *Chem Phys.* (2002) 284:67–90. doi: 10.1016/S0301-0104(02)00537-2
- West BJ. *Fractional Calculus View of Complexity: Tomorrow's Science.* London: CRC Press (2016).
- Saxton MJ, Jacobson K. Single-particle tracking: applications to membrane dynamics. *Annu Rev Biophys Biomol Struct.* (1997) 26:373–99. doi: 10.1146/annurev.biophys.26.1.373
- Wirtz D. Particle-tracking microrheology of living cells: principles and applications. *Annu Rev Biophys.* (2009) 38:301–26. doi: 10.1146/annurev.biophys.050708.133724
- Gal N, Lechtman-Goldstein D, Weiss D. Particle tracking in living cells: a review of the mean square displacement method and beyond. *Rheol Acta* (2013) 52:425–43. doi: 10.1007/s00397-013-0694-6
- Hozé N, Holcman D. Statistical methods for large ensembles of super-resolution stochastic single particle trajectories in cell biology. *Annu Rev Stat Appl.* (2017) 4:189–223. doi: 10.1146/annurev-statistics-060116-054204
- Höfling F, Franosch T. Anomalous transport in the crowded world of biological cells. *Rep Prog Phys.* (2013) 76:046602. doi: 10.1088/0034-4885/76/4/046602
- Manzo C, Garcia-Parajo MF. A review of progress in single particle tracking: from methods to biophysical insights. *Rep Prog Phys.* (2015) 78:124601. doi: 10.1088/0034-4885/78/12/124601
- Shen H, Tauzin LJ, Baiyasi R, Wang W, Moringo N, Shuang B, et al. Single particle tracking: from theory to biophysical applications. *Chem Rev.* (2017) 117:7331–76. doi: 10.1021/acs.chemrev.6b00815
- Zagato E, Forier K, Martens T, Neyts K, Demeester J, De Smedt S, et al. Single-particle tracking for studying nanomaterial dynamics: applications and fundamentals in drug delivery. *Nanomedicine* (2014) 9:913–27. doi: 10.2217/nnm.14.43
- Kaerger J, Ruthven DM, Theodorou DN. *Diffusion in Nanoporous Materials.* Weinheim: Wiley-VCH Verlag (2012). doi: 10.1002/9783527651276
- Kaerger J, Ruthven D. Diffusion in nanoporous materials: fundamental principles, insights and challenges. *New J Chem.* (2016) 40:4027–48. doi: 10.1039/C5NJ02836A
- Caputo M, Fabrizio M. A new definition of fractional derivative without singular kernel. *Progr Fract Differ Appl.* (2015) 1:73–85. doi: 10.12785/pfda/010201
- Atangana A, Baleanu D. New fractional derivative with non-local and non-singular kernel. *Therm Sci.* (2016) 20:763–9. doi: 10.2298/TSCI16011018A
- Gómez-Aguilar JF, Atangana A. Fractional Hunter-Saxton equation involving partial operators with bi-order in Riemann-Liouville and Liouville-Caputo sense. *Eur Phys J Plus* (2017) 132:100. doi: 10.1140/epjp/i2017-11371-6
- Hristov J. Transient heat diffusion with a non-singular fading memory: from the Cattaneo constitutive equation with Jeffrey's Kernel to the Caputo-Fabrizio time-fractional derivative. *Therm Sci.* (2016) 20:757–62. doi: 10.2298/TSCI160112019H
- Gómez-Aguilar JF. Space time fractional diffusion equation using a derivative with nonsingular and regular kernel. *Phys A* (2017) 465:562–72. doi: 10.1016/j.physa.2016.08.072

51. Abdeljawad T, Baleanu D. Integration by parts and its applications of a new nonlocal fractional derivative with Mittag-Leffler nonsingular kernel. *J Nonlin Sci Appl.* (2017) **10**:1098–107. doi: 10.22436/jnsa.010.03.20
52. Podlubny I. *Fractional Differential Equations*. San Diego, CA; Academic Press (1999).
53. Hristov J. Derivation of fractional Dodson's equation and beyond: transient mass diffusion with a non-singular memory and exponentially fading-out diffusivity. *Progr Fract Differ Appl.* (2017) **3**:255–70. doi: 10.18576/pfda/030402
54. Hristov J. Derivatives with Non-Singular kernels from the Caputo - Fabrizio definition and beyond: appraising analysis with emphasis on diffusion models. In: Bhalekar S, editor. *Frontiers in Fractional Calculus*. Sharjah: Bentham Science Publishers (2017), p. 235–95.
55. Méndez V, Campos D. Characterization of stationary states in random walks with stochastic resetting. *Phys Rev E* (2016) **93**:022106. doi: 10.1103/PhysRevE.93.022106
56. Shkilev VP. Continuous-time random walk under time-dependent resetting. *Phys Rev E* (2017) **96**:012126. doi: 10.1103/PhysRevE.96.012126
57. Mathai AM, Saxena RK, Haubold HJ. *The H-Function: Theory and Applications*. New York, NY: Springer (2009).
58. Mo J, Simha A, Raizen MG. Broadband boundary effects on Brownian motion. *Phys Rev E* (2015) **92**:062106. doi: 10.1103/PhysRevE.92.062106
59. Berry H, Chaté H. Anomalous diffusion due to hindering by mobile obstacles undergoing Brownian motion or Ornstein-Uhlenbeck process. *Phys Rev E* (2014) **89**:022708. doi: 10.1103/PhysRevE.89.022708
60. Koslover EF, de la Rosa MD, Spakowitz AJ. Crowding and hopping in a protein's diffusive transport on DNA. *J Phys A Math Theor.* (2017) **50**:074005. doi: 10.1088/1751-8121/aa53ee
61. Evans MR, Majumdar SN. Diffusion with stochastic resetting. *Phys Rev Lett.* (2011) **106**:160601. doi: 10.1103/PhysRevLett.106.160601
62. Skalski GT, Gilliam JF. Modeling diffusive spread in a heterogeneous population: a movement study with stream fish. *Ecology* (2000) **81**:1685–1700. doi: 10.1890/0012-9658(2000)081[1685:MDSIAH]2.0.CO;2
63. Upadhyaya A, Rieub JP, Glaziera JA, Sawadac Y. Anomalous diffusion and non-Gaussian velocity distribution of Hydra cells in cellular aggregates. *Phys A* (2001) **293**:549–58. doi: 10.1016/S0378-4371(01)00009-7
64. Mieruszynski S, Digman MA, Gratton E, Jones MR. Characterization of exogenous DNA mobility in live cells through fluctuation correlation spectroscopy. *Sci Rep.* (2015) **5**:13848. doi: 10.1038/srep13848
65. Ribeiro HV, Tateishi AA, Alves LGA, Zola RS, Lenzi EK. Investigating the interplay between mechanisms of anomalous diffusion via fractional Brownian walks on a comb-like structure. *New J Phys.* (2014) **16**:093050. doi: 10.1088/1367-2630/16/9/093050
66. Alves LG, Scariot DB, Guimaraes RR, Nakamura CV, Mendes RS, Ribeiro HV. Transient superdiffusion and long-range correlations in the motility patterns of trypanosomatid flagellate protozoa. *PLoS ONE* (2016) **11**:e0152092. doi: 10.1371/journal.pone.0152092
67. Tateishi AA, Lenzi EK, da Silva LR, Ribeiro HV, Picoli S Jr, Mendes RS. Different diffusive regimes, generalized Langevin and diffusion equations. *Phys Rev E* (2012) **85**:011147. doi: 10.1103/PhysRevE.85.011147
68. Chechkin AV, Gorenflo R, Sokolov IM. Retarding subdiffusion and accelerating superdiffusion governed by distributed-order fractional diffusion equations. *Phys Rev E* (2002) **66**:046129. doi: 10.1103/PhysRevE.66.046129
69. Lenzi EK, Mendes RS, Tsallis C. Crossover in diffusion equation: anomalous and normal behaviors. *Phys Rev E* (2003) **67**:031104. doi: 10.1103/PhysRevE.67.031104
70. Bian X, Kimb C, Karniadakis GE. 111 years of Brownian motion. *Soft Matt.* (2016) **12**:6331. doi: 10.1039/C6SM01153E
71. Huang R, Chavez I, Taute KM, Lukić B, Jeney S, Raizen MG, Florin EL. Direct observation of the full transition from ballistic to diffusive Brownian motion in a liquid. *Nat Phys.* (2011) **7**:576–80. doi: 10.1038/nphys1953
72. Caputo M. Mean fractional-order-derivatives differential equations and filters. *Ann Univ Ferrara Sez* (1995) **41**:73–84.
73. Caputo M. Distributed order differential equations modelling dielectric induction and diffusion. *Fract Calc Appl Anal.* (2001) **4**:421–42.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Tateishi, Ribeiro and Lenzi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.