



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The role of global dietary transitions for safeguarding biodiversity

Citation for published version:

Henry, R, Alexander, P, Rabin, S, Anthoni, P, Rounsevell, M & Arneith, A 2019, 'The role of global dietary transitions for safeguarding biodiversity', *Global Environmental Change*, vol. 58.
<https://doi.org/10.1016/j.gloenvcha.2019.101956>

Digital Object Identifier (DOI):

[10.1016/j.gloenvcha.2019.101956](https://doi.org/10.1016/j.gloenvcha.2019.101956)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Global Environmental Change

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 **The role of global dietary transitions for safeguarding biodiversity.**

2
3

4 **Abstract**

5

6 Diets lower in meat could reduce agricultural expansion and intensification thereby reducing
7 biodiversity impacts. However, land use requirements, associated with alternate diets, in
8 biodiverse regions across different taxa are not fully understood. We use a spatially explicit
9 global food and land system model to address this gap. We quantify land-use change in locations
10 important for biodiversity across taxa and find diets low in animal products reduce agricultural
11 expansion and intensity in regions with high biodiversity. Reducing ruminant meat
12 consumption alone however was not sufficient to reduce fertiliser and irrigation application in
13 biodiverse locations. The results differed according to taxa, emphasising that land-use change
14 effects on biodiversity will be taxon specific. The links shown between global meat consumption
15 and agricultural expansion and intensification in the biodiverse regions of the world indicates
16 the potential to help safeguard biodiverse natural ecosystems through dietary change.

17 **Key words:** biodiversity, land use change, diet, consumption

18

19

20

21 **1. Introduction**

22

23 Expansion of agricultural land, together with intensified management practices are some of the
24 greatest threats to the conservation of terrestrial ecosystems and biodiversity (Machovina et al.,
25 2015; Machovina and Feeley, 2014; Marchal et al., 2011; Newbold et al., 2015; Ripple et al.,
26 2014a). Over 35% of the Earth's permanent ice-free land surface is currently used for food
27 production (Foley et al., 2005), with the expansion of agricultural land for food production in
28 the last 300 years having reduced natural grasslands by up to fifty percent and natural forests
29 by one third (Goldewijk, 2001). The associated loss of natural ecosystems has had negative
30 consequences for biodiversity (Gibson et al., 2011; Pereira et al., 2012; Pimm et al., 2014).
31 Agricultural intensification that increases yields can reduce the area of land needed for
32 production, but can also harm biodiversity through fertiliser and pesticide pollution (Flohre et
33 al., 2011; Gibbs et al., 2009; Kleijn et al., 2009) as well as irrigation abstraction on ecological
34 river flows (De Frutos et al., 2015; Yamaguchi and Blumwald, 2005). Land-use change models
35 have demonstrated that biodiverse regions will be significantly threatened by future
36 agricultural expansion and intensification (Delzeit et al., 2017; Kehoe et al., 2017, 2015).
37 Protected areas can be an effective contribution to prevent agricultural expansion (Pringle,
38 2017), but conservation efforts that focus on food demand will also play a role.

39

40 Meat production has been associated with higher land and water use, and higher GHG
41 emissions, per unit of energy or protein than other foods (Machovina et al., 2015; Poore and
42 Nemecek, 2018; Tilman and Clark, 2014a). In particular, heavily managed and densely stocked
43 pastures pose serious threats to biodiversity (Machovina and Feeley, 2014; Ripple et al., 2014a).
44 65% of agricultural expansion in recent decades has been associated with increased production
45 of animal products (Alexander et al., 2015), and land-use changes associated with animal
46 husbandry account for roughly 30% of current global biodiversity loss (Westhoek et al., 2011).
47 Livestock production is increasing most rapidly in tropical regions with high biodiversity
48 (Machovina et al., 2015). The tropics are also experiencing the highest rates of species
49 extinction (Dirzo et al., 2014), at a time when global extinction rates have been estimated to be
50 1000 times the geological background rate (Pimm et al., 2014, 1995). Much future human
51 population growth is expected to occur in these biodiverse tropical nations, and as incomes
52 continue to rise in developing countries, animal product consumption is expected to increase
53 further (Machovina et al., 2015; Stoll-Kleemann and Schmidt, 2017). If current trends in animal
54 product consumption continue, and if industrialised countries do not reduce high rates of meat
55 consumption, it is estimated that one billion additional hectares of natural land will be cleared
56 for agriculture by 2050 (Tilman et al., 2011, 2001).

57

58 Reducing meat consumption would not only improve global human health—consumption of
59 meat in industrialised countries is currently double the amount that is deemed healthy
60 (Wellesley et al., 2015)—but the Intergovernmental Panel on Climate Change (IPCC) also
61 identified it as an important focus for climate change mitigation (Pachauri et al., 2014).
62 Modelling studies have quantified land-use changes associated with dietary shifts,
63 demonstrating that demand-side reductions in meat consumption could reduce GHG emissions
64 and deforestation (Bajželj et al., 2014; Erb et al., 2016; Popp et al., 2010; Stehfest et al., 2009;
65 Tilman and Clark, 2014a; Wirsenius et al., 2010). However, fewer studies (Kok et al., 2018;
66 Tilman et al., 2017; Visconti et al., 2016) have considered the effects of diet on biodiversity, and

67 none have explored the spatial impacts across multiple taxa. The spatial nature of biodiversity
68 and variations in distributions between taxa means that spatially explicit analyses are required
69 to understand the impact of dietary choices on biodiversity.

70

71 Here we address this critical gap in understanding the environmental consequences of food-
72 system changes. We use a global food-system model (PLUMv2/LPJ-GUESS, Alexander et al.,
73 2018) to explore land use and agricultural intensity change until 2100 under three alternative
74 dietary scenarios: Business-as-usual (BAU), 95% reduction in ruminant product consumption
75 (LOW-R), 95% reduction in animal product consumption (LOW-AP). This work is unique in
76 considering the spatially disaggregated consequences of future dietary scenarios for high
77 biodiversity locations across different taxa. We also, for the first time, consider the role of
78 nitrogen and irrigation intensity changes on locations important for biodiversity.

79

80 2. Methods

81

82 2.1. Modelling framework

83 PLUMv2 is a global land use and food-system model that combines spatially-explicit,
84 biophysically-derived yield responses with socio-economic scenario data to project future
85 demand, land use, and management inputs (Alexander et al., 2018). For each country and time-
86 step, the agricultural land use and level of imports or exports is determined through a least-cost
87 optimisation that meets the demand for food and bioenergy commodities in each country. Food
88 demand is projected based on log-linear relationships with per-capita income using GDP and
89 populations from the Shared Socioeconomic Pathway (SSP) scenarios (O'Neill et al., 2014).
90 Demand for food and bioenergy commodities is projected at a country level for six commodity
91 groups: cereals, oilcrops, pulses, starchy roots, ruminant products, and monogastric products.
92 Demand for dedicated energy crops (i.e., second-generation bioenergy) is specified as a global
93 trajectory with all production locations determined endogenously. Food and bioenergy demand
94 is met by in-country expansion or intensification of crops or from imports from the global
95 market. ~~Over~~ Production of commodities in ~~excess of a~~ country's domestic demand are
96 exported to the global market. ~~In PLUMv2 supply and demand in t~~The global market is not
97 constrained to ~~be in~~ equilibrium, ~~with~~ over- or under- supply of commodities ~~can be~~ buffered
98 through explicitly modelled stocks. ~~For each commodity a single tariff free price exists in each~~
99 ~~time step and the initial price was set exogenously but subsequently adjusted in each of the~~
100 ~~following time periods according to under- or oversupply in the global market. Prices are~~
101 ~~updated for the next year based on the aggregate imbalance of imports and exports in that year.~~
102 ~~For example over supply of a commodity on the global market decreases the price; this reduces~~
103 ~~the benefits from its export and reduces the cost of importing it, creating a tendency to correct~~
104 ~~for the oversupply. For each commodity a single tariff free price exists in each time step, which~~
105 ~~is adjusted for transport costs and other barriers, e.g. tariffs, to obtain country specific prices.~~
106 ~~For example over supply of a commodity decreases the price; this reduces the benefits from its~~
107 ~~export and reduces the cost of importing it, creating a tendency to correct for the oversupply.~~

108 Crop yield responses used in PLUMv2 are provided on a 0.5° grid by a dynamic global
109 vegetation model, LPJ-GUESS (Smith et al., 2014), for a range of fertilisation rates and rain-fed
110 vs irrigated conditions. Other management practices (e.g. pesticide application, machinery
111 stock, reseedling of grassland) are represented in PLUMv2 by a "management intensity" factor.
112 Natural land cover here is comprised of forested primary land, non-forested primary land and
113 abandoned agricultural land. **In the grid cells four decision variables (i.e. area, fertiliser,**

Commented [HR1]: In general I've added to the methods section to address the following comment

'The authors refer to a recent model development paper but to stand at its own more detailed information is needed.'

Commented [HR2]: REVIEWER COMMENT: How is trade dealt with within the modelling framework? It is mentioned in line 93-4 but not expanded on.

115 irrigation and other intensity) for each of the eight land use types (seven crop types plus
116 pasture) are determined during the optimisation, resulting in more than ??,000 decision
117 variables in each year. To achieve this PLUMv2 uses the spatially specific crop yield responses
118 to intensity inputs, various land use costs, such as land conversion costs and input costs, and
119 trade costs. This ultimately determine land use solutions that meet country level demand.

Commented [AP3]: What 'this'? Just "PLUMv2 uses...?"

Commented [AP4]: Again, unclear what 'this' refers to.

121 Socioeconomic parameters are in line with the "middle of the road" SSP scenario (SSP2), which
122 also provided population and GDP trajectories (Dellink et al., 2017; Jones and O'Neill, 2016). The
123 SSPs describe alternative global societal pathways through the 21st century (O'Neill et al., 2015,
124 2014). SSP2 is the middle of the road pathways with trends largely exhibiting historic patterns.

Commented [HR5]: Reviewer comment: **Line 309-311**
"plum chooses not to abandon existing cropland" - is
this a results or is PLUMv2 hard-coded into the model
to do this?

125 Population and GDP trajectories were taken from SSP2 using World Bank projections (IIASA,
126 2014), and demand for food commodities were taken from FAOSTAT (FAOSTAT, 2015a, 2015b).
127 The climate and atmospheric CO₂ forcing scenario RCP 6.0 was used as it is considered the
128 Representative Concentration Pathway (van Vuuren et al., 2011) most consistent with SSP2
129 (Engström et al., 2016). Forcings were taken from the 1850–2100 IPSL-CM5A-MR outputs from
130 the Fifth Coupled Model Intercomparison Project (CMIP5; 71). First- and second-generation
131 bioenergy demand trajectories are specified exogenously to represent a business-as-usual
132 scenario with no specific climate change mitigation policies. Demand for first-generation
133 bioenergy is modelled from an observed baseline level of demand (Alexander et al., 2015;
134 FAOSTAT, 2015a) adjusted to double by 2030 from the 2010 level and thereafter remain
135 constant. Global demand for dedicated second-generation bioenergy crops increases to 4000 Mt
136 DM/year by 2100, in line with the SSP2 demand with baseline assumptions (Popp et al., 2016).
137 For parameter settings that were not specified exogenously from available data on existing and
138 future trends, for example technology change rates, expert judgement was used to align
139 quantitative parameter settings with the qualitative SSP2 storyline. Scenario elements of the
140 SSP2 narrative that were assumed to influence changes in the PLUMv2 input parameters were
141 identified. Qualitative changes in parameters were estimated based on an interpretation of the
142 SSP2 storyline (Engström et al., 2016). These qualitative estimates of parameters and
143 uncertainty levels were translated into quantitative values characterised by a uniform
144 distribution. Each parameter therefore had a range defined by a range of 50% above and below
145 the central parameter values. A Monte Carlo approach to explore uncertainty associated with
146 input parameters was used and parameters were sampled using a Sobol sequence method with
147 n = 30 (Chalaby et al., 2015); the central parameter values used in each of the scenarios can be
148 found in Appendix C, Table C2. This approach allowed us to capture the uncertainty within the
149 model framework.

150

151 2.2. Protected areas

152 The proportion of protected land with a status of "designated" and IUCN category I–VI within a
153 grid cell is calculated using the WDPA database (UNEP-WCMC., 2016). Within each grid cell,
154 natural land designated as protected is prevented from conversion to any form of agricultural
155 use. Within each grid cell, a minimum fraction (5%) of primary unprotected natural land is also
156 prevented from agricultural use due to assumed limits to agricultural production, e.g. field
157 boundaries, roads/tracks, and other farm infrastructure. Slope constraints (IIASA/FAO, 2010)
158 also prevent agricultural use in regions of high altitude. In cases where agricultural land already
159 exceeds the area specified as protected, no further agricultural expansion can occur. China's
160 National Forest Protection Program is implemented as an annual limit to deforestation of 1.1%
161 in China. (Ren et al., 2015).

162

163

164 2.3. Scenario description

165

166 2.3.1. Business as usual (BAU)

167 This scenario assumes that the shift in consumption away from staples, such as pulses and
168 starchy roots, and towards animal products continues as incomes rise. The relationship
169 between rising income and increasing consumption of commodities such as meat, milk, and
170 refined sugars has been observed historically and therefore in line with the SSP2 pathway, we
171 assume future consumption trends in the baseline largely exhibit historic patterns (Keyzer et
172 al., 2005; Tilman et al., 2011).

173 2.3.2. 95% reduction in ruminant product consumption (LOW-R)

174 This scenario represents a major shift in world consumption patterns of ruminant products, this
175 could be potentially driven by increasing meat prices induced by stricter climate and health
176 policies, consumer awareness and increasing land and animal feed expenses. This scenario
177 assumes that the consumption of ruminant products decreases steadily from 2010 to 2100 until
178 crop products replace 95% of ruminant product consumption. Ruminant products are replaced
179 by a mixture of cereals, starchy roots, pulses, and oilcrops; however, the same calorie intake is
180 maintained. An example of the dietary changes in terms of per capita consumption and the
181 proportions of substitution and are given in Appendix C, Table C1 and Table C2 respectively.
182 95% was chosen for the stylised scenarios to demonstrate the potential effects a very large, but
183 not total, reduction in ruminant product consumption would have on global land use.

184 2.3.3. 95% reduction in animal product consumption (LOW-AP)

185 This scenario is similar to the above, but assumes that non-meat commodities replace both
186 ruminant and monogastric consumption. As with the LOW-R scenario 95% was chosen for to
187 demonstrate the potential effects a very large, but not total, reduction in animal product
188 consumption would have on global land use.

189 2.4. Exploring the consequences of dietary change for biodiversity

190

191 2.4.1. Conservation International (CI) biodiversity hotspots

192

193 The 35 CI hotspots cover 2.3% of the land surface but support 50% of the world's endemic
194 plant species and 43% of vertebrate endemic species. To qualify as a hotspot, a region must
195 be threatened —i.e. contain at most 30% of its original natural vegetation—yet contain at
196 least 1500 endemic vascular plants. The CI biodiversity hotspot database is used to identify
197 particular regions of importance for biodiversity (Mittermeier et al., 2004; Myers et al.,
198 2000). CI hotspot shapefile data are converted to 0.5° raster maps. Any 0.5° cell containing CI
199 hotspot polygon data is classified as a CI hotspot irrespective of hotspot size. The CI map is
200 therefore binary and cells are classified as either a CI hotspot or not.

201

202 2.4.2. Vertebrate species richness maps

203

204 Criteria for the biodiversity hotspots database only account for vascular plant species
205 richness. Thus, we also consider maps of vertebrate species richness, small-range vertebrate
206 species richness, and threatened species richness (Jenkins et al., 2013; Pimm et al., 2014).
207 The resolution of the vertebrate species richness maps was decreased from 0.1° to 0.5°
208 resolution to match PLUMv2; the mean species richness was calculated for each grid cell. For
209 all taxa, the distribution of species richness across grid cells is right-skewed: most cells contain
210 a few species while there are a few cells with a large number of species. For each taxon's map
211 we therefore convert the mean species richness values of grid cells into percentile values

Commented [AP6]: Might read better for each of the scenarios to phrase first sentence as.

"The business-as-usual scenario ..." or equivalent.

Not essential

Commented [HR7]: REVIEWER COMMENT: On the scenario description (section 2.3) it would be good to have more contextual information used in the justification of the settings chosen for the modelling. Were they taken from existing trends or entirely taken from the expert judgment of the modelling team. To what extent are the settings chosen plausible?

I have also added a section to the results/discussion about limitations related to this.

212 (richness index). We assume that ‘species-rich regions’ comprise cells with a richness index \geq
213 0.9, i.e. the 90th percentile of grid cells and therefore, similar to the CI hotspots, we focus on
214 those regions with the greatest biodiversity (appendix A, figures A1-3).

215
216 We explore land use change, agricultural expansion, and intensification projected by PLUMv2 in
217 CI hotspots and in vertebrate species-rich regions for the different dietary scenarios. We
218 consider the loss of natural land, forests, and natural grasslands and changes in input intensities
219 such as fertiliser and irrigation in grid cells classed either as CI hotspots or with a richness index
220 \geq 0.9. From this, we identify regions of threat using a threat index: regions with high
221 biodiversity that overlap with areas of projected agricultural expansion. We calculate this
222 overall threat index for all species in each 0.5° grid cell. This is the proportion of natural land
223 projected to be lost by 2100 multiplied by the summed richness index of birds, mammals, and
224 amphibians for the median PLUMv2 parameter simulation run. For the threat index we
225 therefore assume each species is equally important regardless of taxon.
226

227 3. Results and discussion

228

229 3.1. Land cover change in biodiverse regions


230 In agreement with results from previous modelling studies (Delzeit et al., 2017; Kehoe et al.,
231 2017, 2015), the most threatened regions—locations with high biodiversity under pressure
232 from agricultural expansion—are in the tropics under BAU scenarios (Figure 1). Scenarios of
233 lower animal product consumption (LOW-R and LOW-AP) greatly reduce agricultural expansion
234 in regions of high biodiversity compared to the BAU scenario (Figure 1 & 2). By 2100, 9% (984
235 Mha) of global natural land is lost, of which 95% is in the tropics—equivalent to 24% of natural
236 land in these latitudes (Figure 2). Conversely, reduced animal product consumption (LOW-AP)
237 resulted in a 7% (703 Mha) increase in global natural land between 2010–2100 (Figure 2,
238 Appendix B Figure B1) with lower losses across the tropics (Figure 1, Figure 2) and increases in
239 natural land across the northern hemisphere (Figure 2). Deforestation and land clearing for
240 agriculture have been identified as the leading causes of biodiversity decline (Gibson et al.,
241 2011). Therefore, the potential for dietary change to reduce global agricultural expansion by
242 approximately 1687 Mha (11% of global land area) is an important finding for biodiversity
243 conservation (Laurance et al., 2012; Pereira et al., 2012).

244 Species-rich regions across the different taxa are largely found in the tropics (Appendix A,
245 Figure A1-3) and the greatest loss of natural land in species-rich regions occurs in BAU (Figure
246 3). In BAU, on average, 98% of global pasture expansion takes place in the tropics as demand
247 for ruminant products in tropical countries increases with increasing population and income
248 (Appendix C, Figure C2). As incomes increase, consumption shifts from staples such as starchy
249 roots and pulses to commodities such as meat, milk, and refined sugars (Keyzer et al., 2005;
250 Tilman et al., 2011). However, the rate of increasing consumption of animal products slows and
251 plateaus with any further rise in income (Cole and Mccoskey, 2013), which is also represented
252 in the log-linear relationships with per-capita income used in our model (Alexander et al.,
253 2018). Consequently, in developing tropical countries, the transition from low incomes to high
254 incomes results in greater demand of ruminant products (Appendix C, Figure C2), and pasture
255 expands at the expense of natural land. In contrast, income and the animal product consumption
256 in developed countries outside the tropics are already high, with large areas of existing pasture
257 meeting demand for ruminant products. Given the relationship between income and
258 consumption, increases in income in developed countries do not lead to further large increases
259 in demand for animal products (Appendix C, Figure C2). Under LOW-AP and LOW-R,

260 abandonment of existing pasture in developed countries leads to large increases of natural land
261 at a global level (Figure 2). This does not coincide with large increases in natural land in
262 species-rich regions (Figure 3), however, because it largely takes place in locations that are not
263 here classified as species-rich — i.e. those outside the tropics. This result can be seen when
264 comparing Figures A1–A3 in Appendix A (spatial distribution of species-rich regions for the
265 different taxa) with Figure 2. In species-rich regions the LOW-R and LOW-AP scenarios reduce
266 pasture expansion rather than increase natural land. Although existing pasture in the tropics is
267 also abandoned, this is offset by cropland expansion (see below); therefore natural land area in
268 species-rich regions under LOW-R and LOW-AP is relatively stable, compared to BAU (Figure 3).

269 Cropland expands by 28% in the tropics under BAU to produce crops for food and as animal
270 feed as demand for animal products grows in the developing world. Under LOW-R, cropland
271 expands by 38% in the tropics, this is greater than under BAU because demand for food and
272 feed for monogastrics is the same while additional crops are required to replace ruminant
273 products. Under LOW-AP, despite reduced demand for feed for animals, existing cropland area
274 in 2010 is not sufficient to produce enough crops to replace animal products and meet food
275 demand of a growing population; consequently, cropland still expands by 27%. The greater
276 cropland requirements under LOW-R explain the marginally greater losses of natural land in
277 species-rich regions (Figure 3) in LOW-R compared to LOW-AP. However, on average, the
278 amounts of water and nitrogen applied to cropland in the tropics under LOW-AP are 42% and
279 68% less, respectively, than under BAU. Therefore, while the total area of cropland remains the
280 same in the tropics, the intensity of agricultural inputs declines under LOW-AP with the
281 reduction in demand for animal feed.


282
283
284



285

286 *Figure 1: Spatial distribution of regions of threat; regions with high biodiversity under pressure*
 287 *from agricultural expansion. The left column (a,c,e) is the BAU scenario and the right column*
 288 *(b,d,f) is the LOW-AP scenario for the different types of species richness. Blue dashed lines*
 289 *delineate the tropics.*

290




291 *Figure 2: Change in natural land cover fraction between 2010-2100 for (a) BAU (b) LOW-*
 292 *AP (c) LOW-R. Dotted lines delineate the tropics.*
 293

294
 295
 296 Tilman et al., (2017) investigated the biodiversity of mammals and birds in a non-spatial,
 297 country-level approach and found dietary change reduced extinction risk. However, previous
 298 spatially explicit studies typically only consider single taxon with amphibians particularly
 299 underrepresented. For example, Visconti et al., (2016), considering only mammals, found
 300 consumption change could reduce extinction risk. The locations classified here as species-rich
 301 differ between mammals, birds and amphibians (see Appendix A, Figures A1–3). Differences
 302 regarding the impacts of land-use change therefore arise between and within taxa, and are
 303 important when considering conservation targets (Ceballos and Ehrlich, 2006; Jenkins et al.,
 304 2013; Orme et al., 2005; Pimm et al., 2014; Possingham and Wilson, 2005). For example,
 305 natural land loss in parts of Ecuador overlap to a greater extent with regions of threatened bird
 306 species-richness than with regions of threatened mammal-species-richness. While broad
 307 patterns are similar across CI hotspots and taxa—for example, the greatest loss of natural land
 308 occurs in BAU while the LOW-R and LOW-AP scenarios result in smaller losses or increasing
 309 natural land cover (Figure 3)—important differences remain. For example, with LOW-AP, while
 310 the small ranged species-rich regions shows little change or decreases in natural land cover
 311 by 2100, the grid cells within the threatened species-rich region show increasing natural land
 312 cover by 2100. Changing dietary patterns may therefore have the greatest benefits for
 313 threatened species in term of habitat recovery. Measures of total species richness are
 314 important when considering threats to overall range size and ecosystem functioning related
 315 to population sizes. However, the richness of small-ranged species and/or threatened species
 316 are often regarded as more appropriate measures when planning conservation to prevent
 317 extinctions (Ceballos and Ehrlich, 2006). Visconti et al., (2015), for example, highlighted the
 318 importance of considering the status of taxa from a protected area perspective: Targeting
 319 protection towards threatened species had positive effects on suitable habitat for terrestrial
 320 mammals, while expanding protected areas according to ecoregion targets had negative effects.
 321 Furthermore, a number of studies have demonstrated that hotspots of species richness
 322 between taxa and classifications of taxa are often incongruent (Jenkins et al., 2013; Orme et

Commented [HR8]: REVIEWER COMMENT: The directions of the trends discussed aren't really surprising, but perhaps there is something in the magnitudes or spatial patterns that could be picked out to highlight the novelty more keenly?

Commented [AP9R8]: Hard to know what's been added and therefore to assess how well it deals with the point raised.

323 al., 2005; Pimm et al., 2014). The differences between CI hotspots and type of taxa here
 324 further support the argument that no single metric is sufficient when considering threats to,
 325 and the conservation of, biodiversity (Ceballos and Ehrlich, 2006; Jenkins et al., 2013; Orme
 326 et al., 2005; Pimm et al., 2014; Possingham and Wilson, 2005).



327
 328 *Figure 3: Projected natural land change by 2100 in (a) bird-, (b) mammal-, and (c) amphibian-*
 329 *species-rich regions, and (d) CI hotspots for the different dietary scenarios. Species-rich regions are*
 330 *comprised of cells with a richness index ≥ 0.9 . Colours in a–c represent the different types of*
 331 *species-rich regions: all species (blue), small-ranged species (orange), and threatened species*
 332 *(red). Boxplots distributions generated with $n=30$.*

333

335 The type and level of agricultural management has an important role in the impact on different
 336 taxa, for example, at a European scale Flohre et al., (2011) found that while agricultural
 337 intensity negatively affected the species richness of birds it did not affect carabid beetles (De
 338 Frutos et al., 2015; Flohre et al., 2011; Gibbs et al., 2009; Kleijn et al., 2009; Yamaguchi and
 339 Blumwald, 2005). ~~It has been shown that~~ increasing nitrogen, in particular, reduces plant
 340 biodiversity (Bobbink et al., 2010; Reich, 2009; Stevens et al., 2004) with consequences for
 341 faunal biodiversity (Nijssen et al., 2017). ~~In their review~~ Bobbink et al., (2010) highlighted that
 342 the negative effects of nitrogen accumulation on biodiversity has occurred across a wide range
 343 of ecosystems and geographic areas. Nijssen et al., (2017) ~~recently~~ identified ten pathways
 344 through which increased nitrogen alters faunal biodiversity. N-driven faunal decline has been
 345 demonstrated, for example, in some rare bird species where elevated nitrogen reduced
 346 vegetation heterogeneity and/or preferred habitat with consequences for prey abundance (de
 347 Vries et al., 2011). However, ~~while pathways identified by Nijssen et al., (2017) were supported~~
 348 ~~by peer reviewed literature~~ there remains knowledge gaps regarding the mechanisms that drive
 349 observed biodiversity changes. ~~In terms of~~ ~~Similarly, irrigation~~ there is a wide body of evidence
 350 that demonstrates the negative effects of water extraction on natural ecosystems, ~~with and~~
 351 ~~agricultural water use is one of the leading causes of the majority of~~ freshwater withdrawal ~~in~~
 352 ~~the world globally used in agriculture~~. The disruption of water flows and river regulation has, for
 353 example, altered floodplain forests resulting in their dieback globally. Such forests are
 354 ecologically important due to their high biodiversity, ~~with and~~ climate change induced droughts
 355 are likely to further exacerbate forest mortality (Horner et al., 2009). In Spain, the expansion of
 356 irrigated agriculture has coincided with the disappearance of up to 61% of biodiverse wetlands
 357 over the last fifty years (Fuentes-Rodríguez et al., 2013). Furthermore, intensive livestock
 358 farming that involves irrigation has also been found to substantially alter water chemistry of
 359 nearby rivers with potential consequences for both aquatic and riparian species diversity
 360 (Martín-Queller et al., 2010). Given the ~~widespread~~ implications ~~for biodiversity~~ of increasing
 361 nitrogen and irrigation use ~~for biodiversity~~ the need to consider such consequences are
 362 apparent. However, no previous land use modelling studies have explored changes in irrigation
 363 and nitrogen fertiliser intensities that are associated with reductions in meat consumption in
 364 biodiverse regions. Agricultural intensity is typically represented in land-use models in a
 365 stylised and spatially aggregated manner, making the evaluation of their impacts challenging
 366 (Lotze-Campen et al., 2008; Nelson et al., 2014). Our analysis addresses this gap and by
 367 including spatially specific crop responses to different inputs in our modelling framework, ~~We~~
 368 are able to show the effects of dietary changes on ~~intensity~~ inputs ~~intensity, with a focus on in~~
 369 species-rich regions ~~where biodiversity impacts are likely to be most acute~~.

370 In the LOW-R scenario, demand for monogastric feed crops is unchanged from BAU, while
 371 demand for food crops increases to replace ruminant products (Appendix C, Figure C1). This net
 372 increase in crop demand results in crop area expansion relative to BAU. However average global
 373 fertiliser and irrigation intensities (on a per-area basis) change similarly to BAU; the median
 374 increase in nitrogen and irrigation, in CI hotspots and species-rich regions, under LOW-R are
 375 similar to BAU (Figure 4, Figure 5). LOW-AP decreases demand relative to BAU for
 376 monogastrics, as well as ruminants, and consequently decreases demand for crops as feed
 377 (Figure C1). Rather than reduce cropland area this results in reduced nitrogen and water inputs
 378 in these locations. From a fertiliser and irrigation perspective, reduced feed production
 379 therefore has the greatest potential to reduce inputs and replacing pasture-fed ruminant
 380 products alone may not have substantial benefits for biodiversity.

Commented [AP10]: Compared to what else?

Commented [AP11]: Two sentences of the form "A said <something>". Could at least one, perhaps both be rephrased so avoid that rather boring structure.

Commented [AP12]: Delete? Seems to add little.

Commented [AP13]: 70% from memory!? Worth a reference too. Aquastat?


Commented [AP14]: Right word? "retreat" "reduction in area". Might need a reordering of phrase.

Commented [AP15]: Rather weak evidence! Sure lots of things have coincided with that. Coincided with increased call for Catalanian independence as well!

Commented [HR16]: REVIEWER COMMENT: The link between nitrogen application and biodiversity is clear, but could also do with a little more justification. The link between irrigation and biodiversity is less clear-cut. What are the thoughts / concerns here?

382 The differences in nitrogen and water inputs between alternate dietary scenarios highlight the
 383 need to consider fertiliser and irrigation individually. Any potential intensity changes associated
 384 with dietary change will require scrutiny as measures to reduce agricultural expansion may not
 385 necessarily reduce intensification. We do not consider a scenario with a 'livestock revolution'—
 386 a shift away from pasture-based production toward industrialised production that requires
 387 crop-based feeds (Delgado et al., 2001; Naylor et al., 2005; Swain et al., 2018)—which could
 388 similarly reduce the rate of agricultural expansion, but with increased intensity. There is an
 389 inherent trade-off between agricultural intensification and expansion. Intensification is more
 390 polluting, but requires less land, while expansion is less polluting, but requires more land.
 391 Ultimately, both can have negative consequences for biodiversity and thus managing this trade-
 392 off is complex. For example, the recent IPBES Regional Assessment for Europe & Central Asia
 393 recommends that Europe reduce agricultural intensity to conserve European biodiversity
 394 (IPBES, 2018). However, this could displace food production and the associated consequences
 395 for biodiversity, through imports, to other parts of the world.

396




397

398 *Figure 4: Projected change in irrigation water use by 2100 in (a) bird-, (b) mammal-, and (c)*
 399 *amphibian-species-rich regions and (d) CI hotspots for the different dietary scenarios. Species-rich*
 400 *regions are comprised of cells with a richness index ≥ 0.9 . Colours in a–c represent the different*

401 types of species-rich regions: all species (blue), small-ranged species (orange), threatened species
 402 (red). Boxplots distributions generated with n=30.

403



404

405 *Figure 5: Projected nitrogen fertiliser intensity change by 2100 in (a) bird-, (b) mammal-, and (c)*
 406 *amphibian-species-rich regions and (d) CI hotspots for the different dietary scenarios. Species-rich*
 407 *regions are comprised of cells with a richness index ≥ 0.9 . Colours in a-c represent the different*
 408 *types of species-rich regions: all species (blue), small-ranged species (orange), threatened species*
 409 *(red). Boxplots distributions generated with n=30.*

410 Within scenarios the intensity results show large differences across species-rich regions,
 411 establishing the need to consider land expansion jointly with land management when assessing
 412 biodiversity impacts of land-use change, and to provide these analyses for individual taxa of
 413 different status. For example, increases in irrigation water applied in locations rich in small-
 414 ranged amphibians are greater compared to locations rich in small-ranged birds or mammals
 415 (Figure 4). Without separating out taxa, such a finding could be overlooked, despite the
 416 probable greater importance of irrigation water withdrawal for amphibian populations. The
 417 intensity change results are heterogeneous between the different regions of species richness
 418 because food demand, the crops grown and yield response to agricultural inputs are location-

419 specific (see methods). We find, for example, nitrogen and irrigation application in bird-species-
420 rich regions increases over the period 2010–2100 (Figure 5a). Conversely, nitrogen and
421 irrigation application declines in threatened mammal- and amphibian-species-rich regions. In
422 the threatened mammal and amphibian locations by 2100 under BAU agricultural area expands
423 (Figure 3) and consequently agricultural production has increased sufficiently to meet demand
424 such that less nitrogen and water are required. In the LOW-AP scenarios, in the threatened
425 mammal and amphibian locations, reduced nitrogen and water use reduces with shrinking
426 agricultural area (Figure 3). Changing dietary demand may therefore have the greatest benefits
427 for threatened species through the reduction of both agricultural land area and agricultural
428 inputs in regions of high biodiversity.

429
430
431

432 3.3. Uncertainty and limitations

433 The stylized scenarios here assume high substitution rates of animal products, 95%, similar to
434 other studies that have assumed shifts towards complete vegetarianism (Stehfest et al., 2009;
435 Tilman and Clark, 2014a) or large reductions, e.g. Visconti et al., (2016) assumptions imply
436 reducing meat and egg consumption in all regions by 76–88%. Such scenarios are useful for
437 illustrating the effects of dietary transitions on land use changes, however, arguably such large
438 scale shifts will face barriers as dietary choices that individuals make are influenced by a
439 number of factors such as culture, price, availability, taste and convenience. Taking such
440 factors into account may reduce the potential for large scale dietary change. Lower rates of
441 animal product substitution would inevitable result in lower environmental benefits in this
442 study and others. E.g. for example, Stehfest et al., (2009) found a healthy diet that included some
443 level of animal product consumption resulted in greater land use and GHG emissions than
444 scenarios that reduced meat or animal product consumption entirely.

445 We explore land use change in regions with the greatest levels of biodiversity by include CI
446 hotspots and grid cells that are in the 90th percentile for species richness in our analysis. Our
447 regions of interest are therefore largely have a focused in the tropics. Using absolute species
448 richness loss has the advantage of highlighting particularly biodiverse regions at risk, with land
449 use change in these areas will therefore potentially have having a disproportionate effect on
450 global biodiversity loss. Similar to other studies we find these highly diverse regions, such as
451 sub-Saharan Africa and Latin America, are suitable for large scale agricultural expansion further
452 highlighting their importance in terms of conservation. However the choice to focus on areas
453 with the highest biodiversity inevitably means land use change and the consequences for
454 temperate or other climate zones are not well addressed in this study here. Changing dietary
455 patterns will undoubtedly affect such regions too for example reducing animal product
456 consumption reduced natural land loss in the south east of the United States, Kazakhstan and
457 Mongolia (Figure 2). However, while these regions do not necessary harbour high levels of
458 biodiversity, and are therefore not in our 90th percentile, they nevertheless may contain species
459 of significant conservation or cultural importance. Therefore while it was out with the scope of
460 our study to consider all geographic regions the consideration of temperate zones should not be
461 overlooked.

462 Land use changes to meet demand in PLUMv2 arise through a complex decision making process
463 that involves assessing spatially explicit crop yield responses, a variety of agricultural costs and
464 trade related costs. The parameter settings used in this study produced benchmarking results in
465 line with historical data (Alexander et al., 2018). However agricultural costs may change with

Commented [AP17]: Not sure I understand. Or the next sentence.

Commented [HR18]: REVIEWER COMMENT: One striking finding is that threatened species behave markedly different in Figure 4 and 5 – it would be worth investigating this more. Why is this the case and what are possible conservation implications here?

Commented [HR19]: REVIEWER COMMENT: Also, a more critical discussion of underlying uncertainties would be very useful. How robust are these results and how would the biodiversity implications change if the spatial patterns looked different?

REVIEWER COMMENT: There is more space for a discussion of the modelling as a whole within the broader context of the literature and particularly for some reflection on the limitations of the approach (especially the impacts of input choices on final results).

REVIEWER COMMENT: I suggest its publication, after minor revision. In particular I don't find clearly an estimation of the uncertainties linked to the models used for the analysis.

Commented [AP20]: Long and hard to follow sentence

Commented [HR21]: REVIEWER COMMENT: On the scenario description (section 2.3) it would be good to have more contextual information used in the justification of the settings chosen for the modelling. Were they taken from existing trends or entirely taken from the expert judgment of the modelling team. To what extent are the settings chosen plausible?

Commented [AP22]: Reading this paragraph makes it seem like only the tropics is included. That's not right either. Ok, so the tropics is over-represented, but don't think it's quite as extreme as implied overall in this text (perhaps I'm wrong through).

Commented [AP23]: Would it not be good to explicitly make the point here that we are considering changes in diets globally (including from these temperate regions) even those the focus is on the biodiversity in tropical regions.

Commented [HR24]: REVIEWER COMMENT: The choice of focusing on only areas in the top 10% of species richness numbers is not quite clear. This way, large (e.g. virtually the whole temperate zone) areas are excluded ex ante from further analysis. Obviously changing diets will also have a huge impact on them. It would be nice to see this also addressed in the study.

466 future economic development and policies. While we use a Monte-Carlo approach to
467 incorporate uncertainty of the input parameters, rates of agricultural expansion would reduce if
468 we increased the cost of agricultural expansion. Similarly if agricultural input costs, such as
469 fertiliser, were to rise the intensity of fertiliser use would decrease. Both outcomes would likely
470 have beneficial effects for biodiversity in the BAU scenario, thereby lessening the magnitude of
471 area and intensity savings with dietary change. The assumptions regarding future socio-
472 economic and climate condition as based on SSP2 and RCP 6.0 respectively. Further analysis
473 under a range of SSP trajectories may alter the land use patterns we find. For example changing
474 GDP, population size or political shifts, such as increasing protectionism under SSP3, would
475 alter the baseline food demand projections and change food supply requirements with
476 consequences for land use. SSPs that therefore project more demand than SSP2 used here,
477 particularly in developing tropical countries, will likely result in greater agricultural expansion
478 in biodiverse regions. Similarly alternative climate pathways may have consequences for
479 projected intensity use in biodiverse regions. Increased atmospheric CO₂ levels are linked to
480 higher yield potentials, reduced nitrogen losses and greater water use efficiency. In previous
481 work this leads to lower fertiliser and irrigation inputs in PLUMv2 (Alexander et al., 2018).
482 Therefore, while lower climate forcing's could be beneficial for climate change, they may have
483 unexpected negative effects such that more intensity inputs are required in agriculture to
484 achieve desired yields to meet demand. Global agricultural cropland projections could also
485 potentially shift as cropland area in regions negatively affected by climate change is reduced
486 while cropland area in regions with increased crop potential grows. Changing both the SSP and
487 RCP trajectory used in modelling studies of biodiversity may therefore alter the spatial patterns
488 of threats to biodiversity. Indeed recent modelling studies comparing SSPs and RCPs found that
489 stronger mitigation scenarios, corresponding to lower RCPs, had greater benefits for
490 biodiversity (Chaudhary and Mooers, 2018; Newbold et al., 2015).
491
492

493 *3.4. Conclusions and perspectives*

494
495 We find diets low in animal products reduce agricultural expansion and intensity in regions
496 with high biodiversity and the magnitude of change differed according to taxa, emphasising that
497 land-use change effects on biodiversity will be taxon specific. Numerous tropical countries with
498 high biodiversity have rates of increasing per capita meat production, and several are projected
499 to require up to 30% more agricultural land by 2050 (e.g. Ecuador, Brazil, and China)
500 (Machovina et al., 2015). Our results also demonstrate the importance of developing countries,
501 particularly those in the tropics, for biodiversity. The transition from low incomes to high
502 incomes and the associated increase in animal product consumption in developing countries
503 drives large losses of agricultural land across the tropics and in species-rich-regions under BAU.
504 In many developing countries, access to sufficient protein is limited and demand-side measures
505 such as global dietary interventions could be detrimental to the welfare of populations and thus
506 not ethical. Efforts to preserve biodiversity and ecosystem functioning will therefore require
507 scrutiny to ensure that changes are complementary to food security goals in developing
508 countries (including nutritional requirements) and respectful of cultural heritage. Land use
509 change in our BAU scenario is comparable to socio-economic conditions within the shared
510 socio-economic pathway (SSP) scenario SSP2 ('middle of the road', Popp et al., 2016). Likewise,
511 the scenarios of reduced meat consumption have been uniformly applied across countries.
512 Future scenarios of land-use change associated with alternative diets should encapsulate
513 aspects of fairness and equity (Tilman and Clark, 2014b). For example, a reduction of animal
514 product consumption in developed countries combined with the sustainable trade of meat into
515 countries with animal-protein deficits could simultaneously increase the health of industrialised
516 countries and prevent the destruction of natural land in tropical regions.
517

518 Dietary change, will be most effective if implemented as part of a suite of demand-side and
519 supply-side measures to reduce biodiversity loss (Tilman et al., 2017; Visconti et al., 2016). In a
520 modelling approach that combined increasing vegetarianism with reduction of food waste, by
521 2030, agricultural land decreased to a greater extent than we find here under LOW-AP
522 (Wirsenius et al., 2010). Policy screening scenarios similarly found that reaching any
523 biodiversity target will require a combination of strategies: for example, dietary change
524 combined with waste reduction and more efficient agricultural practices (Marchal et al., 2011;
525 Ten Brink et al., 2010). Reducing global meat consumption, and other demand-side measures
526 such as reducing food waste, will be socially and politically complex. It has been suggested that
527 large-scale dietary change will require incentives or regulations (Ripple et al., 2014b).
528 Furthermore, global diet alterations will need to complement food security goals and address
529 global food inequalities. However, biodiversity is an essential component of ecosystem
530 functioning, as well as human well-being, e.g. via provisioning of ecosystem services (IPBES,
531 2018; Naeem et al., 2016). Efforts to preserve biodiversity are, therefore, of the utmost
532 importance and may require dietary change.
533

534

535 4. References

- 536 Alexander, P., Rabin, S., Anthoni, P., Henry, R., Pugh, T., Rounsevell, M., Arneth, A., 2018.
537 Adaptation of global land use and management intensity to changes in climate and
538 atmospheric carbon dioxide. *Glob. Chang. Biol.*
- 539 Alexander, P., Rounsevell, M.D.A., Dislich, C., Dodson, J.R., Engström, K., Moran, D., 2015. Drivers
540 for global agricultural land use change: The nexus of diet, population, yield and bioenergy.
541 *Glob. Environ. Chang.* 35, 138–147.
542 doi:<http://dx.doi.org/10.1016/j.gloenvcha.2015.08.011>
- 543 Bajželj, B., Richards, K.S., Allwood, J.M., Smith, P., Dennis, J.S., Curmi, E., Gilligan, C.A., 2014.
544 Importance of food-demand management for climate mitigation. *Nat. Clim. Chang.* 4, 924–
545 929. doi:10.1038/nclimate2353
- 546 Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., Bustamante, M.,
547 Cinderby, S., Davidson, E., Dentener, F., Emmett, B., Erisman, J.-W., Fenn, M., Gilliam, F.,
548 Nordin, A., Pardo, L., De Vries, W., 2010. Global assessment of nitrogen deposition effects
549 on terrestrial plant diversity: a synthesis. *Ecol. Appl.* 20, 30–59. doi:10.1890/08-1140.1
- 550 Ceballos, G., Ehrlich, P.R., 2006. Global mammal distributions, biodiversity hotspots, and
551 conservation. *Proc. Natl. Acad. Sci.* 103, 19374–19379. doi:10.1073/pnas.0609334103
- 552 Chalaby, Y., Dutang, C., Savicky, P., Wuertz, D., 2015. Toolbox for Pseudo and Quasi Random
553 Number Generation and RNG Tests Version. Comprehensive R Archive Network (CRAN).
- 554 Chaudhary, A., Mooers, A., 2018. Terrestrial Vertebrate Biodiversity Loss under Future Global
555 Land Use Change Scenarios. *Sustainability* 10, 2764.
- 556 Cole, J.R., Mccoskey, S., 2013. Does global meat consumption follow an environmental Kuznets
557 curve? *Sustain. Sci. Pract. Policy* 9, 26–36.
- 558 De Frutos, A., Olea, P.P., Mateo-Tomás, P., 2015. Responses of medium- and large-sized bird
559 diversity to irrigation in dry cereal agroecosystems across spatial scales. *Agric. Ecosyst.*
560 *Environ.* 207, 141–152. doi:<https://doi.org/10.1016/j.agee.2015.04.009>
- 561 de Vries, W., Erisman, J.W., Spranger, T., Stevens, C.J., van den Berg, L., 2011. Nitrogen as a threat
562 to European terrestrial biodiversity, in: Sutton, M.A., Howard, C.M., Erisman, J.W., Billen, G.,
563 Bleeker, A., Grennfelt, P., Van Grinsven, H., Grizzetti, B. (Eds.), *The European Nitrogen*
564 *Assessment: Sources, Effects and Policy Perspectives*. Cambridge University Press, pp.

565 436–494.

566 Delgado, C., Rosegrant, M., Steinfeld, H., Ehui, S., Courbois, C., 2001. Livestock to 2020: the next
567 food revolution. *Outlook Agric.* 30, 27–29.

568 Dellink, R., Chateau, J., Lanzi, E., Magné, B., 2017. Long-term economic growth projections in the
569 Shared Socioeconomic Pathways. *Glob. Environ. Chang.* 42, 200–214.

570 Delzeit, R., Zabel, F., Meyer, C., Václavík, T., 2017. Addressing future trade-offs between
571 biodiversity and cropland expansion to improve food security. *Reg. Environ. Chang.* 17,
572 1443. doi:10.1007/s10113-016-0944-0

573 Dirzo, R., Young, H.S., Galetti, M., Ceballos, G., Isaac, N.J.B., Collen, B., 2014. Defaunation in the
574 Anthropocene. *Science* (80-.). 345, 401 LP-406.

575 Engström, K., Olin, S., Rounsevell, M.D.A., Brogaard, S., Van Vuuren, D.P., Alexander, P., Murray-
576 Rust, D., Arneth, A., 2016. Assessing uncertainties in global cropland futures using a
577 conditional probabilistic modelling framework. *Earth Syst. Dyn.* 7, 893–915.
578 doi:10.5194/esd-7-893-2016

579 Erb, K.-H., Lauk, C., Kastner, T., Mayer, A., Theurl, M.C., Haberl, H., 2016. Exploring the
580 biophysical option space for feeding the world without deforestation. *Nat. Commun.* 7.

581 FAOSTAT, 2015a. Commodity Balances/Crops Primary Equivalent (2015-12-16). Food and
582 Agriculture Organization of the United Nations, Rome, Italy.

583 FAOSTAT, 2015b. Commodity Balances/Livestock and Fish Primary Equivalent (2015-12-16).
584 Food and Agriculture Organization of the United Nations, Rome, Italy.

585 Flohre, A., Fischer, C., Aavik, T., Bengtsson, J., Berendse, F., Bommarco, R., Ceryngier, P., Clement,
586 L.W., Dennis, C., Eggers, S., Emmerson, M., Geiger, F., Guerrero, I., Hawro, V., Inchausti, P.,
587 Liira, J., Morales, M.B., Oñate, J.J., Pärt, T., Weisser, W.W., Winqvist, C., Thies, C., Tschardtke,
588 T., 2011. Agricultural intensification and biodiversity partitioning in European landscapes
589 comparing plants, carabids, and birds. *Ecol. Appl.* 21, 1772–1781. doi:10.1890/10-0645.1

590 Foley, J.A., DeFries, R., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin, F.S., Coe, M.T.,
591 Daily, G.C., Gibbs, H.K., 2005. Global consequences of land use. *Science* (80-.). 309, 570–
592 574.

593 Fuentes-Rodríguez, F., Juan, M., Gallego, I., Lusi, M., Fenoy, E., León, D., Peñalver, P., Toja, J., Casas,
594 J.J., 2013. Diversity in Mediterranean farm ponds: trade-offs and synergies between
595 irrigation modernisation and biodiversity conservation. *Freshw. Biol.* 58, 63–78.
596 doi:10.1111/fwb.12038

597 Gibbs, K.E., Mackey, R.L., Currie, D.J., 2009. Human land use, agriculture, pesticides and losses of
598 imperiled species. *Divers. Distrib.* 15, 242–253.

599 Gibson, L., Lee, T.M., Koh, L.P., Brook, B.W., Gardner, T.A., Barlow, J., Peres, C.A., Bradshaw, C.J.A.,
600 Laurance, W.F., Lovejoy, T.E., 2011. Primary forests are irreplaceable for sustaining
601 tropical biodiversity. *Nature* 478, 378.

602 Goldewijk, K.K., 2001. Estimating global land use change over the past 300 years: The HYDE
603 Database. *Global Biogeochem. Cycles* 15, 417–433. doi:10.1029/1999GB001232

604 Horner, G.J., Baker, P.J., Mac Nally, R., Cunningham, S.C., Thomson, J.R., Hamilton, F., 2009.
605 Mortality of developing floodplain forests subjected to a drying climate and water
606 extraction. *Glob. Chang. Biol.* 15, 2176–2186. doi:10.1111/j.1365-2486.2009.01915.x

607 IIASA, 2014. SSP Database (version 0.93). International Institute for Applied Systems Analysis,
608 Laxenburg, Austria, International Institute for Applied Systems Analysis, Laxenburg,

609 Austria.

610 IIASA/FAO, 2010. Global Agroecological Zones (GAEZ v3.0).

611 IPBES, 2018. Summary for policymakers of the regional assessment report on biodiversity and
612 ecosystem services for Europe and Central Asia of the Intergovernmental Science-Policy
613 Platform on Biodiversity and Ecosystem Services. IPBES secretariat, Bonn, Germany.

614 Jenkins, C.N., Pimm, S.L., Joppa, L.N., 2013. Global patterns of terrestrial vertebrate diversity and
615 conservation. *Proc. Natl. Acad. Sci.* 110, E2602–E2610. doi:10.1073/pnas.1302251110

616 Jones, B., O'Neill, B.C., 2016. Spatially explicit global population scenarios consistent with the
617 Shared Socioeconomic Pathways. *Environ. Res. Lett.* 11, 84003.

618 Kehoe, L., Kuemmerle, T., Meyer, C., Levers, C., Václavík, T., Kreft, H., 2015. Global patterns of
619 agricultural land-use intensity and vertebrate diversity. *Divers. Distrib.* 21, 1308–1318.
620 doi:10.1111/ddi.12359

621 Kehoe, L., Romero-Muñoz, A., Polaina, E., Estes, L., Kreft, H., Kuemmerle, T., 2017. Biodiversity at
622 risk under future cropland expansion and intensification. *Nat. Ecol. Evol.* 1, 1129–1135.
623 doi:10.1038/s41559-017-0234-3

624 Keyzer, M. a., Merbis, M.D., Pavel, I.F.P.W., van Wesenbeeck, C.F. a., 2005. Diet shifts towards
625 meat and the effects on cereal use: can we feed the animals in 2030? *Ecol. Econ.* 55, 187–
626 202. doi:10.1016/j.ecolecon.2004.12.002

627 Kleijn, D., Kohler, F., Baldi, A., Batary, P., Concepcion, E., Clough, Y., Diaz, M., Gabriel, D.,
628 Holzschuh, A., Knop, E., Kovacs, A., Marshall, E.J., Tschardtke, T., Verhulst, J., 2009. On the
629 relationship between farmland biodiversity and land-use intensity in Europe. *Proc. R. Soc.*
630 *B Biol. Sci.* 276, 903–909. doi:10.1098/rspb.2008.1509

631 Kok, M.T.J., Alkemade, R., Bakkenes, M., van Eerd, M., Janse, J., Mandryk, M., Kram, T., Lazarova,
632 T., Meijer, J., van Oorschot, M., Westhoek, H., van der Zagt, R., van der Berg, M., van der Esch,
633 S., Prins, A.-G., van Vuuren, D.P., 2018. Pathways for agriculture and forestry to contribute
634 to terrestrial biodiversity conservation: A global scenario-study. *Biol. Conserv.* 221, 137–
635 150. doi:https://doi.org/10.1016/j.biocon.2018.03.003

636 Laurance, W.F., Useche, D.C., Rendeiro, J., Kalka, M., Bradshaw, C.J.A., Sloan, S.P., Laurance, S.G.,
637 Campbell, M., Abernethy, K., Alvarez, P., 2012. Averting biodiversity collapse in tropical
638 forest protected areas. *Nature* 489, 290.

639 Lotze-Campen, H., Müller, C., Bondeau, A., Rost, S., Popp, A., Lucht, W., 2008. Global food demand,
640 productivity growth, and the scarcity of land and water resources: A spatially explicit
641 mathematical programming approach. *Agric. Econ.* 39, 325–338. doi:10.1111/j.1574-
642 0862.2008.00336.x

643 Machovina, B., Feeley, K.J., 2014. Taking a bite out of biodiversity. *Science* (80-.). 343, 838 LP-
644 838.

645 Machovina, B., Feeley, K.J., Ripple, W.J., 2015. Biodiversity conservation: The key is reducing
646 meat consumption. *Sci. Total Environ.* 536, 419–431. doi:10.1016/j.scitotenv.2015.07.022

647 Marchal, V., Dellink, R., Van Vuuren, D., Clapp, C., Chateau, J., Magné, B., van Vliet, J., 2011. OECD
648 environmental outlook to 2050. *Organ. Econ. Co-operation Dev.*

649 Martín-Queller, E., Moreno-Mateos, D., Pedrocchi, C., Cervantes, J., Martínez, G., 2010. Impacts of
650 intensive agricultural irrigation and livestock farming on a semi-arid Mediterranean
651 catchment. *Environ. Monit. Assess.* 167, 423–435. doi:10.1007/s10661-009-1061-z

652 Mittermeier, R., Robles Gil, P., Hoffmann, M., Pilgrim, J., Brooks, T., Goettsch Mittermeier, C.,

653 Lamoreux, J., Fonseca, G., 2004. Hotspots revisited. earth's biologically richest and most
654 endangered terrestrial ecoregions., Conservation International.

655 Myers, N., Mittermeier, R.A., Mittermeier, C.G., Da Fonseca, G.A.B., Kent, J., 2000. Biodiversity
656 hotspots for conservation priorities. *Nature* 403, 853.

657 Naeem, S., Chazdon, R., Duffy, J.E., Prager, C., Worm, B., 2016. Biodiversity and human well-
658 being: an essential link for sustainable development. *Proc. R. Soc. B* 283, 20162091.

659 Naylor, R., Steinfeld, H., Falcon, W., Galloway, J., Smil, V., Bradford, E., Alder, J., Mooney, H., 2005.
660 Losing the links between livestock and land. *Science* (80-.). 310, 1621–1622.

661 Nelson, G.C., Valin, H., Sands, R.D., Havlík, P., Ahammad, H., Deryng, D., Elliott, J., Fujimori, S.,
662 Hasegawa, T., Heyhoe, E., Kyle, P., Von Lampe, M., Lotze-Campen, H., Mason d’Croz, D., van
663 Meijl, H., van der Mensbrugge, D., Müller, C., Popp, A., Robertson, R., Robinson, S., Schmid,
664 E., Schmitz, C., Tabeau, A., Willenbockel, D., 2014. Climate change effects on agriculture:
665 economic responses to biophysical shocks. *Proc. Natl. Acad. Sci.* 111, 3274–9.
666 doi:10.1073/pnas.1222465110

667 Newbold, T., Hudson, L.N., Hill, S.L.L., Contu, S., Lysenko, I., Senior, R.A., Börger, L., Bennett, D.J.,
668 Choimes, A., Collen, B., Day, J., De Palma, A., Díaz, S., Echeverria-Londoño, S., Edgar, M.J.,
669 Feldman, A., Garon, M., Harrison, M.L.K., Alhousseini, T., Ingram, D.J., Itescu, Y., Kattge, J.,
670 Kemp, V., Kirkpatrick, L., Kleyer, M., Correia, D.L.P., Martin, C.D., Meiri, S., Novosolov, M.,
671 Pan, Y., Phillips, H.R.P., Purves, D.W., Robinson, A., Simpson, J., Tuck, S.L., Weiher, E., White,
672 H.J., Ewers, R.M., MacE, G.M., Scharlemann, J.P.W., Purvis, A., 2015. Global effects of land use
673 on local terrestrial biodiversity. *Nature* 520, 45–50. doi:10.1038/nature14324

674 Nijssen, M.E., WallisDeVries, M.F., Siepel, H., 2017. Pathways for the effects of increased nitrogen
675 deposition on fauna. *Biol. Conserv.* 212, 423–431.
676 doi:https://doi.org/10.1016/j.biocon.2017.02.022

677 O’Neill, B.C., Kriegler, E., Ebi, K.L., Kemp-Benedict, E., Riahi, K., Rothman, D.S., van Ruijven, B.J.,
678 van Vuuren, D.P., Birkmann, J., Kok, K., 2015. The roads ahead: narratives for shared
679 socioeconomic pathways describing world futures in the 21st century. *Glob. Environ.*
680 *Chang.*

681 O’Neill, B.C., Kriegler, E., Riahi, K., Ebi, K.L., Hallegatte, S., Carter, T.R., Mathur, R., van Vuuren,
682 D.P., 2014. A new scenario framework for climate change research: The concept of shared
683 socioeconomic pathways. *Clim. Change* 122, 387–400. doi:10.1007/s10584-013-0905-2

684 Orme, C.D.L., Davies, R.G., Burgess, M., Eigenbrod, F., Pickup, N., Olson, V.A., Webster, A.J., Ding,
685 T.-S., Rasmussen, P.C., Ridgely, R.S., 2005. Global hotspots of species richness are not
686 congruent with endemism or threat. *Nature* 436, 1016.

687 Pachauri, R.K., Allen, M.R., Barros, V.R., Broome, J., Cramer, W., Christ, R., Church, J.A., Clarke, L.,
688 Dahe, Q., Dasgupta, P., 2014. Climate change 2014: synthesis report. Contribution of
689 Working Groups I, II and III to the fifth assessment report of the Intergovernmental Panel
690 on Climate Change. IPCC.

691 Pereira, H.M., Navarro, L.M., Martins, I.S., 2012. Global biodiversity change: The bad, the good,
692 and the unknown. *Annu. Rev. Environ. Resour.* 37, 25–50. doi:10.1146/annurev-environ-
693 042911-093511

694 Pimm, S.L., Jenkins, C.N., Abell, R., Brooks, T.M., Gittleman, J.L., Joppa, L.N., Raven, P.H., Roberts,
695 C.M., Sexton, J.O., 2014. The biodiversity of species and their rates of extinction,
696 distribution, and protection. *Science* (80-.). 344. doi:10.1126/science.1246752

697 Pimm, S.L., Russell, G.J., Gittleman, J.L., Brooks, T.M., 1995. The future of biodiversity. *Science*
698 (80-.). 269, 347 LP-350.

699 Poore, J., Nemecek, T., 2018. Reducing food's environmental impacts through producers and
700 consumers. *Science* (80-.). 360, 987 LP-992.

701 Popp, A., Calvin, K., Fujimori, S., Havlik, P., Humpenöder, F., Stehfest, E., Bodirsky, B.L., Dietrich,
702 J.P., Doelmann, J.C., Gusti, M., Hasegawa, T., Kyle, P., Obersteiner, M., Tabeau, A., Takahashi,
703 K., Valin, H., Waldhoff, S., Weindl, I., Wise, M., Kriegler, E., Lotze-Campen, H., Fricko, O.,
704 Riahi, K., van Vuuren, D.P., 2016. Land-use futures in the shared socio-economic pathways.
705 *Glob. Environ. Chang.* 42, 331–345.
706 doi:<http://dx.doi.org/10.1016/j.gloenvcha.2016.10.002>

707 Popp, A., Lotze-Campen, H., Bodirsky, B., 2010. Food consumption, diet shifts and associated
708 non-CO2 greenhouse gases from agricultural production. *Glob. Environ. Chang.* 20, 451–
709 462. doi:10.1016/j.gloenvcha.2010.02.001

710 Possingham, H.P., Wilson, K.A., 2005. Biodiversity: Turning up the heat on hotspots. *Nature* 436,
711 919.

712 Pringle, R.M., 2017. Upgrading protected areas to conserve wild biodiversity. *Nature* 546, 91–99.
713 doi:10.1038/nature22902

714 Reich, P.B., 2009. Elevated CO2 reduces losses of plant diversity caused by nitrogen deposition.
715 *Science* (80-.). 326, 1399 LP-1402. doi:10.1126/science.1178820

716 Ren, G., Young, S.S., Wang, L., Wang, W., Long, Y., Wu, R., Li, J., Zhu, J., Yu, D.W., 2015.
717 Effectiveness of China's National Forest Protection Program and nature reserves. *Conserv.*
718 *Biol.* 29, 1368–1377. doi:10.1111/cobi.12561

719 Ripple, W.J., Estes, J.A., Beschta, R.L., Wilmers, C.C., Ritchie, E.G., Hebblewhite, M., Berger, J.,
720 Elmhagen, B., Letnic, M., Nelson, M.P., Schmitz, O.J., Smith, D.W., Wallach, A.D., Wirsing, A.J.,
721 2014a. Status and ecological effects of the world's largest carnivores. *Science* (80-.). 343.

722 Ripple, W.J., Smith, P., Haberl, H., Montzka, S.A., McAlpine, C., Boucher, D.H., 2014b. Ruminants,
723 climate change and climate policy. *Nat. Clim. Chang.* 4, 2–5. doi:10.1038/nclimate2081

724 Smith, B., Wärlind, D., Arneth, a., Hickler, T., Leadley, P., Siltberg, J., Zaehle, S., 2014. Implications
725 of incorporating N cycling and N limitations on primary production in an individual-based
726 dynamic vegetation model. *Biogeosciences* 11, 2027–2054. doi:10.5194/bg-11-2027-2014

727 Stehfest, E., Bouwman, L., Van Vuuren, D.P., Den Elzen, M.G.J., Eickhout, B., Kabat, P., 2009.
728 Climate benefits of changing diet. *Clim. Change* 95, 83–102.

729 Stevens, C.J., Dise, N.B., Mountford, J.O., Gowing, D.J., 2004. Impact of Nitrogen Deposition on the
730 Species Richness of Grasslands. *Science* (80-.). 303, 1876 LP-1879.
731 doi:10.1126/science.1094678

732 Stoll-Kleemann, S., Schmidt, U.J., 2017. Reducing meat consumption in developed and transition
733 countries to counter climate change and biodiversity loss: a review of influence factors.
734 *Reg. Environ. Chang.* 17, 1261–1277. doi:10.1007/s10113-016-1057-5

735 Swain, M., Blomqvist, L., McNamara, J., Ripple, W.J., 2018. Reducing the environmental impact of
736 global diets. *Sci. Total Environ.* 610, 1207–1209.

737 Taylor, K.E., Stouffer, R.J., Meehl, G.A., 2012. An overview of CMIP5 and the experiment design.
738 *Bull. Am. Meteorol. Soc.* 93, 485–498.

739 Ten Brink, B., van der Esch, S., Kram, T., Van Oorschot, M., Arets, E., 2010. Rethinking Global
740 Biodiversity Strategies: Exploring structural changes in production and consumption to
741 reduce biodiversity loss. Netherlands Environmental Assessment Agency.

742 Tilman, D., Balzer, C., Hill, J., Befort, B.L., 2011. Global food demand and the sustainable

743 intensification of agriculture. *Proc. Natl. Acad. Sci.* 108, 20260–4.
744 doi:10.1073/pnas.1116437108

745 Tilman, D., Clark, M., 2014a. Global diets link environmental sustainability and human health.
746 *Nature* 515, 518–522.

747 Tilman, D., Clark, M., 2014b. Global diets link environmental sustainability and human health.
748 *Nature* 515, 518–522.
749 doi:http://www.nature.com/nature/journal/v515/n7528/full/nature13959.html

750 Tilman, D., Clark, M., Williams, D.R., Kimmel, K., Polasky, S., Packer, C., 2017. Future threats to
751 biodiversity and pathways to their prevention. *Nature* 546, 73–81.
752 doi:10.1038/nature22900

753 Tilman, D., Fargione, J., Wolff, B., D'Antonio, C., Dobson, A., Howarth, R., Schindler, D.,
754 Schlesinger, W.H., Simberloff, D., Swackhamer, D., 2001. Forecasting agriculturally driven
755 global environmental change. *Science* (80-.). 292, 281 LP-284.

756 UNEP-WCMC., I.&, 2016. The World Database on Protected Areas (WDPA). [WWW Document].

757 van Vuuren, D.P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G.C., Kram,
758 T., Krey, V., Lamarque, J.F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S.J., Rose, S.K.,
759 2011. The representative concentration pathways: An overview. *Clim. Change* 109, 5–31.
760 doi:10.1007/s10584-011-0148-z

761 Visconti, P., Bakkenes, M., Baisero, D., Brooks, T., Butchart, S.H.M., Joppa, L., Alkemade, R., Di
762 Marco, M., Santini, L., Hoffmann, M., Maiorano, L., Pressey, R.L., Arponen, A., Boitani, L.,
763 Reside, A.E., van Vuuren, D.P., Rondinini, C., 2016. Projecting global biodiversity indicators
764 under future development scenarios. *Conserv. Lett.* 9, 5–13. doi:10.1111/conl.12159

765 Visconti, P., Bakkenes, M., Smith, R.J., Joppa, L., Sykes, R.E., 2015. Socio-economic and ecological
766 impacts of global protected area expansion plans. *Philos. Trans. R. Soc. B Biol. Sci.* 370,
767 20140284. doi:10.1098/rstb.2014.0284

768 Wellesley, L., Happer, C., Froggatt, A., 2015. Changing climate, changing diets: pathways to lower
769 meat consumption. Royal Institute of International Affairs, Chatham House.

770 Westhoek, H., Rood, T., van den Berg, M., Janse, J., Nijdam, D., Reudink, M., Stehfest, E., Lesschen,
771 J.P., Oenema, O., Woltjer, G.B., 2011. The protein puzzle: the consumption and production of
772 meat, dairy and fish in the European Union. Netherlands Environmental Assessment
773 Agency.

774 Wirsenius, S., Azar, C., Berndes, G., 2010. How much land is needed for global food production
775 under scenarios of dietary changes and livestock productivity increases in 2030? *Agric.*
776 *Syst.* 103, 621–638.

777 Yamaguchi, T., Blumwald, E., 2005. Developing salt-tolerant crop plants: challenges and
778 opportunities. *Trends Plant Sci.* 10, 615–620.

779

780

781

782

783

784