

8-1981

# The Role of Metals in Carcinogenesis: Biochemistry and Metabolism

K Wetterhahn Jennette  
*Dartmouth College*

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>

 Part of the [Chemistry Commons](#)

---

## Recommended Citation

Jennette, K Wetterhahn, "The Role of Metals in Carcinogenesis: Biochemistry and Metabolism" (1981). *Open Dartmouth: Faculty Open Access Articles*. 3563.  
<https://digitalcommons.dartmouth.edu/facoa/3563>

This Article is brought to you for free and open access by Dartmouth Digital Commons. It has been accepted for inclusion in Open Dartmouth: Faculty Open Access Articles by an authorized administrator of Dartmouth Digital Commons. For more information, please contact [dartmouthdigitalcommons@groups.dartmouth.edu](mailto:dartmouthdigitalcommons@groups.dartmouth.edu).

# The Role of Metals in Carcinogenesis: Biochemistry and Metabolism

by K. Wetterhahn Jennette\*

The oxyanions of vanadium, chromium, molybdenum, arsenic, and selenium are stable forms of these elements in high oxidation states which cross cell membranes using the normal phosphate and/or sulfate transport systems of the cell. Once inside the cell, these oxyanions may act as phosphate and/or sulfate analogs, inhibiting enzymes involved in catalyzing phosphoryl or sulfuryl transfer reactions. Often the oxyanions serve as alternate enzyme substrates but form ester products which are hydrolytically unstable compared with the sulfate and phosphate esters and, therefore, decompose readily in aqueous solution. Arsenite and selenite are capable of reacting with sulfhydryl groups in proteins. Some cells are able to metabolize redox active oxyanions to forms of the elements in other stable oxidation states. Specific enzymes may be involved in the metabolic processes. The metabolites of these elements may form complexes with small molecules, proteins and nucleic acids which inhibit their ability to function properly.

The divalent ions of beryllium, manganese, cobalt, nickel, cadmium, mercury, and lead are stable forms of these elements which may mimic essential divalent ions such as magnesium, calcium, iron, copper, or zinc. These ions may complex small molecules, enzymes, and nucleic acids in such a way that the normal activity of these species is altered. Free radicals may be produced in the presence of these metal ions which damage critical cellular molecules.

The biochemistry and metabolism of inorganic species involved in carcinogenesis encompasses transport of inorganic species across the cell membrane, enzymatic and chemical transformation of redox active inorganic species within the cell, coordination of inorganic species to cellular small molecules and macromolecules, and inhibition, activation or change of specificity of cellular enzymes by inorganic species. Inorganic species may mimic forms of essential elements in these processes. The kinetic and thermodynamic properties of inorganic species may be substantially different from those of essential elements and, therefore, may ultimately determine the manifestation of cellular damage, including carcinogenic and cocarcinogenic effects. For example, in order for an inorganic compound to be a mutagen it must enter the cell; however, an inorganic compound may be able to act as a cocarcinogen without penetration into the cell by interacting with the cell surface. Once inside the cell the inorganic species originally transported

across the cell membrane may be transformed into a different form by reaction with cellular components. The transformation may result in a complex which is easily excreted, hence deactivated. Alternatively, the element may be activated and bind to enzymes and/or nucleic acids, altering their ability to function properly in the cell. The ultimate disposition of the element will depend on the chemical properties of the individual inorganic species involved.

## Oxide Complexes

The early transition elements have a tendency to exist in high oxidation states as oxy or hydroxy complexes in aqueous solution (1). The nature of the oxy species depends on concentration and pH. For example, in neutral aqueous solution at concentrations less than  $10^{-4}M$ , vanadium (V) exists mainly as  $H_2VO_4^-$ ; chromium (VI) exists as chromate,  $CrO_4^{2-}$ ; manganese (VII) exists as permanganate,  $MnO_4^-$ ; molybdenum (VI) exists as molybdate,  $MoO_4^{2-}$ . As the concentration of these species is increased, the mononuclear ions aggregate to form polyanions.

\* Department of Chemistry, Dartmouth College, Hanover, New Hampshire 03755.

In addition to the early transition elements, certain main group elements also tend to exist as oxyanions (1). At pH 7, phosphorus (V) exists as an equilibrium mixture of  $\text{H}_2\text{PO}_4^-$  and  $\text{HPO}_4^{2-}$ ; arsenic (V) exists as an equilibrium mixture of  $\text{H}_2\text{AsO}_4^-$  and  $\text{HAsO}_4^{2-}$ ; sulfur (VI) exists as sulfate,  $\text{SO}_4^{2-}$ ; and selenium (VI) as selenate,  $\text{SeO}_4^{2-}$ . All of these oxyanions have a tetrahedral geometry (Fig. 1).

Other stable oxy-complexes of several of these elements also exist (1). Vanadium (IV) complexes are generally five- or six-coordinate and typically contain the vanadyl ion,  $\text{VO}^{2+}$ . Pyramidal structures are found for arsenic (III) which exists in neutral aqueous solution mainly as protonated arsenite,  $\text{As}(\text{OH})_3$ , sulfur (IV) which exists as sulfite,  $\text{SO}_3^{2-}$ , and selenium (IV) which exists as selenite,  $\text{SeO}_3^{2-}$ .

## Transport

Since phosphate and sulfate are known to cross cell membranes by passive transport and mediated permeation (2), the other oxyanions may use these transport systems to enter the cells. Arsenate was found to enter yeast cells by the phosphate transport system (3). Arsenate was a competitive inhibitor of

phosphate uptake into 3T3 mouse fibroblast cells, however, chromate and vanadate showed very little inhibition of the phosphate transport system (4). Vanadate crossed the red blood cell membrane by the anion-exchange system responsible for phosphate transport (5). Vanadate uptake was inhibited by dinitrostilbene disulfonate, a known inhibitor of anion exchange (5). Vanadate inhibited phosphate transport into red cells (5).

Chromate was able to cross the membranes and enter red blood cells (6), Ehrlich ascites carcinoma cells (7), mastocytoma cells (8) and EL4 ascites leukemia lymphoblasts (9). The velocity of chromate uptake by human leukocytes was slightly inhibited by sulfate, tungstate and molybdate; metavanadate was a competitive inhibitor of chromate transport; arsenite and *p*-chloromercuribenzenesulfonate were potent inhibitors of chromate influx (10). Transport of molybdate across rat intestinal membranes was inhibited by sulfate, selenate and chromate, but not phosphate (11). Sulfate was a competitive inhibitor of molybdate transport in everted sacs of rat and ovine small intestines (12-24). Tungstate, selenate and thiosulfate also inhibited sulfate transport by rat ileum *in vitro* (12, 13). Sulfate inhibited the absorption of molybdate




FIGURE 1. Oxyanions and oxo-complexes stable in dilute neutral aqueous solution.

in isolated chick intestinal loop segments (15). Active sulfate uptake in the yeast *Saccharomyces cerevisiae* was inhibited by chromate, molybdate, selenate, thiosulfate, sulfite, and tellurite (16). The activity of the sulfate transport system in the filamentous fungus *Penicillium notatum* was strongly inhibited by selenate, molybdate, thiosulfate, chromate, and sulfite (17). An active transport system for molybdate was characterized for the molybdenum-requiring nitrogen-fixing bacterium *Clostridium pasteurianum* (18). Tungstate and sulfate were competitive inhibitors of molybdate uptake in these organisms, however, thiosulfate and vanadate were not inhibitors of molybdate uptake (18). The transport of sulfate into the bacterium *Salmonella typhimurium* was strongly inhibited by sulfite, chromate, thiosulfate and selenate and was moderately inhibited by molybdate, vanadate, and tungstate (19).

In summary, oxyanions of chromium, vanadium, molybdenum, tungsten, arsenic, and selenium, readily permeated the cell membranes of prokaryotes and eukaryotes. The oxyanions entered the cells by using the normal active transport systems for phosphate and sulfate.

## Enzyme Interaction

The oxyanions have been shown to affect the activities of phosphotransferases and phosphohydrolases. Human liver and wheat germ acid phosphatases were competitively inhibited by arsenate, vanadate, molybdate, and tungstate (20). Arsenate (21), vanadate, and vanadyl ion,  $\text{VO}^{2+}$  (22), were potent competitive inhibitors of *E. coli* alkaline phosphatase; permanganate was an irreversible inhibitor, whereas chromate and perchlorate were not inhibitors (23). Molybdate was a potent inhibitor of phosphoprotein phosphatase isolated from bovine tracheal smooth muscle extract (24). Molybdate was a noncompetitive inhibitor of NADP<sup>+</sup> 2'-nucleotidase from *Hevea brasiliensis* (25). Oxovanadium (IV) ion, and complexes of uridine





FIGURE 2. Proposed trigonal bipyramidal transition state in RNA hydrolysis by ribonuclease A (26).

August 1981



with oxovanadium (IV) and vanadate were competitive inhibitors of ribonuclease A (26). The competitive inhibition of enzyme activity was proposed to be based on the ability of these ions to form trigonal bipyramidal complexes of the type shown in Figure 2, which act as transition-state analogs in reactions involving phosphate esters (20, 26). The irreversible inhibition of alkaline phosphatase activity by permanganate appeared to be due to oxidation of the enzyme (16, 20).

Vanadate was found to be a potent inhibitor of  $(\text{Na}^+, \text{K}^+)\text{ATPases}$  from dog kidney (27, 28), pig kidney (29), dog brain (28), pig brain (30), eel electroplax (28), red blood cells (5), and squid axons (31), dynein ATPases from sea urchin sperm (30, 32) and *Tetrahymena pyriformis* (32), myosin ATPases from human platelets (32), and rabbit muscle (32, 33), and  $(\text{Ca})\text{ATPases}$  from red blood cells (32) and squid axons (35). Vanadate did not inhibit rabbit muscle sarcoplasmic reticulum  $(\text{Ca})\text{ATPase}$  or beef heart mitochondrial  $\text{F}_1\text{ATPase}$  (28).

Vanadate formed a stable stoichiometric ternary complex with ADP and myosin ATPase [Eq. (1)] and it was proposed that the vanadate was coordinated to nucleophiles in the active site of the enzyme (33). Vanadate interacted strongly with the low affinity ATP site of dog kidney  $(\text{Na}^+, \text{K}^+)\text{ATPase}$  [Eq. (2)] and increased the affinity of the enzyme for  $\text{Mg}^{2+}$  and  $\text{K}^+$  (36). In the presence of potassium, vanadate blocked the conformational change of  $(\text{Na}^+, \text{K}^+)\text{ATPase}$  ( $\text{E}_2\text{K} \rightarrow \text{E}_1\text{K}$ ) which normally follows the hydrolysis of the phosphoenzyme (37). It was suggested that inhibition of  $(\text{Na}^+, \text{K}^+)\text{ATPase}$  activity was due to the formation of a stable ternary complex of enzyme, vanadate, magnesium, and potassium (37). It was proposed that vanadate's ability to act as a phosphate transition-state analog (Fig. 2) was responsible for the high affinity of vanadate for the phosphatase site on the enzyme. The affinity of phosphate and arsenate for this site was approximately  $10^5$ - $10^6$  times lower than that of vanadate. Arsenate also inhibited dog kidney  $(\text{Na}^+, \text{K}^+)\text{ATPase}$  by binding to the weak ATP site (36).



E = myosin ATPase



Vanadate markedly enhanced adenylate cyclase activity in ventricular muscle from rat, rabbit,

guinea pig and cat (38) and in rat fat cells (39), and inhibited stimulation of sodium and water transport by cyclic AMP in frog skin (40). Since vanadate has no effect on purified cyclic AMP phosphodiesterase the mechanism for these effects was proposed to be the inhibition of ATPase or GTPase by vanadate (38-40). Molybdate was a reversible activator of adenylate cyclase from rat liver plasma membranes, rat erythrocyte ghosts and rat cardiac, kidney, and brain homogenates (41). The molybdate did not activate the soluble rat testicular adenylate cyclase, and therefore it was suggested that molybdate indirectly activated the enzyme by inhibiting GTPase activity. Chromate and tungstate inhibited rat liver adenylate cyclase activity at concentrations where molybdate markedly stimulated activity (41).

Arsenate prevented the alkaline phosphatase-mediated inactivation of glucocorticoid binding capacity of unbound soluble receptors from L929 mouse fibroblasts (42). Molybdate inhibited the alkaline phosphatase activity of rat thymus pellets and prevented the reduction in glucocorticoid binding capacity of unoccupied rat liver receptors caused by thymus pellets (43). Molybdate also inhibited the inactivation of unbound soluble glucocorticoid receptor from L929 mouse fibroblasts (43), whereas tungstate had no effect on receptor inactivation (44). The transformation of glucocorticoid bound receptor from mouse fibroblasts into the form which binds DNA was blocked by molybdate and tungstate (44). Chick oviduct cytosol phosphatase activity was inhibited by arsenate, chromate, molybdate, tungstate, and vanadate (45). Molybdate, tungstate, and vanadate (but not chromate or arsenate) were potent inhibitors of the activation of progesterone receptor from chick oviduct to the state which binds DNA (45). It was proposed that molybdate interacted directly with the steroid receptors, possibly by forming a phosphate complex with the phosphorylated receptor and thereby preventing its binding to DNA (44, 45).

Vanadate and arsenate markedly stimulated phosphate transfer to glucose by phosphoglucomutase, whereas tungstate, molybdate, and niobate had no effect (46). The half-maximal rate of phosphate transfer occurred at pH 9.8 and was shifted to 9.5 in the presence of arsenate and to 7.7 in the presence of vanadate. It was concluded that arsenate and vanadate bind to phosphoglucomutase and lower the  $pK_a$  of the active site tyrosine hydroxyl group.



Arsenate, sulfate, molybdate, and tungstate were competitive inhibitors of 6-phosphogluconate dehydrogenase with respect to the substrate 6-phosphogluconate and were noncompetitive inhibitors with respect to the  $NADP^+$  cofactor (47). Permanganate, periodate, perchlorate, and chromate irreversibly

inactivated 6-phosphogluconate dehydrogenase, and in the case of permanganate the inactivation was ascribed to the oxidation of cysteine to cysteic acid (47). Ferrate inactivated rabbit muscle phosphorolase b and prevented the binding of 5'-AMP to the enzyme (48). The action of ferrate was ascribed to the oxidation of tyrosine residues at the phosphate binding site. Both arsenate and vanadate replaced phosphate as a substrate in the glyceraldehyde-3-phosphate dehydrogenase reaction (49). Vanadate and arsenate formed analogs of 1,3-diphosphoglyceric acid which were labile in aqueous solution. The normal pathway of formation of ATP from 3-phosphoglyceraldehyde and phosphate through the stable 1,3-diphosphoglycerate intermediate is shown in Eq. (3), where GAPDH denotes glyceraldehyde-3-phosphate dehydrogenase. Substitution of vanadate or arsenate for phosphate [Eq. (4)] results in the formation of unstable acyl vanadate or acyl arsenate intermediates which are quickly hydrolyzed in aqueous solutions (49). Hydrolytic decomposition of the acyl vanadate or acyl arsenate drove the reaction to completion. Arsenate was found to spontaneously form an ester-like product with dihydroxyacetone (50). This arsenate analog of dihydroxyacetone phosphate was recognized as a substrate by glycerol 3-phosphate dehydrogenase. Simple arsenate and vanadate trialkyl esters were found to be very labile compared to phosphate esters and appeared to undergo rapid exchange of alcohol via a five-coordinate transition state (51).

Vanadium (V) in the form of the decavanadate polyanion was a potent inhibitor of adenylate kinase from rat liver mitochondria and rabbit skeletal muscle (49). Sheep heart phosphofructokinase was inhibited by vanadate in a manner similar to ATP inhibition (52).

Molybdate, selenate, sulfate, sulfite, and arsenite were competitive inhibitors of chicken liver mitochondrial pyruvate carboxylase with respect to S-acetylCoA and decreased the rate of inactivation of the enzyme upon incubation at 2°C (53). Arsenate inhibition of pyruvate carboxylase was noncompetitive with respect to S-acetylCoA and it was concluded that unlike the other oxyanions arsenate acted at the nucleotide site rather than the S-acetylCoA site (53).

In addition to acting as phosphate analogs, the oxyanions have also been shown to act as sulfate analogs and, therefore, interfere with sulfur metabolism. The enzyme ATP-sulfurylase catalyzes the formation of adenosine-5'-phosphosulfate (APS) from ATP and sulfate [Eqs. (5)-(8)], the first step in sulfate reduction and in formation of the "active sulfate" donor, adenosine-3'-phosphate-5'-phosphosulfate (PAPS). Sulfate is activated by formation of






adenyl sulfate (APS), which acts as a sulfuryl donor in subsequent enzymatic reactions such as sulfation of polysaccharides [Eqs. (5)-(7)]. Selenate, molybdate, chromate, and tungstate form unstable AMP-anhydrides which are hydrolyzed [Eq. (8)] to form AMP and regenerate the oxyanions (54). Molybdate was an irreversible inhibitor of *Desulfovibrio* ATP-sulfurylase (11). Sulfite, chromate, molybdate, and tungstate caused irreversible cleavage of ATP to AMP by baker's yeast ATP-sulfurylase (54). Molybdate, chromate, tungstate, and selenate inhibited the formation of APS by ATP-sulfurylase purified from *Nitrobacter agilis* (55). Molybdate and selenate inhibited ATP-sulfurylase activity by competing with sulfate for the active site on ATP-sulfurylases purified from *Penicillium chrysogenum* (56) and spinach leaves (57). Molybdate caused the formation of AMP from ATP, however, no AMP was formed with selenate as substrate for the spinach leaf ATP-sulfurylase (57). Selenate was found to act as a substrate for baker's yeast ATP-sulfurylase and form adenosine-5'-phosphoselenate (APSe) which was stable enough to isolate and characterize (54). The APSe was easily hydrolyzed, however, resulting in the formation of AMP and selenate. In contrast, APSe and AMP were not detected when selenate was used as a substrate for purified ATP-sulfurylases from *Astragalus bisulcatus* (a selenium-accumulator species), *Astragalus hamosus* (a nonaccumulator species), and spinach (57, 58). It was proposed that the mechanism of inhibition by these anions [Eqs. (5)-(8)] involved the formation of unstable adenosine-5'-phosphoanion anhydrides which quickly hydrolyzed to form AMP and the anion (54). Kinetic studies on purified ATP-sulfurylases from *Saccharomyces cerevisiae* (59), *Penicillium chrysogenum*

(56), and Furth mouse mastocytoma (60) suggested a sequential mechanism for the enzyme, where both molybdate (or sulfate) and MgATP bind to the enzyme before the products, pyrophosphate and AMP-MoO<sub>4</sub> (or APS), are released.

Molybdate was a reversible competitive inhibitor of sulfate transport in the unicellular red alga, *Porphyridium aeruginosum* (61). Molybdate entered these cells and inhibited the intracellular utilization of the sulfate pool in the synthesis of capsular polysaccharides. Sulfation levels of cell-associated sulfated polysaccharide and the secreted extracellular sulfated polysaccharide were depressed by molybdate (61). Molybdate and selenate at concentrations without effect on growth inhibited the incorporation of sulfate into cellular constituents of *Escherichia coli* (62). Molybdate inhibited sulfate reduction to sulfide by rumen microorganisms without affecting sulfite reduction by these bacteria (63). Molybdate was not toxic in a mutant of *Salmonella typhimurium* lacking ATP-sulfurylase (64). Molybdate and selenate inhibited the synthesis of APS in extracts from tobacco XD cells (65). Molybdate derepressed ATP-sulfurylase in tobacco cells by essentially causing sulfur starvation through inhibition of sulfate uptake and sulfate activation by ATP-sulfurylase. Selenate derepression of ATP-sulfurylase did not occur by a mechanism involving sulfur starvation and it was proposed that derepression was the result of the formation of a seleno-analog which was an antagonist to some product in the sulfate pathway (65).



Arsenite has been shown to inhibit thiol-containing enzymes (66, 67), especially those containing two sulfhydryl groups in close proximity, e.g., chicken and rat liver citrate cleavage enzyme and the lipoic acid-containing pyruvate oxidase system (68). Other



enzymes, such as aldehyde dehydrogenases (69) and glutamine synthetase (70), were inhibited by arsenite only in the presence of exogenous mercaptan. Enzyme activity could be restored by addition of dithiols and in some cases by addition of excess monothiols such as glutathione and cysteine. An antidote used for arsenic poisoning, 2,3-dimercaptopropanol, British antilewisite (BAL), has been shown to react with arsenite through the two sulfhydryl groups [Eqs. (9) and (10)], forming a stable five-membered ring structure (71). The level of mucosal glutathione was immediately depressed after oral administration of arsenic trioxide ( $\text{As}_2\text{O}_3$ ) to rats (72). This effect was attributed to the binding of arsenite to glutathione. Glutathione levels in mucosal cells subsequently rose to twice the normal concentration in response to its depletion by arsenic (72). Arsenite inhibited the activi-

ties of rabbit liver aldehyde oxidase, milk xanthine oxidase and chicken liver xanthine dehydrogenase by binding at the molybdenum active sites in these molybdoenzymes (73). Arsenite was found to reversibly inhibit acetylcholinesterase by a mechanism not involving disulfide or sulfhydryl functions although kinetic data suggested a covalent interaction between the arsenite and enzyme (74).

Selenites have been shown to react with sulfhydryl groups in cysteine, glutathione, dihydrolipoic acid, 2-mercaptoethanol, coenzyme A and in proteins (75-77). Selenite, acting as an oxidant, inhibited enzyme interactions requiring a free sulfhydryl group. Conversion of thiols to disulfides by selenite was shown to involve a selenotrisulfide intermediate [Eqs. (11) and (12)], which could be isolated from the reaction mixture. A selenotrisulfide derivative of glutathione was enzymatically reduced to the



persulfide analog by glutathione reductase (78). The selenopersulfide produced [Eq. (12)] decomposed to glutathione and selenium(0). Selenite reacted with reduced pancreatic ribonuclease A forming a slightly unfolded structure containing intramolecular selenotrisulfide linkages which was devoid of activity (79). Selenite was able to release methylmercury bound to sulfhydryl groups in blood proteins such as albumin in the presence of red blood cells, and liver, kidney and brain homogenates from rats and mice (80). Selenite was effective in removing bound methylmercury from human blood, whereas selenate was ineffective. *In vivo* administration of selenite to mice previously exposed to methylmercury resulted in the release of free methylmercury from blood, liver and kidney tissue, but not from brain tissue (80).

In the presence of glutathione selenite stimulated the swelling of liver mitochondria isolated from selenium-deficient rats (81). Selenate, arsenite, arsenate, sulfite, thiosulfate, molybdate, tellurate, and vanadate had little or no effect on mitochondrial swelling in this system whereas tellurite was slightly active. The swelling effect of selenium was attributed to the ability of selenite to catalyze the reduction of cytochrome c by glutathione (82). Selenocysteine also catalyzed glutathione reduction of cytochrome c, selenate had only a slight effect and selenomethionine was ineffective. It was proposed that the mechanism for selenite catalysis of cytochrome c reduction by glutathione involved the formation of a selenopersulfide intermediate (82). In rat erythrocytes, selenite stimulated reduction of methemoglobin in the presence of glutathione at concentrations which had no effect on NADH-

methemoglobin reductase activity (83). Several proteins involved in electron transport processes, e.g., formate dehydrogenase, glycine reductase, and glutathione peroxidase, have been identified as selenoproteins (84).



In summary, oxyanions of vanadium, chromium, molybdenum, tungsten, arsenic, and selenium inhibited the activity of enzymes involved in phosphate and sulfate metabolism. Selenite and arsenite inhibited processes requiring free sulfhydryl groups. Selenite affected electron transport reactions. Cellular regulatory processes may be affected by these oxyanions through their disturbance of oxidative phosphorylation, inhibition of oxidases and ATPases, and interference with steroid effects on levels of cAMP, e.g., activation of adenylate cyclase activity, and interference with steroid-receptor-DNA interactions.

## Metabolism

The oxyanions, except phosphate, are redox active and have other forms in different oxidation states which are stable in aqueous solution (1). Metabolic processes may be able to convert the high oxidation state metals in oxyanions to stable complexes with the metal in a lower oxidation state.

Liver and kidneys from rats treated with vanadate exhibited an electron paramagnetic resonance (EPR) signal characteristic of vanadium (IV) indicating these tissues were able to reduce vanadate by one electron (85). Lungs contained only small amounts of vanadium (IV) and heart tissue had no detectable V (IV). Toxic levels of vanadate had no





effect on the activities of two molybdenum enzymes, sulfite oxidase and xanthine oxidase. It was suggested that the toxicity of vanadium was related to the one electron reduction of V (V) and subsequent binding of the V (IV) to protein. Intact red blood cells converted vanadate to a form which bound to hemoglobin (5). Red cell ghosts and pure hemoglobin were inactive in vanadate conversion. The vanadium associated with hemoglobin had the EPR spectrum characteristic of vanadyl ion indicating that reduction of vanadate to vanadium (IV) had occurred (86). In contrast to vanadate which markedly inhibited ( $\text{Na}^+$ ,  $\text{K}^+$ ) ATPase, vanadium (IV) was a much less effective inhibitor (86). Rapid oxidation of NADH occurred in the presence of vanadate and purified calf cardiac cell membranes containing ( $\text{Na}^+$ ,  $\text{K}^+$ ) ATPase (87). This effect was attributed to the presence of a heat labile NADH-vanadate reductase activity in cardiac membranes. Catechol, which reduced vanadium (V) to vanadium (IV) and formed a complex with vanadium (IV) reversed the vanadium (V) inhibition of reactivated sea urchin sperm motility, dynein ATPase and ( $\text{Na}^+$ ,  $\text{K}^+$ ) ATPase (30). Norepinephrine reversed vanadate inhibition of dog kidney ( $\text{Na}^+$ ,  $\text{K}^+$ ) ATPase by reduction of vanadium (V) and complexation of the vanadium (IV) formed (88).

Human breast, liver, and thyroid tissues which had been treated with dichromate or with the Cr (VI) containing tissue preservative, Zenker's solution, showed an EPR signal characteristic of chromium (III) (89). The intracellular chromium in chromate treated tumor cells was found as chromium (III) complexed with organelles (nuclei, mitochondria and microsomes), soluble proteins and small molecules (7-9). The intracellular chromium (III) complexes, even in the form of low

molecular weight complexes, were retained in the cell indicating the cell membrane was relatively impermeable to trivalent chromium. Only chromium (III) was detected in BHK21 hamster fibroblasts that had been treated with dichromate (90).

Treatment of red blood cells with chromate resulted in uptake of chromium and recovery of chromium bound to the globin (nonheme) portion of hemoglobin (6). In contrast to the inactivity of purified hemoglobin in vanadate reduction (5) hemoglobin was capable of reducing chromate and binding chromium (III) (6, 91). With both red blood cell-labeled hemoglobin and isolated hemoglobin  $\alpha$ ,  $\beta$ , and  $\gamma$  polypeptides, chromium (III) was found preferentially to bind to the  $\beta$  chains (91). Chromium (III) was not taken up by red blood cells suggesting that the erythrocyte membrane was also impermeable to Cr (III) complexes (6). Chromate inhibited glutathione reductase activity of erythrocytes, whereas chromium (III) had no effect (92). Chromate had no effect on acetylcholinesterase, and glyoxalase activities, and on activities of enzymes of the glycolytic and pentose-phosphate pathways of erythrocytes (92).

Chromium (VI) inhibited benzpyrene (BP) hydroxylase activity in rat lung homogenates (93). Chromium (III), arsenite and selenite had no effect on BP hydroxylase *in vitro* (93). Potassium chromate inhibited hydroxylation of BP by mouse liver homogenates, however, the same study showed that potassium dichromate enhanced BP hydroxylation at low concentrations and inhibited at high concentrations (94). Chromium (III) inhibited BP hydroxylase activity in rat liver microsomes (95).

The microsomal fraction of rat liver cells was found to be capable of metabolizing chromate to

chromium (III) using NADPH or NADH as cofactor (96, 97). The enzymes composing the electron-transport cytochrome P-450 system were shown (98) to be involved in the microsomal reduction of chromate. The NADPH-dependent chromate reductase activity of microsomes was inhibited by carbon monoxide and metyrapone, known inhibitors of cytochrome P-450 activity (98). The ability of cytochrome P-450 to function as a NADH- or NADPH-dependent reductase, rather than a monooxygenase, has been well documented for a number of organic substrates (99). Some of the chromium (III) produced by microsomal reduction remained bound to microsomal protein and the majority was complexed to the  $\text{NAD}^+$  or  $\text{NADP}^+$  cofactor (97). Since chromate is not a strong oxidizing agent at physiological pH, no interaction between the negatively charged chromate ion and the negatively charged DNA polymer was observed in aqueous pH 7.4 solutions *in vitro* (97, 100). However, the presence of a microsomal metabolizing system capable of reducing chromate to chromium (III) caused significant amounts of chromium to bind to DNA (97).

These results lead to an uptake-reduction model (Fig. 3) to explain the carcinogenicity of chromium (VI) (97). Chromium in the form of chromate penetrates the cell membrane using the sulfate transport system whereas chromium (III) does not cross cell membranes and is effectively excluded from the cell. Cellular metabolizing systems, including the microsomal electron-transport cytochrome P-450 system, reduce chromate to chromium (III).

The metabolically produced chromium (III) binds to nucleic acids, proteins and small molecules such as nucleotides. The chromium (III) bound to DNA and/or protein induces damage to the DNA that eventually leads to a mutation and hence cancer. Possible cocarcinogenic effects of chromium may be enzyme inhibition by bound chromium and/or by chromium-small molecule complexes. Extracellular chromium may interfere with normal receptor activities at the membrane and thereby disrupt cellular regulatory processes.

Chromium (III)-nucleotide complexes have been synthesized which have  $\text{Cr}^{3+}$  bound to phosphate groups (101, 102). These  $\text{Cr}^{3+}$ -nucleotides have been used as inhibitors and/or substrates of a number of phosphoryl transfer enzymes, e.g., acetate, pyruvate and 3-phosphoglycerate kinases, hexokinase, glycerokinase, creatine kinase, and phosphofructokinase (103).  $\text{CrADP}$  was shown to compete with  $\text{MgATP}$ , the normal substrate for hexokinase, for binding at the active site of the enzyme, whereas,  $\beta\text{-}\gamma\text{-CrATP}$  was shown to be a substrate for hexokinase (104).

After oral administration of chromium (III) to rats the chromium specifically was bound to the serum protein transferrin (105). The affinity of  $\text{Cr}^{3+}$  for transferrin was close to that of iron (III). Rats injected intravenously with vanadate retained vanadium in the plasma as a vanadium-transferrin complex (106). Both  $\text{Cr}^{3+}$  and  $\text{VO}^{2+}$  have been shown to bind to the  $\text{Fe}^{3+}$  sites of transferrin (107). Transferrin has been shown to bind to receptors on reticulocytes and become internalized by endocytosis




FIGURE 3. Uptake reduction model for the carcinogenicity of chromium (VI).

(108). The affinity of  $\text{Cr}^{3+}$ -saturated transferrin for reticulocytes was about half that of  $\text{Fe}^{3+}$  saturated transferrin (109). The reticulocytes do not take up  $\text{Cr}^{3+}$  from  $\text{Cr}^{3+}$ -transferrin bound at the receptor site (109). It has been suggested that transferrin may function as a carrier for metals other than iron, i.e.,  $\text{Co}^{2+}$ ,  $\text{Cr}^{3+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Mo}^{4+}$ , and  $\text{V}^{4+}$  (107).

Rumen fluid that had been incubated with molybdate exhibited an EPR signal characteristic of Mo (V) (11). Microorganisms such as the yeast *Pichia guilliermondii* and the bacterium *Micrococcus* reduced molybdenum trioxide and molybdate to molybdenum blue which is characteristic of pentavalent molybdenum complexes (110). Mo (V) EPR signals have been detected in bacteria, plant and animal tissues which contain molybdenum enzymes such as xanthine oxidase, sulfite oxidase and formate dehydrogenase (111). Rats were capable of incorporating molybdenum into liver sulfite oxidase and xanthine oxidase when injected with molybdate, however, molybdate was not able to reconstitute the apoenzymes *in vitro* (112). Pentavalent and hexavalent molybdenum were rapidly excreted from the rat after IV injection; however, at high doses (4.6 mg Mo/kg body weight) pentavalent molybdenum was retained to a greater extent than the hexavalent form in blood, liver and jejunioileocecum (113). It was proposed the biliary excretion of hexavalent molybdenum involved the secretory activity of hepatic cells and the liver cells oxidized molybdenum (V) to molybdenum (VI) which was then excreted in bile (113). When cupric molybdate was added to serum, the molybdate remained in solution as the free anion (11). Molybdenum was associated with red cells and plasma in the blood of sheep after oral administration of molybdate (114). All of the molybdenum in blood was dialysable and it was suggested that its form in both plasma and red blood cells was molybdate (114). Incubation of human erythrocytes with molybdenum (V) complexes resulted in labeling of the phosphorylated membrane protein spectrin (115). Incubation of human and rat serum with molybdate resulted in molybdenum bound to  $\alpha_2$ -macroglobulins (116).

Mouse liver extracts enzymatically converted selenite into dimethylselenide (117). Glutathione and S-adenosyl-L-methionine were required for the liver microsomal reduction and methylation of selenite and arsenite was an inhibitor of the reaction (117). Purified yeast glutathione reductase catalyzed the reduction of selenite to hydrogen selenide under anaerobic conditions in the presence of glutathione and NADPH (118). It was proposed that the mechanism of selenite reduction involved the formation of selenodiglutathione from nonenzymatic

reaction of glutathione and selenite which was reduced stepwise by glutathione reductase to glutathione selenopersulfide and then to hydrogen selenide (118). Dimethylselenide was exhaled by rats fed selenate or selenite (119). Trimethylselenonium ion  $(\text{CH}_3)_3\text{Se}^+$ , was the major urinary metabolite in rats injected with selenite, selenate, selenocystine, selenomethionine and methylselenocysteine, or fed seleniferous wheat (120). Hydrogen selenide was released upon acidification of liver homogenates from selenite-treated rats and was mainly associated with the mitochondrial and endoplasmic reticulum fractions (121, 122). In addition to selenide, selenite and higher oxidation states of selenium including organic derivatives were found in the liver of rats treated orally with selenite (122). Selenium-bound proteins were isolated from the plasma of rats injected with selenite (123). Incubation of selenite with plasma or whole blood *in vitro* did not produce the selenium-bound plasma proteins. It was suggested that metabolism of selenite was necessary for incorporation of selenium into these proteins (123). The activity of glutathione peroxidase, a seleno-enzyme, was raised in mouse neuroblastoma cells upon exposure to selenite (124). Selenate produced no increase in enzyme activity and it was suggested that these cells had no mechanism capable of reducing selenate to selenite (124). Selenite was rapidly taken up by mouse lung fibroblasts and incorporated in glutathione peroxidase whereas, selenomethionine was slowly taken up by the cells and its selenium became associated with glutathione peroxidase only after a long lag period (125). Selenocystine has been identified as the form of selenium in rat liver glutathione peroxidase (126).

Selenium was incorporated into prolyl-tRNA of *Clostridium sticklandii* incubated with selenite or selenocystine by a process highly specific for selenium not by selenium simply acting as analog of sulfur (127). Rumen microorganisms were capable of reducing selenate to selenite and metabolizing the selenite to a form (probably selenocystine) which was incorporated into microbial protein (128). No synthesis of selenomethionine by rumen fluid occurred from selenite or selenate and sulfate had no effect on the incorporation of selenium into protein. It was concluded that the metabolism of inorganic selenium by rumen microorganisms was different from that of inorganic sulfur (128). The fungus *Penicillium* produced dimethylselenide, upon incubation with selenite, selenate, or sodium selenide (129). Dimethylselenide, dimethyldiselenide, and dimethylselenone or methyl methylselenite was formed by soil and sewage sludge microorganisms treated with selenite or elemental selenium (130).

Injection of arsenate intravenously in dogs resulted in excretion of some arsenite in the urine (131). Renal reduction of arsenate occurred intracellularly. Very little reduction of arsenite occurred upon incubation with blood or urine *in vitro*. Renal tubular cells were permeable to both arsenite and arsenate and phosphate was an inhibitor of transport of both ions (131). Liver tissue from cows fed arsenate showed only pentavalent arsenic, indicating no reduced form of arsenic was retained in the tissues (132). The same results were found in a similar study with arsenite (133). More than 50% of arsenic excreted in the urine of dogs and cows fed arsenate or arsenite was methanearsonate (134). Livers of rats fed arsenate showed only pentavalent arsenic (135). Rats were unique among animals tested in their ability to retain large quantities of arsenic in the blood when fed either pentavalent or trivalent arsenic (132). The arsenic was found to be bound mainly to hemoglobin in the blood (136). Cacodylic acid was not converted into inorganic arsenic by rats administered the compound by intravenous injection, intratracheal instillation or oral gavage (137). The major site of retention of cacodylic acid was in the red blood cells (137).

Arsenic in human urine was predominantly in the form of cacodylic acid,  $\text{As}(\text{CH}_3)_2\text{O}_2\text{H}$ , with lesser amounts of arsenate and small amounts of methanearsonic acid and arsenite present (138). Copper smelter workers exposed mainly to airborne arsenic (III) excreted mainly dimethylarsinic acid (65%) and methylarsonic acid (20%), along with small amounts of arsenite (9%) and arsenate (6%) in their urine (139). Ingestion of wine containing arsenic (III) by humans resulted in urinary excretion of dimethylarsinic acid (50%), methylarsonic acid (14%), arsenite (8%), and arsenate (8%) (140). Arsenate was rapidly excreted in the urine of humans after ingestion of arsenate in well water; slower excretion of small amounts of arsenite and substantial amounts of dimethylarsinic acid and methylarsonic acid was observed (140).

The microorganisms *Scopuloriopsis brevicaulis*, *Candida humicola*, and *Gliocladium roseum* produced volatile trimethylarsine when incubated with arsenic trioxide ( $\text{As}_2\text{O}_3$ ), methylarsonic acid or dimethylarsinic acid (141). The methyl donor used for the biological methylation of arsenic by these microorganisms was *S*-adenosylmethionine (141). Methylcobalamin served as the methyl donor in the methylation of arsenite or arsenate to dimethylarsine by cell extract of *Methanobacillus* strain M. O. H. (142). Bacteria cultured from sea water reduced arsenate to arsenite (143). The marine alga, *Tetraselmis chuii*, and *Daphnia magna* incorporated arsenic from arsenate into the lipid fraction

(144). The arsenic compound was hydrolyzed by phospholipases and was tentatively identified as an arsenic-containing choline. An arsenylated low molecular weight substance which cochromatographed with phosphatidylethanolamine was isolated from arsenate-treated rat liver mitochondria (145).

A number of metabolic processes have been identified to be involved in biological conversions of oxyanions into different forms, e.g., NAD(P)H-dependent redox processes, reactions involving active sulfhydryl groups, and methyltransferase reactions.

## Mutagenicity

Chromate has been detected as a mutagen in the *Salmonella typhimurium* Ames assay whereas chromium (III) was not mutagenic in this system (146). Addition of a complete microsomal activation system decreased the mutagenicity of chromate (147). Since a microsomal system has been shown to convert chromate to chromium (III) (96), the decrease in mutagenicity can be explained by the microsomal conversion of the mutagenic chromium (VI) to the nonmutagenic chromium (III). Arsenite and arsenate were not mutagenic in the Ames assay (146). There are conflicting reports on the mutagenicity of selenite in the Ames assay. In one case both selenite and selenate were found to be mutagenic (148) whereas the other report found mutagenicity with selenate but not with selenite (146). The *Bacillus subtilis* rec-assay showed that both arsenite and arsenate were mutagenic although arsenite was a stronger mutagen (149). The rec-assay showed that chromate was mutagenic but not chromium (III); permanganate was not mutagenic but manganese (II) was positive; molybdate was positive but pentachloromolybdenum (V) was negative; both selenate and selenite were negative (150) at low concentrations but were positive at higher concentrations (144); and both vanadyl chloride and vanadate were positive (150). Arsenite, dichromate and molybdate were mutagenic in *E. coli* strains which were  $\text{recA}^+$  but not in a strain carrying  $\text{recA}^-$  (148). Chromate was mutagenic in *E. coli* WP2 wild strain and in *errA*<sup>-</sup> and *uvrA*<sup>-</sup> strains, whereas chromium (III), and soluble salts of molybdenum and tungsten were not mutagenic (151). Fluctuation tests which use *E. coli* WP2  $\text{Trp}^+$  reversion detected chromate as a mutagen (152). Arsenite decreased the survival of UV-irradiated wild-type *E. coli* WP2 but had no effect on the survival of a *recA* mutant, suggesting arsenite inhibited *recA*-dependent DNA repair (153). Arsenite was not mutagenic to *E. coli* WP2 at low

concentrations (154). Methylated forms of arsenic were not mutagenic in bacterial assays (66).

Chromate and arsenite enhanced the transformation of Syrian hamster embryo cells by simian adenovirus SA7 (155). Chromate transformed BHK21 cells *in vitro* (156); chromate and arsenate transformed Syrian hamster embryo cells, however, tungsten did not cause transformation (157). Chromosomal aberrations were induced in cultured FM3A mouse mammary carcinoma cells by chromium (VI) and permanganate, but not by chromium (III) and only at a low level by manganese (II) (158). Chromate, but not chromium (III), induced chromosomal aberrations and sister chromatid exchanges (SCE) in human fibroblasts (159) and in Chinese hamster ovary cells (160). The number of chromosomal aberrations in Syrian hamster embryo cells caused by chromium (VI) was diminished upon addition of the reducing agent sulfite (161). Arsenate induced dose-dependent SCE's and chromosomal aberrations in normal human lymphocytes (162). Arsenite and acetylarsan caused chromosomal aberrations in human leukocyte cultures, however, no chromosomal aberrations were induced by arsenate, selenite, selenate or metavanadate (163). Arsenite also induced chromosomal aberrations in human diploid fibroblasts (162). In another study, selenium (VI) compounds induced chromosomal aberrations in cultured human leukocytes whereas selenium (IV) compounds were inactive (164). Selenite did not cause SCE's in purified human lymphocyte cultures or xeroderma pigmentosum cells, however, addition of red blood cell lysate with selenite resulted in induction of SCE in both cell types (165). Induction of chromosomal aberrations and DNA-repair synthesis in human diploid fibroblasts by selenite was enhanced by addition of a mouse liver S-9 microsomal fraction and NADPH (166). Selenate induced only a small amount of DNA-repair synthesis which could not be enhanced by addition of the S-9 fraction (166).

Arsenic (V), selenium (IV), and chromium (VI) did not affect the fidelity of *in vitro* DNA synthesis by a variety of DNA polymerases copying a poly[d(A-T)] template (167, 168). Chromium (III) caused a large increase in the error frequency of DNA polymerases copying poly[d(A-T)] or poly[d(G-C)] templates (167, 168). Cellular metabolism of the oxyanions which initially enter cells is obviously important in determining their mutagenic potential.

## Divalent Ions

Most other toxic and/or mutagenic metals exist as divalent metal ion complexes in neutral aqueous

solution (1), e.g.,  $\text{Be}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Pt}^{2+}$ ,  $\text{Hg}^{2+}$ , and  $\text{Pb}^{2+}$ , although higher or lower oxidation states are possible depending on the particular metal and coordinated ligands. The ionic radii of the first row divalent transition metal ions lie between those of  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$  (1). Other divalent metal ions can replace or mimic ions such as magnesium and calcium, which are extremely important in biological systems, and iron, zinc, copper, cobalt, and manganese, which are essential components of many enzymes or coenzymes.

*E. coli* cells were found to be highly permeable to manganese ion which was a potent mutagen in the bacteria (169, 170). Magnesium competed with manganese uptake by the cells and reduced the number of mutations (170). Manganese was able to substitute for magnesium during *in vitro* DNA synthesis by avian myeloblastosis DNA polymerase, however,  $\text{Mn}^{2+}$  increased the incorporation of noncomplementary nucleotides into the complementary strand (171). Subsequently it was found that  $\text{Be}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Ni}^{2+}$ , and  $\text{Pb}^{2+}$  in addition to  $\text{Mn}^{2+}$  decreased the fidelity of DNA synthesis *in vitro* (168). In the case of  $\text{Be}^{2+}$  infidelity was ascribed to the interaction of  $\text{Be}^{2+}$  with the DNA polymerase enzyme (171). Manganese was also found to be mutagenic in the *B. subtilis* "rec-assay" as were cadmium and methyl mercury, however, in this cellular assay system inorganic salts of  $\text{Be}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Hg}^{2+}$ ,  $\text{Ni}^{2+}$ , and  $\text{Pb}^{2+}$  were inactive (149).

Replacement of  $\text{Mg}^{2+}$  by  $\text{Mn}^{2+}$  relaxed the specificity of the restriction enzymes EcoRI and HindIII, causing many more cleavages of DNA to occur (173). Manganese enhanced the inhibition of terminal deoxynucleotidyltransferase (TdT)-catalyzed DNA synthesis by ribonucleoside triphosphates (174).  $\text{Mn}^{2+}$ -dATP inhibited TdT by binding 10-100 times stronger to the substrate binding site than other  $\text{Mn}^{2+}$ -deoxynucleotide triphosphates and therefore blocking the reaction. DNA synthesis by TdT in the presence of  $\text{Mg}^{2+}$  was instantly inhibited upon the addition of  $\text{Mn}^{2+}$  (174).  $\text{Mn}^{2+}$  altered the composition of the RNA products formed *in vitro* by *Euglena gracillius* RNA polymerases I and II from zinc sufficient cells and the single RNA polymerase from zinc deficient cells (174). The ability of  $\text{Mn}^{2+}$  to allow DNA polymerase to synthesize ribosubstituted DNA and to allow RNA polymerase to synthesize deoxysubstituted RNA has been used for nucleotide sequence determination (175). The specificity of calf thymus ribonuclease H was decreased in the presence of  $\text{Mn}^{2+}$  (176). All DNA-RNA hybrid combinations were cleaved with  $\text{Mn}^{2+}$  whereas only hybrids containing purine ribo strands were cleaved with  $\text{Mg}^{2+}$  (176).

Addition of  $Mn^{2+}$  or  $Fe^{3+}$  to mixtures of rat liver microsomes and benzo[a]pyrene (BP) significantly altered the formation of three diols of BP (177). It was suggested that  $Mn^{2+}$  and  $Fe^{3+}$  could enhance the carcinogenicity of BP by altering the specificity of microsomal metabolism (177).  $Cu^{2+}$ ,  $Zn^{2+}$ ,  $Pb^{2+}$ ,  $Ni^{2+}$ ,  $Fe^{2+}$ ,  $Fe^{3+}$ , and  $Mg^{2+}$  inhibited the metabolism of BP by rat liver microsomes;  $Mn^{2+}$  activated metabolism at low concentrations and inhibited it at high concentrations (95). Another study showed  $Co^{2+}$ ,  $Mg^{2+}$ , and  $Al^{3+}$  enhanced the hydroxylation of BP; and  $Zn^{2+}$ ,  $Cu^{2+}$ ,  $Pb^{2+}$ ,  $Mn^{2+}$ ,  $Cd^{2+}$ , and  $Ni^{2+}$  enhanced or inhibited hydroxylation depending upon the concentration of the metal ion (94). An earlier study showed that  $Be^{2+}$ ,  $Fe^{3+}$ , and  $Mn^{2+}$  inhibited BP-hydroxylase activity in rat lung homogenates; and  $Cu^{2+}$ ,  $Mg^{2+}$ ,  $Fe^{2+}$ ,  $Zn^{2+}$ ,  $Ni^{2+}$ , and  $Co^{2+}$  stimulated the enzyme at low concentrations and inhibited activity at high concentrations (93).

$Mn^{2+}$  and  $Cu^{2+}$  enhanced unscheduled DNA synthesis induced by isoniazid and related hydrazines in human fibroblasts whereas  $Fe^{3+}$  had no effect (178). It was suggested that the DNA damage was caused by hydroxyl radicals produced by reaction of hydrogen peroxide with the metal ions (178).  $Cu^{2+}$ ,  $Co^{2+}$ ,  $Mn^{2+}$ ,  $Fe^{2+}$ , and  $Zn^{2+}$  enhanced the ability of thymine hydroperoxide to transform the DNA of *Haemophilus influenzae*, and it was suggested that the damaging agents were free radicals which formed upon reaction of the metal ions with hydroperoxides (178).  $Cu^{2+}$ ,  $Mn^{2+}$ ,  $Fe^{2+}$ , and  $Fe^{3+}$  enhanced induction of chromosomal aberrations in Chinese hamster ovary cells by ascorbate (180). Hydrogen peroxide, which produces reactive free radicals, appeared to be the DNA-damaging agent in these studies (180).

$Pb^{2+}$ ,  $Cd^{2+}$ ,  $Cu^{2+}$ , and  $Mn^{2+}$  decreased the fidelity of DNA-directed RNA synthesis by *E. coli* RNA polymerase, however, these ions stimulated chain initiation at concentrations which inhibited total RNA synthesis (181). It was suggested that these ions promoted RNA initiation at sites on the DNA template not normally recognized as initiation sites (181).

$Mn^{2+}$  replaced  $Mg^{2+}$  at a high affinity binding site on tubulin and promoted tubulin polymerization to microtubules (182).  $Co^{2+}$  and  $Zn^{2+}$ , but not  $Ca^{2+}$ , also bound to the same site, however, tubulin sheets rather than the microtubules formed in their presence indicating that  $Co^{2+}$  and  $Zn^{2+}$  induced a different conformation of tubulin than  $Mn^{2+}$  or  $Mg^{2+}$  (182). Replacement of  $Ca^{2+}$  by  $Mn^{2+}$  in the extracellular media resulted in a decrease in the tryptophan hydroxylase activity of slices of rat brain stem (183). It was concluded that  $Ca^{2+}$  regulated trypto-

phan hydroxylase activity via  $Mn^{2+}$ -sensitive  $Ca^{2+}$  channels in the nerve membrane (183).  $Co^{2+}$  was a potent inhibitor of  $Ca^{2+}$  translocation mediated by the ionophore A23187 into a hydrophobic domain (184).  $Co^{2+}$  aggregated at synaptic sites and on or near microtubules along the neuronal membranes after injection of  $Co^{2+}$  into the neurons of the locust *Schistocerca americana gregaria* (185). Since calmodulin has been localized at these sites it was suggested that  $Co^{2+}$  bound to calcium-binding proteins in or near the membranes and microtubules (185).  $Co^{2+}$  and  $Mn^{2+}$  blocked the calcium channels in nerve cell bodies of *Tritonia diomedea* (186) and in squid axons (187) in a competitive manner.  $Mn^{2+}$ ,  $Co^{2+}$ , and  $Ni^{2+}$  also decreased the  $Na^{+}$  currents in squid axons but had no effect on  $K^{+}$  channels (187).  $Mg^{2+}$ ,  $Ca^{2+}$ ,  $Sr^{2+}$ ,  $Mn^{2+}$  and  $Ni^{2+}$  decreased the current induced by acetylcholine at the neuromuscular junction of the frog cutaneous pectoris muscle, however,  $Ni^{2+}$  and  $Sr^{2+}$  markedly lengthened the current decay rate whereas  $Mg^{2+}$ ,  $Ca^{2+}$ , and  $Mn^{2+}$  had little effect on the rate (188).

Substitution of zinc in zinc-metalloenzymes by other metals resulted in fully active enzymes or enzymes with altered activities. Substitution of  $Ni^{2+}$  for  $Zn^{2+}$  in the regulatory subunits of aspartate transcarbamoylase resulted in a fully active enzyme having allosteric properties of the native Zn-enzyme (189). However, the native enzyme showed a higher degree of cooperativity than the  $Ni^{2+}$  substituted enzyme possibly because the  $Ni^{2+}$  induced a slightly different conformation in the protein chains (189). Substitution of the zinc in carbonic anhydrase by  $Ni^{2+}$  resulted in complete loss of esterase activity, however, the  $Cd^{2+}$ -substituted enzyme showed esterase activity above pH 9 (190).  $Co^{2+}$ -substituted carbonic anhydrase retained full esterase activity, however, substitution of  $Cd^{2+}$ ,  $Mn^{2+}$ ,  $Ni^{2+}$ ,  $Fe^{2+}$ ,  $Cr^{3+}$ , or  $Fe^{3+}$  resulted in loss of activity (191). Substitution of all four atoms of zinc in horse liver alcohol dehydrogenase (192) or in *E. coli* alkaline phosphatase (193) by  $Co^{2+}$  resulted in decreased enzyme activity. Activities of various metal-substituted alkaline phosphatases were correlated with the distance of the metal to a bound water molecule (194). The  $Mn^{2+}$ -alkaline phosphatase which was inactive had the longest  $Mn-H_2O$  distance, 4.0 Å; the  $Cu^{2+}$ -enzyme with 3-5% the native activity had a  $Cu^{2+}-H_2O$  distance of 3.4 Å; and the  $Co^{2+}$ -enzyme had a nearly normal distance of 2.8 Å and retained 10-20% the activity of the native zinc enzyme (194). Cobalt-leucine aminopeptidase had a higher specific activity with L-leucine-*p*-nitroanilide substrate than the native zinc enzyme (195). Substitution of  $Co^{2+}$ ,  $Mn^{2+}$ , or  $Cd^{2+}$  for  $Zn^{2+}$  in carboxypeptidase resulted in mark-

edly different effects on the peptidase and esterase activities of the enzyme (196). The  $\text{Co}^{2+}$ -,  $\text{Mn}^{2+}$ -, and  $\text{Cd}^{2+}$ -carboxypeptidase had essentially the same binding affinity for peptide substrates as the native zinc enzyme, however, the rate of peptide hydrolysis was much slower by the  $\text{Mn}^{2+}$ - and  $\text{Cd}^{2+}$ -enzymes and substantially faster by the  $\text{Co}^{2+}$ -enzyme. In contrast, rates of ester hydrolysis were virtually identical regardless of the metal, however, the  $\text{Zn}^{2+}$ - and  $\text{Co}^{2+}$ -carboxypeptidase had much higher affinity for the ester substrates than the  $\text{Mn}^{2+}$ - or  $\text{Cd}^{2+}$ -enzyme (196).

The tail baseplate of T-even bacteriophages was shown to contain about 5 atoms of  $\text{Zn}^{2+}$  which were necessary for activity of the phage particles (197).  $\text{Co}^{2+}$ ,  $\text{Cd}^{2+}$ , and  $\text{Ni}^{2+}$  were capable of restoring activity to Zn-depleted phage particles whereas little or no reconstitution of phage activity was observed with  $\text{Mn}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Cu}^{2+}$ , or  $\text{Hg}^{2+}$  (197).  $\text{Co}^{2+}$  and  $\text{Ni}^{2+}$  were able to substitute *in vivo* for  $\text{Zn}^{2+}$  in the tail baseplate of bacteriophage T4D and permitted growth of active phage particles on *E. coli* B (198).

$\text{Zn}^{2+}$  inhibited glucagon-stimulated and fluoride-stimulated adenylate cyclase activity in human liver membranes (199).  $\text{Zn}^{2+}$  inhibited acid phosphatase activity in the nuclear and particulate fractions of adult chicken liver homogenates, whereas,  $\text{Mg}^{2+}$  slightly activated the enzyme activity (200).  $\text{Zn}^{2+}$  complexed with human rhinovirus type 1A coat proteins and rendered them inactive as substrates for proteases and prevented their use in the assembly of the intact virus (201).

$\text{Zn}^{2+}$  enhanced the dissociation of human  $\alpha_2\beta_2$  oxy- or carbon monoxyhemoglobin tetramers to  $\alpha\beta$  dimers which contained one zinc bound per dimer (202). It was suggested that Cys 93 $\beta$ , His 143 $\beta$  and His 146 $\beta$  may coordinate  $\text{Zn}^{2+}$  and the resulting charge repulsion may lead to the observed dissociation (202).  $\text{Zn}^{2+}$  was found to be concentrated in the sulfhydryl-rich flagellar accessory fibers of mammalian and cephalopod sperm (203). Dexamethasone, a synthetic glucocorticoid, stimulated the uptake of  $\text{Zn}^{2+}$  in HeLa cells by inducing the synthesis of metallothionein, a cysteine-rich protein with high affinity for  $\text{Zn}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Hg}^{2+}$ ,  $\text{Ag}^+$ , and  $\text{Cu}^+$  (204).  $\text{Cd}^{2+}$  inhibited soluble rat testicular and liver glutathione peroxidase activity *in vitro* (205).

$\text{Cd}^{2+}$  promoted incorporation of delipidated human placental tissue factor into phospholipid vesicles and thereby greatly enhanced recovery of the coagulant activity of tissue factor (206).  $\text{Cu}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Hg}^{2+}$ , or  $\text{Zn}^{2+}$  decreased recovery of coagulant activity,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Sr}^{2+}$  and  $\text{Ba}^{2+}$  had no effect, and  $\text{Mn}^{2+}$  and  $\text{Co}^{2+}$  slightly enhanced activity. It was suggested that the effectiveness of

$\text{Cd}^{2+}$  and  $\text{Mn}^{2+}$  was due to their known ability to induce fusion of vesicles from lipids (206).

$\text{Mn}^{2+}$  was a noncompetitive inhibitor of glutamine synthetase activity in rat liver cytosol extracts (207).  $\text{Hg}^{2+}$  and  $\text{Cu}^{2+}$  strongly inhibited soya meal urease activity, whereas, slight or no inhibition of urease was observed in the presence of  $\text{Ba}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Cr}^{3+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Fe}^{3+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Sn}^{2+}$ ,  $\text{Sr}^{2+}$ , or  $\text{Zn}^{2+}$  (208).  $\text{Pb}^{2+}$  stimulated phosphorylation of electroplax microsomal protein by the ( $\text{Mg}^{2+}$  +  $\text{Na}^+$ )-dependent protein kinase of ( $\text{Na}^+$ ,  $\text{K}^+$ )ATPase, thus inhibiting ATP hydrolysis (209).  $\text{Pb}^{2+}$  replaced  $\text{Na}^+$  in stimulating the kinase activity, whereas  $\text{Fe}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Sr}^{2+}$ ,  $\text{Hg}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Mn}^{2+}$ , and  $\text{Co}^{2+}$  were inactive. Methylmercury and  $\text{Hg}^{2+}$  inhibited rabbit muscle sarcoplasmic reticulum ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ )ATPase and  $\text{Ca}^{2+}$ -transport by blockage of essential sulfhydryl groups (210).  $\text{Hg}^{2+}$  also competed with  $\text{Ca}^{2+}$  for the  $\text{Ca}^{2+}$ -ionophoric site and inhibited  $\text{Ca}^{2+}$ -transport. Methylmercury, but not  $\text{Hg}^{2+}$ , inhibited the binding of acetylcholine to the acetylcholine receptor from the electric organ of *T. californica* (210). It was suggested that methylmercury reacted with sulfhydryl groups of acetylcholine receptor which were located in a hydrophobic environment inaccessible to  $\text{Hg}^{2+}$  (210).

$\text{Pb}^{2+}$  inhibited milk xanthine oxidase activity (211).  $\text{Pb}^{2+}$ ,  $\text{Ag}^+$  and  $\text{Hg}^{2+}$  strongly inhibited rabbit liver guanine aminohydrolase, a sulfhydryl enzyme, whereas  $\text{Cd}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Ca}^{2+}$ ,  $\text{Be}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Sn}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Fe}^{2+}$ , and  $\text{Ni}^{2+}$  exhibited no inhibition of guanase activity (211).  $\text{Pb}^{2+}$  disturbed heme synthesis by inhibiting the cytoplasmic Zn-dependent, sulfhydryl-rich enzyme, aminolevulinic acid dehydratase, and the mitochondrial sulfhydryl enzyme, ferrochelatase (212).  $\text{Cu}^{2+}$ ,  $\text{Hg}^{2+}$ , and  $\text{Ag}^+$  also inhibited aminolevulinic acid dehydratase (212).

Bacteria from marine sediments oxidized Mn (II) to Mn (III) and this oxidation was shown to involve type C cytochrome (213). Manganosuperoxide dismutases found in bacteria and mitochondria were found to have manganese bound in the trivalent state (214). Transferrin was found to bind manganese as  $\text{Mn}^{3+}$  (107).

Elemental mercury vapor readily crossed red cell membranes and was oxidized to divalent mercury by catalase (215). Bacteria from marine sediments methylated  $\text{Hg}^{2+}$  to methylmercury which was accumulated by living organisms much more rapidly and avidly than  $\text{Hg}^{2+}$  (215). Organolead compounds were methylated by microbial components of sediment, however, no stable methylated-derivative was isolated with  $\text{Pb}^{2+}$  (216). Methylation of mercury and lead involved methyl- $\text{B}_{12}$  as the methyl donor (216).

Divalent ions have diverse effects which may contribute to their carcinogenicity or cocarcinogenicity. They may substitute for essential ions such as calcium, magnesium and zinc; they may react with sulfhydryl groups of enzymes; they may alter enzyme and membrane functions; they may be metabolized to more toxic agents; they may be incorporated into metalloenzymes; they may bind to macromolecules or small molecules and alter their normal activity in the cell; or they may catalyze the formation of toxic agents such as free radicals.

This investigation was supported by Grant Number R01-CA22270, awarded by the National Cancer Institute, DHEW, by the donors of The Petroleum Research Fund, administered by the American Chemical Society, and by Biomedical Research Support Grant RR-05392 from the Biomedical Research Support Branch, Division of Research Facilities and Resources, National Institutes of Health.

## REFERENCES

- Cotton, F. A., and Wilkinson, G., *Advanced Inorganic Chemistry*, Interscience Publishers, New York, 1972.
- Carofoli, E., and Crompton, M. Binding proteins and membrane transport. In: *The Enzymes of Biological Membranes*, Vol. 3, Membrane Transport. A. Martonosi, Ed., Plenum Press, New York, 1976, pp. 193-219.
- Jung, C., and Rothstein, A. Arsenate uptake and release in relation to the inhibition of transport and glycolysis in yeast. *Biochem. Pharmacol.* 14: 1093 (1965).
- Hilborn, D. A. Serum stimulation of phosphate uptake into 3T3 cells. *J. Cell Physiol.* 87: 111 (1976).
- Cantley, L. C., Resh, M. D., and Guidotti, G. Vanadate inhibits the red cell ( $\text{Na}^+$ ,  $\text{K}^+$ ) ATPase from the cytoplasmic side. *Nature* 272: 552 (1978).
- Gray, S. J., and Sterling, K. The tagging of red cells and plasma proteins with radioactive chromium. *J. Clin. Invest.* 29: 1604 (1950).
- Rajam, P. C., and Jackson, A.-L. Distribution and valence state of radiochromium in intracellularly labeled Ehrlich mouse ascites carcinoma cells. *Proc. Soc. Exp. Biol. Med.* 99: 210 (1958).
- Sanderson, C. J. The uptake and retention of chromium by cells. *Transplant.* 21: 526 (1976).
- Johnson, P. K. and Mordiney, M. R. Binding studies concerning the interaction of radiolabeled sodium chromate with EL4 leukemia. *Transplant.* 14: 253 (1972).
- Lilien, D. L., Spivak, J. L. and Goldman, I. D. Chromate transport in human leukocytes. *J. Clin. Invest.* 49: 1551 (1970).
- Huisingh, J., and Matrone, G. Interaction of molybdenum in animal nutrition. In: *Molybdenum in the Environment*, Vol. 1, The Biology of Molybdenum, W. R. Chappell and K. K. Peterson, Eds., Marcel Dekker, New York, pp. 125-148 (1976).
- Cardin, C. J., and Mason, J. Sulphate transport by rat ileum. Effect of molybdate and other anions. *Biochim. Biophys. Acta* 394: 46 (1975).
- Cardin, C. J., and Mason, J. Molybdate and tungstate transfer by rat ileum: Competitive inhibition by sulphate. *Biochim. Biophys. Acta* 455: 937 (1976).
- Mason, J., and Cardin, C. J. The competition of molybdate and sulphate ions for a transport system in the ovine small intestine. *Res. Vet. Sci.* 22: 313 (1977).
- Huisingh, J., Gomez, G. G., and Matrone, G. Interactions of copper, molybdenum, and sulfate in ruminant nutrition. *Fed. Proc.* 32: 1921 (1973).
- McCready, R. G. L. and Din, G. A. Active sulfate transport in *Saccharomyces cerevisiae*. *FEBS Lett.* 38: 361 (1974).
- Cuppoletti, J., and Segel, I. H. Transinhibition kinetics of the sulfate transport system of *Penicillium notatum*: analysis based on an iso uni uni velocity equation. *J. Membrane Biol.* 17: 239 (1974).
- Elliott, B. B., and Mortenson, L. E. Transport of molybdate by *Clostridium pasteurianum*. *J. Bacteriol.* 124: 1295 (1975).
- Pardee, A. B., Prestidge, L. S., Whipple, M. B., and Dreyfuss, J. A binding site for sulfate and its relation to sulfate transport into *Salmonella typhimurium*. *J. Biol. Chem.* 241: 3962 (1966).
- Van Etten, R. L., Waymack, P. P. and Rehkop, D. M. Transition metal ion inhibition of enzyme-catalyzed phosphate ester displacement reactions. *J. Am. Chem. Soc.* 96: 6782 (1974).
- Lazdunski, C., and Lazdunski, M. Étude cinétique du mécanisme d'action catalytique de la phosphatase alcaline d'*Escherichia coli*. *Biochim. Biophys. Acta* 113: 551 (1966).
- Lopez, V., Stevens, T., and Lindquist, R. N. Vanadium ion inhibition of alkaline phosphatase-catalyzed phosphate ester hydrolysis. *Arch. Biochem. Biophys.* 175: 31 (1976).
- Ohlsson, J. K., and Wilson, I. B. The inhibition of alkaline phosphatase by periodate and permanganate. *Biochim. Biophys. Acta* 350: 43 (1974).
- Paietta, E., and Sands, H. Phosphoprotein phosphatase in bovine tracheal smooth muscle: Multiple fractions and multiple substrates. *Biochim. Biophys. Acta* 523: 121 (1978).
- Jacob, J.-L., and Sontag, N. Une enzyme nouvelle: la 2'-nucléotidase de latex d'*Hevea brasiliensis*. *Eur. J. Biochem.* 40: 207 (1973).
- Lindquist, R. N., Lynn, J. L., and Lienhard, G. E. Possible transition-state analogs for ribonuclease. The complexes of uridine with oxovanadium (IV) ion and vanadium (V) ion. *J. Am. Chem. Soc.* 95: 8762 (1973).
- Cantley, L. C., Josephson, L., Warner, R., Yanagisawa, M., Lechene, C., and Guidotti, G. Vanadate is a potent ( $\text{Na}$ ,  $\text{K}$ )-ATPase inhibitor found in ATP derived from muscle. *J. Biol. Chem.* 252: 7421 (1977).
- Josephson, L., and Cantley, L. C. Isolation of a potent ( $\text{Na}$ - $\text{K}$ ) ATPase inhibitor from striated muscle. *Biochemistry* 16: 4572 (1977).
- Beaugé, L. A. and Glynn, I. M. A modifier of ( $\text{Na}^+$  +  $\text{K}^+$ ) ATPase in commercial ATP. *Nature* 268: 355 (1977).
- Gibbons, I. R., Cosson, M. P., Evans, J. A., Gibbons, B. H., Houck, B., Martinson, K. H., Sale, W. S., and Tang, W.-J. Y. Potent inhibition of dynein adenosinetriphosphatase and of the motility of cilia and sperm flagella by vanadate. *Proc. Natl. Acad. Sci. (U.S.)* 75: 2220 (1978).
- Beaugé, L., and Dipolo, R. Vanadate selectively inhibits the  $\text{K}_o^+$ -activated  $\text{Na}^+$  efflux in squid axons. *Biochim. Biophys. Acta* 551: 220 (1979).
- Kobayashi, T., Martensen, T., Nath, J., and Flavin, M. Inhibition of dynein ATPase by vanadate, and its possible use as a probe for the role of dynein in cytoplasmic motility. *Biochem. Biophys. Res. Commun.* 81: 1313 (1978).
- Goodno, C. C. Inhibition of myosin ATPase by vanadate ion. *Proc. Natl. Acad. Sci. (U.S.)* 76: 2620 (1979).
- Bond, G. H., and Hudgins, P. M. Kinetics of inhibition of



- red cell membrane Ca-ATPase by pentavalent vanadium. Fed. Proc. 37: 313 (1978).
35. DiPolo, R., Rojas, H. R., and Beaugé, L. Vanadate inhibits uncoupled Ca efflux but not Na-Ca exchange in squid axons. Nature 281: 228 (1979).
  36. Cantley, L. C., Cantley, L. G. and Josephson, L. A characterization of vanadate interactions with the (Na,K)-ATPase. Mechanistic and regulatory implications. J. Biol. Chem. 253: 7361 (1978).
  37. Karlish, S. J. D., Beaugé, L. A., and Glynn, I. M. Vanadate inhibits (Na<sup>+</sup>, K<sup>+</sup>)ATPase by blocking a conformational change of the unphosphorylated form. Nature 282: 333 (1979).
  38. Grupp, G., Grupp, I., Johnson, C. L., Wallick, E. T., and Schwartz, A. Effects of vanadate on cardiac contraction and adenylate cyclase. Biochem. Biophys. Res. Commun. 88: 440 (1979).
  39. Schwabe, U., Puchstein, C., Hannemann, H., and Söchtig, E. Activation of adenylate cyclase by vanadate. Nature 277: 143 (1979).
  40. DeSousa, R. C., and Grosso, A. Vanadate blocks cyclic AMP-induced stimulation of sodium and water transport in amphibian epithelia. Nature 279: 803 (1979).
  41. Richards, J. M., and Swislocki, N. I. Activation of Adenylate cyclase by molybdate. J. Biol. Chem. 254: 6857 (1979).
  42. Nielsen, C. J., Sando, J. J., and Pratt, W. B. Evidence that dephosphorylation inactivates glucocorticoid receptors. Proc. Natl. Acad. Sci. (U.S.) 74: 1398 (1977).
  43. Nielson, C. J., Sando, J. J., Vogel, W. M., and Pratt, W. B. Glucocorticoid receptor inactivation under cell-free conditions. J. Biol. Chem. 252: 7568 (1977).
  44. Leach, K. L., Dahmer, M. K., Hammond, N. D., Sando, J. J., and Pratt, W. B. Molybdate inhibition of glucocorticoid receptor inactivation and transformation. J. Biol. Chem. 254: 11884 (1979).
  45. Nishigori, H., and Toft, D. Inhibition of progesterone receptor activation by sodium molybdate. Biochemistry 19: 77 (1980).
  46. Layne, P. P., and Najjar, V. A. Evidence for a tyrosine residue of the active site of phosphoglucomutase and its interaction with vanadate. Proc. Natl. Acad. Sci. (U.S.) 76: 5010 (1979).
  47. Rippa, M., Signorini, M., Bellini, T., and Dallochio, F. The active site of 6-phosphogluconate dehydrogenase: A phosphate binding site and its surroundings. Arch. Biochem. Biophys. 189: 516 (1978).
  48. Lee, Y. M. and Benisek, W. F. Inactivation of phosphorylase b by potassium ferrate, a new reactive analog of the phosphate group. J. Biol. Chem. 251: 1553 (1976).
  49. De Master, E. G., and Mitchell, R. A. A comparison of arsenate and vanadate as inhibitors or uncouplers of mitochondrial and glycolytic energy metabolism. Biochemistry 12: 3616 (1973).
  50. Jaffe, K., and Apitz-Castro, R. Studies on the mechanism by which inorganic arsenate facilitates the enzymatic reduction of dihydroxyacetone by  $\alpha$ -glycerophosphate dehydrogenase. FEBS Letter 80: 115 (1977).
  51. White, P. J., Kause, M. J., Edwards, J. O., and Rieger, P. H. Kinetics of an associative ligand exchange process: Alcohol exchange with arsenate (V) and vanadate (V) esters. J. Chem. Soc. Chem. Commun. 1976: 429.
  52. Choate, G. L., and Mansour, T. E. Inhibition of sheep heart phosphofructokinase by orthovanadate. Fed. Proc. 37: 1433 (1978).
  53. Scrutton, M. C., and Fung, C.-H. Pyruvate carboxylase from chicken liver: Effects of sulfate and other anions on catalytic activity and structural parameters. Arch. Biochem. Biophys. 150: 636 (1972).
  54. Wilson, L. G. and Bandurski, R. S. Enzyme reactions involving sulfate, sulfite, selenate, and molybdate. J. Biol. Chem. 233: 975 (1958).
  55. Varma, A. K., and Nicholas, D. J. D. Purification and properties of ATP-sulphurylase from *Nitrobacter agilis*. Biochim. Biophys. Acta 227: 373 (1971).
  56. Tweedie, J. W., and Segel, I. H. Adenosine triphosphate sulphyrylase from *Penicillium chrysogenum* II. Physical, kinetic, and regulatory properties. J. Biol. Chem. 246: 2438 (1971).
  57. Shaw, W. H., and Anderson, J. W. Purification, properties and substrate specificity of adenosine triphosphate sulphyrylase from spinach leaf tissue. Biochem. J. 127: 237 (1972).
  58. Shaw, W. H., and Anderson, J. W. Comparative enzymology of the adenosine triphosphate sulphyrylases from leaf tissue of selenium-accumulator and non-accumulator plants. Biochem. J. 139: 37 (1974).
  59. Hawes, C. S., and Nicholas, D. J. D. Adenosine 5'-triphosphate sulphyrylase from *Saccharomyces cerevisiae*. Biochem. J. 133: 541 (1973).
  60. Shoyab, M., and Marx, W. Enzyme-substrate complexes of ATP-sulphyrylase from mouse mastocytoma. Biochim. Biophys. Acta 258: 125 (1972).
  61. Ramus, J. *In vivo* molybdate inhibition of sulfate transfer to *Porphyridium* capsular polysaccharide. Plant Physiol. 54: 945 (1974).
  62. Pasternak, C. A. Sulphate activation and its control in *Escherichia coli* and *Bacillus subtilis*. Biochem. J. 85: 44 (1962).
  63. Huisingh, J., and Matrone, G. Copper-molybdenum interactions with the sulfate-reducing system in rumen microorganisms. Proc. Soc. Exptl. Biol. Med. 139: 518 (1972).
  64. McKillen, M. N., and Spencer, B. Molybdate toxicity in *Salmonella typhimurium*. Biochem. J. 118: 27p (1970).
  65. Reuveny, Z. Derepression of ATP sulphyrylase by the sulfate analogs molybdate and selenate in cultured tobacco cells. Proc. Natl. Acad. Sci. (U.S.) 74: 619 (1977).
  66. Léonard, A., and Lauwerys, R. R. Carcinogenicity, teratogenicity and mutagenicity of arsenic. Mutat. Res. 75: 49 (1980).
  67. Fowler, B. A. Toxicology of environmental arsenic. Adv. Mod. Toxicol. 2: 79 (1977).
  68. Johnstone, R. M. Sulphydryl agents: Arsenicals. In: Metabolic Inhibitors. A comprehensive treatise, Vol. 2, R. M. Hochster, and J. H. Quastel, Eds., Academic Press, New York, (1963), pp. 99-118.
  69. Jakoby, W. B. Aldehyde dehydrogenases. In: The Enzymes, Vol. 7, P. D. Boyer, H. Lardy, and K. Myrbäck, Eds., Academic Press, New York, 1963, pp. 203-221.
  70. Wu, C. Glutamine synthetase VI. Mechanism of the dithioldependent inhibition by arsenite. Biochim. Biophys. Acta 96: 134 (1965).
  71. Oehme, F. W. British antilewisite (BAL), the classic heavy metal antidote. Clin. Toxicol. 5: 215 (1972).
  72. Pisciotto, P. T., and Graziano, J. H. Induction of mucosal glutathione synthesis by arsenic. Biochim. Biophys. Acta 628: 241 (1980).
  73. Coughlan, M. P., Rajagopalan, K. V. and Handler, P. The role of molybdenum in xanthine oxidase and related enzymes. J. Biol. Chem. 244: 2658 (1969).
  74. Wilson, I. B., and Silman, I. Effects of quaternary ligands on the inhibition of acetylcholinesterase by arsenite. Biochemistry 16: 2701 (1977).
  75. Tsen, C. C., and Tappel, A. L. Catalytic oxidation of glutathione and other sulphydryl compounds by selenite. J. Biol. Chem. 233: 1230 (1958).
  76. Ganther, H. E. Selenotrisulfides. Formation by the reac-

- tion of thiols with selenious acid. *Biochemistry* 7: 2898 (1968).
77. Cummins, L. M., and Martin, J. L. Are selenocystine and selenomethionine synthesized *in vivo* from sodium selenite in mammals? *Biochemistry* 6: 3162 (1967).
  78. Ganther, H. E. Reduction of the selenotrisulfide derivative of glutathione to a persulfide analog by glutathione reductase. *Biochemistry* 10: 4089 (1971).
  79. Ganther, H. E. and Corcoran, C. Selenotrisulfides II. Crosslinking of reduced pancreatic ribonuclease with selenium. *Biochemistry* 8: 2557 (1969).
  80. Sumino, K., Yamamoto, R., and Kitamura, S. A role of selenium against methylmercury toxicity. *Nature* 268: 73 (1977).
  81. Levander, O. A., Morris, V. C., and Higgs, D. J. Acceleration of thiol-induced swelling of rat liver mitochondria by selenium. *Biochemistry* 12: 4586 (1973).
  82. Levander, O. A., Morris, V. C., and Higgs, D. J. Selenium as a catalyst for the reduction of cytochrome c by glutathione. *Biochemistry* 12: 4591 (1973).
  83. Iwata, H., Masukawa, T., Kasamatsu, S., and Komenushi, S. Stimulation of methemoglobin reduction by selenium: A comparative study with erythrocytes of various animals. *Experientia* 34: 534 (1978).
  84. Stadtman, T. C. Selenium biochemistry. *Science* 183: 915 (1974).
  85. Johnson, J. L., Cohen, H. J. and Rajagopalan, K. V. Studies of vanadium toxicity in the rat. Lack of correlation with molybdenum utilization. *Biochem. Biophys. Res. Commun.* 56: 940 (1974).
  86. Cantley, L. C., and Aisen, P. The fate of cytoplasmic vanadium. *J. Biol. Chem.* 254: 1781 (1979).
  87. Erdmann, E., Krawietz, W., Philipp, G., Hackbarth, I., Schmitz, W., Scholz, H., and Crane, F. L. Purified cardiac cell membranes with high (Na<sup>+</sup> + K<sup>+</sup>)ATPase activity contain significant NADH-vanadate reductase activity. *Nature* 282: 335 (1979).
  88. Cantley, L. C., Ferguson, J. H., and Kustin, K. Norepinephrine complexes and reduces vanadium (V) to reverse vanadate inhibition of the (Na, K)-ATPase. *J. Am. Chem. Soc.* 100: 5210 (1978).
  89. Gutierrez, P. L., Sarna, T., and Swartz, H. M. Experimental considerations in biological ESR studies. II. Chromium-tissue complexes detected by electron spin resonance. *Phys. Med. Biol.* 21: 949 (1976).
  90. Levis, A. G., Buttignol, M., Bianchi, V., and Sponza, G. Effects of potassium dichromate on nucleic acid and protein syntheses and on precursor uptake in BHK fibroblasts. *Cancer Res.* 38: 110 (1978).
  91. Pearson, H. A. The binding of <sup>51</sup>Cr to hemoglobin. I. *In vitro* studies. *Blood* 22: 218 (1963).
  92. Koutras, G. A., Hattori, M., Schneider, A. S., Ebaugh, F. G., and Valentine, W. N. Studies on chromated erythrocytes. Effect of sodium chromate on erythrocyte glutathione reductase. *J. Clin. Invest.* 43: 323 (1964).
  93. Dixon, J. R., Lowe, D. B., Richards, D. E., Cralley, L. J., and Stokinger, H. E. The role of trace metals in chemical carcinogenesis. Asbestos cancers. *Cancer Res.* 30: 1068 (1974).
  94. Calop, J., Burckhart, M.-F., and Fontanges, R. Étude de l'influence des éléments traces sur l'hydroxylation du benzo(a)pyrene. *Eur. J. Toxicol.* 9: 271 (1976).
  95. Thomson, R., Webster, I., and Kilroe-Smith, T. A. The metabolism of benzo(a)pyrene in rat liver microsomes: The effect of asbestos associated metal ions and pH. *Environ. Res.* 7: 149 (1974).
  96. Gruber, J. E., and Jennette, K. W. Metabolism of the carcinogen chromate by rat liver microsomes. *Biochem. Biophys. Res. Commun.* 82: 700 (1978).
  97. Jennette, K. W. Chromate metabolism in liver microsomes. *Biol. Trace Element Res.* 1: 55 (1979).
  98. Garcia, J. D., and Jennette, K. W. Electron-transport cytochrome P-450 system is involved in the microsomal metabolism of the carcinogen chromate. *J. Inorg. Biochem.*, in press.
  99. Sugiura, M., Iwasaki, K., and Kato, R. Reduced nicotinamide adenine dinucleotide-dependent reduction of tertiary amine N-oxide by liver microsomal cytochrome P-450. *Biochem. Pharmacol.* 26: 489 (1977).
  100. Baetjer, A. M., Damron, C. M., Clark, J. H., and Budacz, V. Reaction of chromium compounds with body tissues and their constituents. *Arch. Ind. Health* 12: 258 (1955).
  101. De Pamphilis, M. L., and Cleland, W. W. Preparation and properties of chromium (III)-nucleotide complexes for use in the study of enzyme mechanisms. *Biochemistry* 12: 3714 (1973).
  102. Dunaway-Mariano, D., and Cleland, W. W. Preparation and properties of chromium (III) adenosine 5'-triphosphate, chromium (III) adenosine 5'-diphosphate, and related chromium (III) complexes. *Biochemistry* 19: 1496 (1980).
  103. Dunaway-Mariano, D., and Cleland, W. W. Investigations of substrate specificity and reaction mechanism of several kinases using chromium (III) adenosine 5'-triphosphate and chromium (III) adenosine 5'-diphosphate. *Biochemistry* 19: 1506 (1980).
  104. Danenberg, K. D., and Cleland, W. W. Use of chromium-adenosine triphosphate and lyxose to elucidate the kinetic mechanism and coordination state of the nucleotide substrate for yeast hexokinase. *Biochemistry* 14: 28 (1975).
  105. Kopkins, L. L. and Schwarz, K. Chromium (III) binding to serum proteins, specifically siderophilin. *Biochim. Biophys. Acta* 90: 484 (1964).
  106. Sabbioni, E., and Marafante, E. Metabolic patterns of vanadium in the rat. *Bioinorg. Chem.* 9: 389 (1978).
  107. Chasteen, N. D. Human serotransferrin: structure and function. *Coord. Chem. Rev.* 22: 1 (1977).
  108. Goldstein, J. L., Anderson, R. G. W., and Brown, M. S. Coated pits, coated vesicles, and receptor-mediated endocytosis. *Nature* 279: 679 (1979).
  109. Kornfeld, S. The effect of metal attachment to human apotransferrin on its binding to reticulocytes. *Biochim. Biophys. Acta* 194: 25 (1969).
  110. Bautista, E. M., and Alexander, M. Reduction of inorganic compounds by soil microorganisms. *Proc. Soil Sci. Soc. Am.* 36: 918 (1972).
  111. Bray, R. C. Molybdenum-containing enzymes. *J. Less-Common Met.* 54: 527 (1977).
  112. Johnson, J. L., and Rajagopalan, K. V. Experimental molybdenum deficiency in the rat. In: *Molybdenum in the Environment, Vol. 1. The Biology of Molybdenum*, W. R. Chappell and K. K. Petersen, Eds., Marcel Dekker, New York, 1976, pp. 201-219.
  113. Lener, J., and Bibr, B. Biliary excretion and tissue distribution of penta- and hexavalent molybdenum in rats. *Toxicol. Appl. Pharmacol.* 51: 259 (1979).
  114. Dick, A. T. Molybdenum and copper relationships in animal nutrition. In: *A Symposium on Inorganic Nitrogen Metabolism: Function of Metallo-flavoproteins*, W. D. McElroy and B. Glass, Eds., Johns Hopkins Press, Baltimore, 1956, pp. 445-473.
  115. Kselikova, M., Bibr, B., Marik, T., and Lener, J. Conditions of molybdenum-99-labeling of proteins for human erythrocyte membrane. *Radiobiol. Radiother.* 20: 701 (1979).
  116. Kselikova, M., Bibr, B., and Lener, J. Interaction of alpha<sub>2</sub>-macroglobulin with molybdenum in human and rat

- serum. *Physiol. Bohemoslov.* 26: 573 (1977).
117. Ganther, H. E. Enzymic synthesis of dimethylselenide from sodium selenite in mouse liver extracts. *Biochemistry* 5: 1089 (1966).
  118. Hsieh, H. S., and Ganther, H. E. Acid-volatile selenium formation catalyzed by glutathione reductase. *Biochemistry* 14: 1632 (1975).
  119. McConnell, K. P., and Portman, O. W. Excretion of dimethyl selenide by the rat. *J. Biol. Chem.* 195: 277 (1952).
  120. Palmer, I. S., Gunsalus, R. P., Halverson, A. W., and Olson, O. E. Trimethylselenonium ion as a general excretory product from selenium metabolism in the rat. *Biochim. Biophys. Acta* 208: 260 (1970).
  121. Diplock, A. T., and Lucy, J. A. The biochemical modes of action of vitamin E and selenium: A hypothesis. *FEBS Lett.* 29: 205 (1973).
  122. Diplock, A. T., Baum, H., and Lucy, J. A. Effect of vitamin E on the oxidation state of selenium in rat liver. *Biochem. J.* 123: 721 (1971).
  123. Burk, R. F. Effect of dietary selenium level on <sup>76</sup>Se binding to rat plasma proteins. *Proc. Soc. Exptl. Biol. Med.* 143: 719 (1973).
  124. Germain, G. S., and Arneson, R. M. Selenium induced glutathione peroxidase activity in mouse neuroblastoma cells. *Biochem. Biophys. Res. Commun.* 79: 119 (1977).
  125. White, C. L. and Hoekstra, W. G. The metabolism of selenite and selenomethionine in mouse fibroblasts grown in tissue culture. *Proc. N. Zealand Semin. Trace Elem. Health*, 2nd., 140 (1979).
  126. Forstrom, J. W., Zakowski, J. J., and Tappel, A. L. Identification of the catalytic site of rat liver glutathione peroxidase as selenocysteine. *Biochemistry* 17: 2639 (1978).
  127. Chen, C.-S., and Stadtman, T. C. Selenium-containing tRNAs from *Clostridium sticklandii*: cochromatography of one species with L-prolyl-tRNA. *Proc. Natl. Acad. Sci. (U.S.A.)* 77: 1403 (1980).
  128. Paulson, G. D., Baumann, C. A., and Pope, A. L. Metabolism of <sup>76</sup>Se-selenite, <sup>76</sup>Se-selenate, <sup>76</sup>Se-selenomethionine, and <sup>35</sup>S-sulfate by rumen microorganisms in vitro. *J. Anim. Sci.* 27: 497 (1968).
  129. Fleming, R. W., and Alexander, M. Dimethylselenide and dimethyltelluride formation by a strain of *Penicillium*. *Appl. Microbiol.* 24: 424 (1972).
  130. Reamer, D. C., and Zoller, W. H. Selenium biomethylation products from soil and sewage sludge. *Science* 208: 500 (1980).
  131. Ginsburg, J. M. Renal mechanism for excretion and transformation of arsenic in the dog. *Am. J. Physiol.* 208: 832 (1965).
  132. Peoples, S. A. Arsenic toxicity in cattle. *Ann. N. Y. Acad. Sci.* 111: 644 (1964).
  133. NAS Committee on Medical and Biologic Effects of Environmental Pollutants. Arsenic, National Academy of Sciences, Washington, D.C., 1977, pp. 80-116.
  134. Lakso, J. U. and Peoples, S. A. Methylation of inorganic arsenic by mammals. *J. Agr. Food Chem.* 23: 674 (1975).
  135. Winkler, W. O. Identification and estimation of the arsenic residue in livers of rats ingesting arsenicals. *J. Assoc. Off. Anal. Chem.* 45: 80 (1962).
  136. Lanz, H., Jr., Wallace, P. W., and Hamilton, J. G. The metabolism of arsenic in laboratory animals using <sup>74</sup>As as a tracer. *Univ. Calif. Pub. Pharmacol.* 2: 263 (1950).
  137. Stevens, J. T., Hall, L. L., Farmer, J. D., DiPasquale, L. C., Chernoff, N., and Durham, W. F. Disposition of <sup>14</sup>C and/or <sup>74</sup>As-cacodylic acid in rats after intravenous, intratracheal, or peroral administration. *Environ. Health Perspect.* 19: 151 (1977).
  138. Braman, R. C., and Foreback, C. C. Methylated forms of arsenic in the environment. *Science* 182: 1247 (1973).
  139. Smith, T. J., Crecelius, E. A. and Reading, J. C. Airborne arsenic exposure and excretion of methylated arsenic compounds. *Environ. Health Perspect.* 19: 89 (1977).
  140. Crecelius, E. A. Changes in the chemical speciation of arsenic following ingestion by man. *Environ. Health Perspect.* 19: 147 (1977).
  141. Cullen, W. R., Froese, C. L., Lui, A., McBride, B. C., Patmore, D. J., and Reimer, M. The aerobic methylation of arsenic by microorganisms in the presence of L-methionine-methyl-d-3. *J. Organomet. Chem.* 139: 61 (1977).
  142. McBride, B. C., and Wolfe, R. S. Biosynthesis of dimethylarsine by methanobacterium. *Biochemistry* 10: 4312 (1971).
  143. Johnson, D. L. Bacterial reduction of arsenate in sea water. *Nature* 240: 44 (1972).
  144. Irgolic, K. J., Woolson, E. A., Stockton, R. A., Newman, R. D., Bottino, N. R., Zingaro, R. A., Kearney, P. C., Pyles, R. A., Maeda, S., McShane, W. J., and Cox, E. R. Characterization of arsenic compounds formed by *Daphnia magna* and *Tetraselmis chuii* from inorganic arsenic. *Environ. Health Perspect.* 19: 61 (1977).
  145. Chan, T.-L., Thomas, B. R., and Wadkins, C. L. The formation and isolation of an arsenylated component of rat liver mitochondria. *J. Biol. Chem.* 244: 2883 (1969).
  146. Löfroth, G., and Ames, B. N. Mutagenicity of inorganic compounds in *Salmonella typhimurium*: arsenic, chromium and selenium. *Mutat. Res.* 53: 65 (1978).
  147. Löfroth, G. The mutagenicity of hexavalent chromium is decreased by microsomal metabolism. *Naturwiss.* 65: S207 (1978).
  148. Noda, W., Takano, T., and Sakurai, H. Mutagenic activity of selenium compounds. *Mutat. Res.* 66: 175 (1979).
  149. Nishioka, H. Mutagenic activities of metal compounds in bacteria. *Mutat. Res.* 31: 185 (1975).
  150. Kanematsu, K., and Kada, T. Mutagenicity of metal compounds. *Mutat. Res.* 53: 207 (1978).
  151. Venitt, S., and Levy, L. S. Mutagenicity of chromates in bacteria and its relevance to chromate carcinogenesis. *Nature* 250: 493 (1974).
  152. Green, M. H. L., and Muriel, W. J. Mutagen testing using Trp<sup>+</sup> reversion in *Escherichia coli*. *Mutat. Res.* 38: 3 (1976).
  153. Rossman, T., Meyn, M. S., and Troll, W. Effects of sodium arsenite on the survival of UV-irradiated *Escherichia coli*: inhibition of a recA dependent function. *Mutat. Res.* 30: 157 (1975).
  154. Rossman, T. G., Meyn, M. S. and Troll, W. Effects of arsenite on DNA repair in *Escherichia coli*. *Environ. Health Perspect.* 19: 229 (1977).
  155. Casto, B. C., Meyers, J., and DiPaolo, J. A. Enhancement of viral transformation for evaluation of the carcinogenic or mutagenic potential of inorganic metal salts. *Cancer Res.* 39: 193 (1979).
  156. Fradkin, A., Janoff, A., Lane, B. P., and Kuschner, M. *In vitro* transformation of BHK21 cells grown in the presence of calcium chromate. *Cancer Res.* 35: 1058 (1975).
  157. DiPaolo, J. A., and Casto, B. C. Quantitative studies of *in vitro* morphological transformation of syrian hamster cells by inorganic metal salts. *Cancer Res.* 39: 1008 (1979).
  158. Umeda, M., and Nishimura, M. Inducibility of chromosomal aberrations by metal compounds in cultured mammalian cells. *Mutat. Res.* 67: 221 (1979).
  159. MacCrae, W. D., Whiting, R. F. and Stich, H. F. Sister chromatid exchanges induced in cultured mammalian cells by chromate. *Chem.-Biol. Interact.* 26: 281 (1979).
  160. Majone, F., and Levis, A. G. Chromosomal aberrations

- and sister-chromatid exchanges in Chinese hamster cells treated *in vitro* with hexavalent chromium compounds. *Mutat. Res.* 67: 231 (1979).
161. Tsuda, H., and Kato, K. Chromosomal aberrations and morphological transformation in hamster embryonic cells treated with potassium dichromate *in vitro*. *Mutat. Res.* 46: 87 (1977).
  162. Zanzoni, F., and Jung, E. G. Arsenic elevates the sister chromatid exchange (SCE) rate in human lymphocytes *in vitro*. *Arch. Dermatol. Res.* 267: 91 (1980).
  163. Paton, G. R., and Allison, A. C. Chromosome damage in human cell cultures induced by metal salts. *Mutat. Res.* 16: 332 (1972).
  164. Nakamuro, K., Yoshikawa, K., Sayato, Y., Kurata, H., Tonomura, M., and Tonomura, A. Studies on selenium-related compounds. V. Cytogenetic effect and reactivity with DNA. *Mutat. Res.* 40: 177 (1976).
  165. Ray, J. H., and Altenburg, L. C. Sister-chromatid exchange induction by sodium selenite: Dependence on the presence of red blood cells or red blood cell lysate. *Mutat. Res.* 54: 343 (1978).
  166. Lo, L. W., Koropatnick, J., and Stich, H. F. The mutagenicity and cytotoxicity of selenite, "activated" selenite and selenate for normal and DNA repair-deficient human fibroblasts. *Mutat. Res.* 49: 305 (1978).
  167. Tkeshelashvili, L. K., Shearman, C. W., Koplitz, R. M., Zakour, R. A., and Loeb, L. A. Fidelity of DNA synthesis: Effects of As, Se and Co. *Proc. Am. Assoc. Cancer Res.* 20: 263 (1979).
  168. Sirover, M. A., and Loeb, L. A. Infidelity of DNA synthesis *in vitro*: Screening for potential metal mutagens or carcinogens. *Science* 194: 1434 (1976).
  169. Demerec, M., and Hanson, J. Mutagenic action of manganese chloride. *Cold Spring Harbor Symp. Quant. Biol.* 16: 215 (1951).
  170. Roberts, R. B., and Aldous, E. Manganese metabolism of *Escherichia coli* as related to its mutagenic action. *Cold Spring Harbor Symp. Quant. Biol.* 16: 229 (1951).
  171. Dube, D. K., and Loeb, L. A. Manganese as a mutagenic agent during *in vitro* DNA synthesis. *Biochem. Biophys. Res. Commun.* 67: 1041 (1975).
  172. Sirover, M. A., and Loeb, L. A. Metal-induced infidelity during DNA synthesis. *Proc. Natl. Acad. Sci. (U.S.)* 73: 2331 (1976).
  173. Hsu, M.-T., and Berg, P. Altering the specificity of restriction endonuclease: Effect of replacing  $Mg^{2+}$  with  $Mn^{2+}$ . *Biochemistry* 17: 131 (1978).
  174. Falchuk, K. H., Hardy, C., Alpino, L., and Vallee, B. L. RNA metabolism, manganese, and RNA polymerases of zinc-sufficient and zinc-deficient *Euglena gracilis*. *Proc. Natl. Acad. Sci. (U.S.)* 75: 4175 (1978).
  175. Paddock, G. V., Heindell, H. C., and Salsler, W. Deoxysubstitution in RNA by RNA polymerase *in vitro*: A new approach to nucleotide sequence determination. *Proc. Natl. Acad. Sci. (U.S.)* 71: 5017 (1974).
  176. Stavrianopoulos, J. G., Gambino-Giuffrida, A., and Chargaff, E. Ribonuclease H of calf thymus: Substrate specificity, activation, inhibition. *Proc. Natl. Acad. Sci. (U.S.)* 73: 1087 (1976).
  177. Currin, A., and Kibroe-Smith, T. A. The effect of manganese and ferric ions on the *in vitro* formation of dihydrodihydroxy metabolites of benzo[a]pyrene. *Chem. Biol. Interact.* 19: 259 (1978).
  178. Whiting, R. F., Wei, L., and Stich, H. F. Enhancement by transition metals of unscheduled DNA synthesis induced by isoniazid and related hydrazines in cultured normal and xeroderma pigmentosum human cells. *Mutat. Res.* 62: 505 (1979).
  179. Thomas, H. F., Herriott, R. M., Hahn, B. S., and Wang, S. Y. Thymine hydroperoxide as a mediator in ionizing radiation mutagenesis. *Nature* 259: 341 (1976).
  180. Stich, H. F., Wei, L., and Whiting, R. F. Enhancement of the chromosome-damaging action of ascorbate by transition metals. *Cancer Res.* 39: 4145 (1979).
  181. Hoffman, D. J., and Niyogi, S. K. Metal mutagens and carcinogens affect RNA synthesis rates in a distinct manner. *Science* 198: 513 (1977).
  182. Buttlair, D. H., Czuba, B. A., Stevens, T. H., Lee, Y. C., and Himes, R. H. Manganese ion binding to tubulin. *J. Biol. Chem.* 255: 2164 (1980).
  183. Boadle-Biber, M. C. Decrease in the activity of tryptophan hydroxylase from slices of rat brain stem incubated in a low calcium or a calcium-free manganese-substituted medium. *Biochem. Pharmacol.* 28: 3487 (1979).
  184. Couturier, E., and Malaisse, W. J. Cobalt-induced inhibition of ionophore-mediated calcium translocation. *IRCS Med. Sci.: Libr. Compend.* 8: 72 (1980).
  185. Phillips, C. E. Intracellularly injected cobaltous ions accumulate at synaptic densities. *Science* 207: 1477 (1980).
  186. Thompson, S. H. Three pharmacologically distinct potassium channels in molluscan neurons. *J. Physiol.* 265: 465 (1977).
  187. Baker, P. F., Meves, H., and Ridgway, E. B. Effects of manganese and other agents on the calcium uptake that follows depolarization of squid axons. *J. Physiol.* 231: 511 (1973).
  188. Magleby, K. L., and Weinstock, M. M. Nickel and calcium ions modify the characteristics of the acetylcholine receptor-channel complex at the frog neuromuscular junction. *J. Physiol.* 299: 203 (1980).
  189. Johnson, R. S. and Schachman, H. K. Propagation of conformational changes in Ni (II)-substituted aspartate transcarbamoylase: Effect of active-site ligands on the regulatory chains. *Proc. Natl. Acad. Sci. (U.S.)* 77: 1995 (1980).
  190. Bauer, R., Limkilde, P., and Johansen, J. T. Low and high pH form of cadmium carbonic anhydrase determined by nuclear quadrupole interaction. *Biochemistry* 15: 334 (1976).
  191. McQuate, R. S. Carbonic anhydrase and metalloderivatives. *J. Chem. Educ.* 54: 645 (1977).
  192. Sytkowski, A. J., and Vallee, B. L. The catalytic metal atoms of cobalt substituted liver alcohol dehydrogenase. *Biochem. Biophys. Res. Commun.* 67: 1488 (1975).
  193. Anderson, R. A., Bosron, W. F., Kennedy, F. S., Vallee, B. L. Role of magnesium in *Escherichia coli* alkaline phosphatase. *Proc. Natl. Acad. Sci. (U.S.)* 72: 2989 (1975).
  194. Zukin, R. S., and Hollis, D. P. Role of metal ions in *Escherichia coli* alkaline phosphatase. *J. Biol. Chem.* 250: 835 (1975).
  195. Thompson, G. A., and Carpenter, F. H. Leucine aminopeptidase (bovine lens). The relative binding of cobalt and zinc to leucine aminopeptidase and the effect of cobalt substitution on specific activity. *J. Biol. Chem.* 251: 1618 (1976).
  196. Auld, D. S. The role of zinc in biochemical processes. In: (ACS Adv. Chem. Ser., 172) American Chemical Society, Washington, D.C., 1979, pp. 112-133.
  197. Kozloff, L. M., and Lute, M. Zinc, an essential component of the baseplates of T-even bacteriophages. *J. Biol. Chem.* 252: 7715 (1977).
  198. Kozloff, L. M. Properties of T4D bacteriophage grown in synthetic media containing  $Zn^{2+}$ ,  $Co^{2+}$  or  $Ni^{2+}$ . *J. Biol. Chem.* 253: 1059 (1978).
  199. Malmquist, J., Israelsson, B., and Ljungqvist, U. Inhibi-

- tion of human liver membrane adenylate cyclase by zinc ions. *Horm. Metab. Res.* 11: 530 (1979).
200. Panara, F., Cirotto, C., Arangi, I., and Barberini, L. Localization of acid phosphatases in the cell fractions of chick liver. *Experientia* 36: 119 (1980).
  201. Korant, B. D., and Butterworth, B. E. Inhibition by zinc of rhinovirus protein cleavage: Interaction of zinc with capsid polypeptides. *J. Virol.* 18: 298 (1976).
  202. Gray, R. D. The effect of H<sup>+</sup>, inositol hexaphosphate, and Zn (II) on the tetramer-dimer equilibrium of liganded hemoglobin. *J. Biol. Chem.* 255: 1812 (1980).
  203. Baccetti, B., Pallini, V., and Burrini, A. G. The accessory fibers of sperm tail II. Their role in binding zinc in mammals and cephalopods. *J. Ultrastruct. Res.* 54: 261 (1976).
  204. Karin, M., and Herschman, H. R. Dexamethasone stimulation of metallothionein synthesis in HeLa cell cultures. *Science* 204: 176 (1979).
  205. Omaye, S. T., and Tappel, A. L. Effect of cadmium chloride on the rat testicular soluble selenoenzyme, glutathione peroxidase. *Res. Commun. Chem. Pathol. Pharmacol.* 12: 695 (1975).
  206. Carson, S. D., and Konigsberg, W. H. Cadmium increases tissue factor (coagulation factor III) activity by facilitating its reassociation with lipids. *Science* 208: 307 (1980).
  207. Suresh, J. K., Bradford, N. M. and McGivan, J. D. Inhibition of glutamine synthetase activity by manganous ions in a cytosol extract of rat liver. *Biochem. J.* 184: 477 (1979).
  208. Burr, R. G. Inhibition of urease by miscellaneous ions and compounds. Implications for the therapy of infection-induced urolithiasis. *Invest. Urol.* 15: 180 (1977).
  209. Siegel, G. J., and Fogt, S. K. Lead ion activates phosphorylation of electroplax Na<sup>+</sup>- and K<sup>+</sup>-dependent adenosine triphosphatase [(NaK)-ATPase] in the absence of sodium ion. *Arch. Biochem. Biophys.* 174: 744 (1976).
  210. Shamoo, A. E., MacLennan, D. H., and Eldefrawi, M. E. Differential effects of mercurial compounds on excitable tissues. *Chem.-Biol. Interact.* 12: 41 (1976).
  211. Farkas, W. R., and Stanawitz, T. Effects of plumbous ion on guanine metabolism. *J. Inorg. Biochem.* 11: 31 (1979).
  212. Hammond, P. B. Exposure of humans to lead. *Ann. Rev. Pharmacol. Toxicol.* 17: 197 (1977).
  213. Arcuri, E. J., and Ehrlich, H. L. Cytochrome involvement in Mn (II) oxidation by two marine bacteria. *Appl. Environ. Microbiol.* 37: 916 (1979).
  214. Fee, J. A., Shapiro, E. R., and Moss, T. H. Direct evidence for manganese (III) binding to the manganese-superoxide dimutase of *Escherichia coli* B. *J. Biol. Chem.* 251: 6157 (1976).
  215. Clarkson, T. W. Factors involved in heavy metal poisoning. *Fed. Proc.* 36: 1634 (1977).
  216. Wood, J. M., Fanchiang, Y.-T., and Craig, P. J. Some bio-inorganic chemical reactions of environmental significance. In: *Trace Metals in Health and Disease*, N. Kharasch, Ed., Raven Press, New York, 1979, pp. 43-53.