



## The role of self-reported olfactory and gustatory dysfunction as a screening criterion for suspected COVID-19

Liang En Wee<sup>1,2</sup> · Yvonne Fu Zi Chan<sup>2</sup> · Neville Wei Yang Teo<sup>3</sup> · Benjamin Pei Zhi Cherng<sup>2</sup> · Siew Yee Thien<sup>2</sup> · Hei Man Wong<sup>2</sup> · Limin Wijaya<sup>2</sup> · Song Tar Toh<sup>3</sup> · Thuan Tong Tan<sup>2</sup>

Received: 14 April 2020 / Accepted: 18 April 2020 / Published online: 24 April 2020  
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

**Keywords** Anosmia · Ageusia · COVID-19

Dear Editor,

We read with interest the paper written by Lechien et al. [1] describing a multi-centre cohort of COVID-19 patients, in which 85.6% had olfactory or taste disorders (OTD). If the association between COVID-19 and OTD is not recognized, the diagnosis may be missed. However, while a large proportion of COVID-19 patients had OTD, [1–3] the actual specificity as a criterion for screening suspected COVID-19 cases is unknown, as other viral infections may also cause OTD. Other studies have reported patients presenting with acute anosmia with or without upper respiratory tract symptoms, [4, 5] but not all patients were tested for COVID-19 [4]. Additionally, information on OTD in Asian cohorts is lacking [1]. In Singapore, the first case of COVID-19 was reported in end-January 2020. At our institution, the largest acute hospital in Singapore, from end-March 2020 all new admissions were screened for self-reported OTD at emergency department (ED) triage. All patients reporting new-onset OTD were admitted to exclude COVID-19. We

report our experience using self-reported OTD as a screening criterion for suspected COVID-19.

Over a 2-week period from 26 March–10 April 2020, given ongoing community transmission and the difficulty in distinguishing COVID-19 from ordinary viral infection, a questionnaire including respiratory symptoms, self-reported OTD, and travel and epidemiological risk factors was administered at ED triage to risk-stratify admissions. Suspect case criteria for COVID-19 were defined as the presence of respiratory symptoms and suspicious epidemiological links or travel history; or new-onset OTD. Testing was not limited to hospitalised inpatients; as part of the national strategy of containment, all patients who fulfilled suspect case criteria for COVID-19 were tested via real-time reverse transcription PCR (rRT-PCR) of oropharyngeal swabs, even if hospitalization was not otherwise required. Inpatient, if patients had respiratory symptoms and a viral prodrome, or OTD, oropharyngeal specimens were also tested for a routine panel of respiratory viruses. As this study was based on aggregated surveillance data, ethics approval was not required under our hospital's Institutional Review Board guidelines.

Over the study period, a total of 870 patients fulfilled suspect case criteria for COVID-19 at ED triage. A minority (5.05%, 44/870) presented with OTD. The majority of suspects (65.3%, 568/870) were well and discharged to self-isolate while awaiting results. Amongst suspected COVID-19 cases, 154 patients (17.9%, 154/870) tested positive. Of those, roughly one-fifth (22.7%, 35/154) had OTD. The presence of OTD had high specificity as a screening criterion for COVID-19 (98.7%, 95% CI 97.6–99.4%), but lower sensitivity (22.7%, 95% CI 16.4–30.2%). This was roughly equivalent to the specificity and sensitivity of a history of close contact with a confirmed COVID-19 case (specificity: 94.8%, 95% CI 93.0–96.3%; sensitivity: 27.3%, 95% CI 20.4–35.0%), Supplementary Table 1. Amongst the 35

---

This comment refers to the article available online at <https://doi.org/10.1007/s00405-020-05965-1>.

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s00405-020-05999-5>) contains supplementary material, which is available to authorized users.

---

✉ Liang En Wee  
ian.wee@mohh.com.sg

- 1 Singhealth Infectious Diseases Residency, Singapore, Singapore
- 2 Department of Infectious Diseases, Singapore General Hospital, Singapore, Singapore
- 3 Department of Otolaryngology, Singapore General Hospital, Singapore, Singapore

COVID-19 positives with concomitant OTDs, three presented with isolated anosmia without other symptoms. Fever was the most common concomitant symptom (21/35, 60.0%), followed by cough (10/35, 28.5%) and rhinorrhea (10/35, 28.5%). Over the same period, a total of 71 admitted inpatients tested positive for other respiratory viruses, of which the most common was rhinovirus (53.5%, 38/71); followed by influenza (21.1%, 15/71), adenoviruses (8.5%, 6/71), other coronaviruses (7.0%, 5/71). Only 2.8% (2/71) self-reported OTD (one rhinovirus, one human coronavirus 229E). Amongst patients with PCR-proven acute respiratory viral infections, those positive for COVID-19 had higher odds of OTD compared to those positive for other respiratory viruses (odds ratio, OR = 10.14, 95% CI 2.37–43.49,  $p < 0.001$ ).

In conclusion, self-reported OTD had high specificity as a screening criterion for COVID-19 in an Asian cohort. Patients with COVID-19 appeared to have higher odds of OTD compared to those positive for other respiratory viruses. Routine screening in patients with new-onset OTD can improve case detection during a COVID-19 outbreak.

**Funding** This work was not grant-funded.

## Compliance with ethical standards

**Conflict of interest** The authors report no conflicts of interest.

## References

1. Lechien JR, Chiesa-Estomba CM, De Siati DR et al (2020) Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study. *Eur Arch Otorhinolaryngol*. <https://doi.org/10.1007/s00405-020-05965-1> (**Epub ahead of print**)
2. Giacomelli A, Pezzati L, Conti F et al (2020) Self-reported olfactory and taste disorders in SARS-CoV-2 patients: a cross-sectional study. *Clin Infect Dis*. <https://doi.org/10.1093/cid/ciaa330>
3. Vaira LA, Salzano G, Deiana G, De Riu G (2020) Anosmia and ageusia: common findings in COVID-19 patients. *Laryngoscope*. <https://doi.org/10.1002/lary.28692> (**Epub ahead of print**)
4. Hopkins C, Surda P, Kumar N (2020) Presentation of new onset anosmia during the COVID-19 pandemic. *Rhinology*. <https://doi.org/10.4193/Rhin20.116> (**Epub ahead of print**)
5. Gane SB, Kelly C, Hopkins C (2020) Isolated sudden onset anosmia in COVID-19 infection: a novel syndrome? *Rhinology*. <https://doi.org/10.4193/Rhin20.114> (**Epub ahead of print**)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.