

---

# The Role of Smad Proteins for Development, Differentiation and Dedifferentiation of Neurons

---

Uwe Ueberham and Thomas Arendt

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/54532>

---


## 1. Introduction

The development of the nervous system, neuralization of ectodermal cells, specification of cell types as well as generation of neurological diseases are closely linked to Smad proteins, which play a central role by integrating TGF $\beta$  and BMP signalling with other essential pathways. Due to new findings on Smad activity in neurons and the nervous system, which comprises new roles for brain plasticity and functions, independent of the canonical signalling pathways, we reconsider their relevance for neuronal differentiation and dedifferentiation processes and as therapeutic targets for treatment of neurological diseases. In this chapter we develop a view at Smad molecules, which attributes them a basic significance and allow proving their specific contextual molecular, cellular and tissue relationships. In order to facilitate the understanding of the complex Smad network in the nervous system an overview of the canonical Smad signalling pathway is briefly summarized in the following paragraph.

Smads are phylogenetic old proteins, which are mediating intracellular signalling of the large group of soluble TGF $\beta$  ligands (Figure 1), containing transforming growth factor  $\beta$ s (TGF $\beta$ s), bone morphogenic proteins (BMPs), growth and differentiation factors (GDFs), Müllerian inhibitory factors (MISs), activins and inhibins [4]. Ligand binding to activated heteromeric receptor complexes, recruited from seven type I and five type II serine/threonine receptors, results in the specific phosphorylation of receptor-associated Smads (R-Smads) at two C-terminal serine residues.

Activin, nodal and TGF $\beta$  activate R-Smad2 and 3, while BMP acts by R-Smad1,5 and 8 phosphorylation. In the cytoplasm non-phosphorylated R-Smads are sequestered by interacting with specific retention proteins e.g. SARA (Smad anchor for receptor activation) [5], endofin [8], tubulin [3], actin, myosin [11] or filamin [12]. Inhibitory Smad(I-Smad)6 and 7 negatively regulate R-Smad signalling by competing for binding to activated type I receptor and inhibiting R-Smad phosphorylation. I-Smads can also prevent R-Smad complexing to the

co-operating Smad(Co-Smad)4, which is required for the nuclear translocation of activated R-Smads to subsequently control Smad sensitive promoter activity. Moreover, I-Smads support recruitment of HECT-type E3-ubiquitin ligases Smurf1/2, which allow type I receptor ubiquitination and its degradation. Members of HECT-type and RING-type E3 ubiquitin ligases have also been implicated in Smad degradation [13]. Regulating Smad activity allows the control of highly complex developmental networks, e.g. the patterning of ventro-lateral mesoderm, the decisive development of epidermal/neural cell lineages including the induction and establishment of neural plate border, the dorso-ventral patterning of the neural tube, or the migration of neural crest cells [18-21].



**Figure 1. Synopsis of canonical Smad signalling.** The upper panel shows the binding of ligands belonging to the TGFβ superfamily to their serine/threonine kinase receptors types I and II, leading to intracellular activation of R-Smads by receptor type I induced phosphorylation of the C-terminal Smad motif SSXS. The left and right receptor complexes represent the TGFβ/activin/nodal linked Smad2/3 signalling and the BMP/GDF linked Smad1/5/8 signalling, respectively. The middle panel also displays a set of relevant binding partners of Smad proteins and shows the generation of heterotrimer or heterodimer Smad complexes consisting of C-terminally phosphorylated R-Smads and Co-Smad4, which are transported into the nucleus to exert their transcriptional activity by binding to susceptible promoter sequences (a). In (b) and (c) examples of the agonist-induced Smad linker phosphorylation mediated by cyclin-dependent kinases are shown, which increases Smad transcriptional activity prior to signal termination and Smad degradation [14, 17]. In contrast, the MAPK triggered linker phosphorylation of Smads in cytoplasm diminishes the receptor type I responsible Smad phosphorylation and favours the cytoplasmic retention and degradation of Smad. More information is provided in the main text. *Note:* Numbers in square brackets indicate relevant references included in the reference list provided at the chapters end. *Abbreviations:* CDK, cyclin-dependent kinase; dynein, dynein light chain km23-1 or km23-2; MAPK, mitogen activated protein kinase; MT, microtubuli; Nedd4L, neuronal precursor cell expressed developmentally down-regulated 4-like ubiquitin ligase; Nuc214, nucleoporin 214; SARA, Smad anchor for receptor activation; Smurf1, HECT-domain ubiquitin ligase Smurf1.

In R-Smads a linker region, located between the highly conserved N-terminal MH1- and the C-terminal MH2 region (MH represents Mad homology), is rich in potential serine/threonine phosphorylation sites. Several kinases (e.g. MAPK, GSK, CDKs, CamKII, SGK1) can phosphorylate the linker region and determine cellular distribution or protein stability of Smads [14,23-26]. Linker phosphorylation by mitogen-activated protein kinases (MAPKs) restricts Smad1 activity by enabling Smurf1 binding, causing polyubiquitination and inhibiting interaction of Smad1 with nuclear translocation factor Nup214, thus leading to Smad1 degradation or cytoplasmic retention [15].

A similar mechanism acts on Smad2 and 3, where ubiquitination is controlled by ubiquitin ligase Nedd4L [17]. A sustained MAPK activation, as reported from dissociated embryonic ectodermal cells, phosphorylates Smad linker and interrupt a continuing BMP mediated Smad1 signalling finally resulting in neutralisation of dissociated *Xenopus* [27]. Remarkably, if linker phosphorylation is performed by extracellular signal regulated kinase (Erk) at nuclearly localized phosphorylated Smad, the duration of Smad targeted gene transcription can be increased [28]. Linker phosphorylation by CDK8 and CDK9 promotes both, the Smad transcriptional action, and the cooperation with YAP, an effector of the Hippo organ size control pathway, to finally suppress neural differentiation of stem cells [14]. The data demonstrate the particular role of the linker region for distinct Smad functions.

Numerous Smad binding proteins (e.g. Ski, SnoN) effectively influence essential steps during neural development mostly by repression of Smad activity [29-33]. The group of Smad interacting proteins (SIP), containing several zinc finger proteins, complex to Smad and can directly or indirectly regulate its transcription efficiency [34]. For example, the induction of neuroectoderm is dependent on SIP1 [35-37], which is also involved in the myelination process and the oligodendrocyte maturation [38].

## 2. The role of Smads during neural/non-neuronal development

### 2.1. Already early development requires Smad expression

During early ontogenesis, development and aging as well as prior to individual cell death Smad proteins differentially affect cellular function, depending on time and local partners. Already at very early stages of postnatal rat development Smads1-7 were detectable in the ovary [39]. At all stages of follicular development (e.g. in primordial, primary and secondary follicles) Smad proteins are present in oocytes as well as in granulosa cells and theca cells, though with different concentrations [39-41]. While Smad3,6 and 8 content did not change, the expression of Smad1,2,4,5,6 and 7 seems to be regulated in ovary [39]. In mice oocytes a clear and persistent nuclear localization of phosphorylated Smad1,5,9 and Smad2,3 demonstrates activated activin/GDF9/nodal/TGF $\beta$  and BMP/MIS pathways, respectively, and confirms oocytes as a major target for Smad signalling pathways [42]. Smad proteins play also important roles already during maturation of human oocytes [43]. In unfertilized oocytes, Smad2 and 3 and TGF $\beta$  receptors I and II are present [44]. After fertilization at the 4-cell and 8-cell stages Smad2 and 3 are also present, while neither of the TGF $\beta$  receptors is detectable. Later on in

the blastocyte stage Smad2 and 3 kept present and TGF $\beta$  receptor I again appears [44]. Human granulosa cells express Smad1-7 and 9, but Smad2,3, and 4 with the highest expression levels [45]. A mutual interaction between oocytes and granulosa cells is necessary for normal folliculogenesis. However, if Smad4 is deleted in granulosa cells infertility results, while deletion of Smad4 in oocytes only marginally effects the primordial follicle stage [46]. However, Smad2 reduction in zebrafish oocytes completely blocks activin A-induced oocyte maturation and a Smad2 knockdown decreases basal and hCG-induced oocyte maturation demonstrating a key role of Smads during oocyte maturation [47].

## 2.2. Gastrulation and neurulation

### 2.2.1. *Suppression of Smad signalling specifies neuroectoderm generation*

Initially, the ectoderm cells in *Xenopus* are pluripotent until gastrulation [48]. Early gastrulation steps are characterized by the formation of mesenchymal cells and the induction of mesoderm in the marginal-zone by an epithelial-to-mesenchymal transition (EMT) process, which involves activation of Smad2,3,4 signalling by TGF $\beta$ /nodal ligands after binding to corresponding receptors [49-53].

Mesoderm induction requires a subtle coordination of the canonical TGF $\beta$ /nodal signalling, with Smad4 as a key player. However, initially it was shown, that a simple ectopic Smad4 expression in the *Xenopus* animal cap was inefficient to induce mesoderm [54], due to its binding to ectodermin, a RING-type ubiquitin ligase, which is highly expressed in the animal cap [53]. Ectodermin limits Smad4 function via antagonizing nuclear accumulation of Smad4 and supporting Smad4 degradation by ubiquitin-proteasome pathway [53]. By this, ectodermin ensures that ectoderm cells do only develop to a mesodermal lineage, but also supports neural differentiation of ectoderm towards neuroectodermal fates by interfering with BMP signalling via limiting Smad4 function [53]. However, the stable presence of soluble BMP which prevents the execution of an intrinsic program of ectodermal cells to form neural tissue can also be undermined by the release of BMP antagonists as noggin, chordin, follistatin or cerberus, which however, allows creation and development of neural cells. This interrelation is basically reflected in the 'default model' of neural induction [55,56]. Confirming this model it was shown, that the simultaneous depletion of BMP2, BMP4 and BMP7 induces massive brain formation [57]. Accordingly, during early gastrulation-stage of chick embryo Smad1,5,8 signalling is undetectable [21] but is activated at neurula stage. So, phosphorylated Smad1 becomes detectable only at late gastrulation stages in the posterior territory of the embryo [21].

The default model has been further modified due to experiments showing that inhibition of BMP/Smad1 signalling, e.g. by Smad6, was insufficient to induce neural markers efficiently [58], while the additional suppression of Smad2 was sufficient [59]. Moreover, stimulation of Smad2 signalling blocks neural induction even at gastrula stages and indicate prospective neural cells to further undergo mesodermal and non-neural fates [59]. These data are supported by experiments in sea urchin embryos showing that Smad2 and 3 suppress neural differentiation in the oral ectoderm [60].

Remarkably, conversely, it has also been suggested from very early studies on Smad2<sup>+/-</sup>; Smad3<sup>-/-</sup> mutants in zebrafish showing anterior truncations [49], that Smad2 and Smad3, which are mainly effectors of TGFβ/nodal pathways, are also positively involved in neural development [61]. Dominant negative Smad3 mutants inhibit the expression of early neural markers sox2 and sox3 at the onset of gastrulation and lead to reduction of anterior and posterior neuroectodermal markers otx2 and hoxb1b, respectively, during late gastrulation. Accordingly, elevated Smad2 and Smad3 activities increased sox2 and sox3 expression, probably at least partly due to its positive impact on chordin expression, which is a BMP antagonist [61]. In mouse embryo BMP signalling also inhibits premature neural differentiation, but in contrast to *Xenopus* [62] and chick [63,64] FGF signalling seems additionally required to induce neural differentiation [65].

In summary, the data indicate that simultaneous suppression of both Smad1,5,8 and Smad2,3 pathways, e.g. suppression of mesoderm- and ectoderm-inducing Smad signals, respectively, are required for neural induction. In principle, the `dual Smad inhibition` method is adopted for induction of efficient neuralization of human (-induced) pluripotent stem cells (hiPSCs, hPSCs) [66,67].

### 2.2.2. Neural tube closure

To establish a precise borderline between neuroectoderm (containing low BMP2/4 levels) and non-neuroectoderm (containing high BMP2/4 levels) regions during early neurulation, specific strategies are required and examined in urchin embryos, where Fez, a zinc finger protein, drops down the BMP controlled pSmad1,5,8 levels [68]. It is probable that Fez acts as a transcriptional repressor of Smad or interferes with genes attenuating intracellular BMP signalling e.g. SIP1, Smurf1, Ski or SnoN.

Once the neuroectoderm forming is initiated a flat neural plate rolls up, the process of neural plate folding proceeds and is finished by neural tube closure [69]. The closure of neural tube is a complex process involving hinge points (HP), representing specialized neural plate cells. The cells of ventral midline/median hinge point (MHP) as well as both dorso-lateral hinge points (DLHPs) are affected by dynamic BMP signalling [70-73]. Folding of neural plate is connected with a two-dimensionally spatiotemporal gradient of phosphorylated Smad1,5,8 (pSmad1,5,8) [71,73]. While initially a mosaic labelling of pSmad1,5,8 was observed at the apical surface of the presumptive neuroectoderm later two intersecting Smad gradients were exerted: a lateromedial gradient with the lowest pSmad1,5,8 expression at the MHP and a second gradient along the apicobasal axis of the presumptive midbrain plate with mosaic expression in apical nuclei [71]. Some of the apical pSmad1,5,8 positive cells proliferate and thus support the neural fold elevation [71]. Disturbances of the phosphorylation state of Smad1,5,8 alter midbrain shape by multiple hinge-like invaginations but do not affect the ventral cell-fate specification [71]. However, the extent of Smad1,5,8 phosphorylation seems to control MHP formation during neurulation in vertebrates. The correct MHP formation is obviously directly linked to the pSmad1,5,8 controlled regulation of apicobasal polarity of neural plates cells. After phosphorylation, Smad1,5,8 interact with the PAR3-PAR6-aPKC complex and stabilizes linked tight junctions, while inversely low pSmad1,5,8 phosphorylation

supports re-arrangement of neural plate cells, a prerequisite for MHP formation and neural fold elevation [72].

### 2.2.3. *Patterning and developing of the spinal cord*

After closure, the neural tube starts to develop the spinal cord pattern along its dorso-ventral axis which is essentially established by the expression of Wnt and BMPs, the main roof plate morphogens, and sonic hedgehog (SHH) and TGF $\beta$ s, the morphogens released from the floor plate [74-76]. On the basis of provided intricated morphogenic gradients functionally distinct neurons are generated.

Both roof plate derived BMP6 and BMP7 are evoking Smad1,5,8 phosphorylation and subsequent induction of distinct dorsal interneurons (dI) via BMP receptor I activation [77-79]. While the BMP7/pSmad1,5 activated induction of dI1, dI3 and dI5 is independent of the patterned expression of progenitor proteins, e.g. Pax7, Gsh1/2 or Olig3, a Smad regulated expression of specific proneural proteins, e.g. cAth, and cAscl1, is required [79]. The patterning of dorsal progenitor proteins could be associated with other BMPs, e.g. BMP4 [80]. Smad1 and Smad5 activity seems to be important for the maintenance of neural cells as committed progenitors, because the loss of Smad1,5 reduced the total number of newly generated neurons and forces cell cycle exit and premature neurogenesis of neural progenitors [79]. The inhibitory Smad6 also promotes the neuronal differentiation in the spinal cord by interfering with Wnt-signalling [81]. Hazen and colleagues demonstrated, that the inhibitory Smad6 and Smad7 function to restrict the action of roof plate released BMPs to distinct dorsal interneurons and participate in the determination of dI4-dI6 spinal neuron number [82].

However, the subsequent orientation of spinal neurons is also controlled by BMP7 but not via activation of Smad signalling instead using phosphoinositide-3-kinase (PI3K) activation by the BMP receptor type II [77]. Nevertheless, Smad6 most potently allows blocking dI1 axon outgrowth [82].

While Smad3 was also expressed in the developing spinal cord, Smad2 was not detected [83]. Contrasting to Smad1,5,8, the expression of Smad3 in the developing spinal cord depends on progenitor proteins, e.g. Nkx6.1, Olig2, Nkx2.2, which directs the Smad3 expression mainly to discrete ventral progenitor domains [83]. Smad3 activity in turn reduces expression of progenitor proteins and promotes activation of neuronal differentiation, e.g. by supporting cell cycle exit via activating of p27<sup>Kip1</sup> expression [83]. The Smad3 expression is sufficient to promote differentiation of ventral and various dorsal interneurons, while differentiation of motor neurons is impaired. Therefore Smad3 expression is excluded in the prospective motor neuron progenitor cells [83].

### 2.2.4. *Induction of neural crest (NC)*

Multipotent embryonic cells of the dorsal region of the neural tube, an area existing immediately before forming the neural plate border (NPB), are the origin of neural crest cells (NC) that migrate to initiate a panel of diverse derivatives including various non-neural but also neural cells e.g. sensory neurons of the peripheral nervous system. The combination of several

signalling pathways is required [84] for a timely and locally well tuned progression. Besides Wnt signalling a major role in NC induction plays canonical BMP signalling [85-90] combined with FGF signalling [21,91]. The inductive step of NPB development during gastrulation requires a concerted action of activated Wnt signals and inhibition of BMP signalling represented by low Smad1,5,8 activity. Later, when neurulation proceeds and NC develops from NPB, activated Wnt signals cooperate with a robust Smad1,5,8 activation in NPB. While during the inductive step of NC gastrulation FGF downregulates Smad1,5,8 activity by triggering MAPK, the catalysed phosphorylation of Smad1,5,8 linker region is nearly completely absent in NPB during neurulation. In consequence, a strongly elevated Smad1,5,8 signalling is observed suggesting a 'two step model of NC development' with respect to activated Smad1,5,8 signalling [21].

### 2.3. Peripheral nervous system (PNS)

The further outcome of NC cells including the generation of the PNS also strongly depends on Smad signalling. Interestingly, Smad expression regulates both the formation of neurons of PNS as well as early stages of peripheral glial development.

Precursors that emigrate from the neural crest to the bowel generates the enteric nervous system (ENS) belonging to PNS. If fetal enteral neural crest-derived cells (ENCDC) of gut are exposed to BMP2 or BMP4 phosphorylated Smad1 translocates to the nucleus and the cells develop processes, indicating an essential role of Smad phosphorylation for neuron induction in the gut [92]. However, Smad phosphorylation alone is not sufficient to direct development of ENCDC towards ENS neurons or glial cells but requires further factors e.g. glial growth factor 2 (GGF2) [93]. Moreover, besides generation, also maturation of enteric neurons (as well as regulation of gliogenesis) during postnatal development is Smad-dependent. This function was identified because mice lacking in the homeodomain interacting protein kinase 2 (HIPK2), which can interact with Smad1, Smad2 and Smad3 [94] and therefore control transcription of subsequent Smad-dependent promoters, are characterized by a progressive loss of enteric neurons and an arrest in synaptic maturation postnatally. Additionally, in the HIPK2<sup>-/-</sup> mice the remaining enteric neurons exhibit an increased number of cells with nuclear Smad1,5,8 phosphorylation [95]. Interestingly, both in the enteric nervous system and in the midbrain [96], HIPK2 reduction severely reduces survival of dopaminergic neurons through interference with Smad signalling pathways, regulated by BMP2/4 or TGFβ, respectively. Whether altered Smad signalling, affected by HIPK2, is also responsible for Parkinson's disease and would allow to develop a therapeutic intervention has to be investigated in the future.

Recent data demonstrate, that Smad1 signalling by BMP2 induction is implicated in differentiation of primary enteric neurons to catecholaminergic (TH expression) but not to cholinergic (ChAT expression) neuronal subtypes [97].

Peripheral glia precursors derived from neural-crest and exhibiting characteristic expression of immature glia markers e.g. nestin, are critically influenced by BMP2/Smad signalling. Using Smad1 activation, neural-crest precursor cells, that begin to develop along glial pathway are kept in an undifferentiated immature multipotent state, thus allowing migration to their postmigratory locations [98] where they later acquire myelinating properties. Smad-linked

activation of GFAP promoter is part of this process [99]. Generally, oligodendroglial differentiation of precursor cells is actively suppressed by BMPs concomitant with the stimulation of astrocytic differentiation [100]. Accordingly, BMP4-activation directs progenitor cells in vivo to commit to the astrocytic rather than the oligodendroglial lineage [101].

### 3. Specific cell types and the role of Smad expression for their induction and maintenance

#### 3.1. Cell type decision, maturation and differentiation

Both generation and differentiation of neurons in the brain (CNS) comprises several different and complex principles. The initial proliferation of neural precursor cells is followed by the decision to specify neuronal or glial outcome. Afterwards, the primary neurogenesis of neuronal progenitors involves migration, settlement and stopping proliferation by cell cycle exit to enter a terminal differentiation pathway. Many of these basic biological routes as well as further differentiation steps, e.g. dendritic and axonal growth and orientating, synapse formation and myelination depend on Smad proteins. The parallel occurrence of long term and acute Smad-dependent processes require a distinct contextual organisation. For example, in the developing brain an important neural stem cell is provided by radial glia cells which can generate both glial cells and neurons. While TGF $\beta$  promoted differentiation of radial glia into astrocytes is mainly regulated by activation of MAPK signalling, neurogenesis is controlled by the interplay of Smad2,3 activity and PI3K activity [102].

##### 3.1.1. Glial cells

Radial glial cells can function as neuronal precursors and also control migration of neurons during cerebral cortex development [103]. Developing neurons are also actively implicated in the radial glial cell fate commitment [104]. Using TGF $\beta$  release, neurons can activate Smad signalling in radial glia cells and support their transformation to astrocytes [105] and also induces astrocytic differentiation and GFAP expression [106-109].

Moreover, endothelial cells also promote astrocytic differentiation by BMP-induced Smad signalling, but inhibit oligodendrocyte differentiation of postnatal cortical progenitors [110], and therefore participate in the sequential order of the two macroglial cell gliogenesis. The function of BMP/Smad signalling to drive precursors to astrocytic fate seems a common feature [111,112] which is also involved in CNS injury processes where oligodendrocyte precursors are driven towards type II astrocytes [113]. Accordingly, inhibition of BMP induced nuclear translocation of phospho-Smad1 by FGF2 activated MAPK activity is linked to Smad4 dissociation from Olig2 promoter and results in upregulation of oligodendrogenesis [114]. The very importance of a spatially and temporally regulated BMP induced Smad activation for the fate of neurogenic precursor cells was shown in chick spinal cord. At embryonic day 5, in neuroepithelial progenitors, astrocyte marker expression was inhibited and at embryonic day 6 it was promoted initiating a gliogenic period [115].



### 3.1.2. Neurons

The commitment of cell fates in the nervous system is strongly dependent on Smad2,3 signalling cascade. However, there have been contradictory results on the role of Smad3 for neuronal precursor proliferation and their differentiation, indicating a complex Smad signalling network dependent on local, temporal and contextual characteristics.

Some reports show, that TGF $\beta$ , which activates Smad3 signalling, diminishes neurogenesis in hippocampus after its chronic increase [116], and in SVZ and DG after intracerebroventricular TGF $\beta$ 1 infusion [117] and promotes neuronal differentiation from hippocampal and cortical progenitors [118]. An clear inhibitory function of Smad3 on neural precursor proliferation was shown in the developing spinal cord [119], where Smad3 also promotes differentiation and influences the fate of selected neurons. The antiproliferative role of Smad3 and Smad4 plays a role during early-postnatal differentiation of cerebellar neurons into postmitotic neurons, where TGF $\beta$  stimulation induces nuclear translocation of phosphorylated Smads and induction of cyclin-dependent kinase inhibitors p21, p27 and markers of neuronal maturity [120].

However, other reports show that TGF $\beta$  increased neurogenesis in DG after adrenalectomy [121] or in SVZ after stroke [122].

To clarify the relevance of Smad3 for adult neurogenesis Smad3 null mice were studied [123,124]. These mice show decreased neurogenesis in the DG and the SVZ and exhibit a thinner and more disorganized rostral migratory stream (RMS) of neuronal precursor cells (NPC). Using RMS NPCs migrate from SVZ to reach the olfactory bulb, where they differentiate into granular and periglomerular neurons. Though a decreased number of proliferating cells demonstrates the requirement of Smad3 for maintaining a proper cell division rate in SVZ, the neuronal fate is not altered by Smad3 deficiency [123].

For differentiation of mesencephalic progenitors into dopaminergic (DA) neurons a concerted interaction of Smad2,3 signalling and p38 MAPK-pathways by TGF $\beta$  receptor stimulation is necessary. Accordingly, treatment of ventral mesencephalic neural progenitors with TGF $\beta$  increased the number of tyrosine-hydroxylase (TH)-positive cells [125]. Though not required for the neurogenesis, the survival of midbrain dopaminergic neurons depends on function of the homeodomain interacting protein kinase 2 (HIPK2), which interacts with R-Smads to activate TGF $\beta$  responsive genes [96]. Loss of HIPK2 increases apoptosis in DA neurons during development. Nevertheless, apoptosis of specific neural progenitors during neural differentiation also involves Smad phosphorylation as shown for Smad1,5,8 [126,127].

In vitro, treatment of ventral mesencephalic cells with BMP5,6 and 7 also significantly increased the number of TH-positive neurons via Smad phosphorylation and nuclear translocation [128], while the neurotrophin growth/differentiation factor 5 (GDF5) induced Smad pathway promotes neuronal but not dopaminergic differentiation [129]. Remarkably, the in vitro generation of functional dopaminergic substantia nigra neurons for transplantation requires a protocol which uses the temporarily and contextually distinct roles of Smad proteins: Firstly, the dual inhibition of Smad signalling in embryonic stem cells is required to allow induction of floor plate cell state which then passes over to midbrain floor plate cell state.

Secondly, an activation of Smad signalling using TGF $\beta$  pathway is necessary to finally obtain dopaminergic neurons suitable to threat Parkinson's like symptoms [130].

In the developing cerebellum, Smad1 is expressed in the external germinal layer and Smad5 is synthesized in newly differentiated granule neurons. In the granule precursor cells, Smad5 overexpression is sufficient to initiate differentiation [131]. In mice lacking Smad4, a decreased number of cerebellar Purkinje cells and parvalbumin-positive interneurons [132] is found while no alterations of proliferation of neuronal precursor cells were detectable. Loss of Smad4 also reduces differentiation of noradrenergic neurons [133].

Basal forebrain cholinergic neurons are essentially involved in the organisation of cortical brain structures, learning and memory. Smad1-5 are expressed in basal forebrain from embryonic day 14 to the adult age [134]. An intact canonical TGF $\beta$ /Smad2,3 cascade is important for the function of cholinergic neurons. Smad signalling mediates rapid inhibition of calcium influx in cholinergic basal forebrain neurons [135]. The sleep/wake circadian rhythm controlled by cholinergic neurons of the pedunculopontine nucleus (PPT) is also dependent on phosphorylation levels of Smad2 and Smad3. Overactivation of Smad2,3 signalling resulting in increased nuclear translocation is linked to sleep/wake circadian rhythm amplitude deficits [136]. BMP9 driven Smad1,5 phosphorylation and their formation of complexes with Smad4 are involved in the induction of the cholinergic phenotype in the basal forebrain [134,137]. Moreover, only in postmitotic cholinergic neurons, this pathway induces NGF expression, an autocrine/paracrine cholinergic trophic factor, which stimulates ACh production [138]. Accordingly, it is of therapeutic relevance for treatment of Alzheimer's disease, that BMP9/Smad1,5 signalling can prevent lesion-evoked impairment of the cholinergic septo-hippocampal neurons in adult mice [139]. The basal forebrain cholinergic neuron specific gene expression pattern is also basically linked to an interaction of Smad and Szn1 (Smad-interacting zinc finger protein), a factor which causes mental retardation if mutated [140,141].

### 3.2. Axon and synapse formation

Induction of axons and orienting of axon responses are controlled by several Smad molecules of both the BMP and the TGF $\beta$ /activin pathways in a timely and spatially organized manner. Several different neuronal/neural cell types use individual aspects of the Smad machinery.

#### 3.2.1. Basic role of Smads for neuritic outgrowth

In PC12 cells, which are initially derived from neural crest cells and are used as model system for neuronal differentiation, BMP2 induces neurite outgrowth [142] by activation of TAK1/p38 kinase signalling pathway [143,144], which is in turn tightly controlled by the simultaneous, also BMP2-induced expression of the inhibitory Smad6 and Smad7 [142] performed via activation of Smad1,5,8 pathway. The inhibitory Smads are considered to inversely inhibit BMP signalling in a concerted action by repressing Smad1,5,8, mediating BMP receptor degradation by Smurf1 [16,145], and by physical interaction with TAK1-binding protein, which finally reduces the p38-mediated neuritic outgrowth [143,144]. Neurotrophin 3 induced

neuritic outgrowth is potentiated by BMP4/6 induced phosphorylation of Smad1,5,8 and their subsequent inhibition of MEK in chicken neurons [146].

Otherwise, BMP2 inhibits neuritic outgrowth and differentiation of motor neuron-like NSC-34 cells by activation of Smad1,5,8-dependent signalling and subsequent Id genes activation, which are main targets of Smad signalling and which negatively regulate differentiation of various cells including neurogenic precursors and motor-neuron precursors [147].

### 3.2.2. *Subtle control of axonal morphogenesis*

The axonal morphogenesis is strongly influenced by the TGF $\beta$ -regulated signalling protein Smad2. Endogenous Smad2 is constitutively activated and its phosphorylated form is nuclearly localized in human and mouse hippocampal and cortical neurons [148,149] and in primary granule neurons of rat cerebellum [150]. In granule neurons Smad2 can form a physical complex with the endogenous transcriptional modulator SnoN, which is also nuclearly localized and strongly enhances axonal growth [151] and neuronal branching [152] by regulating a large number of neuronal genes [153]. SnoN ubiquitination and degradation is controlled by the nuclear ubiquitin ligase Cdh1-anaphase-promoting complex (Cdh1-APC), which physically interact with nuclear SnoN thus suppressing axonal growth. [151,154]. Obviously Smad2 binding to SnoN facilitates the ability of Cdh1-APC to control SnoN degradation. Consequently, if phosphorylation levels of Smad2 are reduced, SnoN levels increased and axonal growth is stimulated which allow therapeutic potential after brain injury [150]. The constitutive neuronal expression of axonal growth inhibitors e.g. Smad2, Smad3 [148] and Cdh1 [155] and the reduced expression of axon growth promoters (SnoN) in terminally differentiated neurons [150] generate an intrinsic axon growth inhibition control, allowing a balance between steady state and neuronal plasticity.

### 3.2.3. *Axonal regeneration*

Axonal regeneration requires the reversal of an age-dependent loss of intrinsic axonal growth capability [156,157]. In dorsal root ganglion cells (DRG), which possess two branches of a initially unipolar axon, (a) a central branch containing in the spinal cord and (b) a peripheral branch innervating sensory targets, the axotomy of the peripheral branch requires function of transcription factor Smad1 for successful regeneration [158]. While in embryonic DRG neurons during the phase of active axon growth Smad1 RNA and protein were strongly expressed and found abundantly C-terminally phosphorylated in the nuclei (while Smad8 expression was low and Smad5 expression was concentrated to the periventricular zone), in adult DRG neurons Smad1 phosphorylation was diminished [159]. However, after axotomy, the induction and nuclear translocation of Smad1 precedes the onset of axonal extension and are maintained over longer time, demonstrating the importance of Smad1 for the perpetuation of the activated axonal growth program [158]. Interestingly, Smad1-dependent axonal growth program is an intrinsic feature of DRG neurons which functions independently of extracellular BMP. Until now, the detailed underlying mechanism for BMP-independent Smad1 phosphorylation is not completely elucidated. It is possible, that endogenous cytoplasmic kinases or intracellularly available BMP/BMP-receptor complexes might be involved. The ligand-independent C-

terminal phosphorylation of the SSXS domain of Smad2 and Smad3 was already reported [160] suggesting a possible extracellular ligand-independent universal strategy. Probably, Smad1 signalling is also involved in the axonogenesis of many other classes of neurons, e.g. Purkinje cells, retinal ganglionic, olfactory and motor neurons which were already found positively labelled for pSmad1 immunoreactivity [159]. Due to the possible linker phosphorylation of Smad1 by several kinases (e.g. MAPK, GSK, CDK), the molecule might serve as an integrator of various pathways relevant for axon growth and regeneration [14,23-26]. However, for clinical and therapeutic relevance, the transduction of adeno-associated vector encoded BMP4 allows stimulation of Smad1 phosphorylation and activation of axonal growth independent of axotomy and also promotes sensory axon regeneration after axotomy [159]. At least partly BMP effects are dependent on its binding to repulsive guidance molecules [161].

While Smad1 positively supports axonal growth, the inhibitory Smad6 can block axon outgrowth, as shown for post-mitotic dI1 axons of the spinal cord [82] and therefore exert roles in spatially limiting the influence of BMP signalling on neurons.

In *Drosophila*, Smad2 is involved in motor neuron axon guidance, as null mutants of Smad2 exhibit axon guidance defects [162]. It is suggested, that a chemoattractant signal of TGF $\beta$  superfamily members provided by muscle cells might guide motor axons to their appropriate innervation sites.

#### 3.2.4. *R-Smads and Co-Smads exert neuronal plasticity*

The development of peripheral tissues is connected to axonal growth of peripheral nerves, linking sensory information to the spinal cord via neuronal cell bodies localized in ganglia. After initial axon extension for trigeminal sensory neurons in mice, a Smad4-dependent retrograde signalling from developing face to the neuronal nuclei was found, which control the expression of genes in neurons of the trigeminal ganglion in dependence on their position. Extrinsic signals released from developing craniofacial tissue, e.g. BMP4, activate phosphorylation of Smad1,5,8 at the axon terminals and selectively retrogradely signals this information to selected trigeminal neuronal cell bodies thus inducing spacially patterned expression of further transcription factors along the dorso-ventral axis of the trigeminal ganglion [163]. A comparable retrograde mechanism based on dSmad2 was found in R7 axons in the *Drosophila* visual system, where activin, secreted from the R7 cells growth cone in an autocrine manner, activates its receptor and initiates intracellular Smad2 phosphorylation. The pSmad2 complexes to the nuclear import adaptor Importin- $\alpha$ 3, shuttles to the nucleus and reduces growth cone motility and synaptogenesis via transcriptional regulation of several target genes [164]. Already in *Drosophila*, a retrograde Smad-based signalling of the BMP homolog Gbb regulates synaptic growth in presynaptic motor neurons [165]. The underlying mechanism for the retrograde Smad-based signalling was recently examined for the BMP4 signalling system. Ji and colleagues (2012) discovered that endosomes carrying the BMP-signalling complex containing phosphorylated Smad1,5,8, are essentially linked to a BDNF-induced axonal or growth cone translation of Smad proteins to retrogradely control transcription in trigeminal neurons [166,167]. This 'two-target-derived signalling' integrates neurotrophin and BMP signals on the level of Smad proteins, which are axonally translated on demand. This process

is involved in the acquisition of positional identity markers during trigeminal ganglia development and selectively works in ophthalmic and maxillary but not in mandibular axons obviously due to the failure of BDNF in the mandibular target field [166]. The underlying mechanism suggests to re-evaluate the role and extent of Smad based transcriptional regulation. (1) Nearly all cellular surfaces/membranes, including dendrites, axons, growth cones and spines might receive information which could be integrated by Smad signalling. (2) The retrograde transport of other ligands of the TGF $\beta$  group was already reported, e.g. TGF $\beta$ 2 in normal and injured motor neurons [168], implicating the question whether axonal Smad signalling is involved in injury-response. (3) The mechanism might work with all those extracellular and also intracellular factors that affect Smad translation, and might at least partly explain neurodegenerative disorders, which are characterized by disturbed axonal transport and/or neurotrophin deficiency at neuronal terminals

The specific structure of axons and the axonal transport of Smad signals [166] as well as simple translocation of receptor phosphorylated Smads from cytoplasmic receptor site into the nucleus require intact microtubule network. Phosphorylated Smad2 is transported using microtubules by support of kinesin-1 and kinesinATPase activity [169] and also requires dynein light chain km23-1 [6], whereas pSmad3 transport was dependent on km23-2 function [170]. Additionally retrograde axonal transport of phosphorylated Smads, which is necessary for transition of axonal synaptic BMP-like ligand mediated phosphorylation of Smads in *Drosophila*, also depends on an intact function of p150Glued protein. P150Glued is a component of the dynactin complex, which is necessarily involved in synapse assembly and stability [171]. However, a p150Glued knockdown induced disruption of synaptic homeostasis can be rescued by activation of Smad signalling at the soma [172,173], confirming the important role of nuclear phosphorylated Smad to synaptic homeostasis.

### *3.2.5. Astrocytic and oligodendrocytic Smad signalling influences neuronal axonal growth*

Astrocytes surrounding synapses are also target of neuronal activity. Glial metabotropic glutamate 2/3 receptor activation by neuronally released glutamate induces astrocytic TGF $\beta$ 1 secretion, leading to GFAP gene activation and astrocyte differentiation involving astrocytic Smad signalling pathways [174].

Chondroitin sulphate proteoglycans (CSPG), mainly synthesized by astrocytes, can inhibit axonal growth and regeneration. After traumatic brain injury or disruptions of the blood brain barrier, mature or fibrinogen-coupled latent TGF $\beta$  is released into the CNS and increases the CSPG expression by a Smad signalling manner in astrocytes [175,176]. Both astrocytic Smad2 and Smad3 expression and phosphorylation leads to inhibition of neuronal outgrowth and is required for astrocytic neurocan synthesis, whereas phosphacan only requires Smad2 [175,176]. Reduced Smad3 levels selectively reduced induction of chondroitin-4- sulphotransferase 1 and the amount of 4-sulfated CSPGs secreted by astrocytes and also promoted axonal growth of neurons which were fed on these astrocytes [175]. Taxol treatment of astrocytes can reduce CSPG expression by interfering with kinesin-1-dependent pSmad transport into the nucleus [177] and improve axon regeneration after spinal cord injury.

The Smad-interacting-protein 1 (Sip1) is an essential modulator for CNS myelination. In oligodendrocytes, it functions in a dual-mode manner by repressing the differentiation inhibitory signals of the BMP-receptor activated Smad1,5,8 activity [178] and activating oligodendrocytes-promoting factors, thus controlling proper myelination in the CNS [38].

### 3.3. Smads contribute to synapse formation and synaptic transmission processes

Synapse formation and remodelling are results of intrinsic programs and environmental insults. Synapses are characterized by close cell-to-cell communications, which also include synaptic transmission and signalling processes mediated by growth factors e.g. members of the TGF $\beta$  superfamily. In *spinster* a mutation of the inhibitory Smad Dad (Daughters against dpp) causes synaptic overgrowth [179]. A well investigated synaptic system is the neuromuscular junction (NMJ) of *Drosophila*, showing the involvement of Smad proteins in synapse function both in pre- and in postsynaptic cells. The release of muscular BMP4-type ligand Gbb, which is required for intact function of a large part of postsynaptic muscles [165], allows its binding to a presynaptic receptor consisting of one type I and one type II receptor, Sax or Tkv and Wit, respectively [180,181], and the subsequent phosphorylation of the R-Smad, Mothers against dpp (Mad) [165]. Activated Mad is responsible for presynaptic nuclear propagation of the postsynaptic muscular BMP signals and finally for regulating synapse growth, morphology and function of motor-neurons at least partly by altering transcription [165,180,182]. The retrograde transport of Mad utilizes microtubule motor proteins, e.g. dynein complexes, and is sensitive to disruption of dynein motor function [165]. The *Drosophila* Co-Smad/Smad4 homolog Medea (Med) is also necessary for presynaptic BMP signalling cascade and essential for synaptic growth [183]. Both, Med and Mad mutants, exhibit defects in neurotransmitter release and synaptic ultrastructure [165,183]. It is suggested that an equilibrium exist between the binding of presynaptic phosphorylated Mad to Med and the binding of Med to the RING finger E3 ubiquitin ligase Hiw [184], which regulate the synaptic growth in NMJ [183].

Additionally to the retrograde neuronal Mad/Med-mediated Gbb signalling, anterogradely released TGF $\beta$  type ligands activate Mad phosphorylation at the postsynaptic density (PSD) zone of postsynaptic muscles and NMJ depolarization supports Mad nuclear transition in the muscles [185]. In summary, a mutual information flow of neuron and muscle on each physiological state and the synaptic cleft situation is essentially controlled by specific R- and Co-Smad signalling in both cellular systems.

Importantly, the activation of Mad phosphorylation and nuclear translocation by the BMP ligand Gbb directs the expression of synaptic homeostasis independently of synaptic growth control in *Drosophila* [173]. However, retrograde transport of Mad, which was phosphorylated at the periphery of the axon terminal due to synaptic Gbb binding, mediates synaptic homeostasis. Remarkably, Gbb induced phosphorylation of Mad at the soma site of the neuron compensate for a disturbed retrograde axonal pMad transport completely [173]. The release of muscular Gbb into the synaptic cleft at the periphery to activate neuronal retrograde pMad transport, can be regulated by activin, which controls the muscular Gbb synthesis by recruiting the *Drosophila* dSmad2 [186]. Cdc42-interacting protein 4 (dCIP4) which is also localized

postsynaptically at the NMJs co-regulates the phosphorylation of neuronal Mad levels by inhibiting postsynaptic Gbb secretion [187].

These data indicate, that both activin and BMP directed R-Smad signalling controls synaptic function at NMJ and affects the development of synaptic homeostasis.

## 4. Role of Smads for neurological disorders

Several neurological diseases are characterized by disturbed cellular or subcellular Smad localization and show artificial Smad sequestration or deposition. Especially in neurodegenerative and motor neuron diseases a disruption of Smad controlled transcriptional machinery was reported.

Neurodegenerative diseases are characterised by selective and progressive loss of specific populations of neurones. Four disease-classes are proposed based on four major affected proteins, tau,  $\beta$ A4-amyloid ( $\beta$ A4),  $\alpha$ -synuclein and TDP-43, and therefore, neurodegenerative diseases can roughly be classified into four main groups: amyloidosis, tauopathies, synucleinopathies and TDP-43 proteinopathies (for review see [188]). Recent data indicate, that in all four disease groups Smad proteins are essentially involved in the disease progression, finally indicating a remarkable commonness of these diseases.

### 4.1. Alzheimer's disease

Alzheimer's disease (AD), representing a mixture of both amyloidosis and tauopathy, is a severe neurodegenerative disorder and the most common cause of dementia in the elderly. Typical clinical symptoms are memory loss, disturbed activities of daily life and deficiency of social competence. Common morphological correlates to the clinical features are extracellular  $\beta$ A4-plaque depositions, intraneuronal tau pathology, neuronal cell death, and cell cycle activation [189].

In AD, we identified a strong disturbance of the normal constitutive nuclear localization of phosphorylated Smad2 and Smad3 in hippocampal and cortical neurons [148], which subsequently was confirmed by others [190,191]. In AD brain, a strong colocalization of pSmad2,3 with intracellular neurofibrillary tangles (NFTs) in neurons and with  $\beta$ A4-amyloid plaques in addition to a sequestration of pSmad2,3 in cytoplasmic granular vesicles is detectable [148]. Additionally a significant reduction of Smad2,3,4, which are involved in activation of cell cycle proteins was described. Smad4 directly controls cyclin-dependent kinase 4 (CDK4) expression in neuronal cells and is involved in cell cycle activation of neurons in AD brain [192]. It is quite possible, that the recently identified TGF $\beta$ 1-induced antiapoptotic factor (TIAF1), which can bind and block Smad4-dependent promoter activation [193], participates in the cytoplasmic Smad sequestration in AD neurons and suppresses Smad-regulated promoter activation. Environmental stress or TGF $\beta$ 1, can induce TIAF aggregation, which in turn removes soluble Smad4, induces apoptosis and activates  $\beta$ A4 generation and its aggregation [193].

A reason for the intraneuronal dislocation of Smads could be due to the disturbed nucleocytoplasmic transport in hippocampal AD neurons [194,195]. The inner nuclear membrane comprises integral proteins, e.g. MAN1, which regulate Smad phosphorylation and nuclear translocation [196], while the outer nuclear membrane is directly continuous with the rough endoplasmic reticulum. Both membranes are separated by nuclear core complexes, which also control nucleocytoplasmic shuttling of Smad2 [197]. Altered nuclear membrane integrity [195] might provoke misrouting of phosphorylated Smads into the associated ER compartment, indicated by the co-localization of phosphorylated Smad2 granules with a luminal marker protein of the ER, protein disulphide isomerase [148,198]. ER containing Smad2 could be secreted and thus explain the extracellular association of pSmad2,3 with amyloid plaques [148,190]. Alternatively, pSmad2 could interact in early endosomes with  $\beta$ A4-peptides which are accumulating in the early stages of AD [199], and there also meet BACE, the  $\beta$ -secretase, generating  $\beta$ -amyloid peptides [200], activated TGF $\beta$  receptors type I and II as well as SARA, a Smad anchor for receptor activation [201-203]. It is also possible, that early endosomes can fuse to the ER thus allowing a direct route of Smad proteins to the lumen of the ER [204,205].

However, activation of cell cycle is a hallmark of AD and (1) probably supports hyperphosphorylation of tau protein which leads to generation of neurofibrillary tangles and (2) increases neuronal apoptosis by phosphorylation of retinoblastoma protein and activation of E2F based apoptotic impacts (for review see [206]). Hyperphosphorylation of tau affects the neuronal Smad2,3 localization, diminishes its nuclear concentration and thus impedes with transcriptional Smad functions as shown after  $\beta$ A4 treatment of primary neurons [207]. NFT formation in AD brain disturbs common intra-neuronal transport mechanisms [208]. This disturbance could also interfere with retrograde Smad signalling which could be suggested for hippocampal neurons but so far has only demonstrated for *Drosophila* motor neurons [173]. Moreover, we also suggest, that the principle of the two-target-derived signalling integrating neurotrophin (BDNF) and Smad pathway found in rat trigeminal neurons [166,167] might also play a role for AD progression, because BDNF is deficient in entorhinal cortex and hippocampus in AD [209,210]. Finally, a disturbed nuclear Smad localization might influence competence of neurons to express synaptic homeostasis and plasticity, both processes are well investigated for *Drosophila* motor neurons.

Pathogenic reduction of neuronal Smads can also support activation of neuronal cell cycle, resulting in neuronal apoptosis [211] e.g. by repression of cyclin-dependent kinase inhibitors (CDIs) p15<sup>Ink4B</sup>, p16<sup>Ink4A</sup>, or p21<sup>Cip</sup>, which expression is controlled by TGF $\beta$ /Smad signalling [212-214]. Though the increased levels of TGF $\beta$ 1 in AD [215,216] does not seem to be able to compensate for the compromised canonical neuronal Smad pathway [148,190], astrocytes were shown responsive to this growth factor, which induces amyloid precursor protein (APP) expression in cooperation with Sp1 [217] and CTCF [218]. Altered APP cleavage by the concerted action of  $\alpha$ -,  $\beta$ - and  $\gamma$ -secretases is a main aspect of AD pathology and results in the generation of the pathogenetic  $\beta$ A4 peptide 1-42, which is neurotoxic and leads to  $\beta$ A4-amyloid plaques. Whether  $\beta$ A4 peptide 1-42 can further suppress neuronal TGF $\beta$ 1/Smad2,3 signalling by activation of the inhibitory Smad7 in neurons as already shown for human monocytes cannot be excluded [219].



Clearance of  $\beta$ A4 peptide was reported by microglia cells [220]. TGF $\beta$  released by neurons or astrocytes stimulates  $\beta$ A4 peptide uptake through Smad3-dependent increased scavenger receptor SR-A expression and increased phagocytosis [221]. Though the microglial capacity to remove  $\beta$ A4-plaques is limited [222], peripheral phagocytes, migrating into the brain if their Smad2,3 signalling is blocked at the expense of Smad1,5,8 phosphorylation, might perform this task successfully [223].

In other tauopathies, e.g. Pick disease, progressive supranuclear palsy, and corticobasal degeneration, cytoplasmic phospho-tau bound pSmad2,3 was also identified, though reduced nuclear pSmad levels were not found [224] suggesting a different impact on Smad signalling in these diseases compared to AD.

#### 4.2. Parkinson's disease

Parkinson's disease (PD) is a progressive neurodegenerative disorder which is characterized by dopaminergic neurodegeneration in the substantia nigra (SN) pars compacta. The closely associated loss of dopaminergic inputs into the striatum results in failure of motor programmes (e.g. voluntary movements) and initiates a striatal plastic change [225,226]. The gradual loss of dopaminergic neurons is linked to an increase of dopamine turnover [227,228].  $\alpha$ -synuclein ( $\alpha$ -syn) is the major protein component of Lewy bodies, a hallmark of PD. It can aggregate to form toxic oligomers and fibrillar structures [226,229].

It was shown, that the TGF $\beta$ 3-Smad-HIPK2 pathway is important for the survival of dopaminergic neurons during development [96,230]. However, recently, a very specific and important role of Smad3 for the nigrostriatal system was explored in a Smad3 null mouse model representing an exciting model of PD [231]. These animals show a reduced number of dopaminergic neurons in the rostral SN, resulting from a postnatal neurodegenerative process. The selective pro-survival effect of Smad3 for SN dopaminergic neurons was emphasized by stereological quantifications showing no alterations of the striatal neuronal number. In Smad3 null mice a diminished trophic support provided by Erk1/2 signalling and a reduced astrocytic support to nigral dopaminergic neurons was detected [231]. Though striatal dopamine levels were not changed in Smad3 null mice, a strong increase of DA metabolism, including elevated monoamine-oxidase (MAO) levels were identified, resembling PD findings [232]. Finally, increased expression and aggregation of  $\alpha$ -syn in neurites and cell bodies of several telencephalic, mesencephalic and rombencephalic brain regions and in fibres of the primary and secondary motor cortex were found [231]. Two possible functional Smad3-based circuits are suggested. Either Smad3 might directly regulate the  $\alpha$ -syn transcription, which further allows modulating DA metabolism, or the Smad3 deficiency caused elevated DA catabolism generates oxidative stress which allow toxic aggregation of  $\alpha$ -syn. The data qualify Smad3 as a possible target for PD therapy. Interestingly, the currently pursued target to re-implant dopaminergic neurons is also based on modification of Smad signalling during their in vitro establishment [130,233].

### 4.3. Motor neuron diseases

As mentioned above, Smads are essentially controlling the plasticity of NMJ, the intra-axonal transport and affect axonal repair processes. Accordingly, selective motor neuron diseases exhibit disturbances of Smad signalling, e.g. spinal muscular atrophy, [234], spinal bulbar muscular atrophy [235] or amyotrophic lateral sclerosis (ALS).

ALS is a progressive neurodegenerative disease, which targets upper and lower motor neurons. In the primary motor cortex and the anterior horn of the spinal cord, motor neurons disappear and the pyramidal tract degenerates [236-238]. The remaining motor neurons exhibit inclusion bodies e.g. Bunina bodies, hyaline and skein-like inclusions. The pathological transactive response DNA-binding protein with a molecular weight of 43 kDa (TDP-43) was shown to be the major disease protein in ALS. Recently, increased nuclear immunoreactivity for pSmad2,3 in motor neurons was reported from sporadic ALS patients in spinal cord [239]. Motor neurons, where a colocalization of TDP-43 and pSmad2,3 in skeine-like and round hyaline inclusions was detected [239], show reduced nuclear pSmad2,3 immunoreactivity [240]. Obviously a disruption of Smad signalling by Smad segregation, comparable to AD or several tauopathies, enhances the loss of motor neuron function. Smad4 seems actively involved in the control of motor function as targeted disruption of Smad4 demonstrates [132]. Probably, increased nuclear pSmad2,3 content in the remaining ALS motor neurons indicates an endogenously initiated approach to a functional improvement. Experiments with an ALS mouse model support this hypothesis, because TGF $\beta$ 2 administration ameliorated the motor performance of the mice [241]. A reduction of TDP-43 containing aggregates by Smad2 overexpression confirms in vitro a protective role of activated Smad signalling in ALS [242].

In summary, initiation and/or progression of many neurological disorders are directly linked to altered Smad signalling, comprising cytoplasmic Smad aggregation/sequestration, and nuclear reduction, disruption of transcriptional machinery and stimulating proapoptotic signalling, therefore disturbing biological processes, which are essential for all phases of nervous system development and homeostasis.

## Acknowledgements

This manuscript was supported by the Project BBZ09: 14494 (University Leipzig) and the AFI-Project 984 000-150.

## Author details

Uwe Ueberham<sup>\*</sup> and Thomas Arendt

<sup>\*</sup>Address all correspondence to: Uwe.Ueberham@medizin.uni-leipzig.de

Department for Molecular and Cellular Mechanisms of Neurodegeneration, Paul Flechsig Institute for Brain Research, University Leipzig, Leipzig, Germany

## References

- [1] Tsukazaki T, Chiang TA, Davison AF, Attisano L, Wrana JL. SARA, a FYVE domain protein that recruits Smad2 to the TGFbeta receptor. *Cell* 1998;95(6) 779-791.
- [2] Shi W, Chang C, Nie S, Xie S, Wan M, Cao X. Endofin acts as a Smad anchor for receptor activation in BMP signaling. *J. Cell Sci.* 2007;120(Pt 7) 1216-1224.
- [3] Dong C, Li Z, Alvarez R, Jr., Feng XH, Goldschmidt-Clermont PJ. Microtubule binding to Smads may regulate TGF beta activity. *Mol. Cell* 2000;5(1) 27-34.
- [4] Santibanez JF, Quintanilla M, Bernabeu C. TGF-beta/TGF-beta receptor system and its role in physiological and pathological conditions. *Clin. Sci (Lond)*. 2011;121(6) 233-251.
- [5] Shi Y, Massague J. Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell* 2003;113(6) 685-700.
- [6] Jin Q, Ding W, Mulder KM. Requirement for the dynein light chain km23-1 in a Smad2-dependent transforming growth factor-beta signaling pathway. *J. Biol. Chem.* 2007;282(26) 19122-19132.
- [7] Jin Q, Gao G, Mulder KM. Requirement of a dynein light chain in TGFbeta/Smad3 signaling. *J Cell Physiol.* 2009;221(3) 707-715.
- [8] Miyazono K. Molecular mechanisms of transforming growth factor-beta signaling and disease: The 59th Fujihara International Seminar, 2010. *Cancer Sci.* 2011;102(6) 1242-1244.
- [9] Kretschmar M, Doody J, Timokhina I, Massague J. A mechanism of repression of TGFbeta/ Smad signaling by oncogenic Ras. *Genes Dev.* 1999;13(7) 804-816.
- [10] Matsuura I, Denissova NG, Wang G, He D, Long J, Liu F. Cyclin-dependent kinases regulate the antiproliferative function of Smads. *Nature* 2004;430(6996) 226-231.
- [11] Luo Q, Nieves E, Kzhyshkowska J, Angeletti RH. Endogenous transforming growth factor-beta receptor-mediated Smad signaling complexes analyzed by mass spectrometry. *Mol. Cell Proteomics.* 2006;5(7) 1245-1260.
- [12] Sasaki A, Masuda Y, Ohta Y, Ikeda K, Watanabe K. Filamin associates with Smads and regulates transforming growth factor-beta signaling. *J. Biol. Chem.* 2001;276(21) 17871-17877.
- [13] Inoue Y, Imamura T. Regulation of TGF-beta family signaling by E3 ubiquitin ligases. *Cancer Sci.* 2008.
- [14] Alarcon C, Zaromytidou AI, Xi Q, Gao S, Yu J, Fujisawa S, et al. Nuclear CDKs drive Smad transcriptional activation and turnover in BMP and TGF-beta pathways. *Cell.* 2009;139(4) 757-769.
- [15] Sapkota G, Alarcon C, Spagnoli FM, Brivanlou AH, Massague J. Balancing BMP signaling through integrated inputs into the Smad1 linker. *Mol Cell.* 2007;25(3) 441-454.

- [16] Zhu H, Kavsak P, Abdollah S, Wrana JL, Thomsen GH. A SMAD ubiquitin ligase targets the BMP pathway and affects embryonic pattern formation. *Nature* 1999;400(6745) 687-693.
- [17] Gao S, Alarcon C, Sapkota G, Rahman S, Chen PY, Goerner N, et al. Ubiquitin ligase Nedd4L targets activated Smad2/3 to limit TGF-beta signaling. *Mol Cell*. 2009;36(3) 457-468.
- [18] Burstyn-Cohen T, Stanleigh J, Sela-Donenfeld D, Kalcheim C. Canonical Wnt activity regulates trunk neural crest delamination linking BMP/noggin signaling with G1/S transition. *Development*. 2004;131(21) 5327-5339.
- [19] Sela-Donenfeld D, Kalcheim C. Regulation of the onset of neural crest migration by coordinated activity of BMP4 and Noggin in the dorsal neural tube. *Development*. 1999;126(21) 4749-4762.
- [20] Alexandrova EM, Thomsen GH. Smurf1 regulates neural patterning and folding in *Xenopus* embryos by antagonizing the BMP/Smad1 pathway. *Dev. Biol.* 2006;299(2) 398-410.
- [21] Stuhlmiller TJ, Garcia-Castro MI. FGF/MAPK signaling is required in the gastrula epiblast for avian neural crest induction. *Development*. 2012;139(2) 289-300.
- [22] Makkar P, Metpally RP, Sangadala S, Reddy BV. Modeling and analysis of MH1 domain of Smads and their interaction with promoter DNA sequence motif. *J Mol Graph. Model*. 2009;27(7) 803-812.
- [23] Eivers E, Demagny H, De Robertis EM. Integration of BMP and Wnt signaling via vertebrate Smad1/5/8 and *Drosophila* Mad. *Cytokine Growth Factor Rev*. 2009;20(5-6) 357-365.
- [24] Scherer A, Graff JM. Calmodulin differentially modulates Smad1 and Smad2 signaling. *J. Biol. Chem*. 2000;275(52) 41430-41438.
- [25] Kretzschmar M, Doody J, Massague J. Opposing BMP and EGF signalling pathways converge on the TGF-beta family mediator Smad1. *Nature* 1997;389(6651) 618-622.
- [26] Fuentealba LC, Eivers E, Ikeda A, Hurtado C, Kuroda H, Pera EM, et al. Integrating patterning signals: Wnt/GSK3 regulates the duration of the BMP/Smad1 signal. *Cell*. 2007;131(5) 980-993.
- [27] Kuroda H, Fuentealba L, Ikeda A, Reversade B, De Robertis EM. Default neural induction: neuralization of dissociated *Xenopus* cells is mediated by Ras/MAPK activation. *Genes Dev*. 2005;19(9) 1022-1027.
- [28] Hough C, Radu M, Dore JJ. Tgf-Beta induced erk phosphorylation of smad linker region regulates smad signaling. *PLoS. ONE*. 2012;7(8) e42513.
- [29] Luo K. Ski and SnoN: negative regulators of TGF-beta signaling. *Curr. Opin. Genet. Dev*. 2004;14(1) 65-70.

- [30] Mizuide M, Hara T, Furuya T, Takeda M, Kusanagi K, Inada Y, et al. Two short segments of Smad3 are important for specific interaction of Smad3 with c-Ski and SnoN. *J. Biol. Chem.* 2003;278(1) 531-536.
- [31] Wu JW, Krawitz AR, Chai J, Li W, Zhang F, Luo K, et al. Structural mechanism of Smad4 recognition by the nuclear oncoprotein Ski: insights on Ski-mediated repression of TGF-beta signaling. *Cell* 2002;111(3) 357-367.
- [32] Deheuninck J, Luo K. Ski and SnoN, potent negative regulators of TGF-beta signaling. *Cell Res.* 2009;19(1) 47-57.
- [33] Wang W, Mariani FV, Harland RM, Luo K. Ski represses bone morphogenic protein signaling in *Xenopus* and mammalian cells. *Proc. Natl. Acad. Sci. U. S. A* 2000;97(26) 14394-14399.
- [34] Conidi A, Cazzola S, Beets K, Coddens K, Collart C, Cornelis F, et al. Few Smad proteins and many Smad-interacting proteins yield multiple functions and action modes in TGFbeta/BMP signaling in vivo. *Cytokine Growth Factor Rev.* 2011;22(5-6) 287-300.
- [35] Chng Z, Teo A, Pedersen RA, Vallier L. SIP1 mediates cell-fate decisions between neuroectoderm and mesendoderm in human pluripotent stem cells. *Cell Stem Cell.* 2010;6(1) 59-70.
- [36] Nitta KR, Tanegashima K, Takahashi S, Asashima M. XSIP1 is essential for early neural gene expression and neural differentiation by suppression of BMP signaling. *Dev. Biol.* 2004;275(1) 258-267.
- [37] Nitta KR, Takahashi S, Haramoto Y, Fukuda M, Tanegashima K, Onuma Y, et al. The N-terminus zinc finger domain of *Xenopus* SIP1 is important for neural induction, but not for suppression of *Xbra* expression. *Int J Dev. Biol.* 2007;51(4) 321-325.
- [38] Weng Q, Chen Y, Wang H, Xu X, Yang B, He Q, et al. Dual-mode modulation of Smad signaling by Smad-interacting protein Sip1 is required for myelination in the central nervous system. *Neuron.* 2012;73(4) 713-728.
- [39] Drummond AE, Le MT, Ethier JF, Dyson M, Findlay JK. Expression and localization of activin receptors, Smads, and beta glycan to the postnatal rat ovary. *Endocrinology.* 2002;143(4) 1423-1433.
- [40] Findlay JK, Drummond AE, Dyson M, Baillie AJ, Robertson DM, Ethier JF. Production and actions of inhibin and activin during folliculogenesis in the rat. *Mol Cell Endocrinol.* 2001;180(1-2) 139-144.
- [41] Xu J, Oakley J, McGee EA. Stage-specific expression of Smad2 and Smad3 during folliculogenesis. *Biol Reprod.* 2002;66(6) 1571-1578.
- [42] Tian X, Halfhill AN, Diaz FJ. Localization of phosphorylated SMAD proteins in granulosa cells, oocytes and oviduct of female mice. *Gene Expr. Patterns.* 2010;10(2-3) 105-112.

- [43] Pangas SA, Rademaker AW, Fishman DA, Woodruff TK. Localization of the activin signal transduction components in normal human ovarian follicles: implications for autocrine and paracrine signaling in the ovary. *J Clin. Endocrinol. Metab.* 2002;87(6) 2644-2657.
- [44] Osterlund C, Fried G. TGFbeta receptor types I and II and the substrate proteins Smad 2 and 3 are present in human oocytes. *Mol Hum. Reprod.* 2000;6(6) 498-503.
- [45] Kuo FT, Fan K, Ambartsumyan G, Menon P, Ketefian A, tsi-Barnes IK, et al. Relative expression of genes encoding SMAD signal transduction factors in human granulosa cells is correlated with oocyte quality. *J Assist. Reprod. Genet.* 2011;28(10) 931-938.
- [46] Li X, Tripurani SK, James R, Pangas SA. Minimal fertility defects in mice deficient in oocyte-expressed Smad4. *Biol Reprod.* 2012;86(1) 1-6.
- [47] Tan Q, Balofsky A, Weisz K, Peng C. Role of activin, transforming growth factor-beta and bone morphogenetic protein 15 in regulating zebrafish oocyte maturation. *Comp Biochem. Physiol A Mol Integr. Physiol.* 2009;153(1) 18-23.
- [48] Snape A, Wylie CC, Smith JC, Heasman J. Changes in states of commitment of single animal pole blastomeres of *Xenopus laevis*. *Dev. Biol.* 1987;119(2) 503-510.
- [49] Dunn NR, Vincent SD, Oxburgh L, Robertson EJ, Bikoff EK. Combinatorial activities of Smad2 and Smad3 regulate mesoderm formation and patterning in the mouse embryo. *Development.* 2004;131(8) 1717-1728.
- [50] Jia S, Ren Z, Li X, Zheng Y, Meng A. smad2 and smad3 are required for mesendoderm induction by transforming growth factor-beta/nodal signals in zebrafish. *J Biol Chem.* 2008;283(4) 2418-2426.
- [51] Nomura M, Li E. Smad2 role in mesoderm formation, left-right patterning and craniofacial development. *Nature.* 1998;393(6687) 786-790.
- [52] Weinstein M, Yang X, Li C, Xu X, Gotay J, Deng CX. Failure of egg cylinder elongation and mesoderm induction in mouse embryos lacking the tumor suppressor smad2. *Proc Natl Acad Sci U S A.* 1998;95(16) 9378-9383.
- [53] Dupont S, Zacchigna L, Cordenonsi M, Soligo S, Adorno M, Rugge M, et al. Germ-layer specification and control of cell growth by Ectodermin, a Smad4 ubiquitin ligase. *Cell.* 2005;121(1) 87-99.
- [54] Howell M, Itoh F, Pierreux CE, Valgeirsdottir S, Itoh S, ten Dijke P, et al. *Xenopus* Smad4beta is the co-Smad component of developmentally regulated transcription factor complexes responsible for induction of early mesodermal genes. *Dev. Biol.* 1999;214(2) 354-369.
- [55] Hemmati-Brivanlou A, Melton D. Vertebrate neural induction. *Annu. Rev Neurosci.* 1997;20:43-60. 43-60.
- [56] Hemmati-Brivanlou A, Melton D. Vertebrate embryonic cells will become nerve cells unless told otherwise. *Cell.* 1997;88(1) 13-17.

- [57] Reversade B, Kuroda H, Lee H, Mays A, De Robertis EM. Depletion of Bmp2, Bmp4, Bmp7 and Spemann organizer signals induces massive brain formation in *Xenopus* embryos. *Development*. 2005;132(15) 3381-3392.
- [58] Linker C, Stern CD. Neural induction requires BMP inhibition only as a late step, and involves signals other than FGF and Wnt antagonists. *Development*. 2004;131(22) 5671-5681.
- [59] Chang C, Harland RM. Neural induction requires continued suppression of both Smad1 and Smad2 signals during gastrulation. *Development*. 2007;134(21) 3861-3872.
- [60] Yaguchi S, Yaguchi J, Burke RD. Sp-Smad2/3 mediates patterning of neurogenic ectoderm by nodal in the sea urchin embryo. *Dev. Biol.* 2007;302(2) 494-503.
- [61] Jia S, Wu D, Xing C, Meng A. Smad2/3 activities are required for induction and patterning of the neuroectoderm in zebrafish. *Dev. Biol.* 2009;333(2) 273-284.
- [62] Delaune E, Lemaire P, Kodjabachian L. Neural induction in *Xenopus* requires early FGF signalling in addition to BMP inhibition. *Development*. 2005;132(2) 299-310.
- [63] Streit A, Berliner AJ, Papanayotou C, Sirulnik A, Stern CD. Initiation of neural induction by FGF signalling before gastrulation. *Nature*. 2000;406(6791) 74-78.
- [64] Wilson SI, Graziano E, Harland R, Jessell TM, Edlund T. An early requirement for FGF signalling in the acquisition of neural cell fate in the chick embryo. *Curr. Biol.* 2000;20;10(8) 421-429.
- [65] Di-Gregorio A, Sancho M, Stuckey DW, Crompton LA, Godwin J, Mishina Y, et al. BMP signalling inhibits premature neural differentiation in the mouse embryo. *Development*. 2007;134(18) 3359-3369.
- [66] Chambers SM, Fasano CA, Papapetrou EP, Tomishima M, Sadelain M, Studer L. Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. *Nat Biotechnol.* 2009;27(3) 275-280.
- [67] Ganat YM, Calder EL, Kriks S, Nelander J, Tu EY, Jia F, et al. Identification of embryonic stem cell-derived midbrain dopaminergic neurons for engraftment. *J Clin. Invest.* 2012;122(8) 2928-2939.
- [68] Yaguchi S, Yaguchi J, Wei Z, Jin Y, Angerer LM, Inaba K. Fez function is required to maintain the size of the animal plate in the sea urchin embryo. *Development*. 2011;138(19) 4233-4243.
- [69] Juriloff DM, Harris MJ. Mouse models for neural tube closure defects. *Hum. Mol Genet.* 2000;9(6) 993-1000.
- [70] Stottmann RW, Berrong M, Matta K, Choi M, Klingensmith J. The BMP antagonist Noggin promotes cranial and spinal neurulation by distinct mechanisms. *Dev. Biol.* 2006;295(2) 647-663.

- [71] Eom DS, Amarnath S, Fogel JL, Agarwala S. Bone morphogenetic proteins regulate neural tube closure by interacting with the apicobasal polarity pathway. *Development*. 2011;138(15) 3179-3188.
- [72] Eom DS, Amarnath S, Fogel JL, Agarwala S. Bone morphogenetic proteins regulate hinge point formation during neural tube closure by dynamic modulation of apicobasal polarity. *Birth Defects Res A Clin. Mol Teratol*. 2012; 10.
- [73] Ybot-Gonzalez P, Gaston-Massuet C, Girdler G, Klingensmith J, Arkell R, Greene ND, et al. Neural plate morphogenesis during mouse neurulation is regulated by antagonism of Bmp signalling. *Development*. 2007;134(17) 3203-3211.
- [74] Chizhikov VV, Millen KJ. Mechanisms of roof plate formation in the vertebrate CNS. *Nat Rev Neurosci*. 2004;5(10) 808-812.
- [75] Caspary T, Anderson KV. Patterning cell types in the dorsal spinal cord: what the mouse mutants say. *Nat Rev Neurosci*. 2003;4(4) 289-297.
- [76] Dessaud E, McMahon AP, Briscoe J. Pattern formation in the vertebrate neural tube: a sonic hedgehog morphogen-regulated transcriptional network. *Development*. 2008;135(15) 2489-2503.
- [77] Perron JC, Dodd J. Inductive specification and axonal orientation of spinal neurons mediated by divergent bone morphogenetic protein signaling pathways. *Neural Dev*. 2011;6:36. 36.
- [78] Perron JC, Dodd J. Structural distinctions in BMPs underlie divergent signaling in spinal neurons. *Neural Dev*. 2012;7:16. 16.
- [79] Le DG, Garcia-Campmany L, Rabadan MA, Ferronha T, Tozer S, Briscoe J, et al. Canonical BMP7 activity is required for the generation of discrete neuronal populations in the dorsal spinal cord. *Development*. 2012;139(2) 259-268.
- [80] Hu Q, Ueno N, Behringer RR. Restriction of BMP4 activity domains in the developing neural tube of the mouse embryo. *EMBO Rep*. 2004;5(7) 734-739.
- [81] Xie Z, Chen Y, Li Z, Bai G, Zhu Y, Yan R, et al. Smad6 promotes neuronal differentiation in the intermediate zone of the dorsal neural tube by inhibition of the Wnt/beta-catenin pathway. *Proc Natl Acad Sci U S A*. 2011;108(29) 12119-12124.
- [82] Hazen VM, Phan KD, Hudiburgh S, Butler SJ. Inhibitory Smads differentially regulate cell fate specification and axon dynamics in the dorsal spinal cord. *Dev. Biol*. 2011;356(2) 566-575.
- [83] Garcia-Campmany L, Marti E. The TGF $\beta$  intracellular effector Smad3 regulates neuronal differentiation and cell fate specification in the developing spinal cord. *Development* 2007;134(1) 65-75.
- [84] Heeg-Truesdell E, LaBonne C. A slug, a fox, a pair of sox: transcriptional responses to neural crest inducing signals. *Birth Defects Res C. Embryo. Today*. 2004;72(2) 124-139.



- [85] Faure S, de Santa BP, Roberts DJ, Whitman M. Endogenous patterns of BMP signaling during early chick development. *Dev. Biol.* 2002;244(1) 44-65.
- [86] Liem KF, Jr., Tremml G, Jessell TM. A role for the roof plate and its resident TGFbeta-related proteins in neuronal patterning in the dorsal spinal cord. *Cell.* 1997;91(1) 127-138.
- [87] Liem KF, Jr., Tremml G, Roelink H, Jessell TM. Dorsal differentiation of neural plate cells induced by BMP-mediated signals from epidermal ectoderm. *Cell.* 1995;82(6) 969-979.
- [88] Linker C, de A, I, Papanayotou C, Stower M, Sabado V, Ghorani E, et al. Cell communication with the neural plate is required for induction of neural markers by BMP inhibition: evidence for homeogenetic induction and implications for *Xenopus* animal cap and chick explant assays. *Dev. Biol.* 2009;327(2) 478-486.
- [89] Patthey C, Edlund T, Gunhaga L. Wnt-regulated temporal control of BMP exposure directs the choice between neural plate border and epidermal fate. *Development.* 2009;136(1) 73-83.
- [90] Patthey C, Gunhaga L. Specification and regionalisation of the neural plate border. *Eur. J Neurosci.* 2011;34(10) 1516-1528.
- [91] Streit A, Stern CD. Establishment and maintenance of the border of the neural plate in the chick: involvement of FGF and BMP activity. *Mech. Dev.* 1999;82(1-2) 51-66.
- [92] Chalazonitis A, D'Autreaux F, Guha U, Pham TD, Faure C, Chen JJ, et al. Bone morphogenetic protein-2 and -4 limit the number of enteric neurons but promote development of a TrkC-expressing neurotrophin-3-dependent subset. *J. Neurosci.* 2004;24(17) 4266-4282.
- [93] Chalazonitis A, D'Autreaux F, Pham TD, Kessler JA, Gershon MD. Bone morphogenetic proteins regulate enteric gliogenesis by modulating ErbB3 signaling. *Dev. Biol.* 2011;350(1) 64-79.
- [94] Harada J, Kokura K, Kanei-Ishii C, Nomura T, Khan MM, Kim Y, et al. Requirement of the co-repressor homeodomain-interacting protein kinase 2 for ski-mediated inhibition of bone morphogenetic protein-induced transcriptional activation. *J. Biol. Chem.* 2003;278(40) 38998-39005.
- [95] Chalazonitis A, Tang AA, Shang Y, Pham TD, Hsieh I, Setlik W, et al. Homeodomain interacting protein kinase 2 regulates postnatal development of enteric dopaminergic neurons and glia via BMP signaling. *J Neurosci.* 2011;31(39) 13746-13757.
- [96] Zhang J, Pho V, Bonasera SJ, Holtzman J, Tang AT, Hellmuth J, et al. Essential function of HIPK2 in TGFbeta-dependent survival of midbrain dopamine neurons. *Nat Neurosci.* 2007;10(1) 77-86.

- [97] Anitha M, Shahnavaz N, Qayed E, Joseph I, Gossrau G, Mwangi S, et al. BMP2 promotes differentiation of nitrergic and catecholaminergic enteric neurons through a Smad1-dependent pathway. *Am. J Physiol Gastrointest. Liver Physiol.* 2010;298(3) G375-G383.
- [98] Dore JJ, Crotty KL, Birren SJ. Inhibition of glial maturation by bone morphogenetic protein 2 in a neural crest-derived cell line. *Dev. Neurosci.* 2005;27(1) 37-48.
- [99] Dore JJ, Dewitt JC, Setty N, Donald MD, Joo E, Chesarone MA, et al. Multiple Signaling Pathways Converge to Regulate Bone-Morphogenetic-Protein-Dependent Glial Gene Expression. *Dev. Neurosci.* 2009.
- [100] Gross RE, Mehler MF, Mabie PC, Zang Z, Santschi L, Kessler JA. Bone morphogenetic proteins promote astroglial lineage commitment by mammalian subventricular zone progenitor cells. *Neuron.* 1996;17(4) 595-606.
- [101] Gomes WA, Mehler MF, Kessler JA. Transgenic overexpression of BMP4 increases astroglial and decreases oligodendroglial lineage commitment. *Dev. Biol.* 2003;255(1) 164-177.
- [102] Stipursky J, Francis D, Gomes FC. Activation of MAPK/PI3K/SMAD Pathways by TGF-beta(1) Controls Differentiation of Radial Glia into Astrocytes in vitro. *Dev. Neurosci.* 2012;34(1) 68-81.
- [103] Malatesta P, Hack MA, Hartfuss E, Kettenmann H, Klinkert W, Kirchhoff F, et al. Neuronal or glial progeny: regional differences in radial glia fate. *Neuron.* 2003;37(5) 751-764.
- [104] Stipursky J, Spohr TC, Sousa VO, Gomes FC. Neuron-Astroglial Interactions in Cell-Fate Commitment and Maturation in the Central Nervous System. *Neurochem. Res.* 2012.
- [105] Stipursky J, Gomes FC. TGF-beta1/SMAD signaling induces astrocyte fate commitment in vitro: implications for radial glia development. *Glia.* 2007;55(10) 1023-1033.
- [106] Gomes FC, Paulin D, Moura N, V. Glial fibrillary acidic protein (GFAP): modulation by growth factors and its implication in astrocyte differentiation. *Braz. J. Med. Biol. Res.* 1999;32(5) 619-631.
- [107] Sousa VO, Romao L, Neto VM, Gomes FC. Glial fibrillary acidic protein gene promoter is differently modulated by transforming growth factor-beta 1 in astrocytes from distinct brain regions. *Eur. J. Neurosci.* 2004;19(7) 1721-1730.
- [108] Gomes FC, Garcia-Abreu J, Galou M, Paulin D, Moura N, V. Neurons induce GFAP gene promoter of cultured astrocytes from transgenic mice. *Glia.* 1999;26(2) 97-108.
- [109] de Sampaio e Spohr T, Martinez R, da Silva EF, Neto VM, Gomes FC. Neuro-glia interaction effects on GFAP gene: a novel role for transforming growth factor-beta1. *Eur. J. Neurosci.* 2002;16(11) 2059-2069.

- [110] Imura T, Tane K, Toyoda N, Fushiki S. Endothelial cell-derived bone morphogenetic proteins regulate glial differentiation of cortical progenitors. *Eur. J Neurosci.* 2008;27(7) 1596-1606.
- [111] Mekki-Dauriac S, Agius E, Kan P, Cochard P. Bone morphogenetic proteins negatively control oligodendrocyte precursor specification in the chick spinal cord. *Development.* 2002;129(22) 5117-5130.
- [112] Voumvourakis KI, Antonelou RC, Kitsos DK, Stamboulis E, Tsiodras S. TGF-beta/BMPs: crucial crossroad in neural autoimmune disorders. *Neurochem. Int.* 2011;59(5) 542-550.
- [113] Hampton DW, Asher RA, Kondo T, Steeves JD, Ramer MS, Fawcett JW. A potential role for bone morphogenetic protein signalling in glial cell fate determination following adult central nervous system injury in vivo. *Eur. J Neurosci.* 2007;26(11) 3024-3035.
- [114] Bilican B, Fiore-Herich C, Compston A, Allen ND, Chandran S. Induction of Olig2 precursors by FGF involves BMP signalling blockade at the Smad level. *PLoS. ONE.* 2008;3(8) e2863.
- [115] Agius E, Decker Y, Soukkaiech C, Soula C, Cochard P. Role of BMPs in controlling the spatial and temporal origin of GFAP astrocytes in the embryonic spinal cord. *Dev. Biol.* 2010;344(2) 611-620.
- [116] Buckwalter MS, Yamane M, Coleman BS, Ormerod BK, Chin JT, Palmer T, et al. Chronically Increased Transforming Growth Factor- $\beta$ 1 Strongly Inhibits Hippocampal Neurogenesis in Aged Mice. *Am. J. Pathol.* 2006;169(1) 154-164.
- [117] Wachs FP, Winner B, Couillard-Despres S, Schiller T, Aigner R, Winkler J, et al. Transforming growth factor-beta1 is a negative modulator of adult neurogenesis. *J. Neuropathol. Exp. Neurol.* 2006;65(4) 358-370.
- [118] Vogel T, Ahrens S, Buttner N, Kriegelstein K. Transforming growth factor beta promotes neuronal cell fate of mouse cortical and hippocampal progenitors in vitro and in vivo: identification of Nedd9 as an essential signaling component. *Cereb. Cortex.* 2010;20(3) 661-671.
- [119] Garcia-Campmany L, Marti E. The TGFbeta intracellular effector Smad3 regulates neuronal differentiation and cell fate specification in the developing spinal cord. *Development.* 2007;134(1) 65-75.
- [120] Lu J, Wu Y, Sousa N, Almeida OF. SMAD pathway mediation of BDNF and TGF beta 2 regulation of proliferation and differentiation of hippocampal granule neurons. *Development.* 2005;132(14) 3231-3242.
- [121] Battista D, Ferrari CC, Gage FH, Pitossi FJ. Neurogenic niche modulation by activated microglia: transforming growth factor beta increases neurogenesis in the adult dentate gyrus. *Eur. J. Neurosci.* 2006;23(1) 83-93.

- [122] Ma M, Ma Y, Yi X, Guo R, Zhu W, Fan X, et al. Intranasal delivery of transforming growth factor-beta1 in mice after stroke reduces infarct volume and increases neurogenesis in the subventricular zone. *BMC. Neurosci.* 2008;9:117. 117.
- [123] Wang Y, Symes AJ. Smad3 deficiency reduces neurogenesis in adult mice. *J Mol Neurosci.* 2010;41(3) 383-396.
- [124] Yang X, Letterio JJ, Lechleider RJ, Chen L, Hayman R, Gu H, et al. Targeted disruption of SMAD3 results in impaired mucosal immunity and diminished T cell responsiveness to TGF-beta. *EMBO J.* 1999;18(5) 1280-1291.
- [125] Roussa E, Wiehle M, Dunker N, Becker-Katins S, Oehlke O, Kriegelstein K. Transforming growth factor beta is required for differentiation of mouse mesencephalic progenitors into dopaminergic neurons in vitro and in vivo: ectopic induction in dorsal mesencephalon. *Stem Cells.* 2006;24(9) 2120-2129.
- [126] Kendall SE, Battelli C, Irwin S, Mitchell JG, Glackin CA, Verdi JM. NRAGE mediates p38 activation and neural progenitor apoptosis via the bone morphogenetic protein signaling cascade. *Mol Cell Biol.* 2005;25(17) 7711-7724.
- [127] Gambaro K, Aberdam E, Virolle T, Aberdam D, Rouleau M. BMP-4 induces a Smad-dependent apoptotic cell death of mouse embryonic stem cell-derived neural precursors. *Cell Death. Differ.* 2006;13(7) 1075-1087.
- [128] Brederlau A, Faigle R, Kaplan P, Odin P, Funa K. Bone morphogenetic proteins but not growth differentiation factors induce dopaminergic differentiation in mesencephalic precursors. *Mol Cell Neurosci.* 2002;21(3) 367-378.
- [129] Toulouse A, Collins GC, Sullivan AM. Neurotrophic effects of growth/differentiation factor 5 in a neuronal cell line. *Neurotox. Res.* 2012;21(3) 256-265.
- [130] Lindvall O. Dopaminergic neurons for Parkinson's therapy. *Nat Biotechnol.* 2012;30(1) 56-58.
- [131] Rios I, Alvarez-Rodriguez R, Marti E, Pons S. Bmp2 antagonizes sonic hedgehog-mediated proliferation of cerebellar granule neurones through Smad5 signalling. *Development.* 2004;131(13) 3159-3168.
- [132] Zhou YX, Zhao M, Li D, Shimazu K, Sakata K, Deng CX, et al. Cerebellar deficits and hyperactivity in mice lacking Smad4. *J Biol Chem.* 2003;278(43) 42313-42320.
- [133] Morikawa Y, Zehir A, Maska E, Deng C, Schneider MD, Mishina Y, et al. BMP signaling regulates sympathetic nervous system development through Smad4-dependent and -independent pathways. *Development.* 2009;136(21) 3575-3584.
- [134] Lopez-Coviella I, Mellott TM, Kovacheva VP, Berse B, Slack BE, Zemelko V, et al. Developmental pattern of expression of BMP receptors and Smads and activation of Smad1 and Smad5 by BMP9 in mouse basal forebrain. *Brain Res.* 2006;1088(1) 49-56.

- [135] Williams S, Souchelnytskyi S, Danik M. TGFbeta2 mediates rapid inhibition of calcium influx in identified cholinergic basal forebrain neurons. *Biochem. Biophys. Res Commun.* 2002;290(4) 1321-1327.
- [136] George O, Parducz A, Dupret D, Kharouby M, Le MM, Piazza PV, et al. Smad-dependent alterations of PPT cholinergic neurons as a pathophysiological mechanism of age-related sleep-dependent memory impairments. *Neurobiol Aging.* 2006;27(12) 1848-1858.
- [137] Lopez-Coviella I, Berse B, Krauss R, Thies RS, Blusztajn JK. Induction and maintenance of the neuronal cholinergic phenotype in the central nervous system by BMP-9. *Science.* 2000;289(5477) 313-316.
- [138] Schnitzler AC, Mellott TJ, Lopez-Coviella I, Tallini YN, Kotlikoff MI, Follettie MT, et al. BMP9 (bone morphogenetic protein 9) induces NGF as an autocrine/paracrine cholinergic trophic factor in developing basal forebrain neurons. *J Neurosci.* 2010;30(24) 8221-8228.
- [139] Lopez-Coviella I, Mellott TJ, Schnitzler AC, Blusztajn JK. BMP9 protects septal neurons from axotomy-evoked loss of cholinergic phenotype. *PLoS. ONE.* 2011;6(6) e21166.
- [140] Cho G, Lim Y, Golden JA. XLMR candidate mouse gene, *Zcchc12* (*Sizn1*) is a novel marker of Cajal-Retzius cells. *Gene Expr. Patterns.* 2011;11(3-4) 216-220.
- [141] Cho G, Lim Y, Zand D, Golden JA. *Sizn1* is a novel protein that functions as a transcriptional coactivator of bone morphogenic protein signaling. *Mol Cell Biol.* 2008;28(5) 1565-1572.
- [142] Yanagisawa M, Nakashima K, Takeda K, Ochiai W, Takizawa T, Ueno M, et al. Inhibition of BMP2-induced, TAK1 kinase-mediated neurite outgrowth by Smad6 and Smad7. *Genes Cells.* 2001;6(12) 1091-1099.
- [143] Iwasaki S, Hattori A, Sato M, Tsujimoto M, Kohno M. Characterization of the bone morphogenetic protein-2 as a neurotrophic factor. Induction of neuronal differentiation of PC12 cells in the absence of mitogen-activated protein kinase activation. *J Biol Chem.* 1996;19;271(29) 17360-17365.
- [144] Iwasaki S, Iguchi M, Watanabe K, Hoshino R, Tsujimoto M, Kohno M. Specific activation of the p38 mitogen-activated protein kinase signaling pathway and induction of neurite outgrowth in PC12 cells by bone morphogenetic protein-2. *J Biol Chem.* 1999;274(37) 26503-26510.
- [145] Ebisawa T, Fukuchi M, Murakami G, Chiba T, Tanaka K, Imamura T, et al. Smurf1 interacts with transforming growth factor-beta type I receptor through Smad7 and induces receptor degradation. *J. Biol. Chem.* 2001;276(16) 12477-12480.
- [146] Althini S, Usoskin D, Kylberg A, Kaplan PL, Ebendal T. Blocked MAP kinase activity selectively enhances neurotrophic growth responses. *Mol Cell Neurosci.* 2004;25(2) 345-354.

- [147] Benavente F, Pinto C, Parada M, Henriquez JP, Osses N. Bone morphogenetic protein 2 inhibits neurite outgrowth of motor neuron-like NSC-34 cells and up-regulates its type II receptor. *J Neurochem.* 2012;122(3) 594-604.
- [148] Ueberham U, Ueberham E, Gruschka H, Arendt T. Altered subcellular location of phosphorylated Smads in Alzheimer's disease. *Eur. J Neurosci.* 2006;24(8) 2327-2334.
- [149] Ueberham U, Lange P, Ueberham E, Bruckner MK, Hartlage-Rubsamen M, Pannicke T, et al. Smad2 isoforms are differentially expressed during mouse brain development and aging. *Int J Dev. Neurosci.* 2009;27(5) 501-510.
- [150] Stegmuller J, Huynh MA, Yuan Z, Konishi Y, Bonni A. TGFbeta-Smad2 signaling regulates the Cdh1-APC/SnoN pathway of axonal morphogenesis. *J Neurosci.* 2008;20;28(8) 1961-1969.
- [151] Stegmuller J, Konishi Y, Huynh MA, Yuan Z, Dibacco S, Bonni A. Cell-intrinsic regulation of axonal morphogenesis by the Cdh1-APC target SnoN. *Neuron.* 2006;50(3) 389-400.
- [152] Huynh MA, Ikeuchi Y, Netherton S, de IT-U, Kanadia R, Stegmuller J, et al. An isoform-specific SnoN1-FOXO1 repressor complex controls neuronal morphogenesis and positioning in the mammalian brain. *Neuron.* 2011;69(5) 930-944.
- [153] Ikeuchi Y, Stegmuller J, Netherton S, Huynh MA, Masu M, Frank D, et al. A SnoN-Ccd1 pathway promotes axonal morphogenesis in the mammalian brain. *J Neurosci.* 2009;29(13) 4312-4321.
- [154] Stegmuller J, Bonni A. Moving past proliferation: new roles for Cdh1-APC in postmitotic neurons. *Trends Neurosci.* 2005;28(11) 596-601.
- [155] Konishi Y, Stegmuller J, Matsuda T, Bonni S, Bonni A. Cdh1-APC controls axonal growth and patterning in the mammalian brain. *Science* 2004;303(5660) 1026-1030.
- [156] Makwana M, Raivich G. Molecular mechanisms in successful peripheral regeneration. *FEBS J.* 2005;272(11) 2628-2638.
- [157] Goldberg JL, Klassen MP, Hua Y, Barres BA. Amacrine-signaled loss of intrinsic axon growth ability by retinal ganglion cells. *Science.* 2002;296(5574) 1860-1864.
- [158] Zou H, Ho C, Wong K, Tessier-Lavigne M. Axotomy-induced Smad1 activation promotes axonal growth in adult sensory neurons. *J Neurosci.* 2009;29(22) 7116-7123.
- [159] Parikh P, Hao Y, Hosseinkhani M, Patil SB, Huntley GW, Tessier-Lavigne M, et al. Regeneration of axons in injured spinal cord by activation of bone morphogenetic protein/Smad1 signaling pathway in adult neurons. *Proc Natl Acad Sci U S A.* 2011;108(19) E99-107.
- [160] Zhu S, Wang W, Clarke DC, Liu X. Activation of Mps1 promotes transforming growth factor-beta-independent Smad signaling. *J Biol Chem.* 2007;282(25) 18327-18338.
- [161] Ma CH, Brenner GJ, Omura T, Samad OA, Costigan M, Inquimbert P, et al. The BMP coreceptor RGMb promotes while the endogenous BMP antagonist noggin reduces

- neurite outgrowth and peripheral nerve regeneration by modulating BMP signaling. *J Neurosci.* 2011;31(50) 18391-18400.
- [162] Serpe M, O'Connor MB. The metalloprotease tolloid-related and its TGF-beta-like substrate Dawdle regulate *Drosophila* motoneuron axon guidance. *Development.* 2006;133(24) 4969-4979.
- [163] Hodge LK, Klassen MP, Han BX, Yiu G, Hurrell J, Howell A, et al. Retrograde BMP signaling regulates trigeminal sensory neuron identities and the formation of precise face maps. *Neuron* 2007;55(4) 572-586.
- [164] Ting CY, Herman T, Yonekura S, Gao S, Wang J, Serpe M, et al. Tiling of r7 axons in the *Drosophila* visual system is mediated both by transduction of an activin signal to the nucleus and by mutual repulsion. *Neuron.* 2007;56(5) 793-806.
- [165] McCabe BD, Marques G, Haghghi AP, Fetter RD, Crotty ML, Haerry TE, et al. The BMP homolog Gbb provides a retrograde signal that regulates synaptic growth at the *Drosophila* neuromuscular junction. *Neuron.* 2003;39(2) 241-254.
- [166] Ji SJ, Jaffrey SR. Intra-axonal translation of SMAD1/5/8 mediates retrograde regulation of trigeminal ganglia subtype specification. *Neuron.* 2012;74(1) 95-107.
- [167] Takatoh J, Wang F. Axonally translated SMADs link up BDNF and retrograde BMP signaling. *Neuron.* 2012;74(1) 3-5.
- [168] Jiang Y, McLennan IS, Koishi K, Hendry IA. Transforming growth factor-beta 2 is anterogradely and retrogradely transported in motoneurons and up-regulated after nerve injury. *Neuroscience* 2000;97(4) 735-742.
- [169] Batut J, Howell M, Hill CS. Kinesin-mediated transport of Smad2 is required for signaling in response to TGF-beta ligands. *Dev. Cell* 2007;12(2) 261-274.
- [170] Jin Q, Gao G, Mulder KM. Requirement of a dynein light chain in TGFbeta/Smad3 signaling. *J. Cell Physiol* 2009;221(3) 707-715.
- [171] Eaton BA, Fetter RD, Davis GW. Dynactin is necessary for synapse stabilization. *Neuron.* 2002;34(5) 729-741.
- [172] Allan DW, St Pierre SE, Miguel-Aliaga I, Thor S. Specification of neuropeptide cell identity by the integration of retrograde BMP signaling and a combinatorial transcription factor code. *Cell* 2003;113(1) 73-86.
- [173] Goold CP, Davis GW. The BMP ligand Gbb gates the expression of synaptic homeostasis independent of synaptic growth control. *Neuron.* 2007;56(1) 109-123.
- [174] Romao LF, Sousa VO, Neto VM, Gomes FC. Glutamate activates GFAP gene promoter from cultured astrocytes through TGF-beta1 pathways. *J Neurochem.* 2008;106(2) 746-756.

- [175] Susarla BT, Laing ED, Yu P, Katagiri Y, Geller HM, Symes AJ. Smad proteins differentially regulate transforming growth factor-beta-mediated induction of chondroitin sulfate proteoglycans. *J Neurochem.* 2011;119(4) 868-878.
- [176] Schachtrup C, Ryu JK, Helmrick MJ, Vagena E, Galanakis DK, Degen JL, et al. Fibrinogen triggers astrocyte scar formation by promoting the availability of active TGF-beta after vascular damage. *J Neurosci.* 2010;30(17) 5843-5854.
- [177] Hellal F, Hurtado A, Ruschel J, Flynn KC, Laskowski CJ, Umlauf M, et al. Microtubule stabilization reduces scarring and causes axon regeneration after spinal cord injury. *Science.* 2011;331(6019) 928-931.
- [178] Dummula K, Vinukonda G, Chu P, Xing Y, Hu F, Maik S, et al. Bone morphogenetic protein inhibition promotes neurological recovery after intraventricular hemorrhage. *J Neurosci.* 2011;31(34) 12068-12082.
- [179] Sweeney ST, Davis GW. Unrestricted synaptic growth in spinster-a late endosomal protein implicated in TGF-beta-mediated synaptic growth regulation. *Neuron.* 2002;36(3) 403-416.
- [180] Rawson JM, Lee M, Kennedy EL, Selleck SB. Drosophila neuromuscular synapse assembly and function require the TGF-beta type I receptor saxophone and the transcription factor Mad. *J Neurobiol.* 2003;55(2) 134-150.
- [181] Aberle H, Haghighi AP, Fetter RD, McCabe BD, Magalhaes TR, Goodman CS. wishful thinking encodes a BMP type II receptor that regulates synaptic growth in Drosophila. *Neuron.* 2002;33(4) 545-558.
- [182] Marques G, Bao H, Haerry TE, Shimell MJ, Duchek P, Zhang B, et al. The Drosophila BMP type II receptor Wishful Thinking regulates neuromuscular synapse morphology and function. *Neuron.* 2002;33(4) 529-543.
- [183] McCabe BD, Hom S, Aberle H, Fetter RD, Marques G, Haerry TE, et al. Highwire regulates presynaptic BMP signaling essential for synaptic growth. *Neuron.* 2004;41(6) 891-905.
- [184] Wan HI, DiAntonio A, Fetter RD, Bergstrom K, Strauss R, Goodman CS. Highwire regulates synaptic growth in Drosophila. *Neuron.* 2000;26(2) 313-329.
- [185] Dudu V, Bittig T, Entchev E, Kicheva A, Julicher F, Gonzalez-Gaitan M. Postsynaptic mad signaling at the Drosophila neuromuscular junction. *Curr. Biol.* 2006;16(7) 625-635.
- [186] Ellis JE, Parker L, Cho J, Arora K. Activin signaling functions upstream of Gbb to regulate synaptic growth at the Drosophila neuromuscular junction. *Dev. Biol.* 2010;342(2) 121-133.
- [187] Nahm M, Kim S, Paik SK, Lee M, Lee S, Lee ZH, et al. dCIP4 (Drosophila Cdc42-interacting protein 4) restrains synaptic growth by inhibiting the secretion of the retrograde Glass bottom boat signal. *J Neurosci.* 2010;30(24) 8138-8150.



- [188] Dickson DW. Neuropathology of non-Alzheimer degenerative disorders. *Int. J. Clin. Exp. Pathol.* 2009;3(1) 1-23.
- [189] Arendt T. Synaptic plasticity and cell cycle activation in neurons are alternative effector pathways: the 'Dr. Jekyll and Mr. Hyde concept' of Alzheimer's disease or the yin and yang of neuroplasticity. *Prog. Neurobiol.* 2003;71(2-3) 83-248.
- [190] Lee HG, Ueda M, Zhu X, Perry G, Smith MA. Ectopic expression of phospho-Smad2 in Alzheimer's disease: uncoupling of the transforming growth factor-beta pathway? *J Neurosci Res.* 2006;84(8) 1856-1861.
- [191] Chalmers KA, Love S. Neurofibrillary tangles may interfere with Smad 2/3 signaling in neurons. *J Neuropathol. Exp. Neurol.* 2007;66(2) 158-167.
- [192] Ueberham U, Hilbrich I, Ueberham E, Rohn S, Glockner P, Dietrich K, et al. Transcriptional control of cell cycle-dependent kinase 4 by Smad proteins-implications for Alzheimer's disease. *Neurobiol. Aging* 2012;33(12) 2827-2840.
- [193] Lee MH, Lin SR, Chang JY, Schultz L, Heath J, Hsu LJ, et al. TGF-beta induces TIAF1 self-aggregation via type II receptor-independent signaling that leads to generation of amyloid beta plaques in Alzheimer's disease. *Cell Death. Dis.* 2010;1:e110. e110.
- [194] Metzals J, Robitaille Y, Houghton S, Gauthier S, Leblanc R. Paired helical filaments and the cytoplasmic-nuclear interface in Alzheimer's disease. *J. Neurocytol.* 1988;17(6) 827-833.
- [195] Sheffield LG, Miskiewicz HB, Tannenbaum LB, Mirra SS. Nuclear pore complex proteins in Alzheimer disease. *J. Neuropathol. Exp. Neurol.* 2006;65(1) 45-54.
- [196] Worman HJ. Inner nuclear membrane and regulation of Smad-mediated signaling. *Biochim. Biophys. Acta* 2006;1761(5-6) 626-631.
- [197] Xu L, Kang Y, Col S, Massague J. Smad2 nucleocytoplasmic shuttling by nucleoporins CAN/Nup214 and Nup153 feeds TGFbeta signaling complexes in the cytoplasm and nucleus. *Mol. Cell* 2002;10(2) 271-282.
- [198] Ellgaard L, Ruddock LW. The human protein disulphide isomerase family: substrate interactions and functional properties. *EMBO Rep.* 2005;6(1) 28-32.
- [199] Cataldo AM, Petanceska S, Terio NB, Peterhoff CM, Durham R, Mercken M, et al. Abeta localization in abnormal endosomes: association with earliest Abeta elevations in AD and Down syndrome. *Neurobiol. Aging* 2004;25(10) 1263-1272.
- [200] Walter J, Fluhrer R, Hartung B, Willem M, Kaether C, Capell A, et al. Phosphorylation regulates intracellular trafficking of beta-secretase. *J. Biol. Chem.* 2001;276(18) 14634-14641.
- [201] Hayes S, Chawla A, Corvera S. TGF beta receptor internalization into EEA1-enriched early endosomes: role in signaling to Smad2. *J. Cell Biol.* 2002;158(7) 1239-1249.

- [202] Penheiter SG, Mitchell H, Garamszegi N, Edens M, Dore JJ, Jr., Leof EB. Internalization-dependent and -independent requirements for transforming growth factor beta receptor signaling via the Smad pathway. *Mol. Cell Biol.* 2002;22(13) 4750-4759.
- [203] Runyan CE, Schnaper HW, Poncelet AC. The role of internalization in transforming growth factor beta1-induced Smad2 association with Smad anchor for receptor activation (SARA) and Smad2-dependent signaling in human mesangial cells. *J. Biol. Chem.* 2005;280(9) 8300-8308.
- [204] Gagnon E, Duclos S, Rondeau C, Chevet E, Cameron PH, Steele-Mortimer O, et al. Endoplasmic reticulum-mediated phagocytosis is a mechanism of entry into macrophages. *Cell* 2002;110(1) 119-131.
- [205] Touret N, Paroutis P, Grinstein S. The nature of the phagosomal membrane: endoplasmic reticulum versus plasmalemma. *J. Leukoc. Biol.* 2005;77(6) 878-885.
- [206] Ueberham U, Arendt T. The expression of cell cycle proteins in neurons and its relevance for Alzheimer's disease. *Curr. Drug Targets. CNS. Neurol. Disord.* 2005;4(3) 293-306.
- [207] Baig S, van HZ, Love S. Tau hyperphosphorylation affects Smad 2/3 translocation. *Neuroscience.* 2009;163(2) 561-570.
- [208] Cowan CM, Bossing T, Page A, Shepherd D, Mudher A. Soluble hyper-phosphorylated tau causes microtubule breakdown and functionally compromises normal tau in vivo. *Acta Neuropathol.* 2010;120(5) 593-604.
- [209] Connor B, Young D, Lawlor P, Gai W, Waldvogel H, Faull RL, et al. Trk receptor alterations in Alzheimer's disease. *Brain Res. Mol. Brain Res.* 1996;42(1) 1-17.
- [210] Hock C, Heese K, Hulette C, Rosenberg C, Otten U. Region-specific neurotrophin imbalances in Alzheimer disease: decreased levels of brain-derived neurotrophic factor and increased levels of nerve growth factor in hippocampus and cortical areas. *Arch. Neurol.* 2000;57(6) 846-851.
- [211] Arendt T, Holzer M, Stobe A, Gartner U, Luth HJ, Bruckner MK, et al. Activated mitogenic signaling induces a process of dedifferentiation in Alzheimer's disease that eventually results in cell death. *Ann. N. Y. Acad. Sci.* 2000;920 249-255.
- [212] Moustakas A, Pardali K, Gaal A, Heldin CH. Mechanisms of TGF-beta signaling in regulation of cell growth and differentiation. *Immunol. Lett.* 2002;82(1-2) 85-91.
- [213] Bachman KE, Park BH. Dual nature of TGF-beta signaling: tumor suppressor vs. tumor promoter. *Curr. Opin. Oncol.* 2005;17(1) 49-54.
- [214] Peng B, Fleming JB, Breslin T, Grau AM, Fojioka S, Abbruzzese JL, et al. Suppression of tumorigenesis and induction of p15(ink4b) by Smad4/DPC4 in human pancreatic cancer cells. *Clin. Cancer Res.* 2002;8(11) 3628-3638.

- [215] Motta M, Imbesi R, Di RM, Stivala F, Malaguarnera L. Altered plasma cytokine levels in Alzheimer's disease: correlation with the disease progression. *Immunol. Lett.* 2007;114(1) 46-51.
- [216] Ueberham U, Ueberham E, Gruschka H, Arendt T. Connective tissue growth factor in Alzheimer's disease. *Neuroscience* 2003;116(1) 1-6.
- [217] Docagne F, Gabriel C, Lebourrier N, Lesne S, Hommet Y, Plawinski L, et al. Sp1 and Smad transcription factors co-operate to mediate TGF-beta-dependent activation of amyloid-beta precursor protein gene transcription. *Biochem. J.* 2004;383(Pt 2) 393-399.
- [218] Burton T, Liang B, Dibrov A, Amara F. Transforming growth factor-beta-induced transcription of the Alzheimer beta-amyloid precursor protein gene involves interaction between the CTCF-complex and Smads. *Biochem. Biophys. Res Commun.* 2002;19;295(3) 713-723.
- [219] Lee EO, Kang JL, Chong YH. The amyloid-beta peptide suppresses transforming growth factor-beta1-induced matrix metalloproteinase-2 production via Smad7 expression in human monocytic THP-1 cells. *J Biol Chem.* 2005;280(9) 7845-7853.
- [220] Wyss-Coray T, Lin C, Yan F, Yu GQ, Rohde M, McConlogue L, et al. TGF-beta1 promotes microglial amyloid-beta clearance and reduces plaque burden in transgenic mice. *Nat. Med.* 2001;7(5) 612-618.
- [221] Tichauer JE, von Bernhardi R. Transforming growth factor-beta stimulates beta amyloid uptake by microglia through Smad3-dependent mechanisms. *J Neurosci Res.* 2012;90(10) 1970-1980.
- [222] Wisniewski HM, Wegiel J, Wang KC, Kujawa M, Lach B. Ultrastructural studies of the cells forming amyloid fibers in classical plaques. *Can. J Neurol Sci.* 1989;16(4 Suppl) 535-542.
- [223] Town T, Laouar Y, Pittenger C, Mori T, Szekeley CA, Tan J, et al. Blocking TGF-beta-Smad2/3 innate immune signaling mitigates Alzheimer-like pathology. *Nat Med.* 2008;14(6) 681-687.
- [224] Chalmers KA, Love S. Phosphorylated Smad 2/3 colocalizes with phospho-tau inclusions in Pick disease, progressive supranuclear palsy, and corticobasal degeneration but not with alpha-synuclein inclusions in multiple system atrophy or dementia with Lewy bodies. *J Neuropathol. Exp. Neurol.* 2007;66(11) 1019-1026.
- [225] Irvani MM, McCreary AC, Jenner P. Striatal plasticity in Parkinson's disease and L-dopa induced dyskinesia. *Parkinsonism. Relat Disord.* 2012;18 Suppl 1:S123-5. S123-S125.
- [226] Wise RA. Roles for nigrostriatal--not just mesocorticolimbic--dopamine in reward and addiction. *Trends Neurosci.* 2009;32(10) 517-524.

- [227] Lee CS, Samii A, Sossi V, Ruth TJ, Schulzer M, Holden JE, et al. In vivo positron emission tomographic evidence for compensatory changes in presynaptic dopaminergic nerve terminals in Parkinson's disease. *Ann. Neurol.* 2000;47(4) 493-503.
- [228] Sossi V, de LF-F, Holden JE, Schulzer M, Ruth TJ, Stoessl J. Changes of dopamine turnover in the progression of Parkinson's disease as measured by positron emission tomography: their relation to disease-compensatory mechanisms. *J Cereb. Blood Flow Metab.* 2004;24(8) 869-876.
- [229] Braithwaite SP, Stock JB, Mouradian MM. alpha-Synuclein phosphorylation as a therapeutic target in Parkinson's disease. *Rev Neurosci.* 2012;23(2) 191-198.
- [230] Krieglstein K, Richter S, Farkas L, Schuster N, Dunker N, Oppenheim RW, et al. Reduction of endogenous transforming growth factors beta prevents ontogenetic neuron death. *Nat. Neurosci.* 2000;3(11) 1085-1090.
- [231] Tapia-Gonzalez S, Giraldez-Perez RM, Cuartero MI, Casarejos MJ, Mena MA, Wang XF, et al. Dopamine and alpha-synuclein dysfunction in Smad3 null mice. *Mol Neurodegener.* 2011;6:72. 72.
- [232] Hoy SM, Keating GM. Rasagiline: a review of its use in the treatment of idiopathic Parkinson's disease. *Drugs.* 2012;72(5) 643-669.
- [233] Kriks S, Shim JW, Piao J, Ganat YM, Wakeman DR, Xie Z, et al. Dopamine neurons derived from human ES cells efficiently engraft in animal models of Parkinson's disease. *Nature.* 2011;480(7378) 547-551.
- [234] Tsang HT, Edwards TL, Wang X, Connell JW, Davies RJ, Durrington HJ, et al. The hereditary spastic paraplegia proteins NIPA1, spastin and spartin are inhibitors of mammalian BMP signalling. *Hum. Mol Genet.* 2009;18(20) 3805-3821.
- [235] Katsuno M, Adachi H, Minamiyama M, Waza M, Doi H, Kondo N, et al. Disrupted transforming growth factor-beta signaling in spinal and bulbar muscular atrophy. *J Neurosci.* 2010;30(16) 5702-5712.
- [236] Mitchell JD, Borasio GD. Amyotrophic lateral sclerosis. *Lancet.* 2007;369(9578) 2031-2041.
- [237] Strong MJ, Kesavapany S, Pant HC. The pathobiology of amyotrophic lateral sclerosis: a proteinopathy? *J Neuropathol. Exp. Neurol.* 2005;64(8) 649-664.
- [238] Redler RL, Dokholyan NV. The complex molecular biology of amyotrophic lateral sclerosis (ALS). *Prog. Mol Biol Transl. Sci.* 2012;107:215-62. 215-262.
- [239] Nakamura M, Ito H, Wate R, Nakano S, Hirano A, Kusaka H. Phosphorylated Smad2/3 immunoreactivity in sporadic and familial amyotrophic lateral sclerosis and its mouse model. *Acta Neuropathol.* 2008;115(3) 327-334.
- [240] Nakamura M, Kaneko S, Wate R, Asayama S, Nakamura Y, Fujita K, et al. Regionally different immunoreactivity for Smurf2 and pSmad2/3 in TDP-43-positive inclusions of amyotrophic lateral sclerosis. *Neuropathol. Appl. Neurobiol.* 2012; 10-2990.

- [241] Day WA, Koishi K, Nukuda H, McLennan IS. Transforming growth factor-beta 2 causes an acute improvement in the motor performance of transgenic ALS mice. *Neurobiol. Dis.* 2005;19(1-2) 323-330.
- [242] Nakamura M, Kaneko S, Ito H, Jiang S, Fujita K, Wate R, et al. Activation of Transforming Growth Factor-beta/Smad Signaling Reduces Aggregate Formation of Mislocalized TAR DNA-Binding Protein-43. *Neurodegener. Dis.* 2012.

