

Review

The roles of annexins and alkaline phosphatase in mineralization process[✉]

Marcin Balcerzak¹, Eva Hamade², Le Zhang², Slawomir Pikula¹, Gérard Azzar², Jacqueline Radisson², Joanna Bandorowicz-Pikula¹ and Rene Buchet^{2✉}

¹*M. Nencki Institute of Experimental Biology, Polish Academy of Sciences, Warszawa, Poland;*

²*Laboratoire de Physico-Chimie Biologique, UMR CNRS 5013, Université Claude Bernard, Lyon 1, UFR de Chimie-Biochimie, F-69622 Villeurbanne, France*

Received: 05 August, 2003; revised: 24 September, 2003; accepted: 7 October, 2003

Key words: matrix vesicle, annexin, alkaline phosphatase, hydroxyapatite, mineralization

In this review the roles of specific proteins during the first step of mineralization and nucleation are discussed. Mineralization is initiated inside the extracellular organelles–matrix vesicles (MVs). MVs, containing relatively high concentrations of Ca²⁺ and inorganic phosphate (P_i), create an optimal environment to induce the formation of hydroxyapatite (HA). Special attention is given to two families of proteins present in MVs, annexins (AnxAs) and tissue-nonspecific alkaline phosphatases (TNAPs). Both families participate in the formation of HA crystals. AnxAs are Ca²⁺- and lipid-binding proteins, which are involved in Ca²⁺ homeostasis in bone cells and in extracellular MVs. AnxAs form calcium ion channels within the membrane of MVs. Although the mechanisms of ion channel formation by AnxAs are not well under-

[✉]This work was supported in part by grant No. 3 P04A 007 22 from the State Committee for Scientific Research (KBN, Poland) and by CNRS. Authors would like to thank Dr. Pascale Chavassieux from Université Claude Bernard, Lyon 1, France, for fruitful discussion and Dr. John Carew for language corrections.

✉ Corresponding author: Rene Buchet, Laboratoire de Physico-Chimie Biologique, UMR CNRS 5013, Université Claude Bernard, Lyon 1, UFR de Chimie-Biochimie, 6 rue Victor Grignard, F-69622 Villeurbanne, France; tel.: (33) 4 72 43 1320; fax: (33) 4 7243 1543; e-mail: rbuchet@univ-lyon1.fr

Abbreviations: 1,25-(OH)₂-D₃, 1 α ,25-dihydroxyvitamin D₃; AnxA, vertebrate annexin; ATPase, ATP hydrolase; ATRA, all-*trans* retinoic acid; BMP-2, bone morphogenetic protein 2; GPI, glycosylphosphatidylinositol; HA, hydroxyapatite; MV, matrix vesicle; NTP, nucleoside triphosphate; PE, phosphatidylethanolamine; PC-1, plasma cell membrane glycoprotein-1 (NTP pyrophosphatase phosphodiesterase isoenzyme); PS, phosphatidylserine; RAR/RXR, receptors of retinoic acid and its derivatives; T3, 3,5,3'-triiodo-L-thyronine; TGF β , transforming growth factor β ; TNAP, tissue-nonspecific alkaline phosphatase.

stood, evidence is provided that acidic pH or GTP contribute to this process. Furthermore, low molecular mass ligands, as vitamin A derivatives, can modulate the activity of MVs by interacting with AnxAs and affecting their expression. AnxAs and other anionic proteins are also involved in the crystal nucleation. The second family of proteins, TNAPs, is associated with P_i homeostasis, and can hydrolyse a variety of phosphate compounds. ATP is released in the extracellular matrix, where it can be hydrolyzed by TNAPs, ATP hydrolases and nucleoside triphosphate (NTP) pyrophosphohydrolases. However, TNAP is probably not responsible for ATP-dependent Ca^{2+} /phosphate complex formation. It can hydrolyse pyrophosphate (PP_i), a known inhibitor of HA formation and a byproduct of NTP pyrophosphohydrolases. In this respect, antagonistic activities of TNAPs and NTP pyrophosphohydrolases can regulate the mineralization process.

SKELETAL TISSUES

The two major skeletal tissues, cartilage and bones, are structurally and functionally different (Heinegard & Oldberg, 1989). Cartilage is highly hydrated and, except at the growth plates of long bones, rarely mineralizes, resulting in a permeable matrix of gel-like consistency. On the other hand, bone matrix routinely mineralizes to form a rigid impermeable matrix (Marks & Popoff, 1988). Proteoglycan and type II collagen are major matrix components in cartilage, while type I collagen is the major part of bone matrix (Marks & Popoff, 1988). In both cartilage and bone, cellular activities include matrix formation, mineralization and resorption.

In each tissue, different cell types (Fig. 1) perform distinct tasks, which sometimes overlap each other. Bone matrix is formed and mineralized by osteoblasts and resorbed by osteoclasts (Fig. 1A). Osteocytes participate in extracellular exchanges between different components of osseous tissue. Osteocytes are also involved in the mechanotransduction (Marks & Popoff, 1988). In cartilage, matrix formation results from the activity of chondrocytes (Marks & Popoff, 1988). Chondrocytes express hypertrophic and non-hypertrophic phenotypes (Fig. 1B). Hypertrophic chondrocytes are characteristic for developing bones and for so-called growth plate. Non-hypertrophic chondrocytes are also found in the growth plate and may participate in the formation of articular cartilage (Fig. 1C) (Poole, 2001).

Growth plate chondrocytes undergo several series of differentiation events, including proliferation and hypertrophy. All these events are required for bone formation during endochondral ossification. Chondrocyte hypertrophy occurs at the expense of adjacent matrix and it requires matrix resorption.

Bone formation takes place in the organism not only during embryonic development (growth plate cartilage in the process of endochondral bone formation) and growth but throughout the life in the process of physiological bone remodeling (Lian & Stein, 1996).

CELL BIOLOGY

Chondrocytes and osteoblasts are of mesenchymal origin. Mesenchymal stem cells are able to generate progenitors with restricted developmental potential. From progenitor cells, various cell types can be differentiated into fibroblasts, adipocytes, chondrocytes and osteoblasts (Fig. 2). The two latter cell types under the influence of growth factors give rise to cells able to form calcified tissues. Hypertrophic chondrocytes and osteoblasts initiate the calcification process by releasing matrix vesicles (MVs) (Anderson, 2003). MVs of growth plate cartilage differ in lipid and protein composition from MVs produced by osteoblasts (Boyan *et al.*, 1988). It has been suggested that MV biogenesis, from growth plate hypertrophic chondrocytes, could be the result of programmed cell death. This would




Figure 1. Regional organization and relationships among bone, growth plate and articular cartilage.

Panel A. Topographic relationship among bone cells. Osteoblasts are located on the lining layer of bone surface, actively producing matrix, which is not yet calcified (osteoid tissue). Osteocyttes are the most mature or terminally differentiated cells of the osteoblast lineage. Osteocyttes are embedded in the bone matrix. Panel B. primary mammalian growth plate showing progressive development of chondroblasts from the proliferative zone to the lower hypertrophic zone, where matrix synthesis stops and the extracellular matrix is calcified. Panel C. Regional organization of articular cartilage. The superficial zone contains thin collagen fibrils arranged parallel with the articular surface. The partly calcified cartilage of the calcified zone is indicated. Adapted from Marks & Popoff (1988) and Poole (2001).

appear not to be the case for MVs released from viable osteoblasts (Anderson, 2003). MVs initiate mineral formation starting from embryonic ossification to bone formation in adults (Hoshi & Ozawa, 2000).

MATRIX VESICLES

Several stages of mineralization were identified. The mineralization of bone and cartilage

requires the presence of extracellular MVs (Anderson, 1995; 2003), since the first step of mineralization is initiated inside these organelles. MVs (in size between one hundred to several hundred nanometers in diameter) serve as a site for Ca^{2+} and P_i accumulation. MVs create a specific environment where deposition of initial amorphous mineral complexes (nucleation) occurs and where hydroxyapatite (HA) e.g. $\text{Ca}_{10}(\text{PO}_4)(\text{OH})_2$, is produced and forms needle-like crystals on

the inner surface of the MV membrane. The extracellular matrix contains sufficiently high levels of Ca^{2+} and P_i concentrations to sustain the nucleation process and to propagate

2003). Although details of the mechanism are still unknown, assembly of mineral complexes depends probably on electrostatic, structural and stereochemical properties at the inor-




Figure 2. Lineage of osteochondrogenitor cells.

Pluripotent stem cells develop from multipotential mesenchymal stem cells. The pluripotent stem cells are progenitors of all indigenous cells of connective tissues: fibroblasts, adipocytes, osteoblasts and chondrocytes. The influences of several physiological factors like transforming growth factor β ($\text{TGF}\beta$), bone morphogenetic protein 2 (BMP-2), all-*trans* retinoic acid (ATRA), $1\alpha,25$ -dihydroxyvitamin D_3 ($1,25\text{-(OH)}_2\text{-D}_3$), and 3,5,3'-triiodo-L-thyronine (T_3) on the lineage of osteoblasts and chondrocytes are indicated in the figure. Osteoblasts exist in two forms as osteoblasts monolayer on the surface of growing bone tissue which synthesize and secrete organic components of matrix and produce mineralization competent matrix vesicles and osteocytes enclosed within bone matrix. Chondrocytes are characterized by two phenotypes: non-hypertrophic characteristic for articular cartilage, and hypertrophic in growth plate. Adapted from Lian & Stein (1996).

the mineralization (Anderson, 2003). Ion channels and transporters present in MV membrane are responsible for Ca^{2+} and P_i uptake into these organelles. After reaching a certain length, the needle-like HA crystals are released from MVs into the extracellular matrix. The mechanisms by which the HA crystals can break the membrane of MVs are not very well understood. One possible explanation is that the activity of phospholipases could be triggered, once HA crystals are formed, and may affect the MV membrane fluidity (Swain *et al.*, 1992; Schwartz & Boyan, 1988).

The second step of mineralization starts with a release of HA crystals. These crystals serve as a template for the formation of crystalline arrays, leading to a tissue calcification (Anderson,

organic-organic interface. Subtle interactions between negatively charged domains of proteins, anionic phospholipids and mineral complexes are crucial in the propagation of arrays of crystals. All processes taking place in MVs require a dynamic but tightly regulated system to maintain Ca^{2+} homeostasis and P_i delivery. Many actors have been identified to date, among them vertebrate annexin (AnxAs), Ca^{2+} - and membrane-binding proteins, as well as alkaline phosphatase.

ANNEXINS IN MINERALIZING TISSUES

From twelve members of the annexin family of proteins present in mammalian organisms,

three were identified in MVs: annexin A2 (AnxA2), AnxA5 and AnxA6 (Cao *et al.*, 1993; Kirsch *et al.*, 1997a; 1997b; Kirsch & Claassen, 2000). Due to the high Ca^{2+} concentration both inside and outside MVs, and the high content of anionic phospholipids, mainly phosphatidylserine (PS), and cholesterol in MV membrane (Harder *et al.*, 1997; Wuthier, 1975; Ayala-Sanmartin, 2001; Ayala-Sanmartin *et al.*, 2001; De Diego *et al.*, 2002;), AnxAs can be associated with both outer and inner leaflets of MV membrane. AnxAs affect membrane stability in a Ca^{2+} -dependent manner (Goossens *et al.*, 1995). In addition, AnxAs could be involved in the Ca^{2+} transport, as ion channels inserted within the MV membrane.

During the first phase of MV-mediated calcification, mineral complexes appear on the inner surface of MV membrane. The high affinity for Ca^{2+} of PS is quite strong in the inner leaflet of the MV membrane enriched with anionic lipids (Majeska *et al.*, 1979; Taylor *et al.*, 1998). Accordingly, AnxA5 exhibiting Ca^{2+} -dependent PS-binding property was isolated with $\text{PS-Ca}^{2+}\text{-P}_i$ complexes from nucleation core of chicken growth plate MVs (Wu *et al.*, 1993; 1996; 1997a). Smaller amounts of AnxA2 and AnxA6, as well as other proteins, were co-purified with AnxA5 (Wu *et al.*, 1997a; 2002a).

AnxAs associated with the outer surface of MV and bone-derived cell membranes may interact with extracellular matrix molecules. AnxA2 and AnxA6 bind chondroitin sulfate in a Ca^{2+} -dependent manner (Ishitsuka *et al.*, 1998; Ishitsuka, 2000; Takagi *et al.*, 2002). AnxA5 binds types II and X collagens and C-propeptide of type II collagen (Kirsch & Pfäffle, 1992; von der Mark & Mollenhauer, 1997; Kirsch *et al.*, 2000a). The above described interactions may influence MV shape, thereby affecting crystal growth. Indeed, chondrosarcoma cells, expressing low quantities of AnxA5, are not able to bind type II collagen. This suggests that AnxA5 is a key molecule to promote extracellular matrix binding, which is essential for cartilage function (King *et al.*, 1997). In chicken growth plate, types II

and X collagens enhance Ca^{2+} influx into MVs, promoting activity of ion channels formed by AnxAs (Kirsch & Wuthier, 1994; Kirsch *et al.*, 1994; 2000a). However, the presence of collagen is not essential for mineralization, as shown with knockout animals (Jacenko *et al.*, 1993), with reconstituted systems (Kirsch *et al.*, 1997a) and with purified MVs (Hsu & Anderson, 1978; Kirsch *et al.*, 2000a; Wang & Kirsch, 2002). Nevertheless, collagens could influence initialization and progression of mineral formation in MVs. In addition, *ANXA5*^{-/-} mice were normal in respect of development of their skeletons (Brachvogel *et al.*, 2003), probably because other AnxAs could replace AnxA5 function in knockout animals.

AnxAs are specific markers of chondrocyte hypertrophy. Articular cartilage cells, in contrast to growth plate chondrocytes, maintain a stable phenotype. The upper zone of the articular cartilage (Fig. 1C) contains thin collagen fibrils and proteoglycan called aggrecan. In this zone, tensile forces connected with daily life are maximally concentrated. In lower zones, as in the middle and deep zones, the cell density decreases, collagen fibers are thicker and aggrecan content is higher. Calcified zones, where chondrocytes develop an hypertrophic phenotype, provide a link between subchondral bone and joint cartilage (Poole, 2001). Articular cartilage, unlike growth plate, usually does not undergo matrix calcification. However, mineralization frequently occurs during osteoarthritis and aging. In osteoarthritis, progressive damage and loss of articular cartilage matrix (especially in superficial zone) are observed. These events are accompanied by cell death and pathological matrix mineralization, leading to bone remodelling and to subchondral bone mass increase. In addition, an inflammatory process occurs, giving rise to pain and movement disabilities. The amount of AnxAs is scarce in normal articular cartilage, while it significantly increases during progression of osteoarthritis (Mollenhauer *et al.*, 1999;

Kirsch *et al.*, 2000b; Pfander *et al.*, 2001). Therefore, AnxAs could be specific markers of differentiation during osteoarthritis. For example, AnxA8, a protein not previously described in the growth plate, is expressed during inappropriate cell differentiation in osteoarthritis (White *et al.*, 2002). Relatively high annexin expression in articular cartilage chondrocytes is characteristic for hypertrophic cells and cells undergoing apoptosis (Kirsch *et al.*, 2000b; Kouri *et al.*, 2000) with the appearance of MVs or apoptotic bodies, respectively (Derfus *et al.*, 1998; Hashimoto *et al.*, 1998; Mollenhauer *et al.*, 1999; Kirsch *et al.*, 2000b). These events lead to mineralization of joint matrix (Gelse *et al.*, 2003) and expression of hypertrophy protein markers, as type X collagen and alkaline phosphatase (Hoyland *et al.*, 1991; Pullig *et al.*, 2000; Kirsch *et al.*, 2000b). MVs are present in articular cartilage from healthy subjects (Einhorn *et al.*, 1985; Derfus *et al.*, 1996). In osteoarthritis, MVs coexist in extracellular matrix with apoptotic bodies which are the products of chondrocytes at the terminal differentiation stage. There are no phagocytic cells in joint cartilage, therefore, apoptotic bodies remain in the cartilage unless the extracellular matrix becomes degraded.

FACTORS AFFECTING ANNEXIN ION CHANNEL FORMATION

The existence of a Ca^{2+} transport system in MVs is not well established. Possible candidates are AnxAs, since ion channels formed by these proteins *in vitro* have been described in literature (Berendes *et al.*, 1993; Arispe *et al.*, 1996; Kourie & Wood, 2000; Kirilenko *et al.*, 2002). To understand how AnxAs can mediate Ca^{2+} influx into MVs, factors affecting annexin activity in the mineralization process should be identified, as for example lipid composition of MV membrane.

MV membrane is distinct from plasma membrane (Wuthier, 1975). It is enriched in PS,

diphosphatidylglycerol and lysophospholipids due to the difference in the rate of phospholipid degradation (Wuthier *et al.*, 1977; 1978). The anionic phospholipid content in calcified cartilage and bone is significantly higher than in non-calcifying cartilage zones (Wuthier, 1968; Wu *et al.*, 2002a). This may indicate that anionic phospholipids are involved in mineral formation. It is in agreement with the results of many experiments indicating that Ca^{2+} -dependent binding of AnxAs to model membranes is enhanced by the content of anionic phospholipids. Maximal Ca^{2+} influx mediated by AnxAs into liposomes occurs when they are prepared from PS and phosphatidylethanolamine (PE) mixture at 9:1 mole/mole (Kirsch *et al.*, 1997a). In addition, PS clustering may be induced by the high cholesterol content in MV membrane (Wuthier, 1975). AnxA5 interacts in a Ca^{2+} -dependent manner with cardiolipin in isolated mitochondria (Megli *et al.*, 1995; 2000). Since cardiolipin is present also in MV membrane (Wuthier, 1975), these interactions may occur in MVs.

AnxA2, AnxA5 and AnxA6 are abundant in acidified organic extracts of MVs (25–40% of extraction of AnxAs from crude preparations, as reported by Genge *et al.*, 1991), suggesting their presence in the hydrophobic core of lipid bilayer. This was also evidenced by using selective labeling of AnxA5 with photoactivable hydrophobic reagent, revealing that this protein inserts into the membrane hydrophobic core at mildly acidic pH (Isas *et al.*, 2000). At low pH, aspartate and glutamate residues of AnxAs are protonated. The protein surface becomes more hydrophobic, facilitating its insertion into lipid bilayer (Kohler *et al.*, 1997; Beermann *et al.*, 1998; Isas *et al.*, 2000; 2003; Golczak *et al.*, 2001a; b).

Whether, the low pH-induced annexin ion channels in MVs may form during mineralization, remains to be elucidated. In fact, it is not clear which population of AnxAs may participate in ion channel formation: AnxAs associated with the external or the

internal leaflet of the MV membrane. It is possible that annexin channels are formed in plasma membrane before MV budding. The pH measurements made in tissue sections indicate that intracellular pH in chicken growth plate is dependent on the zone from which chondrocytes are derived. The lowest pH was observed in the periphery of late hypertrophic and calcifying cells (Wu *et al.*, 1997b). Moreover, protons are byproducts of HA formation in MVs. Low pH can prevent HA formation by increasing solubility of formed mineral for which the optimal pH for crystal formation is in the range of 7.4–7.8 (Valhmu *et al.*, 1990). However, extensive acidification during crystal formation is prevented by type II carbonic anhydrase (Stechschulte *et al.*, 1992; Sauer *et al.*, 1994).

Chondrocytes in the growth plate release NTPs that may regulate cell maturation and matrix mineralization (Hatori *et al.*, 1995; Hung *et al.*, 1997;). NTPs are also released by non-stimulated (Hatori *et al.*, 1995) and by stimulated osteoblasts in response to mechanical activation (Romanello *et al.*, 2001). AnxAs can bind nucleotides under *in vitro* conditions (Kirilenko *et al.*, 2001; 2002; Bandorowicz-Pikula *et al.*, 2001; 2003) but probably, with the exception of AnxA7 (Caohuy *et al.*, 1996), do not hydrolyze nucleotides. GTP in a millimolar concentration range induced AnxA6 channel formation in planar lipid bilayers (Kirilenko *et al.*, 2002). It was also shown that the AnxA5 ion channel activity in MV could be regulated by NTPs (Arispe *et al.*, 1996). However, the mechanism by which these channels are formed in MVs is not yet elucidated.

INTERACTIONS OF ANNEXINS WITH OTHER PROTEINS DURING MINERALIZATION

Changes in extracellular fluid composition, reductions in extracellular pH, increase in ma-

trix synthesis, as well as morphological changes associated with local compaction of matrix around the cells, may affect chondrocyte proliferation and maturation (Buschmann *et al.*, 1995; Quinn *et al.*, 1998; Wu & Chen, 2000). For example, hyperosmotic stimuli was reported to affect protein synthesis in cartilage, as well as Ca^{2+} and H^+ homeostasis (Dascalu *et al.*, 1996; Erickson *et al.*, 2001).

Additional factors that may affect annexin ion channel activity during mineralization are associated with their interaction with other proteins. Mobasheri *et al.* (2002) attributed perception of mechanical signals in cartilage to cell surface membrane mechanoreceptors. These receptors are composed of integrins and stretch activated ion channels. Multiple mechanosensitive ion channels were characterized in osteoblasts and chondrocytes. None of these channels revealed similarities with AnxAs (Davidson *et al.*, 1990; 1996; Duncan & Hruska, 1994; Guilak *et al.*, 1999; Koprowski & Kubalski, 2001; Biggin & Sansom, 2003; Shakibaei & Mobasheri, 2003). In osteoblasts, increase in $[\text{Ca}^{2+}]_{\text{in}}$ by oscillating fluid flow, was attenuated by the addition of anti-AnxA5 antibodies. This suggests that AnxA5 may be involved in mechanotransduction in bone (Yellowley *et al.*, 2002). Recently, it was observed that AnxA5 binds to the cytoplasmic part of $\beta 5$ subunit of bovine integrin $\alpha \nu \beta 5$ (Andersen *et al.*, 2002).

Homodimeric S100A and S100B proteins interact with AnxA5 and AnxA6 at 1 mole S100 dimer per 2 mole annexin stoichiometry (Donato, 2003). It was previously demonstrated by co-immunoprecipitation (Arcuri *et al.*, 2002) and inhibition of annexin-mediated Ca^{2+} fluxes (Garbuglia *et al.*, 1998; 2000). However, annexin-S100 interactions have not been investigated in cell systems able to perform mineralization. It was reported that calbindin D9k, an unusual monomeric member of S100 proteins, is present in MVs (Balmain, 1991; 1992; Balmain *et al.*, 1989; 1991; 1995). Calbindin

D9k is a vitamin D₃-dependent protein and its expression affects dietary Ca²⁺ accumulation in bones (Li *et al.*, 2001). The presence of this protein is important for interstitial Ca²⁺ absorption. In rat epiphyseal chondrocytes, calbindin D9k is highly expressed only in mature and hypertrophic chondrocytes (Balmain *et al.*, 1995). It is postulated that calbindin D9k takes part in mineral nucleation (Balmain, 1991). Besides, calbindin D9k reveals 47% and 37% identity and 64% and 55% homology in primary structure with S100A and S100B proteins, respectively. Such high similarity between proteins supports the hypothesis that calbindin D9k can interact with AnxAs during mineralization.

EFFECT OF RETINOIC ACID ON THE MATURATION OF CHONDROCYTES AND ON THE MINERALIZATION PROCESS

Recent findings reveal that growth plate chondrocytes proliferate and mature faster upon treatment with all-*trans* retinoic acid (ATRA) (De Luca *et al.*, 2000). It is accompanied by terminal differentiation of chondrocytes and production of mineralization competent MVs, rich in AnxAs and alkaline phosphatase (Wang & Kirsch, 2002; Wang *et al.*, 2003). ATRA, an agonist of receptors of retinoic acid and other retinoids (RAR/RXR), stimulates events leading to mineralization and matrix remodeling. In addition, it stimulates cell differentiation and apoptosis, as well expression of metalloproteinases (Nie *et al.*, 1998), type I collagen (expression of proteoglycans and type II and X collagens is inhibited), alkaline phosphatase and AnxAs (Wu *et al.*, 1997c; Wang *et al.*, 2003). Moreover, events characteristic for apoptosis, such as down-regulation of Bcl-2, activation of capsase-3 and DNA fragmentation occur after treatment with ATRA. These events are reversed by simultaneous treat-

ment of cells with ATRA and BAPTA-AM (intracellular Ca²⁺ chelator) or K-201, a 1,4-benzothiazepine derivative that can inhibit ion channel activity of AnxAs (Kaneko *et al.*, 1997a; 1997b; Hofmann *et al.*, 1998; Wang *et al.*, 2003). This may indicate that annexin-mediated Ca²⁺ fluxes are responsible for events related to cell maturation, cell apoptosis and tissue mineralization. Recently, we observed that precursor of ATRA, all-*trans* retinol (vitamin A), binds to AnxA6 *in vitro* (Fig. 3), especially at acidic pH, providing a possible regulatory link with an annexin-mediated mineralization process. Addition of retinoids could promote the mineralization process not only by enhancing annexin expression but by direct interaction with AnxAs or by changing the membrane fluidity (Wang *et al.*, 2003). It has been also shown that 1 α ,25-dihydroxyvitamin D₃ binds to AnxA2 of rat osteoblast-like ROS 24/1 cells, inducing increases in intracellular Ca²⁺ concentration (Baran *et al.*, 2000).

ROLES OF ANNEXINS AND OTHER ANIONIC PROTEINS IN THE NUCLEATION PROCESS

Most non-collagenous proteins involved in initiation and regulation of biological mineral formation are anionic (Boskey, 1996). Among proteins synthesized by osteoblasts are osteonectin, osteopontin, osteocalcin and bone sialoprotein. Cartilage extracellular proteins are similar to bone, while both tissues differ in types of collagens. All these proteins share a high content of aspartic and glutamic acid residues (30–40%) and multiple phosphoryl and sialyl groups. They differ in their abilities to affect the formation of HA *in vitro* (Hunter *et al.*, 1996). Additionally, the phosphoproteins of bone are processed by limited proteolysis, then they are converted into more phosphorylated species that could facilitate mineralization (Suzuki *et al.*, 1996). AnxAs have several putative phosphorylation sites and




Figure 4. Binding of vitamin A (all-*trans* retinol) to AnxA6.

To determine binding of vitamin A to AnxA6, quenching of intrinsic fluorescence of the annexin was measured, using the same method as for the retinol carrier protein (Raghu *et al.*, 2003). Panel A. Human recombinant AnxA6 (1 μ M) was incubated in 150 mM NaCl, 10 mM Tris/HCl, pH 7.4, without (bold line) or with (dashed line) 3 μ M vitamin A added from concentrated stock solution in ethanol (final concentration of ethanol did not exceed 0.5%). Asolectin liposomes were also added in the presence of AnxA6 (protein/lipid ratio of 1:1000, by mole) and vitamin A (dotted line). Panel B. AnxA6 (1 μ M) was incubated in 150 mM NaCl, 10 mM citric buffer, pH 6.0, without (bold line) or with (dashed line) 3 μ M vitamin A. Asolectin liposomes were also added in the presence of AnxA6 (protein/lipid ratio of 1:1000, by mole) and vitamin A (dotted line). Samples were excited at 295 nm and fluorescence emission spectra were recorded at the wavelength range from 320 to 380 nm. All measurements were performed on a Fluorolog 3 spectrophotometer (Jobin Yvon Spex Edison, NJ) with 2-nm slits for both excitation and emission, at 25°C. Quenching of the AnxA6 intrinsic fluorescence by vitamin A is higher at pH 6.0 than at pH 7.4. After liposome addition, the protein fluorescence returns to the basic level only at pH 7.4, probably due to higher affinity of vitamin A for lipids than for AnxA6 (dissociation of protein-vitamin A complex). At pH 6.0, upon addition of liposomes, AnxA6 inserts within the hydrophobic core of the membrane lipid bilayer where it is still able to interact with hydrophobic vitamin A. The result of this experiment suggests that AnxA6 binds vitamin A *in vitro*.

some of them are phosphorylated *in vitro* (Grima *et al.*, 1994). In the case of AnxA6, phosphorylation mimicking mutation resulted in higher Ca²⁺-binding affinity and conformational changes leading to increased

protein flexibility in comparison with wild AnxA6 (Freye-Minks *et al.*, 2003).

It is not known how AnxAs can influence the nucleation sites at the membrane interface and which charged domains are responsible for electrostatic interactions taking place during nucleation. Crystal structures of AnxAs suggest the importance of flexibility for AnxA6 (Avila-Sakar *et al.*, 2000) and AnxA5 (Oling *et al.*, 2000; 2001) in the annexin-phospholipid interactions. Given these findings, it is tempting to suggest that AnxAs may influence molecular organization during nucleation formation, through changes in molecular flexibility or through protein-protein interactions. Such interactions with other MV proteins may favor accumulation of inorganic material.

ALKALINE PHOSPHATASE AND RELATED PROTEINS IN THE MATRIX VESICLES

Alkaline phosphatase is one of the most frequently used biochemical markers of osteoblast activity (Risteli & Risteli, 1993; Garnero & Delmas, 1996; Nawawi *et al.*, 1996; Magnusson *et al.*, 1999). Four genes encoding human alkaline phosphatase have been cloned (Kam *et al.*, 1985; Millán, 1986; Henthorn *et al.*, 1987; Millán & Manes, 1988) corresponding to three specific alkaline phosphatase genes located in chromosome 2 (germ-cell, placenta and intestinal) and one TNAP gene located in chromosome 1 (Moss, 1992). Alkaline phosphatases from all sources are homodimeric metalloenzymes which catalyze the hydrolysis of almost any phosphomonoester with release of P_i and alcohol (Fernley, 1971).

TNAP exists in three forms derived from bone, liver and kidney and differing in carbohydrate groups. Osseous TNAP localized in plasma membrane and in MVs, is a glycosylphosphatidylinositol (GPI)-anchored protein (Noda *et al.*, 1987; Pizauro *et al.*, 1994). Given

the different solubilization of TNAP from osteoblast plasma membrane, obtained from human primary bone cell culture, it was suggested that changes in TNAP activity result from age-related modifications. These changes could be associated with the post-translational modification of TNAP or with the membrane constituents (Radisson *et al.*, 1996; Bourrat *et al.*, 2000). The role of TNAP in mineral formation was evidenced in the case of hypophosphatasia, an inheritable disorder leading to a defective bone formation and characterized by a deficiency in TNAP (Whyte, 1994). Mice deficient in the gene encoding TNAP mimic a severe form of hypophosphatasia, indicating the importance of TNAP in hydrolyzing phosphate substrates, including PP_i , during mineral formation (Narisawa *et al.*, 1997). In addition, several mutations in TNAP occur around a calcium-binding site of the enzyme, not directly associated with the metal-binding site function for hydrolysis. It is suggested that these mutations result in TNAP misfolding (Mornet *et al.*, 2001).

TNAP appears to be a multifunctional enzyme and several of its properties may be important for the mineralization process (Bellows *et al.*, 1991; Hsu, 1992a; 1992b; Rattner *et al.*, 2000). Although TNAP is a well-known biochemical marker of mineralization, the nature of the substrate hydrolyzed by TNAP is not clearly established. It was proposed a long time ago that TNAP may supply P_i by hydrolyzing phosphate substrates (Robison, 1924). This proposal was further substantiated by the observation that supplementation of culture media with β -glycerophosphate, an exogenous TNAP substrate, induced osteogenesis and HA deposition (Tenenbaum, 1981; Ecaot-Chevrier *et al.*, 1983). Addition of levamisole, a specific inhibitor of TNAP activity, prevented β -glycerophosphate-induced mineralization *in vitro* (Tenenbaum, 1987).

TNAP purified from femur of chicken embryos induces the formation of HA in mineralization medium without P_i but containing

Ca^{2+} and phosphate substrates (AMP, creatine phosphate, glucose phosphate and β -glycerophosphate). Under these conditions, addition of ATP does not promote the formation of HA (Hamade *et al.*, 2003). This finding is consistent with the fact that a specific ATPase, rather than TNAP, is responsible for ATP-dependent mineral formation within MVs isolated from bone and/or cartilage (Hsu & Anderson, 1995; 1996; Hsu *et al.*, 1999). The nature of ATPase involved in the ATP-dependent mineral formation is not known and it was proposed that a Ca^{2+} -ATPase could fulfil this role (Hsu & Anderson, 1996). Therefore, not only TNAP but also other enzymes are involved in the P_i homeostasis (Fig. 4). The local concentration of




Figure 4. Production of pyrophosphate and inorganic phosphate and their antagonistic effects on the mineralization process.

PP_i , inhibitors of HA formation, are produced at least partly by plasma cell membrane glycoprotein-1 (NTP pyrophosphatase phosphodiesterase isoenzyme, PC-1) from the hydrolysis of NTPs. The activity of TNAP may boost the formation of HA, by hydrolyzing PP_i and eliminating its inhibitory effect on HA formation. P_i arises from distinct sources, including the hydrolytic activity of TNAP. Accumulation of P_i and Ca^{2+} can induce the formation of HA. Adapted from Hessle *et al.* (2002).

P_i can be increased by the activities of adenine monophosphodiesterases, ATP hydrolases (ATPases) and NTP pyrophosphohydrolases (Bartling & Chong 1999; Anderson, 2003).

Chondrocytes in the growth plate release ATP (Hung *et al.*, 1997). ATP is also released by non-stimulated bone cells (Hatori *et al.*, 1995) or in response to mechanical stimula-

tion (Romanello *et al.*, 2001). PP_i , which could result from the activity of several types of NTP pyrophosphohydrolases (Ho *et al.*, 2000; Huang *et al.*, 1994; Terkeltaub *et al.*, 1994; Johnson *et al.*, 1999a; 1999b) (Fig. 5), and biphosphonates are known inhibitors of HA formation (Tenenbaum, 1987; Skrtic & Eanes, 1994). Thus, it was suggested that TNAP may hydrolyse pyrophosphate groups (Rezende *et al.*, 1994; Camolezi *et al.*, 2002). Heritable deficiencies of the gene encoding NTP pyrophosphohydrolase could play an important role in the etiology of human ossification of the posterior longitudinal ligament of the spine and pathologic soft-tissue ossification, by decreasing the production of PP_i (Okawa *et al.*, 1998; Johnson *et al.*, 1999a; Nakamura *et al.*, 1999). The antagonistic regulation of PP_i concentration by the activities of TNAP and NTP pyrophosphohydrolase was confirmed by the experiments performed on knockout mice null for both TNAP and plasma cell membrane glycoprotein-1 (PC-1, e.g. NTP pyrophosphatase phosphodiesterase isoenzyme) genes (Hessle *et al.*, 2002). The double knockout mice were essentially normal (Hessle, 2002), while TNAP knockout mice mimicked the metabolic disease – hypophosphatasia (Whyte, 2001). These findings suggest that TNAP, together with other hydrolytic enzymes, participate in the P_i homeostasis. Deficiency in PC-1 may result in cartilage calcification, while lack of TNAP expression may result in hypophosphatasia. Both enzymes could be putative therapeutic targets for the treatment of bone mineralization diseases. It was proposed that inhibitors of PC-1 activity could be used for the treatment of hypophosphatasia (Hessle *et al.*, 2002).

P_i arising from extracellular matrix and from the hydrolytic activities of enzymes located either in MVs or in the plasma membrane of chondrocytes or osteoblast cells, is transported into the MVs to initiate the first stage of the mineralization process. Indeed, sodium-dependent P_i transporter responsible

for the P_i uptake inside MVs has been identified (Montesuit *et al.*, 1991; Anderson, 2003). Recent findings indicate that other P_i transporters, not strictly sodium-dependent, are involved in the P_i uptake inside the MV from chondrocytes (Wu *et al.*, 2002b). The regulatory factors on the function of these transporters have not yet been identified.

CONCLUDING REMARKS

The prerequisite for the initial crystalline HA generation and its deposition, requires the continuous supply of Ca^{2+} and P_i inside the MVs. This is accomplished by the activities of several proteins that are involved in Ca^{2+} and P_i homeostasis, among them AnxAs and TNAPs. Although the functions of AnxAs are not well established, an emerging picture suggests that these proteins form calcium ion channels in MV membrane. At our present stage of knowledge, further investigations are needed to substantiate the mechanism of Ca^{2+} fluxes through annexin channels. In addition, AnxAs bind NTPs, probably regulating NTP supply outside of MVs, and providing a possible link to NTP hydrolysing enzymes including alkaline phosphatases. A further link between calcium homeostasis maintained by AnxAs and P_i supply is provided by the calcium-binding property of alkaline phosphatase, recently reported by Mornet and co-workers (Mornet *et al.*, 2001). One may suggest that AnxAs by affecting calcium homeostasis within MVs, directly or indirectly, finely tune up the structure and function of alkaline phosphatase. This is in accordance with the general idea that most MV proteins are multifunctional by nature (Boskey, 1996). Complex interplays between these proteins are necessary to fulfill the highly ordered and tightly controlled mineralization process. An additional regulation can occur at the protein expression level or at the post-translational modification stage in response to stress or aging.

Summarizing, it becomes more obvious now that the interactions between AnxAs, TNAP and their ligands, as well as their respective localization within MVs, are important factors that may influence the calcification of osseous tissues.

REFERENCES

- Andersen MH, Berglund L, Petersen TE, Rasmussen JT. (2002) Annexin-V binds to the intracellular part of the β_5 integrin receptor subunit. *Biochem Biophys Res Commun.*; **292**: 550–7.
- Anderson HC. (1995) Molecular biology of matrix vesicles. *Clin Orthopaed Relat Res.*; **314**: 266–80.
- Anderson HC. (2003) Matrix vesicles and calcification. *Curr Rheumatol Rep.*; **5**: 222–6.
- Arcuri C, Giambanco I, Bianchi R, Donato R. (2002) Annexin V, annexin VI, S100A1 and S100B in developing and adult avian skeletal muscles. *Neuroscience.*; **109**: 371–88.
- Arispe N, Rojas E, Genge BR, Wu LN, Wuthier RE. (1996) Similarity in calcium channel activity of annexin V and matrix vesicles in planar lipid bilayers. *Biophys J.*; **71**: 1764–75.
- Avila-Sakar AJ, Kretsinger RH, Creutz CE. (2000) Membrane-bound 3D structures reveal the intrinsic flexibility of annexin VI. *J Struct Biol.*; **130**: 54–62.
- Ayala-Sanmartin J. (2001) Cholesterol enhances phospholipid binding and aggregation of annexins by their core domain. *Biochem Biophys Res Commun.*; **283**: 72–9.
- Ayala-Sanmartin J, Henry JP, Pradel LA. (2001) Cholesterol regulates membrane binding and aggregation by annexin 2 at submicromolar Ca^{2+} concentration. *Biochim Biophys Acta.*; **1510**: 18–28.
- Balmain N. (1991) Calbindin-D9k. A vitamin-D-dependent, calcium-binding protein in mineralized tissues. *Clin Orthop.*; **265**: 265–76.
- Balmain N. (1992) Identification of calbindin-D9k in matrix vesicles. *Bone Miner.*; **17**: 197–201.
- Balmain N, Hotton D, Cuisinier-Gleizes P, Mathieu H. (1989) Immunoreactive calbindin-D9K localization in matrix vesicle-initiated calcification in rat epiphyseal cartilage: an immunoelectron microscope study. *J Bone Miner Res.*; **4**: 565–75.
- Balmain N, Hotton D, Cuisinier-Gleizes P, Mathieu H. (1991) Immunoreactive calbindin-D9K in bone matrix vesicle. *Histochemistry.*; **95**: 459–69.
- Balmain N, von Eichel B, Toury R, Belquasmi F, Hauchecorne M, Klaus G, Mehls O, Ritz E. (1995) Calbindin-D28K and -D9K and $1,25(\text{OH})_2$ vitamin D_3 receptor immunolocalization and mineralization induction in long-term primary cultures of rat epiphyseal chondrocytes. *Bone.*; **17**: 37–45.
- Bandorowicz-Pikula J, Buchet R, Pikula S. (2001) Annexins as nucleotide-binding proteins: facts and speculations. *BioEssays.*; **23**: 170–8.
- Bandorowicz-Pikula J, Kirilenko A, van Deursen R, Golczak M, Kühnel M, Lancelin J-M, Pikula S, Buchet R. (2003) A putative consensus sequence for nucleotide-binding site of annexin A6. *Biochemistry.*; **42**: 9137–46.
- Baran DT, Quail JM, Ray R, Leszyk J, Honeyman T. (2000) Annexin II is the membrane receptor that mediates the rapid actions of $1\alpha,25$ -dihydroxyvitamin D_3 . *J Cell Biochem.*; **78**: 34–46.
- Bartling PM, Chong KW. (1999) The involvement of phosphohydrolases in mineralization: studies on enzymatic activities extracted from red deer antler. *Calcif Tissue Int.*; **65**: 232–6.
- Beermann ofm cap BB, Hinz HJ, Hofmann A, Huber R. (1998) Acid induced equilibrium unfolding of annexin V wild type shows two intermediate states. *FEBS Lett.*; **423**: 265–9.
- Bellows CG, Aubin JE, Heersche JNR. (1991) Initiation and progression in mineralization of bone nodules formed in vitro: the role of

- alkaline phosphatase and organic phosphate. *Bone Miner.*; **14**: 27–40.
- Berendes R, Burger A, Voges D, Demange P, Huber R. (1993) Calcium influx through annexin V ion channels into large unilamellar vesicles measured with fura-2. *FEBS Lett.*; **317**: 131–4.
- Biggin PC, Sansom MS. (2003) Mechanosensitive channels: stress relief. *Curr Biol.*; **13**: R183–5.
- Boskey AL. (1996) Matrix proteins and mineralization: an overview. *Connect Tissue Res.*; **35**: 357–63.
- Bourrat C, Radisson J, Chavassieux P, Azzar G, Roux B, Meunier PJ. (2000) Activity increase after extraction of alkaline phosphatase from human osteoblastic membranes by nonionic detergents: influence of age and sex. *Calcif Tissue Int.*; **66**: 22–8.
- Boyan BD, Schwartz Z, Swain LD, Carnes DL Jr, Zislis T. (1988) Differential expression of phenotype by resting zone and growth region costochondral chondrocytes *in vitro*. *Bone.*; **9**: 185–94.
- Brachvogel B, Dikschas J, Moch H, Welzel H, von der Mark K, Hofmann C, Poschl E. (2003) Annexin A5 is not essential for skeletal development. *Mol Cell Biol.*; **23**: 2907–13.
- Buschmann MD, Gluzband YA, Grodzinsky AJ, Hunziker EB. (1995) Mechanical compression modulates matrix biosynthesis in chondrocyte/agarose culture. *J Cell Sci.*; **108**: 1497–508.
- Camolezi FL, Daghasanli KRP, Magalhaes PP, Pizauro JM, Ciancaglini P. (2002) Construction of an alkaline phosphatase-liposome system: a tool for mineralization study. *Int J Biochem Cell Biol.*; **34**: 1091–101.
- Cao X, Genge BR, Wu LNY, Buzzi WR, Showman RM, Wuthier RE. (1993) Characterization, cloning and expression of the 67-kDa annexin from chicken growth plate cartilage matrix vesicles. *Biochem Biophys Res Commun.*; **197**: 556–61.
- Caohuy H, Srivastava M, Pollard HB. (1996) Membrane fusion protein synexin (annexin VII) as a Ca^{2+} /GTP sensor in exocytotic secretion. *Proc Natl Acad Sci U S A.*; **93**: 10797–802.
- Dascalu A, Korenstein R, Oron Y, Nevo Z. (1996) A hyperosmotic stimulus regulates intracellular pH, calcium, and S-100 protein levels in avian chondrocytes. *Biochem Biophys Res Commun.*; **227**: 368–73.
- Davidson RM, Lingenbrink PA, Norton LA. (1996) Continuous mechanical loading alters properties of mechanosensitive channels in G292 osteoblastic cells. *Calcif Tissue Int.*; **59**: 500–4.
- Davidson RM, Tatakis DW, Auerbach AL. (1990) Multiple forms of mechanosensitive ion channels in osteoblast-like cells. *Pflugers Arch.*; **416**: 646–51.
- De Diego I, Schwartz F, Siegfried H, Dauterstedt P, Heeren J, Beisiegel U, Enrich C, Grewal T. (2002) Cholesterol modulates the membrane binding and intracellular distribution of annexin 6. *J Biol Chem.*; **277**: 32187–94.
- De Luca F, Uyeda JA, Mericq V, Mancilla EE, Yanovski JA, Barnes KM, Zile MH, Baron J. (2000) Retinoic acid is a potent regulator of growth plate chondrogenesis. *Endocrinology.*; **141**: 346–53.
- Derfus B, Kranendonk S, Camacho N, Mandel N, Kushnaryov V, Lynch K, Ryan L. (1998) Human osteoarthritic cartilage matrix vesicles generate both calcium pyrophosphate dihydrate and apatite *in vitro*. *Calcif Tissue Int.*; **63**: 258–62.
- Derfus BA, Kurtin SM, Camacho NP, Kurup I, Ryan LM. (1996) Comparison of matrix vesicles derived from normal and osteoarthritic human articular cartilage. *Connect Tissue Res.*; **35**: 337–42.
- Donato R. (2003) Interactions of annexins with S100 proteins. In *Annexins: biological implications and annexin-related pathologies*. (Bandorowicz-Pikula J, ed), pp 100–13. Kluwer Academic/Plenum Publishers, New York, NY & Landes Bioscience, Georgetown, TX.
- Duncan RL, Hruska KA. (1994) Chronic, intermittent loading alters mechanosensitive

- channel characteristics in osteoblast-like cells. *Am J Physiol.*; **267**: F909–16.
- Ecaot-Chevrier B, Glorieux FH, van der Rest M, Perreira G. (1983) Osteoblasts isolated from mouse calvaria initiate matrix mineralization in culture. *J Cell Biol.*; **96**: 639–43.
- Einhorn TA, Gordon SL, Siegel SA, Hummel CF, Avitable MJ, Carty RP. (1985) Matrix vesicle enzymes in human osteoarthritis. *J Orthop Res.*; **3**: 160–9.
- Erickson GR, Alexopoulos LG, Guilak F. (2001) Hyper-osmotic stress induces volume change and calcium transients in chondrocytes by transmembrane, phospholipid and G-protein pathways. *J Biomech.*; **34**: 1527–35.
- Fernley HN. (1971) Mammalian alkaline phosphatase. In *The Enzymes*. 3rd edn. (Boyer PD, ed), pp 417–47. Academic Press, New York, London.
- Freye-Minks C, Kretsinger RH, Creutz CE. (2003) Structural and dynamic changes in human annexin VI induced by a phosphorylation-mimicking mutation, T356D. *Biochemistry.*; **42**: 620–30.
- Garbuglia M, Verzini M, Donato R. (1998) Annexin VI binds S100A1 and S100B and blocks the ability of S100A1 and S100B to inhibit desmin and GFAP assemblies into intermediate filaments. *Cell Calcium.*; **24**: 177–91.
- Garbuglia M, Verzini M, Hofmann A, Huber R, Donato R. (2000) S100A1 and S100B interactions with annexins. *Biochim Biophys Acta.*; **1498**: 192–206.
- Garnero P, Delmas PD. (1996) New developments in biochemical markers for osteoporosis. *Calcif Tissue Int.*; **59** (Suppl 1): S2–9.
- Gelse K, Soder S, Eger W, Diemtar T, Aigner T. (2003) Osteophyte development – molecular characterization of differentiation stages. *Osteoarthritis Cartilage.*; **11**: 141–8.
- Genge BR, Wu LNY, Adkisson IV HD, Wuthier RE. (1991) Matrix vesicle annexins exhibit proteolipid-like properties. Selective partitioning into lipophilic solvents under acidic conditions. *J Biol Chem.*; **266**: 10678–85.
- Golczak M, Kicinska A, Bandorowicz-Pikula J, Buchet R, Szewczyk A, Pikula S. (2001a) Acidic pH-induced folding of annexin VI is a prerequisite for its insertion into lipid bilayers and formation of ion channels by the protein molecules. *FASEB J.*; **15**: 1083–5.
- Golczak M, Kirilenko A, Bandorowicz-Pikula J, Pikula S. (2001b) Conformational states of annexin VI in solution induced by acidic pH. *FEBS Lett.*; **496**: 49–54.
- Goossens EL, Reutelingsperger CP, Jongsma FH, Kraayenhof R, Hermens WT. (1995) Annexin V perturbs or stabilises phospholipid membranes in a calcium-dependent manner. *FEBS Lett.*; **359**: 155–8.
- Grima DT, Kandel RA, Pepinsky B, Cruz TF. (1994) Lipocortin 2 (annexin 2) is a major substrate for constitutive tyrosine kinase activity in chondrocytes. *Biochemistry.*; **33**: 2921–6.
- Guilak F, Zell RA, Erickson GR, Grande DA, Rubin CT, McLeod KJ, Donahue HJ. (1999) Mechanically induced calcium waves in articular chondrocytes are inhibited by gadolinium and amiloride. *J Orthop Res.*; **17**: 421–9.
- Hamade E, Azzar G, Radisson J, Buchet R, Roux B. (2003) Chick embryo anchored alkaline phosphatase and mineralization process *in vitro*. Influence of Ca²⁺ and nature of substrates. *Eur J Biochem.*; **270**: 2082–90.
- Harder T, Kellner R, Parton RG, Gruenberg J. (1997) Specific release of membrane-bound annexin II and cortical cytoskeletal elements by sequestration of membrane cholesterol. *Mol Biol Cell.*; **8**: 533–45.
- Hashimoto S, Ochs RL, Rosen F, Quach J, McCabe G, Solan J, Seegmiller JE, Terkeltaub R, Lotz M. (1998) Chondrocyte-derived apoptotic bodies and calcification of articular cartilage. *Proc Natl Acad Sci U S A.*; **95**: 3094–9.
- Hatori M, Teixeira CC, Debolt K, Pacifici M, Shapiro IM. (1995) Adenine nucleotide metabolism by chondrocytes in vitro: role of

- ATP in chondrocyte maturation and matrix mineralization. *J Cell Physiol.*; **165**: 468–74.
- Heinegard D, Oldberg A. (1989) Structure and biology of cartilage and bone matrix noncollagenous macromolecules. *FASEB J.*; **3**: 2042–51.
- Henthorn PS, Raducha M, Edwards YH, Weiss MJ, Slaughter C, Lafferty MA, Harris H. (1987) Nucleotide and amino acid sequences of human intestinal alkaline phosphatase: close homology to placental alkaline phosphatase. *Proc Natl Acad Sci U S A.*; **84**: 1234–8.
- Hessle H, Johnson KA, Anderson HC, Narisawa S, Sali A, Goding JW, Terkeltaub R, Millan JL. (2002) Tissue-nonspecific alkaline phosphatase and plasma cell membrane glycoprotein-1 are central antagonistic regulators of bone mineralization. *Proc Natl Acad Sci U S A.*; **99**: 9445–9.
- Ho AM, Johnson MD, Kingsley DM. (2000) Role of the mouse ank gene in control of tissue calcification and arthritis. *Science.*; **289**: 265–70.
- Hofmann A, Escherich A, Lewit-Bentley A, Benz J, Raguene-Nicol C, Russo-Marie F, Gerke V, Moroder L, Huber R. (1998) Interactions of benzodiazepine derivatives with annexins. *J Biol Chem.*; **273**: 2885–94.
- Hoshi K, Ozawa H. (2000) Matrix vesicles calcification in bones of adult rats. *Calcif Tissue Int.*; **66**: 430–4.
- Hoyland JA, Thomas JT, Donn R, Marriott A, Ayad S, Boot-Handford RP, Grant ME, Freemont AJ. (1991) Distribution of type X collagen mRNA in normal and osteoarthritic human cartilage. *Bone Miner.*; **15**: 151–63.
- Hsu, HHT. (1992a) Further studies on ATP-mediated Ca deposition by isolated matrix vesicles. *Bone Miner.*; **17**: 279–83.
- Hsu HHT. (1992b) In vitro calcium deposition by rachitic rat matrix vesicles: nucleoside triphosphate supported calcium deposition. *Biochim Biophys Acta.*; **1116**: 227–33.
- Hsu HHT, Anderson HC. (1978) Calcification of isolated matrix vesicles and reconstituted vesicles from fetal bovine cartilage. *Proc Natl Acad Sci U S A.*; **75**: 3805–8.
- Hsu HHT, Anderson HC. (1995) A role for ATPase in the mechanisms of ATP-dependent Ca and phosphate deposition by isolated rachitic matrix vesicles. *Int J Biochem Cell Biol.*; **27**: 1349–56.
- Hsu HHT, Anderson HC. (1996) Evidence of the presence of a specific ATPase responsible for ATP-initiated calcification by matrix vesicles isolated from cartilage and bone. *J Biol Chem.*; **271**: 26383–8.
- Hsu HHT, Camacho NP, Anderson HC. (1999) Further characterization of ATP-initiated calcification by matrix vesicles isolated from rachitic rat cartilage. Membrane perturbation by detergents and deposition of calcium pyrophosphate by rachitic matrix vesicles. *Biochim Biophys Acta.*; **1416**: 320–32.
- Huang R, Rosenbach M, Vaughn R, Provvedini D, Rebbe N, Hickman S, Goding J, Terkeltaub R. (1994) Expression of the murine plasma cell nucleotide pyrophosphohydrolase PC-1 is shared by human liver, bone, and cartilage cells. Regulation of PC-1 expression in osteosarcoma cells by transforming growth factor-beta. *J Clin Invest.*; **94**: 560–7.
- Hung CT, Allen FD, Mansfield KD, Shapiro IM. (1997) Extracellular ATP modulates $[Ca^{2+}]_i$ in retinoic acid-treated embryonic chondrocytes. *Am J Physiol.*; **272**: C1611–7.
- Hunter GK, Hauschka PV, Poole AR, Rosenberg LC, Goldberg HA. (1996) Nucleation and inhibition of hydroxyapatite formation by mineralized tissue proteins. *Biochem J.*; **317**: 59–64.
- Isas JM, Cartailier JP, Sokolov Y, Patel DR, Langen R, Luecke H, Hall JE, Haigler HT. (2000) Annexins V and XII insert into bilayers at mildly acidic pH and form ion channels. *Biochemistry.*; **39**: 3015–22.
- Isas JM, Patel DR, Jao C, Jayasinghe S, Cartailier JP, Haigler HT, Langen R. (2003) Global structural changes in annexin 12: The roles of phospholipid, Ca^{2+} and pH. *J Biol Chem.*; **278**: 30227–34.

- Ishitsuka R. (2000) Interaction of annexins with glycosaminoglycans. *Trends Glycosci Glyc.*; **12**: 192–5.
- Ishitsuka R, Kojima K, Utsumi H, Ogawa H, Matsumoto I. (1998) Glycosaminoglycan binding properties of annexin IV, V, and VI. *J Biol Chem.*; **273**: 9935–41.
- Jacenko O, LuValle PA, Olsen BR. (1993) Spondylometaphyseal dysplasia in mice carrying a dominant negative mutation in a matrix protein specific for cartilage-to-bone transition. *Nature.*; **365**: 56–61.
- Johnson K, Moffa A, Chen Y, Pritzker K, Goding J, Terkeltaub R. (1999a) Matrix vesicle plasma cell membrane glycoprotein-1 regulates mineralization by murine osteoblastic MC3T3 cells. *J Bone Miner Res.*; **14**: 883–92.
- Johnson K, Vaingankar S, Chen Y, Moffa A, Goldring MB, Sano K, Jin-Hua P, Sali A, Goding J, Terkeltaub R. (1999b) Differential mechanisms of inorganic pyrophosphate production by plasma cell membrane glycoprotein-1 and B10 in chondrocytes. *Arthritis Rheum.*; **42**: 1986–97.
- Kam W, Clauser E, Kim YS, Kan YW, Rutter W. (1985) Cloning, sequencing, and chromosomal localization of human term placental alkaline phosphatase cDNA. *Proc Natl Acad Sci U S A.*; **82**: 8715–9.
- Kaneko N, Ago H, Matsuda R, Inagaki E, Miyano M. (1997a) Crystal structure of annexin V with its ligand K-201 as a calcium channel activity inhibitor. *J Mol Biol.*; **274**: 16–20.
- Kaneko N, Matsuda R, Toda M, Shimamoto K. (1997b) Inhibition of annexin V-dependent Ca^{2+} movement in large unilamellar vesicles by K201, a new 1,4-benzothiazepine derivative. *Biochim Biophys Acta.*; **1330**: 1–7.
- King KB, Chubinskaya S, Reid DL, Madsen LH, Mollenhauer J. (1997) Absence of cell-surface annexin V is accompanied by defective collagen matrix binding in the Swarm rat chondrosarcoma. *J Cell Biochem.*; **65**: 131–44.
- Kirilenko A, Golczak M, Pikula S, Bandorowicz-Pikula J. (2001) GTP-binding properties of the membrane-bound form of porcine liver annexin VI. *Acta Biochim Polon.*; **48**: 851–65.
- Kirilenko A, Golczak M, Pikula S, Buchet R, Bandorowicz-Pikula J. (2002) GTP-induced membrane binding and ion channel activity of annexin VI: is annexin VI a GTP biosensor? *Biophys J.*; **82**: 2737–45.
- Kirsch T, Pfaffle M. (1992) Selective binding of anchorin CII (annexin V) to type II and X collagen and to chondrocalcin (C-propeptide of type II collagen). Implications for anchoring function between matrix vesicles and matrix proteins. *FEBS Lett.*; **310**: 143–7.
- Kirsch T, Wuthier RE. (1994) Stimulation of calcification of growth plate cartilage matrix vesicles by binding to type II and X collagens. *J Biol Chem.*; **269**: 11462–9.
- Kirsch T, Ishikawa Y, Mwale F, Wuthier RE. (1994) Roles of the nucleational core complex and collagens (types II and X) in calcification of growth plate cartilage matrix vesicles. *J Biol Chem.*; **169**: 20103–9.
- Kirsch T, Nah HD, Demuth DR, Harrison G, Golub EE, Adams SL, Pacifici M. (1997a) Annexin V-mediated calcium flux across membranes is dependent on the lipid composition: implications for cartilage mineralization. *Biochemistry.*; **36**: 3359–67.
- Kirsch T, Nah HD, Shapiro IM, Pacifici M. (1997b) Regulated production of mineralization-competent matrix vesicles in hypertrophic chondrocytes. *J Cell Biol.*; **137**: 1149–60.
- Kirsch T, Claassen H. (2000) Matrix vesicles mediate mineralization of human thyroid cartilage. *Calcif Tissue Int.*; **66**: 292–97.
- Kirsch T, Harrison G, Golub EE, Nah HD. (2000a) The roles of annexins and types II and X collagen in matrix vesicle-mediated mineralization of growth plate cartilage. *J Biol Chem.*; **275**: 35577–83.
- Kirsch T, Swoboda B, Nah H. (2000b) Activation of annexin II and V expression, terminal differentiation, mineralization and apoptosis in human osteoarthritic cartilage. *Osteoarthritis Cartilage.*; **8**: 294–302.

- Kohler G, Hering U, Zschornig O, Arnold K. (1997) Annexin V interaction with phosphatidylserine-containing vesicles at low and neutral pH. *Biochemistry*; **36**: 8189–94.
- Koprowski P, Kubalski A. (2001) Bacterial ion channels and their eukaryotic homologues. *BioEssays*; **23**: 1148–58.
- Kouri JB, Aguilera JM, Reyes J, Lozoya KA, Gonzalez S. (2000) Apoptotic chondrocytes from osteoarthrotic human articular cartilage and abnormal calcification of subchondral bone. *J Rheumatol*; **27**: 1005–19.
- Kourie JI, Wood HB. (2000) Biophysical and molecular properties of annexin-formed channels. *Prog Biophys Mol Biol*; **73**: 91–134.
- Li YC, Bolt MJ, Cao LP, Sitrin MD. (2001) Effects of vitamin D receptor inactivation on the expression of calbindins and calcium metabolism. *Am J Physiol*; **281**: E558–64.
- Lian JB, Stein GS. (1996) *Osteoblast biology*. Academic Press, San Diego.
- Magnusson P, Larsson L, Magnusson M, Davie MWJ, Sharp CA. (1999) Isoforms of bone alkaline phosphatase: Characterization and origin in human trabecular and cortical bone. *J Bone Miner Res*; **14**: 1926–33.
- Majeska RJ, Holwerda DL, Wuthier RE. (1979) Localization of phosphatidylserine in isolated chick epiphyseal cartilage matrix vesicles with trinitrobenzenesulfonate. *Calcif Tissue Int*; **27**: 41–6.
- Marks SC, Popoff SN. (1988) Bone cell biology: the regulation of development, structure and function in the skeleton. *Am J Anat*; **183**: 1–44.
- Megli FM, Selvaggi M, De Lisi A, Quagliariello E. (1995) EPR study of annexin V-cardiolipin Ca-mediated interaction in phospholipids vesicles and isolated mitochondria. *Biochim Biophys Acta*; **1236**: 273–8.
- Megli FM, Mattiazzi M, Di Tullio T, Quagliariello E. (2000) Annexin V binding perturbs the cardiolipin fluidity gradient in isolated mitochondria. Can it affect mitochondrial function? *Biochemistry*; **39**: 5534–42.
- Millán JL. (1986) Molecular cloning and sequence analysis of human placental alkaline phosphatase. *J Biol Chem*; **261**: 3112–5.
- Millán JL, Manes T. (1988) Seminoma-derived Nagao isozyme is encoded by a germ-cell alkaline phosphatase gene. *Proc Natl Acad Sci U S A*; **85**: 3024–8.
- Mobasheri A, Carter SD, Martin-Vasallo P, Shakibaei M. (2002) Integrins and stretch activated ion channels; putative components of functional cell surface mechanoreceptors in articular chondrocytes. *Cell Biol Int*; **26**: 1–18.
- Mollenhauer J, Mok MT, King KB, Gupta M, Chubinskaya S, Koepf H, Cole AA. (1999) Expression of anchorin CII (cartilage annexin V) in human young, normal adult, and osteoarthritic cartilage. *J Histochem Cytochem*; **47**: 209–20.
- Montesuit C, Caverzasio J, Bonjour JP. (1991) Characterization of a Pi transport system in cartilage matrix vesicles: potential role in the calcification process. *J Biol Chem*; **266**: 17791–7.
- Mornet E, Stura E, Lia-Baldini AS, Stigbrand T, Menez A, Le Du MH. (2001) Structural evidence for a functional role of human tissue-nonspecific alkaline phosphatase in bone mineralization. *J Biol Chem*; **276**: 31171–8.
- Moss DW. (1992) Perspectives in alkaline phosphatase research. *Clin Chem*; **38**: 2486–92.
- Nakamura I, Ikegawa S, Okawa A, Okuda S, Koshizuka Y, Kawaguchi H, Nakamura K, Koyama T, Goto S, Toguchida J, Matsushita M, Ochi T, Takaoka K, Nakamura Y. (1999) Association of the human NPPS gene with ossification of the posterior longitudinal ligament of the spine (OPLL). *Hum Genet*; **104**: 492–7.
- Narisawa S, Fröhlander N, Millan JL. (1997) Inactivation of two mouse alkaline phosphatase genes and establishment of a model of infantile hypophosphatasia. *Dev Dyn*; **208**: 432–46.
- Nawawi H, Samson D, Apperley J, Girgis S. (1996) Biochemical markers in patients with

- multiple myeloma. *Clin Chim Acta.*; **253**: 61–77.
- Nie D, Ishikawa Y, Yoshimori T, Wuthier RE, Wu LNY. (1998) Retinoic acid treatment elevates matrix metalloproteinase-2 protein and mRNA levels in avian growth plate chondrocyte cultures. *J Cell Biochem.*; **68**: 90–9.
- Noda M, Yoon K, Rodan GA, Koppel DE. (1987) High lateral mobility of endogenous and transfected alkaline phosphatase: a phosphatidylinositol-anchored membrane protein. *J Cell Biol.*; **105**: 1671–7.
- Okawa A, Nakamura I, Goto S, Moriya H, Nakamura Y, Ikegawa S. (1998) Mutation in Npps in a mouse model of ossification of the posterior longitudinal ligament of the spine. *Nat Genet.*; **19**: 271–3.
- Oling F, Santos JS, Govorukhina N, Mazeret-Dubut C, Bergsma-Schutter W, Oostergetel G, Keegstra W, Lambert O, Lewit-Bentley A, Brisson A. (2000) Structure of membrane-bound annexin A5 trimers: a hybrid cryo-EM-X-ray crystallography study. *J Mol Biol.*; **304**: 561–73.
- Oling F, Bergsma-Schutter W, Brisson A. (2001) Trimers, dimers of trimers, and trimers of trimers are common building blocks of annexin A5 two-dimensional crystals. *J Struct Biol.*; **133**: 55–63.
- Pfander D, Swoboda B, Kirsch T. (2001) Expression of early and late differentiation markers (proliferating cell nuclear antigen, syndecan-3, annexin VI, and alkaline phosphatase) by human osteoarthritic chondrocytes. *Am J Pathol.*; **159**: 1777–83.
- Pizauro JM, Ciancaglini P, Leone FA. (1994) Osseous plate alkaline phosphatase is anchored by GPI. *Braz J Med Res.*; **27**: 453–6.
- Poole AR. (2001) Cartilage in health and disease. In: *Arthritis and allied conditions. A textbook of rheumatology*. 14th edn. Koopman WJ, ed. pp 226–84. Lippincott Williams & Wilkins.
- Pullig O, Weseloh G, Ronneberger D, Kakonen S, Swoboda B. (2000) Chondrocyte differentiation in human osteoarthritis: expression of osteocalcin in normal and osteoarthritic cartilage and bone. *Calcif Tissue Int.*; **67**: 230–40.
- Radisson J, Angrand M, Chavassieux P, Roux B, Azzar G. (1996) Differential solubilization of osteoblastic alkaline phosphatase from human primary bone cell cultures. *Int J Biochem Cell Biol.*; **28**: 421–30.
- Raghu P, Ravinder P, Sivakumar B. (2003) A new method for purification of human plasma retinol-binding protein and transthyretin. *Biotechnol Appl Biochem.*; **38**: 19–24.
- Rattner A, Sabido O, Le J, Vico L, Massoubre C, Frey J, Chamson A. (2000) Mineralization and alkaline phosphatase in collagen lattices populated by human osteoblasts. *Calcif Tissue Int.*; **66**: 35–42.
- Rezende AA, Pizauro JM, Ciancaglini P, Leone FA. (1994) Phosphodiesterase activity is a novel property of alkaline phosphatase from osseous plate. *Biochem J.*; **15**: 517–22.
- Risteli L, Risteli J. (1993) Biochemical markers of bone metabolism. *Ann Med.*; **25**: 385–93.
- Robison R. (1924) The possible significance of hexosephosphoric esters in ossification. *Biochem J.*; **17**: 286–93.
- Romanello M, Pani B, Bicego M, D'Andrea P. (2001) Mechanically induced ATP release from human osteoblastic cells. *Biochem Biophys Res Commun.*; **289**: 1275–81.
- Quinn TM, Grodzinsky AJ, Buschmann MD, Kim YJ, Hunziker EB. (1998) Mechanical compression alters proteoglycan deposition and matrix deformation around individual cells in cartilage explants. *J Cell Sci.*; **111**: 573–83.
- Sauer GR, Genge BR, Wu LN, Donachy JE. (1994) A facilitative role for carbonic anhydrase activity in matrix vesicle mineralization. *Bone Miner.*; **26**: 69–79.
- Schwartz Z, Boyan B. (1988) The effects of vitamin D metabolites on phospholipase A₂ activity of growth zone and resting zone cartilage cells *in vitro*. *Endocrinology.*; **122**: 2191–8.

- Shakibaei M, Mobasheri A. (2003) Beta1-integrins co-localize with Na, K-ATPase, epithelial sodium channels (ENaC) and voltage activated calcium channels (VACC) in mechanoreceptor complexes of mouse limb-bud chondrocytes. *Histol Histopathol.*; **18**: 343-51.
- Skrtic D, Eanes ED. (1994) Effect of 1-hydroxyethylidene-1,1-biphosphonate on membrane-mediated calcium phosphate formation in model liposomal suspensions. *Bone Miner.*; **26**: 219-29.
- Stechschulte DJ Jr, Morris DC, Silvertown SF, Anderson HC, Vaananen HK. (1992) Presence and specific concentration of carbonic anhydrase II in matrix vesicles. *Bone Miner.*; **17**: 187-91.
- Suzuki Y, Kubota T, Koizumi T, Satoyoshi M, Teranaka T, Kawase T, Ikeda T, Yamaguchi A, Saito S, Mikuni-Takagaki Y. (1996) Extracellular processing of bone and dentin proteins in matrix mineralization. *Connect Tissue Res.*; **35**: 223-9.
- Swain LD, Schwartz Z, Boyan BD. (1992) Regulation of matrix vesicle phospholipid metabolism is cell maturation-dependent. *Bone Miner.*; **17**: 192-6.
- Takagi H, Asano Y, Yamakawa N, Matsumoto I, Kimata K. (2002) Annexin 6 is a putative cell surface receptor for chondroitin sulfate chains. *J Cell Sci.*; **115**: 3309-18.
- Taylor MG, Simkiss K, Simmons J, Wu LN, Wuthier RE. (1998) Structural studies of a phosphatidyl serine-amorphous calcium phosphate complex. *Cell Mol Life Sci.*; **54**: 196-202.
- Tenenbaum HC. (1981) Role of organic phosphate in mineralization of bone *in vitro*. *J Dent Res.*; **60**: 1586-9.
- Tenenbaum HC. (1987) Levamisole and inorganic pyrophosphate inhibit β -glycerophosphate induced mineralization of bone formed *in vitro*. *Bone Miner.*; **3**: 13-26.
- Terkeltaub R, Rosenbach M, Fong F, Goding J. (1994) Causal link between nucleotide pyrophosphohydrolase overactivity and increased intracellular inorganic pyrophosphate generation demonstrated by transfection of cultured fibroblasts and osteoblasts with plasma cell membrane glycoprotein-1. Relevance to calcium pyrophosphate dihydrate deposition disease. *Arthritis Rheum.*; **37**: 934-41.
- Valhmu WB, Wu LN, Wuthier RE. (1990) Effects of Ca/P_i ratio, Ca²⁺ x P_i ion product, and pH of incubation fluid on accumulation of ⁴⁵Ca²⁺ by matrix vesicles *in vitro*. *Bone Miner.*; **8**: 195-209.
- Von der Mark K, Mollenhauer J. (1997) Annexin V interactions with collagen. *Cell Mol Life Sci.*; **53**: 539-45.
- Wang W, Kirsch T. (2002) Retinoic acid stimulates annexin-mediated growth plate chondrocyte mineralization. *J Cell Biol.*; **157**: 1061-9.
- Wang W, Xu J, Kirsch T. (2003) Annexin-mediated Ca²⁺ influx regulates growth plate chondrocyte maturation and apoptosis. *J Biol Chem.*; **278**: 3762-9.
- White AH, Watson RE, Newman B, Freemont AJ, Wallis GA. (2002) Annexin VIII is differentially expressed by chondrocytes in the mammalian growth plate during endochondral ossification and in osteoarthritic cartilage. *J Bone Miner Res.*; **17**: 1851-8.
- Whyte MP. (1994) Hypophosphatasia and the role of alkaline phosphatase in skeletal mineralization. *Endocrinol Rev.*; **15**: 439-61.
- Whyte MP. (2001) Hypophosphatasia. In: *The metabolic and molecular bases of inherited diseases*. Scriver CR, Beaudet AL, Sly WS, Valle D, Childs B, Kinzler KW, Vogelstein B, eds, pp 5313-29. McGraw-Hill, New York.
- Wu LN, Yoshimori T, Genge BR, Sauer GR, Kirsch T, Ishikawa Y, Wuthier RE. (1993) Characterization of the nucleational core complex responsible for mineral induction by growth plate cartilage matrix vesicles. *J Biol Chem.*; **268**: 25084-94.
- Wu LN, Genge BR, Sauer GR, Wuthier RE. (1996) Characterization and reconstitution of the nucleational complex responsible for mineral formation by growth plate cartilage ma-

- trix vesicles. *Connect Tissue Res.*; **35**: 309–15.
- Wu LN, Genge BR, Dunkelberger DG, LeGeros RZ, Concannon B, Wuthier RE. (1997a) Physicochemical characterization of the nucleational core of matrix vesicles. *J Biol Chem.*; **272**: 4404–11.
- Wu LN, Wuthier MG, Genge BR, Wuthier RE. (1997b) *In situ* levels of intracellular Ca^{2+} and pH in avian growth plate cartilage. *Clin Orthop.*; **335**: 310–24.
- Wu LNY, Ishikawa Y, Nie D, Genge BR, Wuthier RE. (1997c) Retinoic acid stimulates matrix calcification and initiates type I collagen synthesis in primary cultures of avian weight-bearing growth plate chondrocytes. *J Cell Biochem.*; **65**: 209–30.
- Wu LNY, Genge BR, Kang MW, Arsenault AL, Wuthier RE. (2002a) Changes in phospholipid extractability and composition accompany mineralization of chicken growth plate cartilage matrix vesicles. *J Biol Chem.*; **277**: 5126–33.
- Wu LNY, Guo Y, Genge BR, Ishikawa Y, Wuthier RE. (2002b) Transport of inorganic phosphate in primary cultures of chondrocytes isolated from the tibial growth plate of normal adolescent chickens. *J Cell Biochem.*; **86**: 475–89.
- Wu QQ, Chen Q. (2000) Mechanoregulation of chondrocyte proliferation, maturation, and hypertrophy: ion-channel dependent transduction of matrix deformation signals. *Exp Cell Res.*; **256**: 383–91.
- Wuthier RE. (1968) Lipids of mineralizing epiphyseal tissues in the bovine fetus. *J Lipid Res.*; **9**: 68–78.
- Wuthier RE. (1975) Lipid composition of isolated epiphyseal cartilage cells, membrane and matrix vesicles. *Biochim Biophys Acta.*; **409**: 128–43.
- Wuthier RE, Majeska RJ, Collins GM. (1977) Biosynthesis of matrix vesicles in epiphyseal cartilage. I. *In vivo* incorporation of ^{32}P orthophosphate into phospholipids of chondrocyte membrane, and matrix vesicle fractions. *Calcif Tissue Res.*; **23**: 135–9.
- Wuthier RE, Wians FH Jr, Giancola MS, Dragic SS. (1978) *In vitro* biosynthesis of phospholipids by chondrocytes and matrix vesicles of epiphyseal cartilage. *Biochemistry.*; **17**: 1431–6.
- Yellowley CE, Haut-Donahue T, Jacobs C, Donahue H. (2002) Annexin V contributes to a mechanotransduction mechanism in osteoblasts. *Biophys J.*; **82**: 1306.