

The SCF^{FBW7} ubiquitin ligase complex as a tumor suppressor in T cell leukemia

Benjamin J. Thompson,^{1,2,3} Silvia Buonamici,^{1,2} Maria Luisa Sulis,⁴ Teresa Palomero,⁴ Tomas Vilimas,⁵ Giuseppe Basso,⁶ Adolfo Ferrando,⁴ and Iannis Aifantis^{1,2}

¹Department of Pathology, ²New York University Cancer Institute, New York University School of Medicine, New York, NY 10016

³Medical Scientist Training Program, Committee on Immunology, University of Chicago, Chicago, IL 60637

⁴Institute for Cancer Genetics, Columbia University, New York, NY 10032

⁵Department of Medicine, Northwestern University, Feinberg School of Medicine, Chicago, IL 60611

⁶Hemato-Oncology Laboratory, Department of Pediatrics, University of Padova, 35122 Padova, Italy

Recent studies have shown that activating mutations of NOTCH1 are responsible for the majority of T cell acute lymphoblastic leukemia (T-ALL) cases. Most of these mutations truncate its C-terminal domain, a region that is important for the NOTCH1 proteasome-mediated degradation. We report that the E3 ligase FBW7 targets NOTCH1 for ubiquitination and degradation. Our studies map in detail the amino acid degron sequence required for NOTCH1–FBW7 interaction. Furthermore, we identify inactivating FBW7 mutations in a large fraction of human T-ALL lines and primary leukemias. These mutations abrogate the binding of FBW7 not only to NOTCH1 but also to the two other characterized targets, c-Myc and cyclin E. The majority of the FBW7 mutations were present during relapse, and they were associated with NOTCH1 HD mutations. Interestingly, most of the T-ALL lines harboring FBW7 mutations were resistant to γ -secretase inhibitor treatment and this resistance appeared to be related to the stabilization of the c-Myc protein. Our data suggest that FBW7 is a novel tumor suppressor in T cell leukemia, and implicate the loss of FBW7 function as a potential mechanism of drug resistance in T-ALL.

CORRESPONDENCE

Iannis Aifantis:
iannis.aifantis@med.nyu.edu

Abbreviations used: AML, acute myeloid leukemia; DN, dominant-negative; GSI, γ -secretase inhibitors; HD, heterodimerization; T-ALL, T cell acute lymphoblastic leukemia.

T cell acute lymphoblastic leukemia (T-ALL) is a disease induced by malignant transformation of T lymphocytes that afflicts mainly children and adolescents (1). Although treatment outcome in T-ALL has improved in recent years, patients with relapsed disease continue to have dismal prognosis, despite the use of protocols involving hematopoietic stem cell transplantation. It is thus very important to identify and study the molecular pathways that control both induction of transformation and treatment resistance in this particular type of leukemia. Recent compelling evidence demonstrated that activating mutations in the *NOTCH1* gene are the trigger for cell transformation in the majority of T-ALL patients.

Notch1 is a transmembrane receptor that controls the differentiation of multiple cell types, including cells of the immune system (2). More

specifically, signaling through the NOTCH1 receptor orchestrates the development of T lymphocytes from uncommitted hematopoietic stem cells (3–5). Although NOTCH1 signaling is required for T cell development (6, 7), aberrant activation of the pathway can lead to disease; >50% of T-ALL cases harbor activating mutations in the *NOTCH1* gene (8). Similar mutations were found in several mouse models of T cell leukemia (9–12). Very recent evidence showed that NOTCH1 pathway activation can induce multiple downstream signaling pathways and genes/targets, including the NF- κ B pathway (13–15) and the transcription factor c-Myc (16–18).

The majority of the T-ALL NOTCH1-activating mutations truncate the C terminus of the protein, called the PEST domain because of the high frequency of P-E-S-T amino acids. As the PEST domain is believed to be essential for proteasome-dependent degradation of NOTCH1, these findings suggest that NOTCH1 protein stability could be an important regulator of intracellular signaling thresholds and that abrogation

B.J. Thompson and S. Buonamici contributed equally to this paper.

The online version of this article contains supplemental material.

of the NOTCH1 degradation machinery could predispose cells for transformation. The exact mechanisms by which the PEST domain confers instability to N1-IC remain unclear. Early genetic studies in *Caenorhabditis elegans* identified a candidate silencer of Notch, the E3 ubiquitin ligase SEL-10 (FBW7/hAGO/hCDC4, which is referred to hereafter as FBW7) (19). FBW7 is a potent tumor suppressor (20), targeting proteins such as cyclin E, c-Myc, and c-Jun for proteasomal degradation (21–23). Several reports have demonstrated that FBW7 can also target nuclear Notch1 and Notch4 for ubiquitination and can suppress Notch signaling (24, 25); however, the detailed biochemical nature and the in vivo physiological relevance in development and cell transformation of this interaction remain unknown. Moreover, Fbw7-deficient embryos die around day 10.5 because of defects in vascular development. Two such lines were simultaneously generated by different groups, and whole embryo lysates contained elevated levels of Notch1 protein in one case, but not the other (26, 27), raising questions about the importance of Fbw7 for the degradation of Notch1 or suggesting tissue-specific interactions between the two proteins.

We demonstrate that FBW7 targets NOTCH1, and perform a detailed mapping of their interaction by identifying a degron in the NOTCH1 PEST domain, which is centered on a conserved threonine, T²⁵¹². Alteration of this degron prevents FBW7-mediated ubiquitination of nuclear NOTCH1. Furthermore, we show that loss of FBW7 occurs frequently in T-ALL, as we identify FBW7 mutations in both T-ALL cell lines and primary tumors that carry NOTCH1-activating mutations. These mutations target specific residues in the FBW7-binding pocket and render the molecule incapable of binding NOTCH1, c-Myc, and cyclin E. FBW7 mutations usually pair with NOTCH1-HD mutations, suggesting a functional cooperation between the two events in the induction of NOTCH pathway activation. Finally, we demonstrate that there is an overrepresentation of FBW7 mutations in relapse patient samples, and that all T-ALL cell lines carrying FBW7 mutations are resistant to γ -secretase inhibitor treatment, suggesting that FBW7 inactivation could serve as a mechanism of treatment resistance, potentially caused by the stabilization of the downstream NOTCH1 effector c-Myc.




Figure 1. Physical and functional association between FBW7 and NOTCH1. (A) Schematic depiction of the NOTCH1 (N1) protein domains, and N1-IC WT (N1-IC^{WT}) and 82 and 162 Δ PEST mutants (N1-IC ^{Δ 82} and N1-IC ^{Δ 162}). Asterisk denotes L1601P mutation. The FBW7 degron is highlighted in gray. LNR, LIN-Notch repeat domain; RAM, RBJ- κ -associated module; TAD, transcriptional activation domain; PEST, P-E-S-T-rich domain; HD, heterodimerization domain. (B and C) Coimmunoprecipitation experiments using anti-Flag beads or anti-HA beads, showing interaction between DN FBW7 and N1-IC^{WT}, but DN FBW7 (D) does not interact with N1-IC ^{Δ 82} and N1-IC ^{Δ 162}. (E) Cotransfection with FBW7 α repressed N1^{L1601P} ($P < 0.02$), but not N1^{L1601P- Δ 82}, in reporter assays using a CSL luciferase reporter. pcDNA3.1 represents the control, "empty" expression vector. Error bars represent the SD of duplicate wells.

RESULTS

Physical and functional interaction between FBW7 and oncogenic NOTCH1 mutants

Fig. 1 A schematically displays the NOTCH1 and N1-IC proteins and maps the potential sites of the FBW7 interactions discussed in our article. To confirm that NOTCH1 and FBW7 physically interact, we performed coimmunoprecipitation studies using a dominant-negative (DN) FBW7 that contains the substrate-recognizing WD40 repeats of FBW7, but lacks the F-box domain that recruits ubiquitination machinery (Fig. 3 A). As shown in Fig. 1 (B and C), we detected both N1-IC and DN FBW7 in the same precipitated immunocomplex. Importantly, this interaction required an intact PEST domain, as two different N1-IC mutants lacking the C-terminal 162 or 82 amino acids (N1-IC Δ 162 and N1-IC Δ 82, respectively) failed to coimmunoprecipitate with DN FBW7 (Fig. 1 D).

We then assessed the functionality of this interaction by monitoring the ability of FBW7 (α isoform) to reduce the activation of a NOTCH1 (RBP-J/CSL)-dependent luciferase reporter. To induce NOTCH1 activation, we generated a full-length NOTCH1 construct that contains a leucine-to-proline substitution in the heterodimerization (HD) domain (N1^{L1601P}).

We and others have previously shown that this mutation found in T-ALL patients (8) can induce NOTCH1 activity (15). As evident from Fig. 1 E, FBW7 α significantly ($P < 0.02$) blocked the activation induced by N1^{L1601P}. Similar results were obtained using N1-IC (unpublished data). Cotransfection of DN FBW7 enhanced NOTCH1 activation (Fig. 1 E) and could antagonize the effect of the full-length FBW7 α (not depicted). On the other hand, an L1601P mutant that also lacked the C-terminal 82 amino acids of the PEST domain (N1^{L1601P- Δ 82}) was unaffected by FBW7 α . These observations confirm that FBW7 can target and suppress the activity of NOTCH1-HD mutants found in T-ALL, and suggest that PEST deletion is an efficient “escape” mechanism that promotes cell transformation.

A T²⁵¹²-centered degron controls the interaction of NOTCH1 with FBW7

Specific F-box protein binding sites, termed degrons, have been characterized for several FBW7 substrates. In each confirmed case, the degron contains a crucial threonine residue that must be phosphorylated to allow recognition by FBW7. This in mind, we chose to analyze four threonine-containing




Figure 2. Mutational analysis of the FBW7 NOTCH1 degron. (A) Coimmunoprecipitation experiment using threonine mutants of N1-IC (N1-IC^{T2133A}, N1-IC^{T2467A}, N1-IC^{T2484A}, and N1-IC^{T2512A}) and DN FBW7. Only N1-IC^{T2512A} was unable to coimmunoprecipitate with DN FBW7. (B) Cell lysates from Bosc23 cells transfected with N1^{L1601P} or N1^{L1601P-T2512A} and FBW7 α were used for luciferase reporter assays. N1^{L1601P-T2512A} was not repressed by FBW7 α . (C) Cell lysates from HeLa cells transfected with full-length N1 mutants were used for luciferase reporter assays. N1^{L1601P- Δ 82} and N1^{L1601P-T2512A} induced a 15- and 5-fold increase in transcriptional activity, respectively, compared with N1^{L1601P}. (D) In vivo ubiquitination assay using Bosc23 cells transfected with N1-IC^{WT}, N1-IC Δ 82, or N1-IC^{T2512A} with or without SCF^{FBW7 α} (SKP1, CUL1, ROC1, and FBW7 α) and Ub-HA. SCF^{FBW7 α} enhanced polyubiquitination of N1-IC^{WT}, but not of N1-IC Δ 82 or N1-IC^{T2512A}. (E) A negative charge is required at position 2516 (T+4). Interaction with DN FBW7 is partially preserved with N1-IC^{E2516D}, but abolished with N1-IC^{E2516A}. (F) Mutation of two CDK8 phosphorylation sites prevents N1-IC interaction with FBW7. A mutant lacking both S²⁵¹⁴ and S²⁵¹⁷ (N1-IC^{S2514A-S2517A}) did not coimmunoprecipitate with DN FBW7. Error bars represent the SD of duplicate wells.

potential degrons in N1-IC that could be targeted by FBW7. One site fell outside of the PEST domain (T²¹³³), but was chosen because it was previously suggested as a potential GSK3-β phosphorylation site and FBW7 degron (18). The other three sites were within the PEST domain and included two potential phosphorylation sites for a proline-directed kinase (T²⁴⁸⁴ and T²⁵¹²) and one nonproline-directed site (T²⁴⁶⁷). We generated four N1-IC constructs, in which one of these threonines was replaced with alanine (N1-IC^{T2133A}, N1-IC^{T2467A}, N1-IC^{T2484A}, and N1-IC^{T2512A}), and tested each one's ability to interact with DN FBW7. Of the four mutants, only N1-IC^{T2512A} failed to coimmunoprecipitate with DN FBW7, suggesting that the N1-IC degron recognized by FBW7 depends on phosphorylation of T²⁵¹² (Fig. 2 A).

Next, we verified the functionality of these results using transcriptional reporter assays. In contrast to N1^{L1601P}, FBW7α failed to suppress the activation of a NOTCH1-dependent luciferase reporter by N1^{L1601P} that also contained a T2512A substitution (N1^{L1601P-T2512A}; Fig. 2 B). Furthermore, DN FBW7 was unable to augment NOTCH1 activation induced by that construct in Bosc23 cells (unpublished data). Thus, mutation of a single residue (T²⁵¹²) could recapitulate the effects of a ΔPEST mutation in this specific assay. Finally, we have previously hypothesized that loss of FBW7–NOTCH1 interactions could be a mechanism for enhancing NOTCH1-dependent transcriptional induction. To address this issue, we compared the activity of each mutant and found

that N1^{L1601P-T2512A} produced a five-fold increase in NOTCH1 activity in HeLa cells compared with N1^{L1601P} (Fig. 2 C). We should note that the difference in N1^{L1601P}-induced transcriptional activation in Fig. 2 (B and C) is caused by the utilization of two different cell lines for the luciferase reporter assays.

To confirm that N1-IC^{T2512A} cannot be ubiquitinated by FBW7, we transfected Bosc23 cells with N1-IC^{WT}, N1-IC^{Δ82}, or N1-IC^{T2512A} in the presence or absence of the SCF^{FBW7α} complex. To preserve ubiquitin conjugates, cells were treated with the proteasome inhibitor MG132 6 h before lysis in a buffer containing *N*-ethyl maleimide. HA-ubiquitin immunoprecipitates were further probed for NOTCH1 expression (Fig. 2 D). Cotransfection of SCF^{FBW7α} increased polyubiquitination of N1-IC^{WT}, but not N1-IC^{Δ82} or N1-IC^{T2512A}. The point mutant retained a higher level of polyubiquitination compared with N1-IC^{Δ82} that was not enhanced by SCF^{FBW7α}, suggesting that an additional E3 ligase can promote ubiquitination of N1-IC. This is consistent with a recent report of an FBW7-independent destabilizing element in the PEST domain (28). These experiments identify a conserved FBW7 degron in the N1-IC PEST domain and demonstrate its ability to alter NOTCH1 signaling.

Molecular analysis of the NOTCH1 FBW7 degron

FBW7 degrons in most other substrates contain a serine residue at the T+4 position. Welcker et al. have demonstrated




Figure 3. FBW7 mutations in T-ALL preclude interaction with NOTCH1, c-Myc, and cyclin E. (A) Schematic depiction of the positions of the FBW7 binding pocket and DN FBW7. (B) Chromatograms showing mutations of FBW7 R⁴⁶⁵ and R⁵⁰⁵ found in CEM and P12-ICHIKAWA T-ALL cell lines. DND41 cells express WT FBW7. Arrows indicate mutated nucleotides. (C) Coimmunoprecipitation experiments demonstrating the inability of DN FBW7^{R465C}, DN FBW7^{R479Q}, and DN FBW7^{R505C} to interact with N1-IC. (D and E) The same DN FBW7 mutants as in C do not interact with c-Myc and cyclin E. Asterisk in E indicates a nonspecific band.

that a negative charge at that position—either from a phosphoserine/threonine or an acidic residue—is required for recognition by FBW7 (29, 30). To determine if this is true for recognition of NOTCH1 by FBW7, we substituted E²⁵¹⁶ with either alanine or aspartate (N1-IC^{E2516A} and N1-IC^{E2516D}, respectively). As shown in Fig. 2 E, N1-IC^{E2516A} was unable to bind to DN FBW7; however, we observed partial binding with N1-IC^{E2516D}. Thus, although glutamate appears to be required at T+4 for optimal binding, substituting a negatively charged aspartate at that position permits at least some interaction with FBW7.

The degron we identified in N1-IC contains serine-proline motifs at T+2 and T+5, either of which could be targeted by a proline-directed kinase. Indeed, both S²⁵¹⁴ and S²⁵¹⁷ have been previously suggested to be CDK8 phosphorylation sites (31). To evaluate their importance for recognition by FBW7, we generated N1-IC constructs in which one or both residues were replaced with alanine (N1-IC^{S2514A}, N1-IC^{S2517A}, and N1-IC^{S2514A-S2517A}). Although mutation of S²⁵¹⁷ did not affect binding, mutation of S²⁵¹⁴ impaired it, and binding was completely abolished in the N1-IC mutant lacking both residues (Fig. 2 F). These studies demonstrate that the FBW7 degron in NOTCH1 requires a conserved charged residue and is potentially targeted by CDK8.

PEST mutations in primary T-ALL disrupt the FBW7 degron

To further test the hypothesis that mutations of the FBW7 degron (spanning from L²⁵¹¹ to P²⁵¹⁹) could be a mechanism that promotes transformation in T-ALL, we have studied the pattern of PEST mutations found in T-ALL lines and primary tumors (8). PEST mutations appear to be of great importance, as they have been found in the majority of Notch1-induced T-ALL samples of both human and mouse origin. These mutations fell into three categories relative to the FBW7 degron. The majority of the mutations introduced translational termination codons starting as early as amino acid 2342 (sample T-ALL 44), and thus deleting almost the entire PEST domain. A second category included mutations that introduced termination codons immediately upstream of the L²⁵¹¹ residue (sample T-ALL 63). A third category of at least nine identified mutants included mutations that specifically targeted the FBW7 degron (Fig. S1, available at <http://www.jem.org/cgi/content/full/jem.20070872/DC1>). For example, the mutant found in the KOPTK1 T-ALL sample replaced the WT LTPSPESP degron with a mutated LTP-SRVP sequence. These findings support our hypothesis and suggest that abrogation of the FBW7-controlled NOTCH1 degradation is an important transforming event in T-ALL.

T-ALL cell lines and primary tumors carry conserved FBW7 mutations

Our previous observations suggested an “indirect” way to inactivate FBW7 by deleting the degron on NOTCH1. As PEST mutations occur in a fraction of T-ALL tumors, we wondered whether selective pressure might exist for these cancers to develop FBW7 mutations that abrogate NOTCH1 recognition.

Although FBW7 mutations underlie a wide variety of solid human cancers, it is unknown whether they occur in T-ALL. To address this, we sequenced the F-box and WD40 repeat regions of *FBXW7* from 13 T-ALL cell lines. We found single-nucleotide substitutions that resulted in point mutations in 4 lines (30.8%, Table S1, available at <http://www.jem.org/cgi/content/full/jem.20070872/DC1>). We also found that one T-ALL line (Be13) does not express any FBW7 message, likely because of a chromosomal deletion near the *FBXW7* locus (32). In addition to the cell lines, we sequenced *FBXW7* from 95 primary T-ALL patient samples; 15 of these samples (16%) carried an FBW7 mutation (Table I). Further analysis revealed that 12% (7/58) of the diagnostic and 22% (8/37) of the relapse samples sequenced harbored FBW7 mutations. For 7 of those relapse samples, we were able to locate and sequence their diagnostic counterparts, and we discovered that 4 (57%) of these patients acquired the mutation only at relapse (Table I), suggesting that such mutations could serve as a mechanism of drug resistance in T-ALL.

T-ALL FBW7 mutants lose their ability to bind NOTCH1, c-Myc, and cyclin E

All of the mutations identified affected one of three conserved arginine residues (R⁴⁶⁵, R⁴⁷⁹, and R⁵⁰⁵) in WD40 repeats III

Table I. FBW7 is mutated in a high percentage of primary human T-ALL samples

Patient number	Diagnostic/relapse sample	<i>FBXW7</i> sequence NM_033632.2	Predicted amino acid change
602-783	Diagnosis	C1542T	R465C
603-932	Diagnosis	C1542T	R465C
1	Diagnosis	NA	NA
	Relapse	C1542T (homozygous)	R465C
4	Diagnosis	WT	WT
	Relapse	C1542T	R465C
17	Diagnosis	C1542T	R465C
	Relapse	C1542T	R465C
16	Diagnosis	G1543A	R465H
	Relapse	G1543A	R465H
602-455	Diagnosis	G1543A	R465H
601-268	Diagnosis	G1543A	R465H
27	Diagnosis	WT	WT
	Relapse	G1543A	R465H
4451	Diagnosis	G1585A	R479Q
602-839	Diagnosis	G1585A	R479Q
603-473	Diagnosis	G1585A	R479Q
	Diagnosis	C1662T	R505C
29	Relapse	C1662T	R505C
	Diagnosis	WT	WT
22	Relapse	C1662T	R505C
	Diagnosis	WT	WT
35	Diagnosis	WT	WT
	Relapse	G1663T	R505L

and IV of FBW7 (Table I and Fig. 3, A and B). These three residues (Fig. 3 A) comprise the binding pocket of FBW7 that permits substrate recognition, and all of them contact the phosphothreonine in the consensus degron (33). Nash et al. have shown that mutations of any of these three arginine residues prevent yeast Cdc4 from associating with cyclin E (34). To determine whether the NOTCH1 degron is recognized by FBW7 in the same manner, we generated DN FBW7 mutants in which R⁴⁶⁵, R⁴⁷⁹, or R⁵⁰⁵ was replaced with cysteine (DN FBW7^{R465C} and DN FBW7^{R505C}) or glutamine (DN FBW7^{R479Q}) and tested their abilities to interact with N1-IC. We chose these substitutions because they recapitulated FBW7 mutations found in our T-ALL samples. All three FBW7 mutants failed to bind (Fig. 3 C) and ubiquitinate (not depicted) N1-IC. They also failed to bind to two other characterized FBW7 substrates, c-Myc and cyclin E (Fig. 3, D and E).

Analysis of the mutational status of the NOTCH1 locus in FBW7 mutants

Interestingly, subsequent analysis of the *NOTCH1* locus demonstrated that none of the T-ALL lines bearing FBW7 mutations (0/5) had any NOTCH1 PEST mutations. Indeed, all of them had either NOTCH1-HD mutations (RPMI 8402, P12-ICHIKAWA, and CEM) or expressed WT NOTCH1 alleles (Be13 and Jurkat E6). Furthermore, all sequenced T-ALL lines carrying NOTCH1 PEST mutations expressed WT FBW7. These observations (Table S1) suggested that a NOTCH1 PEST mutation relieves the mutational “pressure”

on the *FBW7* gene. They also suggested that an FBW7 mutation could be a mechanism of signal amplification for the NOTCH1-HD mutants that are weak transcriptional activators. This scenario was supported by our reporter analysis shown in Fig. 2 C, where introduction of a T²⁵¹² mutation abolishes FBW7 binding and induces a fivefold increase in NOTCH1 transcriptional activation. To further prove this hypothesis, we have sequenced the *NOTCH1* locus in the previously described cohort of primary T-ALL patient samples. As evident from Table II, almost 43% (38/89) of the patient samples harbored NOTCH1 mutations. In agreement with our previous observations, none of the FBW7^{mut} samples (0/15) had a PEST mutation. On the other hand, 60% (9/15) of the FBW7^{mut} samples had NOTCH1-HD mutations. The remaining 40% of the FBW7^{mut} samples had no NOTCH1 mutations. These results suggested that cooperation of FBW7 and NOTCH1 mutations can be found in a substantial fraction of T-ALL, and could be critical for disease progression and treatment outcome.

Restoration of FBW7 function decreases activation of the NOTCH1 pathway

If our hypothesis that FBW7 mutations in T-ALL lines contribute to NOTCH1-induced transformation is correct, then restoration of FBW7 expression should suppress NOTCH1 pathway activation in these cells. We thus used retroviral gene transfer to restore WT FBW7 α expression in mutant T-ALL lines. Our initial observations, using both RT-PCR and immunofluorescence detection, was that Flag-FBW7 α




Figure 4. Restoration of FBW7 expression in CEM cells (FBW7^{MUT}) suppresses NOTCH1 target genes. (A and B) Efficient restoration of FBW7^{WT} in CEM T-ALL cells after transduction with a Flag-FBW7 α -expressing retrovirus (MigR1-FBW7 α). Cells were infected with empty MigR1 vector as a control. (A) RT-PCR amplified Flag-FBW7 α only in cells infected with MigR1-FBW7 α . (B) Immunofluorescence staining of FACS-sorted CEM cells infected with MigR1-FBW7 α using FITC-conjugated anti-Flag antibody shows nuclear localization of the construct. (C) Real-time RT-PCR was used to quantify the expression of the NOTCH1 target genes, *DELTEX1*, and *HES1* in CEM cells infected as in A. Infection with MigR1-FBW7 α suppressed the two genes ($P < 0.0001$ for both). Error bars represent the SD of triplicate wells.

Table II. Notch1 mutational analysis of the primary T-ALL patient samples

Patient #	Notch1	FBW7	Patient #	Notch1	FBW7
600-929	WT	WT	13a	WT	WT
601-039	PEST	WT	14a	WT	WT
601-068	PEST	WT	15a	TAD	WT
601-268	HD exon 26	R465H	16a	HD, exon 26	R465H
601-301	PEST	WT	17a	HD, exon 26	R465C
601-464	WT	WT	18a	WT	WT
601-705	WT	WT	19a	WT	WT
601-756	HD exon 26, TAD	WT	20a	TAD	WT
1601-967	WT	WT	21a	WT	WT
1601-991	WT	WT	22a	HD, exon 26	R505C
602-214	WT	WT	26a	HD, exon 26	WT
602-317	PEST	WT	27a	WT	R465H
602-455	WT	R465H	28a	PEST	WT
602-465	HD exon 26	WT	29a	HD, exon 26	R505C
602-774	TAD	WT	30a	WT	WT
602-787	HD exon 26	R465C	31a	HD, exon 26	WT
602-818	HD exon 26, PEST	WT	33a	WT	WT
602-839	HD exon 27	R479Q	34	PEST	WT
603-007	HD exon 26, PEST	WT	34a	HD, exon 26, PEST	WT
603-118	WT	WT	35	WT	WT
603-194	HD exon 26	WT	35a	HD, exon 26	R505L
603-473	WT	R479Q	37a	WT	WT
603-824	WT	WT	38a	WT	WT
603-825	WT	WT	39a	WT	WT
603-885	PEST	WT	40a	WT	WT
603-888	WT	WT	41a	WT	WT
603-895	WT	WT	42a	WT	WT
603-932	WT	R465C	600-361	WT	WT
603-082	PEST	WT	600-619	WT	WT
604-099	WT	WT	600-745	WT	WT
604-198	WT	WT	600-971	HD, exon 27	WT
604-436	WT	WT	600-484	PEST	WT
604-580	WT	WT	602-819	PEST	WT
604-618	PEST	WT	603-829	WT	WT
1a	HD, exon 26	R465C	600-903	WT	WT
2a	WT	WT	2629	WT	WT
3a	PEST	WT	8192	WT	WT
4a	WT	R465C	8024	HD, exon 26	WT
5a	WT	WT	5833	WT	WT
6a	PEST	WT	5182	WT	WT
7a	WT	WT	747	PEST	WT
8a	WT	WT	4451	WT	R479Q
9a	PEST	WT	16a	HD, exon 26	R465H
10a	WT	WT	17a	HD, exon 26	R465C
11a	WT	WT	18a	WT	WT
12a	WT	WT	19a	WT	WT

PEST, HD, and TAD denote the NOTCH1 domains where the mutations are located.

was efficiently expressed and had a characteristic nuclear localization (Fig. 4, A and B). To directly connect targeting of the NOTCH1 pathway by FBW7, we quantified the expression of two well-characterized direct transcriptional targets of NOTCH1, *HES1* and *DELTEX1* (17). As shown by real-time RT-PCR experiments (Fig. 4 C), expression of both genes was significantly ($P < 0.001$) decreased upon FBW7 expression. These observations demonstrate that restoration of FBW7 expression specifically targets the NOTCH1 pathway in T-ALL cells.

c-Myc stabilization in FBW7^{MUT} as a mechanism of resistance to γ -secretase inhibition

In line with the hypothesis that FBW7 mutations could be a novel mechanism of treatment resistance, we have seen that all T-ALL lines (5/5) bearing FBW7 mutations were resistant to treatment with γ -secretase inhibitors (GSIs; unpublished data) (8). The most attractive mechanism of resistance was the stabilization of c-Myc, as c-Myc was previously shown to be a direct target of NOTCH1 activation (16–18). Thus, we decided to study the regulation of c-Myc in T-ALL lines that bear FBW7 mutations. Initially, to demonstrate that c-Myc is regulated by FBW7 in T-ALL, we have shown that the mutated FBW7 proteins fail to interact with c-Myc (Fig. 3, D and E). Moreover, treatment of WT FBW7 (FBW7^{WT}) T-ALL lines with the proteasome inhibitor MG132 for 6 h resulted in the accumulation of c-Myc caused by the inhibition of SCF^{FBW7}-mediated proteolysis. On the other hand, similar treatment of mutant FBW7 (FBW7^{MUT}) T-ALL lines showed no c-Myc protein accumulation (Fig. 5 A). To directly address the mechanism of γ -secretase inhibitor resistance in the FBW7^{MUT} T-ALL lines, we treated KOPTK (FBW7^{WT}), Be13 (FBW7^{MUT}), P12-ICHICAWA (FBW7^{MUT}), and CEM (FBW7^{MUT}) cells with γ -secretase inhibitor (Compound E) for 1–4 d. As shown in Fig. 5 B, N1-IC protein expression was drastically and rapidly down-regulated in all four lines because the inhibition of γ -secretase-mediated cleavage. c-Myc protein declined after 1 d of Compound E treatment in KOPTK (FBW7^{WT}) cells (note the \sim 2-fold increase in protein loading in days 1–4 compared with day 0). In contrast, c-Myc protein levels remained stable in the 3 FBW7^{MUT} lines during the 4 d of treatment. Together, these studies suggest that blocking NOTCH1 signaling alone cannot down-regulate c-Myc protein expression in the context of a coexisting FBW7 mutation. As a result, FBW7-inactivating mutations could provide an “escape” mechanism for GSI-treated T-ALLs caused by the stabilization of c-Myc.

DISCUSSION

The discovery that hyperactivation of the NOTCH1 pathway contributes to the majority of T-ALL cases revolutionized both scientific and clinical perspectives in this field, and has spurred a search for NOTCH1-silencing therapeutic agents. At the core of this search is the need for a more complete understanding of NOTCH1 protein stability regulation, as the majority of the mutations affect proteasome-mediated




Figure 5. FBW7 mutations as a mechanism of resistance to γ -secretase inhibition. (A) T-ALL cell lines were treated with MG132 (20 μ M for 6 h) and analyzed for expression of c-Myc and actin. Treatment with MG132 resulted in accumulation of c-Myc in KOPTK cells, which express WT FBW7, but not in Be13 and P12-ICHICAWA cells, which express no and mutant FBW7, respectively. (B) Expression of N1-IC and c-Myc after treatment with Compound E (CompE) for 1–4 d. c-Myc expression declines only in KOPTK1 cells, which are sensitive to this treatment, but not in Be13 and P12-ICHICAWA cells, which are resistant. N1-IC expression decreased in all of the cell lines, reflecting inhibition of γ -secretase-mediated cleavage of NOTCH1. Actin expression was used as a loading control for both A and B.

degradation. We demonstrate that FBW7 targets NOTCH1 for ubiquitination and map in detail the amino acid degron sequence required for this interaction. Furthermore, we identify inactivating FBW7 mutations in \sim 40% of human T-ALL lines and 16% of primary leukemias. These mutations abolish the ability of FBW7 to interact with its targets, including NOTCH1, c-Myc, and cyclin E.

Initially, using a series of amino acid point mutants, we have mapped the N1-IC–FBW7 interaction to a degron centered around T²⁵¹². We further analyzed the degron, demonstrating that a T+4 acidic glutamate is necessary for interaction with FBW7. The reason for this requirement is not clear. Possibly, a negative charge at position 2516 either facilitates the phosphorylation of T²⁵¹² or directly promotes interaction with FBW7. Current evidence suggests that phosphorylation of N1-IC by CDK8 plays a key role in recognition by FBW7. Work by Fryer et al. showed that MAML1 recruits CDK8 to the N1-IC transcriptional activation complex, and this kinase phosphorylates N1-IC to promote its degradation (31). Two of the three CDK8 phosphorylation sites reported in that study are within the T²⁵¹² degron we identified. Although mutation of either S²⁵¹⁴ or S²⁵¹⁷ by itself allowed at least some interaction with FBW7 (albeit compromised in the case of S²⁵¹⁴), a N1-IC with both residues mutated lost all ability to bind FBW7. Together with the work of Fryer et al., our observations suggest that CDK8 must phosphorylate N1-IC (minimally at S²⁵¹⁴) for it to be efficiently recognized by FBW7, either by priming the site for a different kinase that phosphorylates T²⁵¹², or by directly facilitating physical interaction with FBW7. We do not know the identity of the kinase that phosphorylates T²⁵¹²; CDK8 might be responsible for this, too, but that possibility has not been directly tested. Although the N1-IC degron identified here is both similar to and different from degrons in other FBW7 substrates,

it appears to occupy the canonical FBW7-binding pocket. Crystal structures of Cdc4 bound to a cyclin E peptide show that the phospho-T in the degron makes physical contact with three arginine residues, R⁴⁶⁵, R⁴⁷⁹, and R⁵⁰⁵, in WD40 repeats III and IV (numbering based on FBW7 α isoform) (33). We have found that the FBW7 point mutations identified in T-ALL cell lines and primary leukemia affect these arginine residues, which is consistent with findings in solid human tumors (35, 36). It has been suggested that Fbw7 is a haploinsufficient tumor suppressor in p53^{+/-} mice (37), which supports the idea that decreased gene dosage stabilizes FBW7 targets *in vivo*. However, it is also possible that these mutations act as DN because their inability to interact with the NOTCH1 and c-Myc degrons might create a pool of nonfunctional SCF^{FBW7} complexes.

Although previously detected in certain solid tumors, FBW7-inactivating mutations do not appear to be a general oncogenic trigger in leukemia. Indeed, a recent sequencing of a large number of acute myeloid leukemia (AML) patient samples failed to detect any such mutations (38). Thus, it was a surprise to detect FBW7-inactivating mutations in both T-ALL lines and primary T-ALL samples. These findings suggested that FBW7 mutations could be frequent in T-ALL because of the ability of FBW7 to recognize, bind to, and degrade important oncogenes, including N1-IC and its transcriptional target c-Myc. Interestingly, it seems that FBW7 deficiency could either activate the NOTCH pathway in the absence of NOTCH1 mutations or amplify its signaling strength by cooperating with already existing NOTCH1-HD mutations. If loss of FBW7 is, indeed, a NOTCH1 signal amplifier that facilitates NOTCH1-induced transformation, one would predict that selective pressure to acquire an FBW7 mutation preferentially affects lymphocytes in which the NOTCH1 PEST domain remains intact, as no increase in NOTCH1 signaling capacity could be gained by eliminating FBW7 function in the context of an existing PEST deletion. In agreement with this hypothesis, none of the 13 T-ALL cell lines or 96 primary samples showed any coexisting FBW7 and NOTCH1 PEST mutations. On the other hand, 60% of FBW7^{mut} samples had mutations in the NOTCH1-HD domain. These results suggest that, in T-ALL, the selective pressure to increase NOTCH1 stability and signaling strength is a primary force that drives oncogenesis. However, we do not want to imply that FBW7-mediated transformation depends solely on NOTCH1 activation. Both c-Myc and cyclin E could also be the mediators of this process. However, c-Myc is also a direct target of NOTCH signaling, so it is possible that FBW7 could impinge on both signaling pathways. The specific contribution and the importance of each FBW7 target in the induction of T-ALL remains to be analyzed using *in vivo* genetic approaches.

Moreover, we found that FBW7 mutations were more frequent in relapse samples and that more than half of the patients with FBW7 mutations at relapse did not carry the mutation at diagnosis, suggesting that FBW7 mutation might be an event that confers resistance to treatment. Interestingly,

all T-ALL lines carrying FBW7 mutations were resistant to treatment with GSI. FBW7 is a very attractive treatment resistance candidate because of its ubiquitin ligase function and its ability to target c-Myc protein stability. c-Myc is a direct transcriptional target of NOTCH1, and overexpression of c-Myc can rescue most T-ALL lines treated with GSI (16–18, 39). As c-Myc is a posttranslational target of FBW7 (40), but—at the same time—a transcriptional target of NOTCH activity (16–18), the transforming ability of FBW7 inactivation could be twofold; by losing FBW7 function, a cell should not only stabilize c-Myc but also produce more of it because of increased NOTCH1 signaling. In agreement with this hypothesis, we have shown that T-ALL FBW7 mutants lose their ability to recognize c-Myc. Also, our experiments demonstrate that c-Myc protein remains stable upon GSI treatment of lines carrying FBW7 inhibitors. Although these experiments strongly argue for a role for FBW7, further experimentation, preferably *in vivo*, is required to prove the role of FBW7 deficiency in GSI resistance. Moreover, it is possible that there are multiple resistance mechanisms, as we have found that a T-ALL line (PF382) was FBW7^{WT}, yet resistant to GSI treatment.

We report the identification of FBW7 mutations in human T-ALL and their direct connection to the NOTCH1–c-Myc oncogenic pathway caused by the ability of FBW7 to recognize, bind, and ubiquitinate these target proteins. These mutations target conserved arginines that form the binding pocket of FBW7 and abolish its binding to its targets. We believe that the identification of FBW7 mutations is of unique importance for the design of future molecular therapies for T-ALL. Indeed, our data suggest that these treatments should be multifaceted, covering both NOTCH1 activation and potential escape mechanisms, such as FBW7 mutations, and their effects on the stability of downstream target proteins.

MATERIAL AND METHODS

FBW7 and NOTCH1 mutational analysis. Genomic DNA extracted from cryopreserved lymphoblast samples provided by collaborating institutions in the United States (St. Jude Children's Research Hospital, Memphis, TN) and Canada (Hospital for Sick Children, Toronto, Canada). Analysis of FBW7 mutations in DNA samples from paired diagnostic and relapse T-ALLs was performed in samples from patients enrolled in Associazione Italiana Ematologia Oncologia Pediatrica–Berlin–Frankfurt–Munster Study Group protocols (41).

Recombinant DNA constructs and retrovirus production. Expression vectors carrying Flag-tagged NOTCH1, NOTCH1^{L1601P- Δ 82}, and NOTCH1^{L1601P-T2512A} mutants were created by multistep insertion of PCR-amplified human NOTCH1 sequences into pcDNA3.1 (Invitrogen). The Δ PEST mutants (Δ 82 and Δ 162) were generated by PCR and cloned into the pCS2-N1-IC HA. The Flag-tagged FBW7 α and DN FBW7 expression plasmids were a gift from B. Clurman (Fred Hutchinson Cancer Research Center, Seattle, WA); the cyclin E, Ub-HA, and SCF (CUL1, ROC1, and SKP1) mammalian expression vectors were gifts from M. Pagano (New York University, New York, NY). The Flag-tagged DN FBW7 and N1-IC HA point mutants were generated by site-directed mutagenesis following the manufacturer's protocol (Stratagene). Integrity of all constructs was verified by sequencing and protein expression was confirmed by Western blotting.

Retroviral supernatants were generated as previously described (42), concentrated by 2.5 h centrifugation at 16,500 rpm, and resuspended in ~200 μ l of serum-free medium.

Western blot and coimmunoprecipitation. Bosc23 cells were plated in 6-cm plates and transfected with expression plasmids using FuGENE 6 Transfection Reagent, according to the manufacturer's protocols (Roche). Cells were lysed using TENT buffer (50 mM Tris pH 8, 150 mM NaCl, 2 mM EDTA, 1% Triton X-100, 1 mM PMSF, 1 mM NaF, 1 mM Na_2VO_4 , and 0.2% protease inhibitor cocktail [Sigma-Aldrich]). Total cell extracts were incubated with HA or Flag beads and rocked for 1 h at 4°C. Bead-bound proteins were collected by centrifugation, and after being washed three times they were loaded in 8% SDS-based gels and analyzed by Western blotting using Flag M2 (Sigma-Aldrich), HA probe (Santa Cruz Biotechnology), and Notch1 (C20; Santa Cruz Biotechnology) antibodies. The T-ALL lines cultured at 37°C under 5% CO_2 in RPMI 1640 supplemented with 10% FBS, penicillin, and streptomycin were lysed using TENT buffer after incubation with 1 μ M Compound E or 20 μ M MG132, as specified in the text. Total cell lysates were loaded and run in 12% SDS gels and analyzed by Western blot using antibody against Notch1 Val1744 (Cell Signaling Technology), c-Myc (N-262; Santa Cruz Biotechnology), cyclin E (HE12; Santa Cruz Biotechnology), and actin (CHEMICON International, Inc.).

RT-PCR, sequencing, and real-time RT-PCR analysis. Total cellular RNA treated with DNaseI was extracted as described in the QIAGEN handbook (QIAGEN). cDNA was prepared according to the First Strand cDNA Synthesis kit protocol (Invitrogen). RT-PCR was performed in 25- μ l reactions containing 1.5 μ l cDNA, 1 μ l *Taq* Red DNA Polymerase (Sigma-Aldrich), and 200 nM forward and reverse primers. For detection of Flag-FBW7 α , the following primers were used: *Flag* forward 5'-TACAAGGATGACGATGACAAGC-3' and *FBW7 α* reverse 5'-CACTCTCCTGGTCCATCTCC-3'. For sequencing of FBW7 in human T-ALL lines, ~1.2 kb of cDNA that encodes the FBW7 F-box and WD40 repeats domains was amplified using the FastStart High Fidelity PCR kit (Roche) according to manufacturer's instructions, using the following primers: *FB2* forward 5'-AACAACTTTTGGGGACCTC-3' and *WD1* reverse 5'-ACACAACCTCCCCACTCCC-3'. PCR products were run on 1% Agarose gels, and the appropriate amplicon was purified and sent for direct bidirectional sequencing with the same primers. Sequencing of FBW7 from primary human samples was done by amplifying exons 9 and 10 from genomic DNA with the following primers: *exon 9* forward 5'-TGATGGGATCATTTTATACGGATG-3', *exon 9* reverse 5'-GACAAAACGCTATGGCTTTCC-3', *exon 10* forward 5'-CCC-AACTTCCCATTCCCTTA-3', and *exon 10* reverse 5'-TTTCTTCATGCCAATTTTAACG-3'. The amplicon was sequenced with the following primers: *exon 9seq* forward 5'-TTTAAATCACTTTTCTTTCTACCC-3' and *exon 10seq* forward 5'-TGACTAAATCTACCATGTTTTCTCA. Quantitative RT-PCR was performed in a 50- μ l reaction containing 2 μ l of cDNA, 12.5 μ l of the SYBR Green SuperMix (Bio-Rad Laboratories), 200 nM forward and reverse primers. Real-time RT-PCR amplification was performed on the iCycler iQ detection system, and the data were collected and analyzed using iCycler iQ version 3.1 software (Bio-Rad Laboratories). The primers used were as follows: *DELTEX1* forward 5'-TGGTCACAGCATCAGGCTAC-3' and *DELTEX1* reverse 5'-TGGTCTGGGTATCAGGGAAG-3'; *HES1* forward 5'-TGAGCCAGCTGAAAACACTG-3' and *HES1* reverse 5'-CATTGATCTGGGTCATGCAG-3'; and *ACTIN* forward 5'-TCGTGCGTGACATTAAGGAG-3' and *ACTIN* reverse 5'-AGCACTGTGTTGGCGTACAG-3'.

Immunofluorescence analysis. The CEM T-ALL line cells infected with FBW7 α were washed three times with PBS and fixed in 4% paraformaldehyde in PBS for 15 min. The cell membrane was permeabilized by treatment with 0.5% Triton X-100 for 12 min. The cells were washed with PBS, and nonspecific antibody binding was blocked with 5% bovine serum albumin (Sigma-Aldrich). The cells were treated with anti-Flag-FITC antibody (Sigma-Aldrich) diluted 1:100 for 1 h. After washing with PBS, the cells were stained

with 4,6-diamidino-2-phenylindole, and treated with a prolonged antifading medium (Invitrogen). Images were acquired on a confocal station (Leica) and analyzed using ImageJ (National Institutes of Health) software.

Luciferase assays. Bosc23 or HeLa cells cultured at 37°C under 5% CO_2 in IMDM supplemented with 10% FBS, penicillin, and streptomycin were seeded on 24-well plates at 8×10^4 cells per well and transiently transfected with expression constructs using FuGENE 6 Transfection Reagent (Roche), according to the manufacturer's protocols. Equivalent DNA concentrations were maintained by adding the appropriate amounts of pCDNA3.1 vector. Whole-cell lysates were prepared 48 h after transfection, and luciferase activity was determined using a Dual-Luciferase Reporter Assay System (Promega).

Online supplemental material. Table S1 shows that FBW7 mutations are only present in human T-ALL cell lines with HD mutations or WT NOTCH1. Fig. S1 shows a comparative analysis of NOTCH1 PEST domain mutations found in human T-ALL lines and patients. The online version of this article is available at <http://www.jem.org/cgi/content/full/jem.20070872/DC1>.

We would like to thank Michele Pagano, Piers Nash, William Carroll, Elizabeth Raetz, and Jane Skok for discussions and technical guidance. We also thank Bruce Clurman and Tasuku Honjo for sharing materials, and Mark Minden and A. Thomas Look for valuable clinical samples and communicating unpublished results.

This work was supported by the Fondazione Città Della Speranza (G. Basso), the WOLF Foundation (A. Ferrando), and National Institutes of Health (NIH) grant CA120196 (A. Ferrando). I. Aifantis is supported by the Cancer Research Institute, the Sidney Kimmel Foundation for Cancer Research, the G&P Foundation for Cancer Research, the Penelope London Fund, the Friedman Fund for Childhood Leukemia, and NIH grant R01CA105129. S. Buonamici is supported by the New York University Molecular Oncology and Immunology Training grant. A. Ferrando is a Leukemia and Lymphoma Society Scholar.

The authors declare that they have no competing interests.

Submitted: 1 May 2007

Accepted: 28 June 2007

REFERENCES

- Grabher, C., H. von Boehmer, and A.T. Look. 2006. Notch 1 activation in the molecular pathogenesis of T-cell acute lymphoblastic leukaemia. *Nat. Rev. Cancer*. 6:1–13.
- Artavanis-Tsakonas, S., M.D. Rand, and R.J. Lake. 1999. Notch signaling: cell fate control and signal integration in development. *Science*. 284:770–776.
- Radtke, F., F. Schweisguth, and W. Pear. 2005. The Notch 'gospel'. *EMBO Rep.* 6:1120–1125.
- Maillard, I., T. Fang, and W.S. Pear. 2005. Regulation of lymphoid development, differentiation, and function by the notch pathway. *Annu. Rev. Immunol.* 23:945–974.
- Rothenberg, E.V., and T. Taghon. 2005. Molecular genetics of T cell development. *Annu. Rev. Immunol.* 23:601–649.
- Radtke, F., A. Wilson, G. Stark, M. Bauer, J. van Meerwijk, H.R. MacDonald, and M. Aguet. 1999. Deficient T cell fate specification in mice with an induced inactivation of Notch1. *Immunity*. 10:547–558.
- Ciofani, M., and J.C. Zuniga-Pflucker. 2005. Notch promotes survival of pre-T cells at the beta-selection checkpoint by regulating cellular metabolism. *Nat. Immunol.* 6:881–888.
- Weng, A.P., A.A. Ferrando, W. Lee, J.P. Morris IV, L.B. Silverman, C. Sanchez-Irizarry, S.C. Blacklow, A.T. Look, and J.C. Aster. 2004. Activating mutations of NOTCH1 in human T cell acute lymphoblastic leukemia. *Science*. 306:269–271.
- Reschly, E.J., C. Spaulding, T. Vilimas, W.V. Graham, R.L. Brumbaugh, I. Aifantis, W.S. Pear, and B.L. Kee. 2006. Notch1 promotes survival of E2A-deficient T cell lymphomas through Pre-T cell receptor dependent and independent mechanisms. *Blood*. 107:4115–4121.
- Dumortier, A., R. Jeannot, P. Kirstetter, E. Kleinmann, M. Sellars, N.R. dos Santos, C. Thibault, J. Barths, J. Ghysdael, J.A. Punt, et al. 2006. Notch activation is an early and critical event during T-cell leukemogenesis in Ikaros-deficient mice. *Mol. Cell. Biol.* 26:209–220.

11. Lin, Y.W., R.A. Nichols, J.J. Letterio, and P.D. Aplan. 2006. Notch1 mutations are important for leukemic transformation in murine models of precursor-T leukemia/lymphoma. *Blood*. 107:2540–2543.
12. O'Neil, J., J. Calvo, K. McKenna, V. Krishnamoorthy, J.C. Aster, C.H. Bassing, F.W. Alt, M. Kelliher, and A.T. Look. 2006. Activating Notch1 mutations in mouse models of T-ALL. *Blood*. 107:781–785.
13. Shin, H.M., L.M. Minter, O.H. Cho, S. Gottipati, A.H. Fauq, T.E. Golde, G.E. Sonenshein, and B.A. Osborne. 2006. Notch1 augments NF-kappaB activity by facilitating its nuclear retention. *EMBO J*. 25:129–138.
14. Vilimas, T., J. Mascarenhas, T. Palomero, M. Mandal, S. Buonamici, F. Meng, B. Thompson, C. Spaulding, S. Macaroun, M.L. Alegre, et al. 2006. Targeting the NF-kappaB signaling pathway in Notch1-induced T-cell leukemia. *Nat. Med.* 13:70–77.
15. Vilimas, T., J. Mascarenhas, T. Palomero, M. Mandal, S. Buonamici, F. Meng, B. Thompson, C. Spaulding, S. Macaroun, M.L. Alegre, et al. 2007. Targeting the NF-kappaB signaling pathway in Notch1-induced T-cell leukemia. *Nat. Med.* 13:70–77.
16. Weng, A.P., J.M. Millholland, Y. Yashiro-Ohtani, M.L. Arcangeli, A. Lau, C. Wai, C. Del Bianco, C.G. Rodriguez, H. Sai, J. Tobias, et al. 2006. c-Myc is an important direct target of Notch1 in T-cell acute lymphoblastic leukemia/lymphoma. *Genes Dev.* 20:2096–2109.
17. Palomero, T., W.K. Lim, D.T. Odom, M.L. Sulis, P.J. Real, A. Margolin, K.C. Barnes, J. O'Neil, D. Neuberg, A.P. Weng, et al. 2006. NOTCH1 directly regulates c-MYC and activates a feed-forward-loop transcriptional network promoting leukemic cell growth. *Proc. Natl. Acad. Sci. USA*. 103:18261–18266.
18. Klinakis, A., M. Szabolcs, K. Politi, H. Kiaris, S. Artavanis-Tsakonas, and A. Efstratiadis. 2006. Myc is a Notch1 transcriptional target and a requisite for Notch1-induced mammary tumorigenesis in mice. *Proc. Natl. Acad. Sci. USA*. 103:9262–9267.
19. Hubbard, E.J., G. Wu, J. Kitajewski, and I. Greenwald. 1997. sel-10, a negative regulator of lin-12 activity in *Caenorhabditis elegans*, encodes a member of the CDC4 family of proteins. *Genes Dev.* 11:3182–3193.
20. Nakayama, K.I., and K. Nakayama. 2005. Regulation of the cell cycle by SCF-type ubiquitin ligases. *Semin. Cell Dev. Biol.* 16:323–333.
21. Welcker, M., A. Orian, J. Jin, J.E. Grim, J.W. Harper, R.N. Eisenman, and B.E. Clurman. 2004. The Fbw7 tumor suppressor regulates glycogen synthase kinase 3 phosphorylation-dependent c-Myc protein degradation. *Proc. Natl. Acad. Sci. USA*. 101:9085–9090.
22. Wei, W., J. Jin, S. Schlisio, J.W. Harper, and W.G. Kaelin Jr. 2005. The v-Jun point mutation allows c-Jun to escape GSK3-dependent recognition and destruction by the Fbw7 ubiquitin ligase. *Cancer Cell*. 8:25–33.
23. Koepp, D.M., L.K. Schaefer, X. Ye, K. Keyomarsi, C. Chu, J.W. Harper, and S.J. Elledge. 2001. Phosphorylation-dependent ubiquitination of cyclin E by the SCFFbw7 ubiquitin ligase. *Science*. 294:173–177.
24. Gupta-Rossi, N., O. Le Bail, H. Gonen, C. Brou, F. Logeat, E. Six, A. Ciechanover, and A. Israel. 2001. Functional interaction between SEL-10, an F-box protein, and the nuclear form of activated Notch1 receptor. *J. Biol. Chem.* 276:34371–34378.
25. Oberg, C., J. Li, A. Pauley, E. Wolf, M. Gurney, and U. Lendahl. 2001. The Notch intracellular domain is ubiquitinated and negatively regulated by the mammalian Sel-10 homolog. *J. Biol. Chem.* 276:35847–35853.
26. Tetzlaff, M.T., W. Yu, M. Li, P. Zhang, M. Finegold, K. Mahon, J.W. Harper, R.J. Schwartz, and S.J. Elledge. 2004. Defective cardiovascular development and elevated cyclin E and Notch proteins in mice lacking the Fbw7 F-box protein. *Proc. Natl. Acad. Sci. USA*. 101:3338–3345.
27. Tsunematsu, R., K. Nakayama, Y. Oike, M. Nishiyama, N. Ishida, S. Hatakeyama, Y. Bessho, R. Kageyama, T. Suda, and K.I. Nakayama. 2004. Mouse Fbw7/Sel-10/Cdc4 is required for notch degradation during vascular development. *J. Biol. Chem.* 279:9417–9423.
28. Chiang, M.Y., M.L. Xu, G. Histén, O. Shestova, M. Roy, Y. Nam, S.C. Blacklow, D.B. Sacks, W.S. Pear, and J.C. Aster. 2006. Identification of a conserved negative regulatory sequence that influences the leukemogenic activity of NOTCH1. *Mol. Cell. Biol.* 26:6261–6271.
29. Welcker, M., J. Singer, K.R. Loeb, J. Grim, A. Bloecher, M. Gurien-West, B.E. Clurman, and J.M. Roberts. 2003. Multisite phosphorylation by Cdk2 and GSK3 controls cyclin E degradation. *Mol. Cell*. 12:381–392.
30. Welcker, M., and B.E. Clurman. 2005. The SV40 large T antigen contains a decoy phosphodegron that mediates its interactions with Fbw7/hCdc4. *J. Biol. Chem.* 280:7654–7658.
31. Fryer, C.J., J.B. White, and K.A. Jones. 2004. Mastermind recruits CycC:CDK8 to phosphorylate the Notch ICD and coordinate activation with turnover. *Mol. Cell*. 16:509–520.
32. Galili, U., A. Peleg, Y. Milner, and N. Galili. 1984. Be13, a human T-leukemia cell line highly sensitive to dexamethasone-induced cytolysis. *Cancer Res.* 44:4594–4601.
33. Orlicky, S., X. Tang, A. Willems, M. Tyers, and F. Sicheri. 2003. Structural basis for phosphodependent substrate selection and orientation by the SCFCdc4 ubiquitin ligase. *Cell*. 112:243–256.
34. Nash, P., X. Tang, S. Orlicky, Q. Chen, F.B. Gertler, M.D. Mendenhall, F. Sicheri, T. Pawson, and M. Tyers. 2001. Multisite phosphorylation of a CDK inhibitor sets a threshold for the onset of DNA replication. *Nature*. 414:514–521.
35. Kemp, Z., A. Rowan, W. Chambers, N. Wortham, S. Halford, O. Sieber, N. Mortensen, A. von Herbay, T. Gunther, M. Ilyas, and I. Tomlinson. 2005. CDC4 mutations occur in a subset of colorectal cancers but are not predicted to cause loss of function and are not associated with chromosomal instability. *Cancer Res.* 65:11361–11366.
36. Strohmaier, H., C.H. Spruck, P. Kaiser, K.A. Won, O. Sangfelt, and S.I. Reed. 2001. Human F-box protein hCdc4 targets cyclin E for proteolysis and is mutated in a breast cancer cell line. *Nature*. 413:316–322.
37. Mao, J.H., J. Perez-Losada, D. Wu, R. Delrosario, R. Tsunematsu, K.I. Nakayama, K. Brown, S. Bryson, and A. Balmain. 2004. Fbxw7/Cdc4 is a p53-dependent, haploinsufficient tumour suppressor gene. *Nature*. 432:775–779.
38. Nowak, D., M. Mossner, C.D. Baldus, O. Hopfer, E. Thiel, and W.K. Hofmann. 2006. Mutation analysis of hCDC4 in AML cells identifies a new intronic polymorphism. *Int J Med Sci.* 3:148–151.
39. Sharma, V.M., J.A. Calvo, K.M. Draheim, L.A. Cunningham, N. Hermance, L. Beverly, V. Krishnamoorthy, M. Bhasin, A.J. Capobianco, and M.A. Kelliher. 2006. Notch1 contributes to mouse T-cell leukemia by directly inducing the expression of c-myc. *Mol. Cell. Biol.* 26:8022–8031.
40. Minella, A.C., M. Welcker, and B.E. Clurman. 2005. Ras activity regulates cyclin E degradation by the Fbw7 pathway. *Proc. Natl. Acad. Sci. USA*. 102:9649–9654.
41. Germano, G., L. del Giudice, L. Lo Nigro, K. Polato, E. Giarin, M. Paganin, and G. Basso. 2005. Comparative sequence analysis of incomplete DJH and TCR gene rearrangements in children with relapses of T-ALL. *Leukemia*. 19:1687–1689.
42. Ory, D.S., B.A. Neugeboren, and R.C. Mulligan. 1996. A stable human-derived packaging cell line for production of high titer retrovirus/vesicular stomatitis virus G pseudotypes. *Proc. Natl. Acad. Sci. USA*. 93:11400–11406.