



THE SET OF QUANTUM CORRELATIONS IS NOT CLOSED

WILLIAM SLOFSTRA

Department of Pure Mathematics and Institute for Quantum Computing, University of Waterloo,
Waterloo, ON, Canada;
email: weslofst@uwaterloo.ca

Received 18 October 2017; accepted 2 November 2018

Abstract

We construct a linear system nonlocal game which can be played perfectly using a limit of finite-dimensional quantum strategies, but which cannot be played perfectly on any finite-dimensional Hilbert space, or even with any tensor-product strategy. In particular, this shows that the set of (tensor-product) quantum correlations is not closed. The constructed nonlocal game provides another counterexample to the ‘middle’ Tsirelson problem, with a shorter proof than our previous paper (though at the loss of the universal embedding theorem). We also show that it is undecidable to determine if a linear system game can be played perfectly with a finite-dimensional strategy, or a limit of finite-dimensional quantum strategies.

2010 Mathematics Subject Classification: 81P45 (primary); 20F05 (secondary)

1. Introduction

A two-player nonlocal game \mathcal{G} consists of finite question sets \mathcal{I}_A and \mathcal{I}_B , finite output sets \mathcal{O}_A and \mathcal{O}_B , and a function $V : \mathcal{O}_A \times \mathcal{O}_B \times \mathcal{I}_A \times \mathcal{I}_B \rightarrow \{0, 1\}$. During the game, the two players, commonly called Alice and Bob, are given inputs $x \in \mathcal{I}_A$ and $y \in \mathcal{I}_B$, respectively, and return outputs $a \in \mathcal{O}_A$ and $b \in \mathcal{O}_B$, respectively. The players win if $V(a, b | x, y) = 1$, and lose if $V(a, b | x, y) = 0$. The players know the rules of the game, and can decide ahead of time on their strategy. However, once the game is in progress, they are unable to communicate, meaning they do not know each other’s inputs or subsequent choices. This can make it impossible for the players to win some games with certainty.

© The Author 2019. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike licence (<http://creativecommons.org/licenses/by-nc-sa/4.0/>), which permits noncommercial re-use, distribution, and reproduction in any medium, provided the same Creative Commons licence is included and the original work is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use.

Imagine that the game is played repeatedly. To an outside observer, Alice and Bob’s actions during the game are described by the probability $p(a, b \mid x, y)$ that Alice and Bob output $a \in \mathcal{O}_A$ and $b \in \mathcal{O}_B$ on inputs $x \in \mathcal{I}_A$ and $y \in \mathcal{I}_B$. The collection $\{p(a, b \mid x, y)\} \subset \mathbb{R}^{\mathcal{O}_A \times \mathcal{O}_B \times \mathcal{I}_A \times \mathcal{I}_B}$ is called a *correlation matrix* (or a *behaviour*). Which correlation matrices can be achieved depends on the physical model. For instance, a correlation matrix $\{p(a, b \mid x, y)\}$ is said to be *classical* if it can be achieved using classical shared randomness. Formally, this means that there must be some integer $k \geq 1$, a probability distribution $\{\lambda_i\}$ on $\{1, \dots, k\}$, probability distributions $\{p_a^{ix}\}$ on \mathcal{O}_A for each $1 \leq i \leq k$ and $x \in \mathcal{I}_A$, and probability distributions $\{q_b^{iy}\}$ on \mathcal{O}_B for each $1 \leq i \leq k$ and $y \in \mathcal{I}_B$, such that

$$p(a, b \mid x, y) = \sum_{i=1}^k \lambda_i p_a^{ix} q_b^{iy} \quad \text{for all } (a, b, x, y) \in \mathcal{O}_A \times \mathcal{O}_B \times \mathcal{I}_A \times \mathcal{I}_B.$$

The set of classical correlation matrices is denoted by $C_c(\mathcal{O}_A, \mathcal{O}_B, \mathcal{I}_A, \mathcal{I}_B)$, although we typically write C_c when the output and input sets are clear.

In quantum information, we are interested in what correlations can be achieved with a shared quantum state. Accordingly, a correlation matrix is said to be *quantum* if there are finite-dimensional Hilbert spaces H_A and H_B , a quantum state $|\psi\rangle \in H_A \otimes H_B$, projective measurements $\{M_a^x\}_{a \in \mathcal{O}_A}$ on H_A for every $x \in \mathcal{I}_A$, and projective measurements $\{N_b^y\}_{b \in \mathcal{O}_B}$ on H_B for every $y \in \mathcal{I}_B$, such that

$$p(a, b \mid x, y) = \langle \psi \mid M_a^x \otimes N_b^y \mid \psi \rangle \quad \text{for all } (a, b, x, y) \in \mathcal{O}_A \times \mathcal{O}_B \times \mathcal{I}_A \times \mathcal{I}_B.$$

A projective measurement on a Hilbert space H is a collection $\{P_x\}_{x \in X}$ of self-adjoint operators on H , such that $P_x^2 = P_x$ for all $x \in X$, and $\sum_{x \in X} P_x = \mathbb{1}$. The set X is interpreted as the set of measurement outcomes.

The set of quantum correlation matrices is denoted by $C_q \cong C_q(\mathcal{O}_A, \mathcal{O}_B, \mathcal{I}_A, \mathcal{I}_B)$. There are two natural variations on this definition. We can drop the requirement that H_A and H_B be finite-dimensional, in which case we get another set of correlations often denoted by C_{qs} . We can also look at correlations which can be realized as limits of finite-dimensional quantum correlations; the corresponding correlation set is the closure of C_q , and is typically denoted by C_{qa} . It is well known that $C_{qs} \subseteq C_{qa}$, and consequently C_{qa} is also the closure of C_{qs} [29].

Since $C_{qs} \subseteq C_{qa}$, we get a hierarchy of correlation sets

$$C_c \subseteq C_q \subseteq C_{qs} \subseteq C_{qa}.$$

All the sets involved are convex, and C_c and C_{qa} are both closed. Bell’s celebrated theorem [2] states that $C_c \neq C_q$, and furthermore that the two sets

can be separated by a hyperplane. It has been a longstanding open problem to determine the relationship between the quantum correlation sets, and in particular to determine whether C_q and C_{q_s} are closed (see [3, 9, 32, 33]). Part of the interest in this latter question comes from the resource theory of nonlocal games: $C_q \neq C_{q_a}$ if and only if there is a nonlocal game which can be played optimally (with respect to some probability distribution on inputs) using a limit of finite-dimensional quantum strategies, but cannot be played optimally using any fixed dimension. Numerical evidence has suggested that even very simple nonlocal games might have this property [18, 25]. For variants of nonlocal games (for instance, with quantum questions, or infinite output sets), there are several examples of games with this property [17, 21, 28].

The purpose of this paper is to show that there are indeed nonlocal games (with finite classical input and output sets) that cannot be played optimally using any fixed dimension. A *perfect strategy* for a nonlocal game \mathcal{G} is a correlation matrix $\{p(a, b \mid x, y)\}$ such that Alice and Bob win with probability one on every pair of inputs x and y . Formally, this means that for all $(a, b, x, y) \in \mathcal{O}_A \times \mathcal{O}_B \times \mathcal{I}_A \times \mathcal{I}_B$, if $V(a, b \mid x, y) = 0$, then $p(a, b \mid x, y) = 0$.

THEOREM 1. *There is a nonlocal game with a perfect strategy in C_{q_a} , but no perfect strategy in C_{q_s} .*

In particular, neither C_q or C_{q_s} are closed. The proof is constructive, with the game in question having input sets of size 184 and 235, and output sets of size 8 and 2.

The set C_q is related to the cone of completely positive-semidefinite (cpsd) matrices defined in [16]. An $n \times n$ matrix M is said to be cpsd if there are nonnegative operators P_1, \dots, P_n on some finite-dimensional Hilbert space with $M_{ij} = \text{tr}(P_i P_j)$ for all $1 \leq i, j \leq n$. By a theorem of Sikora and Varvitsiotis [30], the set C_q is an affine slice of the cone of cpsd matrices, so the cone of cpsd matrices is not closed as a consequence of Theorem 1.

The fact that $C_{q_s} \neq C_{q_a}$ also has an interesting reformulation. Let G_i be the n -fold free product $\mathbb{Z}_m * \dots * \mathbb{Z}_m$, where $n = |\mathcal{I}_i|$ and $m = |\mathcal{O}_i|$, for $i = A, B$. Let M_a^x denote the a th spectral projector of the x th factor of G_A in the full group C^* -algebra $C^*(G_A)$ of G_A , and define M_b^y similarly for $C^*(G_B)$. For each $i = A, B$, find a faithful representation ν_i of $C^*(G_i)$ on some Hilbert space H_i . The minimal (or spatial) tensor product $C^*(G_A) \otimes_s C^*(G_B)$ is the norm closure of the image $\nu_A(C^*(G_A)) \otimes \nu_B(C^*(G_B))$ in the C^* -algebra $\mathcal{B}(H_A \otimes H_B)$. A correlation matrix $\{p(a, b \mid x, y)\}$ belongs to C_{q_a} if and only if there is a state ω on the C^* -algebra $C^*(G_A) \otimes_s C^*(G_B)$ with

$$p(a, b \mid x, y) = \omega(M_a^x \otimes N_b^y)$$

for all $(a, b, x, y) \in \mathcal{O}_A \times \mathcal{O}_B \times \mathcal{I}_A \times \mathcal{I}_B$ [9, 29]. On the other hand, the correlation matrix belongs to C_{qs} if and only if there are representations ϕ_i of G_i on H_i , $i = A, B$, and a vector state $|\psi\rangle \in H_A \otimes H_B$, with

$$p(a, b \mid x, y) = \langle \psi \mid \phi_A(M_a^x) \otimes \phi_B(N_b^y) \mid \psi \rangle$$

for all $(a, b, x, y) \in \mathcal{O}_A \times \mathcal{O}_B \times \mathcal{I}_A \times \mathcal{I}_B$. Since $C_{qs} \neq C_{qa}$, there can be states on the minimal tensor product $C^*(G_A) \otimes_s C^*(G_B)$ which do not come from vector states on some tensor product $\phi_A \otimes \phi_B$ of representations ϕ_A and ϕ_B .

There is another candidate set of quantum correlations, the commuting-operator correlations C_{qc} , which contains C_{qa} . Determining whether C_{qc} is equal to C_t for any $t \in \{q, qs, qa\}$ is known as Tsirelson’s problem [7, 32]. A theorem of Ozawa [24] (see also [9, 13]) states that $C_{qa} = C_{qc}$ if and only if there is an affirmative answer to the Connes embedding problem. In a previous paper [31], we showed that $C_{qs} \neq C_{qc}$. By showing that $C_{qs} \neq C_{qa}$, we provide another proof of this fact. The proof that $C_{qs} \neq C_{qc}$ in [31] uses a universal embedding theorem, which states that every finitely presented group embeds in the solution group of a linear system game. In this paper, we follow a similar line, proving a restricted embedding theorem for a subclass of finitely presented groups which we call linear-plus-conjugacy groups. For the proof of this restricted embedding theorem, we use a completely different method from [31], with the result that the proof is much shorter. However, it remains an open problem to prove the universal embedding theorem via the new approach.

If we choose a probability distribution π on questions $\mathcal{I}_A \times \mathcal{I}_B$, then the probability that the players win when their behaviour is described by correlation matrix $\{p(a, b \mid x, y)\}$ is

$$\sum_{x,y,a,b} \pi(x, y) V(a, b \mid x, y) p(a, b \mid x, y).$$

The *quantum value* of the game (with probability distribution π) is the optimal winning probability for the game over correlations in C_{qa} . A fundamental question about nonlocal games is whether it is possible to compute the quantum value, either exactly or approximately. As long as $\pi(x, y) \neq 0$ for every $(x, y) \in \mathcal{I}_A \times \mathcal{I}_B$, then the quantum value of a game is equal to 1 if and only if the game has a perfect strategy in C_{qa} . Thus as a special case of the exact computation problem, we can ask whether it is possible to determine whether a nonlocal game has a perfect quantum strategy. (There seem to be several reasonable ways to formalize the general problem of computing the exact value of a nonlocal game. One way is as the problem of determining, from a given nonlocal game and a rational number a , whether the quantum value of the game is $\geq a$. Another reasonable formalization is the problem of computing,

from a given nonlocal game and positive integer n , the n th decimal digit of the quantum value of the game, where the decimal expansion used must be the unique expansion which does not terminate in a repeating sequence of 9's. Both of these formalizations contain the problem of determining whether a game has a perfect quantum strategy as a special case.)

Recall that a decision problem of the form 'Is $P(x)$ true?' is *decidable* if there is a Turing machine which halts and outputs whether or not $P(x)$ is true on every input x . If no such Turing machine exists, then the problem is said to be *undecidable*. A set \mathcal{L} is *recursive* if the decision problem 'Is $x \in \mathcal{L}$?' is decidable. An easy consequence of the universal embedding theorem is that it is undecidable to determine if a linear system game has a perfect strategy in C_{qc} . In this paper we prove a stronger result by applying our restricted embedding theorem to Kharlampovich's example [14] of a finitely presented solvable group with an undecidable word problem.

THEOREM 2. *There is a (recursive) family of linear system games such that:*

- (a) *it is undecidable to determine if a game in the family has a perfect strategy in C_{qa} ; and*
- (b) *every game in the family has a perfect strategy in C_{qc} if and only if it has a perfect strategy in C_{qa} .*

Theorem 2 can be interpreted as saying that there is no algorithm to compute the quantum value of a nonlocal game exactly.

A function $f: \mathbb{N} \rightarrow Y$ to some set Y is *computable* if there is a Turing machine which halts with output $f(n)$ on every input n , and *computable in $T(n)$ -time* if there is such a Turing machine which halts within $T(n)$ steps on input n . The computation time of a Turing machine is often defined using the size of inputs n , rather than the values of n . However, to avoid introducing an extra parameter for the length, or discuss encodings of inputs, we always just refer to computation time on specific inputs. Kharlampovich's construction has been extended by Kharlampovich, Myasnikov, and Sapir to show that the word problem for finitely presented residually finite groups can be as hard as any computable function [15]. The word problem for finitely presented residually finite groups is always decidable, so this is the best possible lower bound. Using this extension, we can show:

THEOREM 3. *Let $f: \mathbb{N} \rightarrow \mathbb{N}$ be a computable function. Then there is a family of linear system games \mathcal{G}_n , $n \in \mathbb{N}$, such that:*

- (a) *the games \mathcal{G}_n have input sets of size $\exp(O(n))$, and the function $n \mapsto \mathcal{G}_n$ is computable in $\exp(O(n))$ -time;*

(b) for any algorithm accepting the language

$$\{n \in \mathbb{N} : \mathcal{G}_n \text{ has a perfect strategy in } C_q\},$$

the maximum running time over inputs $n \leq N$ is at least $f(N)$ when N is sufficiently large;

(c) \mathcal{G}_n has a perfect strategy in C_{qc} if and only if it has a perfect strategy in C_q .

Theorem 3 has the following corollary.

COROLLARY 4. *It is undecidable to determine if a linear system game has a perfect strategy in C_q .*

Theorems 2 and 3 say nothing about whether it is possible to approximate the quantum value of a nonlocal game. Formally speaking, the approximation problem is to compute a rational number r from a given game and rational number $\epsilon > 0$, such that r is within ϵ of the quantum value of the given game. This problem is complete for the complexity class MIP^* of multi-prover interactive proofs with entangled provers [12]. Determining whether there is an algorithm to approximate the quantum value (or equivalently, whether MIP^* is computable) is a major open problem in quantum complexity theory. Navascués, Pironio, and Acin have shown that such an algorithm would exist for two-player games if the Connes embedding problem has an affirmative answer [23].

2. Group theory preliminaries

2.1. Group presentations. Given a set S , let $\mathcal{F}(S)$ denote the free group generated by S . If H is a group, then homomorphisms $\mathcal{F}(S) \rightarrow H$ can be identified with functions $S \rightarrow H$, and we use these two types of objects interchangeably. If R is a subset of $\mathcal{F}(S)$, then the quotient of $\mathcal{F}(S)$ by the normal subgroup generated by R is denoted by $\langle S : R \rangle$. If $G = \langle S : R \rangle$ and $R' \subset \mathcal{F}(S \cup S')$, then we write $\langle G, S' : R' \rangle$ to mean $\langle S \cup S' : R \cup R' \rangle$. If $G = \langle S : R \rangle$ and $H = \langle S' : R' \rangle$, then any homomorphism $\psi : G \rightarrow H$ lifts to a homomorphism $\Psi : \mathcal{F}(S) \rightarrow \mathcal{F}(S')$ such that for any $w \in \mathcal{F}(S)$, the image of $\Psi(w)$ in H is equal to $\psi(w)$. Lifts of this type are not unique unless $R' \subseteq \{e\}$.

A group G is said to be *finitely presentable* if $G = \langle S : R \rangle$ for some finite sets S and R . A *finitely presented group* is a tuple (G, S, R) , where $G = \langle S : R \rangle$. In other words, a finitely presented group is a finitely presentable group along with a choice of finite presentation.

For the purposes of this paper, a representation of G will always mean a unitary representation, that is a homomorphism from G to the unitary group $\mathcal{U}(H)$ of

some (possibly infinite-dimensional) Hilbert space H . If $G = \langle S : R \rangle$, then a representation is the same thing as a homomorphism $\phi : \mathcal{F}(S) \rightarrow \mathcal{U}(H)$ such that $\phi(r) = \mathbb{1}$ for all $r \in R$.

2.2. Approximate representations. Let $\|\cdot\|$ be the normalized Hilbert–Schmidt norm, so if T is an endomorphism of a finite-dimensional Hilbert space H , then $\|T\| = \sqrt{\text{tr}(T^*T)}/\sqrt{\dim H}$.

DEFINITION 5. Let $G = \langle S : R \rangle$ be a finitely presented group. A *finite-dimensional ϵ -approximate representation* (or *ϵ -representation* for short) of G is a homomorphism $\phi : \mathcal{F}(S) \rightarrow \mathcal{U}(H)$ from $\mathcal{F}(S)$ to the unitary group $\mathcal{U}(H)$ of some finite-dimensional Hilbert space H , such that

$$\|\phi(r) - \mathbb{1}\| \leq \epsilon$$

for all $r \in R$.

The normalized Hilbert–Schmidt norm is invariant under conjugation by unitaries, so the set of ϵ -representations is independent of the cyclic order of the relations $r \in R$. That means that, for instance, we can write the relation $x = y$ without worrying about whether we mean $xy^{-1} = e$ or $y^{-1}x = e$. We use the term *approximate representation* for any homomorphism $\phi : \mathcal{F}(S) \rightarrow \mathcal{U}(H)$, where H is a finite-dimensional Hilbert space. Since the norm is not actually involved, this terminology is somewhat redundant, but it is helpful for situations where we do not want to specify ϵ or H . Note that, in contrast to ordinary representations, we assume that approximate representations are always finite-dimensional (however, both types of representations are always assumed to be unitary). As a result, a 0-representation of G is the same thing as a finite-dimensional representation of G .

EXAMPLE 6. The group \mathbb{Z}_2^k has presentation

$$\langle x_1, \dots, x_k : x_i^2 = e \text{ for all } 1 \leq i \leq k, [x_i, x_j] = e \text{ for all } 1 \leq i \neq j \leq k \rangle,$$

where $[x, y] := xyx^{-1}y^{-1}$. An ϵ -representation of \mathbb{Z}_2^k with this presentation is a homomorphism

$$\phi : \mathcal{F}(\{x_1, \dots, x_k\}) \rightarrow \mathcal{U}(\mathbb{C}^d)$$

for some d , such that

$$\|\phi(x_i)^2 - \mathbb{1}\| \leq \epsilon \quad \text{for all } 1 \leq i \leq k \tag{1}$$

and

$$\|\phi(x_i)\phi(x_j)\phi(x_i)^*\phi(x_j)^* - \mathbb{1}\| \leq \epsilon \quad \text{for all } 1 \leq i \neq j \leq k. \tag{2}$$

Since the homomorphism ϕ is determined by its values on $\{x_1, \dots, x_k\}$, we can think of an ϵ -representation concretely as a tuple of unitaries $(\phi(x_1), \dots, \phi(x_k))$ in $\mathcal{U}(\mathbb{C}^d)$ satisfying Equations (1) and (2).

There are several different notions of approximate representations in the literature. The notion we are using comes from the study of stable relations of C^* -algebras (see, for instance, [19, Section 4.1]). For the purposes of this paper, we could also use the closely related notion of approximate homomorphisms as in [4, Section II]. However, Definition 5 is very convenient for working with examples, as we frequently do in this paper. The main disadvantage of this definition is that it depends on the choice of presentation. We can work around this using the following easy lemma.

LEMMA 7. *Let $\psi : G \rightarrow H$ be a homomorphism, where $G = \langle S : R \rangle$ and $H = \langle S' : R' \rangle$ are finitely presented groups. If $\Psi : \mathcal{F}(S) \rightarrow \mathcal{F}(S')$ is a lift of ψ , then there is a constant $C > 0$ such that if ϕ is an ϵ -representation of H , then $\phi \circ \Psi$ is a $C\epsilon$ -representation of G .*

Proof. Since Ψ is the lift of a homomorphism, for any $r \in R$ we have that $\Psi(r) = a_1 r_1^{b_1} a_1^{-1} \dots a_k r_k^{b_k} a_k^{-1}$ for some $k \geq 0$, $a_1, \dots, a_k \in \mathcal{F}(S')$, $r_1, \dots, r_k \in R'$, and $b_1, \dots, b_k \in \{\pm 1\}$. Suppose that ϕ is an ϵ -representation of H . The normalized Hilbert–Schmidt norm is invariant under right and left multiplication by unitaries (that is $\|AU\| = \|UA\| = \|A\|$ for any unitary U), so $\|\phi(a_i r_i^{b_i} a_i^{-1}) - \mathbb{1}\| = \|\phi(r_i) - \mathbb{1}\| \leq \epsilon$ for all $i = 1, \dots, k$. Hence

$$\begin{aligned} \|\phi \circ \Psi(r) - \mathbb{1}\| &= \|\phi(a_1 r_1 a_1^{-1} \dots a_k r_k a_k^{-1}) - \mathbb{1}\| \\ &\leq \|(\phi(a_1 r_1 a_1^{-1}) - \mathbb{1}) \phi(a_2 \dots a_k^{-1})\| \\ &\quad + \|(\phi(a_2 r_2 a_2^{-1}) - \mathbb{1}) \phi(a_3 \dots a_k^{-1})\| \\ &\quad + \dots + \|\phi(a_k r_k a_k^{-1}) - \mathbb{1}\| \leq k\epsilon, \end{aligned}$$

where the last inequality again uses the invariance of the norm under right multiplication by unitaries. Since R is finite, we can take C to be the maximum k across all $r \in R$. \square

We record two other simple lemmas for later use.

LEMMA 8. *Let $G = \langle S : R \rangle$, and let M be the length of the longest relation in R . If ϕ is an ϵ -representation of G , and ψ is an approximate representation of G with*

$$\|\psi(x) - \phi(x)\| \leq \delta$$

for all $x \in S$, then ψ is an $(M\delta + \epsilon)$ -representation.

Given approximate representations $\phi : \mathcal{F}(S) \rightarrow \mathcal{U}(H)$ and $\psi : \mathcal{F}(S) \rightarrow \mathcal{U}(H')$ of $G = \langle S : R \rangle$, we can form new approximate representations $\phi \oplus \psi : \mathcal{F}(S) \rightarrow \mathcal{U}(H \oplus H')$ and $\phi \otimes \psi : \mathcal{F}(S) \rightarrow \mathcal{U}(H \otimes H')$.

LEMMA 9. Suppose ϕ and ψ are ϵ - and ϵ' -representations of G , respectively. Then $\phi \oplus \psi$ is a $\max(\epsilon, \epsilon')$ -representation, and $\phi \otimes \psi$ is an $(\epsilon + \epsilon')$ -representation.

A group G is said to be residually finite-dimensional if every nontrivial element of G is nontrivial in some finite-dimensional representation. More generally, the set of elements which are trivial in finite-dimensional representations forms a normal subgroup of G . We let G^{fin} denote the quotient of G by this normal subgroup (alternatively, G^{fin} is the image of G in its profinite completion). Any homomorphism $\phi : G \rightarrow H$ descends to a homomorphism $G^{\text{fin}} \rightarrow H^{\text{fin}}$.

DEFINITION 10. A homomorphism $\phi : G \rightarrow H$ is a *fin-embedding* if the induced map $G^{\text{fin}} \rightarrow H^{\text{fin}}$ is injective, and a *fin*-embedding* if ϕ is both injective and a *fin-embedding*.

Equivalently, ϕ is a *fin-embedding* if $\phi(g)$ is nontrivial in finite-dimensional representations whenever $g \in G$ is nontrivial in finite-dimensional representations.

We can similarly look at elements which are nontrivial in approximate representations:

DEFINITION 11. Let G be a finitely presentable group. An element $g \in G$ is *nontrivial in (finite-dimensional) approximate representations* if there is a finite presentation $G = \langle S : R \rangle$, a representative $w \in \mathcal{F}(S)$ for g , and some constant $\delta > 0$ such that, for all $\epsilon > 0$, there is an ϵ -representation ϕ of G with $\|\phi(w) - \mathbb{1}\| > \delta$.

Alternatively, if $g \in G = \langle S : R \rangle$, let

$$\ell^{fa}(g) := \limsup_{\epsilon \rightarrow 0^+} \sup_{\phi} \|\phi(w) - \mathbb{1}\|,$$

where w is a representative for g , and the supremum is across ϵ -representations ϕ of G . It is easy to see that the right-hand side is independent of the choice of representative w . By Lemma 7, if $\psi : G \rightarrow H$ is a homomorphism, then $\ell^{fa}(g) \geq \ell^{fa}(\psi(g))$. Consequently, $\ell^{fa}(g)$ is independent of the chosen presentation $\langle S : R \rangle$, and g is nontrivial in approximate representations if and

only if $\ell^{fa}(g) > 0$. This makes it apparent that the choice of presentation $\langle S : R \rangle$ and representative w in Definition 11 is arbitrary.

Standard amplification arguments show that the constant δ in Definition 11 is also somewhat arbitrary; in particular, any number in $(0, \sqrt{2})$ will work. Since we use these amplification arguments in a later section, we record them in the following lemma. If X is a linear operator on a finite-dimensional Hilbert space, let $\tilde{\text{tr}}(X) := \text{tr}(X)/\dim(H)$.

LEMMA 12. *Let $G = \langle S : R \rangle$, and suppose that the image of $w \in \mathcal{F}(S)$ in G is nontrivial in approximate representations of G . Then for every $\epsilon, \tau > 0$ there is an ϵ -representation ϕ with $0 \leq \tilde{\text{tr}}(\phi(w)) \leq \tau$.*

For the proof, we follow [4, Section II.2.2]:

Proof. First, let $2 \geq \delta > 0$ be such that for all $\epsilon > 0$, there is an ϵ -representation ϕ with $\|\phi(w) - \mathbb{1}\| \geq \delta$ (since Definition 11 does not depend on the choice of presentation and representative, such a δ exists).

Next, suppose $\phi : \mathcal{F}(S) \rightarrow \mathcal{U}(\mathbb{C}^d)$ is an ϵ -representation. Let $\bar{\phi}$ be the approximate representation defined by $\bar{\phi}(a) = \overline{\phi(a)}$, the entry-wise complex conjugate of $\phi(a)$ with respect to the standard basis on \mathbb{C}^d . It is not hard to check that $\bar{\phi}$ is also an ϵ -representation. Let γ be the direct sum of $\phi \oplus \bar{\phi}$ with $2d$ copies of the trivial representation of $\mathcal{F}(S)$. Then γ is an ϵ -representation of G by Lemma 9, $\text{tr}(\gamma(w)) = \text{tr}(\phi(w)) + \text{tr}(\bar{\phi}(w)) + 2d \geq 0$, and $\|\gamma(w) - \mathbb{1}\|^2 = \|\phi(w) - \mathbb{1}\|^2 / 2 \geq \delta^2 / 2$. Since

$$\|U - \mathbb{1}\|^2 = 2 - 2 \text{Re} \tilde{\text{tr}}(U) \tag{3}$$

for any unitary U , we see that

$$0 \leq \tilde{\text{tr}}(\gamma(w)) = \text{Re} \tilde{\text{tr}}(\gamma(w)) \leq 1 - \frac{\delta^2}{4}.$$

Finally, suppose we are given $\tau > 0$. Since $2 \geq \delta > 0$, we can find k such that $(1 - \delta^2/4)^k < \tau$. By the previous paragraph, for every $\epsilon > 0$, there is an ϵ/k -representation ϕ such that $0 \leq \tilde{\text{tr}}(\phi(w)) \leq 1 - \delta^2/4$. Thus, we can use the tensor-power trick: Since $\tilde{\text{tr}}(X^{\otimes k}) = \tilde{\text{tr}}(X)^k$, Lemma 9 implies that $\phi^{\otimes k}$ is an ϵ -representation of G with

$$0 \leq \tilde{\text{tr}}(\phi^{\otimes k}(w)) \leq \left(1 - \frac{\delta^2}{4}\right)^k \leq \tau,$$

as required. □

By Equation (3), if $\tilde{\text{tr}}(\phi(w)) \leq \tau$, then $\|\phi(w) - \mathbb{1}\| \geq \sqrt{2 - 2\tau}$. Thus Lemma 12 immediately implies that we can take any $\delta \in (0, \sqrt{2})$ in Definition 11. Equivalently, we can say that $\ell^{fa}(g)$ never takes values in $(0, \sqrt{2})$.

Let $\mathbb{R}_{>0}$ be the set of positive reals. Recall from [27] that a group G is hyperlinear if for every $\epsilon > 0, \delta \in (0, \sqrt{2})$, and finite subset $F \subseteq G$, there is a function $f : G \rightarrow \mathcal{U}(H)$, where H is a finite-dimensional Hilbert space, such that:

- $f(e) = \mathbb{1}$;
- $\|f(gh) - f(g)f(h)\| \leq \epsilon$ for all $g, h \in F$; and
- $\|f(g) - \mathbb{1}\| \geq \delta$ for all $g \in F \setminus \{e\}$.

The proof of Lemma 12 can be used to show:

LEMMA 13. A finitely presentable group G is hyperlinear if and only if every element of $G \setminus \{e\}$ is nontrivial in approximate representations.

Proof. Fix a finite presentation $G = \langle S : R \rangle$. We start with the claim that G is hyperlinear if and only if for every $\epsilon > 0, \delta \in (0, \sqrt{2})$, and finite subset $F_0 \subset \mathcal{F}(S)$ such that every word in F_0 has nontrivial image in G , there is an ϵ -representation ϕ with $\|\phi(w) - \mathbb{1}\| \geq \delta$. To prove one direction of the claim, suppose that G is hyperlinear, and that we are given $\epsilon > 0, \delta \in (0, \sqrt{2})$, and a finite subset $F_0 \subset \mathcal{F}(S)$ such that every word in F_0 has nontrivial image in G . Let $M = \max\{|r| : r \in R \cup F_0\} \cup \{2\}$, where $|w|$ denotes the length of $w \in \mathcal{F}(S)$, and let F be the image in G of all words in $\mathcal{F}(S)$ of length $< M$. Given a function $f : G \rightarrow \mathcal{U}(H)$, we can define $\phi : \mathcal{F}(S) \rightarrow \mathcal{U}(H)$ by setting $\phi(s) = f(s)$ for all $s \in S$. If $\|f(gh) - f(g)f(h)\| \leq \epsilon$ for all $g, h \in F$, then

$$\|\phi(s_1 \cdots s_k) - f(s_1 \cdots s_k)\| = \|f(s_1) \cdots f(s_k) - f(s_1 \cdots s_k)\| \leq (k - 1)\epsilon$$

for all $2 \leq k \leq M$. In particular, if $f(e) = \mathbb{1}$ then ϕ will be an $(M - 1)\epsilon$ -representation of G , and

$$\|\phi(w) - \mathbb{1}\| \geq \|f(w) - \mathbb{1}\| - \|\phi(w) - f(w)\| \geq \|f(w) - \mathbb{1}\| - (M - 1)\epsilon$$

for all $w \in F_0$. It follows from the definition of hyperlinearity that for every $\epsilon > 0$ and $\delta \in (0, \sqrt{2})$, there is an ϵ -representation ϕ with $\|\phi(w) - \mathbb{1}\| \geq \delta$ for all $w \in F_0$. The other direction of the claim is similar, and we leave it to the reader.

Now suppose that G is hyperlinear, and $g \in G \setminus \{e\}$. Applying the claim with $F_0 = \{w\}$, where $w \in \mathcal{F}(S)$ is some representative of g , we see that g is nontrivial in approximate representations of G . For the converse, suppose that every element of $G \setminus \{e\}$ is nontrivial in approximate representations, and that $F_0 \subset \mathcal{F}(S)$ is a finite set of words such that every element of F_0 is nontrivial in G . By Lemma 12, for every $\epsilon > 0$, $\delta \in (0, \sqrt{2})$, and $w \in F$ there is an ϵ -representation ϕ with $\|\phi(w) - \mathbb{1}\| \geq \delta$. By taking direct sums of these ϵ -representations, we can conclude that for every $\epsilon > 0$ and $\delta \in (0, \sqrt{2})$, there is an ϵ -representation ϕ such that $\|\phi(w) - \mathbb{1}\| \geq \delta/\sqrt{|F_0|}$ for every $w \in F_0$. To improve this lower bound and remove the dependence on $|F_0|$, note that the amplification procedure in the proof of Lemma 12 does not depend on w , and thus can be used to improve the lower bound on $\|\phi(w) - \mathbb{1}\|$ for all $w \in F_0$ simultaneously. It follows that for every $\epsilon > 0$ and every $\delta \in (0, \sqrt{2})$, there is an ϵ -representation ϕ with $\|\phi(w) - \mathbb{1}\| \geq \delta$ for all $w \in F_0$. \square

Clearly $\ell^{fa}(g) \geq 0$ for all $g \in G$, and it is easy to see that

$$\ell^{fa}(gh) \leq \ell^{fa}(g) + \ell^{fa}(h)$$

and $\ell^{fa}(hgh^{-1}) = \ell^{fa}(g)$ for all $g, h \in G$. Thus the set of elements of G which are trivial in approximate representations (that is for which $\ell^{fa}(g) = 0$) forms a normal subgroup of G . Let G^{fa} be the quotient of G by this normal subgroup. Because ℓ^{fa} is decreasing via homomorphisms, any homomorphism $\phi : G \rightarrow H$ between finitely presentable groups descends to a homomorphism $G^{fa} \rightarrow H^{fa}$.

DEFINITION 14. A homomorphism $\phi : G \rightarrow H$ is an *fa-embedding* if the induced map $G^{fa} \rightarrow H^{fa}$ is injective, and an *fa*-embedding* if ϕ is injective, a *fin-embedding*, and an *fa-embedding*.

Equivalently, ϕ is an *fa-embedding* if $\phi(g)$ is nontrivial in approximate representations whenever $g \in G$ is nontrivial in approximate representations.

If ϕ and ψ are approximate representations, then we say that ϕ is a *direct summand* of ψ if $\psi = \phi \oplus \phi'$ for some other approximate representation ϕ' . We use the following simple trick to construct *fa**-embeddings.

LEMMA 15. Let $G = \langle S : R \rangle$ and $H = \langle S' : R' \rangle$ be two finitely presented groups, and let $\Psi : \mathcal{F}(S) \rightarrow \mathcal{F}(S')$ be a lift of a homomorphism $\psi : G \rightarrow H$.

- (a) Suppose that for every representation (resp. finite-dimensional representation) ϕ of G , there is a representation (resp. finite-dimensional representation) γ of H such that ϕ is a direct summand of $\gamma \circ \psi$. Then ψ is injective (resp. a *fin-embedding*).

- (b) Suppose that there is an integer $N > 0$ and a real number $C > 0$ such that for every d -dimensional ϵ -representation ϕ of G , where $\epsilon > 0$, there is an Nd -dimensional $C\epsilon$ -representation γ of H such that ϕ is a direct summand of $\gamma \circ \Psi$. Then ψ is an fa -embedding.

Proof. Part (a) is clear, so we prove (b). Suppose ϕ is an ϵ -representation of G , where $\epsilon > 0$. If $\gamma \circ \Psi = \phi \oplus \phi'$, where ϕ is d -dimensional and ϕ' is $(N - 1)d$ -dimensional, then

$$\|\gamma(\Psi(w)) - \mathbb{1}\| = \|\phi(w) \oplus \phi'(w) - \mathbb{1}\| \geq \frac{1}{\sqrt{N}} \|\phi(w) - \mathbb{1}\|$$

for all $w \in \mathcal{F}(S)$. So $\ell^{fa}(\psi(g)) \geq \ell^{fa}(g)/\sqrt{N}$, and ψ is an fa -embedding. \square

In our applications it will be possible to check parts (a) and (b) of Lemma 15 simultaneously, in which case ψ will be an fa^* -embedding.

2.3. Groups over \mathbb{Z}_2 . For convenience, we use the following definition from [31]: A group over \mathbb{Z}_2 is a pair (G, J) , where J is a central element of G of order two. Note that J is allowed to be the identity element. Typically we drop the pair notation, and just use the symbol J (or J_G where necessary) to refer to the special element of a group G over \mathbb{Z}_2 , in the same way that we use e to refer to the identity element. If G and H are groups over \mathbb{Z}_2 , then a morphism $G \rightarrow H$ over \mathbb{Z}_2 is a group homomorphism $G \rightarrow H$ which sends $J_G \mapsto J_H$.

If a group G over \mathbb{Z}_2 is finitely presentable, then it has a finite presentation $\langle S : R \rangle$ where $J \in S$, and R includes the relations $J^2 = e$ and $[J, s] = e$ for every $s \in S \setminus \{J\}$. We use presentations of this form often enough that it is helpful to have some notation for them. Suppose that S_0 is a set of indeterminates, and $R_0 \subset \mathcal{F}(S_0 \cup \{J\})$. Then we set

$$\langle S_0 : R_0 \rangle_{\mathbb{Z}_2} := \langle S_0 \cup \{J\} : R_0 \cup \{[J, s] = e : s \in S_0\} \cup \{J^2 = e\} \rangle,$$

and call $\langle S_0 : R_0 \rangle_{\mathbb{Z}_2}$ a presentation over \mathbb{Z}_2 . As with ordinary presentations, if $G = \langle S : R \rangle$ or $\langle S : R \rangle_{\mathbb{Z}_2}$, then $\langle G, S' : R' \rangle_{\mathbb{Z}_2} := \langle S \cup S' : R \cup R' \rangle_{\mathbb{Z}_2}$.

EXAMPLE 16. Consider the group \mathbb{Z}_2^3 , presented as in Example 6. Then

$$\langle \mathbb{Z}_2^3, y : x_1x_2x_3y = J \rangle_{\mathbb{Z}_2}$$

is the finitely presented group

$$\begin{aligned} \langle x_1, x_2, x_3, y, J : x_i^2 = [x_i, J] = e \text{ for all } 1 \leq i \leq 3, \\ [x_i, x_j] = e \text{ for all } 1 \leq i \neq j \leq 3, \\ [y, J] = J^2 = e, \text{ and } x_1x_2x_3y = J \rangle. \end{aligned}$$

3. Linear system games and solution groups

Let $Ax = b$ be an $m \times n$ linear system over \mathbb{Z}_2 . To the system $Ax = b$, we can associate a nonlocal game, called a *linear system game*, as follows. For each $1 \leq i \leq m$, let $V_i = \{j : A_{ij} \neq 0\}$ be the set of indices of variables appearing in the i th equation. Let $S_i \subset \mathbb{Z}_2^{V_i}$ be the set of assignments to variables $x_j, j \in V_j$ satisfying the i th equation, that is $\underline{a} \in \mathbb{Z}_2^{V_i}$ belongs to S_i if and only if $\sum_{j \in V_j} a_j = b_i$. Then Alice receives an equation as input, represented by an integer $1 \leq i \leq m$, and must output an element $\underline{a} \in S_i$. Bob receives a variable, represented by an integer $1 \leq j \leq n$, and must output an assignment b for x_j . The players win if either $j \notin V_i$, or $j \in V_i$ and $a_j = b$, that is, Alice’s and Bob’s outputs are consistent.

A *quantum strategy (presented in terms of measurements)* for a linear system game consists of:

- (1) a pair of Hilbert spaces H_A and H_B ;
- (2) a projective measurement $\{N_b^j\}_{b \in \mathbb{Z}_2}$ on H_B for every integer $1 \leq j \leq n$;
- (3) a projective measurement $\{M_{\underline{a}}^i\}_{\underline{a} \in S_i}$ on H_A for every integer $1 \leq i \leq m$; and
- (4) a quantum state $|\psi\rangle \in H_A \otimes H_B$.

The strategy is *finite-dimensional* if H_A and H_B are finite-dimensional. The associated quantum correlation matrix $\{p(\underline{a}, b | i, j)\}$ is defined by

$$p(\underline{a}, b | i, j) = \langle \psi | M_{\underline{a}}^i \otimes N_b^j | \psi \rangle, \quad 1 \leq i \leq m, \quad 1 \leq j \leq n, \quad \underline{a} \in S_i, \quad b \in \mathbb{Z}_2.$$

As in the introduction, we also use the term *strategy* to refer to the correlation matrix $\{p(\underline{a}, b | i, j)\}$. If $j \in V_i$, then the probability that Alice and Bob win on inputs i and j is

$$p_{ij} := \sum_{\underline{a}, b : a_j = b} p(\underline{a}, b | i, j).$$

A strategy is *perfect* if and only if $p_{ij} = 1$ for all $1 \leq i \leq m$ and $j \in V_i$.

For linear system games, it is often convenient to work with strategies presented in terms of ± 1 -valued observables—self-adjoint operators which square to the identity—rather than measurement operators. A *quantum strategy (presented in terms of observables)* consists of

- (a) a pair of Hilbert spaces H_A and H_B ;
- (b) a collection of self-adjoint operators $X_j, 1 \leq j \leq n$, on H_B such that $X_j^2 = \mathbb{1}$ for every $1 \leq j \leq n$;

(c) a collection of self-adjoint operators Y_{ij} , $1 \leq i \leq m$, $j \in V_i$ on H_A such that:

- (i) $Y_{ij}^2 = \mathbb{1}$ for every $1 \leq i \leq m$ and $j \in V_i$,
- (ii) $\prod_{j \in V_i} Y_{ij} = (-\mathbb{1})^{b_i}$ for every $1 \leq i \leq m$, and
- (iii) $Y_{ij}Y_{il} = Y_{il}Y_{ij}$ for every $1 \leq i \leq m$ and $j, l \in V_i$; and

(d) a quantum state $|\psi\rangle \in H_A \otimes H_B$.

Given a quantum strategy presented in terms of measurements, we can get a quantum strategy presented in terms of observables by setting $X_j = N_0^j - N_1^j$ for every $1 \leq j \leq n$, and

$$Y_{ij} = \sum_{\underline{a} \in \mathcal{S}_i} (-1)^{a_j} M_{\underline{a}}^i$$

for $1 \leq i \leq m$ and $j \in V_i$. Conversely, given a quantum strategy in terms of observables, we can recover the measurement presentation using the spectral decomposition of the observables. So the two notions of strategy are equivalent. Note that if $j \in V_i$, then

$$\begin{aligned} \langle \psi | Y_{ij} \otimes X_j | \psi \rangle &= \langle \psi | \sum_{\underline{a} \in \mathcal{S}_i} (-1)^{a_j} M_{\underline{a}}^i \otimes \sum_{b \in \mathbb{Z}_2} (-1)^b N_b^j | \psi \rangle \\ &= \sum_{\underline{a} \in \mathcal{S}_i, b \in \mathbb{Z}_2} (-1)^{a_j+b} p(\underline{a}, b | i, j) \\ &= 2 \left[\sum_{\underline{a}, b : a_j=b} p(\underline{a}, b | i, j) \right] - 1 = 2p_{ij} - 1, \end{aligned} \tag{4}$$

where p_{ij} is, again, the probability that Alice and Bob win on inputs i and j . The quantity $2p_{ij} - 1$ is called the *winning bias* on inputs i and j .

To every linear system, we can also associate a finitely presented group over \mathbb{Z}_2 , as follows.

DEFINITION 17. Let $Ax = b$ be an $m \times n$ linear system. The *solution group* of this system is the group

$$\begin{aligned} \Gamma(A, b) &:= \left\langle x_1, \dots, x_n : x_j^2 = e \text{ for all } 1 \leq j \leq n, \right. \\ &\quad \left. \prod_{j=1}^n x_j^{A_{ij}} = J^{b_i} \text{ for all } 1 \leq i \leq m, \text{ and} \right. \\ &\quad \left. x_j x_k = x_k x_j \text{ if } j, k \in V_i \text{ for some } 1 \leq i \leq m \right\rangle_{\mathbb{Z}_2}. \end{aligned}$$

We say that a group over \mathbb{Z}_2 is a *solution group* if it has a presentation over \mathbb{Z}_2 of this form.

Solution groups and linear system games are related as follows.

THEOREM 18 ([6], see also [5]). *Let \mathcal{G} be the linear system game associated to a system $Ax = b$. Then the following are equivalent:*

- (a) \mathcal{G} has a perfect strategy in C_{qs} .
- (b) \mathcal{G} has a perfect strategy in C_q .
- (c) J_Γ is nontrivial in some finite-dimensional representation of $\Gamma = \Gamma(A, b)$.

Although we have not defined the set of commuting-operator correlations C_{qc} , we can work with C_{qc} through the following result.

THEOREM 19 [5]. *The linear system game associated to a system $Ax = b$ has a perfect strategy in C_{qc} if and only if J_Γ is nontrivial in $\Gamma = \Gamma(A, b)$.*

The main point of this section is to prove an analogue of one direction of Theorem 18 for approximate representations.

PROPOSITION 20. *Let $\Gamma = \Gamma(A, b)$ be a solution group. If J_Γ is nontrivial in finite-dimensional approximate representations of Γ then the linear system game associated to $Ax = b$ has a perfect strategy in C_{qa} .*

The proof of Proposition 20 is a straightforward application of a number of easy stability lemmas. We start by pinning down what we want to prove.

LEMMA 21. *The linear system game associated to $Ax = b$ has a perfect strategy in C_{qa} if and only if, for all $\epsilon > 0$, there is a finite-dimensional quantum strategy (presented in terms of observables) $\{Y_{ij}\}, X_j, |\psi\rangle$ such that*

$$\langle \psi | Y_{ij} \otimes X_j | \psi \rangle \geq 1 - \epsilon \quad \text{for all } 1 \leq i \leq n, j \in V_i.$$

Proof. Since C_{qa} is the closure of C_q , the linear system game associated to $Ax = b$ has a perfect strategy in C_{qa} if and only if, for every $\epsilon > 0$, there is a finite-dimensional quantum strategy such that the winning probability $p_{ij} \geq 1 - \epsilon/2$ for every $1 \leq i \leq m$ and $j \in V_i$. But $p_{ij} \geq 1 - \epsilon/2$ if and only if the winning bias $2p_{ij} - 1 \geq 1 - \epsilon$, so the lemma follows from Equation (4). \square

Next, we come to the stability lemmas, which will allow us to turn approximate representations of the solution group Γ into quantum strategies. The following lemmas are all likely well known to experts (see, for instance, [8, 10]); we include the proofs for completeness.

LEMMA 22. For any diagonal matrix X , there is a diagonal matrix D with $D^2 = \mathbb{1}$ and

$$\|D - X\| \leq \left(1 + \frac{1}{\sqrt{2}}\right) \|X^2 - \mathbb{1}\|.$$

Proof. Suppose X is a $d \times d$ matrix, and let $D_{ii} = \text{sgn Re } X_{ii}$ for all $1 \leq i \leq d$, where $\text{sgn } x = 1$ if $x \geq 0$ and -1 if $x < 0$. To show that the desired inequality holds, consider a complex number $\alpha = a + bi$. Then

$$\begin{aligned} |\alpha^2 - 1|^2 &= |a^2 - b^2 - 1 + 2abi|^2 = [(a^2 - 1) - b^2]^2 + 4a^2b^2 \\ &= (a^2 - 1)^2 + 2b^2 + 2a^2b^2 + b^4. \end{aligned}$$

In particular, this implies that $|\alpha^2 - 1|^2$ is greater than or equal to $(a^2 - 1)^2$ and $2b^2$. Consequently,

$$\begin{aligned} \|(\text{Re } X)^2 - \mathbb{1}\| &= \sqrt{\frac{1}{d} \sum_j [(\text{Re } X_{jj})^2 - 1]^2} \\ &\leq \sqrt{\frac{1}{d} \sum_j |X_{jj}^2 - 1|^2} = \|X^2 - \mathbb{1}\|, \end{aligned}$$

and

$$\begin{aligned} \|\text{Re } X - X\| = \|\text{Im } X\| &= \sqrt{\frac{1}{d} \sum_j |\text{Im } X_{jj}|^2} \leq \sqrt{\frac{1}{2d} \sum_j |X_{jj}^2 - 1|^2} \\ &= \frac{1}{\sqrt{2}} \|X^2 - \mathbb{1}\|. \end{aligned}$$

By considering the cases $a \geq 0$ and $a < 0$ separately, we see that

$$|a^2 - 1| = |1 + a||1 - a| = (1 + |a|)|\text{sgn } a - a| \geq |\text{sgn } a - a|$$

for all $a \in \mathbb{R}$. Thus, as above, $\|D - \text{Re } X\| \leq \|(\text{Re } X)^2 - \mathbb{1}\|$, and the lemma follows. \square

LEMMA 23. Suppose X_1, \dots, X_n are commuting unitary matrices, with $X_i^2 = \mathbb{1}$ for all $1 \leq i \leq n$, and Y is a unitary matrix such that $Y^2 = \mathbb{1}$ and Y commutes

with X_i for all $1 \leq i \leq n - 1$. Then there is a unitary matrix Z such that $Z^2 = \mathbb{1}$, Z commutes with X_i for all $1 \leq i \leq n$, and

$$\|Z - Y\| \leq \left(1 + \frac{1}{2\sqrt{2}}\right) \|X_n Y - Y X_n\|.$$

Proof. Let $Z_0 = \frac{1}{2}(Y + X_n Y X_n)$. Clearly Z_0 commutes with X_i for all $1 \leq i \leq n - 1$. Since $X_n^2 = \mathbb{1}$, we also have that $X_n Z_0 = \frac{1}{2}(X_n Y + Y X_n) = Z_0 X_n$. Since $Y^2 = \mathbb{1} = (X_n Y X_n)^2$ as well, we have that

$$\begin{aligned} \|Z_0^2 - \mathbb{1}\| &= \frac{1}{4} \|Y X_n Y X_n + X_n Y X_n Y - 2\mathbb{1}\| \\ &\leq \frac{1}{4} \|Y X_n Y X_n - \mathbb{1}\| + \frac{1}{4} \|X_n Y X_n Y - \mathbb{1}\| = \frac{1}{2} \|X_n Y - Y X_n\|. \end{aligned}$$

Since X_n and Y are self-adjoint, Z_0 is self-adjoint, so we can simultaneously diagonalize X_1, \dots, X_n and Z_0 . Hence by Lemma 22, there is a matrix Z such that $Z^2 = \mathbb{1}$, Z commutes with X_i for all $1 \leq i \leq n$, and

$$\|Z - Z_0\| \leq \left(1 + \frac{1}{\sqrt{2}}\right) \|Z_0^2 - \mathbb{1}\| \leq \left(\frac{1}{2} + \frac{1}{2\sqrt{2}}\right) \|X_n Y - Y X_n\|.$$

Finally,

$$\|Y - Z_0\| = \frac{1}{2} \|Y - X_n Y X_n\| = \frac{1}{2} \|X_n Y - Y X_n\|,$$

so the lemma follows. □

LEMMA 24. Consider \mathbb{Z}_2^k as a finitely presented group with presentation

$$\langle x_1, \dots, x_k : x_i^2 = e, [x_i, x_j] = e \text{ for all } i \neq j \rangle.$$

Then there is a constant $C > 0$, depending on k , such that if ϕ is an ϵ -representation of \mathbb{Z}_2^k on a Hilbert space H , then there is a representation ψ of \mathbb{Z}_2^k on H with

$$\|\psi(x_i) - \phi(x_i)\| \leq C\epsilon$$

for all $1 \leq i \leq k$.

Proof. Suppose ψ is an ϵ -representation of \mathbb{Z}_2^k such that the following properties hold for some $1 \leq l \leq k - 1$:

- (a) $\psi(x_i)^2 = \mathbb{1}$ for all $1 \leq i \leq k$; and
- (b) $\psi(x_i)$ commutes with $\psi(x_j)$ for all $1 \leq i \leq l - 1$ and $1 \leq j \leq k$.

In particular, property (b) requires that $\psi(x_1), \dots, \psi(x_l)$ pairwise commute. Then by Lemma 23, for each $l < j \leq k$ there is a unitary matrix X_j such that $X_j^2 = \mathbb{1}$, X_j commutes with $\psi(x_i)$ for all $1 \leq i \leq l$, and

$$\|X_j - \psi(x_j)\| \leq C_0 \|\psi(x_l)\psi(x_j) - \psi(x_j)\psi(x_l)\| \leq C_0\epsilon,$$

where $C_0 = 1 + 1/(2\sqrt{2})$. Define an approximate representation ψ' of \mathbb{Z}_2^k by $\psi'(x_i) = \psi(x_i)$ if $i \leq l$ and $\psi'(x_i) = X_i$ if $i > l$. Then $\psi'(x_i)^2 = \mathbb{1}$ for all $1 \leq i \leq k$, and $\psi'(x_i)$ commutes with $\psi'(x_j)$ for all $1 \leq i \leq l$ and $1 \leq j \leq k$. In other words, ψ' satisfies properties (a) and (b) with l replaced by $l + 1$. Finally, $\|\psi'(x_i) - \psi(x_i)\| \leq C_0\epsilon$ for all $1 \leq i \leq k$, so ψ' is a $(4C_0 + 1)\epsilon$ -representation by Lemma 8.

Now suppose that ϕ is any ϵ -representation of \mathbb{Z}_2^k . By Lemma 22, there is an approximate representation ψ_1 of \mathbb{Z}_2^k with $\psi_1(x_i)^2 = \mathbb{1}$ and $\|\psi_1(x_i) - \phi(x_i)\| \leq C_1\epsilon$ for all $1 \leq i \leq k$, where $C_1 = (1 + 1/(\sqrt{2}))$. By Lemma 8, ψ_1 is a $(4C_1 + 1)\epsilon$ -representation. Clearly, ψ_1 satisfies conditions (a) and (b) with $l = 1$. Using the argument in the previous paragraph, we can then iteratively define approximate representations $\psi_2, \dots, \psi_{k-1}$, where ψ_j satisfies conditions (a) and (b) with $l = j$ for all $1 \leq j \leq k - 1$. Let $\epsilon_l = (4C_0 + 1)^{l-1}(4C_1 + 1)\epsilon$, so ψ_1 is an ϵ_1 -representation. It is not hard to check that ψ_l is an ϵ_l -representation, and furthermore that

$$\|\psi_l(x_i) - \psi_1(x_i)\| \leq \frac{1}{4}((4C_0 + 1)^{l-1} - 1)\epsilon_1 = \frac{1}{4}((4C_0 + 1)^{l-1} - 1)(4C_1 + 1)\epsilon$$

for all $1 \leq i \leq k$. Since ψ_{k-1} is an exact representation, we can take

$$C = \frac{1}{4}((4C_0 + 1)^{k-2} - 1)(4C_1 + 1) + C_1. \quad \square$$

LEMMA 25. Suppose $G = \langle S_0 : R_0 \rangle_{\mathbb{Z}_2}$, where R_0 includes the relations $s^2 = e$ for all $s \in S_0$. If J_G is nontrivial in finite-dimensional approximate representations of G , then for every $\epsilon > 0$ there is an ϵ -representation ϕ of G such that $\phi(J) = -\mathbb{1}$, and $\phi(s)^2 = \mathbb{1}$ for all $s \in S_0$.

Proof. Suppose A is an $m \times n$ matrix, and let $S = S_0 \cup \{J\}$. If J is nontrivial in approximate representations, then there is a $\delta > 0$ such that for all $\epsilon > 0$, there is an ϵ -representation ϕ with $\|\phi(J) - \mathbb{1}\| > \delta$.

By Lemmas 8, 22, and 23, there are constants $C, C' > 0$ such that if ϕ is an ϵ -representation, then there is a $C'\epsilon$ -representation ψ such that:

- (1) $\psi(x)^2 = \mathbb{1}$ for all $x \in S$;
- (2) $\psi(s)$ and $\psi(J)$ commute for all $s \in S_0$; and
- (3) $\|\psi(J) - \phi(J)\| \leq C\epsilon$.

We can take $C = (1 + 1/\sqrt{2})$, while C' will depend on the length of the longest defining relation of G . If $\|\phi(J) - \mathbb{1}\| > \delta$, and $\epsilon < \delta/(2C)$, then

$$\delta < \|\phi(J) - \mathbb{1}\| \leq \|\phi(J) - \psi(J)\| + \|\psi(J) - \mathbb{1}\| \leq \frac{\delta}{2} + \|\psi(J) - \mathbb{1}\|,$$

so $\|\psi(J) - \mathbb{1}\| \geq \delta/2$. Thus we conclude that for all $\epsilon > 0$, there is an ϵ -representation ψ satisfying conditions (1) and (2), and with $\|\psi(J) - \mathbb{1}\| > \delta/2$.

Suppose ψ is an ϵ -representation satisfying conditions (1) and (2), and with $\|\psi(J) - \mathbb{1}\| > \delta/2$. Choose a basis with $\psi(J) = \mathbb{1}_{d_0} \oplus (-\mathbb{1}_{d_1})$. Since $\psi(s)$ commutes with $\psi(J)$ for all $s \in S_0$, we must have $\psi = \psi_0 \oplus \psi_1$, where ψ_a is an approximate representation of dimension d_a , and $\psi_a(J) = (-\mathbb{1})^a$, $a = 0, 1$. Since $\psi(s)^2 = \mathbb{1}$, we also have $\psi_a(s)^2 = \mathbb{1}$ for all $s \in S_0$, $a = 0, 1$. To finish the proof, we just need to show that ψ_1 is a $C''\epsilon$ -representation for some constant C'' independent of ψ . If $w \in \mathcal{F}(S)$, then

$$\|\psi(w) - \mathbb{1}\|^2 = \frac{d_0}{d_0 + d_1} \|\psi_0(w) - \mathbb{1}\|^2 + \frac{d_1}{d_0 + d_1} \|\psi_1(w) - \mathbb{1}\|^2.$$

If $w = J$, then $\|\psi_0(w) - \mathbb{1}\| = 0$ and $\|\psi_1(w) - \mathbb{1}\| = \|-2\mathbb{1}\| = 4$, so we conclude that

$$\frac{\delta^2}{4} < \|\psi(J) - \mathbb{1}\|^2 = \frac{4d_1}{d_0 + d_1},$$

so $d_1/(d_0 + d_1) > \delta^2/16$. On the other hand, if $w = r$ is one of the defining relations of G , then

$$\epsilon^2 \geq \|\psi(r) - \mathbb{1}\|^2 \geq \frac{d_1}{d_0 + d_1} \|\psi_1(r) - \mathbb{1}\|^2 > \frac{\delta^2}{16} \|\psi_1(r) - \mathbb{1}\|^2.$$

Thus ψ_1 is a $4\epsilon/\delta$ -representation with $\psi_1(J) = -\mathbb{1}$ and $\psi_1(s)^2 = \mathbb{1}$ for all $s \in S_0$. Since δ is a constant, the lemma follows. □

Proof of Proposition 20. For this proof, we use the notation $O(\epsilon)$ to hide constants which are independent of ϵ , ϕ , and so on. The constants can still depend on the linear system $Ax = b$, however. Suppose J is nontrivial in finite-dimensional approximate representations of Γ . Given $\epsilon > 0$, let ϕ be an ϵ -representation of Γ with $\phi(J) = -\mathbb{1}$ and $\phi(x_j)^2 = \mathbb{1}$ for all $1 \leq j \leq n$, as in Lemma 25. Suppose ϕ has dimension d , and let $|v\rangle$ be the maximally entangled state on $\mathbb{C}^d \otimes \mathbb{C}^d$. For each $1 \leq j \leq n$, set $X_j = \phi(x_j)$. For each $1 \leq i \leq m$, let j_i be the maximal element of V_i , and set $W_i := V_i \setminus \{j_i\}$. The restriction of ϕ to the subgroup $\langle x_j : j \in W_i \rangle$ is an ϵ -representation of $\mathbb{Z}_2^{W_i}$, and by Lemma 24, there is a representation ψ_i of $\mathbb{Z}_2^{W_i}$ with $\|\psi_i(x_j) - \phi(x_j)\| \leq O(\epsilon)$. Set $Y_{ij} := \psi_i(x_j)^T$

(the transpose of $\psi_i(x_j)$ in a Schmidt basis for $|v\rangle$) for all $j \in W_i$, and set $Y_{ij} := (-1)^{b_i} \prod_{j \in W_i} Y_{ij}$.

Suppose $j \in W_i$ for some $1 \leq i \leq m$. Since Y_{ij} and X_j are self-adjoint, we have that

$$2 - \frac{2}{d} \operatorname{tr}(Y_{ij}^T X_j) = \|Y_{ij}^T - X_j\|^2 = \|\psi(x_j) - \phi(x_j)\|^2 \leq O(\epsilon^2),$$

so $(1/d) \operatorname{tr}(Y_{ij}^T X_j) \geq 1 - O(\epsilon^2)$. For the remaining variable in V_i , we have that

$$\begin{aligned} \|Y_{ii}^T - X_{j_i}\| &= \left\| (-1)^{b_i} \prod_{j \in W_i} \psi_i(x_j) - \phi(x_{j_i}) \right\| \\ &\leq \left\| (-1)^{b_i} \prod_{j \in W_i} \phi(x_j) - \phi(x_{j_i}) \right\| + |W_i| \epsilon \\ &= \left\| (-1)^{b_i} \prod_{j \in V_i} \phi(x_j) - \mathbb{1} \right\| + |W_i| \epsilon \leq O(\epsilon), \end{aligned}$$

where the last equality uses the fact that $\phi(x_{j_i})^2 = \mathbb{1}$. Because the Y_{ij} 's commute for all $j \in W_i$, Y_{ii} is also self-adjoint, so once again we conclude that

$$2 - \frac{2}{d} \operatorname{tr}(Y_{ii}^T X_{j_i}) = \|Y_{ii}^T - X_{j_i}\|^2 \leq O(\epsilon^2)$$

or in other words that $(1/d) \operatorname{tr}(Y_{ii}^T X_{j_i}) \geq 1 - O(\epsilon^2)$.

Now clearly $\{Y_{ij}\}, \{X_j\}, |v\rangle$ is a strategy for the linear system game associated to $Ax = b$. If A and B are any two $d \times d$ matrices, it follows from the definition of maximally entangled states that

$$\langle v | A \otimes B | v \rangle = \frac{1}{d} \operatorname{tr}(A^T B).$$

We conclude that $\langle v | Y_{ij} \otimes X_j | v \rangle = (1/d) \operatorname{tr}(Y_{ij}^T X_j) \geq 1 - O(\epsilon^2)$ for all $j \in V_i$, $1 \leq i \leq m$. The proposition follows from Lemma 21. □

4. Linear-plus-conjugacy groups

The goal of the next two sections is to show that there is a solution group Γ such that J_Γ is trivial in finite-dimensional representations, but nontrivial in approximate representations. In this section, we start by showing that it suffices to construct more general types of groups with these properties. Specifically,

in Section 4.1 we show that a certain class of groups, which we call linear-plus-conjugacy groups, fa^* -embeds over \mathbb{Z}_2 in solution groups. In Section 4.2 we show that if we drop the requirement that the embeddings be over \mathbb{Z}_2 , then a more general class of groups, which we call extended homogeneous-linear-plus-conjugacy groups, can also be fa^* -embedded in solution groups.

4.1. Embeddings over \mathbb{Z}_2 . Given an $m \times n$ linear system $Ax = b$, we once again let $V_i = V_i(A) := \{1 \leq j \leq n : A_{ij} \neq 0\}$.

DEFINITION 26. Suppose $Ax = b$ is an $m \times n$ linear system over \mathbb{Z}_2 , and $\mathcal{C} \subseteq [n] \times [n] \times [n]$, where $[n] = \{1, \dots, n\}$. Let

$$\Gamma(A, b, \mathcal{C}) := \langle \Gamma(A, b) : x_i x_j x_i = x_k \text{ for all } (i, j, k) \in \mathcal{C} \rangle_{\mathbb{Z}_2}.$$

Lacking a better term, we say that a group over \mathbb{Z}_2 is a *linear-plus-conjugacy group* if it has a presentation over \mathbb{Z}_2 of this form.

The conjugacy part of the name comes from the fact that since x_i is an involution, the relation $x_i x_j x_i = x_k$ is equivalent to the relation $x_i x_j x_i^{-1} = x_k$, so $\Gamma(A, b, \mathcal{C})$ can be thought of as a solution group with additional conjugacy relations. In the context of linear-plus-conjugacy and related groups, we use the term *conjugacy relations* as a convenient shorthand for relations of the form $xyx = z$. We also use the term *linear relation* $x_1 \cdots x_n = e$ to refer to the set of relations

$$\{x_1 \cdots x_n = e\} \cup \{[x_i, x_j] = e : 1 \leq i \neq j \leq n\}.$$

Finally, observe that there are two ways to make generators x_i and x_j commute in a linear-plus-conjugacy group: we can add a conjugacy relation $x_i x_j x_i = x_j$, or add an additional generator x_{n+1} and a linear relation $x_i x_j x_{n+1} = e$. We pick and choose from these two methods based on what is convenient.

The main point of this section is to prove:

PROPOSITION 27. *Let G be a linear-plus-conjugacy group. Then there is an fa^* -embedding $G \rightarrow \Gamma$ over \mathbb{Z}_2 , where Γ is a solution group.*

We prove Proposition 27 by first showing that linear-plus-conjugacy groups can be embedded in linear-plus-conjugacy groups of a certain form.

DEFINITION 28. A linear-plus-conjugacy group is *nice* if it has a presentation of the form $\Gamma(A, b, \mathcal{C})$, where A is an $m \times n$ matrix over \mathbb{Z}_2 , $b \in \mathbb{Z}_2^m$, and $\mathcal{C} \subseteq [n] \times [n] \times [n]$ is such that if $(i, j, k) \in \mathcal{C}$, then $j, k \in V_l$ for some $1 \leq l \leq m$.

This means that if $x_i x_j x_i = x_k$ is a defining relation of a nice linear-plus-conjugacy group, then $x_j x_k = x_k x_j$ will also be a defining relation.

LEMMA 29. *Let G be a linear-plus-conjugacy group. Then there is an $f a^*$ -embedding $G \rightarrow K$ over \mathbb{Z}_2 , where K is a nice linear-plus-conjugacy group.*

Proof. Suppose $G = \Gamma(A, b, \mathcal{C})$, where A is an $m \times n$ matrix. Let

$$\begin{aligned}
 K := \langle & \Gamma(A, b), w_j, y_j, z_j \text{ for } 1 \leq j \leq n \text{ and } f : \\
 & f^2 = e, y_j^2 = z_j^2 = w_j^2 = e \text{ for all } 1 \leq j \leq n, \\
 & x_j = y_j z_j = f w_j \text{ and } f y_j f = z_j \text{ for all } 1 \leq j \leq n, \\
 & y_j z_k = z_k y_j \text{ for all } (i, j, k) \in \mathcal{C}, \text{ and} \\
 & w_i y_j w_i = z_k \text{ for all } (i, j, k) \in \mathcal{C} \setminus \mathbb{Z}_2.
 \end{aligned}$$

Since the generators are involutions, note that the relations imply that $f w_j = w_j f$, $y_j z_j = z_j y_j$, and $f z_j f = y_j$ for all $1 \leq j \leq n$. If $(i, j, k) \in \mathcal{C}$, then

$$w_i z_j w_i = w_i f y_j f w_i = f w_i y_j w_i f = f z_k f = y_k,$$

so

$$\begin{aligned}
 x_i x_j x_i &= f w_i y_j z_j f w_i = (f w_i y_j w_i f)(f w_i z_j w_i f) \\
 &= (f z_k f)(f y_k f) = y_k z_k = x_k
 \end{aligned}$$

in K . Thus there is a homomorphism $\psi : G \rightarrow K$ sending $x_i \mapsto x_i$.

Suppose ϕ is an ϵ -representation of G , where $\epsilon > 0$. Define an approximate representation γ of K by

$$\begin{aligned}
 \gamma(x_i) &= \begin{pmatrix} \phi(x_i) & 0 \\ 0 & \phi(x_i) \end{pmatrix}, & \gamma(J) &= \begin{pmatrix} \phi(J) & 0 \\ 0 & \phi(J) \end{pmatrix}, \\
 \gamma(y_i) &= \begin{pmatrix} \phi(x_i) & 0 \\ 0 & \mathbb{1} \end{pmatrix}, & \gamma(z_i) &= \begin{pmatrix} \mathbb{1} & 0 \\ 0 & \phi(x_i) \end{pmatrix}, \\
 \gamma(w_i) &= \begin{pmatrix} 0 & \phi(x_i) \\ \phi(x_i) & 0 \end{pmatrix}, & \text{and } \gamma(f) &= \begin{pmatrix} 0 & \mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}.
 \end{aligned}$$

It is straightforward to check that γ is an ϵ -representation of K . If Ψ is the lift of ψ sending $x_i \mapsto x_i$, then $\gamma \circ \Psi = \phi \oplus \phi$. When ϕ is an exact representation of dimension d (possibly infinite), the same construction gives an exact representation γ of dimension $2d$. By Lemma 15, ψ is an $f a^*$ -embedding.

Finally, we observe that K is a nice linear-plus-conjugacy group. Indeed, since the relation $x_i = y_i z_i$ forces y_i and z_i to commute, this relation is equivalent to

the relations

$$x_i y_i z_i = e = [x_i, y_i] = [x_i, z_i] = [y_i, z_i],$$

which means that we can make $x_i = y_i z_i$, and similarly $x_i = f w_i$, part of the ‘linear’ relations. By adding ancilla variables g_{jk} , the commuting relations $y_j z_k = z_k y_j$ can also be replaced with equivalent linear relations $g_{jk} y_j z_k = e$. The conjugacy relations $f y_j f = z_j$ and $w_i y_j w_i = z_k$ will then satisfy the requirements of Definition 28. □

Proof of Proposition 27. By Lemma 29, we can assume that G is a nice linear-plus-conjugacy group. Let $G = \Gamma(A, b, C)$ be a presentation satisfying the conditions of Definition 28. Augment the linear system $Ax = b$ by adding additional variables y_{Ij} for each $I \in \mathcal{C}$ and $1 \leq j \leq 7$, and additional relations

$$\begin{aligned} x_i + y_{i1} + y_{i2} &= 0, & x_j + y_{j2} + y_{j3} &= 0, & y_{l3} + y_{l4} + y_{l5} &= 0 \\ x_i + y_{i5} + y_{i6} &= 0, & x_k + y_{k6} + y_{k7} &= 0, & y_{l1} + y_{l4} + y_{l7} &= 0 \end{aligned}$$

for every $I = (i, j, k) \in \mathcal{C}$. Let Γ be solution group of this augmented linear system, so

$$\Gamma = \langle \Gamma(A, b), y_{Ij} \text{ for } I \in \mathcal{C}, 1 \leq j \leq 7 : R \rangle_{\mathbb{Z}_2},$$

where R consists of the new relations (now written in multiplicative form)

$$x_i y_{i1} y_{i2} = x_j y_{j2} y_{j3} = y_{l3} y_{l4} y_{l5} = x_i y_{i5} y_{i6} = x_k y_{k6} y_{k7} = y_{l1} y_{l4} y_{l7} = e \quad (1)$$

for every $I = (i, j, k) \in \mathcal{C}$, as well as the corresponding commutation relations. In Γ , we have that

$$x_i x_j x_i = (y_{i1} y_{i2}) (y_{i2} y_{i3}) (y_{i5} y_{i6}) = y_{i1} (y_{i3} y_{i5}) y_{i6} = y_{i1} y_{i4} y_{i6} = y_{i7} y_{i6} = x_k$$

for every $I = (i, j, k) \in \mathcal{C}$. So once again we get a homomorphism $\psi : G \rightarrow \Gamma$ sending $x_i \mapsto x_i$.

Suppose ϕ is an ϵ -representation of G . Define an approximate representation γ of Γ by

$$\begin{aligned} \gamma(x_i) &= \begin{pmatrix} \phi(x_i) & 0 \\ 0 & \phi(x_i) \end{pmatrix}, & \gamma(y_{i1}) &= \begin{pmatrix} 0 & \phi(x_i) \\ \phi(x_i) & 0 \end{pmatrix}, \\ \gamma(y_{i2}) &= \begin{pmatrix} 0 & \mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}, & \gamma(y_{i3}) &= \begin{pmatrix} 0 & \phi(x_j) \\ \phi(x_j) & 0 \end{pmatrix}, \\ \gamma(y_{i4}) &= \begin{pmatrix} 0 & \phi(x_j x_i) \\ \phi(x_i x_j) & 0 \end{pmatrix}, & \gamma(y_{i5}) &= \begin{pmatrix} \phi(x_j x_i x_j) & 0 \\ 0 & \phi(x_i) \end{pmatrix}, \\ \gamma(y_{i6}) &= \begin{pmatrix} \phi(x_j x_k) & 0 \\ 0 & \mathbb{1} \end{pmatrix}, & \text{and } \gamma(y_{i7}) &= \begin{pmatrix} \phi(x_j) & 0 \\ 0 & \phi(x_k) \end{pmatrix} \end{aligned}$$

for all $I = (i, j, k) \in \mathcal{C}$. It is straightforward to show that γ is a $C\epsilon$ -representation of Γ , where C is a positive constant ≤ 15 . For instance, consider the relation $y_{I5}^2 = e$. To show that $\gamma(y_{I5})^2 \approx \mathbb{1}$, we need to show that $\phi(x_j x_i x_j)^2 \approx \mathbb{1}$. Write $X \approx_\epsilon Y$ to mean that $\|X - Y\| \leq \epsilon$. Since $\phi(x_i)^2 \approx_\epsilon \mathbb{1}$ and $\phi(x_j)^2 \approx \mathbb{1}$, we have $\phi(x_i x_k x_i)^2 \approx_{3\epsilon} \mathbb{1}$. We can conclude from this that $\gamma(y_{I5})^2 \approx_{3\epsilon} \mathbb{1}$ (we can do slightly better by averaging over the blocks of $\gamma(y_{I5})$, but we ignore this to simplify the analysis). We can similarly show that $\gamma(y_{Ij})^2 \approx_{3\epsilon} \mathbb{1}$ for all $1 \leq j \leq 7$, and that the linear relations in Equation (1) hold to within 3ϵ .

This leaves the commuting relations. Consider the relation $y_{I3}y_{I4}y_{I5} = e$. We want to show that $\gamma(y_{I3})$, $\gamma(y_{I4})$, and $\gamma(y_{I5})$ approximately commute. But since $\gamma(y_{I3})\gamma(y_{I4})\gamma(y_{I5}) \approx_{3\epsilon} \mathbb{1}$ and $\gamma(y_{Ij})^2 \approx_{3\epsilon} \mathbb{1}$, we conclude that

$$\gamma(y_{I4})\gamma(y_{I5}) \approx_{3\epsilon} \gamma(y_{I3})^* \approx_{3\epsilon} \gamma(y_{I3}) \approx_{3\epsilon} \gamma(y_{I5})^* \gamma(y_{I4})^* \approx_{6\epsilon} \gamma(y_{I5})\gamma(y_{I4}),$$

or in other words, $\gamma(y_{I4})\gamma(y_{I5}) \approx_{15\epsilon} \gamma(y_{I5})\gamma(y_{I4})$. The other commuting relations follow similarly.

Let Ψ be the lift of ψ sending $x_i \mapsto x_i$. Then $\gamma \circ \Psi = \phi \oplus \phi$. Once again, the same construction applies when ψ is an exact representation, so ψ is an $f a^*$ -embedding by Lemma 15. □

Note that if $j = k$ in a relation $x_i x_j x_i = x_k$, then the system in Equation (1) is precisely the Mermin–Peres magic square [22, 26]. The magic square has previously been used by Ji to show that linear system games can require a (finite but) arbitrarily high amount of entanglement to play perfectly [11].

The proof of Proposition 27 has several interesting features:

REMARK 30. Let $G = \Gamma(A, b, \mathcal{C})$ be an $m \times n$ linear-plus-conjugacy group, and let $\Gamma' = \Gamma'(A', b')$ be the solution group constructed in the proof of Proposition 27. Then, accounting for Lemma 29, the system $A'x = b'$ has $11n + 8c + 1$ variables and $8n + m + 7c$ equations, where $c = |\mathcal{C}|$ is the number of conjugacy relations. A presentation for Γ' can be constructed in polynomial time in m, n , and c .

The proofs of Lemma 29 and Proposition 27 show that there is a constant $C > 0$, and a lift Ψ of the homomorphism $G \rightarrow \Gamma'$ to the defining free groups, such that for any d -dimensional ϵ -representation ϕ of G , there is a $4d$ -dimensional $C\epsilon$ -representation ψ of Γ' with $\psi \circ \Psi = \phi^{\oplus 4}$. Taking into account the fact that we have to change the presentation of the group K in the proof of Lemma 29, we can take the constant $C \leq 75$. The lift Ψ can be chosen to send the generators of G to generators of Γ' (although not every generator of Γ' will lie in the image of Ψ).

4.2. Embeddings not over \mathbb{Z}_2 . It is important for our argument that the fa^* -embedding in Proposition 27 is over \mathbb{Z}_2 . However, we can go a little further in what type of groups can be embedded if we drop this requirement.

DEFINITION 31. Suppose A is an $m \times n$ matrix over \mathbb{Z}_2 , and $\mathcal{C} \subseteq [n] \times [n] \times [n]$. Let

$$\Gamma_0(A, \mathcal{C}) := \left\langle x_1, \dots, x_n : x_n^2 = e \text{ for all } 1 \leq j \leq n, \right. \\ \left. \prod_{j=1}^n x_j^{A_{ij}} = e \text{ for all } 1 \leq i \leq m, \right. \\ \left. x_j x_k = x_k x_j \text{ if } j, k \in V_i(A) \text{ for some } 1 \leq i \leq m, \text{ and} \right. \\ \left. x_i x_j x_i = x_k \text{ for all } (i, j, k) \in \mathcal{C} \right\rangle.$$

We say that a group G is a *homogeneous-linear-plus-conjugacy group* if it has a presentation of this form.

Since $\Gamma_0(A, \mathcal{C})$ is not presented over \mathbb{Z}_2 , a homogeneous-linear-plus-conjugacy group is *not* a linear-plus-conjugacy group. However, the two types of groups are closely related, as $\Gamma_0(A, \mathcal{C}) \times \mathbb{Z}_2 = \Gamma(A, 0, \mathcal{C})$.

DEFINITION 32. Suppose A is an $m \times n$ matrix over \mathbb{Z}_2 , $\mathcal{C}_0 \subseteq [n] \times [n] \times [n]$, $\mathcal{C}_1 \subseteq [\ell] \times [n] \times [n]$, and L is an $\ell \times \ell$ lower-triangular matrix with nonnegative integer entries. Let

$$E\Gamma_0(A, \mathcal{C}_0, \mathcal{C}_1, L) := \langle \Gamma_0(A, \mathcal{C}_0), y_1, \dots, y_\ell : y_i x_j y_i^{-1} = x_k \text{ for all } (i, j, k) \in \mathcal{C}_1, \\ \text{and } y_i y_j y_i^{-1} = y_j^{L_{ij}} \text{ for all } i > j \text{ with } L_{ij} > 0 \rangle.$$

We refer to the generators x_i in this presentation as *involutory generators*, and to the generators y_j as *noninvolutory generators*. We say that a group G is an *extended homogeneous-linear-plus-conjugacy group* if it has a presentation of this form.

PROPOSITION 33. Let $G = E\Gamma_0(A, \mathcal{C}_0, \mathcal{C}_1, L)$ as in Definition 32, where A is an $m \times n$ matrix. Then there is an $m \times n'$ matrix A' and a set $\mathcal{C}' \subset [n'] \times [n'] \times [n']$, where $n \leq n'$, such that there is an fa^* -embedding $\psi : G \rightarrow \Gamma_0(A', \mathcal{C}')$ with $\psi(x_i) = x_i$ for all $1 \leq i \leq n$.

Proof. Suppose G has ℓ noninvolutory generators, and let

$$G' = \langle G, z, w : z^2 = w^2 = e, y_1 = zw, zy_i = y_i z \text{ for } i = 2, \dots, \ell \rangle.$$

We claim that the natural morphism $\psi : G \rightarrow G'$ is an fa^* -embedding. Indeed, let $\Psi : \mathcal{F}(S) \rightarrow \mathcal{F}(S \cup \{z, w\})$ be the natural inclusion, where $S = \{x_1, \dots, x_n, y_1, \dots, y_\ell\}$. Given an ϵ -representation ϕ of G , define an approximate representation γ of G' by

$$\begin{aligned} \gamma(x_i) &= \begin{pmatrix} \phi(x_i) & 0 \\ 0 & \mathbb{1} \end{pmatrix}, & \gamma(z) &= \begin{pmatrix} 0 & \mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}, \\ \gamma(w) &= \begin{pmatrix} 0 & \phi(y_1)^* \\ \phi(y_1) & 0 \end{pmatrix}, & \gamma(y_1) &= \begin{pmatrix} \phi(y_1) & 0 \\ 0 & \phi(y_1)^* \end{pmatrix}, & \text{and} \\ \gamma(y_i) &= \begin{pmatrix} \phi(y_i) & 0 \\ 0 & \phi(y_i) \end{pmatrix} & \text{for } i = 2, \dots, \ell. \end{aligned}$$

Because L is lower-triangular, G' has no defining relations of the form $y_i y_1 y_i^{-1} = y_i^{L_{ii}}$. Suppose $L_{i1} > 0$, so that $\phi(y_i)\phi(y_1)\phi(y_i)^* \approx_\epsilon \phi(y_1)^{L_{i1}}$, where once again $X \approx_\epsilon Y$ means that $\|X - Y\| \leq \epsilon$. Then $\phi(y_i)\phi(y_1)^*\phi(y_i)^* \approx_\epsilon \phi(y_1)^{-L_{i1}}$, so $\psi(y_i)\psi(y_1)\psi(y_i)^* \approx_\epsilon \psi(y_1)^{L_{i1}}$. It is easy to see that the remaining defining relations of G' hold to within ϵ , so ψ is an ϵ -representation of G' . Since ϕ is a direct summand of $\gamma \circ \Psi$, we can apply Lemma 15 with $N = 2$ and $C = 1$ to see that ψ is an fa -embedding. The same construction for exact representations shows that ψ is an fa^* -embedding.

Next, observe that G' is an extended homogeneous-linear-plus-conjugacy group with $\ell - 1$ noninvolutive generators. Indeed, suppose $(1, j, k) \in \mathcal{C}_1$. Then the defining relation $y_1 x_j y_1^{-1} = x_k$ is equivalent to the relation $z w x_j w z = x_k$. By adding an ancilla variable Z_{jk} with $Z_{jk}^2 = e$, we can replace this relation with the two conjugacy relations $w x_j w = Z_{jk}$ and $z Z_{jk} z = x_k$. Similarly, suppose $L_{i1} > 0$. Then the relation $y_i y_1 y_i = y_i^{L_{i1}}$ is equivalent to the relation $y_i w y_i^{-1} = w (z w)^{L_{i1}-1}$. Once again, we can replace this relation with a sequence of conjugacy relations by adding ancilla variables. For instance, if $L_{i1} = 3$, then we would add ancilla variables W_{i0} and W_{i1} with $W_{i0}^2 = W_{i1}^2 = e$, and conjugacy relations $z w z = W_{i0}$, $w W_{i0} w = W_{i1}$, and $y_i w y_i^{-1} = W_{i1}$. After making these replacements, the only relation containing y_1 is $y_1 = z w$, so we can remove y_1 from the set of generators. The commuting relations added in G' are equivalent to $y_i z y_i^{-1} = z$ for all $2 \leq i \leq \ell$, so G' is an extended homogeneous-linear-plus-conjugacy group. The additional variables (including the ancilla) are involutive generators, so G' has $\ell - 1$ noninvolutive generators.

Iterating this construction, we get a sequence of fa^* -embeddings terminating in a homogeneous-linear-plus-conjugacy group, as desired. \square

The reason the above argument does not apply for groups over \mathbb{Z}_2 is that, if we set $\gamma(J) = \phi(J) \oplus \mathbb{1}$, then $\gamma(J)$ would not commute with $\gamma(z)$ and $\gamma(w)$, while

if we set $\gamma(J) = \phi(J) \oplus \phi(J)$, then any linear relations containing J would not be satisfied.

REMARK 34. The above proof shows that, in Proposition 33, we can take

$$n' = n + 2\ell + \binom{\ell}{2} + |\mathcal{C}_1| + \text{sum}(L)$$

and

$$|\mathcal{C}'| = |\mathcal{C}_0| + 2|\mathcal{C}_1| + 2\binom{\ell}{2} + \text{sum}(L) + \#(L),$$

where ℓ is the number of noninvolutive generators, $\text{sum}(L)$ is the sum of the entries of L , and $\#(L)$ is the number of nonzero entries of L . The matrix A' and set \mathcal{C}' can be constructed in polynomial time in $m, n, \ell, |\mathcal{C}_0|, |\mathcal{C}_1|$, and $\text{sum}(L)$.

Since a homogeneous-linear-plus-conjugacy group can be turned into a linear-plus-conjugacy group by taking a product with \mathbb{Z}_2 , Propositions 27 and 33 imply that every extended homogeneous-linear-plus-conjugacy group can be fa^* -embedded in a solution group. When using these embedding theorems in the next two sections, we start with an extended homogeneous-linear-plus-conjugacy group of interest, and embed it in a homogeneous-linear-plus-conjugacy group using Proposition 33. We then turn this homogeneous-linear-plus-conjugacy group into a linear-plus-conjugacy group by taking the product with \mathbb{Z}_2 and adding relations involving J . Finally, we embed this linear-plus-conjugacy group into a solution group over \mathbb{Z}_2 using Proposition 27.

5. Proof of Theorem 1

The point of this section is to prove the following proposition, and hence finish the proof of Theorem 1.

PROPOSITION 35. *There is a solution group Γ for which J is trivial in finite-dimensional representations, but nontrivial in finite-dimensional approximate representations.*

For the proof of Proposition 35, it is convenient to work with sofic groups. We do not need to know the definition of soficity, just that the class of sofic groups has the following properties:

- (1) Amenable groups are sofic.
- (2) Sofic groups are hyperlinear.

(3) If H is an amenable subgroup of a sofic group G , and $\alpha : H \rightarrow G$ is injective homomorphism, then the HNN extension

$$\langle G, t : tht^{-1} = \alpha(h) \rangle \quad (\text{where } t \text{ is a new indeterminate})$$

of G by α is sofic.

An expository treatment of sofic groups can be found in [4]. In particular, the last ‘closure property’ can be found in [4, Section II.4].

We need one more general-purpose lemma before proceeding to the proof.

LEMMA 36. Suppose $G = \langle S : R \rangle$ is a finitely presented group, where R contains the relation $a^2 = e$ for some $a \in S$. Let

$$\widehat{G} := \langle G, t : t^2 = e, tat = Ja \rangle_{\mathbb{Z}_2},$$

where $J, t \notin S$. If a is nontrivial in approximate representations of G , then J is nontrivial in approximate representations of \widehat{G} .

Note that \widehat{G} is the ‘ \mathbb{Z}_2 -HNN extension’ of $G \times \mathbb{Z}_2$, where J is the generator of the \mathbb{Z}_2 factor, by the order-two automorphism sending $a \mapsto Ja$ and $J \mapsto J$.

Proof. Recall that if X is a linear operator on a finite-dimensional Hilbert space H , then $\widetilde{\text{tr}}(X) := \text{tr}(X)/\dim H$. Suppose ϕ is an ϵ -representation of G with $\phi(a)^2 = 1$ and $\text{tr}(\phi(a)) \geq 0$ (we explain how to fulfil these hypotheses shortly). Because the eigenvalues of $\phi(a)$ belong to $\{\pm 1\}$, we can choose a basis so that $\phi(a) = \mathbb{1}_{d_0} \oplus (-\mathbb{1})_{d_0} \oplus \mathbb{1}_{d_1}$, where $d_1 = \text{tr}(\phi(a))$. Define an approximate representation ψ of \widehat{G} by

$$\psi(x) = \phi(x) \quad \text{for all } x \in S, \psi(J) = -\mathbb{1}, \quad \text{and} \quad \psi(t) = \begin{pmatrix} 0 & \mathbb{1}_{d_0} & 0 \\ \mathbb{1}_{d_0} & 0 & 0 \\ 0 & 0 & \mathbb{1}_{d_1} \end{pmatrix}.$$

Clearly $\|\psi(r) - \mathbb{1}\| = \|\phi(r) - \mathbb{1}\| \leq \epsilon$ for all relations $r \in R$, $\psi([J, s]) = \mathbb{1}$ for all $s \in S \cup \{t\}$, and $\psi(t)^2 = \psi(J)^2 = \mathbb{1}$. For the remaining relation,

$$\|\psi(tat) - \psi(Ja)\| = \|\mathbb{0}_{2d_0} \oplus 2\mathbb{1}_{d_1}\| = 2\sqrt{\frac{d_1}{2d_0 + d_1}} = 2\sqrt{\widetilde{\text{tr}}(\phi(a))}.$$

So ψ will be a $\max(\epsilon, 2\sqrt{\widetilde{\text{tr}}(\phi(a))})$ -representation with $\|\psi(J) - \mathbb{1}\| = 2$.

Suppose a is nontrivial in approximate representations of G , and fix $\tau > 0$. By Lemma 12, for every $\epsilon > 0$ there is an ϵ -representations γ of G with

$0 \leq \widetilde{\text{tr}}(\gamma(a)) \leq \tau$. We want to show that it is possible to find ϵ -representations γ of this form with $\gamma(a)^2 = \mathbb{1}$. In the proof of Lemma 12, the approximate representations γ can be constructed from any family of ϵ -representations ϕ_ϵ , $\epsilon > 0$, such that $\|\phi_\epsilon(a) - \mathbb{1}\| \geq \delta$ for all $\epsilon > 0$, where $\delta > 0$ is some fixed constant. By Lemmas 8 and 22, it is possible to find such a family with $\phi_\epsilon(a)^2 = \mathbb{1}$ for all $\epsilon > 0$. The approximate representations γ are then constructed by taking tensor powers of direct sums of the approximate representations $\phi_\epsilon, \overline{\phi_\epsilon}$, and copies of the trivial representation, and will also satisfy $\gamma(a)^2 = \mathbb{1}$.

Taking $\tau = \epsilon^2/4$, the argument of the first paragraph gives ϵ -representations ψ of \widehat{G} with $\|\psi(J) - \mathbb{1}\| = 2$, so J is nontrivial in approximate representations of \widehat{G} as desired. \square

We are now ready to prove Proposition 35. Note that any hyperlinear but nonresidually finite group has an element which is trivial in finite-dimensional representations, but nontrivial in approximate representations. To prove Proposition 35, we show that

$$K = \langle x, y, a, b : a^2 = b^2 = e, ab = ba, yay^{-1} = a, yby^{-1} = ab, xyx^{-1} = y^2 \rangle$$

is an extended homogeneous-linear-plus-conjugacy group which is hyperlinear but nonresidually finite. Indeed, to see that K has a presentation as in Definition 32, we can introduce a third variable c with $c^2 = e$ and $c = ab$. Then K is equivalent to the extended homogeneous-linear-plus-conjugacy group with three involutory generators a, b, c , one linear relation $abc = e$ (along with the corresponding commuting relations), two noninvolutory generators x and y , and three conjugacy relations $yay^{-1} = a$, $yby^{-1} = c$, and $xyx^{-1} = y^2$. For the remainder of this section, K will refer to this group.

LEMMA 37. *K is sofic, and the element $a \in K$ is nontrivial.*

Proof. $K_1 := \langle y, a, b : a^2 = b^2 = e, ab = ba, yay^{-1} = a, yby^{-1} = ab \rangle$ is isomorphic to a semidirect product $\mathbb{Z} \ltimes (\mathbb{Z}_2 \times \mathbb{Z}_2)$, and in particular is solvable (hence amenable). The group K is the HNN extension of K_1 by the injective endomorphism of $\langle y \rangle \cong \mathbb{Z}$ sending $y \mapsto y^2$. Hence K is sofic by properties (1) and (3) of sofic groups above. In addition, the natural morphism $K_1 \rightarrow K$ is injective. Since a is clearly nontrivial in K_1 , we conclude that a is nontrivial in K . \square

The following lemma comes from discussions with Tobias Fritz.

LEMMA 38. *The element $a \in K$ is trivial in all finite-dimensional representations of K .*

Proof. By a theorem of Mal'cev [20], it suffices to show that a is trivial in finite representations, rather than finite-dimensional representations. So let $\phi : G \rightarrow H$ be a homomorphism from G to a finite group H . Now the order k of $\phi(x)$ is finite, so $\phi(y) = \phi(x)^k \phi(y) \phi(x)^{-k} = \phi(y)^{2^k}$. It follows that the order $m = |\phi(y)|$ of $\phi(y)$ divides $2^k - 1$, and in particular is odd. Since $\phi(y)\phi(b)\phi(y)^{-1} = \phi(ab)$ and $\phi(y)\phi(ab)\phi(y)^{-1} = \phi(b)$, we conclude that $\phi(b) = \phi(y)^m \phi(b) \phi(y)^{-m} = \phi(ab)$. Consequently $\phi(a) = 1$ as desired. \square

Proof of Proposition 35. By Proposition 33, there is an fa -embedding of K to a homogeneous-linear-plus-conjugacy group $G = \Gamma_0(A, C)$, in which $a \in K$ is mapped to a generator x_i of G . Let

$$\widehat{G} = \langle G, t : t^2 = e, tx_it = Jx_i \rangle_{\mathbb{Z}_2}.$$

The relation $tx_it = Jx_i$ can be replaced with the relations $tx_it = Z$ and $Zx_i = J$, where Z is an ancilla variable with $Z^2 = e$. With this presentation, \widehat{G} is a linear-plus-conjugacy group. By Proposition 27, there is an fa -embedding over \mathbb{Z}_2 of \widehat{G} to a solution group Γ .

By Lemma 37, a is nontrivial in approximate representations of K , and hence x_i is nontrivial in approximate representations of G . By Lemma 36, $J_{\widehat{G}}$ is nontrivial in approximate representations of \widehat{G} , and we conclude that J_{Γ} is nontrivial in approximate representations of Γ .

Finally, there is a morphism from K to \widehat{G} which sends a to x_i , so x_i will be trivial in all finite-dimensional representations of \widehat{G} by Lemma 38. But since $J_{\widehat{G}} = [t, x_i]$, this means that $J_{\widehat{G}}$ (and hence J_{Γ}) is trivial in all finite-dimensional representations of \widehat{G} . \square

Proof of Theorem 1. Let Γ be the solution group from Proposition 35, and let \mathcal{G} be the associated game. Since J is trivial in finite-dimensional representations, Theorem 18 implies that \mathcal{G} does not have a perfect strategy in C_{qs} . But since J is nontrivial in approximate representations, Proposition 20 implies that \mathcal{G} has a perfect strategy in C_{qa} . \square

REMARK 39. By Remarks 30 and 34, the linear system constructed in the proof of Theorem 1 will have 235 variables and 184 equations.

6. Proofs of Theorems 2 and 3

To prove Theorem 2, we want to find a hyperlinear group with an undecidable word problem, which fa -embeds in a solution group. For Theorem 3, we want to

find a family of residually finite groups with arbitrarily hard (albeit computable) word problems, which *fin*-embed in solution groups. Fortunately, such groups are provided by Kharlampovich [14] and Kharlampovich, Myasnikov, and Sapir [15]. Since the presentations are rather complicated, we do not repeat them here. Instead, we summarize some points of the construction from [15] in the following theorem.

It is helpful to use the following notation: given $S_0 \subseteq S_1$, let $\mathcal{N}(S_0, S_1)$ denote the normal subgroup generated by S_0 in the free group $\mathcal{F}(S_1)$. Note that if $S_1 \subseteq S$, then $\mathcal{N}(S_0, S_1)$ is a (not necessarily normal) subgroup of $\mathcal{F}(S)$ in a natural way. Also, if x, y are group elements, recall that $[x, y] = xyx^{-1}y^{-1}$, and $x^y = yxy^{-1}$. (This is the reverse of the convention in [15], where $[x, y] = x^{-1}y^{-1}xy$ and $x^y = y^{-1}xy$.) Finally, recall that a set $X \subseteq \mathbb{N}$ is said to be *recursively enumerable* if there is a Turing machine which takes integers n as input, and halts with output true if and only if $n \in X$ (if $n \notin X$ then the machine can either halt with output false, or run forever).

THEOREM 40 ([15], see also [14]). *Let $X \subseteq \mathbb{N}$ be recursively enumerable. Then there is a finitely presented solvable group $K_X = \langle S : R \rangle$ with the following properties:*

- (1) *The set S is divided into three subsets $L_i, i = 0, 1, 2$.*
- (2) *The relations in R come in three types:*
 - (a) *R contains the relations $x^2 = e$ for all $x \in L_0 \cup L_1$.*
 - (b) *R also contains commuting relations of the form $xy = yx$, for certain pairs $x, y \in S$.*
 - (c) *For every other relation $r \in R$, there are some subsets $S_1 \subseteq S$ and $S_0 \subseteq (L_0 \cup L_1) \cap S_1$ such that $r \in \mathcal{N}(S_0, S_1)$, and the image of $\mathcal{N}(S_0, S_1)$ in K_X is abelian.*
- (3) *The image of $\mathcal{N}(L_0, S)$ in K_X is abelian.*
- (4) *There are elements $z_0, z_1 \in L_0, A_1, A_2 \in L_1$, and $a, a' \in L_2$, such that $n \in X$ if and only if*

$$[A_2, [A_1, w(2^n)]] = [A_2, [A_1, z_0]]$$

in K_X , where $w(m)$ is defined by

$$w(m) := \begin{cases} z_1 & m = 0, \\ w(m-1)w(m-1)^{a^{-1}}w(m-1)^aw(m-1)^{a'} & m \geq 1. \end{cases}$$

- (5) *If X is recursive, then K_X is residually finite.*

Note that there is some overlap between relations of type (2b) and (2c). Indeed, if $[x, y] = e$ is a relation, then the image of $\mathcal{N}(\{x\}, \{x, y\})$ in K_X is equal to $\langle x \rangle$, and in particular is abelian. Since $[x, y]$ belongs to $\mathcal{N}(\{x\}, \{x, y\})$, any relation $[x, y] = e$ of type (2b) with $x \in L_0 \cup L_1$ can also be regarded as a relation of type (2c).

Proof. For the convenience of the reader, we explain how to recover Theorem 40 from [15]. First, if X is recursively enumerable, then by [15, Theorem 2.7] there is a 2-glass deterministic Minsky machine MM_2 enumerating X , in the sense that $n \in X$ if and only if the machine takes input configuration $(1; 2^n, 0)$ to the accept configuration $(0; 0, 0)$ (this and other unexplained terminology and notation is as in [15]). Furthermore, if X is recursive, then we can take MM_2 to be a sym-universally halting 2-glass Minsky machine deciding X . We let K_X be the finitely presented group $G(MM_k)$ defined in relations (G1)–(G8) of [15, Section 4.1], with parameter $p = 2$. Then part (1) of the theorem is exactly the notation used in [15], part (3) follows immediately from [15, Lemma 4.5], and part (5) is [15, Theorem 4.18].

It remains to show that K_X satisfies parts (2) and (4) of the theorem. The relations in part (2a) follow from the relations in (G1). For the rest of part (2), we need to show that every other relation in (G1)–(G8) falls under one of (2b) or (2c). It is immediate that the relations in (G1) (excepting the relations from (2a)) and the relations in (G2) fall under (2b). The relations in (G3) and (G4) are written down in parts (1)–(3) of [15, Lemma 4.1]. The relations in part (1) of this lemma are just commuting relations, and hence fall under (2b). Using the notation of this lemma, if we set $S_0 = X$ and $S_1 = X \cup F \cup F'$, then the relations in parts (2) and (3) of Lemma 4.1 belong to $\mathcal{N}(S_0, S_1)$. The fact that $\mathcal{N}(S_0, S_1)$ is abelian is the conclusion of Lemma 4.1. The specific set X and sets F, F' used in (G3) and (G4) are contained in L_1 and L_2 , respectively, so these relations fall under (2b) and (2c). Finally, the relations in (G5)–(G8) belong to $\mathcal{N}(L_0, S)$, and hence fall under (2c) by part (3) of the theorem.

For part (4), note that the element $w(m)$ belongs to $\mathcal{N}(L_0, S)$, which is abelian. Since $z_1^2 = e$ by part (2a), we can conclude by induction that $w(m)^2 = e$ for all $m \geq 0$. Using the notation of [15, Theorem 4.3(b)], let $z_0 = x(q_0 A_0)$, $z_1 = x(q_1 A_1)$, $a = a_1$, $a' = a'_1$, and A_1, A_2 be the symbols of the same name. Using again that $\mathcal{N}(L_0, S)$ is abelian, it follows that $w(m)$ is the element denoted in [15] by $x(q_1 A_0) * a_1^{(m)}$, that $[A_2, [A_1, w(m)]]$ is the element $x(q_1 A_0) * a_1^{(m)} * A_1 * A_2$, and that $[A_2, [A_1, z_0]]$ is the element $x(q_0 A_0) * A_1 * A_2$. Thus [15, Theorem 4.3(b) and Section 3.1] imply that $[A_2, [A_1, w(m)]] = [A_2, [A_1, z_0]]$ if and only if MM_2 takes input configuration $(1; m, 0)$ to accept configuration $(0; 0, 0)$. Hence part (4) of the theorem follows from the definition of MM_2 . \square

We show that every group with a presentation as in parts (1) and (2) of Theorem 40 is an extended homogeneous-linear-plus-conjugacy group (as in Definition 32). We first illustrate this with an example.

EXAMPLE 41. Let

$$K = \langle x_1, x_2, y_1, y_2 : x_1^2 = x_2^2 = [x_1, x_2] = [x_2, y_1] = [y_1, y_2] = x_1x_2x_1^{y_1}x_1^{y_1^{-1}} = e \rangle.$$

This presentation of K satisfies parts (1) and (2) of Theorem 40 with $L_0 \cup L_1 = \{x_1, x_2\}$ and $L_2 = \{y_1, y_2\}$. Indeed, all the defining relations fall under (2a) and (2b) except the last, which belongs to $\mathcal{N}(S_0, S_1)$ with $S_0 = \{x_1, x_2\}$, $S_1 = \{x_1, x_2, y_1\}$. Since $x_1^{y_1}, x_1^{y_1^{-1}}$, and x_1x_2 are all involutions in K , the last relation implies that $x_1^{y_1}, x_1^{y_1^{-1}}$, and x_1x_2 commute in K . But x_2 commutes with both x_1 and y_1 , and hence commutes with $x_1^{y_1}$ and $x_1^{y_1^{-1}}$, so x_1 also commutes with $x_1^{y_1}$ and $x_1^{y_1^{-1}}$. It follows that the image of $\mathcal{N}(S_0, S_1)$ is abelian in K , and hence the last defining relation falls under (2c).

To construct a presentation of K as an extended homogeneous-linear-plus-conjugacy group, note that the generators split into involutory generators x_1, x_2 and noninvolutory generators y_1, y_2 . The order in which we list the involutory generators does not matter, but for the noninvolutory generators the order is significant. Most of the defining relations for K are commuting relations, and these can be replaced with the conjugacy relations $x_1x_2x_1 = x_2, y_1x_2y_1^{-1} = x_2$, and $y_2y_1y_2^{-1} = y_1$. Note that we could choose $x_2x_1x_2 = x_1$ in place of $x_1x_2x_1 = x_2$, but that we are forced to pick $y_2y_1y_2^{-1} = y_1$ over $y_1y_2y_1^{-1} = y_2$ by the ordering on y_1 and y_2 . For the last defining relation, add two new involutory generators x_3 and x_4 , along with relations $y_1x_1y_1^{-1} = x_3$ and $y_1x_4y_1^{-1} = x_1$. The relation $x_1x_2x_1^{y_1}x_1^{y_1^{-1}} = e$ can then be replaced with the linear relation $x_1x_2x_3x_4 = e$. Note that, following the convention introduced in Section 4.1, this last linear relation is really the set of relations $\{x_1x_2x_3x_4 = e\} \cup \{[x_i, x_j] = e \text{ for } i \neq j\}$. However, the additional relations $[x_i, x_j] = e$ do not change the group, since $x_1^{y_1}, x_1^{y_1^{-1}}, x_1$, and x_2 commute in K .

We conclude that if $A = (1 \ 1 \ 1 \ 1)$, $\mathcal{C}_0 = \{(1, 2, 2)\}$, $\mathcal{C}_1 = \{(1, 2, 2), (1, 1, 3), (1, 4, 1)\}$, and $L = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, then the natural map

$$K \rightarrow E\Gamma_0(A, \mathcal{C}_0, \mathcal{C}_1, L) : x_i \mapsto x_i, i = 1, 2 \quad \text{and} \quad y_i \mapsto y_i, i = 1, 2$$

is an isomorphism. Note that the relation $x_1x_2x_1 = x_2$ coming from including $(1, 2, 2)$ in \mathcal{C}_0 is actually redundant, since this relation is also implied by the linear relation, but we include it to illustrate how to handle this type of commuting relation in general.

We now turn to the general case.

LEMMA 42. *Suppose $K = \langle S : R \rangle$ is a finitely presented group satisfying properties (1) and (2) of Theorem 40. Then K is an extended homogeneous-linear-plus-conjugacy group. Furthermore, if $S_0 \subseteq S_1 \subseteq S$ are two subsets such that $S_0 \subseteq L_0 \cup L_1$, and the image of $\mathcal{N}(S_0, S_1)$ in K is abelian, then for every $w \in \mathcal{N}(S_0, S_1)$, there is a presentation of K as an extended homogeneous-linear-plus-conjugacy group in which w is equal in K to one of the involutory generators x_j .*

Proof. The proof is exactly as in Example 41. The generators S of K split into involutory generators $L_0 \cup L_1 = \{x_1, \dots, x_{n_0}\}$ and noninvolutory generators $L_2 = \{y_1, \dots, y_\ell\}$, where we choose an arbitrary order on each set (although only the order on L_2 is significant). We show that there are $m \geq 0, n \geq n_0$, an $m \times n$ matrix A over \mathbb{Z}_2 , subsets $\mathcal{C}_0 \subseteq [n] \times [n] \times [n]$ and $\mathcal{C}_1 \subseteq [\ell] \times [n] \times [n]$, and an $\ell \times \ell$ lower-triangular matrix L , such that the natural inclusion

$$K \rightarrow E\Gamma_0(A, \mathcal{C}_0, \mathcal{C}_1, L) : \begin{aligned} x_i &\mapsto x_i & \text{for } 1 \leq i \leq n_0 & \text{ and} \\ y_i &\mapsto y_i & \text{for } 1 \leq i \leq \ell \end{aligned}$$

is an isomorphism which sends the element w to some x_j . Indeed, in property (2) of Theorem 40, the defining relations for K split into three types, type (2a), (2b), and (2c). The relations of type (2a) simply say that the generators $L_0 \cup L_1$ are involutions, and the commuting relations in type (2b) are equivalent to conjugacy relations (where, for relations $y_i y_j = y_j y_i$, we choose either $y_i y_j y_i^{-1} = y_j$ or $y_j y_i y_j^{-1} = y_i$ depending on whether $i > j$ or $i < j$). Thus the only difficulty in constructing the presentation $E\Gamma_0(A, \mathcal{C}_0, \mathcal{C}_1, L)$ is handling the relations of type (2c). For these relations, we prove the following claim: if $S_0 \subseteq S_1 \subseteq S$ are subsets such that $S_0 \subseteq L_0 \cup L_1$, and the image of $\mathcal{N}(S_0, S_1)$ in K is abelian, then there is a set of indeterminates $S_w = \{x_{n_0+1}, \dots, x_q\}$, $q \geq n_0$, and a set of relations R_w , such that:

- (i) R_w consists of linear relations $x_{i_1} \cdots x_{i_r} = e$, $1 \leq i_1 < \dots < i_r \leq q$ (which, as in Section 4.1, include the commuting relations $[x_{i_j}, x_{i_k}] = e$ for all $1 \leq j, k \leq r$), conjugacy relations $x_i x_j x_i = x_k$, $1 \leq i, j, k \leq q$, and conjugacy relations $y_i x_j y_i^{-1} = x_k$, $1 \leq i \leq \ell$ and $1 \leq j, k \leq q$;
- (ii) the relations

$$\tilde{R}_w := R_w \cup \{s^2 = e : s \in L_0 \cup L_1 \cup S_w\}$$

imply that w is equal to an element of $S_0 \cup S_w$; and

(iii) the added generators S_w and relations R_w do not change the group, that is the inclusion

$$K \rightarrow \langle K, S_w : R_w \cup \{s^2 = e : s \in S_w\} \rangle$$

is an isomorphism.

To prove the claim, we use induction on the length of w in $\mathcal{F}(S_1)$. The claim is trivially true if $w \in S_0 \cup S_0^{-1}$. Suppose $w = zw'z^{-1}$, where $w' \in \mathcal{N}(S_0, S_1)$ has length less than w , and $z \in S_1$. By induction, there is a set of ancilla variables $S_{w'} = \{x_{n_0+1}, \dots, x_{q'}\}$ and relations $R_{w'}$ satisfying properties (i)–(iii) for w' . In particular, the relations $\tilde{R}_{w'}$ imply that w' is equal to some $X \in S_0 \cup S_{w'}$. Then we can set $q := q' + 1$, $S_w := S_{w'} \cup \{x_q\}$, and $R_w := R_{w'} \cup \{x_q = zXz\}$ or $R_{w'} \cup \{x_q = zXz^{-1}\}$ depending on whether $z \in L_0 \cup L_1$ or $z \in L_2$. If $w = z^{-1}w'z$, then we do the same thing, but using $zx_qz^{-1} = X$ in place of $x_q = zXz^{-1}$. In both cases, the relations \tilde{R}_w imply that w is equal to x_q , so properties (i)–(iii) hold for R_w . Similarly, suppose that $w = w_1 \cdots w_k$, where each $w_i \in \mathcal{N}(S_0, S_1)$ has smaller length than w . By induction, there are sets $S_{w_i} = \{x_{q_{i-1}+1}, \dots, x_{q_i}\}$, where $n_0 = q_0 \leq q_1 \leq \dots \leq q_k$, and relations \tilde{R}_{w_i} implying that w_i is equal to some $X_i \in S_0 \cup S_{w_i}$. We then set $q := q_k + 1$, $S_w := \bigcup S_{w_i} \cup \{x_q\}$, and

$$R_w := \bigcup R_{w_i} \cup \{x_q X_1 \cdots X_k = e = [x_q, X_i] = [X_i, X_j] \text{ for all } 1 \leq i, j \leq k\}$$

(in other words R_w contains all relations from R_{w_i} and the single linear relation $x_q X_1 \cdots X_k = e$). Since the image of $\mathcal{N}(S_0, S_1)$ in K is abelian, adding the relations R_w does not change K , so properties (i)–(iii) hold again, finishing the proof of the claim.

Now suppose that K has a defining relation r of type (2c), so $r \in \mathcal{N}(S_0, S_1)$ for some $S_0 \subseteq L_0 \cup L_1$ and $S_1 \subseteq S$ such that the image of $\mathcal{N}(S_0, S_1)$ is abelian in K . If $r = zr'z^{-1}$ for some $r' \in \mathcal{N}(S_0, S_1)$ and $z \in S_1 \cup S_1^{-1}$, then r can be replaced with the simpler relation r' . Hence we can assume without loss of generality that $r = r_1 \cdots r_n$, where each $r_i \in \mathcal{N}(S_0, S_1)$. By the claim, we can add ancilla variables and relations as in property (i) such that each r_i is equal to an involutory generator X_i in K , and the relation r can be replaced with the linear relation $X_1 \cdots X_n = e$. By repeatedly applying the claim to all relations of type (2c), we can construct a presentation of K as an extended homogeneous-linear-plus-conjugacy group. Furthermore, the claim immediately implies that we can construct such a presentation sending a specified element w to some involutory generator, as required. \square

We now come to the main result of this section.

PROPOSITION 43. *Let $X \subseteq \mathbb{N}$ be a recursively enumerable set. Then there is a family of solution groups $\Gamma_n = \Gamma(A^{(n)}, b^{(n)})$, $n \geq 1$, such that:*

- (a) $A^{(n)}x = b^{(n)}$ is an $\exp(O(n)) \times \exp(O(n))$ linear system;
- (b) the function $n \mapsto (A^{(n)}, b^{(n)})$ is computable in $\exp(O(n))$ -time;
- (c) J_{Γ_n} is nontrivial in Γ_n if and only if $n \in X$;
- (d) if J_{Γ_n} is nontrivial in Γ_n , then J_{Γ_n} is nontrivial in approximate representations; and
- (e) if X is recursive and J_{Γ} is nontrivial in Γ_n , then J_{Γ_n} is nontrivial in finite-dimensional representations.

Before giving the proof, we need the following exact version of Lemma 36.

LEMMA 44. Suppose $G = \langle S : R \rangle$ is a finitely presented group, where R contains the relation $a^2 = e$ for some $a \in S$. Let

$$\widehat{G} := \langle G, t : t^2 = e, tat = Ja \rangle_{\mathbb{Z}_2},$$

where $J, t \notin S$. If a is nontrivial in finite-dimensional representations of G , then J is nontrivial in finite-dimensional representations of \widehat{G} .

Proof. Suppose a is nontrivial in finite-dimensional representations of G . A theorem of Baumslag states that the free product of two residually finite groups amalgamated over a finite subgroup is residually finite [1]. Let $\widetilde{G} := G \times \mathbb{Z}_2$, where the generator of the \mathbb{Z}_2 factor is denoted by J , and let

$$H = \langle t, a : t^2 = a^2 = e, tat = aJ \rangle_{\mathbb{Z}_2} \cong \mathbb{Z}_2 \rtimes \mathbb{Z}_2 \times \mathbb{Z}_2.$$

Then \widehat{G} is isomorphic to the amalgamated free product of \widetilde{G} and H over $\langle a, J \rangle \cong \mathbb{Z}_2 \times \mathbb{Z}_2$, a finite group. While \widetilde{G} is not necessarily residually finite, the group $\widetilde{G}^{\text{fin}}$ is residually finite by definition, and there is natural map from \widehat{G} to the amalgamated free product of $\widetilde{G}^{\text{fin}}$ and H over $\mathbb{Z}_2 \times \mathbb{Z}_2$. The image of $J_{\widehat{G}}$ is nontrivial in $\widetilde{G}^{\text{fin}}$, and hence in the amalgamated product of $\widetilde{G}^{\text{fin}}$ and H . So J is nontrivial in finite-dimensional representations of \widehat{G} by Baumslag’s result. \square

Proof of Proposition 43. Given a recursively enumerable subset $X \subseteq \mathbb{N}$, let $K_X = \langle S : R \rangle$ be the associated group from Theorem 40. Using the notation from property (4) of Theorem 40, let $c(n) = [A_2, [A_1, w(2^n)]] [A_2, [A_1, z_0]]^{-1}$, so that $c(n) = e$ in K_X if and only if $n \in X$. Since $c(n)$ belongs to $\mathcal{N}(L_0, S)$, Lemma 42 and property (3) of Theorem 40 implies that K_X has a presentation $ET_0(A^{(n)}, \mathcal{C}_0^{(n)}, \mathcal{C}_1^{(n)}, L^{(n)})$ as an extended homogeneous-linear-plus-conjugacy group, in which $c(n)$ is equal to some involutory generator x_i . For the purposes of this

proof, define the *size* of a presentation $E\Gamma_0(A, C_0, C_1, L)$ to be the maximum of the dimensions of A , the sizes $|C_0|$ and $|C_1|$, and the sum $\sum_{i,j} L_{ij}$ of the entries of L . Since the presentation $\langle S : R \rangle$ is fixed, the size of $E\Gamma_0(A^{(n)}, C_0^{(n)}, C_1^{(n)}, L^{(n)})$ depends only on the number of ancilla generators and linear and conjugacy relations needed to set $c(n)$ equal to one of the involutory generators. Inspection of the argument from Lemma 42 reveals that we need to add $4m$ ancilla generators and the same number of linear and conjugacy relations to set $w(m)$ to an involutory generator. Thus $E\Gamma_0(A^{(n)}, C_0^{(n)}, C_1^{(n)}, L^{(n)})$ will have size $O(2^n)$, and the function $n \mapsto (A^{(n)}, C_0^{(n)}, C_1^{(n)}, L^{(n)})$ can be computed in $O(2^n)$ -time.

By Proposition 33, there is an fa^* -embedding from $E\Gamma_0(A^{(n)}, C_0^{(n)}, C_1^{(n)}, L^{(n)})$ to a homogeneous-linear-plus-conjugacy group G_n , in which $c(n)$ is mapped to some generator x_i . As in the proof of Proposition 35, let

$$\widehat{G}_n = \langle G_n, t : t^2 = e, tx_it = Jx_i \rangle_{\mathbb{Z}_2}.$$

Then \widehat{G}_n is a linear-plus-conjugacy group, and by Proposition 27, there is an fa^* -embedding of \widehat{G}_n in a solution group $\Gamma_n = \Gamma(A^{(n)}, b^{(n)})$. By Remarks 30 and 34, $A^{(n)}$ and $b^{(n)}$ can be constructed in time polynomial in the size of $E\Gamma_0(A^{(n)}, C_0^{(n)}, C_1^{(n)}, L^{(n)})$, so $A^{(n)}$ and $b^{(n)}$ satisfy parts (a) and (b) of the proposition.

Suppose $c(n)$ is nontrivial. Since K_X is solvable, it is hyperlinear, so $c(n)$ is nontrivial in approximate representations. By Lemma 36, J_{Γ_n} will be nontrivial in approximate representations. If X is recursive, then K_X will be residually finite by property (5) of Theorem 40, and hence J_{Γ_n} will be nontrivial in finite-dimensional representations by Lemma 44 (this uses the fact that fa^* -embeddings are also *fin*-embeddings). On the other hand, if $c(n)$ is trivial then J_{Γ_n} will be trivial. Hence parts (c)–(e) of the proposition follow from property (4) of Theorem 40. □

Proof of Theorem 2. Let $X \subseteq \mathbb{N}$ be a recursively enumerable but nonrecursive set, and take the family $\{G_n : n \in \mathbb{N}\}$ of games associated to the solution groups $\{\Gamma_n : n \in \mathbb{N}\}$ constructed in Proposition 43. By Theorem 19 and part (c) of Proposition 43, \mathcal{G}_n will have a perfect strategy in C_{qc} if and only if $n \in X$. By Proposition 20 and part (d) of Proposition 43, \mathcal{G}_n will have a perfect strategy in C_{qc} if and only if it has a perfect strategy in C_{qa} . Because the function $n \mapsto \mathcal{G}_n$ is computable by part (b) of Proposition 43, it is undecidable to determine if the games in this family have perfect strategies in C_{qa} . □

Proof of Theorem 3. Given a computable function $f(n)$, let $X \subseteq \mathbb{N}$ be a recursive subset such that for any Turing machine accepting X , the running time

over inputs $n \leq N$ is at least $f(N)$ when N is sufficiently large. (As mentioned in the introduction, often when talking about the running time, we look at the maximum running time over inputs of size $\leq N$, rather than value $\leq N$. However, thinking of the running time in terms of the values of the inputs does not change the fact that such sets X exist.) As in the proof of Theorem 2, we can take the family of games $\{\mathcal{G}_n : n \in \mathbb{N}\}$ associated to the solution groups $\{\Gamma_n : n \in \mathbb{N}\}$ from Proposition 43. Then part (a) of Theorem 3 follows from parts (a) and (b) of Proposition 43, while parts (b) and (c) of Theorem 3 follow from parts (c) and (e) of Proposition 43, as well as Theorems 18 and 19. \square

Proof of Corollary 4. Suppose there is an algorithm to decide if a linear system game has a perfect strategy in C_q . Let $g(n)$ be the maximum running time of this algorithm on games coming from linear systems with at most n rows and columns. Note that $g(n)$ is an increasing function. Let $f(n)$ be any computable function such that

$$f(n) > g(2^{n^2}) + 2^{n^2}$$

for all $n \geq 1$. Let \mathcal{G}_n be the family of games associated to $f(n)$ as in Theorem 3. Then there is a constant C such that \mathcal{G}_n has size $\leq 2^{Cn}$ for all $n \geq 1$, and the function $n \mapsto \mathcal{G}_n$ is computable in time 2^{Cn} . Plugging \mathcal{G}_n into the algorithm to decide whether a linear system game has a perfect strategy in C_q , we get an algorithm for the language

$$X = \{n \in \mathbb{N} : \mathcal{G}_n \text{ has a perfect strategy in } C_q\}$$

with running time at most $g(2^{CN}) + 2^{CN}$ on inputs $n \leq N$. But by part (b) of Theorem 3, when N is sufficiently large the maximum running time on inputs $n \leq N$ for any algorithm for X must be at least $f(N)$. Since N^2 will eventually be larger than CN , we get a contradiction. Thus there is no algorithm to decide if a linear system game has a perfect strategy in C_q . \square

Acknowledgements

Acknowledgements I thank Jason Crann, Richard Cleve, Tobias Fritz, Li Liu, Martino Lupini, Narutaka Ozawa, Vern Paulsen, Mark Sapir, Jamie Sikora, and Thomas Vidick for helpful comments and conversations. This work was partially supported by NSERC grant number 2018-03968.

References

- [1] G. Baumslag, ‘On the residual finiteness of generalised free products of nilpotent groups’, *Trans. Amer. Math. Soc.* **106**(2) (1963), 193–209.
- [2] J. S. Bell, ‘On the Einstein Podolsky Rosen paradox’, *Physics* **1**(3) (1964), 195–200.

- [3] S. Burgdorf, M. Laurent and T. Piovosan, ‘On the closure of the completely positive semidefinite cone and linear approximations to quantum colorings’, *Electron. J. Linear Algebra* **32** (2017), 15–40.
- [4] V. Capraro and M. Lupini, *Introduction to Sofic and Hyperlinear Groups and Connes’ Embedding Conjecture*, Lecture Notes in Mathematics, 2136 (Springer, Switzerland, 2015).
- [5] R. Cleve, L. Liu and W. Slofstra, ‘Perfect commuting-operator strategies for linear system games’, *J. Math. Phys.* **58** 012202 (2017).
- [6] R. Cleve and R. Mitta, ‘Characterization of binary constraint system games’, in *Automata, Languages, and Programming*, Lecture Notes in Computer Science, 8572 (Springer, Heidelberg, 2014), 320–331.
- [7] K. J. Dykema and V. Paulsen, ‘Synchronous correlation matrices and Connes’ embedding conjecture’, *J. Math. Phys.* **57**(1) 015214 (2016).
- [8] N. Filonov and I. Kachkovskiy, ‘A Hilbert–Schmidt analog of Huaxin Lin’s theorem’. Preprint, 2010, [arXiv:1008.4002](https://arxiv.org/abs/1008.4002).
- [9] T. Fritz, ‘Tsirelson’s problem and Kirchberg’s conjecture’, *Rev. Math. Phys.* **24**(05) 1250012 (2012).
- [10] L. Glebsky, ‘Almost commuting matrices with respect to normalized Hilbert–Schmidt norm’. Preprint, 2010, [arXiv:1002.3082](https://arxiv.org/abs/1002.3082).
- [11] Z. Ji, ‘Binary constraint system games and locally commutative reductions’. Preprint, 2013, [arXiv:1310.3794](https://arxiv.org/abs/1310.3794) [quant-ph].
- [12] Z. Ji, ‘Compression of quantum multi-prover interactive proofs’, in *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2017* (ACM, New York, 2017), 289–302.
- [13] M. Junge, M. Navascués, C. Palazuelos, D. Pérez-García, V. B. Scholz and R. F. Werner, ‘Connes’ embedding problem and Tsirelson’s problem’, *J. Math. Phys.* **52**(1) 012102 (2011).
- [14] O. G. Kharlampovich, ‘A finitely presented solvable group with unsolvable word problem’, *Math. USSR Izvestija* **19**(1) (1982), 151–169.
- [15] O. G. Kharlampovich, A. Myasnikov and M. Sapir, ‘Algorithmically complex residually finite groups’, *Bull. Math. Sci.* **7**(2) (2017), 309–352.
- [16] M. Laurent and T. Piovosan, ‘Conic Approach to quantum graph parameters using linear optimization over the completely positive semidefinite cone’, *SIAM J. Optim.* **25**(4) (2015), 2461–2493.
- [17] D. Leung, B. Toner and J. Watrous, ‘Coherent state exchange in multi-prover quantum interactive proof systems’, *Chic. J. Theoret. Comput. Sci.* **19**(1) (2013), 1–18.
- [18] Debbie Leung and Bingjie Wang, in preparation.
- [19] T. A. Loring, *Lifting Solutions to Perturbing Problems in C^* -algebras*, Fields Institute monographs (American Mathematical Society, Providence, RI, 1997).
- [20] A. I. Mal’cev, ‘On the faithful representations of infinite groups of matrices’, *Amer. Math. Soc. Transl. Ser. 2* **45** (1965), 1–18.
- [21] L. Mančinska and T. Vidick, ‘Unbounded entanglement can be needed to achieve the optimal success probability’, in *Automata, Languages, and Programming*, Lecture Notes in Computer Science, 8572 (Springer, Heidelberg, 2014), 835–846.
- [22] N. David Mermin, ‘Simple unified form for the major no-hidden-variables theorems’, *Phys. Rev. Lett.* **65**(27) (1990), 3373–3376.
- [23] M. Navascués, S. Pironio and A. Acín, ‘A convergent hierarchy of semidefinite programs characterizing the set of quantum correlations’, *New J. Phys.* **10**(7) 073013 (2008).
- [24] N. Ozawa, ‘About the Connes embedding conjecture’, *Jpn. J. Math.* **8**(1) (2013), 147–183.

- [25] K. F. Pál and T. Vértesi, ‘Maximal violation of a bipartite three-setting, two-outcome Bell inequality using infinite-dimensional quantum systems’, *Phys. Rev. A* **82**(2) 022116 (2010).
- [26] A. Peres, ‘Incompatible results of quantum measurements’, *Phys. Lett. A* **151**(3) (1990), 107–108.
- [27] V. G. Pestov, ‘Hyperlinear and sofic groups: a brief guide’, *Bull. Symbolic Logic* **14**(4) (2008), 449–480.
- [28] O. Regev and T. Vidick, ‘Quantum XOR games’, *ACM Trans. Comput. Theory* **7**(4) (2015), 15:1–15:43.
- [29] V. B. Scholz and R. F. Werner, ‘Tsirelson’s problem’. Preprint, 2008, [arXiv:0812.4305](https://arxiv.org/abs/0812.4305).
- [30] J. Sikora and A. Varvitsiotis, ‘Linear conic formulations for two-party correlations and values of nonlocal games’, *Math. Program.* **162**(1–2) (2017), 431–463.
- [31] W. Slofstra, ‘Tsirelson’s problem and an embedding theorem for groups arising from non-local games’. Preprint, 2016, [arXiv:1606.03140](https://arxiv.org/abs/1606.03140).
- [32] B. S. Tsirelson, Bell inequalities and operator algebras. *Problem statement for website of open problems at TU Braunschweig*, 2006, available at <http://web.archive.org/web/20090414083019/http://www.imaph.tu-bs.de/qi/problems/33.html>.
- [33] S. Wehner, M. Christandl and A. C. Doherty, ‘Lower bound on the dimension of a quantum system given measured data’, *Phys. Rev. A* **78**(6) 062112 (2008).