

The Silkworm Z Chromosome Is Enriched in Testis-Specific Genes

K. P. Arunkumar,* Kazuei Mita[†] and J. Nagaraju*¹

*Centre of Excellence for Genetics and Genomics of Silkmoths, Laboratory of Molecular Genetics, Centre for DNA Fingerprinting and Diagnostics, Tuljaguda, Nampally, Hyderabad 500001, India and [†]National Institute of Agrobiological Sciences, Owashi 1-2, Tsukuba, Ibaraki 305-8634, Japan

Manuscript received December 22, 2008
Accepted for publication March 22, 2009

ABSTRACT

The role of sex chromosomes in sex determination has been well studied in diverse groups of organisms. However, the role of the genes on the sex chromosomes in conferring sexual dimorphism is still being experimentally evaluated. An unequal complement of sex chromosomes between two sexes makes them amenable to sex-specific evolutionary forces. Sex-linked genes preferentially expressed in one sex over the other offer a potential means of addressing the role of sex chromosomes in sexual dimorphism. We examined the testis transcriptome of the silkworm, *Bombyx mori*, which has a ZW chromosome constitution in the female and ZZ in the male, and show that the Z chromosome harbors a significantly higher number of genes expressed preferentially in testis compared to the autosomes. We hypothesize that sexual antagonism and absence of dosage compensation have possibly led to the accumulation of many male-specific genes on the Z chromosome. Further, our analysis of testis-specific paralogous genes suggests that the accumulation on the Z chromosome of genes advantageous to males has occurred primarily by translocation or tandem duplication.

IN the silkworm, *Bombyx mori*, as in other organisms, differentiation of the testis and formation of sperm are under the control of a large number of tissue- and stage-specific genes. In previous studies, a few testis-specific genes involved in silkworm spermatogenesis, such as the genes coding for BmAHAI (MIYAGAWA *et al.* 2005), BmDmc1 (KUSAKABE *et al.* 2001), and testis-specific tektin (OTA *et al.* 2002), have been reported. However, no comprehensive data on the testis transcriptomes are available for *B. mori*. Among insects, the testis transcriptome has been studied in detail only in *Drosophila melanogaster* (ANDREWS *et al.* 2000; PARISI *et al.* 2003; RANZ *et al.* 2003; MIKHAYLOVA *et al.* 2008).

Recent high-throughput genomics projects have focused on the construction, annotation, and analysis of cell-specific and tissue-specific transcriptomes, providing fundamental insights into biological processes (FIZAMES *et al.* 2004; URUSHIHARA *et al.* 2006). Data gathered from expressed sequence tags (ESTs) and microarray-based gene expression profiles are important resources for discovery of novel genes expressed in a tissue-specific manner. In this context, we analyzed the *B. mori* testis transcriptome to identify testis-specific genes and their functions. Further, we annotated the testis-specific ESTs by using protein homology data and

Gene Ontology and deduced gene structures of a few testis-specific genes using whole-genome shotgun (WGS) sequence data and ESTs.

Another rewarding exercise of analyzing the testis transcriptome is to examine whether there is any correlation between the sex-specific expression of the genes and their chromosomal location. In eukaryotes, an unequal complement of sex-determining chromosomes exists between the two sexes. Unlike mammals and dipteran insects, which have an XX/XY system of sex chromosome composition, lepidopteran insects have a ZZ/ZW or a ZZ/ZO system of sex chromosome composition. Two current hypotheses propose a contradictory fate for the genes that reside on the sex chromosomes and are preferentially or exclusively expressed in one sex. Rice's hypothesis (RICE 1984) proposes that genes with sex-biased or sex-specific expression are overrepresented on the X chromosome and has been tested in human (SAIFI and CHANDRA 1999; VALLENDER and LAHN 2004) and mouse (WANG *et al.* 2001). An alternative hypothesis suggests that the X becomes feminized because the X chromosome is present in females 50% more frequently than in males, providing evolution with more opportunity to act on genes that benefit females (REINKE *et al.* 2000); thus, genes with female-biased expression should reside preferentially on the X chromosome. So which hypothesis is correct? Or are the two hypotheses mutually compatible? Studies by WANG *et al.* (2001) on mouse male germ cells and by SAIFI and CHANDRA (1999) and LERCHER *et al.* (2003) on the human X chromosome concluded that genes

Supporting information is available online at <http://www.genetics.org/cgi/content/full/genetics.108.099994/DC1>.

¹Corresponding author: Laboratory of Molecular Genetics, Centre for DNA Fingerprinting and Diagnostics, Tuljaguda, Nampally, Hyderabad 500001, India. E-mail: jnagaraju@cdfd.org.in

expressed specifically in males are predominantly X-linked, far in excess of male-specific genes on the autosomes, thus corroborating the Rice hypothesis. In contrast, in a study of sex-biased expression in mouse, KHIL *et al.* (2004) showed that genes involved in spermatogenesis are relatively underrepresented on the X chromosome whereas female-biased genes are overrepresented on it. Further, they showed that meiotic sex chromosome inactivation (MSCI), in which sex chromosomes become heterochromatic and transcriptionally inactive, accounts for the depletion of testis-expressed genes on the X chromosome. This was inferred because the genes expressed before MSCI are overrepresented on the X chromosome. In sharp contrast, male-biased genes in *Drosophila* (PARISI *et al.* 2003; RANZ *et al.* 2003) and genes expressed in spermatogenic and oogenic cells in *Caenorhabditis elegans* (REINKE *et al.* 2004) are underrepresented on the X chromosome.

In this study we tested this phenomenon in *B. mori*, a lepidopteran model system. In silkworm, the female is the heterogametic sex and has one Z and one W chromosome, whereas the male has two Z chromosomes. We analyzed the distribution of 1104 testis-specific genes, identified by microarray analysis in an earlier study (XIA *et al.* 2007), and 1984 full-length testis-specific cDNAs and ESTs, which were mapped onto their specific locations on *B. mori* chromosomes. Our results showed that the Z chromosome (linkage group 1) harbors a significantly higher number of testis-specific genes than the autosomes. We suggest that lack of dosage compensation and sexual antagonism could have led to the accumulation of male advantageous genes on the Z chromosome. In the course of evolution, proteins that are required in higher amounts in males than in females would have been favored on the Z chromosome to facilitate phenotypic sexual dimorphism. Finally, initial analysis of testis-specific paralogs suggests that male-advantageous genes are accumulated on the Z chromosome by translocation from autosomes or tandem duplication.

MATERIALS AND METHODS

Sequence source: More than 100,000 ESTs are available for *B. mori* in the NCBI dbEST (MITA *et al.* 2003; XIA *et al.* 2004). We downloaded 9614 testis ESTs and a total of 96,051 ESTs derived from tissues other than testis (supporting information, Table S1).

Sequence analysis: Since many ESTs may be derived from the same gene, the sequences were assembled into clusters with the TGICL program (PERTEA *et al.* 2003). A cluster is defined as a unique set of sequences that have sequence similarity. A cluster that has only one sequence is termed a singleton.

On the basis of Gene Ontology (GO) annotation of closely related homologs, ESTs were assigned with molecular functions, biological process, and cellular component from the GO database (ASHBURNER *et al.* 2000) (see File S1, Figure S1, Figure S2, and Figure S3 for details).

To identify putative homologies to known proteins, 3622 clusters were subjected to BLASTx searches against the non-redundant (nr) database of the NCBI, using a cutoff *E*-value of 1×10^{-5} for parsing output files (ALTSCHUL *et al.* 1990). An *E*-value of 1×10^{-20} was used as an upper limit to assign significant homology. In cases where no significant homology was found, an *E*-value limit of 1×10^{-5} was used to assign weak homology. We found this additional category useful for data mining as many clusters do not represent full-length sequences, making it possible that only a highly divergent region of a gene sequence is available in our collection. The category of weak homology allowed us to find potential homologs in such situations.

Digital differential display: We carried out *in silico* differential display to identify genes expressed specifically in the testis. This was done by carrying out MegaBLAST with a cutoff score ≥ 50 , percentage of similarity ≥ 94 , and *E*-value ≤ 1 , against the nonredundant EST set obtained by clustering and assembly of *B. mori* ESTs from tissues other than testis and using nonredundant testis ESTs as queries. Testis ESTs that did not show any similarity with ESTs from other tissues were considered as putative testis-specific genes.

Experimental validation of a few evolutionarily conserved testis-specific genes: To confirm the testis specificity of genes predicted by digital differential display, semiquantitative RT-PCR was carried out for a select set of 15 transcripts (Table S2) using total RNA from six different tissues: midgut, fatbody, head, silk gland, epidermis, and gonads of fifth instar larvae as templates. Sex-limited silkworm strains carrying translocations of a region of chromosome 2 that harbors the gene for larval markings to the W chromosome were utilized for this purpose. Male and female larvae of this silkworm stock can be distinguished from the third instar onward by the presence of markings on the thoracic segment in females but not in males. Total RNA was isolated separately from all tissues using Trizol (Invitrogen, Carlsbad, CA), followed by DNase (Invitrogen) treatment to remove genomic DNA.

PCR was carried out using a thermal cycler (Eppendorf, Germany) under the following conditions: initial denaturation at 94° for 2 min; 30 cycles of 94° for 30 sec, 58° for 30 sec, and 72° for 2 min; and a final elongation step at 72° for 10 min. Silkworm cytoplasmic β -actin cDNA was amplified as an endogenous control. The PCR reaction components included $1 \times$ buffer, 100 μ M dNTPs, 1.5 mM $MgCl_2$, 0.5 unit Taq polymerase (MBI), and 0.5 μ M primers.

Physical mapping of testis-specific genes and analysis of testis-specific gene paralogs: In general, tissue-specific gene expression is considered as an indicator or a predictor of a tissue-specific function. Using the criterion that the intensity of expression of a gene in a particular tissue exceeds twice that in other tissues, a total of 1104 genes were identified as being testis specific in *B. mori* (XIA *et al.* 2007).

Testis-specific genes validated in this manner by microarray results were assigned chromosomal positions on the *B. mori* genome to examine their distribution on different chromosomes. Mapping was carried out using the *B. mori* physical chromosome map implemented in KAIKOBLAST and the University of Tokyo Genome Browser (UTGB). In brief, all of the gene sequences were queried against the physical map database using the BLAST program. The results were then manually parsed to determine the exact location of genes. The total number of genes on each chromosome and the number of genes per megabase of each chromosome were calculated using the map data. We also mapped 465 genes from other somatic tissues identified in XIA *et al.* (2007) through microarray analysis onto chromosomes to compare their distribution with that of testis-specific genes.

From the testis-specific genes identified through microarray analysis, paralogs were selected by combining BLAST param-

eters and homology to published annotated genes. Translated BLAST (tBLASTx) was performed for each of the testis-specific genes against the complete set of testis-specific genes. The BLAST result was parsed to obtain information on hits with a score >100. The parsed data were manually checked to group the paralogs and assigned chromosomal locations. Each group was then analyzed by carrying out BLAST against the NCBI protein nr database. Only groups that had the same functional annotation for the genes included in a group were regarded as genuine paralogs.

To strengthen our findings, we clustered 20,000 full-length cDNAs (fl-cDNAs) generated from testis (K. MITA, unpublished data) and 9614 testis ESTs (MITA *et al.* 2003; XIA *et al.* 2004) to obtain a nonredundant set of sequences. From these sequences we identified 2559 testis-specific genes by removing the sequences that were present in EST libraries derived from other tissues. Further, after excluding the microarray-validated testis-specific genes, 1857 testis-specific genes were identified as new ones and were mapped onto *B. mori* chromosomes to study their distribution on the Z chromosome and the autosomes.

RESULTS AND DISCUSSION

In the study reported here *in silico* analysis of 9614 ESTs derived from *B. mori* fifth instar larval testis revealed that several testis-specific genes are evolutionarily conserved from insects to mammals. These findings are consistent with the idea that the testis expresses a complex set of transcripts, as observed in *Drosophila* (ANDREWS *et al.* 2000). The present study also identified several families of testis-specific genes, such as those that encode tektins, dyneins, kinases, and tubulins, involved in silkworm spermatogenesis.

High gene diversity in the testis transcriptome: The availability of 9× coverage of the *B. mori* genome sequence and abundant EST resources for this insect facilitated the identification of a subset of testis-specific genes. Clustering and assembly of 9614 testis ESTs resulted in a total of 3622 unique clusters containing 1112 contigs and 2510 singletons. We also obtained 24,857 unique clusters (8819 contigs and 16,038 singletons) by clustering and assembly of the ESTs derived from other tissues.

To identify putative homologs and orthologs, we subjected the clusters to BLASTx searches against the nr database of the NCBI. A total of 2385 (66%) unique sequences showed homology with known proteins and could be assigned a putative identity. Of the 3622 clusters, 2.6% matched proteins with an *E*-value of $\leq 1e-99$ and were considered to be genuine orthologs. Thirty percent of the clusters found a hit with an *E*-value between $1e-20$ and $1e-99$ and were assigned significant homology. Finally, 16% of these clusters had a first hit with an *E*-value between $1e-19$ and $1e-05$ and were assigned weak homology to a protein from the nr database.

***In silico* differential display of *B. mori* testis ESTs:** Several interesting features about the testis transcriptome were revealed by *in silico* subtraction of testis-

derived nonredundant transcripts from the total transcripts from other somatic tissues. More than 900 of the 3622 unique clusters of testis originally identified to be of testis origin were found to be expressed only in the testis. These were regarded as putative testis-specific genes. Similar results were reported by Xia and co-workers (XIA *et al.* 2007), who found 1104 testis-specific genes using microarray analysis.

Several testis-specific genes are conserved across phyla: Using the EST database, we identified four β -*tubulin* and three α -*tubulin* genes that could be classified into at least three distinct subfamilies: ubiquitously expressed, developmentally regulated, and testis specific. Among them we identified a previously unknown α -*tubulin* gene, which we have named *bmtua4* (accession no. Bmo.6186). This gene has a single exon, similar to the three genes reported previously (KAWASAKI *et al.* 2003). *bmtua4* showed ~76% similarity to *bmtua1* and *bmtua2* and ~71% similarity to *bmtua3* genes. Whereas the β -*tubulin* (*bmtub4*) (MITA *et al.* 1995) and α -*tubulin* (*bmtua3*) genes reported earlier (KAWASAKI *et al.* 2003) were testis specific, *bmtua4* showed enhanced expression in testis. In this respect silkworm spermatogenesis has the same distinctive properties as other species in that many gene families include paralogs such as *tubulin*, *dynein light chain* genes that are expressed solely in spermatogenic cells (EDDY and O'BRIEN 1998), and others are expressed in somatic and germinal cells.

Through BLAST analysis we found 34 testis-specific genes (Table S3) that are known to have specialized functions in testis. All these genes have testis-specific homologs in species as diverse as mammals (*Homo sapiens*, *Mus musculus*, *Rattus norvegicus*, *Canis familiaris*, *Bos taurus*), birds (*Gallus gallus*), amphibia (*Xenopus laevis*), fishes (*Tetraodon nigroviridis*, *Danio rerio*), ascidians (*Ciona intestinalis*), and sea urchins (*Strongylocentrotus purpuratus*). To confirm their testis specificity, RT-PCR expression analysis of 15 of these evolutionarily conserved genes was carried out. Of these, 9, including a 717-bp testis-specific splice product of *Sperm mitochondria associated protein*, were found to be expressed exclusively in the testis; the remaining 6 genes showed male specificity with enhanced expression in the testis. Only 1 gene, a homolog of *TEGT* (*Testis Enhanced Gene Transcript*), showed equal expression in all the tissues examined (Figure 1). Expression analysis using RT-PCR was consistent with the tissue distribution of their ESTs as revealed by *in silico* analysis. Expression analysis also revealed that most of the testis-enhanced genes were male specific. Conservation of several testis-specific genes from insects to mammals suggests that the proteins they encode are probably indispensable for spermatogenesis.

Expression analysis confirmed the presence of predicted alternative splice forms of *Sperm mitochondria associated protein* and *Serine protease* genes. The *Sperm mitochondria associated protein* gene showed three tran-




FIGURE 1.—RT-PCR analysis of expression of 15 predicted testis-specific genes. β -Actin was used as an internal control. Details of accession numbers and tissue specificity as predicted *in silico* are given in Table S3. Primer sequences are given in Table S2.

scripts with amplicons of 800, 717, and 417 bp, whereas the *Serine protease* gene showed two products of 575 and 331 bp. We found one testis-specific transcript (accession no. BP127445), having no apparent similarity to any protein from the nr database of the NCBI, and it was confirmed to be testis specific through RT-PCR (Figure 1).

Physical mapping reveals abundance of testis-specific genes on the Z chromosome: An increasing body of evidence suggests that the evolutionary significance of sex chromosomes probably involves not only the mechanism of sex determination but also the evolution of sexually dimorphic traits (CHARLESWORTH 1996; RICE 1996; REINHOLD *et al.* 1998; GOTTER *et al.* 1999; HURST and RANDERSON 1999; GIBSON *et al.* 2002).

Analyzing the distribution of sex-specific genes between sex chromosomes and autosomes could uncover the underlying molecular mechanisms of sexual dimorphism since the factors governing the molecular evolution of sex-linked genes differ in several ways compared to those affecting the evolution of autosomal genes. In humans, the X chromosome has more than its fair share of genes involved in sex and reproduction (SAIFI and CHANDRA 1999; VALLENDER and LAHN 2004) compared to *Drosophila* (PARISI *et al.* 2003; RANZ *et al.* 2003) and *C. elegans* (REINKE *et al.* 2004), where the X chromosome is deficient in male-biased genes.

The biased representation of sex-specific genes on sex chromosomes is attributed to two main reasons, sexual antagonism and dosage compensation. According to the hypothesis of sexual antagonism (RICE 1984; HURST and RANDERSON 1999), an unusual homogametic sex chromosome gene content reflects a nonrandom accumulation of sexually antagonistic mutations on this chromosome. The other hypothesis concerns the epigenetic modifications of the sex chromosomes associated with meiotic sex chromosome inactivation and dosage compensation (PARISI *et al.* 2003; ROGERS *et al.* 2003; REINKE *et al.* 2004; KHIL *et al.* 2005). However, the exact role of these mechanisms still remains elusive, as all the studies have been carried out in male heterogametic sex chromosome systems (XX/XY system).

Alternatively, the mechanisms responsible for the nonrandom representation of sex-biased and sex-specific genes on the homogametic sex chromosome may be clarified by analyzing the gene content of the Z chromosome, a homogametic sex chromosome in heterogametic female organisms. Unlike the X chromosome, the Z chromosome spends more time in male individuals. If sexually antagonistic selection were the primary mechanism affecting the sex chromosome gene content, the opposite trend would be expected in the representation of sex-biased and sex-specific




FIGURE 2.—Distribution of testis-specific genes identified through microarray analysis (top panel) and through EST and fl-DNA sequence analysis (bottom panel) on different chromosomes of *B. mori*. There is a significant difference ($P < 0.001$) between the number of testis-specific genes present on the Z chromosome and on autosomes in both the cases. Among 1104 microarray-validated testis-specific genes, 1029 were successfully mapped onto *B. mori* chromosomes. The average number of testis-specific genes on autosomes was calculated to be 35 ± 2 , which is indicated by a black horizontal line on the histogram. Of 1984 testis-specific genes identified through analysis of testis-derived fl-cDNAs and ESTs, 1857 genes could be mapped onto *B. mori* chromosomes. The average number of genes on autosomes was calculated to be 63 ± 4 (green horizontal line).

genes on the X and Z chromosomes (STORCHOVA and DIVINA 2006). In a recent study in chicken, which is a female heterogametic system like *B. mori*, the Z chromosome was found to be significantly enriched in male-biased genes expressed in brain compared to female-biased genes. However, the study did not find any bias in distribution of testis-specific genes on the Z chromosome (STORCHOVA and DIVINA 2006).

Unlike in *Drosophila*, the recent findings show that there is no dosage compensation in *B. mori* (KOIKE *et al.* 2003) and other lepidopterans (JOHNSON *et al.* 1979; CHARLESWORTH 1996; RICE 1996; REINHOLD *et al.* 1998; GOTTER *et al.* 1999; HURST and RANDERSON 1999; GIBSON *et al.* 2002).

To address the question whether or not the sex-specific genes are distributed in a biased manner on the

Z chromosome, we investigated the genomic distribution of *B. mori* genes that are preferentially expressed only in testis. For this purpose, we assayed two data sets, one set comprising 1104 microarray-validated testis-specific genes (XIA *et al.* 2007) and the second comprising 1984 testis-specific genes derived from fl-cDNAs and ESTs. These were assigned chromosomal positions using the *B. mori* physical chromosome map implemented in KAIKOBLAST and UTGB. We were able to map 1029 genes and 1857 genes in the first and second data sets, respectively, onto their respective chromosomal locations (Figure 2). Physical mapping revealed many interesting features about the frequency and chromosomal distribution of testis-specific genes. Surprisingly, the Z chromosome harbored the highest number of testis-specific genes (82 and 151 from the




FIGURE 3.—Distribution of tissue-specific genes (excluding testis and ovary) on different chromosomes of *B. mori*. Of 501 other tissue-specific genes, 465 were successfully mapped. The average number of other tissue-specific genes on autosomes was calculated to be 16, which is indicated by the black horizontal line on the histogram.

first and second data sets, respectively) as compared to the average number of testis-specific genes on autosomes (35 ± 2 and 63 ± 4 ; Figure 2). A Student's *t*-test revealed that the Z chromosome harbored a significantly higher ($P < 0.001$) number of genes compared to that on autosomes. These results suggest that the locus on the Z chromosome is more than two times more likely to have testis-specific expression than an autosomal locus.

We also assessed the allocation of genes expressed exclusively in a single tissue excluding testis and ovary, between autosomes and the Z chromosome. For this purpose, we mapped 465 additional tissue-specific genes (XIA *et al.* 2007), which showed no apparent chromosome bias in their distribution (Figure 3). These results indicate that testis-specific genes are more often located on the Z chromosome than is expected by chance.

To study the density of testis-specific genes on different chromosomes, we calculated the number of genes per megabase for each chromosome. The Z showed a significantly higher frequency of testis-specific genes (Student's *t*-test, $P < 0.001$), with 11.45 genes per megabase of chromosome compared to 6.54 ± 0.18 genes on autosomes (Table 1). These results provide further evidence that overrepresentation of testis-specific genes on the Z is not due to its sheer size or to a higher frequency of genes on the Z chromosome. To determine the presence of any clusters of testes-specific genes, we constructed a physical map using Karyoview (<http://www.ensembl.org/index.html>). The map showed the spatial distribution of genes on all *B. mori* chromosomes (Figure 4). It was evident from the map that genes are quite evenly and densely distributed on the Z chromosome. However, the analysis confirmed the presence of several clusters of testis-specific genes on different autosomes, suggesting an apparent bias in their distribution compared to the Z. Information on the exact chromosomal location of each gene can be found in Table S4.

Why is a certain category of genes abundant on the Z chromosome? Previous studies have shown that in butterflies a disproportionate number of genes related to sexuality, reproduction, and speciation are located on the Z chromosome, which forms approximately one-sixtieth of the genome in females. Female mate-selection behavior, male courtship signals, female limitation of color polymorphism, and mimicry are thought to result largely from interactions between autosomal genes and uncompensated Z-linked regulatory genes (STEHR 1959; SHEPPARD 1961; COOK 1964; GRULA and TAYLOR 1980; SPERLING 1994). In *B. mori*, four muscle protein genes, *Bmkettin*, *Bmtitin1*, *Bmtitin2*, and *Bmprojectin*, and another gene involved in locomotor behavior, *Bmhig*, are Z-linked and are not dosage compensated (SUZUKI *et al.* 1998, 1999; KOIKE *et al.* 2003). These genes are functionally conserved between *B. mori* and *Drosophila*. In silkworm, although adult moths have lost flight during the course of domestication, male moths flap their wings more vigorously than females to approach sedentary females. In this context, Z-linkage of uncompensated muscle proteins encoding genes ensures a higher quantity of these proteins in males. It is speculated that the Z chromosome has evolved through a process of genome shuffling to accumulate genes whose products are required at higher levels in males (KOIKE *et al.* 2003). Considering these results together, we argue that the absence of dosage compensation and non-random distribution of male-biased genes on the Z chromosome have ensured accumulation of male advantageous genes in silkworm.

Our results are consistent with the Rice thesis (RICE 1984), which suggests that sex-linked genes are likely to evolve sex-related functions. The beneficial mutations will be selected naturally if they are located on sex chromosomes rather than on autosomes. The evolutionary dynamics of male-benefit mutations were considered when they first appear as rare alleles on X chromosomes in contrast with autosomes. When male-

TABLE 1
Number and frequency of testis-specific genes on different chromosomes of the silkworm, *B. mori*

Chromosome	Size (Mb)	No. of testis-specific genes (full-length cDNA and ESTs, and microarray-validated genes)	No. of testis-specific genes/Mb
Chromosome 1 (Z chromosome)	20.35	233	11.45
Chromosome 2	7.94	50	6.30
Chromosome 3	14.68	135	9.20
Chromosome 4	17.87	125	6.99
Chromosome 5	18.37	146	7.95
Chromosome 6	15.91	114	7.17
Chromosome 7	12.75	78	6.12
Chromosome 8	15.36	125	8.14
Chromosome 9	16.47	86	5.23
Chromosome 10	17.22	108	6.27
Chromosome 11	19.58	135	6.89
Chromosome 12	16.47	102	6.19
Chromosome 13	17.45	130	7.45
Chromosome 14	12.60	64	5.08
Chromosome 15	17.97	139	7.74
Chromosome 16	13.70	60	4.38
Chromosome 17	14.34	107	7.46
Chromosome 18	15.16	94	6.20
Chromosome 19	13.19	89	6.75
Chromosome 20	10.61	65	6.12
Chromosome 21	14.92	79	5.29
Chromosome 22	17.36	117	6.74
Chromosome 23	20.08	141	7.01
Chromosome 24	13.96	97	6.95
Chromosome 25	14.14	104	7.36
Chromosome 26	10.76	63	5.86
Chromosome 27	10.44	50	4.79
Chromosome 28	10.33	50	4.84
Total	419.98	2886	
Autosomal average \pm SE	14.80 \pm 0.47	98 \pm 6	6.54 \pm 0.18

benefit mutations are rare, autosomal recessive alleles offer no advantage to (heterozygous) males and thus would be unlikely to spread in the population. By contrast, X-linked recessive alleles would confer a benefit to hemizygous males, greatly favoring the alleles' likelihood of permeating the population. Eventually, as an allele's frequency increased in the population, female fitness would be diminished by the detrimental effects of homozygosity. This would generate adaptive pressure to limit the gene's expression to males through additional mutations. Therefore, it was postulated that X chromosomes should evolve to harbor a disproportionate share of male-specific genes functioning in male differentiation (RICE 1984; WANG *et al.* 2001).

By contrast, Z linkage can facilitate the increase of traits that also favor the homogametic sex. Dominant mutations favoring the homogametic sex have a greater chance to be fixed on the Z chromosome where they are exposed to selection, because two-thirds of the Z chromosomes reside in homogametic individuals com-

pared to only one-third in heterogametic individuals and due to absence of dosage compensation. In the Z chromosome, dominant sexually antagonistic genes that favor the homogametic sex are expressed at a higher rate in the sex where they are positively selected. Our findings are consistent with this prediction.

Possible translocation of male advantageous genes onto Z chromosomes from autosomes: Duplicated genes may acquire novel functions and altered expression patterns (OHNO *et al.* 1968) and thus contribute to diversification of tissues during development (MIKHAYLOVA *et al.* 2008). Analysis of such paralogs provides a unique opportunity to study genome evolution as it sheds light on the history of gene duplication and gene trafficking between chromosomes through translocation events. In *Drosophila*, a number of testis-specific genes have been reported to be generated by gene duplications, where the duplicated gene is specifically expressed in the male reproductive system while the parental gene is ubiquitous (NURMINSKY *et al.* 1998; BETRAN and LONG 2003).




FIGURE 4.—Physical map showing the distribution of 1029 microarray-validated testis-specific genes on *B. mori* chromosomes.

In the present study we have used the data on paralogous genes that express in *B. mori* testis to investigate the evolutionary causes that lead to the enrichment of testis-specific genes on the Z chromosome. Through BLAST analysis we obtained 30 groups of paralogs (comprising 74 of 1104 testis-specific genes), and among them only 12 were found in clusters on different chromosomes. In contrast, the remaining 18 did not show any clustering, and although some paralogs were mapped onto the same chromosome, they were located far apart from each other. Duplications were more frequent among genes coding for dynein proteins (3 of 30 paralogous groups), probably because of their requirement in large amounts in sperm for motility (Moss *et al.* 1992). None of the clustered coexpressed testis-specific genes showed similarity to transposable elements, in contrast with nonclustered paralogous groups, where 6 paralogous groups were found to code for transposable elements (*e.g.*, reverse transcriptase, transposon polyprotein, and transposase).

One or more genes in the six nonclustered paralogous groups were located on the Z chromosome (Table S5), corresponding to ~30% of the nonclustered paralogs, the highest for any chromosome. Our specu-

lation is that during the course of evolution male advantageous genes were translocated onto the Z chromosome and selected positively. This may explain the preponderance of one of the copies of testis-specific nonclustered paralogs on the Z chromosome. Also, 3 (25%) of the 12 coexpressed clusters were present on the Z chromosome, again the highest number for any chromosome. Analysis of expression patterns of paralogous genes in testes would provide valuable information to clarify the functional relationships between these genes (MIKHAYLOVA *et al.* 2008). On the basis of these results we surmise that testis-specific paralogs are more concentrated on Z chromosomes either by clustering or by the location of one of the copies of nonclustered paralogs. Such translocations followed by fixation may be responsible for the enrichment of testis-specific genes on the Z chromosome. This argument is in line with the proposal by CHARLESWORTH and CHARLESWORTH (1980) that translocations from the autosomes to the X chromosome would be favored if the translocated region harbored sexually antagonistic genes.

Conclusions: *In silico* analyses of the *B. mori* transcriptome provide a foundation for further in-depth analysis of activities in male germ cells. Information produced by the sequencing of *B. mori* cDNA libraries and identification of testis-specific genes in the present study provide new insights into the structure, diversity, and molecular evolution of genes involved in spermatogenesis in lepidopterans in particular and insects in general. Our study of the distribution of testis-specific genes on *B. mori* chromosomes has shown that testis-specific genes are accumulated on the Z chromosomes either by translocation from other chromosomes or by tandem duplication on the Z chromosome.

We thank two anonymous reviewers and Marian R. Goldsmith for their comments on the manuscript. This work was supported by a Centre of Excellence grant from the Department of Biotechnology, Government of India to J.N. K.P.A. is a recipient of a fellowship from the Council of Scientific and Industrial Research, India.

LITERATURE CITED

- ALTSCHUL, S. F., W. GISH, W. MILLER, E. W. MYERS and D. J. LIPMAN, 1990 Basic local alignment search tool. *J. Mol. Biol.* **215**: 403–410.
- ANDREWS, J., G. G. BOUFFARD, C. CHEADLE, J. LU, K. G. BECKER *et al.*, 2000 Gene discovery using computational and microarray analysis of transcription in the *Drosophila melanogaster* testis. *Genome Res.* **10**: 2030–2043.
- ASHBURNER, M., C. A. BALL, J. A. BLAKE, D. BOTSTEIN, H. BUTLER *et al.*, 2000 Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat. Genet.* **25**: 25–29.
- BETRAN, E., and M. LONG, 2003 *Dntf-2i*, a young *Drosophila* retroposed gene with specific male expression under positive Darwinian selection. *Genetics* **164**: 977–988.
- CHARLESWORTH, B., 1996 The evolution of chromosomal sex determination and dosage compensation. *Cult. Biol.* **6**: 149–162.
- CHARLESWORTH, D., and B. CHARLESWORTH, 1980 Sex differences in fitness and selection for centric fusions between sex-chromosomes and autosomes. *Genet. Res.* **35**: 205–214.
- COOK, A. G., 1964 Dosage compensation and sex-chromatin in non-mammals. *Genet. Res.* **5**: 354–365.

- EDDY, E. M., and D. A. O'BRIEN, 1998 Gene expression during mammalian meiosis. *Curr. Top. Dev. Biol.* **37**: 141–200.
- FIZAMES, C., S. MUNOS, C. CAZETTES, P. NACRY, J. BOUCHEREZ *et al.*, 2004 The Arabidopsis root transcriptome by serial analysis of gene expression. Gene identification using the genome sequence. *Plant Physiol.* **134**: 67–80.
- GIBSON, J. R., A. K. CHIPPINDALE and W. R. RICE, 2002 The X chromosome is a hot spot for sexually antagonistic fitness variation. *Proc. Biol. Sci.* **269**: 499–505.
- GOTTER, A. L., J. D. LEVINE and S. M. REPERT, 1999 Sex-linked period genes in the silkworm, *Antheraea pernyi*: implications for circadian clock regulation and the evolution of sex chromosomes. *Neuron* **24**: 953–965.
- GRULA, J. W., and O. R. TAYLOR, JR., 1980 The effect of X-chromosome inheritance on mate-selection behavior in the sulfur butterflies, *Colias eurytheme* and *C. philodice*. *Evolution* **34**: 688–695.
- HURST, L. D., and J. P. RANDERSON, 1999 An eXceptional chromosome. *Trends Genet.* **15**: 383–385.
- JOHNSON, L. D., M. BINDER and R. A. LAZZARINI, 1979 A defective interfering vesicular stomatitis virus particle that directs the synthesis of functional proteins in the absence of helper virus. *Virology* **99**: 203–206.
- KAWASAKI, H., K. SUGAYA, G. X. QUAN, J. NOHATA and K. MITA, 2003 Analysis of alpha- and beta-tubulin genes of *Bombyx mori* using an EST database. *Insect Biochem. Mol. Biol.* **33**: 131–137.
- KHIL, P. P., N. A. SMIRNOVA, P. J. ROMANIENKO and R. D. CAMERINI-OTERO, 2004 The mouse X chromosome is enriched for sex-biased genes not subject to selection by meiotic sex chromosome inactivation. *Nat. Genet.* **36**: 642–646.
- KHIL, P. P., B. OLIVER and R. D. CAMERINI-OTERO, 2005 X for intersection: retrotransposition both on and off the X chromosome is more frequent. *Trends Genet.* **21**: 3–7.
- KOIKE, Y., K. MITA, M. G. SUZUKI, S. MAEDA, H. ABE *et al.*, 2003 Genomic sequence of a 320-kb segment of the Z chromosome of *Bombyx mori* containing a kettin ortholog. *Mol. Genet. Genomics* **269**: 137–149.
- KUSAKABE, T., Y. KAWAGUCHI, T. MAEDA and K. KOGA, 2001 Role of interaction between two silkworm RecA homologs in homologous DNA pairing. *Arch. Biochem. Biophys.* **388**: 39–44.
- LERCHER, M. J., A. O. URRUTIA and L. D. HURST, 2003 Evidence that the human X chromosome is enriched for male-specific but not female-specific genes. *Mol. Biol. Evol.* **20**: 1113–1116.
- MIKHAYLOVA, L. M., K. NGUYEN and D. I. NURMINSKY, 2008 Analysis of the *Drosophila melanogaster* testes transcriptome reveals coordinate regulation of paralogous genes. *Genetics* **179**: 305–315.
- MITA, K., M. NENOI, M. MORIMYO, H. TSUJI, S. ICHIMURA *et al.*, 1995 Expression of the *Bombyx mori* beta-tubulin-encoding gene in testis. *Gene* **162**: 329–330.
- MITA, K., M. MORIMYO, K. OKANO, Y. KOIKE, J. NOHATA *et al.*, 2003 The construction of an EST database for *Bombyx mori* and its application. *Proc. Natl. Acad. Sci. USA* **100**: 14121–14126.
- MIYAGAWA, Y., J. M. LEE, T. MAEDA, K. KOGA, Y. KAWAGUCHI *et al.*, 2005 Differential expression of a *Bombyx mori* AHA1 homologue during spermatogenesis. *Insect Mol. Biol.* **14**: 245–253.
- MOSS, A. G., W. S. SALE, L. A. FOX and G. B. WITMAN, 1992 The alpha subunit of sea urchin sperm outer arm dynein mediates structural and rigor binding to microtubules. *J. Cell Biol.* **118**: 1189–1200.
- NURMINSKY, D. I., M. V. NURMINSKAYA, D. DE AGUIAR and D. L. HARTL, 1998 Selective sweep of a newly evolved sperm-specific gene in *Drosophila*. *Nature* **396**: 572–575.
- OHNO, S., U. WOLF and N. B. ATKIN, 1968 Evolution from fish to mammals by gene duplication. *Hereditas* **59**: 169–187.
- OTA, A., T. KUSAKABE, Y. SUGIMOTO, M. TAKAHASHI, Y. NAKAJIMA *et al.*, 2002 Cloning and characterization of testis-specific tektin in *Bombyx mori*. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **133**: 371–382.
- PARISI, M., R. NUTTALL, D. NAIMAN, G. BOUFFARD, J. MALLEY *et al.*, 2003 Paucity of genes on the *Drosophila* X chromosome showing male-biased expression. *Science* **299**: 697–700.
- PERTEA, G., X. HUANG, F. LIANG, V. ANTONESCU, R. SULTANA *et al.*, 2003 TIGR gene indices clustering tools (TGICL): a software system for fast clustering of large EST datasets. *Bioinformatics* **19**: 651–652.
- RANZ, J. M., C. I. CASTILLO-DAVIS, C. D. MEIKLEJOHN and D. L. HARTL, 2003 Sex-dependent gene expression and evolution of the *Drosophila* transcriptome. *Science* **300**: 1742–1745.
- REINHOLD, K., M. D. GREENFIELD, Y. JANG and A. BROCE, 1998 Energetic cost of sexual attractiveness: ultrasonic advertisement in wax moths. *Anim. Behav.* **55**: 905–913.
- REINKE, V., H. E. SMITH, J. NANCE, J. WANG, C. VAN DOREN *et al.*, 2000 A global profile of germline gene expression in *C. elegans*. *Mol. Cell* **6**: 605–616.
- REINKE, V., I. S. GIL, S. WARD and K. KAZMER, 2004 Genome-wide germline-enriched and sex-biased expression profiles in *Caenorhabditis elegans*. *Development* **131**: 311–323.
- RICE, W. R., 1984 Sex chromosomes and the evolution of sexual dimorphism. *Evolution* **38**: 735–742.
- RICE, W. R., 1996 Sexually antagonistic male adaptation triggered by experimental arrest of female evolution. *Nature* **381**: 232–234.
- ROGERS, D. W., M. CARR and A. POMIANKOWSKI, 2003 Male genes: X-pelled or X-cluded? *BioEssays* **25**: 739–741.
- SAIFI, G. M., and H. S. CHANDRA, 1999 An apparent excess of sex- and reproduction-related genes on the human X chromosome. *Proc. Biol. Sci.* **266**: 203–209.
- SHEPPARD, P. M., 1961 Some contributions to population genetics resulting from the study of the Lepidoptera. *Adv. Genet.* **10**: 165–216.
- SPELRLING, F. A. H., 1994 Sex-linked genes and species differences in Lepidoptera. *Can. Entomol.* **126**: 807–818.
- STEHR, G., 1959 Hemolymph polymorphism in a moth and the nature of sex-controlled inheritance. *Evolution* **13**: 537–560.
- STORCHOVA, R., and P. DIVINA, 2006 Nonrandom representation of sex-biased genes on chicken Z chromosome. *J. Mol. Evol.* **63**: 676–681.
- SUZUKI, M. G., T. SHIMADA and M. KOBAYASHI, 1998 Absence of dosage compensation at the transcription level of a sex-linked gene in a female heterogametic insect, *Bombyx mori*. *Heredity* **81** (Pt. 3): 275–283.
- SUZUKI, M. G., T. SHIMADA and M. KOBAYASHI, 1999 Bm kettin, homologue of the *Drosophila* kettin gene, is located on the Z chromosome in *Bombyx mori* and is not dosage compensated. *Heredity* **82**(Pt. 2): 170–179.
- URUSHIHARA, H., T. MORIO and Y. TANAKA, 2006 The cDNA sequencing project. *Methods Mol. Biol.* **346**: 31–49.
- VALLENDER, E. J., and B. T. LAHN, 2004 How mammalian sex chromosomes acquired their peculiar gene content. *BioEssays* **26**: 159–169.
- WANG, P. J., J. R. MCCARREY, F. YANG and D. C. PAGE, 2001 An abundance of X-linked genes expressed in spermatogonia. *Nat. Genet.* **27**: 422–426.
- XIA, Q., Z. ZHOU, C. LU, D. CHENG, F. DAI *et al.*, 2004 A draft sequence for the genome of the domesticated silkworm (*Bombyx mori*). *Science* **306**: 1937–1940.
- XIA, Q., D. CHENG, J. DUAN, G. WANG, T. CHENG *et al.*, 2007 Microarray-based gene expression profiles in multiple tissues of the domesticated silkworm, *Bombyx mori*. *Genome Biol.* **8**: R162.

GENETICS

Supporting Information

<http://www.genetics.org/cgi/content/full/genetics.108.099994/DC1>

Silkworm Z Chromosome is Enriched in Testis-Specific Genes

K. P. Arunkumar, Kazuei Mita and J. Nagaraju

Copyright © 2009 by the Genetics Society of America

DOI: 10.1534/genetics.108.099994

File S1**Gene Ontology annotation**

GO annotation generates a dynamic controlled vocabulary that can be applied to all organisms, even while knowledge of gene and protein roles in cells is still accumulating and changing. To this end, the Seqdblite FASTA sequence flat file was downloaded from the GO database. By running BLAST against Seqdblite, closest homologue was identified. From BLAST output, molecular functions, biological processes and cellular localisation were parsed by building an in-house GO database in MySQL from the GO-term-database flat file, downloaded from Gene Ontology Database Downloads (<http://www.godatabase.org/dev/>). The Perl-DBI was used to interface with MySQL, to extract the parent terms of each individual GO term that are obtained by parsing BLAST output. The output was then represented graphically.

All ESTs were assigned a biological process, molecular function and cellular component using Gene Ontology (GO) database. The closest annotated homologue in the GO database was used for assigning these categories. The results of the GO annotation are graphically represented in Figures S1-3.

Many of the gene products were found to be localized in cell (42%). In cell, gene products were predominant in intracellular region (78%) which comprised of localizations in intracellular organelle (38%) and cytoplasm (29%). The other localizations were organelle (29%) followed by protein complex (18%) (Figure S1). The most abundant category of molecular function was binding (47%) wherein many of them were protein binding (69%). The next category was catalytic activity (17%) (Figure S2). In case of biological processes, majority of ESTs showed physiological process (36%) in which cellular physiological process was about 75%. The next category was cellular process (35%). The other processes were development (10%), regulation of biological process (6%). ESTs showing similarity to proteins involved in reproduction accounted for only 4% of total ESTs. This category consists of genes involved in male gamete formation, germ cell migration, fertilization etc. (Figure S3).




FIGURE S1.—Annotated GO terms of *B. mori* testis EST libraries. (A) Gene Ontology electronic annotation in the category 'cellular component'. The largest proportion of annotated ESTs was found to be located in 'cell' (42.2%). In the cell, gene products were abundant in (B) intracellular region (78%) which comprised of localizations in intracellular organelle (38%) and cytoplasm (29%).




FIGURE S2.—Categorization of *B. mori* testis ESTs based on (A) molecular function. Majority of the ESTs showed binding activity. (B) The binding function was further classified into different sub-classes.




FIGURE S3.—Functional classification of *B. mori* testis ESTs based on (A) biological processes. Around 35 % of ESTs were involved in cellular processes. (B) Cellular processes were further classified into cellular physiological process, cell differentiation, cell communication, membrane fusion and regulation of cellular processes.

TABLE S1

Number of *B. mori* ESTs derived from different tissues at different developmental stages, downloaded from dbEST

Library	ESTs downloaded from dbEST
Fatbody	19,792
Embryo	17,544
Hemocyte	13,451
Ovary	12,928
Silk gland	9,680
Midgut	7,252
Testis	9,614
Compound Eye	5,390
Brain	3,403
Pheromone gland	3,042
Malpighian tubules	1,697
Wing disc	1,059
Prothoracic gland	813
Total	1,05,665

TABLE S2**Details of primers used in RT-PCR expression analysis of 15 predicted testis specific genes**

Sl. No.	Gene name	Forward primer	Reverse primer
1	<i>Bm-meichroacidin</i>	GATACGAAGGCGATTGGAAA	TGGTCAACAGGTGAAATCCA
2	<i>Bmtektin-1</i>	AGCGAATGCTTTCTCTGGAA	TGTCGCTCCAATCAAATTCA
3	<i>Bmtektin-2</i>	AATGCAAGCTTTTCGGAAACA	TTCTGTGTTTTGCTGCCTTG
4	<i>Outer dense fiber of sperm tails</i>	GAACAAGCTGAGGGAGTTGG	GCCTCATCTGAGTCGCTTTC
5	<i>Male specific sperm protein</i>	GCGAGAGGTGCTTATGGTTC	TATTTATGCGCACCATGGAC
6	<i>BLu protein</i>	ACGATGCTGGATCAAATTC	TTTTTGCAAGCCTTCTCTCC
7	<i>Serine protease</i>	CGGATACGGTCAATCAGCTT	GGCTTCGTACACGGAGAGAG
8	<i>bmtua4</i>	CATACCGCCAACCTTTTCCAC	GCTGCATCTGAATGTTCCAA
9	<i>Testis protein</i>	ATCGCTTCAGTCGTGGATCT	CCTCAATCGAGAGCCAAAAG
10	<i>Sperm mitochondria associated protein</i>	AGGAATCACCCACGTGCTAC	CTCGTCGTAGGGATCAGCTC
11	<i>Bm-tegt</i>	ACGGTTTGGTTCCCTCCTTCT	CTCTGCTGCTGCTATGCTTG
12	<i>Novel testis specific transcript</i>	GCACGTGTAAATGGTGATGC	TTGTTTCACGGGAGAAGGAC
13	<i>Mage-D1</i>	ACCTTGCTACCCTGGATGTG	CACGAACCAAATGAAGGACA
14	<i>Dynein light chain</i>	GGTTTGGGCAAAAAGACAGA	CACTGGCGAAACCAAAAAGAT
15	<i>Male germ cell-associated kinase</i>	TCACAAACCGTACTTTACAGAGA	TTTTTCCCCTGTGTCTCGTC

TABLE S3

Testis- specific gene homologues identified by BLAST analyses of *B. mori* testis ESTs that are conserved in different animal phyla

Unigene / dbEST ID	Gene description	Tissue specificity
Bmo.521	<i>Bm-Meichroacidin</i>	Testis specific
CK537614	<i>Dynein light chain 3 (BmDLC3a)</i>	Testis specific
Bmo.7465	<i>Dynein light chain</i>	Testis specific
Bmo.1163	<i>Tubulin beta chain (bmtub4)</i>	Testis specific
Bmo.525	<i>Tubulin alpha chain (bmtua3)</i>	Testis specific
Bmo.764	<i>Bmtektin-1 (BA3388)</i>	Testis specific
CK537544	<i>Bmtektin-2</i>	Testis specific
CK536102	<i>Serine protease-6</i>	Testis specific
CK537191	<i>Serine kinase 1</i>	Testis specific
Bmo.5916	Sperm mitochondria-associated cysteine-rich protein	Testis specific
CK536677	<i>Testis protein</i>	Testis specific
Bmo.110	<i>BLu protein</i>	Testis specific
Bmo.709	<i>ATPase inhibitor (BAB39164)</i>	Testis specific
Bmo.5017	<i>Male germ cell-associated kinase</i>	Testis specific
Bmo.514	<i>Mage-d1</i>	Testis specific
CK534071	<i>Bm-tesmin</i>	Testis specific
CK533275	<i>Bm TPX1</i>	Testis specific
Bmo.4264	<i>Serine/threonine protein kinase MAK</i>	Testis specific
CK533681	<i>Structural sperm protein</i>	Testis specific
CK533950	<i>Male sterility protein 2-like protein</i>	Testis specific
CK537063	<i>Sperm ion channel</i>	Testis specific
CK536763	<i>Sperm associated antigen 6</i>	Testis specific
CK533294	<i>Sperm associated antigen 9</i>	Testis enhanced
Bmo.6386	<i>Outer dense fiber of sperm tails 2</i>	Testis enhanced
Bmo.2817	<i>Outer dense fiber of sperm tails 3</i>	Testis enhanced
Bmo.2777	<i>Sperm nuclear basic protein</i>	Testis enhanced
CK537618	<i>Channel, sperm associated 4</i>	Testis enhanced
Bmo.956	<i>Dynein light chain 3 (BmDLC3)</i>	Testis enhanced
Bmo.6404	<i>Tubulin alpha chain (bmtua4)</i>	Testis enhanced
Bmo.6412	<i>Testis intracellular mediator protein</i>	Testis enhanced
Bmo.2293	<i>Bm-tegt</i>	Equal expression
BP127933	<i>Testis specific 10</i>	Equal expression
Bmo.5282	<i>Outer dense fiber of sperm tails 1</i>	Equal expression
Bmo.6298	<i>Motile sperm domain containing 3</i>	Equal expression

Note: Many clusters did not have a corresponding unigene ID. Such clusters are represented here with the dbEST ID of first EST in the cluster.

TABLE S4

Chromosomal location of each gene

Please see the Excel file available at

<http://www.genetics.org/cgi/content/full/genetics.108.099994/DC1>