

The small glycine-rich RNA binding protein *AtGRP7* promotes floral transition in *Arabidopsis thaliana*

Corinna Streitner¹, Selahattin Danisman^{1,†}, Franziska Wehrle^{1,‡}, Jan C. Schöning¹, James R. Alfano² and Dorothee Staiger^{1,*}

¹Department of Molecular Cell Physiology, University of Bielefeld, Bielefeld, Germany, and

²Center for Plant Science Innovation and Department of Plant Pathology, University of Nebraska, Lincoln, NE, USA

Received 25 February 2008; revised 14 May 2008; accepted 21 May 2008; published online 1 August 2008.

*For correspondence (fax ++49 521 106 6410; e-mail dorothee.staiger@uni-bielefeld.de).

†Present address: Plant Research International, Wageningen, The Netherlands.

‡Present address: Herbamed AG, CH-9055 Bühler, Switzerland.

Summary

The RNA binding protein *AtGRP7* is part of a circadian slave oscillator in *Arabidopsis thaliana* that negatively autoregulates its own mRNA, and affects the levels of other transcripts. Here, we identify a novel role for *AtGRP7* as a flowering-time gene. An *atgrp7-1* T-DNA mutant flowers later than wild-type plants under both long and short days, and independent RNA interference lines with reduced levels of *AtGRP7*, and the closely related *AtGRP8* protein, are also late flowering, particularly in short photoperiods. Consistent with the retention of a photoperiodic response, the transcript encoding the key photoperiodic regulator *CONSTANS* oscillates with a similar pattern in *atgrp7-1* and wild-type plants. In both the RNAi lines and in the *atgrp7-1* mutant transcript levels for the floral repressor *FLC* are elevated. Conversely, in transgenic plants ectopically overexpressing *AtGRP7*, the transition to flowering is accelerated mainly in short days, with a concomitant reduction in *FLC* abundance. The late-flowering phenotype of the RNAi lines is suppressed by introducing the *flc-3* loss-of-function mutation, suggesting that *AtGRP7* promotes floral transition, at least partly by downregulating *FLC*. Furthermore, vernalization overrides the late-flowering phenotype. Retention of both the photoperiodic response and vernalization response are features of autonomous pathway mutants, suggesting that *AtGRP7* is a novel member of the autonomous pathway.

Keywords: *Arabidopsis*, flowering time, circadian clock, post-transcriptional regulation, autonomous pathway, RNA binding protein.

Introduction

The appropriate timing of the transition from vegetative to reproductive growth is controlled by a suite of signaling pathways responding to endogenous cues and tracking environmental signals, such as ambient temperature and light quality (Corbesier and Coupland, 2006; Kobayashi and Weigel, 2007; Putterill *et al.*, 2004; Simpson and Dean, 2002). *Arabidopsis thaliana* flowers earlier in long photoperiods than in short photoperiods. This photoperiodic flower induction is mediated by the endogenous circadian clock. The zinc-finger protein *CONSTANS* (CO) plays a critical role in interpreting day length to initiate floral transition. CO mRNA oscillates with a circadian rhythm, and peaks at the end of the daily light phase in long days (LD), but after the light–dark transition in short days (SD) (Suarez-Lopez *et al.*, 2001). In light, the CO protein is stabilized, and thus accu-

mulates to a level sufficient to induce flowering (Valverde *et al.*, 2004). CO, in turn, directly activates *FLOWERING LOCUS T* (*FT*) (Abe *et al.*, 2005; An *et al.*, 2004; Wigge *et al.*, 2005). Movement of FT protein from phloem cells in the leaves to the apex induces flower formation (Corbesier *et al.*, 2007; Jaeger and Wigge, 2007; Mathieu *et al.*, 2007).

FT and *SUPPRESSOR OF CONSTANS* (*SOC1*), encoding two of the earliest targets of the photoperiodic pathway (Borner *et al.*, 2000; Samach *et al.*, 2000), are also negatively regulated by the MADS-box protein *FLOWERING LOCUS C* (*FLC*), which is a key repressor of flowering (Lee *et al.*, 2000; Michaels and Amasino, 1999; Michaels *et al.*, 2005; Sheldon *et al.*, 1999). The repressive effects of FLC can be overcome by prolonged cold treatment (vernalization) to ensure that flowering occurs when winter is over. During this process,

FLC is silenced through chromatin modification (Bastow *et al.*, 2004; Sung and Amasino, 2004).

Genes of the autonomous pathway promote floral transition independently of temperature and daylength, also via the suppression of *FLC*. For example, *FVE* is a component of a histone deacetylase complex (He *et al.*, 2003), and inhibits *FLC* expression by promoting deacetylation of *FLC* chromatin (Ausin *et al.*, 2004). FLOWERING LOCUS D (FLD) is a homolog of human lysine-specific demethylase 1 (LSD1), but has also been found to affect the histone acetylation state at the *FLC* locus (He *et al.*, 2003; Shi *et al.*, 2004). Additionally, RNA processing appears to play a crucial role in the autonomous pathway. *FCA* and *FPA* harbor multiple RNA recognition motifs (RRMs), and FLOWERING LOCUS K (FLK) has three K homology domains implicated in RNA binding (Lim *et al.*, 2004; Macknight *et al.*, 1997; Schomburg *et al.*, 2001). *FY* encodes a protein with similarity to the yeast polyadenylation factor *Psf2p* (Simpson *et al.*, 2003). *FCA* negatively regulates its own expression in concert with *FY* by promoting the cleavage and polyadenylation at a poly(A) site within its third intron (Quesada *et al.*, 2003; Simpson *et al.*, 2003). This leads to the formation of a truncated transcript, *FCA-β*, limiting the production of fully spliced transcript *FCA-γ*, which encodes the full-length functional protein (Macknight *et al.*, 2002). This auto regulation adjusts the level of active *FCA*, and thus ultimately determines the *FLC* mRNA accumulation.

Negative autoregulation at the post-transcriptional level is also the hallmark of the *AtGRP7* (*Arabidopsis thaliana* glycine-rich RNA binding protein 7) protein that influences the amplitude of its own circadian transcript oscillations through alternative splicing (Schöning *et al.*, 2007; Staiger *et al.*, 2003b). *AtGRP7* has a single RRM and a glycine-rich stretch. The transcript oscillates with a peak at the end of the daily light phase. Constitutive over expression of *AtGRP7* favors the use of a cryptic intronic 5' splice site, leading to a short-lived splice form and damping of the *AtGRP7* transcript oscillations. This feedback loop is under the control of the *Arabidopsis* core clockwork, and in turn influences other transcripts including *AtGRP8* encoding a closely related glycine-rich RRM protein (Schöning *et al.*, 2007; Staiger *et al.*, 2003b).

Here, we identify a previously unrecognised task for *AtGRP7* in the network regulating floral transition. An *atgrp7* T-DNA mutant, as well as transgenic plants with reduced *AtGRP7* expression, as a result of RNA interference, have a late-flowering phenotype, whereas transgenic plants ectopically overexpressing *AtGRP7* flower early. Nevertheless, these plants with altered *AtGRP7* levels retain a photoperiodic response. The effect on flowering is in large parts mediated by *FLC*, and the late-flowering phenotype can be overcome by vernalization, implicating *AtGRP7* in the autonomous pathway.

Results

An atgrp7 T-DNA insertion mutant is late flowering

Because post-transcriptional control emerges as an important mechanism in flowering time control, we asked whether *AtGRP7* plays a role in floral transition. First, we investigated how the loss of *AtGRP7* impacts on the flowering time in the *atgrp7-1* mutant from the SALK collection with a T-DNA insertion in the 5' region (Fu *et al.*, 2007). Under SDs, *atgrp7-1* plants formed about 61 leaves at the onset of bolting, compared with 54 leaves in wild-type (WT) plants, and under LDs, *atgrp7-1* plants flowered with 15 leaves compared with WT flowering (13 leaves) (Table 1). Student's *t*-tests revealed that this small increase in leaf number was significant. To monitor the floral transition at the molecular level, we investigated the floral meristem identity gene *APETALA1* (*AP1*). Whereas in LD-grown WT plants *AP1* transcript levels were increased by about sixfold from day 10 to day 21, the expression level rose more slowly in *atgrp7-1* (Figure 1a). Thus, in plants that lack *AtGRP7*, the transition to flowering is weakly delayed with long photoperiods, and is more strongly delayed with short photoperiods.

The response to inductive LDs is determined by the phase of the key regulator *CO* (Suarez-Lopez *et al.*, 2001). In *atgrp7-1* plants, diurnal oscillations of the *CO* transcript were very similar to WT, with respect to phase and amplitude, both in LDs and SDs (Figure 1b,c), which is consistent with their retaining a photoperiodic response.


Because the lack of *AtGRP7* in *atgrp7-1* had a relatively mild effect on flowering time, we examined in detail the expression of *AtGRP8*, which shows 77% sequence identity at the amino acid level and oscillates more or less in phase with *AtGRP7*.

Notably, whereas the *AtGRP7* transcript was undetectable, as expected (Fu *et al.*, 2007), the *AtGRP8* transcript level was elevated in *atgrp7-1* plants compared with WT (Figure 1d). To be able to distinguish between the highly similar *AtGRP7* and *AtGRP8* proteins, anti peptide antibodies specifically

Table 1 The *atgrp7-1* mutant is late-flowering

	leaf number	<i>n</i>	<i>P</i>
SD			
Col	54.82 ± 4.65	34	0.000012
<i>grp7-1</i>	61.57 ± 5.92	30	
LD			
Col	12.91 ± 1.54	34	0.000011
<i>grp7-1</i>	15.38 ± 2.30	32	

Leaf number of short-day (SD) and long-day (LD) grown *atgrp7-1* and wild-type (WT) plants ± SD at an inflorescence height of 0.5 cm: the total number of plants *n* and the *P* value determined by a Student's *t*-test are indicated. A representative experiment of five independent replicates with *n* ≥ 30 each is shown.



directed against *AtGRP7* or *AtGRP8* were generated (see Experimental procedures). The specificity of the antibodies was tested using recombinant *AtGRP7* and *AtGRP8*, respectively (Figure 1e). Weak cross-reactivity was only observed for the *AtGRP7* antibody when as much as 2 µg of pure recombinant *AtGRP8* protein was blotted. Whereas *AtGRP7* protein was absent in *atgrp7-1* plants, the *AtGRP8* protein level was elevated compared with WT (Figures 1e and S1). This indicates that the release from negative regulation in the absence of *AtGRP7* leads to higher *AtGRP8* accumulation.

Figure 1. Characterization of the *atgrp7-1* T-DNA insertion line.

(a) Developmental changes in *AP1* levels in long-day (LD) grown Col-0 and *atgrp7-1* plants.

Plants were harvested at days 10, 16, 19 and 21 after the release from stratification. *AP1* levels were assayed with real-time PCR with respect to the *PTB* reference gene. The relative expression in wild-type (WT) plants at day 10 was set to 1.

(b, c) Diurnal *CO* transcript pattern in *atgrp7-1*.

Col-0 WT and *atgrp7-1* mutants were harvested at 4-h intervals on day 12 (LD) (b) and day 19 (short-day grown, SD) (c), respectively. *CO* levels were assayed with real-time PCR with respect to the *PTB* reference gene. The relative expression in WT at zeitgeber time 0 (zt0; lights on) was set to 1.

(d) The *atgrp7-1* mutant and WT plants were harvested at zt12. The RNA gel blot was hybridized with the gene-specific *AtGRP7* probe (left), and with the gene-specific *AtGRP8* probe (right). The ethidium-bromide stained gel confirms equal loading.

(e) The immunoblot with 40 µg of total protein was incubated with the antibodies against *AtGRP7* (left) and *AtGRP8* (right). To demonstrate the preference of the antibodies, recombinant *AtGRP7* and *AtGRP8* proteins were loaded in the quantities indicated. Incubation with an LHCP antibody served as a loading control.

Thus, it remains possible that the elevated *AtGRP8* levels may mask part of the effect of *AtGRP7* on flowering time. To substantiate the floral-promotive function of *AtGRP7* we decided to analyze flowering time in transgenic plants with reduced levels of *AtGRP7* and *AtGRP8* on the one hand, and with constitutively elevated *AtGRP7* levels on the other hand.

Molecular analysis of *AtGRP7* and *AtGRP8* RNAi lines

To obtain transgenic Arabidopsis plants with reduced *AtGRP7* or *AtGRP8* expression, we performed RNA interference using the pKannibal system encoding a Cauliflower mosaic virus (CaMV) promoter-driven hairpin (hp) RNA, consisting of an inverted repeat of the gene fragment separated by the PdK intron (Wesley *et al.*, 2001). Two types of constructs were designed: short (s) constructs comprising only the part encoding the N-terminal RNA recognition motif of *AtGRP7* or *AtGRP8*, and long (l) constructs comprising additionally the part encoding the glycine-rich C terminus (Figure 2). The short construct directed against *AtGRP7* could not be stably propagated in *Agrobacterium tumefaciens*, and thus was not used further (not shown).

Transformed plants were selected on phosphinotricin, and were checked for the presence of the transgene. Seedlings harboring the intact hp constructs were raised to maturity, and were then surveyed for endogenous RNA and protein levels in the next generation. To test whether the simultaneous presence of *AtGRP7i* and *AtGRP8i* hp constructs would further reduce the protein levels, crosses were performed between selected *AtGRP7i* and *AtGRP8i* lines (Table S1).

AtGRP7 and *AtGRP8* transcript levels were monitored in LD-grown plants harvested at zt12 (zeitgeber time 12, 12 h after lights on), with gene- and strand-specific probes derived from their 5'-untranslated regions (UTRs) that do not cross-hybridize with the transgenic mRNA (Figures 3




Figure 2. Map of the *AtGRP7* and *AtGRP8* RNAi constructs. Numbers indicate nucleotide positions relative to the ATG start codon. (a) Construct *AtGRP7i-l* comprises 523 bps, including 22 nucleotides of the 5' untranslated region (UTR), the RRM and the glycine stretch. (b) Construct *AtGRP8i-s* comprises 218 bps spanning the RNA recognition motif (RRM). (c) Construct *AtGRP8i-l* comprises 496 bps spanning the RRM and part of the glycine stretch.




Figure 3. Molecular characterization of the *AtGRP7i* and *AtGRP8i* lines. Plants harbouring the Cambia (C) vector, the constructs *AtGRP7i-l*, *AtGRP8i-l* or *AtGRP8i-s*, and the offspring of crosses of selected lines (*AtGRP7i* × *8i*) were grown on phosphinotricin-containing plates along with Col-0 wild-type (WT) plants, and were harvested at zt12. The RNA gel blot was hybridized with the gene-specific *AtGRP7* probe (a), gene-specific *AtGRP8* probe (c) and *ACTIN* to confirm equal loading (b, d). Immunoblots of the corresponding total protein extract were incubated with antipeptide antibodies against *AtGRP7* (e), *AtGRP8* (g) and an antibody directed against LHCP as the loading control (f, h).

and S1). In *AtGRP7i-l* lines the endogenous *AtGRP7* transcript was strongly reduced compared with WT plants and Cambia (C) plants harboring the empty vector (Figure 3a). The *AtGRP8* transcript was also weakly affected (Figure 3c). Similarly in *AtGRP8i-l* and *AtGRP8i-s* lines, the endogenous

AtGRP8 transcript was almost undetectable (Figure 3c), and the *AtGRP7* transcript was reduced (Figure 3a). In all *AtGRP7i-l* lines, the *AtGRP7* protein level was strongly reduced or almost undetectable compared with WT and Cambia lines (Figure 3e), and in the *AtGRP8i* lines, the *AtGRP8* protein was almost undetectable (Figures 3g and S1). Again, in the *AtGRP7i-l* lines, the *AtGRP8* protein was only weakly affected, or remained at WT levels, whereas in the *AtGRP8i* lines the *AtGRP7* protein level was clearly lower than in WT or Cambia lines (Figure 3g). Taken together, these data suggest that the *AtGRP7i* or *AtGRP8i* constructs are effective against endogenous *AtGRP7* or *AtGRP8*, respectively, but also show some cross-regulation. The fact that in the *AtGRP8i* plants the *AtGRP7* protein is also strongly affected, whereas in the *AtGRP7i* plants the *AtGRP8* protein remains more or less at WT levels may indicate that the *AtGRP8i* constructs are more effective against *AtGRP7* than vice versa. Alternatively, the effect of the *AtGRP7i* construct on *AtGRP8* may be partially masked by enhanced *AtGRP8* accumulation as a consequence of *AtGRP7* down-regulation. In the crosses containing both hp constructs simultaneously, a further reduction of the *AtGRP7* and *AtGRP8* levels was only observed in line *AtGRP7i* × *8i*-21, compared with the parental lines (Figure 3e,g).

AtGRP7i and *AtGRP8i* plants are late flowering

Lines of all three genotypes *AtGRP7i-l*, *AtGRP8i-l* and *AtGRP8i-s*, and of the respective crosses, developed more leaves before flowering than WT or Cambia plants in SDs (Figure 4a). Also, in LDs, the RNAi lines formed between one and three leaves more than the control plants (Figure 4b). One-way ANOVA followed by a post-hoc Dunnett test showed that these differences are statistically significant (Table S2). Taken together, in the RNAi lines the delay in the transition to flowering is more pronounced than in the *atgrp7-1* mutant. Again, the late-flowering phenotype is more pronounced in LDs than in SDs.

Vernalization response of the RNAi and *atgrp7-1* plants

The retention of a photoperiodic response is a feature of mutants in the autonomous pathway that flower late in LDs, like mutants in the photoperiodic pathway, but flower even later in SDs, i.e. irrespective of the photoperiod. Additionally, their late-flowering phenotype can be corrected by vernalization. To determine whether the RNAi lines and *atgrp7-1* also show this feature, their response to extended periods of cold in the early seedling stage was assessed (see Experimental procedures). Again, non-vernalized RNAi lines and *atgrp7-1* flowered with more leaves than WT or C plants (Figure 5 and Table S3). After vernalization, RNAi lines flowered with a similar leaf number as the WT plants. *atgrp7-1* mutants flowered even at a slightly reduced leaf




Figure 4. Flowering time of *AtGRP7* and *AtGRP8* lines grown in short-day (SD) and long-day (LD) conditions.

(a) Wild type (WT), Cambia-1 (C-1), *AtGRP7*-148 and *AtGRP7*-168, *AtGRP8*-171, *AtGRP7*x8i-21 and *AtGRP7*x8i-50 (left), and WT, C-5, *AtGRP7*-11 and *AtGRP7*-154, *AtGRP8*-114, *AtGRP8*-s7, *AtGRP7*x8i-58 and *AtGRP7*x8i-109 (right), respectively, were grown in SDs.

(b) WT, C-5, *AtGRP7*-11, *AtGRP7*-154 and *AtGRP8*-114 and *AtGRP8*-s7 (left) and WT, C-5, *AtGRP7*x8i-21 and *AtGRP7*x8i-50 (right), respectively, were grown in LDs.

The number of rosette leaves (LN) produced at bolting is given as means \pm SD.

An ANOVA followed by a Dunnett test was performed to show statistical significance (see Table S2).




Figure 5. Effect of vernalization on flowering time of wild-type (WT), Cambia-5 (C-5), *AtGRP7* and *AtGRP8* lines and *atgrp7-1*.

Flowering time was measured as the number of rosette leaves (LN) produced at bolting in non-vernalized plants (white bar) and vernalized plants (black bar). Values represent the means \pm SD (see Table S3).

number compared with WT. Thus, vernalization completely overcomes the late-flowering phenotype of the RNAi lines and *atgrp7-1* mutant.

Influence of *AtGRP7* on *FLC*

The autonomous pathway acts in parallel with vernalization to constitutively repress *FLC* (Michaels and Amasino, 2001). Because vernalization bypasses the requirement of *AtGRP7* in the RNAi lines and the *atgrp7-1* mutant, we investigated whether the floral promotive effect of *AtGRP7* was mediated by *FLC*.

To this end, we first determined *FLC* levels by real-time PCR (Figure 6a). In the RNAi lines and in *atgrp7-1*, *FLC* was elevated by more than fivefold relative to WT or C plants. Vernalization strongly reduced the *FLC* levels in WT. Also, in the RNAi lines and *atgrp7-1*, *FLC* levels were reduced to a basal level similar to WT and C plants (Figure S2). Concomitantly, the floral integrator *SOC1*, which is negatively controlled by *FLC*, was upregulated upon vernalization both in WT and the RNAi lines (not shown). Thus, plants with reduced *AtGRP7* levels or complete loss of *AtGRP7* remain fully responsive to vernalization, indicating that *FLC* down-regulation in response to cold treatment does not require *AtGRP7*.

If the late flowering effect seen in plants with reduced *AtGRP7* levels entirely resulted from a lack of *FLC* repression, it should be suppressed in a genetic background lacking *FLC*. Therefore, we crossed the late-flowering RNAi lines *AtGRP7*x8i-21 and *AtGRP7*x8i-50 with *flc-3* mutants lacking active *FLC* because of a 107-bp deletion around the ATG (Michaels and Amasino, 1999). In the F_2 generation we identified lines homozygous for both the RNAi construct and the *flc-3* allele, by monitoring phosphinotricin resistance and PCR genotyping. Flowering time was investigated in the F_3 generation. Again, *AtGRP7*x8i-21 and *AtGRP7*x8i-50 flowered later than WT and C plants both in SDs and LDs (Figure 6b,c and Table S4). This late-flowering phenotype was eliminated in *AtGRP7*x8i-*flc-3* plants, which flowered with a leaf number comparable with *flc-3*. As previously noted, *flc-3* flowered with fewer leaves than WT in SDs (Michaels and Amasino, 2001). These data indicate that the lack of *AtGRP7* does not manifest itself in the absence of *FLC*.

Constitutive overexpression of *AtGRP7* promotes flowering

Because of the partial off-target effect of the *AtGRP7*i and *AtGRP8*i hp constructs on the *AtGRP8* and *AtGRP7* abundance, respectively, and the relief of *AtGRP8* repression in the *atgrp7-1* T-DNA line, the floral promotive effect could only be assigned tentatively to *AtGRP7*. Therefore, an effect of ectopic *AtGRP7* overexpression on flowering time was investigated in a complementary approach. The *AtGRP7*-ox lines D and G express highly elevated levels of *AtGRP7* protein, resulting in strongly reduced *AtGRP8* protein levels (Figure 7a). In SDs, these plants formed about 10 leaves less than WT at bolting (Figure 7b). This clearly shows that *AtGRP7* by itself is able to promote flowering. In the *At*




Figure 6. The effect of AtGRP7 on flowering time is largely mediated by *FLC*. (a) *FLC* expression in 6-week-old short-day (SD) grown Cambia-5 (C-5), *AtGRP8i-s7* and *AtGRP7ix-109* lines (left), and wild type (WT) and *atgrp7-1* (right). Five plants were pooled for RNA isolation and *FLC* levels were determined by real-time PCR in duplicates for two biological replicates. Expression levels were normalized to *PPR*, and the relative *FLC* level in WT was set to 1. (b, c) The *flc-3* null allele eliminates the late-flowering phenotype of *AtGRP7ix8i*. WT, C-5 and the parental lines *AtGRP7ix8i-21*, *AtGRP7ix8i-50* and *flc-3* were grown in parallel with *AtGRP7ixAtGRP8i-21 flc-3* and *AtGRP7ix8i-50 flc-3* in SDs (b) or LDs (c). The numbers 1, 2 and 3 represent F₃ plants of independent crosses. The leaf number (LN) at bolting ± SD is shown. This experiment was performed twice with similar results.

GRP7-ox lines, *FLC* was downregulated to 10–20% of the level in WT (Figure 7c). After vernalization, *AtGRP7-ox* and WT plants formed about the same number of leaves at bolting (Table S5). In LDs, the *AtGRP7-ox* lines flowered only




Figure 7. Influence of *AtGRP7* overexpression on flowering time. (a) *AtGRP7* and *AtGRP8* protein levels were analyzed in short-day (SD) grown *AtGRP7-ox* and wild-type (WT) plants using specific antibodies. The reaction with an LHCP antibody or the unspecific reaction of the *AtGRP8* antibody with the small subunit of Rubisco served as the loading control. (b) *AtGRP7* overexpression causes early flowering in SDs. Independent *AtGRP7-ox* lines were grown along with WT plants. The leaf number (LN) at the onset of bolting is shown ± SD. (c) *FLC* levels are lower in *AtGRP7-ox* plants. *FLC* levels were determined by real-time PCR, as described in the legend to Fig. 6a, and were normalized to *PTB*. The level in WT was set to 1.

slightly earlier than WT plants (Table S5), indicating that the promotive effect of *AtGRP7* has a stronger impact under non-inductive photoperiods.

To address the question of whether *AtGRP7* may act via other autonomous pathway components, we analyzed selected transcripts encoding proteins associated with RNA metabolism, and their respective alternative splice

forms (<http://www.plantgdb.org/ASIP/>) in the AtGRP7 gain-of-function and loss-of-function plants. For *FPA* encoding a protein with three RRM (Schomburg *et al.*, 2001), a transcript retaining the 129-nt (nucleotide) intron 4 was detected, in addition to the spliced mRNA (Figure S3a). The ratio between the longer and shorter variants, and their levels, were not changed in the *atgrp7-1* T-DNA line or representative RNAi lines compared with WT or the C line. Also, constitutive overexpression of AtGRP7 did not affect intron 4 retention. Furthermore, retention of a 99-nt intron located 40-nt downstream of the stop codon in the *FPA* 3'-UTR was found for all genotypes (Figure S3b).

For the 3' end processing factor, *FY*, the use of an alternative acceptor site at intron 8 leads to the inclusion of four additional nt in intron 8, causing a frameshift and premature in-frame termination codon. Using flanking primers, a fragment of the same size as in WT and C lines was detected on a polyacrylamide gel for the RNAi lines, the *atgrp7-1* T-DNA line and the AtGRP7-ox line (Figure S3d). Also for *FCA*, no obvious variations were detected for the *FCA-γ* mRNA, encoding active FCA protein, and the alternatively spliced, prematurely polyadenylated *FCA-β* form (not shown). The autonomous pathway component LUMINIDEPENDENS (LD) is a homeodomain protein that may interact with DNA or RNA (Aukerman *et al.*, 1999). The steady-state abundance of LD is not significantly different from WT levels in the RNAi lines, the *atgrp7-1* mutant and the AtGRP7-ox lines (Figure S3e).

Discussion

The RNA-binding protein AtGRP7 promotes flowering

Here, we demonstrate the participation of a small glycine-rich RNA binding protein with a single RRM in the regulation of flowering time in Arabidopsis. The loss of function of AtGRP7 delays the transition to flowering, whereas the gain of function through ectopic overexpression of AtGRP7 promotes flowering.

The *atgrp7-1* T-DNA line flowers later than WT plants. In LDs, the delay is small, yet statistically significant. These data are supported by the observation that the rise in *AP1*, indicative of floral induction, occurs later in development than in the WT. The retardation becomes more pronounced in SDs, with a reduced impact of the photoperiodic inductive signal.

The well-described prominent regulation of AtGRP7 by the circadian clock may have pointed to a role in the photoperiodic pathway. Photoperiodic mutants like *co* or *gi*, however, very strongly delay flowering in LDs, in contrast to *atgrp7-1* (Koorneef *et al.*, 1991). Thus, AtGRP7 is dispensable for measuring day length. Accordingly, diurnal oscillations of the key photoperiodic regulator *CO*, the presence of

which during the light phase initiates floral transition, are similar to WT.

The subtle effect that the loss of AtGRP7 has on the transition to flowering presumably has precluded its identification as a floral promoter in conventional screens for flowering time mutants. Alternatively, one may envisage that AtGRP8, the closest homolog of AtGRP7 that shares much of its regulatory properties, may act redundantly. In fact, by taking advantage of specific antipeptide antibodies, we uncovered an elevated AtGRP8 protein level in the *atgrp7-1* mutant compared with WT. Also, the AtGRP8 transcript level is elevated in the absence of AtGRP7 protein. This is consistent with our previous observation that AtGRP7 negatively regulates AtGRP8 oscillations (Schöning *et al.*, 2007; Staiger *et al.*, 2003b). Thus, it remains possible that the elevated AtGRP8 level in *atgrp7-1* may partly obscure the loss of AtGRP7 as a result of overlapping functions. Therefore, we aimed to generate plants with reduced AtGRP7 and AtGRP8 levels through RNA interference, as well as plants ectopically overexpressing AtGRP7.

Using hp constructs targeted against the respective RRM, or the entire coding region, we obtained a series of transgenic plants in which AtGRP7 and AtGRP8 are down-regulated to a varying degree. These RNAi lines showed a more pronounced late-flowering phenotype in LDs, and particularly in SDs, compared with *atgrp7-1*. On the other hand, plants constitutively overexpressing AtGRP7 flower with 10 leaves less than WT plants in SDs. In LDs, the advance is only very small, but is nevertheless significant. These data clearly indicate that AtGRP7 by itself promotes flowering, as AtGRP8 is almost completely downregulated in AtGRP7-ox plants as a result of the generation of an unproductively spliced transcript with a premature termination codon, which rapidly decays via an UPF1- and UPF3-dependent pathway (Schöning *et al.*, 2007).

Nevertheless, it remains possible that AtGRP8 also promotes floral transition to some degree. For example, the flowering of *AtGRP7-148*, which displays WT levels of AtGRP8 but almost no AtGRP7, is only weakly retarded, and *AtGRP7ix8i-50*, with a weak reduction of AtGRP7, but strong reduction of AtGRP8, flowers very late. So plants with selective reduction of AtGRP8 or strong ectopic overexpression of AtGRP8 will be needed to unequivocally resolve this issue.

AtGRP7 is a novel autonomous pathway component

Because the lack of AtGRP7 does not affect the photoperiodic response, we investigated whether AtGRP7 may share other features with components of the autonomous pathway. Three lines of evidence indeed place AtGRP7 in the autonomous pathway.

(i) The late flowering of AtGRP7 loss-of-function plants in SDs correlates with an elevated *FLC* level, whereas the early

flowering of *AtGRP7* gain-of-function plants correlates with a reduction in *FLC*.

(ii) Whereas *FLC* is also a target of the vernalization pathway, in *AtGRP7* loss-of-function plants, prolonged cold treatment in the young seedling stage leads to a reduction of *FLC*, to a level similar to WT plants. Thus, vernalization downregulates *FLC* in the absence of *AtGRP7*, and, in fact, vernalization completely overrides the late-flowering phenotype of *atgrp7-1* and the RNAi lines in SDs, as previously observed for mutants in the autonomous pathway (Koornneef *et al.*, 1991; Michaels and Amasino, 2001; Sheldon *et al.*, 2000).

(iii) Moreover, crosses between RNAi lines and *f1c-3* mutants lacking active *FLC* provide genetic evidence that *AtGRP7* influences the floral transition through *FLC*, as the plants flower with a similar number of leaves as *f1c-3* mutants. However, *AtGRP7* may also have *FLC*-independent effects, as observed for other flowering-time genes (Doyle *et al.*, 2005). So far we have not detected consistent changes in the alternative splicing pattern and/or steady-state abundance of transcripts encoding other autonomous pathway proteins implicated in RNA processing. Consistent with this, transcript profiling of LD-grown *AtGRP7*-ox lines and WT plants using the ATH1 GeneChip did not reveal significant changes in *FCA*, *FY* or *LD* steady-state abundance, whereas the *FLC* levels were lowered in *AtGRP7*-ox plants (C. Streitner, F. Rudolf and D. Staiger, unpublished data). Thus, based on the present data, impaired expression of *FCA*, *FY*, *FPA* or *LD* does not seem to be the major cause for the influence of *AtGRP7* on floral transition. To further define the position of *AtGRP7* within the autonomous pathway, however, we must analyze whether a lack of *AtGRP7*, and/or *AtGRP8*, impacts on the late-flowering phenotype of the respective autonomous pathway components in double mutant combinations. For the other RNA binding proteins of the autonomous pathway, it had been suggested that they act in parallel, and ultimately control *FLC* levels independently (Quesada *et al.*, 2005).

RNA binding proteins in the regulation of flowering time

Several autonomous pathway genes encode RNA binding proteins or RNA processing factors, comprising different types of RNA binding modules and associated domains implicated in protein–protein interaction (Quesada *et al.*, 2005). Although these loci had been identified early on because of their late-flowering phenotype when mutated (Koornneef *et al.*, 1991), it is now thought that they regulate additional processes in the plant. In a microarray analysis, several transcripts with abnormal expression pattern in *fca* mutants were identified in which expression was also changed in *fy* mutants (Marquardt *et al.*, 2006). This points to a more general function of *FCA* and *FY*.

Other proteins involved in various aspects of RNA metabolism have also been associated recently with flowering. The *ABA HYPERSENSITIVE 1* (*abh1*) mutant defective in the large subunit of the CAP-binding complex, CBP80, flowers early in SDs and LDs (Bezerra *et al.*, 2004; Hugouvieux *et al.*, 2001). The *HUA2* protein involved in the processing of *AGAMOUS* intron 2 is required for the correct regulation of *FLC* and the related repressors *FLM1/MAF1* or *MAF2* (Cheng *et al.*, 2003; Doyle *et al.*, 2005). Mutants defective in the tetratricopeptide repeat protein AT PRP39-1, with similarity to a yeast pre-mRNA processing protein, are late flowering (Wang *et al.*, 2007). Notably, all these RNA binding proteins, like the well-known autonomous pathway components and *AtGRP7*, affect flowering time in a large part by influencing *FLC* levels. Based on the domain structure of these proteins, it has been inferred that they impact on *FLC* levels by a post-transcriptional mechanism (Kuhn *et al.*, 2007; Quesada *et al.*, 2005). Recently, however, the downregulation of *FLC* abundance by *FCA* has been shown to be dependent on *FLD* (Liu *et al.*, 2007). Nascent *FLC* transcripts accumulate to higher levels in *fca* and *fld* mutants, suggesting that *FCA* and *FLD* actually silence *FLC* expression at the transcriptional level through H3K4 demethylation.

AtGRP7 was originally identified on the basis of its clock regulation and cold responsiveness (Carpenter *et al.*, 1994; Heintzen *et al.*, 1994). The *AtGRP7* feedback loop is a molecular slave oscillator within the circadian system, regulating both rhythmic and non-rhythmic target transcripts (Heintzen *et al.*, 1997; Schöning *et al.*, 2007). Furthermore, *AtGRP7* is involved in pathogen defense (Fu *et al.*, 2007), and has been implicated in ABA and stress signaling, as an *atgrp7* mutant accumulated higher levels of the ABA- and stress-inducible *RD29A* transcript (Cao *et al.*, 2006).

Presumably *AtGRP7* plays a more general role in pre-mRNA processing, and the flowering phenotype we identify here reflects the dependence of floral transition on fine-tuned *FLC* levels. The relatively mild phenotype, particularly in LDs, may have precluded the identification of *AtGRP7* in screens for flowering-time genes so far.

It has long been suggested that the circadian clock is an integral part of the photoperiodic sensory device (Bünning, 1936). Experimental proof came from the identification of mutants in *Arabidopsis* with disturbances in both circadian and photoperiodic timekeeping (Fowler *et al.*, 1999; Hicks *et al.*, 1996; Park *et al.*, 1999; Schaffer *et al.*, 1998; Somers *et al.*, 1998; Staiger *et al.*, 2003a; Suarez-Lopez *et al.*, 2001; Wang and Tobin, 1998).

Following the observation that *FLC* lengthens the period of the circadian clock at 27°C, the autonomous pathway mutants *fca*, *ld* and *fve* have been investigated for clock phenotypes, and have been found to moderately increase the period of leaf movement rhythms (Edwards *et al.*, 2006; Salathia *et al.*, 2006).

Our results now establish a link between the slave oscillator component AtGRP7 operating downstream of the circadian clock and floral transition, through the autonomous pathway. This points to extensive crosstalk between the circadian system and the floral-promoting network, beyond photoperiodic timekeeping.

Experimental procedures

Plant materials

The *flc-3* mutant was kindly provided by Dr Amasino (Michaels and Amasino, 1999). The *atgrp7-1* T-DNA mutant was kindly provided by Drs Fu and Guo (Fu *et al.*, 2007). Seeds were obtained from the Nottingham Arabidopsis Stock Centre (<http://arabidopsis.info>).

Constructs for hairpin RNA-mediated silencing of AtGRP7 and AtGRP8

Fragments of *AtGRP7* and *AtGRP8* cDNAs were amplified using forward primers, while simultaneously introducing *XhoI* and *XbaI* sites, and reverse primers, introducing *Clal* and *EcoRI* sites. The amplification products were inserted into pTOPOII (Invitrogen, <http://www.invitrogen.com>), verified by sequencing and moved to pKannibal in the antisense orientation after *XhoI-EcoRI* restriction, and in the sense orientation after *Clal-XbaI* restriction (Wesley *et al.*, 2001). The entire cassette was cut out using *NotI*, blunted by Klenow fill-in and subcloned into *SmaI*-cut pCAMBIA3300.

The following primers were used: for construct *AtGRP7i-1*, comprising part of the 5'-UTR, the RRM and the glycine-rich part (from position -22 to position -504, relative to the ATG start codon), p7i-F (5'-CTCGAGTCTAGATCTTCTTT-3') and p7i-523R (5'-ATC-GATGAATTCGGTAACCTCC-3'); for constructs *AtGRP8i-s*, comprising the RRM (positions 13–231 relative to the ATG), p8i-F (5'-TTTCTAGACTCGAGTACCGG-3') and p8i-218R (5'-TTGAATTCA-TCGATGACACG-3'); for constructs *AtGRP8i-l*, comprising the RRM and the glycine-rich part (positions 13–507 relative to the ATG) p8i-F and p8i-496R (5'-TTATCGATGAATTCAGCGCC-3').

Transgenic plants

The RNAi constructs and the Cambia vector were introduced into *A. thaliana* L. *Columbia* by *Agrobacterium*-mediated vacuum infiltration (Bechthold *et al.*, 1993). Primary transformants were selected on plates with agar-solidified half-strength MS medium (Duchefa, <http://www.duchefa.com>) with 0.5% sucrose, adjusted to pH 5.7, and containing 25 mg l⁻¹ phosphinotricin.

The genomic DNA of transformants was isolated according to the protocol of the Wisconsin ko facility (Sussman *et al.*, 2000). The presence of the hp constructs was confirmed by PCR using primers for the *OCS* terminator (Table S6).

Determination of flowering time

Plants were grown in a randomized fashion on soil in LDs (16-h light) or SDs (8-h light) at 20°C in Percival AR66-L3 incubators (CLF Laboratories, <http://www.clf.de>). The flowering time was determined by counting the rosette leaves once the bolt was 0.5-cm tall. Mean values \pm SD were calculated.

For vernalization treatments, seeds were stratified at 4°C for 3 days in the dark. Germinated seeds were transferred to SDs at

20°C for 7 days, returned to 4°C in SDs for 46 days, and were subsequently transferred back to SDs at 20°C. Control plants were germinated at 4°C for 3 days and were immediately transferred to SDs at 20°C.

Statistical analysis of flowering time data

Statistical analysis was performed using STATISTICA 6.0 (<http://www.statsoft.com>). Mean values and standard deviation were calculated for each data set. $P < 0.05$ was considered significant. In experiments analyzing differences between two lines, a Student's *t*-test was used when normal distribution and homogeneity of variances were proven by the Kolmogorov–Smirnov and the *F*-test, respectively. If normal distribution was not given, the Mann–Whitney *U*-test was used to analyze the significance of differences. If a normal distribution, but not the homogeneity of variances, was given, Welch's *t*-test was used for analysis. When analysing means of more than two different samples, an ANOVA was performed. If significant differences were shown, the Dunnett test helped us to consider which samples were different from the control lines.

RNA analysis

Total RNA was isolated using the Trizol reagent. Hybridizations of RNA gel blots with ³²P-labeled gene-specific antisense probes covering the 5'-UTR of *AtGRP7* and *AtGRP8*, respectively, were performed as described by Heintzen *et al.* (1997).

For real-time PCR, duplicate samples were analyzed in an MJ research Opticon DNA Engine (<http://www.bio-rad.com>). Total RNA was treated with DNaseI and reverse-transcribed using Superscript II (Invitrogen). A 20-ng portion of retrotranscribed RNA was amplified with the Eppendorf Real MasterMix kit (Eppendorf, <http://www.eppendorf.com>) using an initial denaturation step of 2 min, followed by 45 cycles of 20 sec at 94°C, 30 sec at 60°C and 40 sec at 68°C. Threshold cycle (C_T) values were determined, and relative expression levels for the analyzed transcripts were calculated based on non-equal efficiencies for each primer pair (Czechowski *et al.*, 2004; Pfaffl, 2001). Data were normalized to a transcript encoding the translation initiation factor eIF-4A-1 (At3g13920), PTB (At3g01150) and PPR (At5g55840) (Czechowski *et al.*, 2005). The absence of amplification products from genomic DNA was confirmed in a non-retrotranscribed control.

For semiquantitative RT-PCR, retrotranscribed RNA was amplified with Taq Polymerase. To determine the linear range of amplification for each primer pair, samples were withdrawn after 24, 26, 28, 30, 32 and 34 cycles. PCR products were separated on agarose gels, or polyacrylamide gel in the case of *FY*, and were visualized by ethidium-bromide staining.

Generation of antibodies against AtGRP7 and AtGRP8, and protein analysis

Antibodies were raised against synthetic polypeptides spanning amino acids 22–31 of AtGRP7 and amino acids 20–29 of AtGRP8, respectively, which are divergent between the two proteins (Pineda Antikörper Service, <http://pineda-abservice.com>). The specificity of the antibodies for AtGRP7 and AtGRP8 was monitored by immunoblots against recombinant glutathione-S-transferase fusion proteins after the release of the AtGRP7 and AtGRP8 moiety by PreScission protease (GE Healthcare, <http://www.gehealthcare.com>) cleavage.

Protein extraction and immunoblots with chemiluminescence detection were performed as previously described (Heintzen *et al.*, 1997).

Acknowledgements

We thank K. Neudorf and E. Detring for expert technical assistance, and D. Balten and U. Büchner for care of the plants in the greenhouse. We are grateful to Dr R. Amasino for making available the *flc-3* mutant, and to Dr M. Guo for identifying the *atgrp7-1* mutant. We thank Dr Judith Gomez-Porras for her critical comments on the manuscript, and Drs George Coupland and Lars Hennig for useful discussions. The pKannibal vector was kindly provided by Drs Wesley, Helliwell and Waterhouse, CSIRO, Australia. We acknowledge the Nottingham Stock Centre for providing the seeds. This work was supported by the DFG (STA 653/2 and SFB 613), the Swiss National Foundation and the US National Institutes of Health Award No. 1R01AI069146-01A2 to JRA.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Figure S1. Molecular characterization of further *AtGRP7i* and *AtGRP8i* lines.

Figure S2. FLC levels in vernalized RNAi and *atgrp7-1* lines.

Figure S3. Influence of altered *AtGRP7* and *AtGRP8* levels on autonomous pathway components.

Table S1. Crosses between *AtGRP7i* and *AtGRP8i* lines.

Table S2. Flowering time of *AtGRP7i* and *AtGRP8i* lines in short-day (SD) and long-day (LD) growth conditions.

Table S3. Effect of vernalization on the flowering time of the wild type (WT), Cambia, *AtGRP7i* and *AtGRP8i* lines and *atgrp7-1*.

Table S4. Flowering time in crosses between *flc-3* and *AtGRP7ix8i* lines.

Table S5. Flowering time in *AtGRP7-ox* lines.

Table S6. Primers used for PCR.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

References

- Abe, M., Kobayashi, Y., Yamamoto, S., Daimon, Y., Yamaguchi, A., Ikeda, Y., Ichinoki, H., Notaguchi, M., Goto, K. and Araki, T. (2005) FD, a bZIP protein mediating signals from the floral pathway integrator FT at the shoot apex. *Science*, **309**, 1052–1056.
- An, H., Rousot, C., Suarez-Lopez, P. *et al.* (2004) CONSTANS acts in the phloem to regulate a systemic signal that induces photoperiodic flowering of *Arabidopsis*. *Development*, **131**, 3615–3626.
- Aukerman, M.J., Lee, I., Weigel, D. and Amasino, R.M. (1999) The *Arabidopsis* flowering-time gene LUMINIDEPENDENS is expressed primarily in regions of cell proliferation and encodes a nuclear protein that regulates LEAFY expression. *Plant J.* **18**, 195–203.
- Ausin, I., Alonso-Blanco, C., Jarillo, J.A., Ruiz-Garcia, L. and Martinez-Zapater, J.M. (2004) Regulation of flowering time by FVE, a retinoblastoma-associated protein. *Nat. Genet.* **36**, 162–166.
- Bastow, R., Mylne, J.S., Lister, C., Lippman, Z., Martienssen, R.A. and Dean, C. (2004) Vernalization requires epigenetic silencing of FLC by histone methylation. *Nature*, **427**, 164–167.
- Bechthold, N., Ellis, J. and Pelletier, G. (1993) In planta *Agrobacterium*-mediated gene transfer by infiltration of adult *Arabidopsis thaliana* plants. *Life Sci.* **316**, 1194–1199.
- Bezerra, I.C., Michaels, S.D., Schomburg, F.M. and Amasino, R.M. (2004) Lesions in the mRNA cap-binding gene ABA HYPERSENSITIVE 1 suppress FRIGIDA-mediated delayed flowering in *Arabidopsis*. *Plant J.* **40**, 112–119.
- Borner, R., Kampmann, G., Chandler, J., Gleissner, R., Wisman, E., Apel, K. and Melzer, S. (2000) A MADS domain gene involved in the transition to flowering in *Arabidopsis*. *Plant J.* **24**, 591–599.
- Bünning, E. (1936) Die endogene Tagesrhythmik als Grundlage der photoperiodischen Reaktion. *Ber. Dtsch. Bot. Ges.* **54**, 590–607.
- Cao, S., Jiang, L., Song, S., Jing, R. and Xu, G. (2006) AtGRP7 is involved in the regulation of abscisic acid and stress responses in *Arabidopsis*. *Cell Mol. Biol. Lett.* **11**, 526–535.
- Carpenter, C.D., Kreps, J.A. and Simon, A.E. (1994) Genes encoding glycine-rich *Arabidopsis thaliana* proteins with RNA-binding motifs are influenced by cold treatment and an endogenous circadian rhythm. *Plant Physiol.* **104**, 1015–1025.
- Cheng, Y., Kato, N., Wang, W., Li, J. and Chen, X. (2003) Two RNA binding proteins HEN4 and HUA1, act in the processing of AGAMOUS pre-mRNA in *Arabidopsis thaliana*. *Dev. Cell*, **4**, 53–66.
- Corbesier, L. and Coupland, G. (2006) The quest for florigen: a review of recent progress. *J. Exp. Bot.* **57**, 3395–3403.
- Corbesier, L., Vincent, C., Jang, S. *et al.* (2007) FT protein movement contributes to long-distance signaling in floral induction of *Arabidopsis*. *Science*, **316**, 1030–1033.
- Czechowski, T., Bari, R.P., Stitt, M., Scheible, W.R. and Udvardi, M.K. (2004) Real-time RT-PCR profiling of over 1400 *Arabidopsis* transcription factors: unprecedented sensitivity reveals novel root- and shoot-specific genes. *Plant J.* **38**, 366–379.
- Czechowski, T., Stitt, M., Altmann, T., Udvardi, M.K. and Scheible, W.R. (2005) Genome-wide identification and testing of superior reference genes for transcript normalization in *Arabidopsis*. *Plant Physiol.* **139**, 5–17.
- Doyle, M.R., Bizzell, C.M., Keller, M.R., Michaels, S.D., Song, J., Noh, Y.S. and Amasino, R.M. (2005) HUA2 is required for the expression of floral repressors in *Arabidopsis thaliana*. *Plant J.* **41**, 376–385.
- Edwards, K.D., Anderson, P.E., Hall, A., Salathia, N.S., Locke, J.C., Lynn, J.R., Straume, M., Smith, J.Q. and Millar, A.J. (2006) FLOWERING LOCUS C mediates natural variation in the high-temperature response of the *Arabidopsis* circadian clock. *Plant Cell*, **18**, 639–650.
- Fowler, S., Lee, K., Onouchi, H., Samach, A., Richardson, K., Morris, B., Coupland, G. and Putterill, J. (1999) GIGANTEA: a circadian clock-controlled gene that regulates photoperiodic flowering in *Arabidopsis* and encodes a protein with several possible membrane-spanning domains. *EMBO J.* **18**, 4679–4688.
- Fu, Z.Q., Guo, M., Jeong, B.R., Tian, F., Elthon, T.E., Cerny, R.L., Staiger, D. and Alfano, J.R. (2007) A type III effector ADP-ribosylates RNA-binding proteins and quells plant immunity. *Nature*, **447**, 284–288.
- He, Y., Michaels, S.D. and Amasino, R.M. (2003) Regulation of flowering time by histone acetylation in *Arabidopsis*. *Science*, **302**, 1751–1754.
- Heintzen, C., Melzer, S., Fischer, R., Kappeler, S., Apel, K. and Staiger, D. (1994) A light- and temperature-entrained circadian clock controls expression of transcripts encoding nuclear proteins with homology to RNA-binding proteins in meristematic tissue. *Plant J.* **5**, 799–813.

- Heintzen, C., Nater, M., Apel, K. and Staiger, D. (1997) AtGRP7, a nuclear RNA-binding protein as a component of a circadian-regulated negative feedback loop in *Arabidopsis thaliana*. *Proc. Natl Acad. Sci. USA*, **94**, 8515–8520.
- Hicks, K.A., Millar, A.J., Carre, I.A., Somers, D.E., Straume, M., Meeks-Wagner, D.R. and Kay, S.A. (1996) Conditional circadian dysfunction of the *Arabidopsis* early-flowering 3 mutant. *Science*, **274**, 790–792.
- Hugouvieux, V., Kwak, J.M. and Schroeder, J.I. (2001) An mRNA cap binding protein, ABH1, modulates early abscisic acid signal transduction in *Arabidopsis*. *Cell*, **106**, 477–487.
- Jaeger, K.E. and Wigge, P.A. (2007) FT protein acts as a long-range signal in *Arabidopsis*. *Curr. Biol.* **17**, 1050–1054.
- Kobayashi, Y. and Weigel, D. (2007) Move on up, it's time for change: mobile signals controlling photoperiod-dependent flowering. *Genes Dev.* **21**, 2371–2384.
- Koornneef, M., Hanhart, C.J. and van der Veen, J.H. (1991) A genetic and physiological analysis of late flowering mutants in *Arabidopsis thaliana*. *Mol. Gen. Genet.* **229**, 57–66.
- Kuhn, J.M., Breton, G. and Schroeder, J.I. (2007) mRNA metabolism of flowering-time regulators in wild-type *Arabidopsis* revealed by a nuclear cap binding protein mutant, abh1. *Plant J.* **50**, 1049–1062.
- Lee, H., Suh, S.S., Park, E., Cho, E., Ahn, J.H., Kim, S.G., Lee, J.S., Kwon, Y.M. and Lee, I. (2000) The AGAMOUS-LIKE 20 MADS domain protein integrates floral inductive pathways in *Arabidopsis*. *Genes Dev.* **14**, 2366–2376.
- Lim, M.H., Kim, J., Kim, Y.S., Chung, K.S., Seo, Y.H., Lee, I., Hong, C.B., Kim, H.J. and Park, C.M. (2004) A new *Arabidopsis* gene, *FLK*, encodes an RNA binding protein with K homology motifs and regulates flowering time via *FLOWERING LOCUS C*. *Plant Cell*, **16**, 731–740.
- Liu, F., Quesada, V., Crevillen, P., Baurle, I., Swiezewski, S. and Dean, C. (2007) The *Arabidopsis* RNA-binding protein FCA requires a lysine-specific demethylase 1 homolog to downregulate FLC. *Mol. Cell*, **28**, 398–407.
- Macknight, R., Bancroft, I., Page, T.L. *et al.* (1997) *FCA*, a gene controlling flowering time in *Arabidopsis*, encodes a protein containing RNA-binding domains. *Cell*, **89**, 737–745.
- Macknight, R., Duroux, M., Laurie, R., Dijkwel, P., Simpson, G. and Dean, C. (2002) Functional significance of the alternative transcript processing of the *Arabidopsis* floral promoter FCA. *Plant Cell*, **14**, 877–888.
- Marquardt, S., Boss, P.K., Hadfield, J. and Dean, C. (2006) Additional targets of the *Arabidopsis* autonomous pathway members, FCA and FY. *J. Exp. Bot.* **57**, 3379–3386.
- Mathieu, J., Warthmann, N., Kuttner, F. and Schmid, M. (2007) Export of FT protein from phloem companion cells is sufficient for floral induction in *Arabidopsis*. *Curr. Biol.* **17**, 1055–1060.
- Michaels, S.D. and Amasino, R.M. (1999) FLOWERING LOCUS C encodes a novel MADS domain protein that acts as a repressor of flowering. *Plant Cell*, **11**, 949–956.
- Michaels, S.D. and Amasino, R.M. (2001) Loss of FLOWERING LOCUS C activity eliminates the late-flowering phenotype of FRIGIDA and autonomous pathway mutations but not responsiveness to vernalization. *Plant Cell*, **13**, 935–941.
- Michaels, S.D., Himelblau, E., Kim, S.Y., Schomburg, F.M. and Amasino, R.M. (2005) Integration of flowering signals in winter-annual *Arabidopsis*. *Plant Physiol.* **137**, 149–156.
- Park, D.H., Somers, D.E., Kim, Y.S., Choy, Y.H., Lim, H.K., Soh, M.S., Kim, H.J., Kay, S.A. and Nam, H.G. (1999) Control of circadian rhythms and photoperiodic flowering by the *Arabidopsis* GIGANTEA gene. *Science*, **285**, 1579–1582.
- Pfaffl, M.W. (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* **29**, e45.
- Putterill, J., Laurie, R. and Macknight, R. (2004) It's time to flower: the genetic control of flowering time. *Bioessays*, **26**, 363–373.
- Quesada, V., Macknight, R., Dean, C. and Simpson, G.G. (2003) Autoregulation of FCA pre-mRNA processing controls *Arabidopsis* flowering time. *EMBO J.* **22**, 3142–3152.
- Quesada, V., Dean, C. and Simpson, G.G. (2005) Regulated RNA processing in the control of *Arabidopsis* flowering. *Int. J. Dev. Biol.* **49**, 773–780.
- Salathia, N., Davis, S.J., Lynn, J.R., Michaels, S.D., Amasino, R.M. and Millar, A.J. (2006) FLOWERING LOCUS C-dependent and -independent regulation of the circadian clock by the autonomous and vernalization pathways. *BMC Plant Biol.* **6**, 10.
- Samach, A., Onouchi, H., Gold, S.E., Ditta, G.S., Schwarz-Sommer, Z., Yanofsky, M.F. and Coupland, G. (2000) Distinct roles of CONSTANS target genes in reproductive development of *Arabidopsis*. *Science*, **288**, 1613–1636.
- Schaffer, R., Ramsay, N., Samach, A., Putterill, J., Carre, I.A. and Coupland, G. (1998) The late elongated hypocotyl mutation of *Arabidopsis* disrupts circadian rhythms and the photoperiodic control of flowering. *Cell*, **93**, 1219–1229.
- Schomburg, F.M., Patton, D.A., Meinke, D.W. and Amasino, R.M. (2001) *FPA*, a gene involved in floral induction in *Arabidopsis*, encodes a protein containing RNA-recognition motifs. *Plant Cell*, **13**, 1427–1436.
- Schöning, J.C., Streitner, C., Page, D.R., Hennig, S., Uchida, K., Wolf, E., Furuya, M. and Staiger, D. (2007) Autoregulation of the circadian slave oscillator component AtGRP7 and regulation of its targets is impaired by a single RNA recognition motif point mutation. *Plant J.* **52**, 1119–1130.
- Sheldon, C.C., Burn, J.E., Perez, P.P., Metzger, J., Edwards, J.A., Peacock, W.J. and Dennis, E.S. (1999) The FLF MADS box gene: a repressor of flowering in *Arabidopsis* regulated by vernalization and methylation. *Plant Cell*, **11**, 445–458.
- Sheldon, C.C., Rouse, D.T., Finnegan, E.J., Peacock, W.J. and Dennis, E.S. (2000) The molecular basis of vernalization: the central role of FLOWERING LOCUS C (FLC). *Proc. Natl Acad. Sci. USA*, **97**, 3753–3758.
- Shi, Y., Lan, F., Matson, C., Mulligan, P., Whetstone, J.R., Cole, P.A., Casero, R.A. and Shi, Y. (2004) Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. *Cell*, **119**, 941–953.
- Simpson, G.G. and Dean, C. (2002) *Arabidopsis*, the rosetta stone of flowering time? *Science*, **296**, 285–289.
- Simpson, G.G., Dijkwel, P.P., Quesada, V., Henderson, I. and Dean, C. (2003) FY is an RNA 3' end-processing factor that interacts with FCA to control the *Arabidopsis* floral transition. *Cell*, **113**, 777–787.
- Somers, D.E., Webb, A.A., Pearson, M. and Kay, S.A. (1998) The short-period mutant, *toc1-1*, alters circadian clock regulation of multiple outputs throughout development in *Arabidopsis thaliana*. *Development*, **125**, 485–494.
- Staiger, D., Allenbach, L., Salathia, N., Fiechter, V., Davis, S.J., Millar, A.J., Chory, J. and Fankhauser, C. (2003a) The *Arabidopsis* *SRR1* gene mediates phyB signaling and is required for normal circadian clock function. *Genes Dev.* **17**, 256–268.
- Staiger, D., Zecca, L., Wiczeorek, K., Kirk, D.A., Apel, K. and Eckstein, L. (2003b) The circadian clock regulated RNA-binding protein AtGRP7 autoregulates its expression by influencing alternative splicing of its own pre-mRNA. *Plant J.* **33**, 361–371.
- Suarez-Lopez, P., Wheatley, K., Robson, F., Onouchi, H., Valverde, F. and Coupland, G. (2001) CONSTANS mediates between the circadian clock and the control of flowering in *Arabidopsis*. *Nature*, **410**, 1116–1120.

- Sung, S. and Amasino, R.M.** (2004) Vernalization in *Arabidopsis thaliana* is mediated by the PHD finger protein VIN3. *Nature*, **427**, 159–164.
- Sussman, M.R., Amasino, R.M., Young, J.C., Krysan, P.J. and Austin-Phillips, S.** (2000) The *Arabidopsis* Knockout Facility at the University of Wisconsin-Madison. *Plant Physiol.* **124**, 1465–1467.
- Valverde, F., Mouradov, A., Soppe, W., Ravenscroft, D., Samach, A. and Coupland, G.** (2004) Photoreceptor regulation of CONSTANS protein in photoperiodic flowering. *Science*, **303**, 1003–1006.
- Wang, Z.Y. and Tobin, E.M.** (1998) Constitutive expression of the CIRCADIAN CLOCK ASSOCIATED1 (CCA1) gene disrupts circadian rhythms and suppresses its own expression. *Cell*, **93**, 1207–1217.
- Wang, C., Tian, Q., Hou, Z., Mucha, M., Aukerman, M. and Olsen, O.A.** (2007) The *Arabidopsis thaliana* AT PRP39-1 gene, encoding a tetratricopeptide repeat protein with similarity to the yeast pre-mRNA processing protein PRP39, affects flowering time. *Plant Cell Rep.* **26**, 1357–1366.
- Wesley, S.V., Helliwell, C.A., Smith, N.A. et al.** (2001) Construct design for efficient, effective and high-throughput gene silencing in plants. *Plant J.* **27**, 581–590.
- Wigge, P.A., Kim, M.C., Jaeger, K.E., Busch, W., Schmid, M., Lohmann, J.U. and Weigel, D.** (2005) Integration of spatial and temporal information during floral induction in *Arabidopsis*. *Science*, **309**, 1056–1059.