

The STAR project: context, objectives and approaches

Mike Furse^{1,*}, Daniel Hering², Otto Moog³, Piet Verdonschot⁴, Richard K. Johnson⁵, Karel Brabec⁶, Kostas Gritzalis⁷, Andrea Buffagni⁸, Paulo Pinto⁹, Nikolai Friberg¹⁰, John Murray-Bligh¹¹, Jiri Kokes¹², Renate Alber¹³, Philippe Usseglio-Polatera¹⁴, Peter Haase¹⁵, Roger Sweeting¹⁶, Barbara Bis¹⁷, Krzysztof Szoszkiewicz¹⁸, Hanna Soszka¹⁹, Gunta Springe²⁰, Ferdinand Sporka²¹ & Il'ja Krno²²

¹Centre for Ecology and Hydrology, CEH Dorset, Winfrith Technology Centre, Winfrith Newburgh, Dorchester, Dorset DT2 8ZD, UK

²Institute of Hydrology, University of Duisburg-Essen, Universitaetsstr. 5, 45117 Essen, Germany

³Institute for Hydrobiology and Aquatic Ecosystem Management, University of Natural Resources and Applied Life Sciences Vienna, Max Emanuel Strasse 17, A-1180 Vienna, Austria

⁴Department of Ecology and Environment, Alterra, Droevendaalsesteeg 3, 6700 AA Wageningen, The Netherlands

⁵Department of Environmental Assessment, Swedish University of Agricultural Sciences, P.O. Box 7050, S-750 07 Uppsala, Sweden

⁶Department of Zoology and Ecology, Masaryk University, Kotlářská, 611 37 Brno, Czech Republic

⁷Hellenic Centre for Marine Research, Institute of Inland Waters, 46.7 km Athens-Sounion Avenue, 190 13 Anavyssos, Greece

⁸CNR-Water Research Institute, Via della Mornera, 25 I-20047 Brugherio (Milano), Italy

⁹Centre of Applied Ecology, University of Evora, Apartado 94, Lago dos Colegiais 2, 7002–554 Evora, Portugal

¹⁰Department of Freshwater Ecology, NERI, National Environmental Research Institute, Vejlsovej 25, P.O. Box 314, DK-8600 Silkeborg, Denmark

¹¹South West Region, Manley House, Kestrel Way, Environment Agency, EX2 7LQ Exeter, Devon, UK

¹²Vyzkumny Ustav Vodohospodarsky T.G. Masayka, Drevarska 12, 657 57 Brno, Czech Republic

¹³LABBIO, Unterbergstrasse 2, 39055 Laives, Italy

¹⁴Centre of Ecotoxicology, Biodiversity and Environmental Health, University of Metz, Campus Bridoux, Rue de Général, 57070 Metz, France

¹⁵Senckenbergische Naturforschende Gesellschaft, Lochmuehle 2, D-63599 Biebergemünd, Germany

¹⁶Freshwater Biological Association, The Ferry House, Far Sawrey, LA22 0LP Cumbria, UK

¹⁷Institute of Ecology and Nature Protection, Department of Applied Ecology, University of Łódź, Banacha 12/16, 90-237 Łódź, Poland

¹⁸Department of Ecology and Environmental Protection, Agricultural University of August Cieszkowski, ul. Piątkowska 94C, 61-691 Poznan, Poland

¹⁹Lake Protection Laboratory, Instytut Ochrony Środowiska, Kolektorska 4, 01-692 Warsaw, Poland

²⁰Institute of Biology, University of Latvia, Miera 3, 2169 Salaspils, Latvia

²¹Institute of Zoology, Department of Hydrobiology, Slovak Academy of Sciences, Dubravska cesta 9, 84206 Bratislava, Slovakia

²²Faculty of Science, Department of Ecology, Comenius University Bratislava, Mlynská dolina B-2, 842 15 Bratislava, Slovakia

(* Author for correspondence: E-mail: mtf@ceh.ac.uk)

Key words: Water Framework Directive, ecological status, biological quality elements, intercalibration, uncertainty, software

Abstract

STAR is a European Commission Framework V project (EVK1-CT-2001-00089). The project aim is to provide practical advice and solutions with regard to many of the issues associated with the Water

Framework Directive. This paper provides a context for the STAR research programme through a review of the requirements of the directive and the Common Implementation Strategy responsible for guiding its implementation. The scientific and strategic objectives of STAR are set out in the form of a series of research questions and the reader is referred to the papers in this volume that address those objectives, which include: (a) Which methods or biological quality elements are best able to indicate certain stressors? (b) Which method can be used on which scale? (c) Which method is suited for early and late warnings? (d) How are different assessment methods affected by errors and uncertainty? (e) How can data from different assessment methods be intercalibrated? (f) How can the cost-effectiveness of field and laboratory protocols be optimised? (g) How can boundaries of the five classes of Ecological Status be best set? (h) What contribution can STAR make to the development of European standards? The methodological approaches adopted to meet these objectives are described. These include the selection of the 22 stream-types and 263 sites sampled in 11 countries, the sampling protocols used to sample and survey phytobenthos, macrophytes, macroinvertebrates, fish and hydromorphology, the quality control and uncertainty analyses that were applied, including training, replicate sampling and audit of performance, the development of bespoke software and the project outputs. This paper provides the detailed background information to be referred to in conjunction with most of the other papers in this volume. These papers are divided into seven sections: (1) typology, (2) organism groups, (3) macrophytes and diatoms, (4) hydromorphology, (5) tools for assessing European streams with macroinvertebrates, (6) intercalibration and comparison and (7) errors and uncertainty. The principal findings of the papers in each section and their relevance to the Water Framework Directive are synthesised in short summary papers at the beginning of each section. Additional outputs, including all sampling and laboratory protocols and project deliverables, together with a range of freely downloadable software are available from the project website at www.eu_star.at.

Context

The Water Framework Directive

Europe has a hundred years of experience of using biological assemblages to assess the condition of streams and rivers. The first procedures were developed early in the 20th century in central Europe and were based on the concept of saprobity (Sladeczek, 1973). Saprobian systems varied in their design and application but could use both micro- and macroscopic plant and animal communities in order to evaluate sites. A wide diversity of techniques blossomed throughout the 20th century (Hellowell, 1978, 1986) and, whilst a range of different biological groups continued to be used, the use of benthic macroinvertebrates became by far the commonest approach (Metcalf, 1989; Metcalfe-Smith, 1994). Each country or, sometimes, region of a country tended to develop their own methodological procedures (Knoben et al., 1995). These incorporated a common internal

approach to sampling, sample processing, indexation and quality classifications (Birk & Hering, 2002).

Whilst a range of specific monitoring traditions was evolving in individual states, the formation of the European Union resulted in a growing convergence of the legislative infrastructure of its Member States and the strategies adopted to implement this legislation. The mechanism commonly used to implement common community practices has been the issue of a directive from the European parliament. In the 1990's pressure grew for the rationalisation of these 'water quality' directives into a single overarching directive to meet this objective (Mandl, 1992). The resultant directive, commonly known as the Water Framework Directive or WFD, was published in 2000 (European Commission, 2000).

Significantly, the directive embraced the concept of the 'Reference Condition' (Hughes, 1995) as a unifying concept for aiding the harmonization of results obtained in a variety of different

countries/regions using a variety of their own 'traditional' assessment protocols. This concept had already been applied successfully in the United Kingdom through the development and application of RIVPACS (Wright et al., 1989, 2000) and had subsequently been taken up outside Europe in Australia (Norris, 1994) and Canada (Reynoldson et al., 1995, 2000; Rosenberg et al., 2000).

The WFD recognised type specific biological reference conditions based on a physical and chemical typology of surface water bodies in each European eco-region sensu Illies (1978). For this purpose Member States were expected to develop a reference network for each stream type containing a sufficient number of sites of high ecological status to provide a sufficient level of confidence about the values for the reference condition.

The term 'Ecological Status' was the overarching term coined by the WFD to represent the 'quality of the structure and functioning of aquatic ecosystems associated with surface waters'. Five categories of Ecological Status are recognised by the directive; High, Good, Moderate, Poor and Bad. The WFD provides normative definitions of the biological community structure associated with the High, Good and Moderate status classes. (European Commission, 2000). Member States are required to implement programmes of measures in order that all surface water bodies achieve at least 'good Ecological Status' within a defined timetable.

Whereas only macroinvertebrate data were required for the application of most prediction and assessment systems, the WFD required the sampling and interpretation of data on a broader suite of 'biological quality elements' (BQEs). These included phytoplankton, other aquatic flora, macroinvertebrates and fish. Parameters to be considered for each element are the composition and abundance of its biotic assemblages. In addition the age structure of fish populations shall be taken into consideration.

In common with systems such as RIVPACS (Wright et al., 2000), the WFD required that observed metric values for BQEs in a water body undergoing monitoring were mathematically compared with expected values for reference condition sites based on predictive modelling, hind-casting or expert judgement. The WFD presumed

that the ratios so-calculated would be in the range 0–1 and the numerical value derived by such a comparison was termed the Ecological Quality Ratio (EQR). The division of the value range of an EQR into classes provides a mechanism for categorising the ecological status of sites.

The precise BQEs to be monitored will be dependant on the type of monitoring to be undertaken. The WFD recognises three forms of monitoring: surveillance (to provide an assessment of the overall surface water status within each catchment), operational (to establish the status of water bodies identified as being at risk of failing to meet environmental objectives) and investigative (the source and magnitude of a specific pollutant). In surveillance monitoring, parameters indicative of all biological elements shall be monitored except where it is not possible to establish reference conditions for a particular element due to that element's high degree of natural variability in the water body being monitored. In contrast, operational and investigative monitoring may be restricted to one or two BQEs.

In addition to the direct monitoring of the biological assemblages, the other quality elements to be monitored for the classification of Ecological Status comprise hydromorphological, chemical and physiochemical elements supporting the biological elements.

Common Implementation Strategy

The WFD sets the framework for future monitoring of surface waters and sets out the mechanisms for reporting on the results of monitoring programmes and the formulation of river basin management plans, based upon the information gathered by monitoring and other sources. However, it is not prescriptive of the methodologies to be used to collect and process biological samples nor the specific metrics or multi-metrics to be used to calculate the Ecological Quality Ratios or the class value limits of these EQRs for each of the five classes of Ecological Status. It also provides no specific guidance on how the results of monitoring of the many and diverse quality elements shall be integrated in order to provide a single classification of the water body's status nor on how estimates of the required level of confidence and precision should be made.

For these reasons (European Commission, 2001), a Common Implementation Strategy (CIS) was established in order to develop common understanding of the technical and scientific implications of the directive and, in so doing, to achieve its harmonised implementation. Amongst the many guidance documents emanating from CIS working groups are reports on the establishment of the Intercalibration Network and on the intercalibration exercise (European Commission, 2002), on establishing reference conditions and ecological status class boundaries (European Commission, 2003a), on monitoring for the WFD (European Commission, 2003b) and on the overall approach to the classification of Ecological Status (European Commission, 2003c).

A series of Geographical Intercalibration Groups (GIGS) have been set up to agree on the intercalibration strategy to be adopted in discrete geographical areas of the European Union. Fifteen GIGS have been established including five river groups for the regions Mediterranean, Central, Alpine, Eastern Continental and Northern. A defined number of countries comprise each GIG but individual countries may belong to more than one GIG if the variation in the river types within its borders qualifies it to do so.

Supportive European Commission research projects

AQEM

In support of the technical activities associated with the implementation of the WFD, the European Union has commissioned a series of research projects designed to provide scientific support for the technical processes. The first of these projects specifically concerned with the assessment of Ecological Status was the AQEM project (EVK1-CT1999-00027). The structure and objectives of the project and the main scientific findings and applied outputs are described in a special issue of *Hydrobiologia* (Hering et al., 2004b).

The AQEM project established a standard macroinvertebrate sampling protocol, the AQEM method, and a common field protocol for recording hydromorphological, physical, chemical and geographical information concerning the study sites and their upstream, downstream and riparian environs (Hering et al., 2004a). Outputs of the

project include a database (AQEMDip) for the orderly storage and retrieval of macroinvertebrate and environmental data and a river assessment program (now termed ASTERICS) for calculating the values of almost 200 biological metrics and selected national multi-metric systems.

Whilst the AQEM project addressed many of the key questions associated with the use of macroinvertebrate data for assessing the Ecological Status of surface waters, the directive also required the integration of other biological quality elements together with the hydromorphological, chemical and physical elements that support the biological elements.

STAR

STAR is a European Commission Framework V project (EVK1-CT-2001-00089) with the full title of 'Standardisation of river classifications: Framework method for calibrating different biological survey results against ecological quality classifications to be developed for the Water Framework Directive'. The project is categorised as 'Pre-normative, co-normative research and standardisation'. It therefore seeks to provide practical solutions to some of the additional problems associated with the implementation of the directive. Issues addressed include comparison of macroinvertebrate sampling methods, the effectiveness of the use of different organism groups in different stream types and for different stressors, variation and uncertainty in the collection and interpretation of biological data, the inter-calibration of assessment methods for the allocation of Ecological Status, the formulation of drafts for the relevant CEN bodies, and the development of a decision support system to assist water managers in applying the project findings. In this paper the objectives of the STAR project and the methodological approach adopted to achieve these aims will be described. It will provide the background for the remaining papers that comprise this special issue of *Hydrobiologia*.

FAME and REBECCA

Clustered with the STAR project and collaborating closely with it has been another EC Framework V project, FAME (EVK1-CT-2001-00094). This project has developed a specific system for the assessment of the Ecological

Status of surface waters based on the fish communities that they support (Noble & Cowx, 2002). The STAR project is also working collaboratively with the EC Framework VI project, REBECCA (SSP1-CT-2003-502158), that aims to provide new interpretations of the relationships between Chemical and Ecological Status of surface waters in order to support the implementation of the WFD.

Objectives

The central objectives of STAR and the papers in this volume that address them are:

- *Which methods or biological quality elements are best able to indicate certain stressors?*

The varying responses to stressors of different biological quality elements will allow WFD monitoring data to be interpreted in a diagnostic manner in order to identify the pressures operating on aquatic systems. Advice on the selection of the most appropriate BQEs for specific objectives and in specific regions is provided by Johnson et al. (2006a, b) and Pinto et al. (2006). Other authors consider specific techniques (Kokeš et al., 2006 – PERLA) or taxonomic groups and stressors (Szoszkiewicz et al., 2006b – macrophytes and organic pollution; O'Hare et al., 2006 – macrophytes and habitat alteration). In addition the application of River Habitat Survey Techniques (Raven et al., 1998) to the evaluation of the hydromorphological condition of watercourses is evaluated by Erba et al. (2006) and Szoszkiewicz et al. (2006a).

- *Which method can be used on which scale?*

The organism groups that the WFD require to be considered in assessing the Ecological Status of waterbodies indicate environmental change on different scales. The issue of scale has been considered by Springe et al. (2006) and Verdonschot (2006a).

- *Which methods are suited for early and late warnings?*

Besides the spatial dimension, different organism groups indicate change on different temporal dimensions, thus providing different signals of

early or late warning. Johnson et al. (2006b) address this issue for all BQEs except phytoplankton.

- *How are different assessment methods affected by errors and how can 'signal' be distinguished from 'noise'?*

STAR has investigated a range of factors that confound the ability of bioassessment procedures to detect change and many papers in this volume address the issue of uncertainty. These include Besse-Lotoskaya et al. (2006) who investigate uncertainty associated with diatom sampling and interpretation, Clarke et al. (2006a, b) and Lorenz & Clarke (2006) who look at the impact of sampling variation on macro-invertebrate assessments, Haase et al. (2006) who consider the effects of macro-invertebrate sorting and identification errors; Staniszewski et al. (2006) and Baattrup-Pedersen et al. (2006) who examine uncertainty associated with macrophyte surveys and Johnson et al. (2006a) who explore the incidence and effects of Type I and Type II errors for most BQEs. One factor that may influence the evaluation of sites is the method used to define reference conditions. Davy-Bowker et al. (2006) consider the implications of using type specific conditions based on a physical/chemical typology with those site specific reference conditions produced by predictive systems such as RIVPACS and PERLA (Kokeš et al., 2006).

- *How can data from different assessment methods and taxonomic groups be compared and inter-calibrated and how can the results of the STAR programme be used to assist the WFD intercalibration exercise?*

A central problem, for the implementation of the WFD, is how biological data collected using different national protocols and biological quality elements (BQEs) can be compared and integrated in order to derive comparable allocations of sites to standard European classes of environmental degradation. Friberg et al. (2006) compare the main macroinvertebrate sampling procedures used in Europe, whilst alternative mechanisms for inter-calibration are discussed by Birk & Hering (2006 – macroinvertebrates), Birk et al. (2006 – macrophytes) and Buffagni et al. (2006 – general but principally macroinvertebrates).

- *How can the cost-effectiveness of field and laboratory protocols for the collection and processing of macroinvertebrate samples be optimised?*

Methodologically, standardisation must also take a balanced account of the relative costs and ecological effectiveness of different field and laboratory procedures. Experimental field studies were devised to consider a spectrum of relevant issues (Šporka et al., 2006; Vlek et al., 2006) whilst Verdonschot (2006b) examined the significance of varying levels taxonomic precision on the biological typology of European streams and rivers.

- *Can species trait analysis provide a unifying procedure for the establishment of reference conditions and the assessment of Ecological Status?*

An aim of STAR was to test the applicability of a species trait analysis as a unifying theme for the derivation of functionally based reference conditions and, as a result, for the assessment of Ecological Status. These results are presented elsewhere including as Deliverable N2 (Bis & Usseglio-Polatera, 2004) on the STAR website – www.eu-star.at

- *How can boundaries of the five classes of Ecological Status recognised by the WFD be best set?*

On the basis of the field and laboratory protocols and metrics that will be tested in STAR, an aim of the project is investigate and to elaborate standard procedures for the determination of European class boundaries of Ecological Status. Mechanisms for setting and inter-calibrating class boundaries are considered by Birk & Hering (2006), Birk et al. (2006) and Buffagni et al. (2006).

- *How can the results of the STAR programme be used to make recommendations for common European standards?*

The STAR consortium have suggested outline standards, on methodological issues related to the implementation of the WFD that are being considered by CEN (Comité Européen de Normalisation) for adoption as full standards. These include multi-habitat sampling for invertebrates, the construction of multi-metric assessment

systems and the selection of the best suited organism groups for specific monitoring purposes. Methodologies for developing multi-metric indices are elaborated in this volume by Hering et al. (2006). An additional standard tool for the use of the European water industry and academia is a pan-European macro-invertebrate ecological database and taxa inventory described here by Schmidt-Kloiber et al. (2006).

Approaches: site selection

Research framework

The STAR consortium comprised 22 partners from 14 countries including four countries who were candidate states, the Czech Republic, Slovakia, Poland and Latvia, that acceded to the European Union during the course of the project on 1st May 2004.

The project was divided into 19 discrete but inter-linked workpackages (Table 1). Most workpackages (WPs) could be allocated to one or other of two loose groupings. There were ten core WPs in which most partners worked collaboratively on a common activity and nine that were specific research programmes contributed to by a small minority of the partners and predominantly engaged in by a dominant leading institute (Table 1).

Stream types studied

The central components of the STAR project were the two WPs devoted to the collection of new biological, hydromorphological and other environmental data (WP7 and WP8).

WP7 (Table 1) involved the selection and monitoring of sites in two core groups of stream types (Table 2). The variables and their ranges used to define each group were those involved in the system A approach to surface water body typology given in the WFD (European Commission, 2000). Sites in core group 1 were defined as ‘Small, shallow, upland streams’ in early STAR project documentation. In WFD system A terms they are sites with a ‘small’ catchment situated in the lower 60% of the ‘mid-altitude’ range. Core group 2 sites were defined in early STAR

Table 1. The 19 STAR project workpackages (WPs)

No.	Theme
1	<i>Project co-ordination</i>
2	<i>Project homepage</i>
3	<i>Review of data on reference conditions and existing assessment methods using benthic invertebrates, fish, phytobenthos, macrophytes and river habitat surveys, national standards on sampling, analysis and quality evaluation, related national projects and existing databases</i>
4	<i>Acquisition of existing data</i>
5	Selection of sampling sites
6	Sampling workshops to standardise the understanding and application of sampling protocols between participants and to undertake replicate sampling programmes for diatoms and macroinvertebrates
7	Investigation of core stream types 1 (small, shallow, mountain streams) and 2 (medium-sized, deeper, lowland streams)
8	Investigation of additional stream types
9	Audit of performance in the processing and identification of macroinvertebrate and diatom samples
10	Generation and hosting of the project database
11	Comparison and linking assessment systems based on invertebrates
12	Linking of assessment systems working with different organism groups
13	Linking of the project database and the database of existing data
14	Recommendations for standardisation to support CEN in its development of appropriate standard methods for the WFD
15	<i>Elaboration of a decision support system, implemented through a DSS computer program, to provide practical guidance in the application of monitoring programmes necessary to meet the terms and objectives of the Water Framework Directive</i>
16	<i>Examination of the effectiveness of and relative cost-efficiency of different field and laboratory protocols for the collection and processing of macroinvertebrate samples</i>
17	<i>Examination of the value of species trait analysis as a unifying system for the establishment of functionally based reference conditions and the assessment of Ecological Status</i>
18	<i>Spatial scale analyses</i>
19	<i>Study of errors and variation associated with field protocols for the collection and application of macrophyte and hydro-morphological data in the implementation of the WFD</i>

The 10 core collaborative WPs are shown in regular font.

documentation as ‘Medium-sized, deeper lowland streams’. In WFD system A terms they have ‘medium’ catchment sizes and are situated at ‘lowland’ altitudes.

WP8 (Table 1) involved the selection and sampling of a group of ‘additional’ stream types. Additional streams types were not prescriptively allocated to any WFD system A typology and could include sites whose combination of altitude and catchment size characteristics might or might

not fit the definition of either core stream type groups 1 or 2. In general terms they were confined to either the system A mid-altitude or lowland categories and to the system A small, medium or, very occasionally, large catchment size categories. Initially the additional stream types were selected to fulfil four specific roles. These were to:

- allow new, characteristic sites of individual states to be included in the analysis;

Table 2. Definitions of the two STAR core stream type groups

Core stream type		Theoretical value range of typological variables		
No.	Description	Altitude	Catchment size	Geology
1	Small, shallow, upland streams	200–500 m	10–100 km ²	Calcareous or siliceous
2	Medium-sized, deeper lowland streams	<200 m	>100–1000 km ²	Calcareous, siliceous or organic

- provide an opportunity to extend the range of sites in existing European assessment systems;
- extend the range of sites at which the specific field methods are compared;
- provide an opportunity to test alternative sampling/assessment methods of specific importance to individual consortium Member States.

However, the data for core and additional stream types were used jointly in most analyses. Core and additional stream types could also be defined as either calcareous, siliceous or, occasionally, organic but, with a few exceptions, sites within specific site sets (see the following section) were all in the same geological category.

Selection of site sets

Each participating partner in WP7 and/or WP8 selected a minimum of one and a maximum of three sets of sites to sample. Each set of sites was in one of the three basic stream type groups (core 1, core 2 or additional) described in the previous section. Partners with two or more site sets selected these sets to be either in the same or different stream type groups. Sets of sites defined by their stream type group, eco-region or sub-eco-region and, optionally, other geographical criteria are termed 'stream types'. The definition of stream types used here is that established by the AQEM project and is "an artificially delineated but potentially ecologically meaningful entity with limited internal biotic (taxa composition) and abiotic (chemical and hydromorphological) variation and a biotic and abiotic discontinuity toward other types" (Hering et al., 2004a). Selection of specific stream types within the three stream type groups defined in the previous section took account of many of the criteria for stream typology in System B of the WFD.

In total, 22 stream types were selected for study as part of either WP7 or WP8 (Table 3). In addition, two other stream types in Italy (small-sized calcareous streams in the Southern Apennines and medium-sized calcareous streams in the Northern Apennines) and three other stream types in Greece (small-sized siliceous streams in Northern Greece, medium-sized calcareous streams in Southern Greece and small-sized siliceous streams on the Aegean Islands) were sampled for other national purposes connected with the STAR project.

For each stream type, a minimum of ten and a maximum of 24 sites were sampled (Table 4). For each stream type, sites were selected to represent a gradient of degradation usually due to a pre-identified dominant stressor (Table 4). For the purpose of site selection, these dominant stressors were divided into three broad categories: organic pollution (including eutrophication), toxic pollution (including acidification) and habitat degradation. In one case, (stream type I06 – Italy) a single dominant stressor could not be identified and the category 'general' stressors was applied. In a few other cases (see Table 4) different dominant stresses applied to specific sites within a stream type and some of these only became apparent during the sampling programme.

In general, approximately 25% of sites in each site set were selected to be likely to be of 'high', 25% of 'good', 25% of 'moderate' and 25% of 'poor'/'bad' Ecological Status. The 'high' status sites were selected to represent the reference condition for their particular stream type. Reference condition sites were selected through a combination of site visits, cartographic information and information derived from new biological sampling or existing sample data held by internal (i.e. partner's own) or external (e.g. national monitoring organisations) sources. Where adequate data were available, all biological quality elements and hydromorphological and chemical quality elements were considered. However, in many cases the most important elements considered were macroinvertebrates, hydromorphology and nutrient status. In order to aid the process of reference site selection a list of criteria was developed (Table 5) based on Hering et al. (2003) but modified in response to the ongoing discussions of the REFCOND group.

In many cases, e.g. some lowland stream types or larger streams, no reference sites meeting all of the criteria above were available. For these stream types the 'best available' existing sites were selected. However, where possible, the description of reference communities of these types could be supplemented by evaluation of historical data and possibly the biotic composition of comparable stream types, e.g. streams of a similar size but located in different ecoregions.

The remaining sites, other than reference sites, were pre-classified using the same sources of information but with particular attention to

Table 3. The 22 stream types sampled as part of WP7 (core streams) or WP8 (additional streams)

Country	Core stream type group 1		Core stream type group 2		Additional stream type	
	No.	Name	No.	Name	No.	Name
Austria	A05	Small-sized, shallow mountain streams			A06	Small-sized, crystalline streams of the ridges of the Central Alps
Czech Republic	C04	Small-sized, shallow mountain streams			C05	Small-sized streams in the Central sub-alpine Mountains
Denmark			K02	Medium-sized lowland streams		
France					F08	Small-sized, shallow headwater streams in Eastern France
Germany	D04	Small-sized, shallow mountain streams	D03	Medium-sized lowland streams	D06	Small-sized, Buntsandstein streams
Greece					H04	Small-sized, calcareous mountain streams in Western, Central and Southern Greece
Italy					I05	Small-sized streams in the southern calcareous Alps
Latvia			L02	Medium-sized lowland streams	I06	Small-sized, calcareous streams in the Central Apennines
Poland			O02	Medium-sized lowland streams (Ecoregion 14)		
			O03	Medium-sized lowland streams (Ecoregion 16)		
Portugal					P04	Medium-sized streams in lower mountainous areas of Southern Portugal
Slovakia	V01	Small-sized, calcareous mountain streams in the Eastern Carpathians				
	V02	Small-sized, siliceous mountain streams in the Western Carpathians				
Sweden			S05	Medium-sized lowland streams	S06	Medium-sized streams on calcareous soils
United Kingdom			U23	Medium-sized lowland streams	U15	Small-sized, shallow, lowland streams

Table 4. The number of WP7 and WP8 sites sampled for each biological quality element, where the qualifying criterion is that sites must have been fully sampled for macroinvertebrates (except Slovakia, where regular 'national' sampling was not undertaken)

Country	Site type	Number of sites	Geology	Degradation	Number of sites sampled for each quality element				
					Diatoms	Macrophytes	Macroinvertebrates	Fish	Hydromorphology
Austria	A05	15	Siliceous	Stream morphology	11	11	13	12	13
	A06	21	Siliceous	Stream morphology	7	8	8	0	8
Czech Republic	C04	14	11 siliceous and 3 calcareous	Organic pollution	14	14	14	14	14
	C05	10	Siliceous	Stream morphology	10	10	10	10	10
Denmark	K02	12	9 calcareous and 3 siliceous	Stream morphology	11	11	11	11	11
	F08	12	Calcareous	Organic pollution	12	12	12	12	12
Germany	D03	13	Siliceous	Stream morphology	10	12	12	9	12
	D04	12	Siliceous	Stream morphology	10	12	12	10	12
Greece	D06	10	Siliceous	Stream morphology	6	6	6	6	6
	H04	10	Calcareous	Organic pollution	10	10	10	10	10
Italy	I05	10	Calcareous	Stream morphology	10	0	10	10	10
	I06	11	Calcareous	General	11	11	11	11	11
Latvia	L02	24	16 siliceous, 6 calcareous and 2 siliceous/calcareous	Organic pollution	24	24	24	24	24
	O02	13	12 siliceous and 1 siliceous/calcareous	Organic pollution	13	13	13	13	13
Poland	O03	12	8 organic, 2 siliceous and 2 organic/siliceous	Organic pollution	11	11	11	11	11
	P04	10	Siliceous	Organic pollution	10	10	10	10	10
Slovakia	V01	12	Calcareous	Organic pollution	12	12	12	12	12
	V02	12	Siliceous	Organic pollution	12	12	12	12	12
Sweden	S05	16	Siliceous	Acidification/toxic (5 sites), organic (4 sites), reference (3 sites), undefined (4 sites)	14	16	16	16	16
	S06	11	Calcareous	Organic pollution	9	11	11	11	11
United Kingdom	U15	13	Calcareous	Organic pollution	13	13	13	13	13
	U23	12	11 calcareous and 1 siliceous	Organic pollution	12	12	12	12	12
Totals		285			252	251	263	249	263

Table 5. Criteria for reference site selection

<p><i>General</i></p> <ul style="list-style-type: none"> • The reference condition must be politically palatable and reasonable • <u>A reference site, or process for determining it, must hold or consider important aspects of 'natural' conditions</u> • <u>The reference conditions must reflect only minimal anthropogenic disturbance</u> <p><i>Land use practices in the catchment area</i></p> <ul style="list-style-type: none"> • The degree of urbanisation, agriculture and silviculture should be as low as possible for a site to serve as a reference site • Least-influenced sites with the most natural vegetation are to be chosen <p><i>River channel and habitats</i></p> <ul style="list-style-type: none"> • The reference site floodplain should not be cultivated. If possible, it should be covered with natural climax vegetation and/or unmanaged forest • <u>Coarse woody debris should not be removed (minimum demand: presence of coarse woody debris)</u> • <u>Stream bottoms and stream margins must not be fixed</u> • <u>Spawning habitats for the natural fish population (e.g. gravel bars, floodplain ponds connected to the stream) should be present</u> • Preferably, there should be no migration barriers (affecting the bed load transport and/or the biota of the sampling site) • In stream types in which naturally anadromous fish species would occur, the accessibility of the reference site from downstream is an important aspect for the site selection • Only moderate influence due to flood protection measures can be accepted <p><i>Riparian vegetation and floodplain</i></p> <ul style="list-style-type: none"> • <u>Natural riparian vegetation and floodplain conditions must still exist</u> • Lateral connectivity between the stream and its floodplain should be possible • The riparian buffer zone should be greater or equal to 3× channel width <p><i>Hydrologic conditions and regulation</i></p> <ul style="list-style-type: none"> • No alterations of the natural hydrograph and discharge regime should occur • There should be no or only minor upstream impoundments, reservoirs, weirs and reservoirs retaining sediment; no effect on the biota of the sampling site should be recognisable • There should be no effective hydrological alterations such as water diversion, abstraction or pulse releases <p><i>Physical and chemical conditions</i></p> <ul style="list-style-type: none"> • <u>No point sources of pollution or nutrient input affecting the site</u> • <u>No point sources of eutrophication affecting the site</u> • No sign of diffuse inputs or factors which suggest that diffuse inputs are to be expected • 'Normal' background levels of nutrient and chemical base load, which reflect a specific catchment area • <u>No sign of acidification</u> • No liming activities • No impairments due to physical conditions • Thermal conditions must be close to natural • <u>No local impairments due to chemical conditions; especially no known point-sources of significant pollution, all the while considering near-natural pollution capacity of the water body</u> • <u>No sign of salinity</u> <p><i>Biological conditions</i></p> <ul style="list-style-type: none"> • No significant impairment of the indigenous biota by introduction of fish, crustaceans, mussels or any other kind of plants and animals • No significant impairment of the indigenous biota by fish farming • No intensive management, e.g. of the fish population 	<hr/> <p>Underlined criteria were mandatory and as many of the other criteria as possible were met.</p>
--	---

reported known sources of stress as supplied by national and regional agencies with responsibility for monitoring water quality and hydromorphology.

The relative emphasis placed on individual quality elements varied from partner to partner with some placing more emphasis on biological data and less

on hydromorphological and chemical elements than others in establishing their site pre-classification. However, each partner used their specific approach to establish a set of sites with a marked degradation gradient according to their chosen dominant stressor.

In total, excluding the Italian and Greek sites sampled for other purposes, 285 sites were selected for possible sampling for WP7 or WP8. Of these, 263 (Fig. 1) were sampled for macroinvertebrates in each sampling season using both the AQEM and, with the single exception of Slovakia, a second, mainly 'national' sampling protocol in each sampling season. These were the sites included in most of the central analyses undertaken in the project. All of these 263 sites were also subject to hydromorphological surveys and 252 were sampled for phytobenthos, 251 for macrophytes and 249 for fish. A final total of 233 were fully sampled for all biological quality elements.

Selection of quality elements

Three biological quality elements were sampled in all or almost all of the sites contributing to the central project analyses. These were 'aquatic flora', 'benthic invertebrate fauna' and 'fish fauna'. The aquatic flora was subdivided into phytobenthos

and macrophytes for the purposes of this project. The only component of the flora not sampled was phytoplankton because this element was considered not to be a significant component of the biota of the small to medium-sized, often fast-flowing streams that predominated in the STAR sampling programme.

At least two survey protocols were used to record components of the hydromorphological quality element supporting the biological elements. Information on chemical and physico-chemical quality elements supporting the biological elements was, in all cases collected both from direct field sampling and surveys and also, in some cases, from data collected by the national water quality monitoring agencies.

Selection of sampling reach

Prior to starting field sampling and surveying, the study reach at each site was selected. The reach was 500 m long and was selected as representative of the hydromorphological conditions of the stretch of river under investigation. A stretch of river is a continuous section of river without any significant tributaries or point sources of pollution likely to modify its Chemical Status (equivalent to a 'water body' as defined by the WFD). The



Figure 1. The location of the 263 sites sampled for macro-invertebrates.

selection of the sampling and survey reach normally followed the completion of the AQEM site protocol (see below).

Wherever possible, at each site, a common monitoring strategy was adopted in relation to the relative positions of the different sampling/surveying points. Field surveyors were provided with a conceptual diagram of this strategy (Fig. 2).

A preliminary reach was first located and, within this, the STAR-AQEM invertebrate sampling area was selected. This was a length of river of up to 100 m, depending on stream width (see the STAR-AQEM section below) at which all of the common representative habitat types of that river stretch were present, including both erosional ('riffle') and depositing ('pool') areas if possible. The centre of this sampling length was taken to be River Habitat Survey (RHS) spot check 9 (Raven et al., 1998) and was used to

define the exact position of the whole 500 m RHS survey reach.

The 'national' invertebrate sampling and the phytobenthos sampling were undertaken in the same 100 m section as the STAR-AQEM method. Care was taken to minimise the disturbance to the river by each sampling method and overlap between the different precise sampling locations.

The macrophyte survey was undertaken in the 100 m reach immediately upstream of the invertebrate and diatom sampling reach and after these elements had been sampled. Where all three elements were sampled on the same day this spatial separation and sequence of sampling was designed to minimise any trampling of plants resulting from the sampling of the other two elements.

Fish sampling took place over a ≥ 100 m section immediately upstream of the macrophyte survey area. Fish sampling was normally undertaken on a




Figure 2. The conceptual locations of the STAR sampling areas for each of the five recorded quality elements at each site, as provided to field surveyors.

separate date to the other sampling. Chemical sampling was from within this 500 m reach of river and avoided any disturbance to the sediment caused by biological sampling.

Whilst this strategy represented the ideal, on occasions local conditions required variations in the general pattern of sampling. Such departures from the optimal were kept to the minimum.

Approaches: field and laboratory protocols

Phytobenthos – diatoms

Diatoms were sampled once only at the WP7 and WP8 sites. Samples were collected during periods of stable stream flow and at least four weeks after a period of extreme conditions like a major storm or drought. The time of these stable conditions varied from region to region. Subject to this criterion, spring was the preferred season for sampling as diatoms dominate the phytobenthos during this season (Moore, 1977).

The location selected for STAR-AQEM invertebrate samples (Fig. 2) and the criteria for its selection, ensured that it also had the most suitable available substrata for sampling benthic diatoms. This ranged from stones to macrophytes and to mineral sediments, depending on the type of river. Selection criteria also ensured that the sampled section also combined riffles and pools and thus enabled the sampling of a good variety of natural substrata. Bank side areas were avoided during sampling with samples taken at least 10% of the river width away from the river edge.

In general terms, the sampling and processing protocols used followed those of Kelly et al. (1998) and Winter & Duthie (2000). Methods conform to the CEN standards EN 13946 and EN 14407. The full STAR protocol for the sampling, processing and audit of diatom samples was prepared by Alterra and is available from the STAR website (www.eu-star.at).

Phytobenthos – non-diatoms

Collection of non-diatom phytobenthos was voluntary and not all partners collected information on this taxonomic group. Where partners did collect and process material they adopted the

methods described in the project protocol for sampling, processing and audit of non-diatom benthic algal samples, which was also prepared by Alterra and is available from the STAR website.

Macrophytes

Macrophytes were surveyed once only at the WP7 and WP8 sites. Surveys were undertaken using a slightly adapted form of the Mean Trophic Rank (MTR) field protocol developed in the United Kingdom (Holmes et al., 1999). Most surveys were carried out between mid-June and mid-September after several days of low flow or low-normal flow as opposed to high flow/spate.

The MTR survey procedure is based on the presence and abundance of species of aquatic macrophytes, where a macrophyte is defined as ‘any plant observable with the naked eye and nearly always identifiable when observed’ (Holmes & Whitton, 1977). This definition includes all higher aquatic plants, vascular cryptogams and bryophytes, together with groups of algae which can be seen to be composed predominantly of a single species.

Survey techniques conformed to the CEN standard EN 14184. The full STAR survey protocol for macrophytes (Guidance for the field assessment of macrophytes of rivers within the STAR project) was prepared for STAR by the Centre for Ecology and Hydrology and is available from the STAR website.

Macroinvertebrates

All 263 WP7 and WP8 sites listed in the macroinvertebrates column of Table 4 were sampled using a modified form of the AQEM method (AQEM consortium, 2002; Hering et al., 2004a), known as the STAR-AQEM method. With the exception of all Slovakian sites in stream type V01 and six sites in V02, all sites were also sampled using a current national method of the country (Table 6). Where no consistent national sampling method existed for the country, either the RIVP-ACS (Austria, Germany and Greece) or PERLA (Slovakia) methods were used instead (Table 6). With the exception of the three non-UK RIVP-ACS users, the national methods used were

Table 6. 'National' sampling methods applied in each STAR country participating in project workpackages 7 and 8

Country	Methods applied	Reference
Denmark	Danish Stream Fauna Index (DSFI)	Danish Environmental Protection Agency (1998)
Italy	Indice Biotico Estesio	Ghetti (1997)
France	Indice Biologique Global Normalisé (IBGN)	GAY, Cabinet en Environnement (1994)
Latvia	LVS 240:1999	Unpublished (see 'Protocols' on www.eu-star.at)
Czech Republic	PERLA	Kokeš et al. (2006)
Slovakia	PERLA	
Poland	Polish national method	Unpublished (see 'Protocols' on www.eu-star.at)
Portugal	Portuguese national method (PMP)	Unpublished (see 'Protocols' on www.eu-star.at)
Austria	River In-Vertebrate Prediction And Classification System (RIVPACS)	Murray-Bligh et al. (1997)
Germany	River In-Vertebrate Prediction And Classification System (RIVPACS)	
Greece	River In-Vertebrate Prediction And Classification System (RIVPACS)	
United Kingdom	River In-Vertebrate Prediction And Classification System (RIVPACS)	
Sweden	Swedish national method	Swedish Environmental Protection Agency (1996)

assumed to be the methods likely to be adopted by their countries for implementing the WFD.

Immediately prior to sampling, the length of river to be sampled was surveyed as part of the AQEM field protocol and the proportions of the different habitats present at the river bottom were estimated (Hering et al., 2004a). This knowledge was used to establish the precise STAR-AQEM sampling area and the proportions of micro-habitats to be sampled (Hering et al., 2004a). Normally, the STAR-AQEM sample was the first to be collected followed by the 'national' sample. For the national sample, care was taken to avoid the specific locations at which the STAR-AQEM sample was collected.

Most samples were fixed and/or preserved in the field using a fixative/preservative of the partner's choice which was normally either formaldehyde solution or ethanol of varying strength. Exceptions to this generalisation were the Italian IBE and most Latvian LVS 240:1999 samples that were sorted at the bankside, and some Portuguese 'national' samples that were sorted live in the laboratory within 48 h of collection. Prior to preservation and/or transport to the laboratory, large and easily identified specimens and identifiable

specimens of taxa of known conservation importance or particular fragility to damage were recorded and returned live to the river.

The laboratory sample processing techniques were specific to the particular field protocols and differed between the STAR-AQEM and 'national' samples and between the different 'national' field protocols. However, all partners were trained to collect and process STAR-AQEM samples in a consistent and prescriptive manner.

In all cases taxa were identified to the best achievable level, according to the expertise of the partner and the availability of adequate national keys. Most partners achieved species level identification for most groups but this was not possible in Latvia, where only some groups could be identified to this level, nor in Greece, Italy or Portugal where most identifications were to family level.

It is not possible to describe each field and laboratory protocol here but the key features of each method are provided Friberg et al. (2006). For further details the reader is directed to the references given in Table 6, to the 'Protocols' section of the STAR website (www.eu-star.at), the AQEM website (www.aqem.de) for the key principles of the AQEM method that formed the basis

of the STAR-AQEM procedure and to the 'Water-view' database developed during the STAR project (Birk & Hering, 2002) and accessible via the 'Review' section of the STAR website. LVS 240:1999, the Latvian national sampling protocol (Latvian Standard Ltd., 1999) and can also be accessed via www.lvs.lv/en/services/services_EP.html.

In all cases where hand-net sampling was employed sampling and equipment specifications were consistent with CEN standard EN 27828. Where Surber sampling was used, as in the case of the French IBGN method and, occasionally, in the STARAQEM method, sampling and equipment specifications were consistent with CEN standard EN 28265.

Fish

The fishing strategy used conformed to CEN standard EN 14011 and was developed following discussions with STAR's cluster project FAME (<http://fame.boku.ac.at>). Almost all STAR sites were, on average, less than one metre deep and, in these circumstances, the STAR protocol was based on the section of EN 14011 relating to electric fishing of wadeable rivers.

Where possible, fishing was carried out using direct current (dc) fields. However where this was not possible, due to high conductivity water, variable electrical characteristics of stream topography or poor fish response to dc field pulsed, direct current (pdc) fields were used. In all cases, fields were adjusted to the minimum voltage gradient and current density concomitant with efficient fish capture.

Optimally, the length of river fished was a minimum of 100 m and located in the centre of the RHS survey area (Fig. 2). Normally the full width of the river was surveyed over this length. However, in a small minority of cases the fishing reach was slightly shorter than 100 m for logistical reasons (Table 7). The relative position of the fishing area within the survey area was also sometimes varied for practical reasons. Wherever possible the fishing area was demarcated by upstream and downstream stop nets (Table 7). Net mesh sizes were suitable for preventing fish >5 cm from escaping. A minimum of two fishing runs was undertaken at most sites.

In the small number of cases where sites were not wadeable, fishing was undertaken from a boat (Table 7) and at a series of spot locations within the RHS survey area. In such circumstances, stop nets were not used, the sites were normally >10 m wide and the length of river sampled was often less than 100 m. Some wide, wadeable sites in Sweden were also sampled discontinuously without stop nets.

All or most of the following elements of the fish population in the sample area were recorded:

- Number of species
- Species composition (percentage of each species by number)
- Fish density by species (number of fish per m²) of individuals other than young of the year. There was no requirement to measure or age fish
- Young of the year per species (qualitative assessment by class, e.g. abundant, common or rare)
- Ratio between number of phytophils and limnophils (fish species grouped by reproductive guild (Balon, 1975; Mann, 1996)
- Number of intolerant or sensitive species in terms of functionally descriptive fish species (i.e., salmonids for water quality, migratory species for connectivity, etc.)
- Number of endemic species (species which are only present in the river basin under study)
- Number of native species (species known to be present in the watercourses of the country for a long period of time i.e. >200 years)
- Subjective assessment of degree of infestation of external parasites or other diseases

Final population estimates, capture efficiency and standard errors of population numbers were also determined. Two catch estimates were based on the Seber & LeCren (1967) method but where more than two capture runs were undertaken values were calculated using the Exact Maximum Likelihood methodology.

Hydromorphology

At least two standard site assessment protocols, River Habitat Survey and the AQEM site protocol, were conducted at each STAR WP7 and WP8 site. Only one RHS survey was undertaken in each

Table 7. Fish sample procedures operated by STAR partners in each country participating in WP7 and WP8

Parameter	Units /categories	United Kingdom	Germany	Austria	Sweden	Czech Republic	Greece (L.ABBIO)	Italy (L.ABBIO)	Portugal	Denmark	France	Poland	Latvia (Eastern)	Slovakia (Central)	Slovakia (Eastern)	
Normal length river fished at each site	Metres	100	280	100	50	100	100	100	100	100	100	100	70	100	100	
Maximum length of river fished at each site	Metres	100	810	130	50	123	100	100	100	100	100	100	80	120	100	
Minimum length of river fished at each site	Metres	80	100	30	50	63	100	100	100	100	80	100	50	50	30	
Most common depth category at sites	A = <25 cm B = 25- <50 cm C = 50- <100 cm D = ≥1 m	B	B	B	B	A	B	A	B/C	C	B/C	C	B	B	A	A
Use of boat to undertake fishing	A = Always B = Sometimes C = Never	C	B	C	C	C	C	C	C	B	C	C	C	C	C	C
Use of stop nets to enclose the fishing area	A = Always B = Sometimes C = Never	A	B	A	C	B	A	C	A	C	A	A	C	B	B	B
Multiple fishing runs at each sampling site	A = Always B = Sometimes C = Never	A	B	A	A	B	B	A	A	A	A	A	A	A	A	A
Maximum number of fishing runs at a single site	Number	3	5	3	3	2	2	2	3	3	2	3	3	2	2	2
Season most commonly sampled	A = Spring only B = Summer only C = Summer & autumn D = Autumn only	D	C	C	D	C	C	D	A	B	C	D	C	B	B	B

reach and normally in the period July to September. However, the AQEM site protocol comprised both time variant and time invariant variables. Time invariant variables were recorded during the first site visit only but time variant variables were recorded at the time of each macro-invertebrate sampling.

RHS is a system for assessing the quality of rivers based on their physical structure. The technique and associated interpretation comprise a standard field survey, a bespoke database and indexation systems for Habitat Quality Assessment (HQA) and Habitat Modification Score (HMS). River Habitat Survey (RHS) was developed and applied in the United Kingdom (Raven et al., 1998) but has also been applied in other European countries and has been specially adapted for use in southern Europe (Buffagni & Kemp, 2002). The southern European modifications to the system were principally, but not exclusively, to cater for the braided channels that commonly occur there (Buffagni & Kemp, 2002). The procedures conform to the evolving CEN standard prEN 14614.

The AQEM site protocol, as its name implies, was developed for the AQEM project and subsequently modified and simplified for the STAR project. A brief outline of the method is provided by Hering et al. (2004a) and fuller details of the original AQEM system and the modified STAR version are given on the two project websites (www.aqem.de and www.eu-star.at).

In addition separate site assessment protocols for phytobenthos, MTR and 'national' macroinvertebrate sampling were completed at most sites. Field measurements were complemented by cartographic information assembled for most of the protocols. The fish surveys were also complemented by a standard suite of site information required for the Fides database and site indexation system developed by the FAME project. These including field measurements, a broad suite of cartographic information and information on the biological, hydromorphological and chemical condition of the site collated from published data and from data supplied by national monitoring agencies.

The various site protocols adopted at STAR sites are each too complex to document in detail here and the reader is generally referred to the cited literature for a more complete understanding

of the components and implementation of the RHS and AQEM site protocol techniques.

Approaches: quality control and uncertainty

A specific requirement of the WFD is that Member States support their assessments of Ecological Status of water bodies with estimates of the level of confidence and precision of the results of the monitoring programmes. The sources of uncertainty associated with results will include components resulting from each of the sampling and surveying process, the sorting of samples, identification of the sorted material, data logging and the precision of the models, hind casting or other procedures used to set the reference condition to calculate EQR values.

The assessment of many of these sources of uncertainty has been researched by Clarke (2000) and Clarke et al. (2002, 2003). In the STAR project, focus was on the uncertainty associated with sampling, sample sorting and the identification of specimens, where errors of data logging were considered together with identification errors.

A presumption of a well-implemented monitoring programme is that the persons responsible for carrying out each stage in the process are well trained and competent in the tasks that they are undertaking. In the STAR project partners carrying out the sampling process initially had variable experience of the tasks that they were required to perform. Even where they were experienced and proficient in parts of some tasks, such as collecting macroinvertebrates using their specific national method, they were often less well trained and experienced in collecting STAR-AQEM samples.

Sampling and survey training and identification courses

Prior to any sampling, extensive training course were arranged in the field and laboratory procedures to be used. Representatives of all partners were trained in diatom sampling and preservation, MTR procedures and River Habitat Survey. Particular emphasis was based on consistent application of the STAR-AQEM site and sampling procedures since these were the common standards against which other sampling protocols were to be

compared. An initial week's training course in France included training sessions in sampling of diatoms, macrophytes and macroinvertebrates, including RIVPACS sampling training for the Austrian, German and Greek partners, who were using it as their 'national' method. The specialist diatom and macrophyte trainers were international experts Martyn Kelly and Nigel Holmes respectively. Macroinvertebrate sampling training was provided by highly experienced STAR internal partners.

The French training course also included the requisite three day RHS accreditation course that required all participants to pass a rigorous exam in the application of the method before they were able to undertake it in STAR. The training and accreditation was led by Helena Parsons of the UK Environment Agency.

The French course was supplemented by additional macroinvertebrate sampling training in Denmark and Poland. A separate diatom and macrophyte training course also took place in Poland and included the three day RHS training course. Additionally specialist training courses in the processing and identification of diatoms and identification courses on Oligochaeta, Plecoptera and Trichoptera were organised by expert STAR partners for other STAR partners and external scientists.

Diatom ring test, replicate sampling programme and audit

A diatom ring test was undertaken during the French training course. It was used to compare the results of simultaneous sampling of diatoms by the STAR personnel responsible for sampling this biological quality element during the main sampling programme. Parameters compared included intra- and inter-substratum variability and intra- and inter-operator variability in the type and relative proportions of taxa collected and identified and the indices derived from the results of sampling.

For the ring test, samples were collected from two locations on the Plaine River, in the Vosges region of France. Samples were collected from three different habitat types; stones, macrophytes and sediments. Sampling methods followed the STAR sampling protocol (see above). Each

partner collected three samples from each of two substrata at each of the two test sites. The participants who collected the samples also prepared the samples in their respective laboratories and identified and counted a minimum of 300 valves. The results of the ring test, including identification checks were evaluated by specialists at STAR partner Alterra (Besse-Lotoskaya et al., 2006).

An additional replicate sampling programme was also carried out by partners in the Czech Republic, France, Greece, Portugal, Sweden and the United Kingdom (Table 8). All samples were prepared and identified by the organisation that collected them.

All partners collecting and processing diatom samples for WP7 and/or WP8 were subject to auditing of their taxon counts and identifications. Thirty-eight percent of all core and additional stream samples were subject to audit by experts at Alterra. The samples to be re-analysed were selected randomly from all the samples taken by each partner. Therefore, all samples were numbered and the numbers of samples to be audited were selected using a list of random digits.

The identification of taxa was initially to the most precise taxonomic level that was achievable (species or variety/forma). Subsequently, following discussions amongst the STAR partners, the level of identification of some difficult taxa was made less rigorous. This provided a more consistent level of achievable identification but one that remained compatible with the metrics to be used for Ecological Status assessments. After resolving nomenclatural differences, the results of taxa and counts obtained by the primary analysts and the auditor were compared to determine the error rates (Besse-Lotoskaya et al., 2006).

Macroinvertebrate replicate sampling programme and audit

A replicate macroinvertebrate sampling programme was undertaken by all partners involved in WP7 and/or WP8 except in Slovakia. Replicate sampling was undertaken at 80 sites (Table 7). At each replicate site two STAR-AQEM samples and two national samples were collected in the same 100 m section of the river on the same visit in each of the two macroinvertebrate sampling

Table 8. The number of sites subject to replicate diatom and/or macroinvertebrate sampling in the main WP7 and WP8 programmes

Country	Stream type	Number of replicate sites	
		Diatoms: one sample per site in one sampling season	Macroinvertebrates: two samples per method per site in each of two sampling seasons
Austria	A05	0	3
	A06	0	3
Czech Republic	C04	3	3
	C05	3	3
Denmark	K02	0	6
France	F08	3	6
Germany	D03	0	2
	D04	0	2
	D06	0	4
Greece	H04	6	6
Italy	I05	0	0
	I06	0	6
Latvia	L02	0	6
Poland	O02	0	3
	O03	0	3
Portugal	P04	6	6
Slovakia	V01	0	0
	V02	0	6
Sweden	S05	1	3
	S06	5	3
United Kingdom	U15	3	3
	U23	3	3
Totals		33 (33 samples)	80 (320 samples)

seasons. In the case of STAR-AQEM samples, two separate blind estimates of the proportion of the habitats present were normally made and the number of sample units on each habitat in each sample was based on their respective substratum recording forms. Additional replicate sampling was undertaken at some sites as itemised by Clarke et al. (2006a, 2006b).

Sorting and processing of all main and replicate samples were undertaken by the partner collecting the samples. Each partner also conducted a separate replicate sub-sampling programme for their STAR-AQEM samples where feasible. Replicate sub-sampling was attempted on all of the 160 replicate STAR-AQEM samples. In this programme a standard STAR-AQEM sub-sample was processed for the replicate sample, involving the sorting of either five cells or the number of cells needed to obtain the 700+ specimens required. On

completion of this sub-sample a second sub-sample was processed using the material from five, or more if necessary, of the remaining cells from the sample tray grid. Replicate sub-sampling could not be undertaken in specimen-poor sites, particularly in Greece, where two sub-samples of 700+ specimens could not be achieved.

An audit programme was undertaken involving re-sorting and re-identification of samples collected and first processed by partners. Audits were undertaken on replicate samples collected as part of the replicate sampling programme. A single replicate sample was audited for each method at each site with half the audited samples being collected in spring and half in the second sampling season. The single exception was Italy where it was only possible to audit spring samples for operational reasons. For STAR-AQEM samples the first sub-sample of the replicate sub-sampling programme was

normally selected for auditing but occasionally the second sub-sample was audited instead.

Partners were not informed of which samples had been selected for auditing, nor that only replicate samples would be chosen, until all project samples had been processed. Prior to notification of which samples were to be audited, partners were required to provide the auditors with copies of their full taxon lists for all samples. This prevented modification of results prior to dispatching the notified samples for audit.

The audit programme was in two parts. The first part was the sorting audit to record any families of macroinvertebrates that had not been removed by the partner of origin of the sample. Representative specimens of all taxa present, including those not found by the original partner were removed from the sample and retained for further audit. Sorting audits were all undertaken by the Centre for Ecology and Hydrology except for their own samples, which were audited by the University of Duisburg-Essen.

The second stage of the audit process was the identification audit. This involved re-identification of all the taxa removed from the sample and identified by the partner of origin plus the identification of all of the additional specimens, including any new families, removed from the sample during the sorting audit process. Identification audits were shared amongst most STAR partners with the auditing partner being in the same, adjacent or similar eco-region to that of the audited partner and therefore familiar with the majority of taxa in the audited partner's region.

The output of the auditing process was a list of the families gained to samples by the sorting audit, which were genuine errors, plus a comparison of the two different lists of taxa separately identified by the audited and auditing partner. The latter differences were perceived to be differences of opinion since no arbitration or consensus of identifications were attempted. The audit therefore compared the uncertainty involved in two different experts both identifying the same set of specimens.

Approaches: software development

The project data were curated, managed and analysed using a series of bespoke database and

software products developed by STAR or modified from existing software created within the AQEM project or by the Centre for Ecology and Hydrology (Table 9). The STAR website (www.eu-star.at) provided an internal discussion forum, data repository and portal to the outside world for information, ideas and reports developed during the project and the protocols used in the project for data collection. Access routes to the databases and software products in the public domain, including the Decision Support System, MONSTAR, developed as a project deliverable, are provided in Table 9.

In addition to the software developed in the STAR, AQEM and Euro-limpacs projects, phyto-benthos metric values were calculated using the version 3.2 of the Omidia software (Lecoite et al., 1993). Fish data collected in the STAR project were stored and retrieved using the Fides software developed by the FAME project. Fides software was used to calculate metric values including the new European Fish Index and classification system (EFI) developed by FAME. FAME software may be accessed from <http://fame.boku.ac.at/downloads.htm>.

Approaches: output

STAR has been one of the central research projects contributing to the implementation of the Water Framework Directive. The STAR research programme has made significant contributions to this process through membership of, or formal advice to the Common Implementation Strategy (CIS) Working Groups, including ECOSTAT and the Geographical Inter-calibration Groups and to the Comité Européen de Normalisation (CEN) responsible for producing the methodological standards for use in conjunction with the WFD monitoring programmes.

The detailed programme of work undertaken by the group, the project deliverables, in the form of reports, data and software and much other information, including the Waterview database (Birk & Hering, 2002) are available from the project website (www.eu-star.at). The objective of this special issue of *Hydrobiologia* has been to make the major findings of the project available to a wider audience via a series of individual papers.

Table 9. STAR bespoke databases and software

Product	Brief product description	Access route
AQEMDip	An Access [®] -based program for storage and retrieval of macroinvertebrate and phytobenthos data in standard input and output formats. Output macroinvertebrate data are supplied in the input format structure for ASTERICS. AQEMDip was developed using the platform of the Austrian assessment program ECOPROF. AQEMDip and ECOPROF were developed by Softwarehaus Graf and Partner	AQEMDip software and manual are available for download from the STAR website (www.eu-star.at). ECOPROF is available from www.ecoprof.at
STARMTRI	A database for the storage and retrieval of macrophyte data collected using Mean Trophic Rank survey and similar procedures. The Access [®] -based program has functions for standardising (adjusting) taxonomic levels of surveys prior to analysis. Standard taxonomic adjustments are provided but user defined adjustments may also be created. Mean Trophic Rank scores may be calculated in order to assess the degree to which a stream is impacted by organic pollution or enrichment. The STAR software is an extension of a program first developed by the Centre for Ecology and Hydrology	STARMTRI software, with a detailed internal help system, is available for download from the STAR website (www.eu-star.at).
STARRHSI	An Access [®] -based program for storage and retrieval of survey data collected using the River Habitat Survey (RHS) methodology. The software is compatible with the 1997, 2001 and 2003 versions of the British RHS system and also the 2001 southern European modification (Buffagni & Kemp, 2002). STARRHSI can be used to calculate the Habitat Quality Assessment (HQA) and Habitat Modification Score (HMS) indices. The STAR software is an extension of a program first developed by the Centre for Ecology and Hydrology	STARRHSI software, with a detailed internal help system, is available for download from the STAR website (www.eu-star.at)
ASTERICS	A program for calculating the values of approximately 200 macroinvertebrate metrics used to assess the Ecological Status of running waters. The program, referred to as the AQEM River Assessment Program can also be used to calculate the values of multi-metric systems developed for specific stream, stressor and/or countries during the AQEM project (Hering et al., 2004a) and several national projects. The program was developed by Alterra Green World Research and by the University of Duisburg-Essen, using funds from the European Union (AQEM and STAR) and national funding from Germany (Umweltbundesamt, Länderarbeitsgemeinschaft Wasser)	Software and manual are available from: www.eu-star.at , www.aqem.de and www.fliessgewaesserbewertung.de

Macroinvertebrate Taxa and Autecology Database

A database of approximately 10 000 macroinvertebrate taxa occurring in Europe and the associated taxon codes applied to each taxon in each of six different European systems. The database was principally developed during the AQEM project, extended in the STAR project and is now being further developed and enhanced with comprehensive autecological data in the EC PPVI project Euro-limpacs (GOCE-CT-2003-505540). The database has been developed under the management of the University of Duisburg-Essen and BOKU, the University of Natural Resources and Applied Life Sciences, Vienna

STARBUGS

STARBUGS is an Excel[®]-based system for estimating the impact of known sources and levels of uncertainty on the values of single and multi-metrics and on the confidence and precision with which metric values can be assigned to individual classes of Ecological Status. STARBUGS may be used to test the development of multi-metrics. The system was designed for running waters but is portable to other surface waters or even to the terrestrial environment. STARBUGS was developed by the Centre for Ecology and Hydrology

MONSTAR

MONSTAR is a software product that provides guidance in the design of monitoring programmes necessary to meet the terms and objectives of the Water Framework Directive (WFD). MONSTAR gives insight in the consequences of different choices in terms of costs and results that enables users to optimize their monitoring programme. MONSTAR specifically focus on running waters and addresses stream types and stressors studied in the STAR project with a strong focus on macro-invertebrates. However, MONSTAR provides a framework for including additional information related to other stream types, stressors and quality elements. MONSTAR was developed by Alterra Green World Research

The database taxon lists and codes can be accessed on the Euro-limpacs website at www.freshwaterecology.info

STARBUGS and its associated manual are available for download from the STAR website (www.eu-star.at)

Each of these papers directly or indirectly addresses particular practical issues faced by the CIS and by those charged with implementing the WFD. For this purpose, the volume has been divided into a series of sections devoted to specific generic issue and each comprising a set of two or more papers. The seven component sections are (1) typology, (2) organism groups, (3) macrophytes and diatoms, (4) hydromorphology, (5) tools for assessing European streams with macroinvertebrates, (6) intercalibration and comparison and (7) errors and uncertainty. This structure mirrors the sequence of practical considerations that need to be addressed in delivering a coherent monitoring programme for evaluating the Ecological Status of streams and rivers within the European Union.

The objective of this paper, as its title implies, has been to introduce the project and to set out the context, objectives and approaches taken to provide some of the scientific information needed to best implement the WFD. In doing so it provides a reference point for many of the individual papers contained within this volume and obviates the necessity to repeat this information elsewhere. In order to assimilate the key findings of the research programme, as presented here, each of the seven separate sections of this volume is prefaced by a summary paper drawing out the key results of the component papers and highlighting the recommendations of these papers and their practical contribution to the implementation process.

Acknowledgements

STAR was funded by the European Commission, 5th Framework Program, Energy, Environment and Sustainable Development, Key Action Water, contract no. EVK1-CT-2001-00089. AQEMDip and ASTERICS (formerly the AQEM River Assessment Program) were initially developed during the European Commission, 5th Framework Project AQEM, contract No. EVK1-CT1999-00027. ASTERICS was co-funded by Umweltbundesamt, Länderarbeitsgemeinschaft Wasser. The authors acknowledge the support of all their colleagues, too numerous to mention, who have contributed to the work described here. We are grateful to Stefan Schmutz and his colleagues in the FAME project for the use of the Fides

database and their EFI software. We also wish to thank Hartmut Barth, Mogens Gadeberg and Elena Domínguez, EC Scientific Officers for STAR for their constant help and encouragement.

References

- AQEM Consortium, 2002. Manual for the application of the AQEM system. A comprehensive method to assess European streams using benthic macroinvertebrates, developed for the purpose of the Water Framework Directive. Version 1.0, February 2002.
- Balon, E. K., 1975. Reproductive guilds of fishes: a proposal and definition. *Journal of the Fisheries Research Board of Canada* 32: 821–864.
- Baatrup-Pedersen, A., K. Szoszkiewicz, R. Nijboer, M. O'Hare & T. Ferreira, 2006. Macrophyte communities in unimpacted European streams: variability in assemblage patterns, abundance and diversity. *Hydrobiologia* 566: 179–196.
- Besse-Lotoskaya, A., P. F. M. Verdonchot & J. A. Sinkeldam, 2006. Uncertainty in diatom assessment: Sampling, identification and counting variation. *Hydrobiologia* 566: 247–260.
- Birk, S. & D. Hering, 2002. Waterview Web-Database: a comprehensive review of European assessment methods for rivers. *FBA News* 20: 4.
- Buffagni, A., S. Erba, M. Cazzola, J. Murray-Bligh, H. Soszka & P. Genoni, 2006. The STAR common metrics approach to the WFD intercalibration process: Full application for small, lowland rivers in three European countries. *Hydrobiologia* 566: 379–399.
- Buffagni, A. & L. J. Kemp, 2002. Looking beyond the shores of the United Kingdom: addenda for the application of River Habitat Survey in southern European rivers. *Journal of Limnology* 61: 199–214.
- Birk, S. & D. Hering, 2006. Direct comparison of assessment methods using benthic macroinvertebrates: a contribution to the EU Water Framework Directive intercalibration exercise. *Hydrobiologia* 566: 401–415.
- Birk, S., T. Korte & D. Hering, 2006. Intercalibration of assessment methods for macrophytes in lowland streams: direct comparison and analysis of common metrics. *Hydrobiologia* 566: 417–430.
- Bis, B. & P. Usseglio-Polatera, 2004. Species traits analysis. STAR deliverable N2 to the European Commission, 148 pp.
- Clarke, R. T., 2000. Uncertainty in estimates of river quality based on RIVPACS. In Wright, J. F., D. W. Sutcliffe & M. T. Furse (eds), *Assessing the Biological Quality of Freshwaters: RIVPACS and Similar Techniques*. Freshwater Biological Association, Ambleside, 39–54.
- Clarke, R. T., J. Davy-Bowker, L. Sandin, N. Friberg, R. K. Johnson & B. Bis, 2006a. Estimates and comparisons of the effects of sampling variation using 'national' macroinvertebrate sampling protocols on the precision of metrics used to assess ecological status. *Hydrobiologia* 566: 477–503.
- Clarke, R. T., M. T. Furse, R. J. M. Gunn, J. M. Winder & J. F. Wright, 2002. Sampling variation in macroinvertebrate

- data and implications for river quality indices. *Freshwater Biology* 47: 1735–1751.
- Clarke, R. T., A. Lorenz, L. Sandin, A. Schmidt-Kloiber, J. Strackbein, N. T. Kneebone & P. Haase, 2006b. Effects of sampling and sub-sampling variation using the STAR-AQEM sampling protocol on the precision of macroinvertebrate metrics. *Hydrobiologia* 566: 441–459.
- Clarke, R. T., J. F. Wright & M. T. Furse, 2003. RIVPACS models for predicting the expected macroinvertebrate fauna and assessing the ecological quality of rivers. *Ecological Modelling* 160: 219–233.
- Danish Environmental Protection Agency, 1998. Biological Assessment of Watercourse Quality. Guidelines, No. 5. – Danish Environmental Protection Agency, Ministry of Environment and Energy, Copenhagen.
- Davies, P. E., 2003. Development of a national river bioassessment system (AUSRIVAS) in Australia. In Wright, J. F., D. W. Sutcliffe & M. T. Furse (eds), *Assessing the Biological Quality of Freshwaters: RIVPACS and Similar Techniques*. Freshwater Biological Association, Ambleside, 113–124.
- Davy-Bowker, J., R. T. Clarke, R. K. Johnson, J. Kokes, J. F. Murphy & S. Zahrádková, 2006. A comparison of the European Water Framework Directive physical typology and RIVPACS-type models as alternative methods of establishing reference conditions for benthic macroinvertebrates. *Hydrobiologia* 566: 91–105.
- Erba, S., A. Buffagni, N. Holmes, M. O'Hare, P. Scarlett & A. Stenico, 2006. Preliminary testing of River Habitat Survey features for the aims of the WFD hydro-morphological assessment: an overview from the STAR Project. *Hydrobiologia* 566: 281–296.
- European Commission, 2000. Directive of the European Parliament and of the Council 2000/60/EC establishing a framework for Community action in the field of water policy. European Commission PE-CONS 3639/1/00 REV 1, Luxembourg.
- European Commission, 2001. Common implementation strategy for the Water Framework Directive (2000/60/EC). Strategic document as agreed by the Water Directors under Swedish presidency, 2 May 2001. European Commission, 81 pp.
- European Commission, 2002. Water Framework Directive (WFD) Common Implementation Strategy Working Group 2.5. Intercalibration: Towards a guidance on establishment of the intercalibration network and on the process of the intercalibration exercise. European Commission, 50 pp.
- European Commission, 2003a. Water Framework Directive (WFD) Common Implementation Strategy Working Group 2.3 Reference conditions for inland surface waters (REFCOND). Guidance on establishing reference conditions and Ecological Status class boundaries for inland surface waters. Final version, 30 April 2003. European Commission, 86 pp.
- European Commission, 2003b. Water Framework Directive Common Implementation Strategy Working Group 2.7 Monitoring. Guidance on monitoring for the Water Framework Directive. Final version. 23 January 2003. European Commission, 170 pp.
- European Commission, 2003c. Water Framework Directive Common Implementation Strategy Working Group 2.A Ecological Status (ECOSTAT). Overall approach to the classification of Ecological Status and Ecological Potential. European Commission, 86 pp.
- GAY, Cabinet en Environnement, 1994. Indice Biologique Global Normalisé. NF T 90-350. Guide Technique. Agences de l'eau et Ministère de l'Environnement, Paris.
- Friberg, N., L. Sandin, M. T. Furse, S. E. Larsen, R. T. Clarke & P. Haase, 2006. Comparison of macroinvertebrate sampling methods in Europe. *Hydrobiologia* 566: 365–378.
- Ghetti, P. E., 1997. Manuale di Applicazione. Indice Biotico Esteso (I. B. E.). I Macroinvertebrati Nell Controllo Della Qualità Degli Ambienti di Acque Correnti. – Provincia Autonoma di Trento, Agenzia Provinciale per la Protezione dell'Ambiente, Trento.
- Haase, P., J. Murray-Bligh, S. Lohse, S. Pauls, A. Sundermann, R. Gunn & R. Clarke, 2006. Assessing the impact errors in sorting and identifying macroinvertebrate samples. *Hydrobiologia* 566: 505–521.
- Hellawell, J. M., 1978. *The Biological Surveillance of Rivers: A Biological Monitoring Handbook*. Water Research Centre, Stevenage.
- Hellawell, J. M., 1986. *Biological Indicators of Freshwater Pollution and Environmental Management*. Pollution Monitoring Series. Elsevier Applied Science, London, New York.
- Hering, D., A. Buffagni, O. Moog, L. Sandin, M. Sommerhäuser, I. Stubauer, C. Feld, R. K. Johnson, P. Pinto, N. Skoulikidis, P. F. M. Verdonchot & S. Zahrádková, 2003. The development of a system to assess the ecological quality of streams based on macroinvertebrates – design of the sampling programme within the AQEM project. *Internationale Revue der gesamten Hydrobiologie* 88: 345–361.
- Hering D., C. K. Feld, O. Moog & T. Ofenböck, 2006. Cookbook for the development of a Multimetric-Index for biological condition of aquatic ecosystems: experiences from the European AQEM and STAR projects and related initiatives. *Hydrobiologia* 566: 311–324.
- Hering, D., O. Moog, L. Sandin & P. F. M. Verdonchot, 2004a. Overview and application of the AQEM assessment system. *Hydrobiologia* 516: 1–21.
- Hering, D., P. F. M. Verdonchot, O. Moog & L. Sandin (eds), 2004b. Integrated assessment of running waters in Europe. *Hydrobiologia*. 516 pp.
- Holmes, N. T. H., J. R. Newman, S. Chadd, K. J. Rouen, L. Sharp & F. H. Dawson, 1999. Mean Trophic Rank: A Users' Manual. R&D Technical Report No. E38, Environment Agency, Bristol.
- Holmes, N. T. H. & B. A. Whitton, 1977. Macrophytic vegetation of the River Swale, Yorkshire. *Freshwater Biology* 7: 545–558.
- Hughes, R. M., 1995. Defining acceptable status by comparing with reference conditions. In Davis, W. S. & T. P. Simon (eds), *Biological Assessment and Criteria. Tools for Water Resource Planning and Decision Making*. Lewis Publishers, Boca Raton, FL, 31–47.
- Illies, J. (ed.), 1978. *Limnofauna Europaea*, 2nd edn. Gustav Fischer Verlag, Stuttgart, New York; Swets and Zeitlinger B. V., Amsterdam.
- Johnson, R. K., D. Hering, M. T. Furse & R. T. Clarke, 2006a. Detection of ecological change using multiple organism groups: metrics and uncertainty. *Hydrobiologia* 566: 115–137.

- Johnson, R. K., D. Hering, M. T. Furse & P. F. M. Verdonchot, 2006b. Indicators of ecological change: comparison of the early response of four organism groups to stress gradients. *Hydrobiologia* 566: 139–152.
- Kelly, M. G., A. Cazaubon, E. Coring, A. Dell'Uomo, L. Ector, B. Goldsmith, H. Guasch, J. Hürlimann, A. Jarlman, B. Kawecka, J. Kwadrans, R. Laugaste, E.-A. Lindström, M. Leitao, P. Marvan, J. Padisák, E. Pipp, J. Prygiel, E. Rott, S. Sabater, H. van Dam & J. Vizinet, 1998. Recommendations for the routine sampling of diatoms for water quality assessments in Europe. *Journal of Applied Phycology* 10: 215–224.
- Knoben, R. A. E., C. Roos & M. C. van Oirschot, 1995. Biological assessment methods for watercourses. UN/ECE Task Force on Monitoring and Assessment, 86 pp.
- Kokeš, J., S. Zahradková, D. Němejcová, J. Hodovský, J. Jarkovský & T. Soldán, 2006. The PERLA system in the Czech Republic: a multivariate approach for assessing the ecological status of running waters. *Hydrobiologia* 566: 343–354.
- Latvian Standard Ltd., 1999 LVS 240:1999 Water quality – operative evaluation biological quality of small stream by saprobity index of macrozoobenthos community. In Catalogue of Latvian standards, Riga, Latvian Standard Ltd, 1999: Group 13.060, 1(11)–11.
- Lecointe, C., M. Coste & J. Prygiel, 1993. “OMNIDIA” software for taxonomy, calculation of diatom indices and inventories management. *Hydrobiologia* 269/270: 509–513.
- Lorenz, A. & R. T. Clarke, 2006. Sample coherence – a field study approach to assess similarity of macroinvertebrate samples. *Hydrobiologia* 566: 461–476.
- Mandl, V., 1992. Draft EC directive on ecological quality of surface waters. In Newman, P. J., M. A. Piavaux & R. A. Sweeting (eds), *River Water Quality. Ecological Assessment and Control*. Publication EUR 14606 EN-FR. Commission of the European Communities, Luxembourg, 18.
- Mann, R. H. K., 1996. Environmental requirements of European non-salmonid fish in rivers. *Hydrobiologia* 323: 223–235.
- Metcalf, J. L., 1989. Biological water quality assessment of running waters based on macroinvertebrate communities: history and present status in Europe. *Environmental Pollution* 60: 101–139.
- Metcalf-Smith, J. L., 1994. Biological water quality assessment of rivers: use of macroinvertebrate communities. In Calow, P. & G. E. Petts (eds), *The Rivers Handbook*, Vol. II. Blackwell Scientific Publications, London, 144–170.
- Moore, W. W., 1977. Seasonal succession of algae in a eutrophic stream in southern England. *Hydrobiologia* 53: 181–192.
- Murray-Bligh, J. A. D., M. T. Furse, F. H. Jones, R. J. M. Gunn, R. A. Dines & J. F. Wright, 1997. Procedure for collecting and analysing macroinvertebrate samples for RIVPACS. Joint publication by the Institute of Freshwater Ecology and the Environment Agency, 162 pp.
- Noble, R. & I. Cowx, 2002. Development, evaluation & implementation of a standardised fish-based assessment method for the Ecological Status of European rivers – a contribution to the water framework directive (FAME). A report to the European Commission, 100 pp.
- Norris, R. H., 1994. Rapid biological assessment, natural variability and selecting reference sites. Classification of rivers and environmental health indicators. In Uys, M. C. (ed.), *Proceedings of a Joint South African/Australian Workshop*, Cape Town, South Africa. Water Research Commission, Report No. TT/63/94: 129–166.
- O'Hare, M. T., A. Baatrup-Pedersen, R. Nijboer, K. Szoszkiewicz & T. Ferreira, 2006. Macrophyte communities of European streams with altered physical habitat. *Hydrobiologia* 566: 197–210.
- Pinto, P., M. Morais, M. Ilhéu & L. Sandin, 2006. Relationships among biological elements (macrophytes, macroinvertebrates and ichthyofauna) for different river types across Europe at two different spatial scales. *Hydrobiologia* 566: 75–90.
- Raven, P. J., N. T. H. Holmes, F. H. Dawson, P. J. A. Fox, M. Everard, I. R. Fozzard & K. J. Rouen, 1998. *River Habitat Quality – The Physical Character of Rivers and Streams in the UK and Isle of Man*. River Habitat Survey Report Number 2. Environment Agency, Bristol: Scottish Environment Protection Agency, Stirling: Environment and Heritage Service, Belfast, 1–84.
- Reynoldson, T. B., R. C. Bailey, K. E. Day & R. H. Norris, 1995. Biological guidelines for freshwater sediment based on Benthic Assessment of Sediment (the BEAST) using a multivariate approach for predicting biological state. *Australian Journal of Ecology* 20: 198–219.
- Reynoldson, T. B., K. E. Day & T. Pascoe, 2000. The development of the BEAST: a predictive approach for assessing sediment quality in the Great Lakes. In Wright, J. F., D. W. Sutcliffe & M. T. Furse (eds), *Assessing the Biological Quality of Freshwaters: RIVPACS and Similar Techniques*. Freshwater Biological Association, Ambleside, 165–180.
- Rosenberg, D. M., T. B. Reynoldson & V. H. Resh, 2000. Establishing reference conditions in the Fraser River catchment, British Columbia, Canada, using the BEAST (Benthic Assessment of Sediment) predictive model. In Wright, J. F., D. W. Sutcliffe & M. T. Furse (eds), *Assessing the Biological Quality of Freshwaters: RIVPACS and Similar Techniques*. Freshwater Biological Association, Ambleside, 181–194.
- Schmidt-Kloiber, A., W. Graf, A. Lorenz & O. Moog, 2006. The AQEM/STAR taxalist – a pan-European macroinvertebrate ecological database and taxa inventory. *Hydrobiologia* 566: 325–342.
- Seber, G. A. F. & E. D. Le Cren, 1967. Estimating population parameters from catches large relative to the population. *Journal of Animal Ecology* 36: 631–643.
- Sladeczek, V., 1973. System of water quality from the biological point of view. *Archiv für Hydrobiologie Ergebnisse der Limnologie* 7: 1–218.
- Šporka, F., H. E. Vlek, E. Bulánková & I. Krno, 2006. Influence of seasonal variation on bioassessment of streams using macroinvertebrates. *Hydrobiologia* 566: 543–555.
- Springe, G., L. Sandin, A. Briede & A. Skuja, 2006. Biological quality metrics: their variability and appropriate scale for assessing streams. *Hydrobiologia* 566: 153–172.
- Staniszewski, R., K. Szoszkiewicz, J. Zbierska, J. Lesny, S. Jusik & R. T. Clarke, 2006. Assessment of sources of uncertainty in macrophyte surveys and the consequences for river classification. *Hydrobiologia* 566: 235–246.

- Swedish Environmental Protection Agency, 1996. Bottenfauna i sjöars litoral och i vattendrag – tidsserier. [In Swedish: Benthic fauna in lake litoral and running waters – time series]. Swedish EPA monitoring handbook, Fresh waters. [Published digitally at: www.naturvardsverket.se].
- Szoszkiewicz, K., A. Buffagni, J. Davy-Bowker, J. Lesny, B. H. Chojnicki, J. Zbierska, R. Staniszewski & T. Zgola, 2006a. Occurrence and variability of River Habitat Survey features across Europe and the consequences for data collection and evaluation. *Hydrobiologia* 566: 267–280.
- Szoszkiewicz, K., T. Ferreira, T. Korte, A. Baattrup-Pedersen, J. Davy-Bowker & M. O'Hare, 2006b. European river plant communities: the importance of organic pollution and the usefulness of existing macrophyte metrics. *Hydrobiologia* 566: 211–234.
- Verdonschot, P. F. M., 2006a. Evaluation of the use of Water Framework Directive typology descriptors, reference sites and spatial scale in macroinvertebrate stream typology. *Hydrobiologia* 566: 39–58.
- Verdonschot, P. F. M., 2006b. Data composition and taxonomic resolution in macroinvertebrate stream typology. *Hydrobiologia* 566: 59–74.
- Vlek, H. E., F. Šporka & I. Krno, 2006. Influence of macroinvertebrate sample size on bioassessment of streams. *Hydrobiologia* 566: 523–542.
- Winter, J. G. & H. C. Duthie, 2000. Stream epilithic, epipellic and epiphytic diatoms: habitat fidelity and use in biomonitoring. *Aquatic Ecology* 34: 345–353.
- Wright, J. F., P. D. Armitage & M. T. Furse, 1989. Prediction of invertebrate communities using stream measurements. *Regulated Rivers: Research and Management* 4: 147–155.
- Wright, J. F., D. W. Sutcliffe & M. T. Furse (eds), 2000. *Assessing the Biological Quality of Freshwaters: RIVPACS and Similar Techniques*. Freshwater Biological Association, Ambleside.