

The Structure of Maximum Subsets of $\{1, \dots, n\}$ with No Solutions to $a + b = kc$

Andreas Baltz*

Mathematisches Seminar
University of Kiel, D-24098 Kiel, Germany
aba@numerik.uni-kiel.de

Peter Hegarty, Jonas Knape, Urban Larsson

Department of Mathematics
Chalmers University of Technology
Göteborg, Sweden
hegarty@math.chalmers.se {md9jonas,md0larur}@mdstud.chalmers.se

Tomasz Schoen **

Wydział Matematyki i Informatyki
Adam Mickiewicz University
Poznań, Poland
schoen@amu.edu.pl

Submitted: Nov 9, 2004; Accepted: Apr 22, 2005; Published: Apr 28, 2005

MR Subject Classifications: 05D05, 11P99

Abstract

If k is a positive integer, we say that a set A of positive integers is k -sum-free if there do not exist a, b, c in A such that $a + b = kc$. In particular we give a precise characterization of the structure of maximum sized k -sum-free sets in $\{1, \dots, n\}$ for $k \geq 4$ and n large.

1 Introduction

A set of positive integers is called k -sum-free if it does not contain elements a, b, c such that

$$a + b = kc,$$

*supported by DFG, Grant SR 7/9 – 2

**research partially supported by KBN Grant 2 PO3A 007 24

where k is a positive integer. Denote by $f(n, k)$ the maximum cardinality of a k -sum-free set in $\{1, \dots, n\}$. For $k = 1$ these extremal sets are well-known: Deshouillers, Freiman, Sós, and Temkin [1] proved in particular that the maximum 1-sum-free sets in $\{1, \dots, n\}$ are precisely the set of odd numbers and the “top half” $\{\lceil \frac{n+1}{2} \rceil, \dots, n\}$. For $n > 8$ even $\{\frac{n}{2}, \dots, n-1\}$ forms the only additional extremal set. The famous theorem of Roth [4] gives $f(n, 2) = o(n)$. Chung and Goldwasser [2] solved the case $k = 3$ by showing that the set of odd integers is the unique extremal set for $n > 22$. For $k \geq 4$ they gave an example of a k -sum-free set [3] of cardinality $\frac{k(k-2)}{k^2-2}n + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)}n + \mathcal{O}(1)$, which implies $\lim_{n \rightarrow \infty} \frac{f(n, k)}{n} \geq \frac{k(k-2)}{k^2-2} + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)}$, and they conjectured that this lower bound is the actual value. Moreover they conjectured that extremal k -sum-free sets consist of three intervals of consecutive integers with slight modifications at the end-points if n is large.

In this paper we prove that the first conjecture is true, and we expose a structural result which is very close to the second. Our proof is elementary. In fact it is based on two simple observations:

Suppose we are given a k -sum-free set A . Then

- $kx - y \notin A$ for all $x, y \in A$
(Otherwise we could satisfy the equation $kx = (kx - y) + y$ in A .)
- for all $y \in A$ any interval centered around $\frac{ky}{2}$ cannot share more than half of its elements with A .
(Otherwise we would find a pair $\lfloor \frac{ky}{2} \rfloor - d, \lceil \frac{ky}{2} \rceil + d$ in A , giving $(\lfloor \frac{ky}{2} \rfloor - d) + (\lceil \frac{ky}{2} \rceil + d) = ky$.)

2 Preparations

Let $n \in \mathbb{N}$ be large and let $k \in \mathbb{N}_{\geq 4}$. We start by agreeing on some notations.

Notations

Let $A \subseteq \{1, \dots, n\}$ be a set of positive integers. Denote by

$$s_A := \min A \text{ and } m_A := \max A$$

the smallest and the largest elements of A respectively.

For $l, r \in \mathbb{R}$ let

$$\begin{aligned} (l, r] &:= \{x \in \mathbb{N} \mid l < x \leq r\} \\ [l, r) &:= \{x \in \mathbb{N} \mid l \leq x < r\} \\ (l, r) &:= \{x \in \mathbb{N} \mid l < x < r\} \\ [l, r] &:= \{x \in \mathbb{N} \mid l \leq x \leq r\} \end{aligned}$$

abbreviate intervals of integers. Continuous intervals will be indicated by the subscript \mathbb{R} .

Furthermore for any $y \in \mathbb{N}$ and $d \in \mathbb{N}_0(:= \mathbb{N} \cup \{0\})$ put

$$I_y^d := \left[\frac{ky - 1}{2} - d, \frac{ky + 1}{2} + d \right].$$

Note that if ky is even then $I_y^d = \left\{ \frac{ky}{2} - d, \frac{ky}{2} - d + 1, \dots, \frac{ky}{2} + d \right\}$ and $|I_y^d| = 2d + 1$, while if ky is odd we have $I_y^d = \left\{ \frac{ky-1}{2} - d, \dots, \frac{ky+1}{2} + d \right\}$ and $|I_y^d| = 2d + 2$.

The first Lemma restates our introductory observations.

Lemma 1 *Let $A \subseteq [1, n]$ be a k -sum-free set. If $x, y \in A$ then $kx - y \notin A$. If $y \in A$ and $d \in \mathbb{N}_0$ then $|I_y^d \setminus A| \geq d + 1$.*

Suppose A' is a k -sum-free set consisting of intervals $(l_i, r_i]$. The interval $(l_i, r_i]$ is k -sum-free if $l_i \geq \frac{2r_i}{k}$. Moreover we observe that reasonably large consecutive intervals $(l_{i+1}, r_{i+1}], (l_i, r_i]$ (where we assume $r_{i+1} < l_i$) should satisfy $kr_{i+1} \leq l_i + s_{A'}$. This leads to the following definition, describing a successive transformation of an arbitrary k -sum-free set A into a k -sum-free set of intervals.

Definition 1 *Let $n \in \mathbb{N}$ and let $A \subseteq [1, n]$ be k -sum-free with smallest element $s := s_A$. Define sequences (r_i) , (l_i) , (A_i) by:*

$$\begin{aligned} A_0 &:= A, \quad r_1 := n, \\ l_i &:= \left\lfloor \frac{2r_i}{k} \right\rfloor, \quad r_{i+1} := \left\lfloor \frac{l_i + s}{k} \right\rfloor, \\ A_i &:= (A_{i-1} \setminus (r_{i+1}, l_i]) \cup (l_i, r_i] \cap [s, n] \text{ for } i \geq 1. \end{aligned}$$

The letter $t = t_A$ will be reserved to denote the least integer such that $r_{t+1} < s$. Observe that, for all $i \geq t$,

$$A_i = A_t = [\alpha, r_t] \cup \left(\bigcup_{j=1}^{t-1} (l_j, r_j] \right), \tag{1}$$

where $\alpha = \alpha_A := \max\{l_t + 1, s\}$.

3 The structure of maximum k -sum-free sets

To obtain the structural result we consider the successive transformation of an arbitrary k -sum-free set A into a set A_t of intervals as in (1). Our plan is to show that each member of the transformation sequence (A_i) is k -sum-free and has size greater than or equal to $|A|$. For n sufficiently large, depending on k , and a maximum sized k -sum-free subset A of $[1, n]$, it will turn out that A_t consists of three intervals only, i.e.: that $t = 3$. This observation will do to determine $f(n, k)$, and we conclude our proof by showing that A

could be enlarged if it did not contain (nearly) the whole interval $(l_3, r_3]$ and consequently almost all elements from $(l_2, r_2]$ and $(l_1, r_1]$, so that in fact almost nothing happens during the transformation of an extremal set.

Lemma 2 *Let $A \subseteq [1, n]$ be k -sum-free. Let $i \in \mathbb{N}$.*

a) A_i is k -sum-free.

b) $|A_i| \geq |A_{i-1}|$.

Proof. a) Clearly, it is enough to prove the claim for $i \leq t$, so we may assume that $s \leq r_i$. Suppose there are $a, b, c \in A_i$ with $a + b = kc$. A_i is of the form

$$A_i = A_{i-1} \cap [s, r_{i+1}] \cup (l_i, r_i] \cap [s, n] \cup (l_{i-1}, r_{i-1}] \cup \dots \cup (l_1, r_1].$$

If $c \in (l_1, r_1]$, then $kc > 2n$, which is impossible. If $i \geq 2$ and $c \in (l_j, r_j]$ for some $j \in [2, i]$, then $kc \in (2r_j, l_{j-1} + s]$ and the larger one of a, b must be in $(r_j, l_{j-1}]$. But $(r_j, l_{j-1}] \cap A_i = \emptyset$ by construction. Hence $c \in A_{i-1} \cap [s, r_{i+1}]$. Now, $kc \leq kr_{i+1} \leq l_i + s$. Since $(r_{i+1}, l_i] \cap A_i = \emptyset$, both a and b have to be in $A_{i-1} \cap [s, r_{i+1}] = A \cap [s, r_{i+1}]$. But A is k -sum-free, a contradiction.

b) The inequality is trivial for $i \geq t$. For $1 \leq i < t$ we have that $l_i \geq s$ and hence

$$A_i = (A_{i-1} \cap [1, r_{i+1}]) \cup (l_i, r_i] \cup \left(\bigcup_{j=1}^{i-1} (l_j, r_j] \right).$$

Thus it suffices to prove that

$$|A_{i-1} \cap [1, r_i]| \leq |A_{i-1} \cap [1, r_{i+1}]| + \left\lceil \frac{(k-2)r_i}{k} \right\rceil.$$

Clearly, then, it suffices to prove the inequality for $i = 1$, i.e.: to prove that, for any $n > 0$, and any k -sum-free subset A of $[1, n]$ with smallest element s_A , we have

$$|A| \leq |A \cap [1, r_{2,A}]| + \left\lceil \frac{(k-2)n}{k} \right\rceil, \tag{2}$$

where

$$r_{2,A} := \left\lceil \frac{\lfloor 2n/k \rfloor + s_A}{k} \right\rceil.$$

The proof is by induction on n . The result is trivial for $n = 1$. So suppose it holds for all $1 \leq m < n$ and let A be a k -sum-free subset of $[1, n]$. Note that the result is again trivial if $s_A > 2n/k$, so we may assume that $s_A \leq 2n/k$, which implies that $r_{2,A} \leq n/k$, since $k \geq 4$.

First suppose that there exists $x \in A \cap (n/k, 2n/k]$. Then $1 \leq kx - n \leq n$ and the

map $f : y \mapsto kx - y$ is a 1-1 mapping from the interval $[kx - n, n]$ to itself. For each y in this interval, at most one of the numbers y and $f(y)$ can lie in A , since A is k -sum-free. To simplify notation, put $w := kx - n - 1$. Then our conclusion is that

$$|A \cap (w, n]| \leq \frac{1}{2}(n - w). \quad (3)$$

If $w = 0$ or if $A \cap [1, w] = \emptyset$, then we are done (since $k \geq 4$). Put $B := A \cap [1, w]$. Then we may assume $B \neq \emptyset$, hence $s_B = s_A$. Applying the induction hypothesis to B , we find that

$$|B| = |A \cap [1, w]| \leq |B \cap [1, r_{2,B}]| + \left\lceil \frac{(k-2)w}{k} \right\rceil. \quad (4)$$

But $s_B = s_A$ implies that $r_{2,B} \leq r_{2,A}$, hence that $B \cap [1, r_{2,B}] \subseteq A \cap [1, r_{2,A}]$. Thus (3) and (4) yield the inequality

$$|A| \leq |A \cap [1, r_{2,A}]| + \left\lceil \frac{(k-2)w}{k} \right\rceil + \frac{1}{2}(n - w),$$

which in turn implies (2), since $|A|$ is an integer. Thus we are reduced to completing the induction under the assumption that $A \cap (n/k, 2n/k] = \emptyset$. Suppose $x \in A \cap (r_{2,A}, n/k]$. Then $\lfloor 2n/k \rfloor + s_A < kx \leq n$ and $kx - s_A \notin A$. In other words, we can pair off elements in $A \cap (r_{2,A}, 2n/k]$ with elements in $(2n/k, n] \setminus A$. This immediately implies (2), and the proof of Lemma 2 is complete. \square

We have seen so far that any k -sum-free set A can be turned into a k -sum-free set A_t having overall size at least $|A|$. The set A_t is a union of intervals, as given by (1), though note that the final interval $[\alpha, r_t]$ may consist of a single point, since $r_t = s$ is possible. The proof of the following Lemma uses a fact shown in [3] by Chung and Goldwasser, to prove that t must be equal to three if $|A|$ is maximum.

Lemma 3 *Let A be a maximum k -sum-free subset of $[1, n]$, where $n > n_0(k)$ is sufficiently large. Let $s := s_A$ and let $t := \max\{i \in \mathbb{N} \mid r_i \geq s\}$. Then $t = 3$.*

Proof. Let A_t be the set of positive integers given by (1). In a similar manner we now define a k -sum-free subset A'_t of $(0, 1]_{\mathbb{R}}$.

Put $c := s/n$ and, for $i = 1, \dots, t$ define real numbers R_i, L_i as follows :

$$R_1 := 1, \quad L_i := \frac{2R_i}{k}, \quad R_{i+1} := \frac{L_i + c}{k}.$$

Then we put

$$A'_t := [\alpha', R_t)_{\mathbb{R}} \cup \left(\bigcup_{j=1}^{t-1} [L_j, R_j)_{\mathbb{R}} \right),$$

where $\alpha' := \max\{L_t, c\}$. That A'_t is k -sum-free is shown in [3]. One sees easily that

$$|A_t| \leq n \cdot \mu(A'_t) + t, \tag{5}$$

where μ denotes the Lebesgue-measure. Now suppose that $t \neq 3$. It is shown in [3] that there exists a constant $c_k > 0$, depending only on k , such that in this case

$$|\mu(A'_t)| \leq \frac{k(k-2)}{k^2-2} + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)} - c_k. \tag{6}$$

In fact, in the notation of page 8 of [3], an explicit value for c_k (which we will use later) is given by

$$c_k = \frac{2}{k}(R(3) - R(4)),$$

which by definition of R amounts to

$$c_k = \frac{8(k^4 - 4k^2 - 4)(k - 2)}{(k^6 - 2k^4 - 4k^2 - 8)(k^4 - 2k^2 - 4)k}. \tag{7}$$

Now (5) and (6) would imply that

$$|A| \leq \frac{k(k-2)}{k^2-2}n + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)}n - c_k n + t.$$

But we have seen in the introduction that $|A| \geq \frac{k(k-2)}{k^2-2}n + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)}n + \mathcal{O}(1)$ and, since $t = \mathcal{O}(\log_k n)$, we thus have a contradiction for sufficiently large n . Hence t must equal three, for large enough n , as required. \square

Now we are nearly in a position to determine $f(n, k)$. We want to calculate the cardinality of an extremal k -sum-free set A via computing $|A_3|$. Since $|A_3|$ depends on s_A , the following lemma will be helpful :

Lemma 4 *Let $n > n_0(k)$ be sufficiently large. If A is a maximal k -sum-free subset of $[1, n]$, then $S - 2k \leq s_A \leq S + 3$, where $S := \lfloor \frac{8n}{k^5 - 2k^3 - 4k} \rfloor$.*

Proof. Set $s := s_A$. By Lemma 3, for $n > n_0(k)$ we have $r_4 < s$. Since A is maximal we have $|A| = |A_3|$. Now, for a fixed n , the cardinality of A_3 is a function of $s \in [1, n]$ only. So we need to show that $|A_3(s)|$ attains its maximum value only for some $s \in [S - 2k, S + 3]$. Define

$$s' := \min\{s \in [1, n] : l_3(s) < s\}.$$

A tedious computation (see the Appendix below) yields that $s' = S + 1$ if k is even and $s' = S$ or $S + 1$ if k is odd. Hence

$$s' \in [S, S + 1]. \tag{8}$$

Clearly,

$$|A_3(s)| = \begin{cases} \lceil \frac{(k-2)n}{k} \rceil + r_2(s) - l_2(s) + r_3(s) - s + 1, & \text{if } s \geq s', \\ \lceil \frac{(k-2)n}{k} \rceil + r_2(s) - l_2(s) + r_3(s) - l_3(s), & \text{if } s < s'. \end{cases} \quad (9)$$

How does $|A_3(s)|$ change (ignoring its maximality for a while) if we alter s ?

First suppose $s \geq s'$. If s increases by one, then $|A_3|$ will decrease by one unless either r_2 or r_3 increases. Now r_2 can only increase (by one) once in $k(\geq 4)$ times. Almost the same is true of r_3 , though its dependence on l_2 makes things a little more complicated. However, it is not hard to see that we encounter an irreversible decrease in the cardinality of $|A_3|$ after at most 3 steps of increment of s . Hence $|A_3(s)| < |A_3(s')|$ if $s \geq s' + 3$.

Next suppose $s < s'$. If we decrease s , then $|A_3|$ cannot increase at all, since l_i will not decrease unless r_i does. Moreover, $|A_3|$ will become smaller if the size of any interval is diminished. So we can focus our attention on $(l_2, r_2]$. While r_2 decreases once in k times, l_2 does so no more than once in $k \lfloor \frac{k}{2} \rfloor \geq 2k$ times. Thus $|A_3(s)| < |A_3(s' - 1)|$ if $s \leq s' - 1 - 2k$.

We have now shown that, as a function of $s \in [1, n]$, the cardinality of A_3 attains its maximum only for some $s \in [s' - 2k, s' + 2]$. This, together with (8), completes the proof of the lemma. \square

Now we can prove the first conjecture of Chung and Goldwasser.

Theorem 1

$$\lim_{n \rightarrow \infty} \frac{f(n, k)}{n} = \frac{k(k-2)}{k^2-2} + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)}.$$

Proof. Let A be a maximum k -sum-free set in $[1, n]$, with n sufficiently large. From Lemma 4 we have $\frac{s_A}{n} = \frac{S^*}{n} + o(1)$, where $S^* = \frac{8n}{k^5-2k^3-4k}$. Thus we can estimate

$$\begin{aligned} \frac{f(n, k)}{n} &= \frac{|A_3|}{n} = \frac{r_1 - l_1 + r_2 - l_2 + r_3 - S^* + 1}{n} + o(1) \\ &= \frac{1}{n} \left(n - \frac{2n}{k} + \frac{2n + kS^*}{k^2} - \frac{4n + 2kS^*}{k^3} + \frac{4n + 2kS^* + k^3S^*}{k^4} - S^* \right) + o(1) \\ &= \frac{k^4 - 2k^3 + 2k^2 - 4k + 4}{k^4} + \frac{S^*}{nk^3}(2k^2 - 2k + 2 - k^3) + o(1) \\ &= \frac{k^4 - 2k^3 + 2k^2 - 4k + 4}{k^4} + \frac{8(2k^2 - 2k + 2 - k^3)}{(k^5 - 2k^3 - 4k)k^3} + o(1) \\ &= \frac{k^5 - 2k^4 - 4k + 8}{(k^4 - 2k^2 - 4)k} + o(1) \\ &= \frac{k(k-2)}{k^2-2} + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)} + o(1), \end{aligned}$$

and the claim follows by taking the limit. \square

We can now show the main result.

Theorem 2 Let $k \in \mathbb{N}_{\geq 4}$ and $n > n_1(k)$. Let S and s' be as in Lemma 4. Let $A \subseteq \{1, \dots, n\}$ be a k -sum-free set of maximum cardinality, with smallest element $s = s_A$. Then $s \in [S, S + 3]$ and $A = \mathcal{I}_3 \cup \mathcal{I}_2 \cup \mathcal{I}_1$, where

$$\begin{aligned} \mathcal{I}_3 &\in \begin{cases} \{[s, r_3], [s, r_3 + 1]\}, & \text{if } s \geq s' \\ \{[s, r_3], [s, r_3] \setminus \{r_3 - 1\}\}, & \text{if } s < s', \end{cases} \\ \mathcal{I}_2 &\in \begin{cases} \{[l_2 + 2, r_2], [l_2 + 2, r_2 + 1]\}, & \text{if } r_3 + 1 \in A \\ \{(l_2, r_2), (l_2, r_2 + 1), [l_2, r_2], [l_2, r_2] \setminus \{r_2 - 1\}\}, & \text{if } r_3 + 1 \notin A, \end{cases} \\ \mathcal{I}_1 &\in \begin{cases} \{[l_1 + 2, n]\}, & \text{if } r_2 + 1 \in A \\ \{[l_1, n], (l_1, n], [l_1, n] \setminus \{n - 1\}\}, & \text{if } r_2 + 1 \notin A, \end{cases} \end{aligned}$$

If k is even, then $\mathcal{I}_i \neq [l_i, r_i] \setminus \{r_i - 1\}$ for $1 \leq i \leq 3$.

Remark. Note that Theorem 2 does not precisely determine the k -sum-free subsets of $\{1, \dots, n\}$ of maximum size, for every $n > n_1(k)$. With n and k fixed, one first needs to determine for which value(s) of $s \in [S, S + 3]$ the quantity $|A_3(s)|$, as given by (9), is maximized. The result will depend on n and k . Even then, for a fixed s , not all the possibilities for $\mathcal{I}_3 \cup \mathcal{I}_2 \cup \mathcal{I}_1$ need be k -sum-free. See Section 4 below for further discussion.

Proof. We have already seen that $|A_3| = |A|$. Our first aim is to show by comparing A_3 with A_2 that almost the whole interval $(l_3, r_3]$ must be in A . Having achieved this, we infer by Lemma 1 that $(r_3, l_2] \cap A$ is nearly empty. Comparing A_2 with A_1 will then reveal that most of $(l_2, r_2]$ is contained in A . Again Lemma 1 will help us to see that A cannot share many elements with $(r_2, l_1]$ and a final comparison of A_1 with A will conclude the proof.

(I) The first aim is easily reached if $s := s_A \geq l_3 + 1$. Simply note that

$$A_2 = (A \cap [s, r_3]) \cup (l_2, r_2] \cup (l_1, r_1] \subseteq [s, r_3] \cup (l_2, r_2] \cup (l_1, r_1] = A_3.$$

The maximality of $|A_2|$ gives $A_2 = A_3$ and hence $[s, r_3] \subseteq A$. Observe that $s > l_3$ together with Lemma 4 and (8) give $S \leq s \leq S + 3$.

Assume now that $s \leq l_3$. We want to show that in this case $s = l_3$. Suppose $s < l_3$ and let $B = [S - 2k, l_3] \cap A$. Define

$$C := I_{s_B}^1 \cup \bigcup_{b \in B \setminus \{s_B\}} I_b^0.$$

Clearly $C \subseteq (l_3, r_3]$ for all $n \gg 0$. Then since C is the union of disjoint intervals, Lemma 1 gives that $|C \setminus A| > |B|$. Hence we get the contradiction $|A_3| = |(A_2 \setminus B) \cup (l_3, r_3]| \geq |(A_2 \setminus B) \cup (C \setminus A)| > |A_2| - |B| + |B| = |A_2|$. Therefore we are left with $s = l_3$, and this implies

$$|A_2| = |A_3| \iff |A \cap [s, r_3]| = |(l_3, r_3] \cap [s, r_3]| = |(s, r_3]|. \quad (10)$$

If $r_3 \notin A$ we can infer from (10) that

$$A \cap [s, r_3] = [s, r_3 - 1] = [l_3, r_3 - 1].$$

If $r_3 \in A$, Lemma 1 gives $kl_3 - r_3 \notin A$, so $-k + 1 \leq kl_3 - 2r_3 \leq -1$. If $kl_3 - 2r_3 \leq -2$ we get $I_{l_3}^1 \subseteq (l_3, r_3]$ and $|I_{l_3}^1 \setminus A| \geq 2$, which is impossible since this would imply $|A_3| > |A_2|$. Hence $kl_3 - 2r_3 = -1$ and k is odd. Using (10) one obtains

$$A \cap [s, r_3] = [l_3, r_3] \setminus \{r_3 - 1\}.$$

Suppose now that $s = l_3$ and $r_3 + 1 \in A$. Then $kl_3 - (r_3 + 1) \notin A$ and

$$r_3 - k \leq kl_3 - (r_3 + 1) \leq r_3 - 1.$$

This contradicts that $[s, r_3 - 2] \subseteq A$ unless $kl_3 - (r_3 + 1) = r_3 - 1$, but then $r_3 \notin A$ and $|A \cap [s, r_3]| = |A \cap [s, r_3 - 2]|$ which contradicts (10). Hence $r_3 + 1 \notin A$ if $s = l_3$.

Finally note that, if $s = l_3$ and $kl_3 \geq 2r_3 - 1$, the latter being a requirement for either of the two possibilities for \mathcal{I}_3 to be k -sum-free, then another computation similar to the one in the Appendix yields that $s \geq S$. Again, using Lemma 4 we obtain

$$S \leq s \leq S + 3, \tag{11}$$

as claimed in the statement of the theorem. This completes the first part of our proof.

(II) For the second part note that we have just shown

$$s \geq l_3. \tag{12}$$

Plugging (11) into the definition of l_3 yields (after a further tedious computation similar to that in the Appendix)

$$S - 1 \leq l_3 \leq S + 1, \tag{13}$$

which implies in view of (12) and (11)

$$l_3 \leq s \leq l_3 + 4. \tag{14}$$

Moreover we have observed that $[s, r_3 - 2] \subseteq A$. Let $\xi_1, \dots, \xi_5 \in \{0, \dots, k - 1\}$ be constants such that

$$kl_1 = 2r_1 - \xi_1 \tag{15}$$

$$kr_2 = l_1 + s - \xi_2 \tag{16}$$

$$kl_2 = 2r_2 - \xi_3 \tag{17}$$

$$kr_3 = l_2 + s - \xi_4 \tag{18}$$

$$kl_3 = 2r_3 - \xi_5. \tag{19}$$

We suppose that n is sufficiently large, so we can be sure that

$$[ks - (r_3 - 2), k(r_3 - 2) - s] \cap A = \emptyset.$$

By (14) we can infer that

$$\begin{aligned} \emptyset &= [k(l_3 + 4) - (r_3 - 2), k(r_3 - 2) - s] \cap A \\ &= [r_3 - \xi_5 + 4k + 2, l_2 - \xi_4 - 2k] \cap A. \end{aligned}$$

Let $J = [r_3 + 2, r_3 - \xi_5 + 4k + 1] \cap A$ and $K = \bigcup_{x \in J} \{kx - (s + 2), kx - (s + 1), kx - s\}$. Then $K \cap A = \emptyset$, $|K| = 3|J|$ and by (18) and (19) we have

$$K \subseteq [l_2 - \xi_4 + 2k - 2, l_2 - \xi_4 - k\xi_5 + 4k^2 + k] \subseteq (l_2 + k - 2, l_2 + 4k^2 + k] \subseteq (l_2 + 2, r_2],$$

if $n \gg 0$. Let $B = [l_2 - \xi_4 - 2k + 1, l_2] \cap A$. If $B \cup J \subseteq \{l_2\}$ then $A \cap [r_3 + 2, l_2 - 1] = \emptyset$. Otherwise, with C as in part (I) if $|B| > 1$ we can verify that $C \subseteq [r_2 - \frac{3k^2 - k + 2}{2}, r_2] \subseteq (l_2 + 1, r_2]$, for $n \gg 0$, and $|C \setminus A| > |B|$. Put $C := \emptyset$ if $|B| \leq 1$. For large n , K and C are disjoint. Hence $|B \cup J| < |(C \setminus A) \cup K|$ and we get

$$|A_2| = |[A_1 \setminus (J \cup B \cup \{r_3 + 1\})] \cup (l_2, r_2]| > |A_1 \setminus \{r_3 + 1\}|.$$

Thus if $r_3 + 1 \notin A$ we get $|A_2| > |A_1|$ so suppose $r_3 + 1 \in A$. Then neither l_2 nor $l_2 + 1$ can be in A_1 . Otherwise, since $(s - \xi_4 + k), s - \xi_4 + k - 1 \in [s, s + k] \subseteq [s, r_3 - 2] \subseteq A$ we get

$$k(r_3 + 1) = l_2 + (s - \xi_4 + k) = (l_2 + 1) + (s - \xi_4 + k - 1),$$

which is impossible. But $l_2 + 1 \in A_2$, so also in this case it follows that $|A_2| > |A_1|$, since $l_2 + 1 \notin K \cup C$ for large n . Again we conclude that $A \cap [r_3 + 2, l_2 - 1] = \emptyset$. Consequently,

$$|A_2| = |A_1| \Leftrightarrow |A \cap ([l_2, r_2] \cup \{r_3 + 1\})| = |[l_2, r_2]|,$$

which gives $A \cap [l_2, r_2] = [l_2 + 2, r_2]$ if $r_3 + 1 \in A$. If $r_3 + 1 \notin A$ and either $l_2 \notin A$ or $r_2 \notin A$, we get $A \cap [l_2, r_2] = (l_2, r_2]$ or $A \cap [l_2, r_2] = [l_2, r_2)$, respectively. In case $r_3 + 1 \notin A$ and both $l_2, r_2 \in A$, we see that $kl_2 - r_2 = r_2 - \xi_3 \notin A$. If $\xi_3 \geq 2$ then $I_{l_2}^1 \subseteq (l_2, r_2]$ and l_2 could be profitably replaced. Hence $\xi_3 = 1$, $A \cap [l_2, r_2] = [l_2, r_2] \setminus \{r_2 - 1\}$ and k is odd.

(III) For the final interval $(l_1, r_1]$ we use Lemma 1 to conclude from

$$[s, r_3 - 2] \subseteq A \text{ and } [l_2 + 2, r_2 - 2] \subseteq A$$

in view of (16) and (17) that, for $n \gg 0$,

$$\begin{aligned} \emptyset &= A \cap [k(l_2 + 2) - (r_2 - 2), k(r_2 - 2) - (l_2 + 2)] \\ &= A \cap [r_2 - \xi_3 + 2k + 2, l_1 + s - \xi_2 - 2k - l_2 - 2], \text{ and} \\ \emptyset &= A \cap [k(l_2 + 2) - (r_3 - 2), k(r_2 - 2) - s] \\ &= A \cap [2r_2 - \xi_3 + 2k - r_3 + 2, l_1 - \xi_2 - 2k] \end{aligned}$$

Let $J = [r_2 + 2, r_2 - \xi_3 + 2k + 1] \cap A$ and $K = \cup_{x \in J} \{kx - s, kx - (s + 1), kx - (s + 2)\}$. From (14) we have

$$K \subseteq [l_1 - \xi_2 + 2k - 2, l_1 - \xi_2 - k\xi_3 + 2k^2 + k] \subseteq (l_1 + k - 2, r_1], \text{ if } n \gg 0.$$

Let $B = [l_1 - \xi_2 - 2k + 1, l_1] \cap A$. If $s_B < l_1$ with C as in (I) we can verify that, for sufficiently large n ,

$$C \subseteq \left[\frac{2r_1 - \xi_1 - k\xi_2 - 2k^2 + k - 5}{2}, r_1 \right] \subseteq (l_1, r_1],$$

$|C \setminus A| > |B|$ and $\max K < s_C$. By analogy with part (II) we get $A \cap [r_2 + 2, l_1 - 1] = \emptyset$ and the rest of the claim follows as before. \square

4 Estimates and Periodicity

We first want to estimate values of $n_i(k)$, $i = 0, 1$, for which Lemmas 3 and 4, and Theorem 2 respectively are valid. The estimates we shall arrive at can probably be improved upon. The example of a k -sum-free set A in [3], referred to in the proof of Lemma 3, satisfies

$$|A| > \frac{k(k-2)}{k^2-2}n + \frac{8(k-2)}{k(k^2-2)(k^4-2k^2-4)}n - 3.$$

Hence the proof of Lemma 3 goes through provided n is sufficiently large so that

$$c_k n - t_0 \geq 3, \tag{20}$$

where $t_0 = t_0(n, k)$ is the largest possible value for t in Definition 1. Now from Definition 1 we easily deduce that, if $i < t$, then $r_{i+1} \leq \left(\frac{4}{k^2}\right)r_i$, and hence that $r_t \leq \left(\frac{4}{k^2}\right)^{t-1}n$. Since $r_t \geq 1$ a priori, we can thus estimate

$$t_0 \leq \frac{1}{2} \log_{k/2} n + 1. \tag{21}$$

Since, by (7), $c_k = \mathcal{O}\left(\frac{1}{k^6}\right)$, we thus deduce from (18) and (19) that one can take $n_0(k) = \mathcal{O}(k^6)$. It is then an easy and tedious exercise to go through the proof of Theorem 2 and check that one can also take $n_1(k) = \mathcal{O}(k^6)$.

Next, we explain what we mean by the word ‘periodicity’ in the title of this section. If $k \geq 4$ is even then, for $n > 0$, we have $s' = S + 1 = \lfloor \frac{8n}{k^5 - 2k^3 - 4k} \rfloor + 1$. Hence for a fixed k , if we regard s' as a function of n , then $s'(n) + 1 = s'(n + p_k)$, where $p_k := \frac{k^5 - 2k^3 - 4k}{8}$. For odd k , we define $p_k := k^5 - 2k^3 - 4k$ and in this case, a little more care is required to check that $s'(n) + 8 = s'(n + p_k)$.

Now for any k and n , let $\mathcal{F}(k, n)$ denote the family of maximal k -sum-free subsets of $\{1, \dots, n\}$. Then for n sufficiently large, as estimated above, and k even (resp. k odd), the map $s \mapsto s + 1$ (resp. $s \mapsto s + 8$) clearly induces a 1-1 correspondence between the sets in $\mathcal{F}(k, n)$ and $\mathcal{F}(k, n + p_k)$. This is what we mean by ‘periodicity’. This observation clearly reduces, for any fixed k , the full classification of all k -sum-free subsets of $\{1, \dots, n\}$, for all n , to a finite computation.

As an example, we now look at $k = 4$. By (7) we compute $c_4 = \frac{47}{48290}$. Then Lemma 3 is valid at least for all n satisfying

$$c_4 n - \frac{1}{2} \log_2 n - 1 \geq 3,$$

which reduces to $n \geq 11008$. One can then check that the proof of Theorem 2 also goes through for all such n . We have $p_4 = 110$. We now present the full classification of all 4-sum-free subsets of $\{1, \dots, n\}$, valid (at least) for all $n \geq 11008$. This was obtained with the help of a computer.

For each $s, n \in \mathbf{N}$ we define the sets $J_x(s)$, $1 \leq x \leq 13$, as follows (the l_i and r_i are functions of s and n as in Definition 1) :

$$\begin{aligned} J_1 &= [S, r_3 - 1] \cup [l_2, r_2 - 1] \cup [l_1, n - 1], \\ J_2 &= [S, r_3 - 1] \cup [l_2, r_2 - 1] \cup [l_1 + 1, n], \\ J_3 &= [S, r_3 - 1] \cup [l_2 + 1, r_2] \cup [l_1, n - 1], \\ J_4 &= [S, r_3 - 1] \cup [l_2 + 1, r_2] \cup [l_1 + 1, n], \\ J_5 &= [S, r_3 - 1] \cup [l_2 + 1, r_2 + 1] \cup [l_1 + 2, n], \\ J_6(s) &= [s, r_3] \cup [l_2, r_2 - 1] \cup [l_1, n - 1], \\ J_7(s) &= [s, r_3] \cup [l_2, r_2 - 1] \cup [l_1 + 1, n], \\ J_8(s) &= [s, r_3] \cup [l_2 + 1, r_2] \cup [l_1, n - 1], \\ J_9(s) &= [s, r_3] \cup [l_2 + 1, r_2] \cup [l_1 + 1, n], \\ J_{10}(s) &= [s, r_3] \cup [l_2 + 1, r_2 + 1] \cup [l_1 + 2, n], \\ J_{11}(s) &= [s, r_3 + 1] \cup [l_2 + 2, r_2] \cup [l_1, n - 1], \\ J_{12}(s) &= [s, r_3 + 1] \cup [l_2 + 2, r_2] \cup [l_1 + 1, n], \\ J_{13}(s) &= [s, r_3 + 1] \cup [l_2 + 2, r_2 + 1] \cup [l_1 + 2, n]. \end{aligned}$$

Note that, by Theorem 2, for a given $n \geq 11008$, every maximal 4-sum-free subset of $\{1, \dots, n\}$ is one of the sets $J_x(s)$, for some $s \in [S, S + 3] = [s' - 1, s' + 2]$. By the remarks above, for each $i \in \{0, \dots, 109\}$, there are natural 1-1 correspondences between the sets in the families $\mathcal{F}(4, n)$ for all $n \equiv i \pmod{110}$. By slight abuse of notation, we denote any such family simply by \mathcal{F}_i . Our computer program yielded the following result :

If $|\mathcal{F}_i| = 1$, then $i = 6, 7, 22, 23, 46, 47, 49, 51, 54, 55, 57, 59, 61, 70, 71, 73, 75, 77, 86, 87, 89$

or 91 and

$$\mathcal{F}_i = \{J_9(s')\},$$

or $i = 36, 37, 100$ or 101 and

$$\mathcal{F}_i = \{J_9(s' + 1)\}.$$

If $|\mathcal{F}_i| = 2$, then \mathcal{F}_i is

$$\{J_9(s'), J_9(s' + 1)\} \quad \text{if } i = 93, 103, 105, 107,$$

$$\{J_4, J_9(s')\} \quad \text{if } i = 9, 11, 13, 25, 27,$$

$$\{J_8(s'), J_9(s')\} \quad \text{if } i = 48, 50, 56, 58, 60, 72, 74, 76, 88, 90$$

$$\{J_7(s'), J_9(s')\} \quad \text{if } i = 63, 65, 67, 79, 81.$$

If $|\mathcal{F}_i| = 3$:

$$\mathcal{F}_8 = \mathcal{F}_{24} = \{J_4, J_8(s'), J_9(s')\},$$

$$\mathcal{F}_{15} = \{J_4, J_7(s'), J_9(s')\},$$

$$\mathcal{F}_{29} = \{J_4, J_9(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{39} = \{J_9(s'), J_{12}(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{62} = \mathcal{F}_{78} = \{J_6(s'), J_7(s'), J_9(s')\},$$

$$\mathcal{F}_{53} = \{J_9(s'), J_{10}(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{83} = \{J_7(s'), J_9(s'), J_9(s' + 2)\},$$

$$\mathcal{F}_{92} = \{J_8(s'), J_9(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{95} = \mathcal{F}_{97} = \{J_7(s'), J_9(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{102} = \{J_9(s'), J_8(s' + 1), J_9(s' + 1)\},$$

$$\mathcal{F}_{109} = \{J_9(s'), J_7(s' + 1), J_9(s' + 1)\}.$$

If $|\mathcal{F}_i| = 4$:

$$\mathcal{F}_1 = \mathcal{F}_3 = \mathcal{F}_{17} = \{J_2, J_4, J_7(s'), J_9(s')\},$$

$$\mathcal{F}_{10} = \mathcal{F}_{12} = \mathcal{F}_{26} = \{J_3, J_4, J_8(s'), J_9(s')\},$$

$$\mathcal{F}_{38} = \{J_9(s'), J_{12}(s'), J_8(s' + 1), J_9(s' + 1)\},$$

$$\mathcal{F}_{41} = \mathcal{F}_{43} = \{J_4, J_9(s'), J_{12}(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{52} = \{J_8(s'), J_9(s'), J_{10}(s'), J_9(s' + 1)\},$$

$$\mathcal{F}_{64} = \mathcal{F}_{66} = \mathcal{F}_{80} = \{J_6(s'), J_7(s'), J_8(s'), J_9(s')\},$$

$$\mathcal{F}_{104} = \mathcal{F}_{106} = \{J_8(s'), J_9(s'), J_8(s' + 1), J_9(s' + 1)\},$$

$$\mathcal{F}_{69} = \{J_7(s'), J_9(s'), J_{10}(s'), J_9(s' + 1)\}.$$

If $|\mathcal{F}_i| = 5$:

$$\begin{aligned}
\mathcal{F}_{14} &= \{J_3, J_4, J_6(s'), J_7(s'), J_9(s')\}, \\
\mathcal{F}_{19} &= \{J_2, J_4, J_7(s'), J_9(s'), J_9(s'+2)\}, \\
\mathcal{F}_{28} &= \{J_3, J_4, J_8(s'), J_9(s'), J_9(s'+1)\}, \\
\mathcal{F}_{31} &= \{J_4, J_7(s'), J_9(s'), J_{12}(s'), J_9(s'+1)\}, \\
\mathcal{F}_{82} &= \{J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_9(s'+2)\}, \\
\mathcal{F}_{94} &= \{J_6(s'), J_7(s'), J_9(s'), J_8(s'+1), J_9(s'+1)\}, \\
\mathcal{F}_{99} &= \{J_7(s'), J_9(s'), J_9(s'+1), J_{10}(s'+1), J_9(s'+2)\}, \\
\mathcal{F}_{108} &= \{J_8(s'), J_9(s'), J_6(s'+1), J_7(s'+1), J_9(s'+1)\}.
\end{aligned}$$

If $|\mathcal{F}_i| = 6$:

$$\begin{aligned}
\mathcal{F}_5 &= \{J_2, J_4, J_7(s'), J_9(s'), J_{10}(s'), J_9(s'+1)\}, \\
\mathcal{F}_{33} &= \{J_2, J_4, J_7(s'), J_9(s'), J_{12}(s'), J_9(s'+1)\}, \\
\mathcal{F}_{45} &= \{J_4, J_9(s'), J_{12}(s'), J_{13}(s'), J_7(s'+1), J_9(s'+1)\}, \\
\mathcal{F}_{68} &= \{J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_{10}(s'), J_9(s'+1)\}, \\
\mathcal{F}_{85} &= \{J_7(s'), J_9(s'), J_{10}(s'), J_9(s'+1), J_{12}(s'+1), J_9(s'+2)\}, \\
\mathcal{F}_{96} &= \{J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_8(s'+1), J_9(s'+1)\}.
\end{aligned}$$

If $|\mathcal{F}_i| = 7$:

$$\begin{aligned}
\mathcal{F}_0 = \mathcal{F}_{16} &= \{J_1, J_2, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s')\}, \\
\mathcal{F}_{40} &= \{J_4, J_8(s'), J_9(s'), J_{11}(s'), J_{12}(s'), J_8(s'+1), J_9(s'+1)\}.
\end{aligned}$$

If $|\mathcal{F}_i| = 8$:

$$\begin{aligned}
\mathcal{F}_2 &= \{J_1, J_2, J_3, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s')\}, \\
\mathcal{F}_{21} &= \{J_2, J_4, J_7(s'), J_9(s'), J_{10}(s'), J_9(s'+1), J_{12}(s'+1), J_9(s'+2)\}, \\
\mathcal{F}_{30} &= \{J_3, J_4, J_6(s'), J_7(s'), J_9(s'), J_{12}(s'), J_8(s'+1), J_9(s'+1)\}, \\
\mathcal{F}_{35} &= \{J_2, J_4, J_7(s'), J_9(s'), J_{12}(s'), J_9(s'+1), J_{10}(s'+1), J_9(s'+2)\}, \\
\mathcal{F}_{42} &= \{J_3, J_4, J_8(s'), J_9(s'), J_{11}(s'), J_{12}(s'), J_8(s'+1), J_9(s'+1)\}, \\
\mathcal{F}_{98} &= \{J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_8(s'+1), J_9(s'+1), J_{10}(s'+1), J_9(s'+2)\}.
\end{aligned}$$

If $|\mathcal{F}_i| = 9$:

$$\begin{aligned}
\mathcal{F}_{18} &= \{J_1, J_2, J_3, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_9(s'+2)\}, \\
\mathcal{F}_{84} &= \{J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_{10}(s'), J_9(s'+1), J_{12}(s'+1), J_8(s'+2), J_9(s'+2)\}.
\end{aligned}$$

If $|\mathcal{F}_i| = 10$:

$$\begin{aligned}
\mathcal{F}_4 &= \{J_1, J_2, J_3, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_{10}(s'), J_9(s'+1)\}, \\
\mathcal{F}_{44} &= \{J_3, J_4, J_8(s'), J_9(s'), J_{11}(s'), J_{12}(s'), J_{13}(s'), J_6(s'+1), J_7(s'+1), J_9(s'+1)\}.
\end{aligned}$$

If $|\mathcal{F}_i| = 11, 13$ or 14 , we get precisely one family for each size:

$$\begin{aligned} \mathcal{F}_{32} &= \{J_1, J_2, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_{11}(s'), J_{12}(s'), J_8(s'+1), J_9(s'+1)\}, \\ \mathcal{F}_{20} &= \{J_1, J_2, J_3, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_{10}(s'), \\ &\quad J_9(s'+1), J_{12}(s'+1), J_8(s'+2), J_9(s'+2)\}, \\ \mathcal{F}_{34} &= \{J_1, J_2, J_3, J_4, J_6(s'), J_7(s'), J_8(s'), J_9(s'), J_{11}(s'), J_{12}(s'), \\ &\quad J_8(s'+1), J_9(s'+1), J_{10}(s'+1), J_9(s'+2)\}. \end{aligned}$$

Note, in particular, that $|\mathcal{F}(4, n)| \leq 14$ for all sufficiently large n . Computer simulations suggest the same may be true for any even k , with a similar result for odd k , but we leave the investigation of this possibility to a subsequent paper.

Appendix

As a prototype for a type of calculation which appears in several places in the paper, we now show, in the notation of Lemma 4, that $s' = S + 1$ when k is even.

We must investigate the condition $l_3(s) < s$. By definition of l_3 this is just

$$\begin{aligned} \left\lfloor \frac{2r_3}{k} \right\rfloor < s &\Leftrightarrow \frac{2r_3}{k} < s \Leftrightarrow r_3 < \frac{ks}{2} \Leftrightarrow \left\lfloor \frac{l_2 + s}{k} \right\rfloor < \frac{ks}{2} \Leftrightarrow \frac{l_2 + s}{k} < \frac{ks}{2} \\ &\Leftrightarrow l_2 < \left(\frac{k^2}{2} - 1\right)s \Leftrightarrow \frac{2r_2}{k} < \left(\frac{k^2}{2} - 1\right)s \Leftrightarrow r_2 < \left(\frac{k^3}{4} - \frac{k}{2}\right)s \\ &\Leftrightarrow \frac{l_1 + s}{k} < \left(\frac{k^3}{4} - \frac{k}{2}\right)s \Leftrightarrow l_1 < \left(\frac{k^4}{4} - \frac{k^2}{2} - 1\right)s \\ &\Leftrightarrow \frac{2n}{k} < \left(\frac{k^4}{4} - \frac{k^2}{2} - 1\right)s \Leftrightarrow n < \left(\frac{k^5}{8} - \frac{k^3}{4} - \frac{k}{2}\right)s \Leftrightarrow s > \frac{8n}{k^5 - 2k^3 - 4k} \\ &\Leftrightarrow s > S. \end{aligned}$$

Thus $s' = S + 1$, as required.

Acknowledgments

We would like to thank the anonymous referee whose detailed comments greatly helped us improving our paper.

References

- [1] J. M. Deshoulliers, G. Freiman, V. Sós, and M. Temkin, On the structure of sum-free sets II, *Asterisque* **258** (1999), 149-161.

- [2] F. R. K. Chung and J. L. Goldwasser, Integer sets containing no solutions to $x+y = 3k$, in “The Mathematics of Paul Erdős”, R. L. Graham and J. Nešetřil eds., Springer Verlag, Heidelberg, (1996).
- [3] F. R. K. Chung and J. L. Goldwasser, Maximum subsets of $(0, 1]$ with no solutions to $x + y = kz$, *Electron. J. Combin.* **3** (1996), R1.
- [4] K. Roth, On certain sets of integers, *J. London Math. Society* **28** (1953), 104-109.