



The TIR1 protein of *Arabidopsis* functions in auxin response and is related to human SKP2 and yeast Grr1p

Max Ruegger, Elizabeth Dewey, William M. Gray, et al.

Genes Dev. 1998 12: 198-207

Access the most recent version at doi:[10.1101/gad.12.2.198](https://doi.org/10.1101/gad.12.2.198)

References

This article cites 40 articles, 15 of which can be accessed free at:
<http://genesdev.cshlp.org/content/12/2/198.full.html#ref-list-1>

Article cited in:

<http://genesdev.cshlp.org/content/12/2/198.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#)

To subscribe to *Genes & Development* go to:
<http://genesdev.cshlp.org/subscriptions>

The TIR1 protein of *Arabidopsis* functions in auxin response and is related to human SKP2 and yeast Grr1p

Max Ruegger,¹ Elizabeth Dewey, William M. Gray, Lawrence Hobbie,² Jocelyn Turner, and Mark Estelle³

Department of Biology, Indiana University, Bloomington, Indiana 47405 USA

Genetic analysis in *Arabidopsis* has led to the identification of several genes that are required for auxin response. One of these genes, *AXR1*, encodes a protein related to yeast Aos1p, a protein that functions to activate the ubiquitin-related protein Smt3p. Here we report the identification of a new gene called *TRANSPORT INHIBITOR RESPONSE 1 (TIR1)*. The *tir1* mutants are deficient in a variety of auxin-regulated growth processes including hypocotyl elongation and lateral root formation. These results indicate that *TIR1* is also required for normal response to auxin. Further, mutations in *TIR1* display a synergistic interaction with mutations in *AXR1*, suggesting that the two genes function in overlapping pathways. The *TIR1* protein contains a series of leucine-rich repeats and a recently identified motif called an F box. Sequence comparisons indicate that *TIR1* is related to the yeast protein Grr1p and the human protein SKP2. Because Grr1p and other F-box proteins have been implicated in ubiquitin-mediated processes, we speculate that auxin response depends on the modification of a key regulatory protein(s) by ubiquitin or a ubiquitin-related protein.

[Key Words: F-box protein; *Arabidopsis*; auxin response; TIR1; human SKP2; yeast Grr1p]

Received September 10, 1997; revised version accepted November 14, 1997.

Physiological studies have implicated the plant hormone indole-3-acetic acid (IAA or auxin) in the regulation of diverse developmental processes including stem elongation, apical dominance, photo- and gravitropism, and lateral root initiation (Klee and Estelle 1991). At the cellular level, auxin acts to regulate these processes through changes in cell division and cell expansion (Evans 1984). Although the mechanism or mechanisms of auxin action are largely unknown, the existing evidence suggests that the hormone acts both at the plasma membrane and within the cell (Vesper and Kuss 1990; Venis and Napier 1995). Auxin treatment results in rapid hyperpolarization of the plasma membrane and induction of specific gene expression (Barbier-Brygoo 1995; Abel and Theologis 1996).

In *Arabidopsis*, genetic studies have resulted in the identification of a number of genes that are required for normal auxin response (Hobbie and Estelle 1994, Leyser 1997). For three of the genes, *AXR1*, *AXR4*, and *AUX1*, recessive mutations result in reduced auxin response as well as an array of auxin-related growth defects (Hobbie and Estelle 1994). By genetic criteria, *AXR1* and *AUX1*

appear to act in distinct pathways. *AUX1* is a membrane protein with similarity to amino acid permeases from plants and fungi (Bennett et al. 1996). Because IAA is an indolic compound structurally related to tryptophan, *AUX1* may function in cellular auxin uptake and not response per se. In contrast, the *axr1* and *axr4* mutations display a synergistic interaction, suggesting that the wild-type genes function in the same or overlapping pathways (Hobbie and Estelle 1995). Genetic experiments indicate that another gene, called *SAR1*, can also be placed in this group. The *sar1* mutants were isolated as suppressors of the *axr1* mutations and have a phenotype that is distinct from that of either *axr1* or wild type (Cernac et al. 1997). In double-mutant combinations, *sar1* is epistatic to *axr1*. This combination of suppression and epistasis suggests that *SAR1* functions after *AXR1*. Thus, *AXR1*, *AXR4*, and *SAR1* form a genetically related group of genes that appear to function together to mediate auxin response.

The *AXR1* gene encodes a protein related to the ubiquitin-activating enzyme (E1), the first enzyme in the ubiquitin conjugation pathway (Leyser et al. 1993). The function of E1 is to form a thiol-ester linkage between the carboxyl terminus of ubiquitin and an internal cysteine. The ubiquitin moiety is subsequently transferred to a ubiquitin-conjugating enzyme (E2). Finally, ubiquitin is attached to the target protein by an isopeptide linkage between a lysine and the carboxyl terminus of ubiq-

Present addresses: ¹Department of Biochemistry, Purdue University, West Lafayette, Indiana 47907 USA; ²Department of Biology, Adelphi University, Garden City, New York 11530 USA.

³Corresponding author.

E-MAIL mestelle@bio.indiana.edu; FAX (812) 855-6705.

uitin. This last step is catalyzed by a protein or protein complex called a ubiquitin-protein ligase (E3). Proteins related to AXR1 have now been identified in humans (Chow et al. 1996) and fungi (Johnson et al. 1997; Shayeghi et al. 1997). These proteins all share extensive sequence similarity with the amino terminus of E1, but lack the active-site cysteine required for E1 activity. Recent studies in *Saccharomyces cerevisiae* indicate that at least one member of this family, called Aos1p, functions as a dimer with a second protein called Uba2p (Johnson et al. 1997). Uba2p is similar to the carboxyl terminus of E1 and includes the region containing the active-site cysteine (Dohmen et al. 1995). The Aos1p/Uba2p dimer forms a thiol-ester linkage with a ubiquitin-related protein called Smt3p. Thus, it is possible that AXR1 and other members of the AXR1 protein family each interact with a second protein to form a functional enzyme. The substrate for each heterodimer-type E1, ubiquitin and/or a ubiquitin-related protein, remains to be determined.

To identify additional genes involved in auxin physiology, we have screened for *Arabidopsis* mutants that are resistant to the growth-inhibiting properties of the auxin-transport inhibitors naphthylphthalamic acid (NPA) and 2-carboxyphenyl-3-phenylpropane-1,2-dione (CPD) (Ruegger et al. 1997). This screen resulted in the recovery of mutants that are altered in auxin transport as well as auxin response. In this report, we show that the *transport inhibitor response 1* (*tir1*) mutants are deficient in auxin response. Molecular characterization indicates that the TIR1 protein is, like AXR1, a member of a family of proteins implicated in ubiquitin-mediated processes.

Results

Genetic characterization of the *tir1* mutants

Previously, we reported the isolation of 16 *tir* mutants by screening for resistance to auxin-transport inhibitors (Ruegger et al. 1997). Genetic analysis indicated that five of these mutations were independent alleles at a locus we called *TIR1*. To determine the genetic basis for resistance, we examined segregation of CPD resistance in F_1 and F_2 populations. In our initial studies, seedlings that were heterozygous for *tir1* alleles were often significantly less affected by CPD than wild-type seedlings, suggesting that the mutations are at least partially dominant. When F_2 populations were analyzed, a graded distribution of phenotypes was observed. An example of a population scored on 5 μ M CPD is shown in Table 1. The genotype of individual F_2 plants was determined by examining the response of F_3 progeny to CPD. This analysis revealed that homozygous *tir1* plants are clearly resistant to the compound and heterozygous plants tended to be somewhat resistant compared with homozygous wild-type plants. To confirm that *tir1* mutations are semidominant, we directly compared populations of wild-type, *tir1/+* and *tir1/tir1* seedlings on medium containing CPD. The results are shown in Table 2. For each

Table 1. Segregation of CPD resistance in a *tir1-2* \times wild-type F_2 population

F_2 root length (mm)	F_2 genotype ¹		
	<i>tir1/tir1</i> ²	<i>tir1/+</i> ²	<i>+/+</i> ²
17	2		
16	5		
15	1		
14			
13			
12		1	
11			
10		6	
9		6	
8		7	6
7		3	4
6		2	4
5			
4		1	
3		1	
Totals	8	27	14

¹Genotype determined by analyzing CPD resistance in F_3 plants.

²Number of seedlings.

of the three alleles tested, heterozygous plants were weakly resistant. The five mutants differ only slightly with respect to CPD response. The *tir1-2* allele is slightly weaker, whereas *tir1-1* and *tir1-3* are slightly stronger than *tir1-6* and *tir1-7*. All of the phenotypic analysis described below was performed with the *tir1-1* mutant.

The *tir1* mutants are affected in auxin response

Treatment of *Arabidopsis* seedlings with auxin-transport inhibitors such as CPD or NPA inhibits root elongation and promotes swelling of the root tip (Ruegger et al. 1997). Both of these effects are thought to be caused by accumulation of auxin in this region as a result of the inhibition of auxin transport away from the root tip (Muday and Haworth 1994; Muday et al. 1995; Ruegger et al. 1997). Because auxin-transport inhibitors affect root growth by increasing auxin levels in the root tip, resistance could result from a change in auxin transport or a

Table 2. CPD resistance in *tir* homozygous and heterozygous plants

Line	Root length (mm)	\pm S.E.	No.
<i>+/+</i>	7.8	0.1	32
<i>tir1-1/tir1-1</i>	15.7	0.3	33
<i>tir1-3/tir1-3</i>	15.8	0.4	34
<i>tir1-7/tir1-7</i>	13.0	0.5	33
<i>tir1-1/+</i>	10.9	0.3	15
<i>tir1-3/+</i>	12.5	0.3	17
<i>tir1-7/+</i>	12.6	0.2	17

Ruegger et al.

reduction in auxin response. To distinguish between these possibilities, we examined the growth of *tir1-1* seedlings on media containing NPA, IAA, or the synthetic auxin, 2,4-dichlorophenoxyacetic acid (2,4-D). The results are shown in Figure 1. As expected, *tir1-1* seedlings display increased root growth on NPA compared with wild type. Mutant seedlings, however, were also less sensitive to the growth-inhibiting effects of the auxins, suggesting that the primary effect of the mutation is on auxin response and not auxin transport. Consistent with this conclusion, polar auxin transport in *tir1-1* stem segments is similar to wild type (Fig. 2). This contrasts with the behavior of the *tir3* mutant, which is resistant to NPA but not auxin and has a clear defect in auxin transport (Ruegger et al. 1997). In addition, the response of *tir1-1* roots to abscisic acid and the cytokinin benzyladenine was similar to wild type.

Cell proliferation in response to auxin-transport inhibitors is reduced in *tir1* seedlings

CPD treatment results in pronounced swelling of the root tip and expansion of the zone of cell division away from the root tip (Ruegger et al. 1997). Root tip swelling is the result of extra periclinal cell divisions in the meristem. To determine whether these events occur in *tir1* roots, we examined cross sections of wild-type and *tir1-1* roots grown on media with or without CPD. The sections shown in Figure 3 were cut 300 μm back from the root tip at a position where the different cell layers of the mature root are clearly defined. The organization of *Arabidopsis* roots is highly regular and consists of the epidermis, cortex, endodermis, pericycle, and vascular tissue. The appearance of untreated *tir1-1* roots was similar to that of untreated wild-type roots. When grown on CPD, wild-type roots exhibited a dramatic increase in cell number as expected (Ruegger et al. 1997). In con-




Figure 1. The roots of wild-type (open bars) and *tir1-1* (solid bars) seedlings are resistant to the growth-inhibiting properties of NPA, IAA, and 2,4-D. Seeds were germinated on nutrient medium. After 3 days, seedlings were transferred to media containing the indicated compound. Five days later, new root growth was measured and plotted as a percentage of root growth on medium without compound. Bars represent standard errors. Absence of bar indicates error less than thickness of line ($n = 12$).




Figure 2. Polar auxin transport is similar in wild-type (solid bars) and *tir1-1* (hatched bars) inflorescence stems. Two and one-half centimeters of stem was excised, and the apical end was placed in a nutrient solution containing $1 \mu\text{M}$ [^{14}C]IAA for the times indicated. The amount of radioactive IAA transported to the basal end of the stem was assayed by liquid scintillation. Each column represents the mean of three replicates; the bar represents the standard error of the mean.

trast, *tir1-1* roots grown on CPD displayed modest changes in cell number and organization, indicating that CPD-induced cell proliferation is reduced in *tir1-1* seedlings.

The TIR1 gene is involved in lateral root formation

Physiological and genetic studies have shown that auxin is required for lateral root initiation and growth (Wightman and Thimann 1980; Celenza et al. 1995; Hobbie and Estelle 1995). Formation of a lateral root primordium starts with a local increase in anticlinal divisions in the xylem-radius pericycle cells. This is followed by radial expansion of these pericycle cells and several periclinal divisions to form a 4-cell-layer dome-shaped primordium (Malamy and Benfey 1997). When seedlings are treated with auxin, these early events occur in a synchronized fashion along the entire length of the root (Laskowski et




Figure 3. Cell proliferation in response to CPD is reduced in the roots of *tir1-1* seedlings. Seedlings were grown for 7 days on nutrient medium plus or minus $5 \mu\text{M}$ CPD, fixed, and embedded in Spurr's (Ruegger et al. 1997). Root cross sections 300 μm from the root tip are shown. Bars, 20 μm .

An F-box protein that functions in auxin response

al. 1995). Examination of seedlings grown on agar medium indicated that *tir1* plants are deficient in lateral root formation. After 10 days of growth, wild-type seedlings had 6.3 ± 0.4 lateral roots ($n = 10$), whereas *tir1-1* seedlings had 1.3 ± 0.4 ($n = 10$). When *tir1-1* seedlings were examined by confocal microscopy, no additional lateral root primordia were observed, suggesting that mutant seedlings are deficient in an early step in formation of a lateral root (data not shown). To determine whether the *TIR1* gene functions in auxin-induced lateral root formation, we exposed excised root segments from wild-type and *tir1-1* seedlings to various concentrations of IAA or 2,4-D. The results for IAA are displayed in Figure 4A. Both compounds induced lateral root formation in wild-type and *tir1-1* root segments. The number of new lateral roots on *tir1-1* segments, however, was reduced compared with wild type. This effect can also be observed in Figure 4B. When grown on medium containing $0.5 \mu\text{M}$ 2,4-D, a continuous row of lateral roots was initiated along the length of the wild-type root. In contrast, initiation of lateral roots on *tir1-1* seedlings is sporadic. These results indicate that the *TIR1* gene functions in auxin induction of lateral root formation.

The tir1 mutants are also deficient in a cell-elongation response

Growth of *Arabidopsis* seedlings at 28°C under moderate light conditions results in a dramatic increase in hypocotyl elongation compared with seedlings grown at 20°C (Fig. 5; W.M. Gray and M. Estelle, unpubl.). After 9 days of growth, the hypocotyls of wild-type seedlings are 4.5-fold longer at 28°C compared with seedlings grown at 20°C . This increase is primarily caused by increased cell elongation because epidermal cells are 3.7-fold longer at




Figure 4. The *tir1-1* mutant is deficient in IAA-induction of lateral roots. (A) Eight-millimeter root segments were excised from 5-day-old wild-type (open bar) and *tir1-1* (solid bar) seedlings grown on nutrient medium and transferred to medium with IAA. Lateral roots were counted after 5 days by use of a dissecting microscope. Bars represent standard errors. Absence of bar indicates error less than thickness of line ($n = 10$). (B) Ten-day-old wild-type (left) and *tir1-1* (right) seedlings grown on $0.5 \mu\text{M}$ 2,4-D. (Insets) Higher magnification images of roots.




Figure 5. Hypocotyl elongation in response to elevated temperature is reduced in *tir1-1* seedlings compared with wild-type seedlings. Seedlings were grown on vertically oriented agar medium for 9 days at the two temperatures.

the higher temperature. We have shown that this response is absent in the *axr1-12* mutant and reduced in plants that are transgenic for the *iaaLys* gene and, consequently, have reduced IAA levels (W.M. Gray and M. Estelle, unpubl.). In contrast, mutants deficient in gibberellin biosynthesis or ethylene response are unaffected. These results indicate that increased hypocotyl elongation at elevated temperature is auxin dependent. To determine whether the *TIR1* gene functions in this response, we grew wild-type and *tir1-1* seedlings at the two temperatures. The results shown in Figure 5 indicate that mutant seedlings are deficient in hypocotyl elongation under these conditions. Mutant hypocotyls are 2.6-fold longer at the higher temperature (compared with 4.5 for wild type). Similarly the epidermal cells of *tir1-1* hypocotyls are 2.2-fold longer at 28°C compared with 20°C (wild-type cells are 3.7-fold longer).

The tir1 and axr1 mutants display a synergistic interaction

The *AXR1* gene is also required for auxin response (Lincoln et al. 1990; Timpte et al. 1995). To explore possible interactions between *AXR1* and *TIR1*, we generated plants that were homozygous for both the *axr1-12* and *tir1-1* mutations and examined their phenotype. When grown on medium containing 0.1 or $0.5 \mu\text{M}$ 2,4-D, *axr1-12 tir1-1* seedlings are more resistant than *axr1-12* seedlings (Fig. 6A). Because this effect occurs in a concentration range where *tir1-1* does not confer resistance by itself, the interaction between the two mutations is synergistic. A similar conclusion was reached when the morphology of double-mutant plants was examined. In most respects, the *tir1-1* rosette and inflorescence are identical in appearance to wild type (Fig. 6B). There was a slight increase in the number of secondary inflorescences growing from the rosette (wild type 1.5 ± 0.2 compared with *tir1* 2.3 ± 0.3 ; $n = 10$), suggesting an effect on apical dominance. In an *axr1-12* background, the *tir1-1*

Ruegger et al.




Figure 6. *tir1-1* and *axr1-12* have synergistic effects on auxin response and plant morphology. (A) The *tir1-1* mutation decreased auxin response in an *axr1-12* background at a concentration at which *tir1-1* has no effect by itself. Seedlings were grown and treated as described for Fig. 1. (Open bars) Wild type; (solid bars) *tir1*; (hatched bars) *axr1*; (stippled bars) *axr1 tir1*. Bars represent standard error and absence of bar indicates error less than thickness of line. (B) Seedlings (top row) were photographed after 12 days of growth on 1 µM 2,4-D. Bar, 2.5 mm. Mature plants (bottom row) were photographed 35 days after germination in soil. Bar, 5 cm.

mutation caused a further reduction in stature (Fig. 6B). This result indicates that, despite the lack of aerial phenotype, the *TIR1* gene functions in aerial structures. Further, it suggests that *TIR1* is involved in *AXR1*-mediated processes.

Isolation of the *TIR1* gene

The *tir1-9* mutant was isolated from a population of T-DNA transformed plants (Ruegger et al. 1997) generated by K. Feldmann and colleagues (Feldmann 1991). Genetic analysis indicated that the *tir1-9* line carried a single T-DNA insert that cosegregated with the *tir1-9* mutation. The left end of the T-DNA insert and flanking *Arabidopsis* sequences were cloned by plasmid rescue. Southern analysis revealed that a 5.5-kb *EcoRI-SalI* restriction fragment recovered in this fashion detected a polymorphism between wild type (WS) and the *tir1-9* mutant, confirming that this *Arabidopsis* DNA lies adjacent to the T-DNA insert (data not shown). The *EcoRI-SalI* restriction fragment was used to isolate cDNA clones. The longest cDNA recovered was 2.2 kb in length. DNA sequence analysis of the insert revealed an open reading frame (ORF) encoding a predicted protein of 594 amino acids (Fig. 7A). To confirm that this ORF corresponds to the *TIR1* gene, two *tir1* alleles were sequenced by RT-PCR. The *tir1-1* mutation is a glycine to aspartate substitution at position 147 and the *tir1-2* mutation is a glycine to aspartate change at position 441. Sequence analysis of a genomic clone spanning the cDNA indicated that the gene contains two introns. The *tir1-9* allele has a T-DNA insert in the first intron. RNA blot experiments with RNA isolated from wild-type and mutant tissues detected a transcript of ~2.2 kb in wild-type tissues. This transcript was absent in RNA from the *tir1-9* mutant (Fig. 7B). The *TIR1* transcript was detected in all tissues examined including roots, rosettes, stems, and flowers (data not shown). The *TIR1* cDNA was

placed adjacent to the CaMV 35S promoter and introduced into *tir1-1* plants. Of five independent transgenic lines examined, auxin response was restored to the wild-type level in four, providing final proof that the candidate cDNA was *TIR1* (Fig. 7C).

To determine the map position of the *TIR1* gene, a restriction fragment length polymorphism (RFLP) was identified between the Columbia and Landsberg ecotypes with the *EcoRI-SalI* fragment containing the *TIR1* gene as a probe. The polymorphism was mapped by use of 44 recombinant inbred lines (Lister and Dean 1993) to position 128 on chromosome 3, close to *nga128*.

The *TIR1* protein contains an F-box domain and leucine rich repeats

Examination of the *TIR1* amino acid sequence revealed several features of interest. First, *TIR1* contains an F-box domain (Fig. 8A; Bai et al. 1996). This motif is present in a variety of regulatory proteins in mammals and yeast including the S-phase kinase-associated protein (SKP2) and Cyclin F proteins from humans and the Cdc4 and Grr1 proteins from yeast (Bai et al. 1996). Each of these proteins binds a second protein called SKP1 in an interaction that requires the F-box (Bai et al. 1996; Li and Johnston 1997). Recent evidence has also shown that the UFO protein from *Arabidopsis* contains an F box and binds *Arabidopsis* orthologs of SKP1 called Ask1 and Ask2 (A. Samach, S. Kohalmi, G. Haughn, and W. Crosby, pers. comm.). Second, *TIR1* has 16 degenerate leucine-rich repeats (LRRs) (Fig. 8B). Similar repeats have been identified in a wide variety of proteins although the repeat consensus sequence, length, and level of redundancy vary widely among different proteins (Kobe and Deisenhofer 1995a,b). Data base searches indicate that the *TIR1* repeats are most similar to those found in SKP2, Grr1p, and an uncharacterized ORF in *Caenorhabditis elegans* called C02F5.7, a protein that also has an F box.




Figure 7. The characterization of three mutant alleles and transformation rescue of the *tir1-1* mutant with a candidate cDNA identify the *TIR1* gene. (A) Predicted amino acid sequence of the TIR1 protein. The position of the *tir1-1* and *tir1-2* mutations, both (G → D), are indicated with an asterisk above the affected residue. The GenBank accession nos. for the cDNA and genomic sequences are AF005048 and AF005047, respectively. (B) RNA blot analysis of *TIR1* transcript in *tir1* mutants. Twenty-five micrograms of total RNA isolated from 13-day-old seedlings was loaded in each lane. (C) Transformation rescue of the *tir1-1* mutant. The *p35S:TIR1* plasmid was introduced into *tir1-1* plants by vacuum infiltration. Transgenic lines were tested for auxin resistance by plating on medium containing 0.085 μ M 2,4-D.

The presence of an F box and the similarity of the LRRs in TIR1, Grr1p, SKP2, and C02F5.7 suggests that these proteins are related. A comparison of the four proteins is shown in Figure 7C. Although the biochemical function of the F-box proteins and their presumed binding partner SKP1 is uncertain, recent studies indicate that these proteins may function together as a ubiquitin-protein ligase complex called an SCF (Skp1-Cdc53-F-box protein) complex (Bai et al. 1996; Feldman et al. 1997; Li and Johnston 1997; Skowrya et al. 1997).

TIR1 is a member of a family of leucine-rich repeat F-box proteins in *Arabidopsis*

In addition to the animal and fungal proteins mentioned above, data base searches revealed the existence of a number of *TIR1*-related genes in *Arabidopsis* and rice. We have called these genes *LRF* for *LEUCINE-RICH REPEAT F-BOX*. Two of the sequences (LRF1 and LRF2) were identified as ESTs. We obtained these clones from the *Arabidopsis* Biological Resource Center (ABRC) and screened cDNA libraries in an attempt to obtain full-length clones. Sequence analysis indicated that for each gene, the longest cDNA we recovered did not contain the entire coding sequence. Because the *LRF1* and *LRF2* transcripts are similar in size to *TIR1* (data not shown), we estimate that ~50 nucleotides of coding sequence is missing from the 5' end of the two cDNAs. The se-

quence of the third gene, *LRF3*, was determined by the *Arabidopsis* Genome Initiative. LRF1 and LRF2 are 69% and 60% identical with TIR1 respectively, whereas LRF3 is 34% identical to TIR1. There is also one highly related rice EST present in the dBest database (accession no. D22807).

Discussion

TIR1 functions in auxin-dependent cell division and cell elongation

The *tir1* mutants are deficient in several auxin-mediated responses including auxin-dependent hypocotyl elongation, auxin inhibition of root elongation, CPD-mediated stimulation of cell proliferation in the root tip, and auxin stimulation of lateral root formation. In addition, *tir1* seedlings have reduced numbers of lateral roots when they are grown on medium without auxin. These defects strongly suggest that *TIR1* functions in auxin response and not auxin transport. Because both cell elongation (hypocotyl elongation) and cell division (cell proliferation at the root tip) are affected, we conclude that *TIR1* has a general role in auxin response.

The nature of the lesion in the *tir1-9* allele, and the absence of detectable full-length *TIR1* RNA in this mutant, strongly suggests that homozygous *tir1-9* seedlings completely lack TIR1 protein. Because this mutant still

participate in the activation of either ubiquitin or a ubiquitin-related protein and TIR1 could function later in the pathway as part of an E3 complex. The nature of the targeted protein is unknown. We hypothesize that auxin response depends on ubiquitin (or ubiquitin-related protein) modification of one or more regulatory proteins. Ubiquitination may then result in degradation by the proteasome (Hochstrasser 1996). This is consistent with a model proposed by Theologis and colleagues in which transcription of auxin-regulated genes is normally repressed by the action of short-lived repressor proteins (Ballas et al. 1995; Abel and Theologis 1996). Auxin may relieve this repression by stimulating degradation of the repressors in an AXR1-TIR1 dependent manner. Alternatively, the modification may alter the activity or cellular localization of a regulatory protein. For example, ligand-dependent ubiquitin modification of the α -factor receptor in yeast results in endocytosis of the receptor (Hicke and Riezman 1996). In mammalian cells, localization of the protein RanGAP1 to the nuclear pore complex depends on modification by the ubiquitin-related protein SUMO1 (Mahajan et al. 1997). Clearly, we will need to identify the substrates of the AXR1-TIR1 pathway(s) to establish a mechanism. One potential substrate is the product of the *SAR1* gene. By genetic criteria, *SAR1* has a negative role in auxin response, and acts downstream of *AXR1* (Cernac et al. 1997).

How auxin affects the pathway is also uncertain, particularly because neither the auxin receptor nor other signal transduction components have been identified. One attractive possibility is suggested by experiments of Li and Johnston (1997) on Grr1p. As mentioned above, Grr1p is required for ubiquitin-mediated degradation of the G₁ cyclins Cln1p and Cln2p in a process that appears to involve an SCF complex that contains Grr1p (Feldman et al. 1997; Li and Johnston 1997; Skowrya et al. 1997). Grr1p also plays a central role in glucose repression and glucose induction of genes encoding glucose transporters, however, a function that also requires Skp1p. When wild-type cells are grown in glucose, there is a substantial increase in the amount of Grr1p-Skp1p complex that is immunoprecipitated relative to cells grown in glycerol even though the levels of Grr1p and Skp1p are similar in the two extracts. Li and Johnston (1997) suggest that this difference may reflect a mechanism by which the cell links nutrient availability to cell cycle regulation. Because TIR1 may bind a SKP1 homolog in *Arabidopsis*, it is possible that auxin affects this interaction in some way, thereby altering the level of ubiquitin modification of one or more targets. We are currently working to identify TIR1-interacting proteins to test this model.

Summary

Genetic studies in *Arabidopsis* have led to the identification of eight genes that are required for normal auxin response (Hobbie and Estelle 1994). To date, three of these genes have been isolated. The structures of the proteins encoded by two of the genes, *AXR1* (Leyser et al. 1993) and *TIR1* (this study), suggest that they function in

a ubiquitin-protein conjugation pathway. Thus, our results provide strong support for a role for ubiquitin (or related protein) modification in auxin response. Recently, D. Xie and J. Turner (University of East Anglia, Norwich, UK) have independently shown that the *LRF3* gene described here is identical to a previously identified gene called *COI1* (J. Turner, pers. comm.). The recessive *coi1* mutants are completely insensitive to jasmonic acid, suggesting that the gene plays a key role in jasmonate signaling or response (Feys et al. 1994). It appears likely that the *TIR1/COI1* genes represent the first members of a novel class of signaling proteins in plants.

Materials and methods

Plant materials and growth conditions

All mutant lines were from the Columbia ecotype. Plants were grown at 21–23°C under continuous fluorescent illumination (100–150 $\mu\text{E}/\text{m}^2$ per sec) in 13-cm clay pots containing Metro-Mix (W.R. Grace & Co.) or an equivalent soilless mixture. During the first 3 weeks of growth, a mineral nutrient solution containing 5 mM KNO₃, 2.5 mM KPO₄ (adjusted to pH 5.8), 2 mM MgSO₄, 2 mM Ca(NO₃)₂, 50 μM Fe-EDTA, 70 μM H₃BO₃, 14 μM MnCl₂, 0.5 μM CuSO₄, 1 μM ZnSO₄, 0.2 μM Na₂MoO₄, 10 μM NaCl, and 10 nM CoCl₂ was supplied to the plants. For many experiments, plants were grown under sterile conditions in petri plates containing the above nutrient solution plus 0.7% agar, 1.0% sucrose (wt/vol), and various hormones or inhibitors as indicated in the text. Before plating, seeds were surface-sterilized by agitation for 10–20 min in 20% commercial bleach and 0.02% Triton X-100, rinsed several times with sterile water, and held at 4°C for 3–4 days to enhance germination. The seeds were dispersed on the growth media with sterile water. The plates were oriented vertically and held in an incubator containing fluorescent lighting (30–50 $\mu\text{E}/\text{m}^2$ per sec; 16 hr photoperiod) and a temperature of 20–21°C. In all experiments where plants are grown, day 0 is considered to be the time when seeds are first placed in the growth conditions described above. For hypocotyl and cell elongation studies, seedlings were grown under 24 hr light at 85 $\mu\text{E}/\text{m}^2$ per sec.

Genetic analysis

For genetic analyses, resistance to inhibition of root growth by CPD (a gift from G.F. Katekar, Commonwealth Scientific and Industrial Research Organization, Canberra City, Australia) was assayed by the following procedure. Surface-sterilized, cold-treated seeds were plated on nutrient medium, as described above, supplemented with 5 μM CPD. On day 7 (unless otherwise stated) the seedlings were straightened with forceps and the root length (distance from the root/hypocotyl junction to the root tip) was measured. Resistance to 0.2 μM 2,4-D was assayed in a similar manner except that roots were scored visually as either 2,4-D-resistant or wild type.

To generate *axr1-12 tir1-1* double mutant plants, crosses between the single mutant lines *axr1-12* and *tir1-1* were performed. The genotypes of individual F₂ plants from these crosses were determined by test crosses to both of the parental lines and tests for CPD or 2,4-D resistance as described above.

Linkage analysis was performed with a 9-kb *EcoRI-SalI* fragment of *Arabidopsis* genomic DNA containing the coding sequences of the *TIR1* gene. This fragment detected a RFLP between *EcoRV*-digested Columbia and Landsberg erecta (Ler) genomic DNA. Mapmaker I (S. Lincoln, M. Daly, J. Abrahamson,

Ruegger et al.

A. Barlow, and L. Newburg; Massachusetts Institute of Technology, Cambridge, MA) was used to compile the results of the DNA blot analysis of 44 RI lines (Lister and Dean 1993) with this fragment as a probe.

Auxin-transport assays

Auxin-transport assays were performed according to the method of Okada et al. (1991). Stem segments (2.5 cm) of primary inflorescence were incubated in a 1.5-ml Eppendorf centrifuge tube containing 30 μ l of nutrient solution with 1 μ M [14 C]IAA (1.74 nCi/ml). The segments were incubated with their apical end in the solution for various times from 2 to 18 hr. After the incubation, a 5-mm section from the basal end of the segment was excised and added to 3 ml of liquid scintillation cocktail (Bio-Safe II; RPI, Mount Pleasant, IL). The samples were shaken at 100 rpm for at least 2 hr and left overnight at room temperature before scintillation counting in a scintillation counter (model LS 6500; Beckman Instruments). The experiment involved three stem segments for each time point. Wild-type and *tir1-1* plants were grown in 24 hr of light (105 μ E/m² per sec) for 36 days. The experiment was repeated three times with similar results. Stems incubated with 15 μ M NPA or with the basal end of the stem segment in the solution transported low levels of [14 C]IAA.

Induction of lateral root formation by IAA

Seedlings were grown on nutrient medium for 5 days. Eight-millimeter root segments were excised with a razor blade from mature root and transferred to nutrient medium supplemented with various concentrations of IAA. The number of lateral roots was determined by use of a dissecting microscope and expressed as the number of lateral roots per millimeter of primary root.

Isolation of sequences flanking the T-DNA insert in *tir1-9*

Plasmid rescue of T-DNA sequences was done following the procedure of Behringer and Medford (1992). Genomic DNA, prepared from rosette leaves of the *tir1-9* mutant line, was digested with either *Sal*I or *Eco*RI to isolate DNA flanking the T-DNA left border or right border, respectively. The digested DNA was ligated with T4 DNA ligase, and electroporated into JS4 *E. coli* (Bio-Rad). Four colonies derived from the *Sal*I-digested DNA (L1–L4) and >100 colonies derived from the *Eco*RI-digested DNA were recovered. Plasmid DNA from L1–L4 and from 14 of the *Eco*RI-derived colonies (R1–R14) was prepared and analyzed by restriction analysis to eliminate clones derived entirely from the T-DNA insert (Behringer and Medford 1992). Two left border isolates (L3 and L4) had identical restriction digest patterns and potentially contained *Arabidopsis* DNA. When used as a probe in DNA blot analysis, a 5.5-kb *Eco*RI–*Sal*I fragment from L3 identified RFLPs between *tir1-9* and wild-type WS genomic DNA.

Isolation and characterization of genomic and cDNA clones

The L3 5.5-kb probe was used to isolate six genomic clones (D1–D6; library, a gift from R. Davis, Stanford University, CA) and four cDNA clones (E1–E4; library, a gift from J. Ecker, University of Pennsylvania, Philadelphia). E4, which contained the largest insert, was subcloned for sequencing (Sequenase kit, U.S. Biochemical). A 9-kb *Eco*RI–*Sal*I fragment from D1 (D109ES), which hybridized to E4, was subcloned into pBluescript (Stratagene). A 1.5-kb *Hinc*II fragment, a 2.1-kb *Sac*I *Clal* fragment, and a 1.5-kb *Hind*III fragment, which together encompassed a

3.8-kb region of D109ES containing the *TIR1* coding region, were subcloned into pBluescript and sequenced (Sequitern, Epicentre Technologies).

Characterization of *tir1* mutant alleles

RT-PCR (Stratagene) was used to amplify the expressed sequences of two *tir1* alleles from total RNA prepared from seedlings grown in liquid culture. The PCR products were separated in a low-melt agarose gel and sequenced directly by use of a modified Sequenase protocol (B. Robertson, University of Colorado). The position of the T-DNA insertion of *tir1-9* was determined by sequencing a subcloned fragment of the L3 plasmid that hybridized to probes made from the *TIR1* cDNA and from the T-DNA left border (plasmid pBSH10).

RNA blot analysis

Total RNA was isolated from 13-day-old seedlings according to Newman et al. (1993) with modifications by Timpte et al. (1995), and blotted as described in Timpte et al. (1995).

Generation of transgenic plants

The 2.2-kb *TIR1* cDNA was cloned into the *pBI121* vector (replacing the *GUS* gene) to generate the *p35S:TIR1* plasmid. This plasmid was introduced into the *Agrobacterium* strain GV3101 by electroporation. *tir1-1* plants were inoculated with this strain by vacuum infiltration according to Bechtold et al. (1993). Transformed seedlings were identified by selection on 50 μ g/ml kanamycin. Transgenic lines were tested for *TIR1* function by plating on medium that contained 0.085 mM 2,4-D.

Acknowledgments

We are grateful to members of the Estelle laboratory and the Indiana University *Arabidopsis* group for stimulating discussion throughout the course of this work, to Roger Innes for careful reading of the manuscript, to the ABRC for cDNAs, and to L. Washington for assistance with DNA sequencing. This research was supported in part by the National Science Foundation (postdoctoral fellowship MCB-9008316 to L.H., IBN-9307134 to M.E.) and National Institutes of Health (U.S. Public Health Service grant-GM43644 to M.E.) and an Eli Lilly/Indiana Institute for Molecular Biology predoctoral fellowship to M.R.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Abel, S. and A. Theologis. 1996. Early genes and auxin action. *Plant Physiol.* **111**: 9–17.
- Bai, C., P. Sen, K. Hofmann, L. Ma, M. Goebel, J.W. Harper, and S.J. Elledge. 1996. *SKP1* connects cell cycle regulators to the ubiquitin proteolysis machinery through a novel motif, the F-box. *Cell* **86**: 263–274.
- Ballas, N., L.-M. Wong, M. Ke, and A. Theologis. 1995. Two auxin-responsive domains interact positively to induce expression of the early indolacetic acid-inducible gene *PS-IAA4/5*. *Proc. Natl. Acad. Sci.* **92**: 3483–3487.
- Barbier-Brygoo, H. 1995. Tracking auxin receptors using functional approaches. *Crit. Rev. Plant Sci.* **14**: 1–25.
- Bechtold, N., J. Ellis, and G. Pelletier. 1993. *In planta Agrobac-*

- terium mediated gene transfer by infiltration of adult *Arabidopsis thaliana* plants. *C.R. Acad. Sci. Ser. III Sci. Vie* **316**: 1194–1199.
- Behringer, F.J. and J.I. Medford. 1992. A plasmid rescue technique for the recovery of plant DNA disrupted by T-DNA insertion. *Plant Mol. Biol. Rep.* **10**: 190–198.
- Bennett, M.J., A. Marchant, H.G. Green, S.T. May, S.P. Ward, P.A. Millner, A.R. Walker, B. Schulz, and K.A. Feldmann. 1996. *Arabidopsis* AUX1 gene: A permease-like regulator of root gravitropism. *Science* **273**: 948–950.
- Celenza, J.L., Jr., P.L. Grisafi, and G.R. Fink. 1995. A pathway for lateral root formation in *Arabidopsis thaliana*. *Genes & Dev.* **9**: 2131–2142.
- Cernac, A., C. Lincoln, D. Lammer, and M. Estelle. 1997. The *SAR1* gene of *Arabidopsis* acts downstream of the *AXR1* gene in auxin response. *Development* **124**: 1583–1591.
- Chow, N.W., J.R. Korenberg, X.N. Chen, and R.L. Neve. 1996. APP-BP1, a novel protein that binds to the carboxyl-terminal region of the amyloid precursor protein. *J. Biol. Chem.* **271**: 11339–11346.
- Dohmen, R.J., R. Stappen, J.P. McGrath, H. Forrova, J. Kolarov, A. Goffeau, and A. Varshavsky. 1995. An essential yeast gene encoding a homolog of ubiquitin activating enzyme. *J. Biol. Chem.* **270**: 18099–18109.
- Evans, M.L. 1984. Functions of hormones at the cellular level of organization. In *Hormonal regulation of development II, encyclopedia of plant physiology* (ed. T.K. Scott), vol. 10, pp. 23–79. Springer-Verlag, Berlin, Germany.
- Feldman, R.M., C.C. Correll, K.B. Kaplan, and R.J. Deshaies. 1997. A complex of Cdc4p, Skp1p, and Cdc53p/cullin catalyzes ubiquitination of the phosphorylated CDK inhibitor Sic1p. *Cell* **91**: 221–230.
- Feldmann, K.A. 1991. T-DNA insertion mutagenesis in *Arabidopsis*: Mutation spectrum. *Plant J.* **1**: 71–82.
- Feys, B.J.F., C.E. Benedetti, C.N. Penfold, and J.G. Turner. 1994. *Arabidopsis* mutants selected for resistance to the phytotoxin coronatine are male sterile, insensitive to methyl jasmonate, and resistant to a bacterial pathogen. *Plant Cell* **6**: 751–759.
- Hicke, L. and H. Riezman. 1996. Ubiquitination of a yeast plasma membrane receptor signals its ligand-stimulated endocytosis. *Cell* **84**: 277–287.
- Hobbie, L. and M.A. Estelle. 1994. Genetic approaches to auxin action. *Plant Cell Environ.* **17**: 525–540.
- . 1995. The *axr4* auxin-resistant mutants of *Arabidopsis thaliana* define a gene important for root gravitropism and lateral root initiation. *Plant J.* **7**: 211–220.
- Hochstrasser, M. 1995. Ubiquitin, proteasomes, and the regulation of intracellular protein degradation. *Curr. Opin. Cell Biol.* **7**: 215–223.
- Johnson, E.S., I. Schwienhorst, R.J. Dohmen, and G. Blobel. 1997. The ubiquitin-like protein Smt3p is activated for conjugation to other proteins by an Aos1p/Uba2p heterodimer. *EMBO J.* **16**: 5509–5519.
- Klee, H. and M. Estelle. 1991. Molecular genetic approaches to the biology of plant hormones. *Annu. Rev. Plant Physiol. Mol. Biol.* **42**: 529–551.
- Kobe, B. and J. Deisenhofer. 1995a. A structural basis of the interactions between leucine-rich repeats and protein ligands. *Nature* **374**: 183–185.
- . 1995b. Proteins with leucine-rich repeats. *Curr. Opin. Struct. Biol.* **5**: 409–416.
- Laskowski, M.J., M.E. Williams, H.C. Nusbaum, and I.M. Sussex. 1995. Formation of lateral root meristems is a two-stage process. *Development* **121**: 3303–3310.
- Leyser, H.M.O., C. Lincoln, C. Timpte, D. Lammer, J. Turner, and M. Estelle. 1993. The auxin-resistance gene *AXR1* of *Arabidopsis* encodes a protein related to ubiquitin-activating enzyme E1. *Nature* **364**: 161–164.
- Leyser, O. 1997. Auxin: Lessons from a mutant weed. *Physiol. Plantarum* **100**: 407–414.
- Li, F.N. and M. Johnston. 1997. Grr1 of *Saccharomyces cerevisiae* is connected to the ubiquitin proteolysis pathway through Skp1: Coupling glucose sensing to gene expression and the cell cycle. *EMBO J.* **16**: 5629–5638.
- Lincoln, C., J.H. Britton, and M. Estelle. 1990. Growth and development of the *axr1* mutants of *Arabidopsis*. *Plant Cell* **2**: 1071–1080.
- Lister, C. and C. Dean. 1993. Recombinant inbred lines for mapping RFLP and phenotypic markers in *Arabidopsis thaliana*. *Plant J.* **4**: 745–750.
- Mahajan, R., C. Delphin, T. Guan, L. Gerace, and F. Melchior. 1997. A small ubiquitin-related polypeptide involved in targeting RanGAP1 to nuclear pore complex protein RanBP2. *Cell* **88**: 97–107.
- Malamy, J.E. and P.N. Benfey. 1997. Organization and cell differentiation in lateral roots of *Arabidopsis thaliana*. *Development* **124**: 33–44.
- Muday, G.K. and P. Haworth. 1994. Tomato root growth, gravitropism and lateral root development: Correlations with auxin transport. *Plant Physiol. & Biochem.* **32**: 193–203.
- Muday, G.K., T.L. Lomax, and D.L. Rayle. 1995. Characterization of the growth and auxin physiology of roots of the tomato mutant, *diageotropica*. *Planta* **195**: 548–553.
- Nadeau, J.A., X.S. Zhang, J. Li, and S.D. O'Neill. 1996. Ovule development: Identification of stage-specific and tissue-specific cDNAs. *Plant Cell* **8**: 213–239.
- Newman, T.C., M. Ohem-takagi, C.B. Taylor, and P.J. Green. 1993. DST sequences, highly conserved among plant SAUR genes, target reporter transcripts for rapid decay in tobacco. *Plant Cell* **5**: 701–714.
- Okada, K., J. Ueda, M.K. Komaki, C.J. Bell, and Y. Shimura. 1991. Requirement of the auxin polar transport system in early stages of *Arabidopsis* floral bud formation. *Plant Cell* **3**: 677–684.
- Ruegger, M., E. Dewey, L. Hobbie, D. Brown, P. Bernasconi, J. Turner, G. Muday, and M. Estelle. 1997. Reduced NPA-binding in the *tir3* mutant of *Arabidopsis* is associated with a reduction in polar auxin transport and diverse morphological defects. *Plant Cell* **9**: 745–757.
- Shayeghi, M., C.L. Doe, M. Tavassoli, and F.Z. Watts. 1997. Characterization of a *Schizosaccharomyces pombe* RAD31, a UBA-related gene required for DNA damage tolerance. *Nucleic Acids Res.* **25**: 1162–1169.
- Skowrya, D., K.L. Craig, M. Tyers, S.J. Elledge, and J.W. Harper. 1997. F-box proteins are receptors that recruit phosphorylated substrates to the SCF ubiquitin ligase complex. *Cell* **91**: 209–219.
- Timpte, C., C. Lincoln, F.B. Pickett, J. Turner, and M. Estelle. 1995. The *AXR1* and *AUX1* genes of *Arabidopsis* function in separate auxin-response pathways. *Plant J.* **8**: 561–569.
- Venis, M.A. and R.M. Napier. 1995. Auxin receptors and auxin binding proteins. *Crit. Rev. Plant Sci.* **14**: 27–47.
- Vesper, M.J. and K.L. Kuss. 1990. Physiological evidence that the primary site of auxin action in maize coleoptiles is an intracellular site. *Planta* **182**: 486–491.
- Wightman, F. and K.V. Thimann. 1980. Hormonal factors controlling the initiation and development of lateral roots. *Physiol. Plant.* **49**: 13–20.
- Zhang, H., R. Kobayashi, K. Galaktionov, and D. Beach. 1995. p19^{Skp1} and p45^{Skp2} are essential elements of the cyclin A-CDK2 S phase kinase. *Cell* **82**: 915–925.