

---

## THE TOXICITY OF HEXANOLIC EXTRACT OF *Xylopi aethiopia* TO LARVAE OF *CULEX* *P. quiuefasciatus*

\*O.A OKE AND A.A.S. AMUSAN

Department of Biological Sciences, College of Natural Sciences,  
Federal University of Agriculture, Abeokuta.

\*Corresponding author: [olubodeoke@yahoo.com](mailto:olubodeoke@yahoo.com) Tel: +2347035432641

---

### ABSTRACT

The hexanolic extract of *Xylopi aethiopia* (Ethiopian pepper) was tested for acute toxicity on the larva of *Culex p. quiuefasciatus* reared in the laboratory of the Department of Biological Science University of Agriculture Abeokuta, Nigeria. Five concentrations: (50ppm, 100ppm, 200ppm, 300ppm and 400ppm) were evaluated for acute toxicity on the larvae and total percentage mortalities recorded at intervals of 1,12,24,48 and 96 hours in each test. Effects of sunlight and ultra violet radiation on stability of the extract's potency at 2,4, and 8 hours respectively were equally examined. The mean lethal concentration LC50 was 257ppm. Toxicity of *Xylopi aethiopia* on *Culex p. quiuefasciatus* larvae was gradual and persisted throughout the test period. Sunlight exposure has no significant effect on the potency of *Xylopi* oil while ultra-violet radiation activated its larvicidal properties.

**Key words:** *Xylopi aethiopia*, acute- toxicity, larvae, *Culex p. quiuefasciatus* and Lethal concentration.

### INTRODUCTION

Insects have attacked plants since the Devonian period. During this prolonged interaction many plants have evolved more sophisticated defence system, largely a complex array of defence chemical produced by the plant themselves (Hedin,1991). Renewed interest in botanical pesticides is motivated by three major objectives: (i) To encourage traditional use of simple formulation of locally available plant materials in pest control. (ii) To identify sources of new botanical pesticides for commercial extraction. (iii) To elucidate the chemical structure of active components of botanicals. This was based on the fact that novel metabolism may serve as model for the chemical synthesis of new pesticides with more desir-

able properties, for example the known structure of natural pyrethroids have provided a basis for the development of a range of insecticides of great practical value (Elliott *et al.*,1978).

However, in spite of the fact that a variety of secondary metabolites of plant origin afford protection against insects, only a few of these have been developed into commercial insecticides. These include margosan-O developed from neem, (Larson, 1987). Black leaf 40" produced from the by- product of tobacco industry (Merck index ,1989), Ryanex or Ryaniacide obtained from *Ryania speciosa*. Other bioactive compounds of plant origin that have been in use in pest control include rotenone from *Derris guinolizidine*,

alkaloids from many leguminosae such as Lupinus, Quassin and Pyrethrum.

It is likely that systematic survey of plant materials with the broad range of bioassay using different test species would lead to the discovery of new active compounds although one may not always come across compounds possessing toxicity against insect pests comparable to the very potent currently available insecticides. Hence there is need for continuous search for new compounds with desirable properties.

Studies on the dried fruit of *Xylopia aethiopica* have led to the isolation of a new diterpene acid, Xylopic acid (Ekong and Organ, 1968) and five other Kaurane diterpenes (Ekong and Ogan, 1968). Xylopia essential oil obtained by direct steam distillation of the derived and crushed fruits gave a strong positive reaction for aldehydes with Tollens reagent. Oji *et al.* (1991) reported that laboratory treatment of maize grain with dust or ethanolic extracts of *P. guineense* and *X. aethiopica* caused significant mortality of adult *Sitophilus zeamais* infesting grain. *X. aethiopica* has not been tested on insect's such as mosquitoes. Majority of the work earlier done on *Xylopia* had been on stored products insects. The present study aims at investigating the insecticidal properties of *Xylopia aethiopica* on larvae of *Cx. P. quiquefasciatus* in the laboratory.

## MATERIALS AND METHODS

### **Plant Extract**

Dry fruits of *Xylopia aethiopica* (Ethiopian pepper) were bought from a local market in Ile-Ogbo, Osun State Nigeria. The dried fruits were washed and later sundried. The dried fruit were ground to a fine powder using electric grinder and later exhaustively extracted with n-hexane using Soxhlet ap-

paratus. The extract was then concentrated with rotary evaporator which removed the hexane component leaving behind a viscous oil required for the analysis. Preparation of volume/volume stock solution of *Xylopia aethiopica* was by measuring out 1cm<sup>3</sup> of the extract, emulsified in about 0.003cm<sup>3</sup> or 3 drops of Tween-80. The emulsified extract was then added up to 1 litre of distilled water to form 100ppm stock solution. From the stock serial concentrations of; 50ppm, 100ppm, 200ppm, 300ppm and 400ppm were prepared.

### **Insect Material**

Sands and debris from breeding sites of *Cx. P. quiquefasciatus* were collected at the peak of dry season (January) and soaked in water. After a day, hatching of viable eggs commenced. Hatched out larvae were then removed in to clean water in a brightly coloured enamel plates using a pipette. Active swimming non feeding pupae were thereafter collected into open bottles and placed in a cage where they were left to emerge.

Mosquitoes cages of about (40 x 40 x 40)cm were made of light wooden frame with sides of mosquito netting and base of wooding board. One side of the cage is provided with sleeve for taking material in and out of the cage. Adult mosquito were allowed to feed on rabbits kept in cages after they have been starved for one to two days. Oviposition commenced after three days of the first blood meal and lasted for about 7 days. Eggs were collected dried and stored away in an incubator at 31±2°C. All the third instars larvae used in the bioassay were hatched out from the stored eggs.

**Evaluation of the effect of *Xylopia aethiopica* fruit oil on *Cx. P. quiquefasciatus* larvae.**

From each concentration 250cm<sup>3</sup> solution was measured and introduced into separate labeled 500cm<sup>3</sup> specimen bottles. Forty larvae of *Cx. P. quiquefasciatus* were then introduced each. The bottles were then covered with a mesh of 1 mm for aeration but prevents entry and exit of any insect. Each treatment was replicated five times. Effects of physicochemical parameters such as sunlight and ultraviolet irradiation were tested on the extract using the method described by Adewumi *et al* (1980). A stock solution of the *X. aethiopica* was exposed to sunlight and U.V lamp (Gallenkamp LH 530) with peak output at 366nm for 2,4 and 8 hours. For the U.V light, stock solution was placed at a distance of about 30cm from the light source. Mortalities were recorded at intervals of 1,12,24,48, and 96 hours. To estimate the 96 hour median le-

thal concentration (LC<sub>50</sub>) of the extract, the 96 hours mortality of the different concentrations were used from these data using probit regression line. Data obtained were subjected to analysis of variance at 5% level of significance and where there was a difference, Duncan's T test was used to determine whether there were significant differences between treated and untreated means.

**RESULTS**

The acute LC<sub>50</sub> of *Xylopia aethiopica* oil extract was 257 ppm (Table 1), very few deaths were recorded in the first 6 hours, percentage mortality increased with increase in time. The highest concentration of 400ppm killed only 5% of the population in 1 hour test period, 3% in 6 hours and 75% in 24 hours test period (Tables 2). Toxicity of *X. aethiopica* was observed to be gradual and lasted throughout the test period.

**Table 1: LC50 at 96 hours of *Xylopia aethiopica* applied oil extract to *Cx. P. quiquefasciatus* larvae**

Treatments	Concentration (ppm)
Untreated	257
2hrs Sunlight	200
4 hours Sunlight	199.53
8 hours Sunlight	316.23
2 hours U.V	63.1
4 hours U. V	95.5
8 hours U.V	63.1

**Table 2: Comparison of treatment mean (Duncan's test)**

Sunlight exposure	Untreated	4.600a
	2 hours	5.250a
	4 hours	4.950a
	8 hours	6.050a
	LSD	2.289
U.V	Untreated	4.600a
	2 hours	6.600a
	4 hours	4.650b
	8 hours	3.35b
	LSD	1.7467

\* Means followed by the same letter are not significantly different.( $p > 0.05$ )

Duncan's test revealed that *X. aethiopica* oil extract exposed to sunlight showed no significant difference between the untreated *X. aethiopica* oil extract and 2,4, and 8 hours sunlight exposure (Table 2). The comparative  $LC_{50}$  value presented in Table 1 revealed little differences between the treated and untreated extracts except for 8 hours treatment.

Within the first 1 hour of treatment no larvae death was recorded. At 2 hours, 60% mortality, was recorded while 47%, and 63% were recorded at 4 and 8 hours exposure respectively, compared to 75% mortality value recorded for the exposed *X. aethiopica*. The result also revealed that at 1 hour test period, 40%, 60% and 45% mortalities were recorded for 2,4 and 8 hours exposure period respectively compared to 5% mortality recorded for exposed *X. aethiopica* oil ex-

tract.

## DISCUSSIONS

Toxicity of *Xylopi aethiopica* hexane extract was gradual and persisted throughout the test period of 96 hours. The toxicity and persistence of *X. aethiopica* hexane extract is very reassuring in the control of *Cx. P. quiquefasciatus* in the field since they lay their eggs on a daily basis when and where breeding exist. The persistence will reduce the need for a daily repeated application. The toxicity of *Xylopi aethiopica* extracts to larvae of *Cx. P. quiquefasciatus* corroborates the report of Oji *et al.* (1991) that *X. aethiopica* showed significant mortality effects on adults *Sitophilus zeamais*.

Ekong and Ogan (1968) isolated a diterpene acid, Xylopic acid, which could be the major insecticide present in *X. aethiopica*.

However the mode of action of xylopic acid is still obscure. Sunlight exposure has no significant effect on the potency of *X. aethiopica* oil extract. This stability was a deviation from Wink's (1993) report that pesticide principles of plant origin are unstable under sunlight. Ultraviolet (U.V) portion of the sunlight was observed to activate the larvicidal properties of *X. aethiopica*. The finding in this study is in line with observations of Graham *et al.* (1980), Wat *et al.* (1979) and Arnason *et al.* (1981) works where sunlight was found to significantly affect the potency of various botanical pesticides.

Since *X. aethiopica* is very common in the local market, it could be very cheap source of a new pesticide, however further research is necessary to identify the active ingredient in *Xylopia aethiopica* as well as its mode of action.

## REFERENCES

- Adewunmi, C.O., Marquis, V.O., Sofowora, E.A.** 1980. Preliminary screening of some plant extracts for molluscicidal activity. *Plant Medica*, 39: 57-65
- Arnason, T., Swain, T., Wat, C.K., Graham, E.A., Partingoto, S., Towers, G.H.N.** 1981. Mosquito larvicidal activity of poly acetylenes from species of the Asteraceae. *Biochem. Systematics Ecol.*, 9: 63-65.
- Ekong, D.E.U., Ogan, A.U.** 1968. Chemistry of the constituents of *Xylopia aethiopica*. The structure of xylopic acid a new diterpene acid. *J. Chem. Soc.*, 3: 311-312.
- Elliot, M., Janes, N.F., Pottes, C.** 1978. The future of pyrethroids in insect control *Ann. Rev. Entomol.*, 23: 443-469.
- Graham, K, Graham, E.A., Towers, G.H.N** 1980. Cercaricidal activity of phenyl heptatriyne and terthienyl naturally occurring compounds in species of Asteraceae (Compositae) *Can.J. Zool.*, 58(11): 1955-1958.
- Hedin, P.A** .1991. Use of natural product in pest control in naturally occurring pest Bio Regulators. ACS Symposium Series No. 449: 162-171.
- Larson, R.O.** 1987. Development of Margosan, O.A pesticide from Neem seed. In : Proceedings 3<sup>rd</sup> int. Neem conf. Nairobi Edited by Schitterer, H. and Ascher, K.R.S Pp. 243-250.
- Oji, O., Madubuike, F.N., Nwaigbo, L.C., Oji, P.O.**1991. Protection of stored Zeamays against *Sitophilus spp* with non-toxic natural products. Potentials of *Xylopia aethiopica guineense*. *Acta agronomica Hungarica*, 41(1-2): 131-135.
- Wat, C.K., Biswas, R.K., Graham, E.A., Bohm, L., Towers, G.H.N** .1979. Ultraviolet mediated Cytotoxic activity of phenyl heptatriyne from *Bidens pilosa*. *J. Nat. Prod.*, 42(1): 103-111.
- Wink, M.** 1993. Production and application of Phytochemicals from an Agricultural Perspective. *Proc. of phytochemical Soc. Europe* 34. Phytochemical and Agriculture (T. A .Van Beek and H. Breteher eds.). June 1993. Pp 11-54.

(Manuscript received: 1st November, 2010; accepted: 20th April, 2011).